

TWISTED MODULES OVER LATTICE VERTEX ALGEBRAS

BOJKO BAKALOV AND VICTOR G. KAC

Dedicated to Professor Ivan Todorov on the occasion of his 70th birthday.

1. INTRODUCTION

A vertex algebra is essentially the same as a chiral algebra in conformal field theory [2, 11]. Vertex algebras arose naturally in the representation theory of infinite-dimensional Lie algebras and in the construction of the “moonshine module” for the Monster finite simple group [3, 10].

If V is a vertex algebra and Γ is a finite group of automorphisms of V , the subalgebra V^Γ of Γ -invariant elements in V is called an *orbifold vertex algebra*. Orbifolds play an important role in string theory; in the physics literature they were introduced in one of the earliest papers on conformal field theory [5, 6]. Recently, there have been numerous mathematical papers on orbifolds. All these papers are concerned in some way or another with the problem of describing the representations of V^Γ in terms of the vertex algebra V and the group Γ . However, the solution is known only in very special cases and it is highly nontrivial.

Let Q be an integral lattice. Then one can construct a vertex algebra V_Q called a *lattice vertex algebra* [3, 10, 13]. If $\bar{\Gamma}$ is a finite group of isometries of Q , its elements can be lifted to automorphisms of V_Q . One obtains a group $\Gamma \subset \text{Aut } V_Q$, which is a central extension of $\bar{\Gamma}$. In [1], we construct a collection of V_Q^Γ -modules, and we compute their characters and modular transformations. Some of these modules play an important role in the attempts of a conformal field theory understanding of the fractional quantum Hall effect (see [4] and the references therein).

The present paper is the first step in the construction of the V_Q^Γ -modules [1]. Here we classify the so-called *twisted V_Q -modules*; our main result is Theorem 4.2. In the case when Q is a root lattice of a simple finite-dimensional Lie algebra and σ is an element of its Weyl group, our results agree with those of [14, 15]. Some of our results were obtained independently in [16], and some special cases were studied earlier in [7, 17].

Acknowledgments. We are grateful to Ivan Todorov for many stimulating discussions and for collaboration on [1]. We thank the organizers of the Varna Workshop for inviting us to present our results and for the inspiring workshop. We also acknowledge the hospitality of the Erwin Schrödinger Institute, where some of this work was done. The first author was supported in part by the Miller Institute for Basic Research in Science. The second author was supported in part by NSF grant DMS-9970007.

2. PRELIMINARIES ON VERTEX ALGEBRAS

In this section we recall the definition of a vertex algebra and some of its properties, following the book [13]. Below z, w, \dots denote formal commuting variables. All vector spaces are over the field \mathbb{C} of complex numbers.

2.1. Local Fields. Let V be a vector space. A *field* on V is a formal power series in z, z^{-1} of the form

$$(2.1) \quad \phi(z) = \sum_{m \in \mathbb{Z}} \phi_{(m)} z^{-m-1}, \quad \phi_{(m)} \in \text{End}(V),$$

such that

$$(2.2) \quad \phi_{(m)} a = 0 \quad \text{for all } a \in V \text{ and } m \gg 0.$$

Note that all fields form a vector space invariant under ∂_z .

Two fields $\phi(z)$ and $\psi(z)$ are called *local* with respect to each other if

$$(2.3) \quad (z-w)^N [\phi(z), \psi(w)] = 0 \quad \text{for } N \gg 0$$

as a formal power series in z, z^{-1}, w, w^{-1} . We will also say that the pair (ϕ, ψ) is local. Obviously, if (ϕ, ψ) is local, then so are (ψ, ϕ) and $(\partial\phi, \psi)$.

The locality (2.3) is equivalent to the commutator formula

$$(2.4) \quad [\phi(z), \psi(w)] = \sum_{n \geq 0} (\phi(w)_{(n)} \psi(w)) \partial_w^{(n)} \delta(z-w),$$

where $\partial_w^{(n)} = \partial_w^n / n!$,

$$(2.5) \quad \delta(z-w) = \sum_{k \in \mathbb{Z}} z^k w^{-k-1}$$

is the formal δ -function, and

$$(2.6) \quad \phi(w)_{(n)} \psi(w) = \text{Res}_z (z-w)^n [\phi(z), \psi(w)], \quad n \geq 0,$$

is the n th *product* of the fields ϕ, ψ . Here, as usual, Res_z denotes the coefficient at z^{-1} . Note that the sum in (2.4) is finite.

One can define n th product of fields for any $n \in \mathbb{Z}$:

$$(2.7) \quad \begin{aligned} \phi(w)_{(n)} \psi(w) &= \text{Res}_z \phi(z) \psi(w) i_{z,w} (z-w)^n \\ &\quad - \text{Res}_z \psi(w) \phi(z) i_{w,z} (z-w)^n. \end{aligned}$$

Here $i_{z,w}$ means that we expand in the domain $|z| > |w|$, i.e.,

$$(2.8) \quad i_{z,w} (z-w)^n = \sum_{k=0}^{\infty} \binom{n}{k} z^{n-k} (-w)^k,$$

while

$$(2.9) \quad i_{w,z} (z-w)^n = \sum_{k=0}^{\infty} \binom{n}{k} z^k (-w)^{n-k}.$$

In particular, $\delta(z-w) = i_{z,w} (z-w)^{-1} - i_{w,z} (z-w)^{-1}$.

The (-1) st product is called the *normally ordered product* and is denoted by $:\phi\psi:$. One has

$$(2.10) \quad :\phi(w)\psi(w): = \phi(w)^+ \psi(w) + \psi(w) \phi(w)^-,$$

where

$$(2.11) \quad \phi(w)^+ = \sum_{m < 0} \phi_{(m)} w^{-m-1}, \quad \phi(w)^- = \sum_{m \geq 0} \phi_{(m)} w^{-m-1}.$$

One also has

$$(2.12) \quad \phi(w)_{(-n-1)} \psi(w) = :(\partial_w^{(n)} \phi(w)) \psi(w):, \quad n \geq 0.$$

2.2. Vertex Algebras.

Definition 2.1. A *vertex algebra* is a vector space V (space of states) endowed with a vector $|0\rangle \in V$ (vacuum vector), an endomorphism T (infinitesimal translation operator) and a linear map from V to the space of fields on V (state-field correspondence)

$$(2.13) \quad a \mapsto Y(a, z) = \sum_{m \in \mathbb{Z}} a_{(m)} z^{-m-1},$$

such that the following properties hold:

locality axiom: all fields $Y(a, z)$ are local with respect to each other,

translation covariance:

$$(2.14) \quad [T, Y(a, z)] = \partial_z Y(a, z),$$

vacuum axioms:

$$(2.15) \quad Y(|0\rangle, z) = \text{id}_V, \quad T|0\rangle = 0,$$

$$(2.16) \quad Y(a, z)|0\rangle - a \in zV[[z]].$$

Here are some corollaries of the definition:

$$(2.17) \quad Ta = a_{(-2)}|0\rangle, \quad Y(Ta, z) = \partial_z Y(a, z).$$

We also have the skew-symmetry

$$(2.18) \quad Y(a, z)b = e^{zT}Y(b, -z)a.$$

The most important property of a vertex algebra V is the following *Borcherds identity* (which along with the vacuum axioms provides an equivalent definition of a vertex algebra):

$$(2.19) \quad \begin{aligned} & \text{Res}_{z-w} Y(Y(a, z-w)b, w) i_{w, z-w} F(z, w) \\ &= \text{Res}_z Y(a, z) Y(b, w) i_{z, w} F(z, w) \\ & \quad - \text{Res}_z Y(b, w) Y(a, z) i_{w, z} F(z, w) \end{aligned}$$

for any $a, b \in V$ and any rational function $F(z, w)$ with poles only at $z = 0$, $w = 0$ or $z = w$.

Let us give some consequences of the Borcherds identity. Taking $F(z, w) = (z-w)^n \delta(u-z)$, viewed as a series in u, u^{-1} , we obtain

$$(2.20) \quad \begin{aligned} & Y(a, u) Y(b, w) i_{u, w} (u-w)^n - Y(b, w) Y(a, u) i_{w, u} (u-w)^n \\ &= \sum_{m=0}^{\infty} Y(a_{(m+n)} b, w) \partial_w^{(m)} \delta(u-w). \end{aligned}$$

For $n = 0$ this gives the commutator formula

$$(2.21) \quad [Y(a, u), Y(b, w)] = \sum_{m=0}^{\infty} Y(a_{(m)} b, w) \partial_w^{(m)} \delta(u-w),$$

(which implies locality). Taking Res_u of (2.20), we get

$$(2.22) \quad Y(a, w)_{(n)} Y(b, w) = Y(a_{(n)} b, w), \quad n \in \mathbb{Z}.$$

2.3. Conformal Vertex Algebras.

Definition 2.2. A vertex algebra V is *conformal* of central charge (or rank) $c \in \mathbb{C}$ if there exists a vector $\nu \in V$ (the conformal vector) with the following properties:

(i) The modes L_n , $n \in \mathbb{Z}$, of the field $L(z) \equiv Y(\nu, z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$ give a representation of the Virasoro algebra with central charge c . (The field $L(z)$ is called the energy-momentum field in CFT.)

(ii) L_{-1} is the infinitesimal translation operator T .

(iii) The operator L_0 is diagonalizable with a non-negative spectrum. (L_0 is called the energy operator or Hamiltonian.)

In a conformal vertex algebra V , (2.21) implies

$$(2.23) \quad [L_0, Y(a, z)] = zY(L_{-1}a, z) + Y(L_0a, z).$$

The operator L_0 defines a gradation of V , $V = \bigoplus_{\Delta \geq 0} V(\Delta)$, such that

$$(2.24) \quad L_0|_{V(\Delta)} = \Delta \text{id}_{V(\Delta)}.$$

For $a \in V(\Delta)$, the field $Y(a, z)$ is of *conformal weight* Δ , i.e., $\deg a_{(m)} = \Delta - m - 1$ for all $m \in \mathbb{Z}$. The field $L(z)$ is of conformal weight 2 and $\nu = L_{-2}|0\rangle$.

3. TWISTED MODULES OVER VERTEX ALGEBRAS

In this section we study twisted modules over vertex algebras (cf. [8, 10]). It seems that some of our results are new even in the untwisted case.

3.1. Definition in Terms of Borchers Identity. Let V be a vertex algebra and σ be an automorphism of V of finite order N . We let $\epsilon = e^{2\pi i/N}$ and $V_j = \{a \in V \mid \sigma a = \epsilon^{-j} a\}$, $0 \leq j \leq N - 1$.

An N -*twisted field* $\phi(z)$ on a vector space M is a formal power series in $z^{1/N}$, $z^{-1/N}$ of the form

$$(3.1) \quad \phi(z) = \sum_{m \in \frac{1}{N}\mathbb{Z}} \phi_{(m)} z^{-m-1}, \quad \phi_{(m)} \in \text{End}(M),$$

such that

$$(3.2) \quad \phi_{(m)} v = 0 \quad \text{for all } v \in M \text{ and } m \gg 0.$$

For any integer k , we will denote by $\phi(e^{2\pi i k} z)$ the field obtained from $\phi(z)$ by substituting $z^{1/N}$ with $\epsilon^k z^{1/N}$, i.e., $\phi(e^{2\pi i k} z) = \sum_{m \in \frac{1}{N}\mathbb{Z}} \phi_{(m)} e^{-2\pi i k m} z^{-m-1}$.

Definition 3.1. A σ -*twisted V -module* is a vector space M endowed with a linear map from V to the space of N -twisted fields on M ,

$$(3.3) \quad a \mapsto Y^M(a, z) = \sum_{m \in \frac{1}{N}\mathbb{Z}} a_{(m)}^M z^{-m-1},$$

such that for all $a \in V$

$$(3.4) \quad Y^M(\sigma a, z) = Y^M(a, e^{2\pi i} z),$$

$$(3.5) \quad Y^M(|0\rangle, z) = \text{id}_M,$$

and the following twisted Borcherds identity holds for any $a \in V_j$, $b \in V$, $0 \leq j \leq N-1$, and any rational function $F(z, w)$ with poles only at $z = 0$, $w = 0$ or $z = w$:

$$(3.6) \quad \begin{aligned} & \operatorname{Res}_{z=w} Y^M(Y(a, z-w)b, w) i_{w, z-w} z^{j/N} F(z, w) \\ &= \operatorname{Res}_z Y^M(a, z) Y^M(b, w) i_{z, w} z^{j/N} F(z, w) \\ & \quad - \operatorname{Res}_z Y^M(b, w) Y^M(a, z) i_{w, z} z^{j/N} F(z, w). \end{aligned}$$

Of course, for $\sigma = 1$, a 1-twisted module is called just a *module*.

Remark 3.1. The Borcherds identity (3.6) is equivalent to the following collection of identities ($a \in V_j$, $b \in V$, $c \in M$, $m \in \frac{j}{N} + \mathbb{Z}$, $n \in \mathbb{Z}$, $k \in \frac{1}{N}\mathbb{Z}$):

$$(3.7) \quad \begin{aligned} \sum_{i=0}^{\infty} \binom{m}{i} (a_{(n+i)} b)_{(m+k-i)}^M c &= \sum_{i=0}^{\infty} (-1)^i \binom{n}{i} a_{(m+n-i)}^M (b_{(k+i)}^M c) \\ & \quad - \sum_{i=0}^{\infty} (-1)^{i+n} \binom{n}{i} b_{(k+n-i)}^M (a_{(m+i)}^M c). \end{aligned}$$

When V is a conformal vertex algebra we will assume that the automorphism σ preserves the conformal vector: $\sigma(\nu) = \nu$. Then, for any σ -twisted V -module M , the modes L_n^M of $Y^M(\nu, z) = \sum_{n \in \mathbb{Z}} L_n^M z^{-n-2}$ give a representation in M of the Virasoro algebra with central charge $c = \operatorname{rank} V$.

Definition 3.2. A σ -twisted module over a conformal vertex algebra V is a σ -twisted V -module M in the sense of Definition 3.1 satisfying the additional requirement that the operator L_0^M be diagonalizable with finite-dimensional eigenspaces.

3.2. Consequences of the Definition. Let M be a σ -twisted V -module. In this subsection we will derive some consequences of Definition 3.1.

First of all, note that, by (3.4), $Y^M(a, z)z^{j/N}$ contains only integer powers of z for $a \in V_j$. For $0 \leq j \leq N-1$, we introduce the N -twisted δ -functions

$$(3.8) \quad \delta_j(z-w) = z^{j/N} w^{-j/N} \delta(z-w) = \sum_{k \in \frac{j}{N} + \mathbb{Z}} z^k w^{-k-1}.$$

Then

$$(3.9) \quad \operatorname{Res}_z Y^M(a, z) \delta_j(w-z) = Y^M(\pi_j a, w),$$

where

$$(3.10) \quad \pi_j = \frac{1}{N} \sum_{k=0}^{N-1} \epsilon^{kj} \sigma^k$$

is the projection onto V_j .

Putting $F(z, w) = z^{-j/N} (z-w)^n \delta_j(u-z)$ in the twisted Borcherds identity (3.6), we obtain a twisted version of (2.20):

$$(3.11) \quad \begin{aligned} & Y^M(a, u) Y^M(b, w) i_{u, w} (u-w)^n - Y^M(b, w) Y^M(a, u) i_{w, u} (u-w)^n \\ &= \sum_{m=0}^{\infty} Y^M(a_{(m+n)} b, w) \partial_w^{(m)} \delta_j(u-w), \quad a \in V_j, n \in \mathbb{Z}. \end{aligned}$$

The collection of identities (3.11) for all $n \in \mathbb{Z}$ is equivalent to (3.6). Note that, as in the untwisted case, all the fields $Y^M(a, z)$ are local with respect to each other,

since, letting $n = 0$ in (3.11), we get the twisted commutator formula ($a \in V_j$):

$$(3.12) \quad [Y^M(a, u), Y^M(b, w)] = \sum_{m=0}^{\infty} Y^M(a_{(m)}b, w) \partial_w^{(m)} \delta_j(u - w).$$

Let us define the n th product ($n \in \mathbb{Z}$) of two twisted fields $Y^M(a, w)$, $Y^M(b, w)$ for $a \in V_j$ as $\text{Res}_u u^{j/N} w^{-j/N}$ of the left hand side of (3.11). For $n = -1$, this definition coincides with (2.10), (2.11), but for $n < -1$ it differs from (2.12). By (3.11), we have for $a \in V_j$

$$(3.13) \quad Y^M(a, w)_{(n)} Y^M(b, w) = \sum_{m=0}^{\infty} \binom{j/N}{m} w^{-m} Y^M(a_{(m+n)}b, w).$$

In particular, letting $b = |0\rangle$ and $n = -2$ in (3.13), we get

$$(3.14) \quad Y^M(Ta, z) = \partial_z Y^M(a, z).$$

When V is a conformal vertex algebra, the commutator formula (3.12) implies:

$$(3.15) \quad [T^M, Y^M(a, z)] = \partial_z Y^M(a, z),$$

$$(3.16) \quad [L_0^M, Y^M(a, z)] = z \partial_z Y^M(a, z) + Y^M(L_0^M a, z),$$

where $T^M := L_{-1}^M$.

3.3. Associativity of Twisted Fields. We will try to invert formula (3.13), i.e., to express $Y^M(a_{(n)}b, w)$ in terms of $Y^M(a, w)_{(n)} Y^M(b, w)$. To this end, we multiply both sides of (3.13) by $z^{-n-1} w^{j/N}$ and sum over $n \in \mathbb{Z}$. Using the properties of the δ -function and Taylor's formula, we get

$$(3.17) \quad \begin{aligned} i_{w,z}(z+w)^{j/N} Y^M(Y(a, z)b, w) &= i_{z,w}(z+w)^{j/N} Y^M(a, z+w) Y^M(b, w) \\ &- \sum_{k=0}^{\infty} Y^M(b, w) a_{(k+j/N)}^M (-\partial_w)^{(k)} \delta(z - (-w)), \quad a \in V_j. \end{aligned}$$

Of course, we cannot divide (3.17) by $(z+w)^{j/N}$. Note, however, that the sum in (3.17) becomes finite when applied to any $v \in M$. Using that $(z+w)^n (-\partial_w)^{(k)} \delta(z - (-w)) = 0$ for $n > k$, we obtain the *associativity* of twisted fields

$$(3.18) \quad \begin{aligned} &i_{w,z}(z+w)^{n+j/N} Y^M(Y(a, z)b, w)v \\ &= i_{z,w}(z+w)^{n+j/N} Y^M(a, z+w) Y^M(b, w)v, \\ &a \in V_j, b \in V, v \in M, n \gg 0. \end{aligned}$$

Here we can take the minimal $n \geq 0$ such that $a_{(k+j/N)}^M v = 0$ for $k \geq n$.

Note that the powers of z and w in both sides of (3.18) are bounded from below. Therefore, we can multiply both sides by $i_{w,z}(z+w)^{-n-j/N}$ to get ($a \in V_j, b \in V, v \in M$):

$$(3.19) \quad \begin{aligned} &Y^M(Y(a, z)b, w)v \\ &= (i_{w,z}(z+w)^{-n-j/N}) i_{z,w}(z+w)^{n+j/N} Y^M(a, z+w) Y^M(b, w)v. \end{aligned}$$

3.4. Definition in Terms of Associativity. We will show that the associativity (3.18), together with (3.4), (3.5), implies the twisted Borcherds identity (3.6).

First let $b = |0\rangle$ in (3.18). Using (3.5) and the identity $Y(a, z)|0\rangle = e^{zT}a$ (cf. (2.18)), we get:

$$i_{w,z}(z+w)^{n+j/N} Y^M(e^{zT}a, w)v = i_{z,w}(z+w)^{n+j/N} Y^M(a, z+w)v.$$

Notice that the right-hand side contains only non-negative integer powers of $z+w$. This implies $Y^M(e^{zT}a, w) = i_{w,z}Y^M(a, z+w)$ and, in particular, Eq. (3.14).

Now let $a \in V_j, b \in V_k, v \in M$, and let $n, p \in \mathbb{Z}$ be such that $a_{(m+j/N)}^M v = 0$ for $m \geq n$, $b_{(m+k/N)}^M v = 0$ for $m \geq p$. Replacing in (3.18) b with $e^{uT}b$, we obtain:

$$(3.20) \quad \begin{aligned} & i_{z,w}i_{w,u}(z+w)^{n+j/N} Y^M(a, z+w)Y^M(b, u+w)v \\ & = i_{w,z}i_{w,u}i_{z,u}(z+w)^{n+j/N} Y^M(Y(a, z-u)b, u+w)v. \end{aligned}$$

Note that, if we multiply the left-hand side by $i_{w,u}(u+w)^{p+k/N}$, it will contain only non-negative integer powers of $u+w$ and hence of w . Therefore it makes sense to put $w = 0$:

$$(3.21) \quad \begin{aligned} & z^{n+j/N}u^{p+k/N} Y^M(a, z)Y^M(b, u)v \\ & = i_{w,z}i_{w,u}i_{z,u}(z+w)^{n+j/N} (u+w)^{p+k/N} \\ & \quad \times Y^M(Y(a, z-u)b, u+w)v \Big|_{w=0}. \end{aligned}$$

Interchanging the roles of a and b and using (2.18), we get:

$$(3.22) \quad \begin{aligned} & z^{n+j/N}u^{p+k/N} Y^M(b, u)Y^M(a, z)v \\ & = i_{w,z}i_{w,u}i_{u,z}(z+w)^{n+j/N} (u+w)^{p+k/N} \\ & \quad \times Y^M(Y(a, z-u)b, u+w)v \Big|_{w=0}. \end{aligned}$$

Notice that

$$i_{w,z}(z+w)^{n+j/N} = i_{w,u} \sum_{i \in \mathbb{Z}_+} \binom{n+j/N}{i} (u+w)^{n-i+j/N} (z-u)^i$$

and

$$i_{z,u}(z-u)^{-m-1} - i_{u,z}(z-u)^{-m-1} = \partial_u^{(m)} \delta(z-u), \quad m \geq 0.$$

Using this and (3.21), (3.22), we get for $l \in \mathbb{Z}$:

$$\begin{aligned} & i_{z,u}(z-u)^l z^{n+j/N} u^{p+k/N} Y^M(a, z)Y^M(b, u)v \\ & \quad - i_{u,z}(z-u)^l z^{n+j/N} u^{p+k/N} Y^M(b, u)Y^M(a, z)v \\ & = i_{w,u} \sum_{\substack{i, m \in \mathbb{Z} \\ i \geq 0, m \geq i+l}} \binom{n+j/N}{i} (u+w)^{n+p-i+(j+k)/N} \\ & \quad \times Y^M(a_{(m)}b, u+w)v \partial_u^{(m-i-l)} \delta(z-u) \Big|_{w=0}. \end{aligned}$$

From here it is easy to deduce (3.11), which implies (3.6).

Therefore, we have the following equivalent definition of a σ -twisted V -module.

Proposition 3.1. *A σ -twisted V -module is the same as a vector space M endowed with a linear map (3.3) from V to the space of N -twisted fields on M , satisfying (3.4), (3.5) and (3.18).*

4. TWISTED MODULES OVER A LATTICE VERTEX ALGEBRA

In the first subsection we introduce the main object of our study: the lattice vertex algebra V_Q . The remainder of the section is devoted to the classification of all σ -twisted V_Q -modules, where σ is an automorphism of the lattice Q .

4.1. Lattice Vertex Algebras. The purpose of this subsection is to fix the notation and review some properties of lattice vertex algebras. For more details, see [13].

Let Q be an integral lattice of rank l . We denote the bilinear form on Q by $(\cdot|\cdot)$, and write $|\alpha|^2 = (\alpha|\alpha)$ for $\alpha \in Q$. We extend the bilinear form to $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{Z}} Q$ by \mathbb{C} -bilinearity. There exists a bimultiplicative function $\varepsilon: Q \times Q \rightarrow \{\pm 1\}$ satisfying

$$(4.1) \quad \varepsilon(\alpha, \alpha) = (-1)^{|\alpha|^2(|\alpha|^2+1)/2}, \quad \alpha \in Q.$$

Then by bimultiplicativity

$$(4.2) \quad \varepsilon(\alpha, \beta)\varepsilon(\beta, \alpha) = (-1)^{(\alpha|\beta)+|\alpha|^2|\beta|^2}, \quad \alpha, \beta \in Q.$$

Introduce the twisted group algebra $\mathbb{C}_\varepsilon[Q]$: it has a basis $\{e^\alpha\}_{\alpha \in Q}$ over \mathbb{C} and multiplication

$$(4.3) \quad e^\alpha e^\beta = \varepsilon(\alpha, \beta)e^{\alpha+\beta}, \quad \alpha, \beta \in Q.$$

Let $\hat{\mathfrak{h}} = \mathfrak{h}[t, t^{-1}] \oplus \mathbb{C}K$ be the *Heisenberg current algebra*; this is a Lie algebra with the bracket

$$(4.4) \quad [ht^m, h't^n] = m\delta_{m,-n}(h|h')K, \quad [ht^m, K] = 0, \quad h, h' \in \mathfrak{h}.$$

It has a unique irreducible representation of level 1 (i.e., with $K = 1$) on the Fock space $S = S(\mathfrak{h}[t^{-1}]t^{-1})$ such that $\mathfrak{h}[t^{-1}]t^{-1}$ acts by multiplication and $\mathfrak{h}[t]1 = 0$. This representation extends to the space $V_Q = S \otimes \mathbb{C}_\varepsilon[Q]$ by

$$(4.5) \quad (ht^m)(s \otimes e^\alpha) = (ht^m + \delta_{m,0}(h|\alpha))s \otimes e^\alpha \quad \text{for } m \geq 0.$$

We define a representation of the algebra $\mathbb{C}_\varepsilon[Q]$ in V_Q by left multiplication:

$$(4.6) \quad e^\gamma(s \otimes e^\alpha) = \varepsilon(\gamma, \alpha)s \otimes e^{\alpha+\gamma}.$$

This gives rise to a representation in V_Q of the associative algebra $\mathcal{A}_Q = U(\hat{\mathfrak{h}}) \otimes \mathbb{C}_\varepsilon[Q]$, which is a “twisted” tensor product of the universal enveloping algebra $U(\hat{\mathfrak{h}})$ of $\hat{\mathfrak{h}}$ and the algebra $\mathbb{C}_\varepsilon[Q]$ by the relation

$$(4.7) \quad e^\alpha(ht^m) = (ht^m - \delta_{m,0}(\alpha|h))e^\alpha.$$

Here and further e^α (respectively ht^m) stands for $1 \otimes e^\alpha$ (respectively $ht^m \otimes 1$). The algebra \mathcal{A}_Q has a \mathbb{Z}_2 -gradation (i.e., it is an associative superalgebra) defined by

$$(4.8) \quad p(u \otimes e^\alpha) = |\alpha|^2 \pmod{2\mathbb{Z}}.$$

This induces a \mathbb{Z}_2 -gradation on V_Q :

$$(4.9) \quad p(s \otimes e^\alpha) = |\alpha|^2 \pmod{2\mathbb{Z}}.$$

Introduce the following fields on V_Q (called *currents*):

$$(4.10) \quad h(z) = \sum_{m \in \mathbb{Z}} (ht^m)z^{-m-1}, \quad h \in \mathfrak{h}.$$

Then the commutation relations (4.4) for $K = 1$ can be rewritten as

$$(4.11) \quad [h(z), h'(w)] = (h|h') \partial_w \delta(z-w), \quad h, h' \in \mathfrak{h}.$$

Hence all the fields $h(z)$ are local with respect to each other.

For $\alpha \in Q$, introduce the *vertex operator*

$$(4.12) \quad \begin{aligned} Y_\alpha(z) &= e^\alpha : \exp \int \alpha(z) : \\ &\equiv e^\alpha z^\alpha \exp \left(\sum_{n < 0} (\alpha t^n) \frac{z^{-n}}{-n} \right) \exp \left(\sum_{n > 0} (\alpha t^n) \frac{z^{-n}}{-n} \right) \end{aligned}$$

of parity $|\alpha|^2 \pmod{2\mathbb{Z}}$. The vertex operator $Y_\alpha(z)$ is a field on V_Q , and it is local with respect to $h(z)$ because

$$(4.13) \quad [h(z), Y_\alpha(w)] = (h|\alpha) Y_\alpha(w) \delta(z-w), \quad h \in \mathfrak{h}, \alpha \in Q.$$

Moreover, the fields $Y_\alpha(z)$ are local among themselves. This follows from (4.2) and the formula

$$(4.14) \quad Y_\alpha(z) Y_\beta(w) = \varepsilon(\alpha, \beta) (z-w)^{(\alpha|\beta)} e^{\alpha+\beta} : \exp(\int \alpha(z) + \int \beta(w)) :.$$

Theorem 4.1. *The fields $Y(ht^{-1}, z) = h(z)$ ($h \in \mathfrak{h}$), of parity 0, and $Y(e^\alpha, z) = Y_\alpha(z)$ ($\alpha \in Q$), of parity $p(e^\alpha) = |\alpha|^2 \pmod{2\mathbb{Z}}$, generate a vertex algebra structure on $V_Q = S \otimes \mathbb{C}_\varepsilon[Q]$ with the vacuum vector $|0\rangle = 1 \otimes 1$ and the operator T defined by*

$$(4.15) \quad [T, ht^m] = -m ht^{m-1}, \quad T e^\alpha = (\alpha t^{-1}) e^\alpha, \quad h \in \mathfrak{h}, \alpha \in Q.$$

This vertex algebra is conformal of rank $l = \text{rank } Q$ with the conformal vector

$$(4.16) \quad \nu = \frac{1}{2} \sum_{i=1}^l (a^i t^{-1})(b^i t^{-1}),$$

where $\{a^i\}, \{b^i\}$ are dual bases of \mathfrak{h} .

Let σ be an automorphism of the lattice Q of finite order N . For $0 \leq j \leq N-1$, let

$$(4.17) \quad \mathfrak{h}_j = \{h \in \mathfrak{h} \mid \sigma h = \epsilon^{-j} h\}, \quad \epsilon = e^{2\pi i/N}.$$

Then \mathfrak{h}_j and \mathfrak{h}_k are orthogonal unless $j+k \equiv 0 \pmod{N\mathbb{Z}}$. Since there exists a unique, up to equivalence, ± 1 -valued 2-cocycle ε on Q satisfying (4.1), the cocycles $\varepsilon(\alpha, \beta)$ and $\varepsilon(\sigma\alpha, \sigma\beta)$ are equivalent. Hence there exists a function $\eta: Q \rightarrow \{\pm 1\}$ such that

$$(4.18) \quad \eta(\alpha)\eta(\beta)\varepsilon(\alpha, \beta) = \eta(\alpha+\beta)\varepsilon(\sigma\alpha, \sigma\beta), \quad \alpha, \beta \in Q.$$

Moreover, η can be chosen in such a way that

$$(4.19) \quad \eta(\alpha) = 1 \quad \text{for } \alpha \in Q \cap \mathfrak{h}_0.$$

Proposition 4.1. *Any automorphism σ of Q can be lifted to an automorphism of the vertex algebra V_Q so that*

$$(4.20) \quad \sigma(ht^m) = \sigma(h)t^m, \quad \sigma(e^\alpha) = \eta(\alpha)^{-1} e^{\sigma\alpha}, \quad h \in \mathfrak{h}, \alpha \in Q.$$

It fixes the conformal vector ν : $\sigma(\nu) = \nu$.

Proof. Follows from the observation that σ defines an automorphism of the associative superalgebra \mathcal{A}_Q . \square

Remark 4.1. If $\sigma: Q \rightarrow Q$ is an automorphism of order N , then its lifting $\sigma: V_Q \rightarrow V_Q$ defined by (4.20) has order N or $2N$.

4.2. Twisted Vertex Operators. Let σ be an automorphism of Q lifted to an automorphism of V_Q , as in Proposition 4.1. We use the notation from Section 3 for $V = V_Q$. Let M be a σ -twisted V_Q -module (see Definition 3.2). We will study the fields $Y^M(ht^{-1}, z)$ and $Y^M(e^\alpha, z)$ for $h \in \mathfrak{h}, \alpha \in Q$.

Comparing the commutator formulas (2.21) and (3.12), we see that Eqs. (4.11), (4.13) immediately imply

$$(4.21) \quad [Y^M(ht^{-1}, z), Y^M(h't^{-1}, w)] = (h|h') \partial_w \delta_j(z-w),$$

$$(4.22) \quad [Y^M(ht^{-1}, z), Y^M(e^\alpha, w)] = (h|\alpha) Y^M(e^\alpha, w) \delta_j(z-w),$$

$$h \in \mathfrak{h}_j, h' \in \mathfrak{h}, \alpha \in Q.$$

These formulas can be restated as:

$$(4.23) \quad [h_{(m)}^M, (h')_{(n)}^M] = (\pi_{Nm} h|h') m \delta_{m,-n},$$

$$(4.24) \quad [h_{(m)}^M, Y^M(e^\alpha, w)] = (\pi_{Nm} h|\alpha) w^m Y^M(e^\alpha, w),$$

$$h, h' \in \mathfrak{h}, m, n \in \frac{1}{N}\mathbb{Z}, \alpha \in Q,$$

where π_j is the projection of \mathfrak{h} onto $\mathfrak{h}_{j \bmod N\mathbb{Z}}$. (Note that this π_j is the restriction to $(\mathfrak{h}t^{-1})|0$ of the π_j defined by (3.10)). In the sequel we will use the notation $h_0 = \pi_0 h$ for $h \in \mathfrak{h}$.

Also note that, for $a = ht^{-1}$, (3.4) is equivalent to

$$(4.25) \quad (\sigma h)_{(m)}^M = h_{(m)}^M e^{-2\pi i m}, \quad h \in \mathfrak{h}, m \in \frac{1}{N}\mathbb{Z}.$$

Lemma 4.1. *There exist operators U_α^M ($\alpha \in Q$) on M such that*

$$(4.26) \quad Y^M(e^\alpha, z) = z^{b_\alpha} U_\alpha^M E_\alpha^M(z)$$

where

$$(4.27) \quad b_\alpha = (|\alpha_0|^2 - |\alpha|^2)/2$$

and

$$(4.28) \quad E_\alpha^M(z) = : \exp \int Y^M(\alpha t^{-1}, z) :$$

$$\equiv z^{\alpha_{(0)}^M} \exp\left(\sum_{n \in \frac{1}{N}\mathbb{Z}_{<0}} \alpha_{(n)}^M \frac{z^{-n}}{-n}\right) \exp\left(\sum_{n \in \frac{1}{N}\mathbb{Z}_{>0}} \alpha_{(n)}^M \frac{z^{-n}}{-n}\right).$$

The operators U_α^M satisfy

$$(4.29) \quad [h_{(m)}^M, U_\alpha^M] = \delta_{m,0} (h_0|\alpha) U_\alpha^M, \quad h \in \mathfrak{h}, m \in \frac{1}{N}\mathbb{Z},$$

$$(4.30) \quad U_{\sigma\alpha}^M = \eta(\alpha) U_\alpha^M e^{2\pi i (b_\alpha + \alpha_{(0)}^M)}.$$

Proof. Define the operators

$$U_\alpha^M(z) = \exp\left(\sum_{n \in \frac{1}{N}\mathbb{Z}_{<0}} \alpha_{(n)}^M \frac{z^{-n}}{n}\right) Y^M(e^\alpha, z) z^{-\alpha_{(0)}^M} \exp\left(\sum_{n \in \frac{1}{N}\mathbb{Z}_{>0}} \alpha_{(n)}^M \frac{z^{-n}}{n}\right).$$

Then, by (4.23), (4.24),

$$[h_{(m)}^M, U_\alpha^M(z)] = \delta_{m,0} (h_0|\alpha) U_\alpha^M(z), \quad h \in \mathfrak{h}.$$

In particular, $U_\alpha^M(z)$ commutes with $\alpha_{(n)}^M$ for $n < 0$; hence, we have

$$(4.31) \quad Y^M(e^\alpha, z) = U_\alpha^M(z)E_\alpha^M(z).$$

We will deduce Eqs. (4.26) and (4.29) once we show that $U_\alpha^M := z^{-b_\alpha}U_\alpha^M(z)$ is independent of z . To this end we use the translation invariance (3.14). By (4.15), we have $Te^\alpha = (\alpha t^{-1})e^\alpha = \alpha_{(-1)}e^\alpha$. Using (3.14), (4.15) and (3.13), we find

$$\begin{aligned} \partial_z Y^M(e^\alpha, z) &= Y^M(Te^\alpha, z) = Y^M(\alpha_{(-1)}e^\alpha, z) \\ &= :Y^M(\alpha t^{-1}, z)Y^M(e^\alpha, z): - \sum_{j=0}^{N-1} \frac{j}{N} (\pi_j \alpha | \alpha) z^{-1} Y^M(e^\alpha, z). \end{aligned}$$

Therefore

$$\partial_z U_\alpha^M(z) = - \sum_{j=0}^{N-1} \frac{j}{N} (\pi_j \alpha | \alpha) z^{-1} U_\alpha^M(z).$$

Using that $(\pi_j \alpha | \alpha) = (\pi_{N-j} \alpha | \alpha)$ for $1 \leq j \leq N-1$, it is easy to see that

$$(4.32) \quad \sum_{j=0}^{N-1} \frac{j}{N} (\pi_j \alpha | \alpha) = -b_\alpha.$$

This proves that $U_\alpha^M = z^{-b_\alpha}U_\alpha^M(z)$ is independent of z .

Finally, to prove (4.30), we apply (3.4) for $a = e^\alpha$, using that $\sigma(e^\alpha) = \eta(\alpha)^{-1}e^{\sigma\alpha}$ and that by (4.25) we have $E_{\sigma\alpha}^M(z) = e^{-2\pi i \alpha_{(0)}^M} E_\alpha^M(e^{2\pi i} z)$. \square

Lemma 4.2. *We have*

$$(4.33) \quad Y^M(e^\alpha, z)Y^M(e^\beta, w) = i_{z,w} f_{\alpha,\beta}(z, w) z^{b_\alpha} w^{b_\beta} U_\alpha^M U_\beta^M E_{\alpha,\beta}^M(z, w),$$

where

$$(4.34) \quad f_{\alpha,\beta}(z, w) = \prod_{k=0}^{N-1} (z^{1/N} - \epsilon^k w^{1/N})^{(\sigma^k \alpha | \beta)},$$

$$(4.35) \quad E_{\alpha,\beta}^M(z, w) = : \exp(\int Y^M(\alpha t^{-1}, z) + \int Y^M(\beta t^{-1}, w)) :.$$

Proof. Standard exercise, using the fact that $e^A e^B e^{-A} = e^{\text{ad } A} e^B = e^{[A,B]} e^B$ for any two operators A, B commuting with $[A, B]$. \square

Lemma 4.3. *We have*

$$(4.36) \quad \begin{aligned} Y^M(Y(e^\alpha, z)e^\beta, w) \\ = \varepsilon(\alpha, \beta) B_{\alpha,\beta}^{-1} i_{w,z} f_{\alpha,\beta}(z+w, w) (z+w)^{b_\alpha} w^{b_\beta} U_{\alpha+\beta}^M E_{\alpha,\beta}^M(z+w, w), \end{aligned}$$

where

$$(4.37) \quad B_{\alpha,\beta} = \frac{f_{\alpha,\beta}(z, w)}{(z-w)^{(\alpha|\beta)}} \Big|_{z^{1/N}=w^{1/N}=1} = N^{-(\alpha|\beta)} \prod_{k=1}^{N-1} (1 - \epsilon^k)^{(\sigma^k \alpha | \beta)}.$$

Proof. We use the same argument as in the proof of Lemma 4.1. First, using (3.12), (4.24) and (4.5), we compute the commutator ($h \in \mathfrak{h}_j$, $n \in \frac{j}{N} + \mathbb{Z}$):

$$\begin{aligned} & [h_{(n)}^M, Y^M(Y(e^\alpha, z)e^\beta, w)] \\ &= \sum_{m=0}^{\infty} \binom{n}{m} w^{n-m} Y^M(h_{(m)}(Y(e^\alpha, z)e^\beta), w) \\ &= \sum_{m=0}^{\infty} \binom{n}{m} w^{n-m} ((h|\alpha)z^m + \delta_{m,0}(h|\beta)) Y^M(Y(e^\alpha, z)e^\beta, w) \\ &= (i_{w,z}(z+w)^n(h|\alpha) + w^n(h|\beta)) Y^M(Y(e^\alpha, z)e^\beta, w). \end{aligned}$$

It follows that

$$Y^M(Y(e^\alpha, z)e^\beta, w) = U_{\alpha,\beta}^M(z, w) E_{\alpha,\beta}^M(z+w, w),$$

where the operator $U_{\alpha,\beta}^M(z, w)$ satisfies

$$[h_{(n)}^M, U_{\alpha,\beta}^M(z, w)] = \delta_{n,0}(h_0|\alpha + \beta) U_{\alpha,\beta}^M(z, w).$$

Next, we note that (by (4.15), (3.14), (2.14))

$$\begin{aligned} Y^M(Y(e^\alpha, z)(\beta_{(-1)}e^\beta), w) &= Y^M(Y(e^\alpha, z)(Te^\beta), w) \\ &= (\partial_w - \partial_z) Y^M(Y(e^\alpha, z)e^\beta, w). \end{aligned}$$

A similar computation as above, using (3.13) for $n = -1$, shows that for $h \in \mathfrak{h}_j$ one has

$$\begin{aligned} Y^M(Y(e^\alpha, z)(h_{(-1)}e^\beta), w) &= :Y^M(ht^{-1}, w)Y^M(Y(e^\alpha, z)e^\beta, w): \\ &\quad - \left(z^{-1}i_{w,z} \left(1 + \frac{z}{w} \right)^{j/N} (h|\alpha) + \frac{j}{N} w^{-1} (h|\beta) \right) Y^M(Y(e^\alpha, z)e^\beta, w). \end{aligned}$$

Therefore

$$\begin{aligned} & (\partial_w - \partial_z) U_{\alpha,\beta}^M(z, w) \\ &= - \sum_{j=0}^{N-1} \left(z^{-1}i_{w,z} \left(1 + \frac{z}{w} \right)^{j/N} (\pi_j \beta|\alpha) + \frac{j}{N} w^{-1} (\pi_j \beta|\beta) \right) U_{\alpha,\beta}^M(z, w). \end{aligned}$$

With some more computation, we see that

$$\begin{aligned} - \sum_{j=0}^{N-1} z^{-1}i_{w,z} \left(1 + \frac{z}{w} \right)^{j/N} (\pi_j \beta|\alpha) &= i_{w,z} i_{w,z+w} \sum_{k=0}^{N-1} \frac{(1/N)\epsilon^k w^{1/N-1} (\sigma^k \alpha|\beta)}{(z+w)^{1/N} - \epsilon^k w^{1/N}} \\ &= i_{w,z} \frac{(\partial_w - \partial_z) f_{\alpha,\beta}(z+w, w)}{f_{\alpha,\beta}(z+w, w)} \end{aligned}$$

On the other hand, by (4.32), we have

$$- \sum_{j=0}^{N-1} \frac{j}{N} w^{-1} (\pi_j \beta|\beta) = (\partial_w - \partial_z) w^{b_\beta} / w^{b_\beta}.$$

It follows that

$$U_{\alpha,\beta}^M(z, w) / i_{w,z} f_{\alpha,\beta}(z+w, w) w^{b_\beta}$$

depends only on $z+w$.

Finally, note that

$$Y^M(Y(e^\alpha, z)e^\beta, w) = z^{(\alpha|\beta)}\varepsilon(\alpha, \beta)Y^M(e^{\alpha+\beta}, w) + \text{higher powers of } z,$$

while

$$i_{w,z}f_{\alpha,\beta}(z+w, w) = z^{(\alpha|\beta)}B_{\alpha,\beta}w^{(\alpha_0-\alpha|\beta)} + \text{higher powers of } z.$$

Since $(\alpha_0 - \alpha|\beta) = b_{\alpha+\beta} - b_\alpha - b_\beta$, this completes the proof. \square

Corollary 4.1. *In any σ -twisted V_Q -module M , one has*

$$(4.38) \quad U_\alpha^M U_\beta^M = \varepsilon(\alpha, \beta)B_{\alpha,\beta}^{-1} U_{\alpha+\beta}^M, \quad \alpha, \beta \in Q.$$

Proof. Follows immediately from (4.33), (4.36) and the associativity (3.18). \square

Remark 4.2. In the proofs of Lemmas 4.2 and 4.3, we used only the commutator formulas (4.23), (4.24), the translation invariance (3.14), and formula (3.13) for $n = -1$, $a = ht^{-1}$, $b = e^\beta$.

4.3. The Heisenberg Pair $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$. The results of the previous subsection motivate the following definitions.

The σ -twisted current algebra $\hat{\mathfrak{h}}_\sigma$ consists of all σ -invariant elements from $\mathbb{C}K \oplus \mathfrak{h}[t^{1/N}, t^{-1/N}]$, where σ acts as

$$(4.39) \quad \sigma(ht^m) = \sigma(h)e^{2\pi im}t^m, \quad \sigma(K) = K, \quad h \in \mathfrak{h}, \quad m \in \frac{1}{N}\mathbb{Z}.$$

In other words, $\hat{\mathfrak{h}}_\sigma$ is spanned over \mathbb{C} by K and the elements ht^m such that $h \in \mathfrak{h}_j$, $m \in \frac{j}{N} + \mathbb{Z}$. This is a Lie algebra with bracket (4.4).

Let $G = \mathbb{C}^\times \times \exp \mathfrak{h}_0 \times Q$ be the set consisting of elements $c e^h U_\alpha$ ($c \in \mathbb{C}^\times$, $h \in \mathfrak{h}_0$, $\alpha \in Q$). We define a multiplication in G by the formulas:

$$(4.40) \quad e^h e^{h'} = e^{h+h'},$$

$$(4.41) \quad e^h U_\alpha e^{-h} = e^{(h|\alpha)} U_\alpha,$$

$$(4.42) \quad U_\alpha U_\beta = \varepsilon(\alpha, \beta)B_{\alpha,\beta}^{-1} U_{\alpha+\beta}.$$

Then G is a group. From (4.42) we get the commutator

$$(4.43) \quad C_{\alpha,\beta} := U_\alpha U_\beta U_\alpha^{-1} U_\beta^{-1} = (-1)^{|\alpha|^2|\beta|^2} \prod_{k=0}^{N-1} (-\epsilon^k)^{-(\sigma^k \alpha|\beta)}.$$

We give another expression for $C_{\alpha,\beta}$ which will be useful in the sequel:

$$(4.44) \quad C_{\alpha,\beta} = (-1)^{|\alpha|^2|\beta|^2} e^{\pi i(\alpha_0|\beta)} e^{2\pi i(\alpha_*|\beta)}$$

for $\alpha = \alpha_0 + (1 - \sigma)\alpha_*$, $\alpha_0 \in \mathfrak{h}_0$, $\alpha_* \in \mathfrak{h}_0^\perp$.

Denote by Q_{ev} the sublattice of Q consisting of all even elements, i.e., α such that $|\alpha|^2 \in 2\mathbb{Z}$.

Lemma 4.4. *The center $Z(G)$ of G consists of all elements of the form $c e^{2\pi i\lambda_0} U_{(1-\sigma)\lambda}$, where $c \in \mathbb{C}^\times$ and $\lambda \in (Q_{\text{ev}})^*$ is such that $\alpha := (1-\sigma)\lambda \in Q$ and $\lambda \in Q^*$ if $\alpha \in Q_{\text{ev}}$.*

Proof. If $e^{2\pi ih} U_\alpha$ is in the center, then (4.41) implies $\alpha_0 = 0$. Then $\alpha = (1-\sigma)\alpha_*$ for some uniquely defined $\alpha_* \in \mathfrak{h}_0^\perp$. Letting $\lambda = h + \alpha_*$, we get $e^{2\pi ih} U_\alpha = e^{2\pi i\lambda_0} U_{(1-\sigma)\lambda}$. Using (4.41) and (4.44), we see that $e^{2\pi i\lambda_0} U_{(1-\sigma)\lambda}$ commutes with U_β iff

$$(4.45) \quad (\lambda|\beta) + |\alpha|^2|\beta|^2/2 \in \mathbb{Z} \quad \text{for } \beta \in Q.$$

Since

$$|\alpha|^2 = ((1 - \sigma)\alpha_* | (1 - \sigma)\alpha_*) = 2(\alpha_* | \alpha_*) - 2(\alpha_* | \sigma\alpha_*) = 2(\alpha_* | \alpha) = 2(\lambda | \alpha),$$

equation (4.45) is equivalent to $\lambda \in Q^*$ if $\alpha \in Q_{\text{ev}}$ and to $\lambda \in (Q_{\text{ev}})^*$ if $\alpha \in Q \setminus Q_{\text{ev}}$. \square

In particular, by Lemma 4.4, all elements of the form $e^{2\pi i \alpha_0} U_{(1-\sigma)\alpha}$ ($\alpha \in Q$) are central in G . We let G_σ be the factor of G over the central subgroup

$$(4.46) \quad N_\sigma := \{\eta(\alpha) U_{\sigma\alpha}^{-1} U_\alpha e^{2\pi i (b_\alpha + \alpha_0)} \mid \alpha \in Q\}.$$

Note that $N_\sigma \cap \mathbb{C}^\times = \{1\}$.

We endow Q with the discrete topology so that G and G_σ are Lie groups with a Lie algebra $\mathbb{C} \oplus \mathfrak{h}_0$. By (4.46), (4.27) and (4.19), we have

$$(4.47) \quad e^{2\pi i \alpha} = 1 \quad \text{in } G_\sigma \text{ for } \alpha \in Q \cap \mathfrak{h}_0.$$

It is easy to see that the connected component of the unit in G_σ is equal to \mathbb{C}^\times times the torus

$$(4.48) \quad T_\sigma := \exp 2\pi i (\mathfrak{h}_0 / Q \cap \mathfrak{h}_0).$$

The group G acts on $\hat{\mathfrak{h}}_\sigma$ by conjugation:

$$(4.49) \quad (ce^h U_\alpha)(h't^m + c'K)(ce^h U_\alpha)^{-1} = h't^m + \delta_{m,0}(h'_0 | \alpha)K + c'K.$$

This action is compatible with the adjoint action of $\mathbb{C} \oplus \mathfrak{h}_0$ on $\hat{\mathfrak{h}}_\sigma$ (which is trivial), hence, $(\hat{\mathfrak{h}}_\sigma, G)$ is a *Heisenberg pair* in the sense of [9]. The same is true for $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ because N_σ acts trivially on $\hat{\mathfrak{h}}_\sigma$. A module M over $\hat{\mathfrak{h}}_\sigma$ or over $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ will be called *restricted* if the action of \mathfrak{h}_0 is diagonalizable and for any $v \in M$, $(ht^m)v = 0$ for $h \in \mathfrak{h}$ and sufficiently large $m \in \frac{1}{N}\mathbb{Z}$.

Now we can summarize the results of Section 4.2 as follows.

Proposition 4.2. *Any σ -twisted V_Q -module M is naturally a restricted module over the Heisenberg pair $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ of level 1 (i.e., both $K \in \hat{\mathfrak{h}}_\sigma$ and $1 \in G_\sigma$ act as 1). Conversely, any restricted $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ -module of level 1 can be endowed with the structure of a σ -twisted V_Q -module. This establishes an equivalence of the corresponding abelian categories.*

Proof. 1. Let M be a σ -twisted V_Q -module. By definition, the action of L_0^M is diagonalizable with finite-dimensional eigenspaces. Since \mathfrak{h}_0 commutes with L_0^M , its action is diagonalizable too. By (4.23), the modes $h_{(m)}^M$ ($h \in \mathfrak{h}$, $m \in \frac{1}{N}\mathbb{Z}$) provide a restricted representation of $\hat{\mathfrak{h}}_\sigma$ of level 1. The action of $U_\alpha \in G_\sigma$ is given by the operator U_α^M , see (4.29), (4.30), (4.38).

2. Conversely, let M be a restricted $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ -module of level 1. Denote the image of ht^m in $\text{End } M$ by $h_{(m)}^M$, and that of U_α by U_α^M . This allows us to define the fields $Y^M(ht^{-1}, z)$ and then $Y^M(e^\alpha, z)$ by (4.26) ($h \in \mathfrak{h}$, $\alpha \in Q$), They satisfy (3.4), (3.12); in particular, they are local. The map Y^M can be extended uniquely to the whole V_Q by applying (3.19) repeatedly for $a \in \mathfrak{h}t^{-1}$. Note that (3.19) implies the translation invariance (3.14) for $a \in \mathfrak{h}t^{-1}$. Then the proof of Lemma 4.1 shows that (3.14) holds for $a = e^\alpha$, and hence for any $a \in V_Q$. It follows from Remark 4.2 that Lemmas 4.2 and 4.3 hold. This implies the associativity (3.18). By Proposition 3.1, M is a σ -twisted V_Q -module. \square

4.4. **The Groups G^\perp and G_σ^\perp .** Before we proceed to the classification of all restricted $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ -modules of level 1, we need to study the groups G and G_σ in more detail.

Let $G^\perp \subset G$ be the subgroup of G consisting of all cU_α with $c \in \mathbb{C}^\times$, $\alpha \in Q \cap \mathfrak{h}_0^\perp$. Clearly, the centralizer of $\hat{\mathfrak{h}}_\sigma$ in G equals $\exp \mathfrak{h}_0 \times G^\perp$ (cf. (4.49)). In other words, G^\perp is the outer centralizer of the torus $\exp \mathfrak{h}_0$ in G .

Denote by G_σ^\perp the image of G^\perp in G_σ . It can be described as the factor of G^\perp over the central subgroup (cf. (4.46), (4.27))

$$(4.50) \quad N_\sigma^\perp := N_\sigma \cap G^\perp = \{\eta(\alpha)(-1)^{|\alpha|^2} U_{\sigma\alpha}^{-1} U_\alpha \mid \alpha \in Q \cap \mathfrak{h}_0^\perp\}.$$

The centralizer of $\hat{\mathfrak{h}}_\sigma$ (and of T_σ) in G_σ is equal to $T_\sigma \times G_\sigma^\perp$. Notice that G_σ^\perp is a central extension (by \mathbb{C}^\times) of the finite abelian group $(Q \cap \mathfrak{h}_0^\perp)/(1-\sigma)(Q \cap \mathfrak{h}_0^\perp)$.

Definition 4.1. (i) Let P_σ be the set of all λ that appear in Lemma 4.4, i.e., the set of all $\lambda \in (Q_{\text{ev}})^*$ such that $(1-\sigma)\lambda \in Q$ and $\lambda \in Q^*$ if $(1-\sigma)\lambda \in Q_{\text{ev}}$. Note that P_σ is a sublattice of $(Q_{\text{ev}})^*$ containing Q .

(ii) Let $Q_\sigma = (1-\sigma)P_\sigma \subset Q$.

(iii) Let $Z_\sigma = P_\sigma/Q$ be the subgroup of $((Q_{\text{ev}})^*/Q)^\sigma$ consisting of classes $\lambda + Q$ such that $\lambda \in Q^*$ if $(1-\sigma)\lambda \in Q_{\text{ev}}$. In particular, when the lattice Q is even, $Z_\sigma = (Q^*/Q)^\sigma$ is the group of σ -invariant elements in Q^*/Q .

Similarly to Lemma 4.4, we can describe the centers of G_σ , G^\perp and G_σ^\perp .

Lemma 4.5. (i) $Z(G_\sigma) \simeq Z(G)/N_\sigma \simeq \mathbb{C}^\times \times Z_\sigma$.

(ii) $Z(G^\perp) = \{cU_\alpha \mid c \in \mathbb{C}^\times, \alpha \in Q_\sigma\} \simeq \mathbb{C}^\times \times Q_\sigma$.

(iii) $Z(G_\sigma^\perp) \simeq Z(G^\perp)/N_\sigma^\perp \simeq \mathbb{C}^\times \times Q_\sigma/(1-\sigma)(Q \cap \mathfrak{h}_0^\perp)$.

Proof. (i) follows from Lemma 4.4, (4.44) and the fact that $N_\sigma \cap \mathbb{C}^\times = \{1\}$.

(ii) A similar argument as in the proof of Lemma 4.4 shows that the center of G^\perp consists of all elements of the form $cU_{(1-\sigma)\lambda}$, where $c \in \mathbb{C}^\times$ and $\lambda \in (Q_{\text{ev}} \cap \mathfrak{h}_0^\perp)^*$ is such that $\alpha := (1-\sigma)\lambda \in Q$ and $\lambda \in (Q \cap \mathfrak{h}_0^\perp)^*$ if $\alpha \in Q_{\text{ev}}$. Next, we use the following lemma.

Lemma 4.6. *Let \mathfrak{h}' be a subspace of \mathfrak{h} such that the restriction of the bilinear form on it is nondegenerate. Denote by π' the orthogonal projection of \mathfrak{h} onto \mathfrak{h}' . Then for any lattice $L \subset \mathfrak{h}$, one has $L^* \cap \mathfrak{h}' = (\pi'L)^{*\prime}$, where $^{*\prime}$ means that the dual is taken in \mathfrak{h}' .*

Proof. Follows from the fact that $(h|\alpha) = (h|\pi'\alpha)$ for $h \in \mathfrak{h}'$, $\alpha \in L$. \square

Now, by Lemma 4.6, $(Q \cap \mathfrak{h}_0^\perp)^* = \pi_\perp(Q^*)$, and similarly for Q_{ev} , where π_\perp is the orthogonal projection from \mathfrak{h} to \mathfrak{h}_0^\perp . Noting that $(1-\sigma)\pi_\perp = 1-\sigma$ completes the proof of part (ii).

Part (iii) follows from part (ii), (4.44) and the fact that $N_\sigma^\perp \cap \mathbb{C}^\times = \{1\}$. \square

Corollary 4.2. *There is a natural exact sequence*

$$(4.51) \quad 1 \rightarrow \exp 2\pi i(Q^* \cap \mathfrak{h}_0) \rightarrow Z(G) \xrightarrow{p} Z(G^\perp) \rightarrow 1,$$

where the homomorphism p is given by

$$(4.52) \quad c e^{2\pi i \lambda_0} U_{(1-\sigma)\lambda} \xrightarrow{p} c U_{(1-\sigma)\lambda}, \quad c \in \mathbb{C}^\times, \lambda \in P_\sigma.$$

The sequence (4.51) splits, so we have a non-canonical isomorphism $Z(G) \simeq \exp 2\pi i(Q^* \cap \mathfrak{h}_0) \times Z(G^\perp)$.

Proof. Clearly, the kernel of p consists of $e^{2\pi i\lambda_0}$ with $\lambda \in Q^*$ such that $(1-\sigma)\lambda = 0$. Let $s: \pi_\perp(P_\sigma) \rightarrow P_\sigma$ be a linear section of the projection π_\perp . For $\lambda' \in \pi_\perp(P_\sigma)$, let $\lambda = s(\lambda') \in P_\sigma$. Then $\lambda' = \pi_\perp(\lambda)$ and $(1-\sigma)\lambda = (1-\sigma)\lambda'$. The map $cU_{(1-\sigma)\lambda'} \mapsto ce^{2\pi i\lambda_0}U_{(1-\sigma)\lambda}$ is a splitting of (4.51). \square

Since G_σ^\perp is a central extension of a finite abelian group, its representations are completely reducible and the irreducible ones are classified by the characters of $Z(G_\sigma^\perp)$. We will consider only representations on which $1 \in G_\sigma^\perp$ acts as the identity operator. The irreducible ones are classified by the finite abelian group $Q_\sigma/(1-\sigma)(Q \cap \mathfrak{h}_0^\perp)$. All of them have the same dimension $d(\sigma)$, which satisfies

$$(4.53) \quad d(\sigma)^2 = |G_\sigma^\perp/Z(G_\sigma^\perp)| = |(Q \cap \mathfrak{h}_0^\perp)/Q_\sigma|.$$

Definition 4.2. The non-negative integer $d(\sigma)$ is called the *defect* of σ (cf. [14]).

4.5. Representations of $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$. In this subsection we show that the category of all restricted $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ -modules of level 1 is semisimple, and we classify the irreducible ones.

Let $\hat{\mathfrak{h}}_\sigma^-$ (resp. $\hat{\mathfrak{h}}_\sigma^+$) be the subalgebra of $\hat{\mathfrak{h}}_\sigma$ consisting of all elements ht^m with $m > 0$ (resp. $m < 0$). It is well known (see, e.g., [12]) that any restricted $\hat{\mathfrak{h}}_\sigma$ -module M of level 1 is induced from its vacuum subspace

$$(4.54) \quad \Omega_M = \{v \in M \mid \hat{\mathfrak{h}}_\sigma^- v = 0\}.$$

More precisely,

$$(4.55) \quad M \simeq \text{Ind}_{\mathfrak{h}_0 \oplus \hat{\mathfrak{h}}_\sigma^-}^{\hat{\mathfrak{h}}_\sigma} \Omega_M \simeq S(\hat{\mathfrak{h}}_\sigma^+) \otimes \Omega_M.$$

The subalgebra \mathfrak{h}_0 acts on Ω_M diagonally, and the $\hat{\mathfrak{h}}_\sigma$ -module M is completely reducible (it is irreducible iff $\dim \Omega_M = 1$).

Now assume that M is a restricted $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ -module of level 1. It follows from (4.49) that Ω_M is a G_σ -module. By definition, $1 \in G_\sigma$ acts as 1 and the torus T_σ acts diagonally (cf. (4.48)). We will call such G_σ -modules *restricted*.

If Ω is a restricted G_σ -module, it has a compatible \mathfrak{h}_0 -action, because \mathfrak{h}_0 is the Lie algebra of the torus T_σ . We let $\hat{\mathfrak{h}}_\sigma^-$ act trivially on Ω and form the induced $\hat{\mathfrak{h}}_\sigma$ -module $M(\Omega) = \text{Ind}_{\mathfrak{h}_0 \oplus \hat{\mathfrak{h}}_\sigma^-}^{\hat{\mathfrak{h}}_\sigma} \Omega$. Using (4.49), we can extend the action of G_σ from Ω to $M(\Omega)$. Then $M(\Omega)$ becomes a restricted $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ -module of level 1.

Proposition 4.3. *The functors $M \mapsto \Omega_M$ and $\Omega \mapsto M(\Omega)$ establish an equivalence of abelian categories between the category of restricted $(\hat{\mathfrak{h}}_\sigma, G_\sigma)$ -modules of level 1 and the category of restricted G_σ -modules.*

Therefore we are left with describing restricted G_σ -modules.

Proposition 4.4. *Any restricted G_σ -module is completely reducible and is determined by the action of the center of G_σ . Isomorphism classes of restricted irreducible G_σ -modules are parameterized by the (finite) set Z_σ .*

Let Ω be a restricted G_σ -module. For $\mu \in \mathfrak{h}_0$, we denote by Ω_μ the weight μ subspace of Ω :

$$(4.56) \quad \Omega_\mu := \{v \in \Omega \mid e^h v = e^{(h|\mu)} v \text{ for } h \in \mathfrak{h}_0\}.$$

Then $(\mu|\alpha) \in \mathbb{Z}$ for $\alpha \in Q \cap \mathfrak{h}_0$, i.e., $\mu \in (Q \cap \mathfrak{h}_0)^* = \pi_0(Q^*)$ by Lemma 4.6.

Lemma 4.7. (i) $U_\alpha \Omega_\mu = \Omega_{\mu+\pi_0\alpha}$ for all $\alpha \in Q$, $\mu \in \pi_0(Q^*)$. In particular, the subgroup $G_\sigma^\perp \subset G_\sigma$ preserves each Ω_μ .

(ii) Let $\Omega_{\mu+\pi_0(Q)} = \sum_{\alpha \in Q} \Omega_{\mu+\pi_0\alpha} \subset \Omega$. Then $\Omega_{\mu+\pi_0(Q)}$ is a G_σ -submodule of Ω .

(iii) The G_σ -module $\Omega_{\mu+\pi_0(Q)}$ is irreducible if and only if the G_σ^\perp -module Ω_μ is irreducible.

Proof. (i) It follows from (4.41) that $U_\alpha \Omega_\mu \subset \Omega_{\mu+\pi_0\alpha}$ for any $\alpha \in Q$. Since by (4.42) U_α^{-1} is proportional to $U_{-\alpha}$, we get $U_\alpha \Omega_\mu = \Omega_{\mu+\pi_0\alpha}$.

(ii) follows from (i) and the definition of G_σ (see (4.40)–(4.42)).

(iii) First, let Ω_μ be an irreducible G_σ^\perp -module. Assume that Λ is a nontrivial G_σ -submodule of $\Omega_{\mu+\pi_0(Q)}$. Using the action of T_σ , we can write $\Lambda = \sum_{\alpha \in Q} \Lambda_{\mu+\pi_0\alpha}$ where each $\Lambda_{\mu+\pi_0\alpha} \subset \Omega_{\mu+\pi_0\alpha}$. Moreover, $\Lambda_{\mu+\pi_0\alpha} = U_\alpha \Lambda_\mu$. In particular, Λ_μ is a G_σ^\perp -submodule of Ω_μ . But Ω_μ is irreducible; hence, $\Lambda_\mu = \Omega_\mu$ and $\Lambda = \Omega_{\mu+\pi_0(Q)}$.

Conversely, assume that the G_σ^\perp -module Ω_μ is not irreducible. Since G_σ^\perp is a central extension of a finite abelian group, its representations are completely reducible. If $\Omega_\mu = \bigoplus_i L_i$ as a G_σ^\perp -module, let $L^i = \sum_{\alpha \in Q} U_\alpha L_i$. Then each L^i is a G_σ -submodule of $\Omega_{\mu+\pi_0(Q)}$ and $\Omega_{\mu+\pi_0(Q)} = \bigoplus_i L^i$ as a G_σ -module. \square

Remark 4.3. $\Omega_{\mu+\pi_0(Q)}$ is isomorphic to the induced module $\text{Ind}_{T_\sigma \times G_\sigma^\perp}^{G_\sigma} \Omega_\mu$.

From Lemma 4.7(iii) and its proof, we see that the G_σ -module $\Omega_{\mu+\pi_0(Q)}$ is completely reducible. Since $\Omega = \bigoplus_{[\mu] \in \pi_0(Q^*)/\pi_0(Q)} \Omega_{[\mu]}$, it is also completely reducible.

Now let Ω be an irreducible G_σ -module. Then $\Omega = \Omega_{\mu+\pi_0(Q)}$ for some $\mu \in \pi_0(Q^*)$ and the G_σ^\perp -module Ω_μ is irreducible. Any irreducible G_σ^\perp -module is completely determined by the action of the center $Z(G_\sigma^\perp)$. Let $\zeta: Z(G_\sigma^\perp) \rightarrow \mathbb{C}^\times$ be the central character of Ω_μ .

We can view Ω as a G -module on which N_σ acts trivially, and similarly Ω_μ as a G^\perp -module with a trivial action of N_σ^\perp . Recall that, by Lemma 4.5(iii), $Z(G_\sigma^\perp) \simeq Z(G^\perp)/N_\sigma^\perp$, so we can extend ζ to a character of $Z(G^\perp)$.

If $\mu' = \mu + \pi_0\alpha$ for some $\alpha \in Q$, then $\Omega_{\mu'} = U_\alpha \Omega_\mu$ and $\Omega_{\mu+\pi_0(Q)} = \Omega_{\mu'+\pi_0(Q)}$. For $v \in \Omega_\mu$, $U_\beta \in Z(G^\perp)$, $\beta \in Q_\sigma$ (see Lemma 4.5(ii)), we have: $U_\beta U_\alpha v = C_{\alpha,\beta}^{-1} U_\alpha U_\beta v = C_{\alpha,\beta}^{-1} \zeta(U_\beta) U_\alpha v$ where $C_{\alpha,\beta}$ is given by (4.43). Hence, two pairs (μ, ζ) and (μ', ζ') correspond to the same irreducible G_σ -module if and only if they are related by:

$$(4.57) \quad \mu' = \mu + \pi_0\alpha, \quad \zeta'(U_\beta) = C_{\alpha,\beta}^{-1} \zeta(U_\beta), \quad \alpha \in Q, \beta \in Q_\sigma.$$

For $\lambda \in P_\sigma$ the element $e^{2\pi i \lambda_0} U_{(1-\sigma)\lambda} \in Z(G)$ acts on Ω_μ as the scalar $e^{2\pi i (\lambda_0 | \mu)} \zeta(U_{(1-\sigma)\lambda})$ (cf. Lemmas 4.4 and 4.5(ii)). Using Corollary 4.2 and Lemma 4.6, it is easy to see that the action of $Z(G)$ on Ω determines uniquely the equivalence class of (μ, ζ) under (4.57), and hence it determines the isomorphism class of the G_σ -module Ω . Conversely, different pairs (μ, ζ) give rise to different actions of $Z(G)$ on the corresponding modules $\Omega_{\mu+\pi_0(Q)}$.

This completes the proof of Proposition 4.4.

4.6. Classification of σ -Twisted V_Q -Modules. Combining Propositions 4.2, 4.3 and 4.4, we obtain the main result of the paper.

Theorem 4.2. *The category of σ -twisted V_Q -modules is a semisimple abelian category with finitely many isomorphism classes of simple objects, parameterized by the set Z_σ .*

Remark 4.4. The irreducible σ -twisted V_Q -module corresponding to $\lambda + Q \in Z_\sigma$ is isomorphic as an $\hat{\mathfrak{h}}_\sigma$ -module to $S(\hat{\mathfrak{h}}_\sigma^+) \otimes e^{\lambda_0} \mathbb{C}[\pi_0 Q] \otimes \mathbb{C}^{d(\sigma)}$, where \mathbb{C} carries the zero action and $d(\sigma)$ is the defect of σ .

REFERENCES

- [1] B. Bakalov, V. G. Kac, and I. T. Todorov, *Orbifolds of lattice vertex algebras*, in preparation.
- [2] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, Nuclear Phys. **B 241** (1984), no. 2, 333–380.
- [3] R. E. Borcherds, *Vertex algebras, Kac–Moody algebras, and the Monster*, Proc. Nat. Acad. Sci. U.S.A. **83** (1986), no. 10, 3068–3071.
- [4] A. Cappelli, L. S. Georgiev, and I. T. Todorov, *A unified conformal field theory description of paired quantum Hall states*, Comm. Math. Phys. **205** (1999), 657–689.
- [5] R. Dijkgraaf, C. Vafa, E. Verlinde, and H. Verlinde, *The operator algebra of orbifold models*, Comm. Math. Phys. **123** (1989), 485–526.
- [6] L. Dixon, J. A. Harvey, C. Vafa, and E. Witten, *String on orbifolds*, Nucl. Phys. **B261** (1985), 620–678; *String on orbifolds. II*, Nucl. Phys. **B274** (1986), 285–314.
- [7] C. Dong, *Twisted modules for vertex algebras associated with even lattices*, J. Algebra **165** (1994), 91–112.
- [8] C. Dong and J. Lepowsky, *Generalized vertex algebras and relative vertex operators*, Progress in Math., 112. Birkhäuser Boston, 1993.
- [9] I. B. Frenkel and V. G. Kac, *Basic representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980/81), no. 1, 23–66.
- [10] I. B. Frenkel, J. Lepowsky and A. Meurman, *Vertex operator algebras and the Monster*, Pure and Appl. Math., vol. 134, Academic Press, Boston, 1988.
- [11] P. Goddard, *Meromorphic conformal field theory*, in “Infinite-dimensional Lie algebras and groups” (Luminy-Marseille, 1988), 556–587, Adv. Ser. Math. Phys., vol. 7, World Sci. Publishing, Teaneck, NJ, 1989.
- [12] V. G. Kac, *Infinite-dimensional Lie algebras*, 3rd edition, Cambridge University Press, Cambridge, 1990.
- [13] V. G. Kac, *Vertex algebras for beginners*, AMS University Lecture Series, vol. 10, 1996. 2nd edition, 1998.
- [14] V. G. Kac and D. H. Peterson, *112 constructions of the basic representation of the loop group of E_8* , Symposium on anomalies, geometry, topology (Chicago, Ill., 1985), 276–298, World Sci. Publishing, Singapore, 1985.
- [15] V. G. Kac and I. T. Todorov, *Affine orbifolds and rational conformal field theory extensions of $W_{1+\infty}$* , Comm. Math. Phys. **190** (1997), 57–111.
- [16] M. Roitman, *On twisted representations of vertex algebras*, Adv. Math. **176** (2003), 53–88.
- [17] X. Xu, *Twisted modules of coloured lattice vertex operators superalgebras*, Quart. J. Math. Oxford Ser. (2) **47** (1996), 233–259.

DEPARTMENT OF MATHEMATICS, NCSU, RALEIGH, NC 27695, USA
E-mail address: bojko_bakalov@ncsu.edu

DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139, USA
E-mail address: kac@math.mit.edu