

# BGG CORRESPONDENCE AND RÖMER'S THEOREM ON AN EXTERIOR ALGEBRA

KOHI YANAGAWA

*To the memory of Professor Tetsushi Ogoma*

ABSTRACT. Let  $E = K\langle y_1, \dots, y_n \rangle$  be the exterior algebra. The (cohomological) distinguished pairs of a graded  $E$ -module  $N$  describe the growth of a minimal graded injective resolution of  $N$ . Römer gave a duality theorem between the distinguished pairs of  $N$  and those of its dual  $N^*$ . In this paper, we show that under Bernstein-Gel'fand-Gel'fand correspondence, his theorem is translated into a natural corollary of local duality for (complexes of) graded  $S = K[x_1, \dots, x_n]$ -modules. Using this idea, we also give a  $\mathbb{Z}^n$ -graded version of Römer's theorem.

## INTRODUCTION

In this section, to introduce a background of the present paper, we summarize results of Aramova-Herzog [2] and Römer [11].

Let  $S = K[x_1, \dots, x_n]$  be the polynomial ring over a field  $K$ , and  $M$  a finitely generated graded  $S$ -module. The  $ij$ th Betti number  $\beta_{i,j}(M) = \dim_K \operatorname{Tor}_i^S(K, M)_j$  of  $M$  is an important invariant. Following Bayer-Charalambous-Popescu [3], we say a Betti number  $\beta_{k,m}(M) \neq 0$  is *extremal*, if  $\beta_{i,j}(M) = 0$  for all  $(i, j) \neq (k, m)$  with  $i \geq k$  and  $j - i \geq m - k$ . This notion has two remarkable properties. First, a homogeneous ideal  $I \subset S$  has the same extremal Betti numbers as its generic initial ideal  $\operatorname{Gin}(I)$  with respect to the reverse lexicographic term order on  $S$ . Another important property is the following.

**Theorem A** (Bayer-Charalambous-Popescu, [3, Theorem 2.8]) *Let  $\Delta \subset 2^{\{1, \dots, n\}}$  be a simplicial complex, and  $K[\Delta] = S/I_\Delta$  the Stanley-Reisner ring. And let  $\Delta^\vee$  be the Alexander dual complex of  $\Delta$ . Then  $\beta_{i,i+j}(K[\Delta])$  is extremal if and only if so is  $\beta_{j,i+j}(I_{\Delta^\vee})$ . Moreover, if this is the case, then  $\beta_{i,i+j}(K[\Delta]) = \beta_{j,i+j}(I_{\Delta^\vee})$ .*

We have  $\beta_{i,n}(K[\Delta]) = \dim_K \tilde{H}_{n-i-1}(|\Delta|; K) =: \tilde{h}_{n-i-1}(|\Delta|)$  by Hochster's formula. If  $\beta_{i,n}(K[\Delta]) \neq 0$  then it is always an extremal Betti number. The equality  $\tilde{h}_{n-i-1}(|\Delta|) = \beta_{i,n}(K[\Delta]) = \beta_{n-i,n}(I_{\Delta^\vee}) = \beta_{n-i+1,n}(K[\Delta^\vee]) = \tilde{h}_{i-2}(|\Delta^\vee|)$  induced by Theorem A corresponds to the usual Alexander duality. More generally, Theorem A gives an Alexander duality for (some) iterated Betti numbers (c.f. [9, 4]).

Let  $E = K\langle y_1, \dots, y_n \rangle$  be the exterior algebra. To understand Theorem A, Aramova-Herzog [2] introduced distinguished pairs for a graded  $E$ -module  $N$ . See Definition 1.6 below. (We use a different convention to describe these pairs. See Remark 1.7.) The distinguished pairs of  $N$  roughly describe the growth of the minimal graded (infinite) injective resolution of  $N$ . Let  $K\{\Delta\} = E/J_\Delta$  be the

exterior face ring of  $\Delta$ . Then [2, Corollary 9.6] states that  $(d, i)$  is a distinguished pair for  $K\{\Delta\}^* := \text{Hom}_E(K\{\Delta\}, E)$  if and only if  $\beta_{d+i-n, d}(K[\Delta])$  is extremal. And the values of extremal Betti numbers can be described by the Cartan cohomologies of  $K\{\Delta\}^*$ .

Römer proved that  $(d, i)$  is distinguished for  $N$  if and only if so is  $(d, 2n - d - i)$  for  $N^*$ . He also gave a duality between certain components of Cartan cohomologies of  $N$  and those of  $N^*$ . Since  $k\{\Delta\}^* = J_{\Delta^\vee}$ , his result implies Theorem A.

Bernstein-Gel'fand-Gel'fand correspondence (BGG correspondence, for short) is a well known theorem which states that the derived category  $D^b(\text{gr } S)$  of finitely generated graded  $S$ -modules is equivalent to the similar category  $D^b(\text{gr } E)$  for  $E$ . In this paper, we give a new proof of the result of Römer using BGG correspondence. More precisely, under this correspondence, Römer's theorem is translated into a statement on  $D^b(\text{gr } S)$  which is a natural consequence of the local duality (Serre duality). A key point is that the duality functor  $\text{Hom}_E(-, E)$  on  $D^b(\text{gr } E)$  corresponds to the duality functor  $R\text{Hom}_S(-, \omega^\bullet)$  on  $D^b(\text{gr } S)$ , where  $\omega^\bullet$  is a dualizing complex of  $S$ . So our proof is (philosophically) simple and sheds new light on the result.

The original paper [3] states Theorem A in the  $\mathbb{Z}^n$ -graded context, while the arguments in [2, 11] are hard to work in this context. But, since BGG correspondence also holds for  $\mathbb{Z}^n$ -graded modules, our method is powerful in this context too. See §2. This part of the present paper is a continuation of the author's previous paper [13].

## 1. $\mathbb{Z}$ -GRADED CASE

Let  $W$  be an  $n$ -dimensional vector space over a field  $K$ , and  $S = \bigoplus_{i \geq 0} \text{Sym}_i W$  the polynomial ring. We regard  $S$  as a graded ring with  $S_i = \text{Sym}_i W$ . Let  $\text{Gr } S$  be the category of graded  $S$ -modules and their degree preserving  $S$ -homomorphisms, and  $\text{gr } S$  the full subcategory of  $\text{Gr } S$  consisting of finitely generated modules. Then there is an equivalence  $D^b(\text{gr } S) \cong D_{\text{gr } S}^b(\text{Gr } S)$ . (For derived categories, consult [8].) So we will freely identify these categories. For  $M = \bigoplus_{i \in \mathbb{Z}} M_i \in \text{Gr } S$  and an integer  $j$ ,  $M(j)$  denotes the shifted module with  $M(j)_i = M_{i+j}$ . For  $M^\bullet \in D^b(\text{Gr } S)$ ,  $M^\bullet[j]$  denotes the  $j$ th translation of  $M^\bullet$ , that is,  $M^\bullet[j]$  is the complex with  $M^\bullet[j]^i = M^{i+j}$ . So, if  $M \in \text{Gr } S$ ,  $M[j]$  is the cochain complex  $\cdots \rightarrow 0 \rightarrow M \rightarrow 0 \rightarrow \cdots$ , where  $M$  sits in the  $(-j)$ th position. If  $M \in \text{gr } S$  and  $N \in \text{Gr } S$ , then  $\text{Hom}_S(M, N)$  has the structure of a graded  $S$ -module with  $\text{Hom}_S(M, N)_i = \text{Hom}_{\text{Gr } S}(M, N(i))$ .

Let  $\omega^\bullet \in D^b(\text{gr } S)$  be a minimal graded injective resolution of  $S(-n)[n]$ . That is,  $\omega^\bullet$  is a graded normalized dualizing complex of  $S$ . Then  $\mathbf{D}_S(-) := \text{Hom}_S^\bullet(-, \omega^\bullet)$  gives a duality functor from  $D^b(\text{gr } S)$  to itself. The  $i$ th cohomology of  $\mathbf{D}_S(M^\bullet)$  is  $\text{Ext}_S^i(M^\bullet, \omega^\bullet)$ . For  $M^\bullet \in D^b(\text{gr } S)$  and  $i \in \mathbb{Z}$ , set  $d_i(M^\bullet) := \dim_S H^i(M^\bullet)$ . Here the Krull dimension of the 0 module is  $-\infty$ .

**Definition 1.1.** We say  $(d, i) \in \mathbb{N} \times \mathbb{Z}$  is a *distinguished pair* for a complex  $M^\bullet \in D^b(\text{gr } S)$ , if  $d = d_i(M^\bullet)$  and  $d_j(M^\bullet) < d + i - j$  for all  $j$  with  $j < i$ .

Let  $M^\bullet \in D^b(\text{gr } S)$  and  $d = d_i(M^\bullet) \geq 0$ . If  $d = \max\{d_j(M^\bullet) \mid j \in \mathbb{Z}\}$ , then  $(d, i)$  is distinguished for  $M^\bullet$ . On the other hand, if  $i = \min\{j \mid H^j(M^\bullet) \neq 0\}$ , then  $(d, i)$  is also distinguished. Thus  $M^\bullet$  has several distinguished pairs in general.

In this paper,  $\deg_S(M)$  denotes the multiplicity of a module  $M \in \text{gr } S$  (i.e.,  $e(M)$  of [5, Definition 4.1.5]).

**Theorem 1.2.** *For  $M^\bullet \in D^b(\text{gr } S)$ , we have the following.*

(1) *A pair  $(d, i)$  is distinguished for  $M^\bullet$  if and only if  $(d, -d - i)$  is distinguished for  $\mathbf{D}_S(M^\bullet)$ .*

(2) *If  $(d, i)$  is a distinguished pair for  $M^\bullet$ , then*

$$\deg_S H^i(M^\bullet) = \deg_S \text{Ext}_S^{-d-i}(M^\bullet, \omega^\bullet).$$

*Proof.* (1) Since the statement is ‘‘symmetric’’, it suffices to prove the direction  $\Rightarrow$ .

From the double complex  $\text{Hom}_S^\bullet(M^\bullet, \omega^\bullet)$ , we have a spectral sequence  $E_2^{p,q} = \text{Ext}_S^p(H^{-q}(M^\bullet), \omega^\bullet) \Rightarrow \text{Ext}_S^{p+q}(M^\bullet, \omega^\bullet)$ . For simplicity, set  $e_r^{p,q} := \dim_S E_r^{p,q}$ . Since  $\text{Ext}_S^i(M, \omega^\bullet) \cong \text{Ext}_S^{n+i}(M, S(-n))$  for  $M \in \text{gr } S$ , the following inequality follows from argument analogous to [5, §8.1, Theorem 8.1.1].

$$(1.1) \quad e_2^{p,q} = \dim_S \text{Ext}_S^p(H^{-q}(M^\bullet), \omega^\bullet) = \begin{cases} -p & \text{if } p = -d_{-q}(M^\bullet), \\ \leq -p & \text{if } -d_{-q}(M^\bullet) < p \leq 0, \\ -\infty & \text{otherwise.} \end{cases}$$

(I) By (1.1), we have  $e_2^{-d,-i} = d$ . On the other hand, we have  $e_2^{p,q} < d$  for all  $(p, q) \neq (-d, -i)$  with  $p + q = -d - i$ . In fact, the assertion follows from (1.1) if  $p > -d$ . So we may assume that  $p < -d$  and  $q = -d - i - p > -i$ . Since  $(d, i)$  is distinguished, we have  $d_{-q}(M^\bullet) < d + i + q = -p$ . Thus  $E_2^{p,q} = 0$  in this case. Anyway, we have  $e_\infty^{p,q} < d$  for all  $(p, q) \neq (-d, -i)$  with  $p + q = -d - i$ .

(II) Since  $d_{i-j+1}(M^\bullet) < d + j - 1 < d + j$  for all  $j \geq 2$ , we have that  $E_2^{-d-j, -i+j-1} = 0$ . So we have  $E_r^{-d-j, -i+j-1} = 0$  for all  $r \geq 2$ . Next we will show that  $d = e_2^{-d,-i} = e_3^{-d,-i} = \dots = e_r^{-d,-i}$  by induction on  $r$ . Recall that  $E_{r+1}^{-d,-i}$  is the cohomology of

$$E_r^{-d-r, -i+r-1} \rightarrow E_r^{-d,-i} \rightarrow E_r^{-d+r, -i-r+1}.$$

But we have seen that  $E_r^{-d-r, -i+r-1} = 0$ . Moreover,  $e_r^{-d+r, -i-r+1} \leq e_2^{-d+r, -i-r+1} \leq d - r < d$  by (1.1), and  $e_r^{-d,-i} = d$  by the induction hypothesis. Thus  $e_{r+1}^{-d,-i} = d$ . Hence  $e_\infty^{-d,-i} = d$ . From this fact and (I), we have that  $\dim_S \text{Ext}_S^{-d-i}(M^\bullet, \omega^\bullet) = d$ .

(III) Finally, we will show that  $\dim_S \text{Ext}_S^{-d-i-j}(M^\bullet, \omega^\bullet) < d + j$  for all  $j > 0$ . To see this, it suffices to show that  $e_2^{p,q} < d + j$  for all  $j > 0$  and all  $(p, q)$  with  $p + q = -d - i - j$ . If  $p > -d - j$ , the assertion is clear. If  $p = -d - j$ , then  $q = -i$  and  $d_{-q}(M^\bullet) = d < -p$ . So  $E_2^{p,q} = 0$  in this case. Hence we may assume that  $p < -d - j$  and  $-q = d + i + j + p < i$ . Since  $(d, i)$  is distinguished,  $d_{-q}(M^\bullet) < d + (i + q) = -j - p < -p$ . So we have  $E_2^{p,q} = 0$  in this case too.

(2) Since  $\deg_S E_r^{-d,-i} = \deg_S E_{r+1}^{-d,-i}$  for all  $r \geq 2$  by the argument in (II) of the proof of (1), we have  $\deg_S E_2^{-d,-i} = \deg_S E_\infty^{-d,-i}$ . So we have

$$\deg_S \text{Ext}_S^{-d-i}(M^\bullet, \omega^\bullet) = \deg_S E_\infty^{-d,-i} = \deg_S E_2^{-d,-i} = \deg_S \text{Ext}_S^{-d}(H^i(M^\bullet), \omega^\bullet),$$

where the first equality follows from (I) and (II).

For a module  $M \in \text{gr } S$  of dimension  $d$ , we have  $\deg_S M = \deg_S \text{Ext}_S^{-d}(M, \omega^\bullet)$ . In fact, for a prime ideal  $\mathfrak{p} \subset S$  with  $\dim S/\mathfrak{p} = d$ , let  $\mathbf{x}$  be a maximal  $S_{\mathfrak{p}}$ -sequence contained in  $\text{Ann}_{S_{\mathfrak{p}}}(M_{\mathfrak{p}})$ , and  $R := S_{\mathfrak{p}}/\mathbf{x}S_{\mathfrak{p}}$  an artinian Gorenstein local ring. Then we have

$$\text{Ext}_S^{-d}(M, \omega^\bullet) \otimes_S S_{\mathfrak{p}} \cong \text{Ext}_{S_{\mathfrak{p}}}^{n-d}(M_{\mathfrak{p}}, S_{\mathfrak{p}}) \cong \text{Hom}_R(M_{\mathfrak{p}}, R).$$

Therefore,

$$l_{S_{\mathfrak{p}}}(\text{Ext}_S^{-d}(M, \omega^\bullet) \otimes_S S_{\mathfrak{p}}) = l_R(\text{Hom}_R(M_{\mathfrak{p}}, R)) = l_R(M_{\mathfrak{p}}) = l_{S_{\mathfrak{p}}}(M_{\mathfrak{p}}).$$

Since  $\dim_S(H^i(M^\bullet)) = d$ , we have  $\deg_S \text{Ext}_S^{-d}(H^i(M), \omega^\bullet) = \deg_S H^i(M^\bullet)$ .  $\square$

*Remark 1.3.* Theorem 1.2 (1) also holds for a noetherian local ring  $A$  admitting a dualizing complex. The part (2) also holds for  $A$ , if we replace  $\deg_S(-)$  by  $l_{A_{\mathfrak{p}}}(- \otimes_A A_{\mathfrak{p}})$  for a prime ideal  $\mathfrak{p} \subset A$  with  $\dim A/\mathfrak{p} = d$ .

Let  $V$  be the dual vector space of  $W$ , and  $E = \bigwedge V$  the exterior algebra. We regard  $E$  as a negatively graded ring with  $E_{-i} = \bigwedge^i V$  (this is the opposite convention from [2, 11]). Let  $\text{gr } E$  be the category of finitely generated graded  $E$ -modules and their degree preserving  $E$ -homomorphisms. Here “ $E$ -module” means a left and right module  $N$  with  $ea = (-1)^{(\deg e)(\deg a)}ae$  for all homogeneous elements  $e \in E$  and  $a \in N$ .

Let  $\{x_1, \dots, x_n\}$  be a basis of  $W$ , and  $\{y_1, \dots, y_n\}$  its dual basis of  $V$ . For a complex  $N^\bullet$  in  $\text{gr } E$ , set  $\mathbf{L}(N^\bullet) = \bigoplus_{i \in \mathbb{Z}} S \otimes_K N^i$  and  $\mathbf{L}(N^\bullet)^m = \bigoplus_{i-j=m} S \otimes_K N_j^i$ . The differential defined by

$$\mathbf{L}(N^\bullet)^m \supset S \otimes_K N_j^i \ni 1 \otimes z \mapsto \sum_{1 \leq l \leq n} x_l \otimes y_l z + (-1)^m (1 \otimes \delta^i(z)) \in \mathbf{L}(N^\bullet)^{m+1}$$

makes  $\mathbf{L}(N^\bullet)$  a cochain complex of free  $S$ -modules. Here  $\delta^i$  is the  $i$ th differential map of  $N^\bullet$ . Moreover,  $\mathbf{L}$  gives a functor from  $D^b(\text{gr } E)$  to  $D^b(\text{gr } S)$ .

For  $M \in \text{gr } S$  and  $i \in \mathbb{Z}$ , we can define a graded  $E$ -module structure on  $\text{Hom}_K(E, M_i)$  by  $(af)(e) = f(ea)$ . Then  $\text{Hom}_K(E, M_i) \cong E(-n) \otimes_K M_i$ . Set  $\mathbf{R}(M) = \text{Hom}_K(E, M)$  and  $\mathbf{R}^i(M) = \text{Hom}_K(E, M_i)$ . The differential defined by

$$\mathbf{R}^i(M) = \text{Hom}_K(E, M_i) \ni f \mapsto [e \mapsto \sum_{1 \leq j \leq n} x_j f(y_j e)] \in \text{Hom}_K(E, M_{i+1}) = \mathbf{R}^{i+1}(M)$$

makes  $\mathbf{R}(M)$  a cochain complex of free  $E$ -modules. We can also construct  $\mathbf{R}(M^\bullet)$  from a complex  $M^\bullet$  in natural way. Then  $\mathbf{R}$  gives a functor from  $D^b(\text{gr } S)$  to  $D^b(\text{gr } E)$ . See [6] for details. The following is a crucial result.

**Theorem 1.4** (BGG correspondence, c.f.[6]). *The functors  $\mathbf{L}$  and  $\mathbf{R}$  give a category equivalence  $D^b(\text{gr } S) \cong D^b(\text{gr } E)$ .*

For  $N \in \text{gr } E$ , then  $N^* := \text{Hom}_E(N, E) \cong \text{Hom}_K(N, K)(n)$  is a graded  $E$ -module again.  $(-)^*$  gives an exact duality functor on  $\text{gr } E$ , and it can be extended to the duality functor  $\mathbf{D}_E$  on  $D^b(\text{gr } E)$ .

**Proposition 1.5.** *For  $N^\bullet \in D^b(\text{gr } E)$ , we have*

$$\mathbf{D}_S \circ \mathbf{L}(N^\bullet) \cong \mathbf{L} \circ \mathbf{D}_E(N^\bullet)(-2n)[2n].$$

*Proof.* Since  $\mathbf{L}(N^\bullet)$  consists of free  $S$ -modules, we have

$$\mathbf{D}_S \circ \mathbf{L}(N^\bullet) \cong \text{Hom}_S^\bullet(\mathbf{L}(N^\bullet), S(-n)[n]).$$

It is easy to see that

$$\text{Hom}_S^m(\mathbf{L}(N^\bullet), S(-n)[n]) \cong \bigoplus_{j-i=m+n} S(-n) \otimes_K (N_j^i)^\vee,$$

where  $(-)^\vee$  means the graded  $K$ -dual. On the other hand,

$$\begin{aligned} \mathbf{L} \circ \mathbf{D}_E(N^\bullet)^m &= \bigoplus_{i-j=m} S \otimes_K \mathbf{D}_E(N^\bullet)_j^i = \bigoplus_{i-j=m} S(n) \otimes_K (N_{-n-j}^{-i})^\vee \\ &= \bigoplus_{j-i=m-n} S(n) \otimes_K (N_j^i)^\vee. \end{aligned}$$

So we can easily construct a quasi-isomorphism  $\mathbf{D}_S \circ \mathbf{L}(N^\bullet) \rightarrow \mathbf{L} \circ \mathbf{D}_E(N^\bullet)(-2n)[2n]$ .  $\square$

For  $N^\bullet \in D^b(\text{gr } E)$ , we have  $H^i(\mathbf{L}(N^\bullet))_j \cong \text{Ext}_E^{j+i}(K, N^\bullet)_j$  by [6, Theorem 3.7]. So the Laurent series  $P_i(t) = \sum_{j \in \mathbb{Z}} (\dim_K \text{Ext}_E^{j+i}(K, N^\bullet)_j) \cdot t^j$  is the Hilbert series of the finitely generated graded  $S$ -module  $H^i(\mathbf{L}(N^\bullet))$ . If  $H^i(\mathbf{L}(N^\bullet)) \neq 0$ , there exists a Laurent polynomial  $Q_i(t) \in \mathbb{Z}[t, t^{-1}]$  such that

$$P_i(t) = \frac{Q_i(t)}{(1-t)^d},$$

where  $d = d_i(\mathbf{L}(N^\bullet)) = \dim_S H^i(\mathbf{L}(N^\bullet))$ . Set  $e_i(N^\bullet) := Q_i(1) = \deg_S H^i(\mathbf{L}(N^\bullet))$ . So  $d_i(\mathbf{L}(N^\bullet))$  and  $e_i(N^\bullet)$  measure the growth of the “ $(-i)$ -linear strand” of a minimal injective resolution of  $N^\bullet$ .

[2, 11] treated  $d_i(\mathbf{L}(N))$  and  $e_i(N)$  for a module  $N \in \text{gr } E$  more or less indirectly. But their approach is very different from ours. They use *Cartan (co)homology* of  $N$ . See [2, 11] for the definition of this (co)homology. Let  $\mathbf{v} = v_1, \dots, v_n$  be a basis of  $V$  which is *generic* with respect to  $N$  in the sense of [2, Definition 4.7]. As [2, 11], we set  $H_i(k) := H_i(v_1, \dots, v_k; N)$  and  $H^i(k) := H^i(v_1, \dots, v_k; N)$  to be Cartan (co)homologies. Note that  $H_i(n) = \text{Tor}_i^E(K, N)$ ,  $H^i(n) = \text{Ext}_E^i(K, N)$  and  $H^i(v_1, \dots, v_k; N^*) \cong H_i(v_1, \dots, v_k; N)^*$ . It follows from the argument in §6 of [2] that the function  $j \mapsto \dim_K H^{i+j}(k)_j$  is a polynomial function for  $j \gg 0$ . Moreover,  $d_i(\mathbf{L}(N)) \leq 0$  if and only if  $H^{i+j}(n)_j = 0$  for  $j \gg 0$  if and only if  $H^{i+j}(k)_j = 0$  for all  $k \leq n$  and  $j \gg 0$ . [2, Proposition 9.4] can be restated as follows: If  $d_i(\mathbf{L}(N)) > 0$ , we have

$$(1.2) \quad d_i(\mathbf{L}(N)) = n + 1 - \min\{k \mid H^{i+j}(k)_j \neq 0 \text{ for all } j \gg 0\}.$$

A (*cohomological*) *distinguished pair* for a module  $N \in \text{gr } E$  was introduced in [11, Definition 3.4] (see also [2]). Here we generalize this notion to a complex.

**Definition 1.6.** Let  $N^\bullet \in D^b(\text{gr } E)$ . We say  $(d, i) \in \mathbb{N} \times \mathbb{Z}$  is a *distinguished pair* for  $N^\bullet$  if and only if it is distinguished for  $\mathbf{L}(N^\bullet)$  (in the sense of Definition 1.1).

*Remark 1.7.* By (1.2), we see that  $(d, i)$  is a distinguished pair for a module  $N \in \text{gr } E$  in the above sense if and only if  $(n+1-d, i)$  is a ‘‘cohomological distinguished pair’’ for  $N$  in the sense of [11]. (Recall that  $E$  is a positively graded ring in [2, 11].) [2] also use the term ‘‘distinguished pair’’. But this is ‘‘homological distinguished pair’’ of [11], and  $(d, i)$  is a distinguished pair for  $N$  in our sense if and only if  $(n+1-d, n-i)$  is a distinguished pair for  $N^*$  in the sense of [2].

**Corollary 1.8** (c.f. [11, Theorem 3.8]). *Let  $N^\bullet \in D^b(\text{gr } E)$ . A pair  $(d, i)$  is distinguished for  $N^\bullet$  if and only if  $(d, 2n-d-i)$  is distinguished for  $\mathbf{D}_E(N^\bullet)$ . If this is the case, we have  $e_i(N^\bullet) = e_{2n-d-i}(\mathbf{D}_E(N^\bullet))$ .*

*Proof.* For the first statement, it suffices to prove the direction  $\Rightarrow$ . By Theorem 1.2,  $(d, -d-i)$  is a distinguished pair for  $\mathbf{D}_S \circ \mathbf{L}(N^\bullet) \cong \mathbf{L} \circ \mathbf{D}_E(N^\bullet)(-2n)[2n]$ . For a complex  $M^\bullet \in D^b(\text{gr } S)$ , we have  $H^j(M^\bullet(-2n)[2n]) = H^{2n+j}(M^\bullet)(-2n)$  and  $d_j(M^\bullet(-2n)[2n]) = d_{2n+j}(M^\bullet)$ . Thus  $(d, 2n-d-i)$  is distinguished for  $\mathbf{L} \circ \mathbf{D}_E(N^\bullet)$ . The last equality follows from Theorem 1.2 (2).  $\square$

For a module  $N \in \text{gr } E$ ,  $d_i(\mathbf{L}(N))$  can be 0 quite often. But we have the following.

**Proposition 1.9.** *Assume that a module  $N \in \text{gr } E$  does not have a free summand. If  $(d, i)$  is a distinguished pair for  $N$ , then we have  $d > 0$ .*

*Proof.* Let  $0 \rightarrow N \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$  (resp.  $\dots \rightarrow I^{-1} \rightarrow I^0 \rightarrow N \rightarrow 0$ ) be a minimal injective (resp. projective) resolution of  $N$ . For  $j \geq 0$ , set  $\Omega_j(N) := (\ker(I^j \rightarrow I^{j+1}))[-j]$ . Obviously,  $0 \rightarrow \Omega_j(N) \rightarrow I^j \rightarrow I^{j+1} \rightarrow \dots$  is a minimal injective resolution. On the other hand, since  $N$  does not have a free summand,  $\dots \rightarrow I^{-1} \rightarrow I^0 \rightarrow \dots \rightarrow I^{j-1} \rightarrow \Omega_j(N) \rightarrow 0$  is a minimal projective resolution. If  $d_i(\mathbf{L}(N)) > 0$ , then  $d_i(\mathbf{L}(\Omega_j(N))) = d_i(\mathbf{L}(N))$  for all  $j \geq 0$ . But, if  $d_i(\mathbf{L}(N)) = 0$ , then  $d_i(\mathbf{L}(\Omega_j(N))) = -\infty$  for  $j \gg 0$ . On the other hand, since a minimal injective resolution of  $N^*$  is the dual of a minimal projective resolution of  $N$ , we have  $d_i(\mathbf{L}(N^*)) = d_i(\mathbf{L}(\Omega_j(N)^*))$  for all  $i$  and all  $j \geq 0$ . So  $N^*$  and  $\Omega_j(N)^*$  have the same distinguished pairs. For a contradiction, we assume that  $(0, i)$  is a distinguished pair for  $N$ . Then  $(0, 2n-i)$  is a distinguished pair for  $N^*$  and  $\Omega_j(N)^*$ . So  $(0, i)$  is a distinguished pair for  $\Omega_j(N)$  for all  $j \geq 0$ . This contradicts the above observation.  $\square$

We say a distinguished pair  $(d, i)$  is *positive*, if  $d > 0$ . Since [2, 11] study a distinguished pair for a module, they only treat a positive one.

*Remark 1.10.* When  $N^\bullet$  is a module, Corollary 1.8 was proved in [11, Theorem 3.8]. On the other hand, for positive distinguished pairs, we can prove the corollary from [11, Theorem 3.8] directly: Let  $I^\bullet$  be an injective resolution of  $N^\bullet$  and  $P^\bullet$  a projective resolution of  $I^\bullet$ . From the quasi-isomorphism  $f : P^\bullet \rightarrow I^\bullet$ , we have the exact complex  $(T^\bullet, \partial^\bullet) := \text{cone}(f)$ . Then  $N := \ker \partial_0$  (resp.  $N^*$ ) has the same positive distinguished pairs as  $N^\bullet$  (resp.  $\mathbf{D}_E(N^\bullet)$ ).

A variant of BGG correspondence gives an equivalence  $\underline{\text{gr}}E \cong D^b(\text{Coh}(\mathbb{P}^{n-1}))$  of triangulated categories, where  $\underline{\text{gr}}E$  is the stable category, and  $\text{Coh}(\mathbb{P}^{n-1})$  is the category of coherent sheaves on  $\mathbb{P}^{n-1} = \text{Proj } S$ . More precisely, the composition of the functor  $\mathbf{L} : \text{gr } E \rightarrow D^b(\text{gr } S)$  and the natural functor  $D^b(\text{gr } S) \rightarrow D^b(\text{Coh}(\mathbb{P}^{n-1}))$  induces this equivalence. Note that the functor  $\text{gr } S \ni M \rightarrow \tilde{M} \in \text{Coh}(\mathbb{P}^{n-1})$  ignores modules of finite length. Hence if  $d_i(M^\bullet) = 0$  then  $H^i(\tilde{M}^\bullet) = 0$ . In this sense, the duality in [11] corresponds to a duality on  $D^b(\text{Coh}(\mathbb{P}^{n-1}))$ .

In the rest of this section, we assume that  $K$  is algebraically closed. Let  $N \in \text{gr } E$ . Following [1], we say  $v \in E_{-1} = V$  is  $N$ -regular if  $\text{Ann}_N(v) = vN$ . It is easy to see that  $v$  is  $N$ -regular if and only if it is  $N^*$ -regular. We say  $V_E(N) = \{v \in V \mid v \text{ is not } N\text{-regular}\}$  is the *rank variety* of  $N$  (see [1]). [1, Theorem 3.1] states that  $V_E(N)$  is an algebraic subset of  $V = \text{Spec } S$ , and  $\dim V_E(N) = \max\{d_i(\mathbf{L}(N)) \mid i \in \mathbb{Z}\}$ . By the above remark,  $V_E(N) = V_E(N^*)$ . We can refine this observation using the grading of  $N$ .

Recall that  $S$  can be seen as the Yoneda algebra  $\text{Ext}_E^*(K, K)$ , and  $\text{Ext}_E^*(K, N)$  has the  $S$ -module structure. By the same argument as [1, Theorem 3.9] (see also the proof of [7, Corollary 3.2 (b)]), we have that

$$V_E(N) = \{v \in V \mid \xi(v) = 0 \text{ for all } \xi \in \text{Ann}_S(\text{Ext}_E^*(K, N))\}.$$

But  $[\text{Ext}_E^{*+i}(K, N)]_* := \bigoplus_{j \in \mathbb{Z}} \text{Ext}_E^{j+i}(K, N)_j$  is an  $S$ -module which is isomorphic to  $H^i(\mathbf{L}(N))$  (see the proof of [6, Proposition 2.3]), and we have  $\text{Ext}_E^*(K, N) \cong \bigoplus_{j \in \mathbb{Z}} [\text{Ext}_E^{*+i}(K, N)]_*$ . Set

$$V_E^i(N) = \{v \in V \mid \xi(v) = 0 \text{ for all } \xi \in \text{Ann}_S([\text{Ext}_E^{*+i}(K, N)]_*)\}.$$

We have  $V_E(N) = \bigcup_i V_E^i(N)$  and  $d_i(\mathbf{L}(N)) = \dim V_E^i(N)$ . For an algebraic set  $X \subset \text{Spec } S$  of dimension  $d$ , set  $\text{Top}(X)$  to be the union of the all irreducible components of  $X$  of dimensions  $d$ .

**Proposition 1.11.** *If  $(d, i)$  is a distinguished pair for  $N \in \text{gr } E$ , then we have  $\text{Top}(V_E^i(N)) = \text{Top}(V_E^{2n-d-i}(N^*))$ .*

*Proof.* By the proof of Theorem 1.2,  $\text{Ann}_S(H^i(\mathbf{L}(N)))$  has the same top dimensional components as  $\text{Ann}_S(H^{-d-i}(\mathbf{D}_S \circ \mathbf{L}(N)))$ .  $\square$

In the above situation, we have  $V_E^i(N) \neq V_E^{2n-d-i}(N^*)$  in general.

## 2. SQUAREFREE CASE

In this section, we regard  $S = K[x_1, \dots, x_n]$  as an  $\mathbb{N}^n$ -graded ring with  $\deg x_i = (0, \dots, 0, 1, 0, \dots, 0)$  where 1 is in the  $i$ th position. Similarly,  $E = K\langle y_1, \dots, y_n \rangle$  is a  $-\mathbb{N}^n$ -graded ring with  $\deg y_i = -\deg x_i$ . Let  ${}^*\text{gr } S$  (resp.  ${}^*\text{gr } E$ ) be the category of finitely generated  $\mathbb{Z}^n$ -graded  $S$ -modules (resp.  $E$ -modules). The functors  $\mathbf{L}$  and  $\mathbf{R}$  defining the BGG correspondence  $D^b(\text{gr } S) \cong D^b(\text{gr } E)$  also work in the  $\mathbb{Z}^n$ -graded context. That is, the functors  $\mathbf{L} : D^b({}^*\text{gr } E) \rightarrow D^b({}^*\text{gr } S)$  and  $\mathbf{R} : D^b({}^*\text{gr } S) \rightarrow D^b({}^*\text{gr } E)$  are defined by the same way as the  $\mathbb{Z}$ -graded case, and they give an equivalence  $D^b({}^*\text{gr } S) \cong D^b({}^*\text{gr } E)$ , see [13, Theorem 4.1]. Note that the dualizing

complex  $\omega^\bullet$  of  $S$  is  $\mathbb{Z}^n$ -graded, and  $\mathbf{D}_S(-) = \text{Hom}_S^\bullet(-, \omega^\bullet)$  is also a duality functor on  $D^b(*\text{gr } S)$ . Similarly,  $\mathbf{D}_E(-) = \text{Hom}_E(-, E)$  is a duality functor on  $D^b(*\text{gr } E)$ . As Proposition 1.5, for  $N^\bullet \in D^b(*\text{gr } E)$ , we have  $\mathbf{D}_S \circ \mathbf{L}(N^\bullet) \cong \mathbf{L} \circ \mathbf{D}_E(N^\bullet)(-2)[2n]$  in  $D^b(*\text{gr } S)$ . Here we set  $\mathbf{j} := (j, j, \dots, j) \in \mathbb{N}^n$  for  $j \in \mathbb{Z}$ .

For  $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}^n$ , set  $\text{supp}(\mathbf{a}) := \{i \mid a_i > 0\} \subset [n] := \{1, \dots, n\}$  and  $|\mathbf{a}| = \sum_{i=1}^n a_i$ . We say  $\mathbf{a} \in \mathbb{Z}^n$  is *squarefree* if  $a_i = 0, 1$  for all  $i \in [n]$ . When  $\mathbf{a} \in \mathbb{Z}^n$  is squarefree, we sometimes identify  $\mathbf{a}$  with  $\text{supp}(\mathbf{a})$ .

**Definition 2.1** ([12]). We say a  $\mathbb{Z}^n$ -graded  $S$ -module  $M$  is *squarefree*, if the following conditions are satisfied.

- (a)  $M$  is  $\mathbb{N}^n$ -graded (i.e.,  $M_{\mathbf{a}} = 0$  if  $\mathbf{a} \notin \mathbb{N}^n$ ) and finitely generated.
- (b) The multiplication map  $M_{\mathbf{a}} \ni y \mapsto (\prod x_i^{b_i}) \cdot y \in M_{\mathbf{a}+\mathbf{b}}$  is bijective for all  $\mathbf{a}, \mathbf{b} \in \mathbb{N}^n$  with  $\text{supp}(\mathbf{a} + \mathbf{b}) = \text{supp}(\mathbf{a})$ .

For a simplicial complex  $\Delta \subset 2^{[n]}$ , the Stanley-Reisner ideal  $I_\Delta := (\prod_{i \in F} x_i \mid F \notin \Delta)$  and the Stanley-Reisner ring  $K[\Delta] := S/I_\Delta$  are squarefree modules. Note that if  $M$  is squarefree then  $M_{\mathbf{a}} \cong M_F$  as  $K$ -vector spaces for all  $\mathbf{a} \in \mathbb{N}^n$  with  $\text{supp}(\mathbf{a}) = F$ . Let  $\text{Sq}_S$  be the full subcategory of  $*\text{gr } S$  consisting of squarefree modules. In  $*\text{gr } S$ ,  $\text{Sq}_S$  is closed under kernels, cokernels and extensions ([12, Lemma 2.3]), and we have that  $D^b(\text{Sq}_S) \cong D_{\text{Sq}_S}^b(*\text{gr } S)$ . If  $M^\bullet \in D^b(\text{Sq}_S)$ , then  $\mathbf{D}_S(M^\bullet) \in D_{\text{Sq}_S}^b(*\text{gr } S)$  (see [13]). So  $\mathbf{D}_S$  gives a duality functor on  $D^b(\text{Sq}_S)$ .

**Definition 2.2** (Römer [11]). A  $\mathbb{Z}^n$ -graded  $E$ -module  $N = \bigoplus_{\mathbf{a} \in \mathbb{Z}^n} N_{\mathbf{a}}$  is *squarefree* if  $N$  is finitely generated and  $N = \bigoplus_{F \subset [n]} N_{-F}$ .

For example, any monomial ideal of  $E$  is a squarefree module. Any monomial ideal of  $E$  is of the form  $J_\Delta = (\prod_{i \in F} y_i \mid F \notin \Delta)$  for some simplicial complex  $\Delta \subset 2^{[n]}$ . We say  $K\{\Delta\} := E/J_\Delta$  is the *exterior face ring* of  $\Delta$ .

Let  $\text{Sq}_E$  be the full subcategory of  $*\text{gr } E$  consisting of squarefree  $E$ -modules. Then there exist functors  $\mathcal{S} : \text{Sq}_E \rightarrow \text{Sq}_S$  and  $\mathcal{E} : \text{Sq}_S \rightarrow \text{Sq}_E$  giving an equivalence  $\text{Sq}_S \cong \text{Sq}_E$ . Here  $\mathcal{S}(N)_F = N_{-F}$  for  $N \in \text{Sq}_E$ , and the multiplication map  $\mathcal{S}(N)_F \ni z \mapsto x_i z \in \mathcal{S}(N)_{F \cup \{i\}}$  for  $i \notin F$  is given by  $\mathcal{S}(N)_F = N_{-F} \ni z \mapsto (-1)^{\alpha(i, F)} y_i z \in N_{-(F \cup \{i\})} = \mathcal{S}(N)_{F \cup \{i\}}$ , where  $\alpha(i, F) = \#\{j \in F \mid j < i\}$ . For example,  $\mathcal{S}(K\{\Delta\}) = K[\Delta]$ . See [11] for further information. Of course,  $\mathcal{S}$  and  $\mathcal{E}$  can be extended to the functors between  $D^b(\text{Sq}_S)$  and  $D^b(\text{Sq}_E)$ .

If  $N \in \text{Sq}_E$ , then  $N^* = \text{Hom}_E(N, E)$  is squarefree again. So  $(-)^*$  gives the duality functor  $\mathbf{D}_E$  on  $D^b(\text{Sq}_E)$ . For example,  $K\{\Delta\}^* = J_{\Delta^\vee}$ , where  $\Delta^\vee = \{F \subset [n] \mid [n] \setminus F \notin \Delta\}$  is the Alexander dual complex of  $\Delta$ . We have the *Alexander duality functor*  $\mathbf{A} := \mathcal{S} \circ \mathbf{D}_E \circ \mathcal{E}$  on  $\text{Sq}_S$  (or  $D^b(\text{Sq}_S)$ ). Of course,  $\mathbf{A}(K[\Delta]) = I_{\Delta^\vee}$ . In general, we have  $\mathbf{A}(H^i(M^\bullet))_F = (H^{-i}(M^\bullet)_{[n] \setminus F})^\vee$ .

An associated prime ideal of  $M \in *\text{gr } S$  is of the form  $P_F = (x_i \mid i \notin F)$  for some  $F \subset [n]$ . Let  $M \in \text{Sq}_S$  be a squarefree module. A monomial prime ideal  $P_F$  is a minimal prime of  $M$  if and only if  $F$  is a maximal element of the set  $\{G \subset [n] \mid M_G \neq 0\}$ . The following is a squarefree version of Definition 1.1.



**Definition 2.3.** We say  $(F, i) \in 2^{[n]} \times \mathbb{Z}$  is a *distinguished pair* for a complex  $M^\bullet \in D^b(\mathrm{Sq}_S)$ , if  $P_F$  is a minimal prime of  $H^i(M^\bullet)$  and  $H^j(M^\bullet)_G = 0$  for all  $j$  with  $j < i$  and  $G \supset F$  with  $|G| < |F| + i - j$ .

**Theorem 2.4.** *Let  $M^\bullet \in D^b(\mathrm{Sq}_S)$ . A pair  $(F, i)$  is distinguished for  $M^\bullet$  if and only if  $(F, -|F| - i)$  is distinguished for  $\mathbf{D}_S(M^\bullet)$ . If this is the case,  $\dim_K H^i(M^\bullet)_F = \dim_K H^{-|F|-i}(\mathbf{D}_S(M^\bullet))_F$ .*

*Proof.* Like the proof of Theorem 1.2, we consider the spectral sequence  $E_2^{p,q} = \mathrm{Ext}_S^p(H^{-q}(M^\bullet), \omega^\bullet) \Rightarrow \mathrm{Ext}_S^{p+q}(M^\bullet, \omega^\bullet)$ . Then  $E_r^{p,q}$  is squarefree for all  $p, q$  and  $r \geq 2$ . When we consider a distinguished pair  $(F, i)$ , we set

$$\dim_F M := \begin{cases} -\infty & \text{if } M_G = 0 \text{ for all } G \supset F \\ \max\{|G| \mid G \supset F, M_G \neq 0\} & \text{otherwise} \end{cases}$$

for  $M \in \mathrm{Sq}_S$ . Set  $d_i(M^\bullet) := \dim_F H^i(M^\bullet)$  and  $e_2^{p,q} := \dim_F \mathrm{Ext}_S^p(H^{-q}(M^\bullet), \omega^\bullet)$  for  $M^\bullet \in D^b(\mathrm{Sq}_S)$ . We also remark that  $\dim_K M_F = l_{S_{P_F}}(M \otimes_S S_{P_F})$  for  $M \in \mathrm{Sq}_S$ . The equation (1.1) holds in this context, and the proof of Theorem 1.2 works verbatim.  $\square$

If  $N^\bullet \in D^b(\mathrm{Sq}_E)$ , then it is easy to see that  $\mathbf{L}(N^\bullet)(-1) \in D^b(\mathrm{Sq}_S)$ . So  $\mathcal{L}(-) := \mathbf{L}(-)(-1)$  gives a functor from  $D^b(\mathrm{Sq}_E)$  to  $D^b(\mathrm{Sq}_S)$ . Moreover, we have  $\mathcal{L} \cong \mathbf{A} \circ \mathbf{D}_S \circ \mathcal{S}$  by [13, Proposition 4.3].

**Definition 2.5.** Let  $N^\bullet \in D^b(\mathrm{Sq}_E)$ . We say  $(F, i)$  is a *distinguished pair* for  $N^\bullet$  if it is a distinguished pair for  $\mathcal{L}(N^\bullet) \in D^b(\mathrm{Sq}_S)$  in the sense of Definition 2.3.

The next result can be proved by the same way as Corollary 1.8 using Theorem 2.4.

**Proposition 2.6.** *Let  $N^\bullet \in D^b(\mathrm{Sq}_E)$ . A pair  $(F, i)$  is distinguished for  $N^\bullet$  if and only if  $(F, 2n - |F| - i)$  is distinguished for  $\mathbf{D}_E(N^\bullet)$ . If this is the case, we have  $\dim_K H^i(\mathcal{L}(N^\bullet))_F = \dim_K H^{2n-|F|-i}(\mathcal{L} \circ \mathbf{D}_E(N^\bullet))_F$ .*

If  $M^\bullet \in D^b(*\mathrm{gr} S)$ , then  $\mathrm{Tor}_i^S(K, M^\bullet) := H^{-i}(K \otimes_E P^\bullet)$  is a  $\mathbb{Z}^n$ -graded module, where  $P^\bullet$  is a graded free resolution of  $M^\bullet$ . Set  $\beta_{i,\mathbf{a}}(M^\bullet) := \dim_K \mathrm{Tor}_i^S(K, M^\bullet)_{\mathbf{a}}$  for  $\mathbf{a} \in \mathbb{Z}^n$ . We say  $\beta_{i,\mathbf{a}}(M^\bullet)$  is the  $(i, \mathbf{a})$ th Betti number of  $M^\bullet$ . If  $M^\bullet \in D^b(\mathrm{Sq}_S)$  and  $\beta_{i,\mathbf{a}}(M^\bullet) \neq 0$ , then  $\mathbf{a}$  is squarefree (see [13]).

**Definition 2.7** (c.f. [3]). A Betti number  $\beta_{i,F}(M^\bullet) \neq 0$  is *extremal* if  $\beta_{j,G}(M^\bullet) = 0$  for all  $(j, G) \neq (i, F)$  with  $j \geq i$ ,  $G \supset F$ , and  $|G| - j > |F| - i$ .

Some of known results and backgrounds of extremal Betti numbers are found in the introduction of the present paper.

**Proposition 2.8** (c.f. [2]). *Let  $M^\bullet \in D^b(\mathrm{Sq}_S)$  and  $N^\bullet := \mathcal{E}(M^\bullet) \in D^b(\mathrm{Sq}_E)$ . A pair  $(F, i)$  is distinguished for  $\mathbf{D}_E(N^\bullet)$  if and only if  $\beta_{i+|F|-n,F}(M^\bullet)$  is an extremal Betti number. If this is the case, then  $\beta_{i+|F|-n,F}(M^\bullet) = \dim_K H^i(\mathcal{L} \circ \mathbf{D}_E(N^\bullet))_F$ .*

*Proof.* For  $j \in \mathbb{Z}$  and  $G \subset [n]$ , we have the following.

$$\begin{aligned} \beta_{j,G}(M^\bullet) &= \dim_K[H^{|G|-j-n}(\mathbf{D}_S \circ \mathbf{A}(M^\bullet))]_{[n] \setminus G} \quad (\text{by [13, Corollary 3.6]}) \\ &= \dim_K[H^{n+j-|G|}(\mathbf{A} \circ \mathbf{D}_S \circ \mathbf{A}(M^\bullet))]_G \\ &= \dim_K[H^{n+j-|G|}(\mathcal{L} \circ \mathcal{E} \circ \mathbf{A}(M^\bullet))]_G \\ &= \dim_K[H^{n+j-|G|}(\mathcal{L} \circ \mathbf{D}_E(N^\bullet))]_G. \end{aligned}$$

The assertion easily follows from this equality.  $\square$

**Corollary 2.9.** *Let  $M^\bullet \in D^b(\text{Sq}_S)$ . A Betti number  $\beta_{i,F}(M^\bullet)$  is extremal if and only if so is  $\beta_{|F|-i,F}(\mathbf{A}(M^\bullet))$ . If this is the case,  $\beta_{i,F}(M^\bullet) = \beta_{|F|-i,F}(\mathbf{A}(M^\bullet))$ .*

*Proof.* If  $\beta_{i,F}(M^\bullet)$  is extremal, then  $(F, n + i - |F|)$  is a distinguished pair for  $\mathbf{D}_E \circ \mathcal{E}(M^\bullet)$  by Proposition 2.8. By Proposition 2.6,  $(F, n - i)$  is a distinguished pair for  $\mathcal{E}(M^\bullet) \cong \mathbf{D}_E \circ \mathcal{E} \circ \mathbf{A}(M^\bullet)$ . So  $\beta_{|F|-i,F}(\mathbf{A}(M^\bullet))$  is extremal. The converse implication can be proved by the same way. The last equality follows from Proposition 2.6.  $\square$

This corollary generalizes results of Bayer-Charalambous-Popescu [3], Römer [11] and Miller [10]. Roughly speaking, the above proof is a “complex version” of [11]. But, his argument itself does not work in the  $\mathbb{Z}^n$ -graded context, since he use a generic base change of  $V = E_{-1}$ .

For  $M^\bullet \in D^b(\text{Sq}_S)$ . Set  $\text{proj. dim}(M^\bullet) = \max\{i \mid \beta_{i,F}(M^\bullet) \neq 0 \text{ for some } F\}$  and  $\text{reg}(M^\bullet) = \max\{|F| - i \mid \beta_{i,F}(M^\bullet) \neq 0\}$ . Since Betti numbers  $\beta_{i,F}(M^\bullet)$  which give  $\text{proj. dim}(M^\bullet)$  or  $\text{reg}(M^\bullet)$  are extremal, the next result follows from Corollary 2.9.

**Corollary 2.10** (c.f.[10, 11]). *If  $M^\bullet \in D^b(\text{Sq}_S)$ , then  $\text{proj. dim}(M^\bullet) = \text{reg}(\mathbf{A}(M^\bullet))$ .*

## REFERENCES

- [1] A. Aramova, L. Avramov and J. Herzog, Resolutions of monomial ideals and cohomology over exterior algebras. *Trans. Amer. Math. Soc.* **352** (2000), 579-594.
- [2] A. Aramova and J. Herzog, Almost regular sequences and Betti numbers, *Amer. J. Math.* **122** (2000), 689-719.
- [3] D. Bayer, H. Charalambous and S. Popescu, Extremal Betti numbers and applications to monomial ideals, *J. Algebra* **221** (1999), 497-512.
- [4] E. Babson, I. Novik and R. Thomas, Symmetric iterated Betti numbers, preprint (math.CO/0206063).
- [5] W. Bruns and J. Herzog, Cohen-Macaulay rings, revised edition, Cambridge University Press, 1998.
- [6] D. Eisenbud, G. Fløystad and F.-O. Schreyer, Sheaf cohomology and free resolutions over exterior algebra, *Trans. Amer. Math. Soc.* **355** (2003), 4397-4426.
- [7] D. Eisenbud, S. Popescu and S. Yuzvinsky, Hyperplane arrangement cohomology and monomials in the exterior algebra, *Trans. Amer. Math. Soc.* **355** (2003), 4365-4383.
- [8] S.I. Gelfand and Y.I. Manin, *Methods of homological algebra*. Springer-Verlag, 1996.
- [9] G. Kalai, Algebraic shifting, in “Computational commutative algebra and combinatorics” (T. Hibi, ed.), pp. 121-163, *Adv. Stud. Pure Math.* **33**, Math. Soc. Japan, 2002.
- [10] E. Miller, The Alexander duality functors and local duality with monomial support, *J. Algebra*. **231** (2000), 180-234.
- [11] T. Römer, Generalized Alexander duality and applications, *Osaka J. Math.* **38** (2001), 469-485.

- [12] K. Yanagawa, Alexander duality for Stanley-Reisner rings and squarefree  $\mathbb{N}^n$ -graded modules, *J. Algebra* **225** (2000), 630–645.
- [13] K. Yanagawa, Derived category of squarefree modules and local cohomology with monomial ideal support, to appear in *J. Math. Soc. Japan*.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY,  
TOYONAKA, OSAKA 560-0043, JAPAN  
*E-mail address:* yanagawa@math.sci.osaka-u.ac.jp