A TRIPLE LACUNARY GENERATING FUNCTION FOR HERMITE POLYNOMIALS

IRA M. GESSEL AND PALLAVI JAYAWANT

ABSTRACT. Some of the classical orthogonal polynomials such as Hermite, Laguerre, Charlier, etc. have been shown to be the generating polynomials for certain combinatorial objects. These combinatorial interpretations are used to prove new identities and generating functions involving these polynomials. In this paper we apply Foata's combinatorial interpretation of the Hermite polynomials as counting matchings of a set to obtain a triple lacunary generating function for the Hermite polynomials. We also give an umbral proof of this generating function.

1. INTRODUCTION

Some of the classical orthogonal polynomials such as Hermite, Laguerre, Charlier, etc. have been shown to be the generating polynomials for certain combinatorial objects ([2], [4], [6]). These combinatorial interpretations can be used to prove new identities and generating functions involving these polynomials. In this paper we focus on the combinatorial interpretation of the Hermite polynomials $H_n(u)$, which may be defined by the exponential generating function

(1)
$$\sum_{n=0}^{\infty} H_n(u) \frac{z^n}{n!} = e^{2uz - z^2}.$$

For our combinatorial interpretation, it is more convenient to take a different normalization, which makes all the coefficients positive, so we work with the polynomials $h_n(u) = \frac{i^n}{2^{n/2}} H_n\left(\frac{-iu}{\sqrt{2}}\right)$, where $i = \sqrt{-1}$, given by the generating function $\sum_{n=0}^{\infty} h_n(u) \frac{z^n}{n!} = e^{uz+z^2/2}.$

All of our formulas for $h_n(u)$ are easily converted into formulas for $H_n(u)$.

Date: March 3, 2004.

The first author was partially supported by NSF Grant DMS-0200596.

Foata [5] gave a combinatorial proof of Doetsch's identity [3] that gives a generating function for $h_{2n}(u)$:

(2)
$$\sum_{n=0}^{\infty} h_{2n}(u) \frac{z^n}{n!} = (1-2z)^{-1/2} \exp\left(\frac{u^2 z}{1-2z}\right).$$

We will prove the following generating function for $h_{3n}(u)$:

(3)
$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = \frac{\exp\left(w^3 z - 9w^4 z^2/2\right)}{\sqrt{1 - 6wz}} \sum_{n=0}^{\infty} \frac{(6n)!}{2^{3n} (3n)! (1 - 6wz)^{3n}} \frac{z^{2n}}{(2n)!},$$

in which $w = (1 - \sqrt{1 - 12uz})/6z = uC(3uz)$, where $C(x) = (1 - \sqrt{1 - 4x})/2x$ is the Catalan number generating function. Note that the formula can be written in terms of hypergeometric series:

$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = \frac{\exp\left(w^3 z - 9w^4 z^2/2\right)}{\sqrt{1 - 6wz}} {}_2F_0\left(\frac{1}{6}, \frac{5}{6}; -; \frac{54z^2}{(1 - 6wz)^3}\right),$$

where $_{2}F_{0}(a, b, -; z) = \sum_{n \ge 0} (a)_{n} (b)_{n} \frac{z^{n}}{n!}$, and

$$(\alpha)_n = \alpha(\alpha+1)\dots(\alpha+n-1)$$

is the rising factorial.

We prove formula (3) by two methods—umbral and combinatorial. In section 2, we define an umbra and study some properties that are useful in the proof of the formula. The umbral proof follows in section 3. An umbral proof of a generating function for $h_{2m+n}(u)$ is given in [7].

To prove (3) combinatorially, we show that both sides enumerate the same weighted objects. To begin with, we describe the combinatorial interpretation of the Hermite polynomials in section 4. In section 5 we begin the combinatorial proof of the generating function (3) for $h_{3n}(u)$ by introducing the objects counted by $w = (1 - \sqrt{1 - 12uz})/6z = uC(3uz)$; section 6 gives the details of the objects counted by both sides of the formula.

By using these methods, it would be possible to give combinatorial proofs of generating functions in two variables for $h_{m+n}(u)$ and $h_{3m+2n+k}(u)$.

2. The umbra and its properties

Rota and Taylor in [8] laid a rigorous foundation for the classical umbral calculus. They consider a vector space of polynomials in several variables or "umbrae" and define the linear functional eval on it. A sequence (a_n) is represented by an umbra A if $eval(A^n) = a_n$ for all n. In practice, the word eval is usually dropped and we simply write $A^n = a_n$ with the understanding that the functional has been applied. When we write f(A) = g(A), we mean eval(f(A)) = eval(g(A)). We consider formal power series f(t) with coefficients in a ring of formal power series $R = \mathbf{Q}[[x, y, z, \cdots]]$.

Definition 2.1. A formal power series $f(t) = \sum_{n=0}^{\infty} f_n t^n$ is admissible if for every monomial $x^i y^j z^k \cdots$ in R, the coefficient of $x^i y^j z^k \cdots$ in f_n is nonzero for only finitely many values of n.

So, for example, $f(t) = e^{xt}$ is an admissible formal power series, while $f(t) = e^t$ is not. Some computations similar to those of this section can be found in section 4 of Gessel [7].

With this brief introduction to umbral calculus, we will define an umbra M that will be shown to be related to the Hermite polynomials in the next section. In this section, we will study some properties of this umbra that we will need. We define the umbra M by $e^{Mz} = e^{z^2/2}$ so that

(4)
$$M^{n} = \begin{cases} \frac{(2k)!}{2^{k}k!}, & \text{if } n = 2k \\ 0, & \text{if } n \text{ is odd.} \end{cases}$$

The following two formulas hold for M.

Lemma 2.2. (i) For any admissible formal power series f, $e^{Mz}f(M) = e^{z^2/2}f(M+z)$.

(*ii*)
$$e^{M^2 z} = \frac{1}{\sqrt{1-2z}}.$$

Proof. (i) We first observe that it is sufficient to prove that the formula holds for $f(t) = e^{tx}$: If the formula is true for $f(t) = e^{tx}$, then comparing coefficients of $x^n/n!$ on both sides we get that the formula holds for $f(t) = t^n$. But this implies by linearity that the formula is true for all admissible formal power series f. To prove the formula for $f(t) = e^{tx}$, we have

$$e^{Mz}e^{Mx} = e^{M(z+x)} = e^{z^2/2 + zx + x^2/2} = e^{z^2/2}e^{zx}e^{Mx} = e^{z^2/2}e^{(M+z)x}.$$

(ii) We have

$$e^{M^2 z} = \sum_{n=0}^{\infty} M^{2n} \frac{z^n}{n!} = \sum_{n=0}^{\infty} \frac{(2n)!}{2^n n!} \frac{z^n}{n!} = \frac{1}{\sqrt{1-2z}}.$$

Corollary 2.3. For any admissible formal power series f,

$$e^{M^2 z} f(M) = \frac{1}{\sqrt{1-2z}} f\left(\frac{M}{\sqrt{1-2z}}\right).$$

Proof. As in the lemma, it is sufficient to prove the formula for $f(t) = e^{tx}$. Here $e^{M^2 z} f(M) = e^{M^2 z + Mx}$. If we apply Lemma 2.2 (i) directly, we cannot eliminate the linear term in M. So we introduce a parameter α and rewrite $e^{M^2 z + Mx}$ as $e^{M\alpha} e^{M^2 z + M(x-\alpha)}$. We will choose a value for α later. Now applying Lemma 2.2 (i) we get

$$e^{M\alpha}e^{M^2z+M(x-\alpha)} = e^{\alpha^2/2}e^{(M+\alpha)^2z+(M+\alpha)(x-\alpha)} = e^{M^2z+M(x-(1-2z)\alpha)+z\alpha^2+x\alpha-\alpha^2/2}.$$

Now we choose the value of α to eliminate the term in M on the right; i.e., we solve $x - (1-2z)\alpha = 0$ and get $\alpha = x/(1-2z)$. Substituting this value of α in the above expression and simplifying, we obtain that $e^{M^2z+Mx} = e^{M^2z}e^{x^2/2(1-2z)}$. By Lemma 2.2 (ii), this is equal to $\frac{1}{\sqrt{1-2z}}e^{x^2/2(1-2z)}$. But applying the definition of M directly gives

$$\frac{1}{\sqrt{1-2z}} f\left(\frac{M}{\sqrt{1-2z}}\right) = \frac{1}{\sqrt{1-2z}} e^{Mx/\sqrt{1-2z}} = \frac{1}{\sqrt{1-2z}} e^{x^2/2(1-2z)}.$$

Applying the corollary to $f(M) = e^{M^3 x}$ gives the formula

(5)
$$e^{M^2 z + M^3 x} = \frac{\exp\left(\frac{M^3 x}{(1 - 2z)^{3/2}}\right)}{\sqrt{1 - 2z}},$$

which we will need in the next section.

3. The umbral proof

To prove formula (3), we first express the Hermite polynomial $h_n(u)$ in terms of the umbra M. We have

$$\sum_{n=0}^{\infty} h_n(u) \frac{z^n}{n!} = e^{uz + z^2/2} = e^{(u+M)z}.$$

Comparing the coefficients of $z^n/n!$ on both sides, we get $h_n(u) = (u+M)^n$. Using this, we get

$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = \sum_{n=0}^{\infty} (u+M)^{3n} \frac{z^n}{n!} = e^{(u+M)^3 z} = e^{(u^3+3Mu^2+3M^2u+M^3)z}.$$

We follow the same procedure as in the proof of Corollary 2.3. We introduce a parameter α and rewrite the last expression as $e^{M\alpha}e^{u^3z+M(3u^2z-\alpha)+3M^2uz+M^3z}$. Now applying Lemma 2.2 (i) we get

(6)
$$e^{M\alpha}e^{u^3z+M(3u^2z-\alpha)+3M^2uz+M^3z} = e^{\alpha^2/2}e^{u^3z+(M+\alpha)(3u^2z-\alpha)+3(M+\alpha)^2uz+(M+\alpha)^3z}$$

= $e^{z\alpha^3+3uz\alpha^2-\alpha^2/2+3u^2z\alpha+u^3z+M(3z\alpha^2+(6uz-1)\alpha+3u^2z)+M^2(3z\alpha+3uz)+M^3z}$.

In order to apply formula (5), we need to eliminate the linear term in M. So we choose a value of α that makes the coefficient of M equal to zero. By solving a quadratic and taking the solution with a power series expansion, we get $\alpha = (1-\sqrt{1-12uz})/6z-u$. We know that the Catalan generating function C(x) is given by $(1-\sqrt{1-4x})/2x$. In terms of this generating function, $\alpha = uC(3uz)-u = w-u$ where w = uC(3uz). Using this value of α to simplify expression (6), we get that

$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = \exp(w^3 z - (w-u)^2/2 + 3M^2 w z + M^3 z).$$

We know that C(x) satisfies the equation $C(x) = 1 + x(C(x))^2$. Substituting x = 3uz and using the fact that C(3uz) = w/u, we obtain $w = u + 3w^2z$; i.e., $w - u = 3w^2z$. This gives us

$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = e^{w^3 z - 9w^4 z^2/2} e^{3M^2 w z + M^3 z}.$$

Applying formula 5 with $f(t) = t^3 z$, we get that

$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = \frac{\exp\left(w^3 z - 9w^4 \frac{z^2}{2}\right)}{\sqrt{1 - 6wz}} \exp\left(\frac{M^3 z}{(1 - 6wz)^{3/2}}\right).$$

Then writing the second exponential function as a series and using (4), we obtain the final result:

$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = \frac{\exp\left(w^3 z - 9w^4 \frac{z^2}{2}\right)}{\sqrt{1 - 6wz}} \sum_{n=0}^{\infty} \frac{(6n)!}{2^{3n}(3n)!(1 - 6wz)^{3n}} \frac{z^{2n}}{(2n)!}.$$

Now we turn to the combinatorial method of proof. We begin with a combinatorial interpretation of the Hermite polynomials that will be used in the combinatorial proofs.

4. Combinatorial interpretation of Hermite Polynomials

We assume that the reader is familiar with enumerative applications of exponential generating functions, as described, for example, in [9, Chapter 5] and [1]. The product formula and the exponential formula for exponential generating functions that appear in [9] play an important role in the combinatorial proofs.

The exponential generating function $uz + z^2/2$ counts sets with one or two elements, where a one-element set is weighted u. Then by the exponential formula, the coefficient of $z^n/n!$ in $e^{uz+z^2/2}$, which is $h_n(u)$, is the generating polynomial for partitions of an *n*-element set into blocks of size one or two, where each block of size one is weighted u. (If we used $H_n(u)$ as Foata did, instead of $h_n(u)$, we would need to attach a weight of -2 to each two-element block and a weight of 2u to each one element block.) It is convenient to represent these partitions as graphs in which the vertices in a two-element block are joined by an edge. We call these graphs, in which every vertex has degree at most one, *matchings*.

Figure 1 shows a matching on the vertex set $\{1, 2, 3, 4, 5, 6\}$ that corresponds to the partition $\{\{1, 3\}, \{2, 6\}, \{4\}, \{5\}\}$. Thus the Hermite polynomial $h_n(u)$ can be



FIGURE 1. Matching

viewed as the generating polynomial for the number of vertices of degree zero over the set of all matchings on n vertices, where each vertex of degree zero is assigned the weight u. With this combinatorial interpretation we will prove the following formula:

(7)

$$\sum_{n=0}^{\infty} h_{3n}(u) \frac{z^n}{n!} = \exp(w^3 z - 9w^4 z^2/2) \frac{1}{(1 - 6wz)^{1/2}} \sum_{n=0}^{\infty} \frac{(6n)!}{2^{3n} (3n)! (1 - 6wz)^{3n}} \frac{z^{2n}}{(2n)!}$$

First we take a look at the graphs counted by w = uC(3uz), where C(x) is the Catalan generating function.

5. w-Trees

The Catalan generating function C(z) can be viewed as the ordinary generating function for rooted binary trees with internal vertices weighted by z. By a rooted binary tree, we mean a rooted tree in which every vertex has zero or two children. A vertex with zero children is called a leaf while a vertex with two children is called an internal vertex. Thus every internal vertex (except the root) has degree three whereas a leaf has degree one. We will think of an edge as being made up of two "half-edges," each half-edge incident with one of the end-vertices. For our purpose (which will become clear in the next section) in any rooted tree we will attach one extra half-edge to the root. Now every internal vertex in a rooted binary tree has degree three; i.e., every internal vertex is trivalent. Figure 2 shows such a tree. Note that in a rooted binary tree, the order of the children at any internal vertex



FIGURE 2. Rooted binary tree

matters. Figure 3 shows two binary trees that are distinct. So if $C(z) = \sum_{n=0}^{\infty} c_n z^n$,



FIGURE 3. Distinct binary trees

then c_n is the number of rooted binary trees (as shown in Figure 2) with n internal vertices.

Definition 5.1. A w-tree is a rooted tree with labeled trivalent internal vertices in which the three half-edges incident with every internal vertex are marked 1, 2 or 3. Moreover, the three marks at each internal vertex are distinct.

Figure 4 shows a *w*-tree. Thus a *w*-tree is similar to a rooted binary tree with



FIGURE 4. w-tree

labeled internal vertices, with one difference—in a w-tree, the order of the children at any internal vertex does not matter. Figure 5 shows two w-trees that are same. A special case of a w-tree is the one with no internal vertices and no marks. It is



FIGURE 5. Same w-trees

simply a leaf with a half-edge attached to it.

5.1. Generating function for *w*-trees. We will find the exponential generating function in z for *w*-trees with each internal vertex weighted with z and each leaf weighted with u. To count *w*-trees with n internal vertices, we will construct such a *w*-tree starting with a rooted binary tree with n internal vertices. Then there are n! ways to label the internal vertices. Now we need to put in the marks on the half-edges at the internal vertices. In order to avoid counting a *w*-tree twice, we will take the canonical representation of a *w*-tree to be the representation in which the two half-edges at an internal vertex connected to the two children are marked in an increasing order from left to right. Then in Figure 5, the canonical representation is the one on the left. Thus while marking the half-edges, there are 3 choices to mark the half-edge at an internal vertex going to the parent (in case of the root, it is the extra half-edge) and then the other marks get decided

automatically. Hence the number of w-trees with n internal vertices is $n! c_n 3^n$. Since a w-tree with n internal vertices has n + 1 leaves, the exponential generating function for w-trees with the appropriate weights is

$$\sum_{n=0}^{\infty} n! c_n 3^n u^{n+1} \frac{z^n}{n!} = uC(3uz) = w.$$

Thus w = uC(3uz) is the exponential generating function for weighted w-trees.

With this interpretation of w, we will begin the combinatorial proof of formula (7).

6. The combinatorial proof

We will describe the graphs enumerated by the left side of formula (7). Then we will describe the same graphs in terms of their connected components and use the product formula for exponential generating functions to complete the proof.

6.1. Graphs counted by the left side. The left side of formula (7) is an exponential generating function for $h_{3n}(u)$. From the interpretation of $h_n(u)$ given above, it follows that $h_{3n}(u)$ is the generating polynomial for the number of vertices of degree zero over the set of all matchings on 3n vertices. But we cannot have 3n labeled vertices in these matchings. Since the power of z in the exponential generating function is n, we have only n labeled vertices in these matchings. We need to modify the matchings to obtain the graphs counted by the left side. We start with n vertices labeled $1, 2, \ldots, n$ and use them to produce 3n vertices using the construction shown in Figure 6 for each labeled vertex i. Each labeled vertex is



FIGURE 6. One vertex produces three vertices

adjacent to three vertices marked 1, 2 and 3 and these marked vertices are distinct for distinct labeled vertices. The n labeled vertices produce 3n marked vertices on which we construct a matching. The graph consisting of the n labeled vertices along with the matching on the 3n marked vertices is a typical graph counted by the left side. Figure 7 shows such a graph.

Let G be the set of all graphs enumerated by the left side of formula (7). We can describe a graph in G as a graph in which every vertex has degree one (monovalent



FIGURE 7. Typical graph counted by left side

vertex), degree two (bivalent vertex), or degree three (trivalent vertex) and the vertices satisfy the following adjacency criteria: A trivalent vertex is adjacent to monovalent or bivalent vertices and has weight z. A monovalent vertex is adjacent to a trivalent vertex and has weight u. A bivalent vertex is adjacent to a trivalent vertex and to another bivalent vertex. This ensures that the monovalent and bivalent vertices constitute a matching. The trivalent vertices have labels and the monovalent and bivalent vertices are marked.

The graphs in G are made up of connected components and we will soon find out that these graphs have three types of connected components. If we show that each term on the right side of formula (7) is the exponential generating function for graphs that contain only one type of connected component, then the product formula for exponential generating functions tells us that the product of the terms enumerates the graphs in G. First we need to determine the various types of connected components of the graphs in G. Then for each type we will find the exponential generating functions for graphs that contain only that type of connected component.

6.2. Connected components of graphs in G. To start with, we eliminate all bivalent vertices from the graphs in G. A bivalent vertex is adjacent to one trivalent vertex and to another bivalent vertex. We transfer the mark on the bivalent vertex to the half-edge at the adjacent trivalent vertex. We also transfer the mark on a monovalent vertex to the half-edge at the adjacent trivalent vertex. Figure 8 shows the graph in Figure 7 with all bivalent vertices eliminated and all the marks transferred to the half-edges.

We see from Figure 8 that this elimination may create some loops at some of the vertices. We think of a loop as contributing two to the degree of a vertex since a loop is an edge made up of two half-edges and each half-edge contributes one to the degree of the vertex. So every vertex in the new graph is monovalent or trivalent. Each of the three edges coming out of a trivalent vertex is incident with another trivalent vertex or to the same vertex or to a monovalent vertex. Each half-edge at a trivalent vertex has a mark and hence every loop and every edge joining two trivalent vertices has two marks on its two half-edges. We can easily recover the original graph by adding a vertex at each half-edge of such an edge or loop. Thus the graphs in G are reduced to graphs (with loops) which have only trivalent or monovalent vertices. Each trivalent vertex is weighted z and has marked half-edges incident with it while each monovalent vertex is weighted u.

The connected components of a graph in G are of three types: components that do not contain any cycles (acyclic), components with only one cycle, and components with more than one cycle. Note that a loop is considered to be a cycle.



FIGURE 8. Graph in Figure 7 with bivalent vertices eliminated

6.3. Graphs with acyclic components. An acyclic connected component is a tree. Figure 9 shows such a tree. We wish to count such unrooted trees. To do



FIGURE 9. Tree component

this, we first count vertex-rooted and edge-rooted versions of these trees. First we count trees rooted at a trivalent vertex. In such a tree, the three half-edges at the root are joined to w-trees. So we can construct such a rooted tree by taking a set of three w-trees, with exponential generating function $w^3/3!$, and a new root vertex with its three marked half-edges, and connected the half-edges of the new root to the half-edges of the three w-trees in 3! ways. Thus the exponential generating for these rooted trees is w^3z .

Next we count versions of these trees rooted at an edge joining two trivalent vertices. With the help of Figure 10 we see that the exponential generating function for such edge-rooted trees is $9w^4z^2/2$. An unrooted tree with *n* trivalent vertices is



FIGURE 10. Edge-rooted tree

counted n times in the generating function for rooted trees as there are n choices for

the root. Since there are n-1 ways to choose an edge-root, such a tree is counted n-1 times in the generating function for edge-rooted trees. In order to get the generating function for unrooted trees, we subtract one generating function from the other so that each tree is counted once. Hence the exponential generating function for graphs whose connected components are only trees is $\exp(w^3z - 9w^4z^2/2)$.

Now let us look at the connected components with one or more cycles. The two components besides the tree in the graph in Figure 8 are such components. First we reduce these components by shrinking all the w-trees present in them. Figure 11 illustrates this process. As we see in the picture, the process of shrinking w-trees



FIGURE 11. Components with cycles with w-trees eliminated

leaves behind vertices weighted with z which are now bivalent. We further reduce the components by eliminating these newly created bivalent vertices. Figure 12 shows what we get after this further reduction. Thus we get either circles with no vertices or components (with multiple edges and loops) in which each vertex is trivalent. We will consider the two cases separately.

6.4. Components with one cycle. If the reduction of a component of a graph in G results in a circle, this means the original component has exactly one cycle and each trivalent vertex on the cycle has a w-tree attached at its third half-edge; i.e, there is a cycle of w-trees. We can see this if we trace the circle component in Figure



FIGURE 12. Graph in Figure 11 with bivalent vertices eliminated

12 back to its original component in Figure 8. The exponential generating function for directed cycles is $\sum_{n=3}^{\infty} z^n/n$ and hence for undirected cycles is $(1/2) \sum_{n=3}^{\infty} z^n/n$. However, in this case we have a cycle of vertices with a *w*-tree attached to each vertex which indicates that we need to replace *z* with *wz* in the generating function. Moreover, the half-edges at each vertex of the cycle have distinct marks and this marking can be done in 3! = 6 ways. So in fact, we need to replace *z* with 6wz. Thus the exponential generating function for undirected cycles of *w*-trees is $(1/2) \sum_{n=3}^{\infty} (6wz)^n/n$. Note that the sum begins with n = 3 because three vertices are needed to form a cycle when no loops and multiple edges are allowed. But the graphs we are counting do have loops and multiple edges. So there can be a cycle with one or two vertices.

Figure 13 shows a cycle with only one vertex. The exponential generating func-



FIGURE 13. Cycle with only one vertex

tion for such cycles is 3wz as there are 3 ways to mark the half-edge which goes to the *w*-tree and the choice of marks for the other two half-edges does not matter since they are connected to each other. Figure 14 shows a cycle with two vertices. The marks for each of the half-edges



With the remaining marks two ways to form the cycle

FIGURE 14. Cycle with two vertices

going to the w-trees can be chosen in 3 ways and then there are two ways to form the cycle. Hence the exponential generating function for these cycles is $18w^2z^2/2!$. Thus the exponential generating function for cycles of w-trees is

$$\frac{1}{2}\sum_{n=3}^{\infty}\frac{(6wz)^n}{n} + 3wz + 18w^2\frac{z^2}{2!} = \frac{1}{2}\sum_{n=1}^{\infty}\frac{(6wz)^n}{n} = \log\left(1 - 6wz\right)^{-1/2}$$

Then the exponential generating function for graphs whose connected components are only cycles of w-trees is $\exp(\log(1-6wz)^{-1/2}) = (1-6wz)^{-1/2}$.

6.5. Components with more than one cycle. The components of graphs in G that reduce to components with only trivalent vertices are the ones with more than one cycle. After reduction, these components can be together thought of as a graph whose connected components have only trivalent vertices. Consider such a graph with m vertices. For the purposes of counting, we add a new vertex at each half-edge of a trivalent vertex. Then the graph looks as shown in Figure 15. It has 3m additional vertices and we can think of it as a complete matching on



FIGURE 15. Complete matching

16

3m vertices. A complete matching is a graph with labeled vertices in which every vertex has degree one; i.e, every vertex is matched with exactly one other vertex. Then it is clear that in a complete matching the number of vertices is even. For 3m to be even, m has to be even. So m = 2n for some n. Then the number of complete matchings on 6n vertices is $(6n)!/(2^{3n}(3n)!)$ and the number of edges in each complete matching is 3n. So the number of graphs with m = 2n vertices whose connected components have only trivalent vertices is $(6n)!/(2^{3n}(3n)!)$.

We can obtain the original components from the reduced graph by introducing an ordered sequence of vertices at each edge. The vertices which are introduced are trivalent vertices with marked half-edges and with a *w*-tree attached to one of the half-edges. We can see this for the component with trivalent vertices in Figure 12 by tracing it back to its original component in Figure 8. Thus we get a term $\sum_{k=0}^{\infty} (6wz)^k$ for each of the 3n edges in the reduced graph, and combining these terms, we get $1/(1-6wz)^{3n}$. As explained in subsection 6.4, we have 6wz instead of simply z. Hence the exponential generating function for components with more than one cycle is

$$\sum_{n=0}^{\infty} \frac{(6n)!}{2^{3n}(3n)!(1-6wz)^{3n}} \frac{z^{2n}}{(2n)!}.$$

We know that a connected component of any graph in G is either acyclic or contains exactly one cycle or has more than one cycle. In each case, we have found the exponential generating function for graphs whose connected components are only of that type. Then by the product formula for exponential generating functions, the exponential generating function for the graphs in G is the product of these generating functions which is

$$\exp(w^3 z - 9w^4 z^2/2) \frac{1}{(1 - 6wz)^{1/2}} \sum_{n=0}^{\infty} \frac{(6n)!}{2^{3n} (3n)! (1 - 6wz)^{3n}} \frac{z^{2n}}{(2n)!}$$

and this is the right side of formula (7).

References

- F. Bergeron, G. Labelle, and P. Leroux, Combinatorial species and tree-like structures, Encyclopedia of Mathematics and its Applications, 67, Cambridge University Press, Cambridge, 1997, Translated from the 1994 French original by Margaret Readdy.
- [2] A. de Médicis, D. Stanton, and D. White, The combinatorics of q-Charlier polynomials, J. Combinatorial Theory Ser. A 69 (1995), 87–114.
- [3] G. Doetsch, Integraleigenshaften der Hermiteschen Polynome, Math. Z. 32 (1930), 557–559.
- [4] D. Foata, A combinatorial proof of the Mehler formula, J. Combinatorial Theory Ser. A 24 (1978), 367–376.
- [5] D. Foata, Some Hermite polynomial identities and their combinatorics, Adv. in Appl. Math. 2 (1981), 250-259.

IRA M. GESSEL AND PALLAVI JAYAWANT

- [6] D. Foata and V. Strehl, Combinatorics of Laguerre Polynomials, Enumeration and design (Waterloo, Ont., 1982), Academic Press, Toronto, ON, 1984, pp. 123–140.
- [7] I. M. Gessel, Applications of the classical umbral calculus, Algebra Universalis 49 (2003), 397–434.
- [8] G.-C. Rota and B. D. Taylor, An introduction to the umbral calculus, Analysis, geometry and groups: a Riemann legacy volume, Hadronic Press, Palm Harbor, FL, 1993, pp. 513–525.
- [9] R. P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge University Press, Cambridge, 1999.

DEPARTMENT OF MATHEMATICS, BRANDEIS UNIVERSITY, WALTHAM MA 02454-9110 *E-mail address*: gessel@brandeis.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, TUCSON AZ 85721

E-mail address: jayawant@math.arizona.edu