AN ARITHMETIC FORMULA FOR CERTAIN COEFFICIENTS OF THE EULER PRODUCT OF HECKE POLYNOMIALS

Xian-Jin Li

ABSTRACT. In 1997 the author [11] found a criterion for the Riemann hypothesis for the Riemann zeta function, involving the nonnegativity of certain coefficients associated with the Riemann zeta function. In 1999 Bombieri and Lagarias [2] obtained an arithmetic formula for these coefficients using the "explicit formula" of prime number theory. In this paper, the author obtains an arithmetic formula for corresponding coefficients associated with the Euler product of Hecke polynomials, which is essentially a product of L-functions attached to weight 2 cusp forms (both newforms and oldforms) over Hecke congruence subgroups $\Gamma_0(N)$. The nonnegativity of these coefficients gives a criterion for the Riemann hypothesis for all these L-functions at once

1. Introduction

We denote by $S_2(N)$ the space of all cusp forms of weight 2 with the principal character modulo N for the Hecke congruence subgroup $\Gamma_0(N)$ of level N. That is, f belongs to $S_2(N)$ if, and only if, f is holomorphic in the upper half-plane, satisfies

$$f\left(\frac{az+b}{cz+d}\right) = (cz+d)^2 f(z)$$

for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$, satisfies the usual regularity conditions at the cusps of $\Gamma_0(N)$, and vanishes at each cusp of $\Gamma_0(N)$.

The Hecke operators T(n), $n = 1, 2, \dots$, are defined by

(1.1)
$$(T(n)f)(z) = \frac{1}{n} \sum_{ad=n, (a,N)=1} a^2 \sum_{0 \le b < d} f\left(\frac{az+b}{d}\right)$$

for any function $f \in S_2(N)$. The Fricke involution W is defined by

$$(Wf)(z) = N^{-1}z^{-2}f(-1/Nz),$$

and the complex conjugation operator K is defined by

$$(Kf)(z) = \bar{f}(-\bar{z}).$$

Set $\overline{W} = KW$. A function $f \neq 0$ in $S_2(N)$ is called a newform if it is an eigenfunction of \overline{W} and of all the Hecke operators T(n).

 $^{1991\} Mathematics\ Subject\ Classification.\ Primary\ 11M26,\ 11M36.$

Key words and phrases. Positivity condition, Weil's explicit formula.

Research supported by National Security Agency MDA904-03-1-0025

Let f be a newform in $S_2(N)$ normalized so that its first Fourier coefficient is 1. Then it have the Fourier expansion

$$f(z) = \sum_{n=1}^{\infty} \lambda(n)e^{2\pi i nz}$$

with the Fourier coefficients equal to the eigenvalues of Hecke operators. The Hecke L-function associated with f is given by

$$L_f(s) = \sum_{n=1}^{\infty} \frac{\lambda(n)}{n^s}$$

for $\Re s > \frac{3}{2}$; see §7.2 in Iwaniec [8]. It has the Euler product

(1.2)
$$L_f(s) = \prod_{p|N} (1 - \lambda(p)p^{-s})^{-1} \prod_{p \nmid N} (1 - \lambda(p)p^{-s} + p^{1-2s})^{-1}.$$

If we denote

(1.3)
$$\xi_f(s) = N^{s/2} (2\pi)^{-s} \Gamma(\frac{1}{2} + s) L_f(\frac{1}{2} + s),$$

then $\xi_f(s)$ is an entire function and satisfies the functional identity

$$\xi_f(s) = w\xi_f(1-s)$$

where $w = (-1)^r$ with r being the vanishing order of $\xi_f(s)$ at s = 1/2.

Let I be the identity map acting on the space $S_2(N)$. For each prime $p \nmid N$, we put

$$H_p(u) = \det(I - T(p)u + pu^2 I)$$

which we call a Hecke polynomial; see Ihara [7]. The Euler product $L_N(s)$ of Hecke polynomials is defined by

(1.4)
$$L_N(s) = \prod_{p \nmid N} H_p(p^{-s})^{-1},$$

which is the main object of study in this paper. Let

(1.5)
$$\xi_N(s) = N^{gs/2} (2\pi)^{-gs} \Gamma^g (\frac{1}{2} + s) L_N (\frac{1}{2} + s)$$

where g denotes the dimension of the space $S_2(N)$. It will be shown (Lemma 4.2) that $\xi_N(s)$ is a product of the $\xi_f(s)$'s over a Hecke eigenbasis of the space $S_2(N)$ times a finite Euler factor, that $\xi_N(s)$ is an entire function, and that zeros of $\xi_N(s)$ in the strip $0 < \Re s < 1$ appear in pairs ρ and $1 - \rho$.

Let

(1.6)
$$\tau_N(n) = \sum_{\rho} \left[1 - \left(1 - \frac{1}{\rho}\right)^{-n}\right]$$

for $n = 1, 2, \dots$, where the sum on ρ runs over all zeros of $\xi_N(s)$ taken in the order given by $|\Im \rho| < T$ for $T \to \infty$ with a zero of multiplicity ℓ appearing ℓ times in the list. If $\rho = 0$ is a zero of $\xi_N(s)$, then $(1 - 1/\rho)^{-n}$ in (1.6) is interpreted to be 0.

In order to state the main result of this paper (Theorem 1.2), we need an explicit formula for the trace $\operatorname{tr} T(p^k)$ of Hecke operators $T(p^k)$ acting on the space $S_2(N)$ for all primes $p \nmid N$ and for $k = 1, 2, \cdots$. This formula is given by the Eichler-Selberg trace formula obtained in Oesterlé [16]. We denote by φ the Euler φ -function. Let

$$\psi(N) = N \prod_{p|N} (1 + 1/p).$$

Lemma 1.1. Let p be a prime with (p, N) = 1. Then we have

$$tr(T(p^k)) = \frac{1 + (-1)^k}{24} \psi(N) + \frac{p^{k+1} - 1}{p - 1}$$

$$- \sum_{t \in \mathbb{Z}, t^2 < 4p^k} \sum_{\substack{m \in \mathbb{Z}^+, m^2 \mid t^2 - 4p^k \\ \frac{t^2 - 4p^k}{m^2} \equiv 0, 1 (\mod 4)}} \frac{h((t^2 - 4p^k)/m^2)}{w((t^2 - 4p^k)/m^2)} \frac{\psi(N)}{\psi(N/(N, m))} \mu_{t, p^k, m}$$

$$- \frac{1}{2} \sum_{0 < d \mid p^k} \min(d, p^k/d) \sum_{(c, \frac{N}{2}) \mid (N, \frac{p^k}{2} - d), c \mid N} \varphi((c, N/c))$$

for $k = 1, 2, \dots$, where $\mu_{t,p^k,m}$ is the number of solutions x modulo N of the equation $x^2 - tx + p^k \equiv 0 \pmod{N(N,m)}$, and where h(f) and w(f) are the class number and the number of units in the ring of integers of the imaginary quadratic field of discriminant f < 0 respectively.

Let $\gamma = 0.5772 \cdots$ be Euler's constant, let

$$\Lambda(k) = \begin{cases} \ln p, & \text{if } k \text{ is a positive power of a prime } p; \\ 0, & \text{otherwise,} \end{cases}$$

and let d(k) be the number of positive divisors of k. For $m = 1, 2, \dots$, we denote by ν_m the dimension of the subspace generated by all newforms in $S_2(m)$.

In this paper we obtain the following arithmetic formula for the $\tau_N(n)$'s, which generalizes an arithmetic formula of Bombieri and Lagarias [2] for the Riemann zeta function.

Theorem 1.2. Let $\tau_N(n)$ be given in (1.6). Then we have

$$\tau_N(n) = \frac{n}{2} \ln(N^{\nu_N} \prod_{\substack{m|N\\1 < m < N}} m^{\nu_m d(N/m)})$$

$$- \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{\substack{m=1\\(m,N)=1}}^{\infty} \frac{\Lambda(m)}{m^{3/2}} B(m) (\ln m)^{l-1}$$

$$- ng (\ln(8\pi) + \gamma - 2) + g \sum_{m=2}^n \binom{n}{m} \sum_{l=1}^{\infty} \frac{(-1)^m}{(l+1/2)^m}$$

for all positive integers n, where $B(p^k) = tr(T(p^k)) - p \, tr(T(p^{k-2}))$ for $p \nmid N$ and where the second term on the right side of the identity is interpreted as the limit

$$\lim_{X \to \infty} \sum_{l=1}^{n} \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{\substack{m < X \\ (m,N)=1}} \frac{\Lambda(m)}{m^{3/2}} B(m) (\ln m)^{l-1}$$

which exists.

In [2], Bombieri and Lagarias generalized a criterion of the author for the Riemann hypothesis [11] and obtained the following useful theorem.

Theorem 1.3. (Bombieri-Lagarias [2]) Let \mathcal{R} be a set of complex numbers ρ , whose elements have positive integral multiplicities assigned to them, such that $1 \notin \mathcal{R}$ and

$$\sum_{\rho} \frac{1 + |\Re \rho|}{(1 + |\rho|)^2} < \infty.$$

Then the following conditions are equivalent:

- (1) $\Re \rho \leq \frac{1}{2} \text{ for every } \rho \text{ in } \mathcal{R};$ (2) $\sum_{\rho} \Re [1 (1 \frac{1}{\rho})^{-n}] \geq 0 \text{ for } n = 1, 2, \cdots.$

As a corollary of Theorem 1.3 we obtain a criterion for the location of all nontrivial zeros of Hecke L-functions associated with all cusp forms which form an orthonormal basis in $S_2(N)$ and which consist of eigenfunctions of all the Hecke operators T(n) with (n, N) = 1.

Corollary 1.4. All zeros of $\xi_N(s)$ in the strip $0 < \Re s < 1$ lie on the critical line $\Re s = 1/2$ if, and only if, $\tau_N(n) \geq 0$ for all positive integers n.

In a subsequent paper [13], the author generalized results of this paper to the space $S_k(N,\chi)$ of cusp forms of weight k for all integers k>2 and of character χ for all Dirichlet characters of modulus N with $\chi(-1) = (-1)^k$.

This paper is organized as follows: The Eichler-Selberg trace formula for Hecke congruence subgroups, which is needed for Theorem 1.2, is given in section 2. An arithmetic formula is obtained in section 3 for a sequence of numbers, whose positivity implies that all zeros of Hecke L-functions associated with newforms lie on the critical line. This formula will be used in the proof of Theorem 1.2. In section 4 we give some preliminary results for the proof of Theorem 1.2. Finally, Theorem 1.2 is proved in section 5.

This research started while the author attended the Workshop on Zeta-Functions and Associated Riemann Hypotheses, New York University, Manhattan, May 29 -June 1, 2002. He wants to thank the American Institute of Mathematics, Brian Conrey, and Peter Sarnak for the support. He wishes to thank William Duke for his encouragement. The author also wishes to thank the referee for his valuable suggestions of improving the presentation of this paper.

2. The Eichler-Selberg trace formula

In this section, we state J. Oesterlé's formula [16] for the trace of Hecke operators acting on the space $S_2(N)$.

Let χ_0 be the principal character modulo N, and let $\chi_0(\sqrt{n}) = 0$ if n is not the square of an integer.

Lemma 2.1. (Théorème 3', [16]) (cf. Cohen [3]) For every positive integer n, the

trace tr(T(n)) of the Hecke operator T(n) acting on the space $S_2(N)$ is given by

$$tr(T(n)) = \frac{1}{12} \chi_0(\sqrt{n}) \psi(N) + \sum_{\substack{0 < d \mid n \\ \gcd(N, n/d) = 1}} d$$

$$- \sum_{t \in \mathbb{Z}, t^2 < 4n} \sum_{\substack{m \in \mathbb{Z}^+, m^2 \mid t^2 - 4n \\ \frac{t^2 - 4n}{m^2} \equiv 0, 1 (\mod 4)}} \frac{h((t^2 - 4n)/m^2)}{w((t^2 - 4n)/m^2)} \mu(t, n, m)$$

$$- \frac{1}{2} \sum_{0 < d \mid n} \min(d, n/d) \sum_{\substack{c \mid N \\ \gcd(c, \frac{N}{c}) \mid \gcd(N, \frac{n}{d} - d)}} \varphi(\gcd(c, N/c)) \chi_0(y)$$

where the integer y is defined modulo $N/\gcd(c,N/c)$ by $y \equiv d \pmod{c}, y \equiv n/d \pmod{N/c}$, where

$$\mu(t, n, m) = \frac{\psi(N)}{\psi(N/\gcd(N, m))} \sum_{\substack{x (\mod N) \\ x^2 - tx + n \equiv 0 (\mod N \gcd(N, m))}} \chi_0(x),$$

and where h(f) and w(f) are respectively the class number and the number of units in the ring of integers of the imaginary quadratic field of discriminant f < 0.

Lemma 2.2. Let $n = p^k$ for a prime $p \nmid N$ and for a positive integer k, let d|n, and let c|N. If y is defined modulo N/(c, N/c) by $y \equiv d \pmod{c}$ and $y \equiv n/d \pmod{N/c}$, then it is unique and (y, N) = 1.

Proof. Suppose that q|(y, N) for a prime q. Then q|c or q|N/c. Assume that q|c. Since $y \equiv d \pmod{c}$, we have q|d. This contradicts to that d is a power of p with (p, N) = 1. The case when q divides N/c can be treated similarly. Thus, we have proved that (y, N) = 1.

Suppose that y_1 is another integer satisfying the hypotheses. Then $y - y_1 \equiv 0 \pmod{c}$ and $y - y_1 \equiv 0 \pmod{N/c}$. Let $\ell = (c, N/c)$. We write $c = \ell c'$, $N/c = \ell N'$ for some integers c', N'. Then $y - y_1 \equiv 0 \pmod{\ell c' N'}$. That is,

$$y \equiv y_1 \pmod{N/\ell}$$
.

Hence, y is unique modulo $N/\gcd(c,N/c)$. \square

Proof of Lemma 1.1. Let $n=p^k$ for a prime $p \nmid N$ and for a positive integer k. Then we have

(2.1)
$$\frac{1}{12}\chi_0(\sqrt{n})\psi(N) = \frac{1+(-1)^k}{24}\psi(N)$$

and

(2.2)
$$\sum_{d>0,(N,n/d)=1,d|n} d = \frac{p^{k+1}-1}{p-1}.$$

Let x be an integer given by the equation $x^2 - tx + n \equiv 0 \pmod{N} \gcd(N, m)$, where m is an integer. Since $n = p^k$ with (p, N) = 1, we have (x, N) = 1. Hence, we have

$$\chi_0(x) = 1.$$

By (2.1), (2.2), (2.3), Lemma 2.1, and Lemma 2.2 we have

$$\operatorname{tr}(T(p^{k})) = \frac{1 + (-1)^{k}}{24} \psi(N) + \frac{p^{k+1} - 1}{p - 1}$$

$$- \sum_{t \in \mathbb{Z}, t^{2} < 4p^{k}} \sum_{\substack{m \in \mathbb{Z}^{+}, m^{2} \mid t^{2} - 4p^{k} \\ \frac{t^{2} - 4p^{k}}{m^{2}} \equiv 0, 1(\mod 4)}} \frac{h((t^{2} - 4p^{k})/m^{2})}{w((t^{2} - 4p^{k})/m^{2})} \frac{\psi(N)}{\psi(N/(N, m))} \mu_{t, p^{k}, m}$$

$$- \frac{1}{2} \sum_{0 < d \mid p^{k}} \min(d, p^{k}/d) \sum_{(c, \frac{N}{c}) \mid (N, \frac{p^{k}}{d} - d), c \mid N} \varphi((c, N/c))$$

where $\mu_{t,p^k,m}$ is the number of solutions x modulo N of the equation $x^2-tx+p^k\equiv 0 \pmod{N(N,m)}$, and where h(f) and w(f) are the class number and the number of units in the ring of integers of the imaginary quadratic field of discriminant f<0 respectively.

This completes the proof of the lemma. \Box

3. An arithmetic formula

In this section, an arithmetic formula is given for the $\tau_f(n)$ (see (3.1) below) which generalizes that of Bombieri and Lagarias [2] for the Riemann zeta function.

Let f be a normalized newform in $S_2(N)$, and let $\xi_f(s)$ be given in (1.3). Put

(3.1)
$$\tau_f(n) = \sum_{\rho} \left[1 - (1 - \frac{1}{\rho})^n\right]$$

for $n = 1, 2, \dots$, where the sum is over all the zeros of $\xi_f(s)$ in the order given by $|\Im \rho| < T$ for $T \to \infty$ with a zero of multiplicity ℓ appearing ℓ times in the list.

Assume that f is a normalized newform in $S_2(N)$. For each prime number p, let α_p and β_p be the two roots of $T^2 - \lambda(p)T + p$ where $\lambda(p)$ is given in (1.2). Put

(3.2)
$$b_f(p^m) = \begin{cases} \lambda(p)^m, & \text{if } p|N; \\ \alpha_p^m + \beta_p^m, & \text{if } (p,N) = 1. \end{cases}$$

The following theorem is essentially obtained in Li [12], where the theorem is presented in the context of L-series of elliptic curves. For the convenience of readers, we give a proof of the following theorem here which is a minor modification of that given in [12].

Theorem 3.1. Assume that f is a normalized newform in $S_2(N)$. If $\tau_f(n)$ is given in (3.1), then we have

$$\tau_f(n) = n \left(\ln \frac{\sqrt{N}}{2\pi} - \gamma \right) - \sum_{j=1}^n \binom{n}{j} \frac{(-1)^{j-1}}{(j-1)!} \sum_{m=1}^\infty \frac{\Lambda(m)}{m^{3/2}} b_f(m) (\ln m)^{j-1} + n \left(-\frac{2}{3} + \sum_{l=1}^\infty \frac{3}{l(2l+3)} \right) + \sum_{j=2}^n \binom{n}{j} (-1)^j \sum_{l=1}^\infty \frac{1}{(l+1/2)^j}$$

for $n = 1, 2, \dots$, where the second term on the right side of the identity is interpreted as the limit

$$\lim_{X \to \infty} \sum_{j=1}^{n} \binom{n}{j} \frac{(-1)^{j-1}}{(j-1)!} \sum_{m < X} \frac{\Lambda(m)}{m^{3/2}} b_f(m) (\ln m)^{j-1}$$

which exists.

Lemma 3.2. (see [14]) Let F(x) be a function defined on \mathbb{R} such that

$$2F(x) = F(x+0) + F(x-0)$$

for all $x \in \mathbb{R}$, such that $F(x) \exp((\epsilon+1/2)|x|)$ is integrable and of bounded variation on \mathbb{R} for a constant $\epsilon > 0$, and such that (F(x) - F(0))/x is of bounded variation on \mathbb{R} . Then

$$\sum_{\rho} \Phi(\rho) = 2F(0) \ln \frac{\sqrt{N}}{2\pi} - \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n} b_f(n) [F(\ln n) + F(-\ln n)] - \int_0^{\infty} \left(\frac{F(x) + F(-x)}{e^x - 1} - 2F(0) \frac{e^{-x}}{x} \right) dx,$$

where the sum on ρ runs over all zeros of $\xi_f(s)$ in the order given by $|\Im \rho| < T$ for $T \to \infty$, and

$$\Phi(s) = \int_{-\infty}^{\infty} F(x)e^{(s-1/2)x}dx.$$

Lemma 3.3. ([5] [6] [15]) Let f be a newform of weight 2 for $\Gamma_0(N)$. Then there an absolute effective constant c > 0 such that $L_f(s)$ has no zeros in the region

$$\{s = \sigma + it : \sigma \ge 1 - \frac{c}{\ln(N+1+|t|)}\},\$$

where $L_f(s)$ is given in (1.2).

Lemma 3.4. (Lemma 2 of [2]) For $n = 1, 2, \dots$, let

$$F_n(x) = \begin{cases} e^{x/2} \sum_{j=1}^n \binom{n}{j} \frac{x^{j-1}}{(j-1)!}, & if -\infty < x < 0; \\ n/2, & if x = 0; \\ 0, & if 0 < x. \end{cases}$$

Then

$$\Phi_n(s) = 1 - \left(1 - \frac{1}{s}\right)^n$$

where Φ_n is related to F_n by the relation

$$\Phi_n(s) = \int_{-\infty}^{\infty} F_n(x)e^{(s-1/2)x}dx.$$

Proof of Theorem 3.1. Since $\xi_f(s)$ is an entire function of order one and satisfies the functional identity $\xi_f(s) = w\xi_f(1-s)$, we have

$$\xi_f(s) = w\xi_f(1) \prod_{\rho} (1 - s/\rho)$$

where the product is over all the zeros of $\xi_f(s)$ in the order given by $|\Im \rho| < T$ for $T \to \infty$. If $\varphi_f(z) = \xi_f(1/(1-z))$, then

(3.3)
$$\frac{\varphi_f'(z)}{\varphi_f(z)} = \sum_{n=0}^{\infty} \tau_f(n+1)z^n$$

where the coefficients $\tau_f(n)$ are given in (3.1).

For a sufficiently large positive number X that is not an integer, let

$$F_{n,X}(x) = \begin{cases} F_n(x), & \text{if } -\ln X < x < \infty; \\ \frac{1}{2}F_n(-\ln X), & \text{if } x = -\ln X; \\ 0, & \text{if } -\infty < x < -\ln X \end{cases}$$

where $F_n(x)$ is given in Lemma 3.4. Then $F_{n,X}(x)$ satisfies all conditions of Lemma 3.2. Let

$$\Phi_{n,X}(s) = \int_{-\infty}^{\infty} F_{n,X}(x)e^{(s-1/2)x}dx.$$

By Lemma 3.2, we obtain that

$$\sum_{\rho} \Phi_{n,X}(\rho) = 2F_{n,X}(0) \ln \frac{\sqrt{N}}{2\pi} - \sum_{k=1}^{\infty} \frac{\Lambda(k)}{k} b_f(k) [F_{n,X}(\ln k) + F_{n,X}(-\ln k)] - \int_0^{\infty} \left(\frac{F_{n,X}(x) + F_{n,X}(-x)}{e^x - 1} - 2F_{n,X}(0) \frac{e^{-x}}{x} \right) dx,$$

where the sum on ρ runs over all zeros of $\xi_f(s)$ in the order given by $|\Im \rho| < T$ for $T \to \infty$. It follows that

$$\lim_{X \to \infty} \sum_{\rho} \Phi_{n,X}(\rho)$$

$$= n \left(\ln \frac{\sqrt{N}}{2\pi} - \gamma \right) - \lim_{X \to \infty} \sum_{j=1}^{n} \binom{n}{j} \frac{(-1)^{j-1}}{(j-1)!} \sum_{k < X} \frac{\Lambda(k)}{k^{3/2}} b_f(k) (\ln k)^{j-1}$$

$$+ n \left(-\frac{2}{3} + \sum_{l=1}^{\infty} \frac{3}{l(2l+3)} \right) + \sum_{j=2}^{n} \binom{n}{j} (-1)^j \sum_{l=1}^{\infty} \frac{1}{(l+1/2)^j}.$$

We have

(3.5)
$$\Phi_{n}(s) - \Phi_{n,X}(s) = X^{-s} \sum_{j=1}^{n} \binom{n}{j} (-1)^{j-1} \sum_{k=0}^{j-1} \frac{(\ln X)^{j-k-1}}{(j-k-1)!} s^{-k-1}$$

$$= \frac{X^{-s}}{s} \sum_{j=1}^{n} \binom{n}{j} \frac{(-\ln X)^{j-1}}{(j-1)!} + O\left(\frac{(\ln X)^{n-2}}{|s|^2} X^{-\Re s}\right).$$

Let ρ be any zero of $\xi_f(s)$. By Lemma 3.3, we have

$$\frac{c}{\ln(N+1+|\rho|)} \le \Re \rho \le 1 - \frac{c}{\ln(N+1+|\rho|)}$$

for a positive constant c. An argument similar to that made in the proof of (3.9) of [2] shows that

$$(3.6) \sum_{\rho} \frac{X^{-\Re \rho}}{|\rho|^2} \ll e^{-c'\sqrt{\ln X}}$$

for a positive constant c'.

Since

$$\sum_{\rho} \frac{X^{-\rho}}{\rho} = \sum_{\rho} \frac{X^{-(1-\rho)}}{1-\rho}$$

$$= -\frac{1}{X} \sum_{\rho} \frac{X^{\rho}}{\rho} + O\left(\sum_{\rho} \frac{X^{-(1-\Re\rho)}}{|\rho|^2}\right)$$

$$= -\frac{1}{X} \sum_{\rho} \frac{X^{\rho}}{\rho} + O\left(e^{-c'\sqrt{\ln X}}\right),$$

and since

$$\lim_{X \to \infty} \frac{(\ln X)^{j-1}}{X} \sum_{\rho} \frac{X^{\rho}}{\rho} = 0$$

for $j = 1, 2, \dots, n$ by Theorem 4.2 and Theorem 5.2 of [15], we have

(3.7)
$$\lim_{X \to \infty} (\ln X)^{j-1} \sum_{\rho} \frac{X^{-\rho}}{\rho} = 0$$

for $j=1,2,\cdots,n$. It follows from (3.5), (3.6) and (3.7) that

(3.8)
$$\lim_{X \to \infty} \sum_{\rho} \Phi_{n,X}(\rho) = \sum_{\rho} \Phi_n(\rho).$$

Since $\xi_f(s)$ is an entire function of order one, the series

$$\sum_{\rho} \Phi_n(\rho)$$

is convergent, where the sum on ρ runs over all zeros of $\xi_f(s)$ in the order given by $|\Im \rho| < T$ for $T \to \infty$. Hence, by (3.4) and (3.8) the limit

$$\lim_{X \to \infty} \sum_{j=1}^{n} {n \choose j} \frac{(-1)^{j-1}}{(j-1)!} \sum_{k < X} \frac{\Lambda(k)}{k^{3/2}} b_f(k) (\ln k)^{j-1}$$

exists.

This completes the proof of the theorem. \Box

4. Preliminary results

In this section, we collect some technical results for the proof of Theorem 1.2.

A fundamental result of Hecke asserts that a basis $\{f_1, f_2, \cdots, f_g\}$ in $S_2(N)$ exists which consists of eigenfunctions of all the Hecke operators T(n) with (n, N) = 1; see Theorem 6.21 in Iwaniec [8]. We can assume that each f_j is either a normalized newform in $S_2(N)$ or coming from a normalized newform in a lower level. For $j = 1, \cdots, g$, we choose $g_j = f_j$ if f_j is a normalized newform in $S_2(N)$, and $g_j = f'_j$ if f_j is an old form in $S_2(N)$ and if f'_j is a normalized newform in $S_2(N')$ for some divisor N' of N such that $f_j(z) = f'_j(dz)$ for some positive integer d|N/N'.

(4.1)
$$\xi_H(s) = \prod_{j=1}^g \xi_{g_j}(s)$$

where $\xi_{g_j}(s)$ is defined as in (1.3). Since $\xi_{g_j}(s)$ is an entire function and satisfies the functional identity $\xi_{g_j}(s) = w_j \xi_{g_j}(1-s)$, where $w_j = (-1)^{r_j}$ with r_j being the vanishing order of ξ_{g_j} at s = 1/2, the function $\xi_H(s)$ is entire and satisfies the functional identity

$$\xi_H(s) = \epsilon \xi_H(1-s),$$

where $\epsilon = \pm 1$. Put

10

(4.3)
$$\tau_H(n) = \sum_{i=1}^g \tau_{g_i}(n),$$

where $\tau_{g_j}(n)$ is defined similarly as in (3.1). If $\varphi_H(z) = \xi_H(1/(1-z))$, then we have

(4.4)
$$\frac{\varphi_H'(z)}{\varphi_H(z)} = \sum_{n=0}^{\infty} \tau_H(n+1)z^n$$

by (3.3).

Lemma 4.1. For all positive integers n with (n, N) = 1, we have

$$(T(n)f_i)(z) = \lambda_{q_i}(n)f_i(z)$$

for $j = 1, 2, \dots, g$, where $\lambda_{g_j}(n)$ is the eigenvalue of T(n) acting on $g_j(z)$.

Proof. If f_i is a newform, the stated identity is trivially true.

Next, we assume that f_j is an old form. Let g_j be a normalized newform in $S_2(N')$ for some divisor N' of N such that $f_j(z) = g_j(dz)$ for some positive integer d|N/N'. Since (n, N) = 1, by (1.1) we have

$$(T(n)f)(z) = \frac{1}{n} \sum_{ad-n} a^2 \sum_{0 \le b \le d} f\left(\frac{az+b}{d}\right)$$

for any function f in $S_2(N)$ or $S_2(N')$. Thus, we have

$$(T(n)f_j)(z) = \frac{1}{n} \sum_{\alpha \delta = n} \alpha^2 \sum_{0 \le \beta \le \delta} g_j \left(\frac{\alpha dz + d\beta}{\delta}\right).$$

Since (n, N) = 1, d|N and $\alpha \delta = n$, we have $(\delta, d) = 1$. Let r_{β} be remainder of $d\beta$ modulo δ . Then $\{r_{\beta} : 0 \leq \beta < \delta\} = \{0, 1, \dots, \delta - 1\}$. Since $g_j \in S_2(N')$, we have

$$g_j\left(\frac{\alpha dz + d\beta}{\delta}\right) = g_j\left(\frac{\alpha dz + r_\beta}{\delta}\right).$$

It follows that

$$(T(n)f_j)(z) = \frac{1}{n} \sum_{\alpha \delta = n} \alpha^2 \sum_{0 \le \beta < \delta} g_j \left(\frac{\alpha dz + \beta}{\delta}\right)$$
$$= (T(n)g_j)(w) = \lambda_{g_j}(n)g_j(w) = \lambda_{g_j}(n)f_j(z)$$

where w = dz.

This completes the proof of the lemma. \Box

Lemma 4.2. Let $\xi_N(s)$ be given in (1.5). Then $\xi_N(s)$ is an entire function, and its zeros in the strip $0 < \Re s < 1$ appear in pairs ρ and $1 - \rho$.

Proof. For $j = 1, 2, \dots, g$, if g_j is a normalized newform in $S_2(N_j)$, then we can write

(4.5)
$$L_{g_j}(s) = \prod_{p \mid N_j} (1 - \lambda_{g_j}(p)p^{-s})^{-1} \prod_{p \nmid N_j} (1 - \lambda_{g_j}(p)p^{-s} + p^{1-2s})^{-1}.$$

Since f_1, f_2, \dots, f_g are eigenfunctions of all the Hecke operators T(n) with (n, N) = 1, by Lemma 4.1 we have

$$\det |1 - T(p)p^{-s} + p^{1-2s}I| = \prod_{j=1}^{g} (1 - \lambda_{g_j}(p)p^{-s} + p^{1-2s})$$

for any prime $p \nmid N$. Let $L_N(s)$ be given in (1.4). Then we have

$$L_N(s) = \prod_{j=1}^g \prod_{p \nmid N} (1 - \lambda_{g_j}(p)p^{-s} + p^{1-2s})^{-1}$$

$$= \prod_{j=1}^g \left(L_{g_j}(s) \prod_{p \mid N_j} (1 - \lambda_{g_j}(p)p^{-s}) \prod_{p \nmid N_j, p \mid N} (1 - \lambda_{g_j}(p)p^{-s} + p^{1-2s}) \right).$$

Since

$$\xi_{g_j}(s) = N_j^{s/2} (2\pi)^{-s} \Gamma(\frac{1}{2} + s) L_{g_j}(\frac{1}{2} + s),$$

we have

$$\xi_H(s) = A^{s/2} N^{gs/2} (2\pi)^{-gs} \Gamma^g \left(\frac{1}{2} + s\right) \prod_{j=1}^g L_{g_j} \left(\frac{1}{2} + s\right)$$

where $A = N^{-g} \prod_{j=1}^{g} N_j$. It follows that (4.6)

$$\xi_N(s) = \xi_H(s) A^{-s/2} \prod_{j=1}^g \left(\prod_{p \mid N_j} (1 - \lambda_{g_j}(p) p^{-s-1/2}) \prod_{p \nmid N_j, p \mid N} (1 - \lambda_{g_j}(p) p^{-s-1/2} + p^{-2s}) \right).$$

This implies that $\xi_N(s)$ is an entire function. Since $|\lambda_{g_j}(p)| = \sqrt{p}$ for $p|N_j$ by ii) in Theorem 3 of Li [10] and since the two roots of the polynomial $1 - \lambda_{g_j}(p)p^{-1/2}z + z^2$ for $p \nmid N_j$ are conjugate complex numbers of absolute value one by the Ramanujan conjecture which was proved in Théorème 8.2 of Deligne [4], zeros of $\xi_N(s)$ in the critical strip $0 < \Re s < 1$ appear in pairs $\rho, 1 - \rho$ by (4.2) and (4.6).

This completes the proof of the lemma. \Box

Proof of Corollary 1.4. Let $\tau_N(n)$ be defined by (1.6) for all positive integers n. Since $\xi_H(s)$ is an entire function of order one, by (4.6) $\xi_N(s)$ is an entire function of order one. This implies that

$$\sum_{\rho} \frac{1 + |\Re \rho|}{(1 + |\rho|)^2} < \infty,$$

where the sum is over all zeros ρ of $\xi_N(s)$. Thus, conditions of Theorem 1.3 are satisfied. Since by (4.6) all zeros of $\xi_N(s)$ outside the strip $0 < \Re s < 1$ lie on the line $\Re s = 0$, Theorem 1.3 implies that all zeros of $\xi_N(s)$ in the critical strip $0 < \Re s < 1$ satisfy $\Re s \le 1/2$ if, and only if, $\tau_N(n) \ge 0$ for all positive integers n. By Lemma 4.2, all zeros of of $\xi_N(s)$ in the critical strip $0 < \Re s < 1$ appear in pairs ρ and $1 - \rho$. Thus, $\tau_N(n) \ge 0$ for all positive integers n if, and only if, $\Re \rho \le 1/2$ and $\Re (1 - \rho) \le 1/2$ for all zeros ρ of $\xi_N(s)$ in the critical strip $0 < \Re s < 1$. That is, all zeros of $\xi_N(s)$ in the strip $0 < \Re s < 1$ lie on the critical line $\Re s = 1/2$ if, and only if, $\tau_N(n) \ge 0$ for all positive integers n.

This completes the proof of the corollary. \Box

Lemma 4.3. Let p be a prime, and let α be a complex number of absolute value one. Then we have

$$1 - \alpha p^{-s} = c_p s^{\epsilon_p} \prod_{\rho} (1 - s/\rho)$$

where the product on ρ is over all nonzero zeros of $1-\alpha p^{-s}$ taken in the order given by $|\rho| < T$ for $T \to \infty$ and where $c_p = 1 - \alpha$, $\epsilon_p = 0$ if $\alpha \neq 1$ and $c_p = \ln p$, $\epsilon_p = 1$ if $\alpha = 1$.

Proof. Since $1-\alpha p^{-s}$ is an entire function of order one, by Hadamard's factorization theorem there is a constant a such that

(4.7)
$$1 - \alpha p^{-s} = c_p e^{as} s^{\epsilon_p} \prod_{\rho} (1 - s/\rho) e^{s/\rho}$$

where the product is over all nonzero zeros of $1-\alpha p^{-s}$. Let $\alpha=e^{it}$ with $0 \le t < 2\pi$. Then the zeros of $1-\alpha p^{-s}$ are $i(t+2k\pi)/\ln p, \ k=0,\pm 1,\pm 2,\cdots$. Since

$$\sum_{k=1}^{\infty} \left(\frac{-i \ln p}{t + 2k\pi} + \frac{-i \ln p}{t - 2k\pi} \right) = 2it \ln p \sum_{k=1}^{\infty} \frac{1}{(2k\pi)^2 - t^2}$$

is absolutely convergent, by using (4.7) we can write

(4.8)
$$1 - \alpha p^{-s} = c_p e^{hs} s^{\epsilon_p} \prod_{\rho} (1 - s/\rho)$$

for a constant h, where the product runs over all nonzero zeros ρ of $1 - \alpha p^{-s}$ taken in the order given by $|\rho| < T$ for $T \to \infty$. By taking logarithmic derivative of both sides of (4.8) with respect to s we get

(4.9)
$$\frac{\alpha \ln p}{p^s - \alpha} = h + \frac{\epsilon_p}{s} + \sum_{\rho} \frac{1}{s - \rho}.$$

By letting $s \to \infty$ in (4.9) we find that h = 0. Then the stated identity follows from (4.8).

This completes the proof of the lemma. \Box

Lemma 4.4. Let α , p be given in Lemma 4.3, and let $s = (1-z)^{-1}$. Then we have

$$\frac{d}{dz}\ln(1-\alpha p^{-s}) = \sum_{n=0}^{\infty} \left(\sum_{\rho} [1-(1-1/\rho)^{-n-1}]\right) z^n$$

for z in a small neighborhood of the origin, where the sum on ρ is over all zeros of $1 - \alpha \rho^{-s}$ taken in the order given by $|\rho| < T$ for $T \to \infty$.

Proof. By Lemma 4.3 we have

(4.10)
$$\frac{d}{dz}\ln(1-\alpha p^{-s}) = \frac{\epsilon_p}{1-z} + \sum_{\rho} \frac{1}{(1-\rho) + \rho z} \frac{1}{1-z}$$

where the sum on ρ is over all nonzero zeros of $1 - \alpha p^{-s}$. Since

$$\frac{1}{(1-\rho)+\rho z} \frac{1}{1-z} = \frac{1}{1-\rho} \left\{ \sum_{k=0}^{\infty} \left(\frac{-\rho}{1-\rho} \right)^k z^k \right\} \left\{ \sum_{l=0}^{\infty} z^l \right\}$$
$$= \sum_{n=0}^{\infty} \left[1 - (1-1/\rho)^{-n-1} \right] z^n$$

for z in a small neighborhood of the origin, the stated identity follows from (4.10). Note that, if $s = \rho = 0$ is a zero of $1 - \alpha p^{-s}$, then $(1 - 1/\rho)^{-n}$ is interpreted to be 0 for all positive integers n.

This completes the proof of the lemma. \Box

Remark 4.5. By (4.6), Lemma 4.4, (3.3) and (4.1) we have

$$\frac{d}{dz}\log\left(A^{s/2}\xi_N(s)\right) = \sum_{n=0}^{\infty} \tau_N(n+1)z^n$$

with $s = (1-z)^{-1}$, where the $\tau_N(n)$'s are given in (1.6). Then, by Lemma 4.2, to prove that all zeros of $\xi_N(s)$ in the strip $0 < \Re s < 1$ lie on the critical line $\Re s = 1/2$ it is enough to find an upper bound for each $\tau_N(n)$ which implies that the above series is analytic for |z| < 1.

Lemma 4.6. Let α , p be given in Lemma 4.3, and let $s = (1-z)^{-1}$. Then we have

$$\frac{d}{dz}\ln(1-\alpha p^{-s}) = \sum_{n=0}^{\infty} \left(\sum_{j=0}^{n} \binom{n+1}{j+1} \frac{(-1)^{j}}{j!} \sum_{k=1}^{\infty} \frac{\ln p}{p^{k}} \alpha^{k} (\ln p^{k})^{j}\right) z^{n}$$

for z in a small neighborhood of the origin.

Proof. Let m be any positive integer. By using mathematical induction on n, we can show that

$$(4.11) \quad \frac{d^n}{dz^n} \left[(1-z)^{-m} p^{-ks} \right]_{|z=0} = p^{-k} \sum_{j=0}^n \binom{n}{j} (n+m-1) \cdots (j+m) (-\ln p^k)^j$$

for $n = 1, 2, \dots$. By using the formula (4.11) with m = 2 we find that

$$\frac{d}{dz}\ln(1-\alpha p^{-s}) = (1-z)^{-2}\ln p \sum_{k=1}^{\infty} \alpha^k p^{-ks}
= \sum_{n=0}^{\infty} \frac{z^n}{n!} \left(\sum_{k=1}^{\infty} \alpha^k \frac{\ln p}{p^k} \sum_{j=0}^n \binom{n}{j} (n+1) \cdots (j+2) (-\ln p^k)^j \right)
= \sum_{n=0}^{\infty} z^n \left(\sum_{j=0}^n \binom{n+1}{j+1} \frac{(-1)^j}{j!} \sum_{k=1}^{\infty} \frac{\ln p}{p^k} \alpha^k (\ln p^k)^j \right).$$

This completes the proof of the lemma. \Box

Lemma 4.7. Let p be a prime, and let α be a complex number of absolute value one. Then we have

$$\sum_{\rho} [1 - (1 - 1/\rho)^{-n}] = \sum_{i=1}^{n} \binom{n}{j} \frac{(-1)^{j-1}}{(j-1)!} \sum_{k=1}^{\infty} \frac{\ln p}{p^k} \alpha^k (\ln p^k)^{j-1}$$

for $n=1,2,\cdots$, where the sum on ρ is over all zeros of $1-\alpha p^{-s}$ taken in the order given by $|\Im \rho| < T$ for $T \to \infty$ with a zero of multiplicity ℓ appearing ℓ times in the list.

Proof. The stated identity follows from Lemma 4.4 and Lemma 4.6. \Box

Lemma 4.8. For $n = 1, 2, \dots$, we have

$$\tau_N(n) = \sum_{j=1}^g \tau_{g_j}(n) + \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{(m,N)>1,m=1}^\infty \frac{\Lambda(m)}{m^{3/2}} (\sum_{j=1}^g b_{g_j}(m)) (\ln m)^{l-1}$$

where $b_{q_i}(m)$ is given as in (3.2).

Proof. Let $\tau_H(n)$ be given in (4.3). By (3.1), (4.1) and (4.6) we have (4.12)

$$\tau_N(n) = \tau_H(n) + \sum_{j=1}^g \{ \sum_{p|N_j} \sum_{\rho_j} [1 - (1 - 1/\rho_j)^{-n}] + \sum_{p \nmid N_j, p|N} \sum_{\alpha_j} [1 - (1 - 1/\alpha_j)^{-n}] \},$$

where the sum on ρ_j is over all zeros of $1 - \lambda_{g_j} p^{-s-1/2}$ with $p|N_j$ and where the sum on α_j is over all zeros of $1 - \lambda_{g_j}(p)p^{-s-1/2} + p^{-2s}$ with $p \nmid N_j, p|N$. By Lemma 4.7 we have

(4.13)
$$\sum_{\rho_j} \left[1 - (1 - 1/\rho_j)^{-n}\right] = \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{k=1}^\infty \frac{\ln p}{p^{3k/2}} b_{g_j}(p^k) (\ln p^k)^{l-1}$$

and

(4.14)
$$\sum_{\alpha_j} \left[1 - (1 - 1/\alpha_j)^{-n}\right] = \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{k=1}^\infty \frac{\ln p}{p^{3k/2}} b_{g_j}(p^k) (\ln p^k)^{l-1}.$$

It follows from (4.13) and (4.14) that

(4.15)
$$\sum_{p|N_j} \sum_{\rho_j} [1 - (1 - 1/\rho_j)^{-n}] + \sum_{p \nmid N_j, p|N} \sum_{\alpha_j} [1 - (1 - 1/\alpha_j)^{-n}]$$

$$= \sum_{p|N} \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{k=1}^\infty \frac{\ln p}{p^{3k/2}} b_{g_j}(p^k) (\ln p^k)^{l-1}.$$

By (4.15) we have

$$\sum_{j=1}^{g} \left(\sum_{p|N_j} \sum_{\rho_j} [1 - (1 - 1/\rho_j)^{-n}] + \sum_{p \nmid N_j, p|N} \sum_{\alpha_j} [1 - (1 - 1/\alpha_j)^{-n}] \right)$$

$$= \sum_{l=1}^{n} \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{(m,N)>1,m=1}^{\infty} \frac{\Lambda(m)}{m^{3/2}} (\sum_{j=1}^{g} b_{g_j}(m)) (\ln m)^{l-1}.$$

The stated identity then follows from (4.12).

This completes the proof of the lemma. \Box

5. Proof of Theorem 1.2

In this section we complete the proof of Theorem 1.2. We define an operator S acting on the space $S_2(N)$ by

(5.1)
$$S(1) = 2I$$

$$S(p) = T(p)$$

$$S(p^m) = T(p^m) - pT(p^{m-2})$$

for $m = 2, 3, \dots$; see Ihara [7].

Lemma 5.1. For each prime $p \nmid N$, the trace $tr(S(p^m))$ of $S(p^m)$ acting on the space $S_2(N)$ is given by

(5.2)
$$tr(S(p^m)) = \sum_{j=1}^{g} b_{g_j}(p^m)$$

for $m = 0, 1, 2, \cdots$.

Proof. Let p be any prime. It follows from the recursion formula (see (6.25) in Iwaniec [8])

$$T(p^{m+1}) = T(p)T(p^m) - pT(p^{m-1})$$

that

(5.3)
$$S(p^{m+1}) = S(p)S(p^m) - pS(p^{m-1})$$

for $m = 1, 2, \cdots$.

Let p is any prime not dividing N. When m=0, we have

$$S(p^m)f_i = 2f_i = b_{a_i}(p^m)f_i$$
.

When m = 1, we have

$$S(p^m)f_j = T(p)f_j = \lambda_{g_i}(p)f_j = b_{g_i}(p^m)f_j$$

by Lemma 4.1. Assume that

$$S(p^m)f_i = b_{q_i}(p^m)f_i$$

for all integers $m \leq k$. When m = k + 1, we have

$$S(p^{m})f_{j} = (S(p)S(p^{k}) - pS(p^{k-1}))f_{j}$$

$$= (b_{g_{j}}(p)b_{g_{j}}(p^{k}) - pb_{g_{j}}(p^{k-1}))f_{j}$$

$$= b_{g_{j}}(p^{m})f_{j}.$$

By mathematical induction the identity

$$S(p^m)f_j = b_{g_j}(p^m)f_j$$

holds for all nonnegative integers m. Since $\{f_1, \dots, f_g\}$ is a basis for $S_2(N)$, we have

$$tr(S(p^m)) = \sum_{j=1}^{g} b_{g_j}(p^m)$$

for $m = 0, 1, 2, \cdots$.

This completes the proof of the lemma. \Box

From Lemma 5.1 and the definition (5.1) we obtain the following corollary.

Corollary 5.2. For each prime $p \nmid N$, we have

$$\sum_{j=1}^{g} b_{g_j}(p^m) = tr(T(p^m)) - p \, tr(T(p^{m-2}))$$

for $m=0,1,2,\cdots$.

Proof of Theorem 1.2. By Lemma 4.8 we have (5.4)

$$\tau_N(n) = \sum_{j=1}^g \tau_{g_j}(n) + \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{(m,N)>1,m=1}^\infty \frac{\Lambda(m)}{m^{3/2}} (\sum_{j=1}^g b_{g_j}(m)) (\ln m)^{l-1}$$

for $n = 1, 2, \dots$. Since g_j is a normalized newform in $S_2(N_j)$, by Theorem 3.1 we have

(5.5)
$$\tau_{g_j}(n) = n \left(\ln \frac{\sqrt{N_j}}{2\pi} - \gamma \right) - \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{m=1}^\infty \frac{\Lambda(m)}{m^{3/2}} b_{g_j}(m) (\ln m)^{l-1} + n \left(-\frac{2}{3} + \sum_{l=1}^\infty \frac{3}{l(2l+3)} \right) + \sum_{m=2}^n \binom{n}{m} (-1)^m \sum_{l=1}^\infty \frac{1}{(l+1/2)^m}$$

By using (5.4) and (5.5) we obtain that (5.6)

$$\tau_N(n) = \frac{n}{2} \ln(N_1 \cdots N_g) - \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{\substack{m=1\\(m,N)=1}}^{\infty} \frac{\Lambda(m)}{m^{3/2}} \{ \sum_{j=1}^g b_{g_j}(m) \} (\ln m)^{l-1} - ng \left(\ln 2\pi + \gamma + \frac{2}{3} - \sum_{l=1}^{\infty} \frac{3}{l(2l+3)} \right) + g \sum_{m=2}^n \binom{n}{m} \sum_{l=1}^{\infty} \frac{(-1)^m}{(l+1/2)^m}.$$

By Theorem 5 of Atkin and Lehner [1], we have

(5.7)
$$\prod_{N_j \neq N, 1 \leq j \leq g} N_j = \prod_{1 < m < N, m \mid N} m^{\nu_m d(N/m)}.$$

Since the dimension of the space $S_2(N)$ is g, which is explicitly given in Proposition 1.40 and Proposition 1.43 of Shimura [17], we have the following recurrence formula for the number of newforms in a basis for $S_2(N)$:

(5.8)
$$\nu_N = g - \sum_{1 < m < N, m \mid N} \nu_m d(N/m).$$

Thus, we have

(5.9)
$$\prod_{N_j = N, 1 \le j \le g} N_j = N^{\nu_N}.$$

By (5.6), (5.7) and (5.9) we have (5.10)

$$\tau_N(n) = \frac{n}{2} \ln \left(N^{\nu_N} \prod_{1 < m < N, m \mid N} m^{\nu_m d(N/m)} \right)$$

$$- \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{\substack{m=1 \ (m,N)=1}}^{\infty} \frac{\Lambda(m)}{m^{3/2}} \{ \sum_{j=1}^g b_{g_j}(m) \} (\ln m)^{l-1}$$

$$- ng \left(\ln 2\pi + \gamma + \frac{2}{3} - \sum_{l=1}^{\infty} \frac{3}{l(2l+3)} \right) + g \sum_{m=2}^n \binom{n}{m} \sum_{l=1}^{\infty} \frac{(-1)^m}{(l+1/2)^m}.$$

Since

$$\ln 2 = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n},$$

we have

$$\ln 2 = \frac{4}{3} - \frac{3}{2} \sum_{l=1}^{\infty} \frac{1}{l(2l+3)}.$$

It follows that

(5.11)
$$\ln 2\pi + \gamma + \frac{2}{3} - \sum_{l=1}^{\infty} \frac{3}{l(2l+3)} = \ln(8\pi) + \gamma - 2.$$

Note that the identity (5.11) is due to B. Conrey. The author wants to thank him for this observation. By (5.10) and (5.11) we have

$$\tau_N(n) = \frac{n}{2} \ln \left(N^{\nu_N} \prod_{1 < m < N, m \mid N} m^{\nu_m d(N/m)} \right)$$
$$- \sum_{l=1}^n \binom{n}{l} \frac{(-1)^{l-1}}{(l-1)!} \sum_{\substack{m=1 \ (m,N)=1}}^{\infty} \frac{\Lambda(m)}{m^{3/2}} \{ \sum_{j=1}^g b_{g_j}(m) \} (\ln m)^{l-1}$$
$$- ng \left(\ln(8\pi) + \gamma - 2 \right) + g \sum_{m=2}^n \binom{n}{m} \sum_{l=1}^{\infty} \frac{(-1)^m}{(l+1/2)^m}.$$

The stated identity then follows from Corollary 5.2.

This completes the proof of the theorem.

The following result for the size of the last term on the right side of the identity in Theorem 1.2 is due to B. Conrey. The author wants to thank him for allowing him to include it here.

Lemma 5.3. (B. Conrey) We have

$$\sum_{m=2}^{n} \binom{n}{m} \sum_{l=1}^{\infty} \frac{(-1)^m}{(l+1/2)^m} = n \ln n + O(n)$$

for all positive integers n.

Proof. We have

(5.12)
$$\sum_{m=2}^{n} \binom{n}{m} \sum_{l=1}^{n} \frac{(-1)^m}{(l+1/2)^m} = \sum_{l=1}^{n} \left[\left(1 - \frac{1}{l+1/2} \right)^n - 1 + \frac{n}{l+1/2} \right]$$
$$= O(n) + n \sum_{l=1}^{n} \frac{1}{l+1/2}$$
$$= O(n) + n \ln n.$$

Since

$$\sum_{l=n+1}^{\infty} \frac{1}{(l+1/2)^m} < \int_n^{\infty} \frac{1}{(t+1/2)^m} dt = \frac{1}{m-1} (n+1/2)^{1-m} < n^{1-m},$$

we have

(5.13)
$$\left| \sum_{m=2}^{n} \binom{n}{m} \sum_{l=n+1}^{\infty} \frac{(-1)^m}{(l+1/2)^m} \right| \le n(1+1/n)^n = O(n).$$

The stated identity then follows from (5.12) and (5.13).

This completes the proof of the lemma. \Box

Remark 5.4. It follows from Lemma 5.3 that the last term on right side of the identity in Theorem 1.2 is asymptotically equal to $gn \ln n + O(n)$ as $n \to \infty$. According to a recent result of Lagarias [9], we would have

$$\lim_{n \to \infty} \frac{\tau_N(n)}{gn \ln n} = 1$$

if all zeros of $\xi_N(s)$ in the strip $0 < \Re s < 1$ lie on the critical line $\Re s = 1/2$.

Remark 5.5. The function $L_N(s)$ in (1.4) is a partial L-function defined only for $p \nmid N$. A question of the referee, which author does not know, is how to define Euler factors for p|N so that the function $\xi_N(s)$ corresponding to a completed L-function $L_N(s)$ satisfies a functional equation of automorphic L-functions.

References

- 1. A. O. L. Atkin and J. Lehner, Hecke operators on $\Gamma_0(m)$, Math. Ann. 185 (1970), 134–160.
- E. Bombieri and J. C. Lagarias, Complements to Li's criterion for the Riemann hypothesis,
 J. Number Theory 77 (1999), 274–287.
- 3. H. Cohen, Trace des opérateurs de Hecke sur $\Gamma_0(N)$, Séminaire de Théorie des Nombres (1976–1977), Exp. No. 4, 5 novembre 1976, Lab. Théorie des Nombres, CNRS, Talence, 1977.
- 4. P. Deligne, La Conjecture de Weil. I, IHES Publ. Math. 43 (1974), 273–307.
- 5. S. S. Gelbart, Automorphic Forms on Adele Groups, Princeton University Press, 1975.
- J. Hoffstein and D. Ramakrishnan, Siegel zeros and cusp forms, Internat. Math. Res. Notices 6 (1995), 279–308.
- Y. Ihara, Hecke polynomials as congruence ζ functions in elliptic modular case, Ann. of Math. 85 (1967), 267–295.
- 8. H. Iwaniec, Topics in Classical Automorphic Forms, Amer. Math. Soc., Providence, RI, 1997.
- 9. J. C. Lagarias, On Li's coefficients for automorphic L-functions, Preprint (2004).
- 10. Wen-Ch'ing W. Li, Newforms and functional equations, Math. Ann. 212 (1975), 285-315.
- Xian-Jin Li, The positivity of a sequence of numbers and the Riemann hypothesis, J. Number Theory 65 (1997), 325–333.
- 12. Xian-Jin Li, Explicit formulas for Dirichlet and Hecke L-functions, Illinois J. Math. (to appear).
- 13. Xian-Jin Li, An explicit formula for Hecke L-functions, Preprint (2004).
- J.-F. Mestre, Formules explictes et minorations de conducteurs de variétés algébriques, Compositio Math. 58 (1986), 209–232.
- C. J. Moreno, Explicit formulas in the theory of automorphic forms, Number Theory Day, Lecture Notes in Math., no. 626, Springer, Berlin, 1977, 73–216.
- J. Oesterlé, Sur la trace des opérateurs de Hecke, Thèse de Docteur 3è cycle, Université de Paris-Sud, Centre d'Orsay, 1977.
- G. Shimura, Introduction to the Arithmetic Theory of Automorphic Functions, Princeton University Press, Princeton, NJ, 1994.

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UTAH 84602 E-mail address: xianjin@math.byu.edu