Guillermo Moreno

Abstract: In this paper, we study the zero set of the Hopf construction map $F_n : \mathbb{A}_n \times \mathbb{A}_n \to \mathbb{A}_n \times \mathbb{A}_0$ given by $F_n(x, y) = (2xy, ||y||^2 - ||x||^2)$ for $n \ge 4$, where \mathbb{A}_n is the Cayley-Dickson algebra of dimension 2^n on \mathbb{R} .

Introduction: Let $f_1 : S^3 \to S^2, f_2 : S^7 \to S^4$ and $f_3 : S^{15} \to S^8$ be the classical **Hopf maps**; these can be defined using **the Hopf construction**: Let $\mathbb{A}_1 = \mathbb{C}, \mathbb{A}_2 = \mathbb{H}$ and $\mathbb{A}_3 = \emptyset$ be the complex, quaternion and octonion numbers respectively and $F_n : \mathbb{A}_n \times \mathbb{A}_n \to \mathbb{A}_n \times \mathbb{R}$ are given by

$$F_n(x,y) = (2xy, ||y||^2 - ||x||^2)$$

for n = 1, 2, 3.

Now write $S^{2^{n+1}-1} = \{(x, y) \in \mathbb{A}_n \times \mathbb{A}_n : ||x||^2 + ||y||^2 = 1\}$. By definition,

$$F_n | S^{2^{n+1}-1} = f_r$$

are the Hopf maps. Since \mathbb{A}_n is a normed real algebra of dimension 2^n , for n = 1, 2, 3 we have that

$$\begin{aligned} ||(2xy, ||y||^2 - ||x||^2)||^2 &= 4||xy||^2 + (||y||^2 - ||x||^2)^2 \\ &= 4||x||^2||y||^2 + ||y||^4 + ||x||^4 - 2||x||^2||y||^2 \\ &= (||x||^2 + ||y||^2)^2, \end{aligned}$$

so if $||x||^2 + ||y||^2 = 1$, then $||F_n(x, y)|| = ||(2xy, ||y||^2 - ||x||^2)|| = 1$. Now by the Cayley-Dickson doubling process ([D]) define

$$\mathbb{A}_{n+1} = \mathbb{A}_n \times \mathbb{A}_n$$

Keywords and phrases: Cayley-Dickson algebras, alternative algebras, zero divisors, flexible algebra,normed algebra.

²⁰⁰⁰ Mathematics Subject Classification: 17A99, 55Q25.

with

$$(a,b)(x,y) = (ax - \overline{y}b, ya + b\overline{x})$$
 for a, b, x and y in \mathbb{A}_n

and

$$\overline{x} = (\overline{x}_1, -x_2)$$
 if $x = (x_1, x_2)$ in $\mathbb{A}_{n-1} \times \mathbb{A}_{n-1}$.

Thus, if $\mathbb{A}_0 = \mathbb{R}$ with $\overline{x} = x$ for x real number, then $\mathbb{A}_1 = \mathbb{C}, \mathbb{A}_2 = \mathbb{H}$ and $\mathbb{A}_3 = \emptyset$, which are normed algebras i.e.; ||xy|| = ||x||||y|| for all x, y in \mathbb{A}_n .

For $n \ge 4$, \mathbb{A}_n is no longer normed and also \mathbb{A}_n has zero divisors (see [K-Y] and [Mo₁]).

Let us define $X_n^{\infty} = \{(x, y) \in \mathbb{A}_n \times \mathbb{A}_n | F_n(x, y) = (0, 0)\}$ and for r nonnegative real number $(x, y) \in X_n^r$ if and only if xy = 0 and ||x|| = ||y|| = r. It is clear that for r > 0 and s > 0 real numbers. X_n^r is homeomorphic to X_n^s . Let us define $X_n := X_n^1$.

The set X_n show up in some important problems in algebraic topology: (1) Cohen's approach to the Arf invariant one problem. (See $[C_1]$ and $[C_2]$). (2) Adem-Lam construction of normed and non-singular bilinear maps.(See [A] and [L]).

In this paper, we will show that for $n \ge 4$, X_n is related to some Stiefel manifolds; using the algebra structure in \mathbb{A}_{n+1} we will construct the chain of inclusions

$$X_n \subset W_{2^{n-1}-1,2} \subset V_{2^n-2,2} \subset V_{2^n-1,2}$$

(see § 2 below) where $V_{m,2}$ and $W_{m,2}$ denote the real and complex Stiefel manifolds of 2-frames in \mathbb{R}^m and \mathbb{C}^m respectively.

In § 3 we show that we can attach to every element in $W_{2^{n-1},2}$ in a canonical way, an eight dimensional vector subspace of \mathbb{A}_{n+1} and that, only for the elements in X_n , such vector subspace is isomorphic, as algebra, to $\mathbb{A}_3 = \emptyset$ (the octonions).

In § 4 we describe X_n as a certain type of algebra monomorphisms from $\mathbb{A}_3 = \emptyset$ to \mathbb{A}_{n+1} for $n \geq 4$.

This paper is a sequel of [Mo1] and we use freely the results of [Sch], we acknowledge with gratitude the hard work made by the reviwer.

§1. Pure and doubly pure elements in \mathbb{A}_{n+1} .

Throughout this paper we use the following notational conventions:

(1) Elements in \mathbb{A}_n will be denoted by Latin characters a, b, c, \ldots, x, y, z . and elements in \mathbb{A}_{n+1} will be denoted by Greek characters $\alpha, \beta, \gamma, \ldots$. For example,

$$\alpha = (a, b) \in \mathbb{A}_n \times \mathbb{A}_n.$$

(2) When we need to represent elements in \mathbb{A}_n as elements in $\mathbb{A}_{n-1} \times \mathbb{A}_{n-1}$ we use subscripts, for instance, $a = (a_1, a_2)$, $b = (b_1, b_2)$, and so on, with a_1, a_2, b_1, b_2 in \mathbb{A}_{n-1} .

Now $\{e_0, e_1, \ldots, e_{2^n-1}\}$ denotes the canonical basis in \mathbb{A}_n . Then by the doubling process

$$\{(e_0, 0), (e_1, 0), \dots, (e_{2^n - 1}, 0), (0, e_0), \dots, (0, e_{2^n - 1})\}$$

is the canonical basis in $\mathbb{A}_{n+1} = \mathbb{A}_n \times \mathbb{A}_n$. By standard abuse of notation, we also denote $e_0 = (e_0, 0), e_1 = (e_1, 0), \dots, e_{2^n-1} = (e_{2^n-1}, 0), e_{2^n} = (0, e_0), \dots, e_{2^{n+1}-1} = (0, e_{2^n-1})$ in \mathbb{A}_{n+1} .

For $\alpha = (a, b) \in \mathbb{A}_n \times \mathbb{A}_n = \mathbb{A}_{n+1}$ we denote $\widetilde{\alpha} = (-b, a)$ (the complexification of α) so $\widetilde{e}_0 = (0, e_0)$ and $\alpha \widetilde{e}_0 = (a, b)(0, e_0) = (-b, a) = \widetilde{\alpha}$. Notice that $\widetilde{\alpha} = -\alpha$.

The **trace** on \mathbb{A}_{n+1} is the linear map $t_{n+1} : \mathbb{A}_{n+1} \to \mathbb{R}$ given by $t_{n+1}(\alpha) = \alpha + \overline{\alpha} = 2$ (real part of α) so $t_{n+1}(\alpha) = t_n(\alpha)$ when $\alpha = (a, b) \in \mathbb{A}_n \times \mathbb{A}_n$.

Definition: $\alpha = (a, b)$ in \mathbb{A}_{n+1} is pure if

$$t_{n+1}(\alpha) = t_n(a) = 0.$$

 $\alpha = (a, b)$ in \mathbb{A}_{n+1} is doubly pure if it is pure and also $t_n(b) = 0$; i.e., $\widetilde{\alpha}$ is pure in \mathbb{A}_{n+1} .

Also $2\langle a, b \rangle = t_n(a\overline{b})$ when $\langle -, - \rangle$ is the inner product in \mathbb{R}^{2^n} (see [A]). Note that for a and b pure elements $a \perp b$ if and only if ab = -ba.

Notation: ${}_{o}\mathbb{A}_{n} = \{e_{o}\}^{\perp} \subset \mathbb{A}_{n}$ is the vector subspace consisting of pure elements in \mathbb{A}_{n} ; i.e., ${}_{o}\mathbb{A}_{n} = \operatorname{Ker}(t_{n}) = \mathbb{R}^{2^{n-1}}$. $\widetilde{\mathbb{A}}_{n+1} = {}_{o}\mathbb{A}_{n} \times {}_{o}\mathbb{A}_{n} = \{e_{0}, \widetilde{e}_{0}\}^{\perp} = \mathbb{R}^{2^{n+1}-2}$ is the vector subspace consisting

 $\mathbb{A}_{n+1} = {}_{o}\mathbb{A}_n \times {}_{o}\mathbb{A}_n = \{e_0, \widetilde{e}_0\}^{\perp} = \mathbb{R}^{2^{n+1}-2}$ is the vector subspace consisting of doubly pure elements in \mathbb{A}_{n+1} .

Lemma 1.1. For a and b in $\widetilde{\mathbb{A}_n}$ we have that

1) $a\tilde{e}_0 = \tilde{a}$ and $\tilde{e}_0 a = -\tilde{a}$.

- 2) $a\widetilde{a} = -||a||^2 \widetilde{e}_0$ and $\widetilde{a}a = ||a||^2 \widetilde{e}_0$ so $a \perp \widetilde{a}$.
- 3) $\widetilde{a}b = -\widetilde{a}b$ with *a* a pure element.
- 4) $a \perp b$ if and only if $\tilde{a}b + \tilde{b}a = 0$.
- 5) $\widetilde{a} \perp b$ if and only if $ab = \widetilde{b}\widetilde{a}$.
- 6) $a \perp b$ and $\tilde{a} \perp b$ if and only if $\tilde{a}b = a\tilde{b}$.

Proof: Note that *a* is pure if and only if $\overline{a} = -a$ and if $a = (a_1, a_2)$ is doubly pure, then $\overline{a}_1 = -a_1$ and $\overline{a}_2 = -a_2$.

- 1) $\tilde{e}_0 a = (0, e_0)(a_1, a_2) = (-\overline{a}_2, \overline{a}_1) = (a_2, -a_1) = -(-a_2, a_1) = -\widetilde{a}.$
- 2) $a\tilde{a} = (a_1, a_2)(-a_2, a_1) = (-a_1a_2 + a_1a_2, a_1^2 + a_2^2) = (0, -||a||^2 e_0) = -||a||^2 \tilde{e}_0.$ Similarly $\tilde{a}a = (-a_2, a_1)(a_1, a_2) = (-a_2a_1 + a_2a_1, -a_2^2 - a_1^2) = ||a||^2 \tilde{e}_0.$

Now, since $-2\langle \widetilde{a}, a \rangle = a\widetilde{a} + \widetilde{a}a = 0$ we have $a \perp \widetilde{a}$.

3) $\widetilde{ab} = (-a_2, a_1)(b_1, b_2) = (-a_2b_1 + b_2a_1, -b_2a_2 - a_1b_1).$ So $\widetilde{ab} = (a_1b_1 + b_2a_2, b_2a_1 - a_2b_1) = (a_1, a_2)(b_1, b_2) = ab$ and then $-\widetilde{ab} = \widetilde{ab}.$

Notice that in this proof we only use that $\overline{a}_1 = -a_1$; i.e., a is pure and b doubly pure.

- 4) $a \perp b \Leftrightarrow ab + ba = 0 \Leftrightarrow ab = -ba \Leftrightarrow \widetilde{ab} = -\widetilde{ba}.$ $\Leftrightarrow -\widetilde{ab} = \widetilde{b}a \Leftrightarrow \widetilde{ab} + \widetilde{b}a = 0$ by (3).
- 5) $\widetilde{a} \perp b \Leftrightarrow \widetilde{\widetilde{a}}b + \widetilde{b}\widetilde{a} = 0$ (by (4)) $\Leftrightarrow -ab + \widetilde{b}\widetilde{a} = 0$.
- 6) If $\widetilde{a} \perp b$ and $a \perp b$, then by (3) and (4) $\widetilde{a}b = -\widetilde{a}b = b\widetilde{a} = -\widetilde{b}a = a\widetilde{b}$. Conversely, put $a = (a_1, a_2)$ and $b = (b_1, b_2)$ in $\mathbb{A}_{n-1} \times \mathbb{A}_{n-1}$ and define $c := (a_1b_1 + b_2a_2)$ and $d := (b_2a_1 - a_2b_1)$ in \mathbb{A}_{n-1} . Then $a\widetilde{b} = (a_1, a_2)(-b_2, b_1) = (-a_1b_2 + b_1a_2, b_1a_1 + a_2b_2)$ so $a\widetilde{b} = (-\overline{d}, \overline{c})$.

Now $ab = (a_1, a_2)(b_1, b_2) = (a_1b_1 + b_2a_2, b_2a_1 - a_2b_1) = (c, d)$, so $\widetilde{ab} = (-d, c)$ and then $\widetilde{ab} = (d, -c)$.

Thus, if $a\tilde{b} = \tilde{a}b$ then $\overline{c} = -c$ and $d = -\overline{d}$. Then

$$t_n(ab) = t_{n-1}(c) = c + \overline{c} = 0 \text{ and } a \perp b$$

 $t_n(\widetilde{a}b) = t_{n-1}(d) = d + \overline{d} = 0 \text{ and } \widetilde{a} \perp b.$
Q.E.D

Corollary 1.2 For each $a \neq 0$ in $\widetilde{\mathbb{A}}_n$ the four dimensional vector subspace generated by $\{e_0, \widetilde{a}, a, \widetilde{e}_0\}$ is a copy of $\mathbb{A}_2 = \mathbb{H}$ we denote it by \mathbb{H}_a .

Proof: We suppose that ||a|| = 1, otherwise we take $\frac{a}{||a||}$. Construct the following multiplication table.

	e_0	\widetilde{a}	a	\widetilde{e}_0
e_0 \widetilde{a}	e_0	\widetilde{a}	a	\widetilde{e}_0
\widetilde{a}	\widetilde{a}	$-e_0$	$+\widetilde{e}_0$	-a
a	a	$-\widetilde{e}_0$	$-e_0$	\widetilde{a}
$a \\ \widetilde{e}_0$	\widetilde{e}_0	a	$-\widetilde{a}$	$-e_0$

By lemma 2.1. $a\widetilde{e}_0 = \widetilde{a}; \ \widetilde{e}_0 a = -\widetilde{a}; \ \widetilde{a}\widetilde{e}_0 = \widetilde{\widetilde{a}} = -a; \ \widetilde{e}_0\widetilde{a} = -\widetilde{\widetilde{a}} = a;$ $a\widetilde{a} = -\widetilde{e}_0 \text{ and } \widetilde{a}a = \widetilde{e}_0.$

Identifying $e_0 \leftrightarrow e_0$, $\tilde{a} \leftrightarrow e_1$, $a \leftrightarrow e_2$ and $\tilde{e}_0 \leftrightarrow e_3$ we have the multiplication table for $\mathbb{A}_2 = \mathbb{H}$

Q.E.D.

§2.- The Stiefel manifold $V_{2^n-1,2}$ in \mathbb{A}_{n+1} and a T^2 -action.

Let $\langle a, b \rangle_n$ denote the standard inner product of a and b in $\mathbb{A}_n = \mathbb{R}^{2^n}$. Now by [A] and [Mo₁]

$$2\langle a,b\rangle_n = (ab+b\overline{a}) = t_n(ab).$$

It is also well known that, for $\alpha = (a, b)$ and $\chi = (x, y)$ in $\mathbb{A}_n \times \mathbb{A}_n = \mathbb{A}_{n+1}$ we have

$$\langle \alpha, \chi \rangle_{n+1} = \langle a, x \rangle_n + \langle b, \overline{y} \rangle_n$$

In particular, if α and χ are doubly pure elements in \mathbb{A}_{n+1} then y and b are pure elements in \mathbb{A}_n , therefore

$$\langle \alpha, \chi \rangle_{n+1} = \langle a, x \rangle_n + \langle b, y \rangle_n.$$

Lemma 2.1 For $\alpha = (a, b)$ in \mathbb{A}_{n+1} define $\hat{\alpha} := (b, a)$.

For $\alpha \in \mathbb{A}_{n+1}$ we have that

i) $\langle \alpha, \hat{\alpha} \rangle_{n+1} = 0$ i.e. $\alpha \perp \hat{\alpha}$ in \mathbb{A}_{n+1} if and only if $\langle a, b \rangle_n = 0$, i.e. $a \perp b$ in \mathbb{A}_n .

ii) $\langle \tilde{\alpha}, \hat{\alpha} \rangle_{n+1} = 0$, i.e. $\tilde{\alpha} \perp \hat{\alpha}$ in \mathbb{A}_{n+1} if and only if ||a|| = ||b|| in \mathbb{A}_n . **Proof:**

i)
$$\langle \alpha, \hat{\alpha} \rangle_{n+1} = \langle (a, b), (b, a) \rangle_{n+1} = 2 \langle a, b \rangle_n.$$

ii) $\langle \widetilde{\alpha}, \hat{\alpha} \rangle_{n+1} = \langle (-b, a), (b, a) \rangle_{n+1} = -\langle b, b \rangle_n + \langle a, a \rangle_n = -||b||^2 + ||a||^2.$
Q.E.D.

By §1, we know that for each $\alpha \neq 0$ in $\widetilde{\mathbb{A}}_{n+1}$. $\mathbb{H}_{\alpha} = \text{Span} \{e_0, \widetilde{\alpha}, \alpha, \widetilde{e}_0\}$ is a copy of \mathbb{A}_2 and that if $\mathbb{H}_{\alpha}^{\perp}$ denotes the orthogonal complement of \mathbb{H}_{α} , then $\mathbb{A}_{n+1} = \mathbb{H}_{\alpha} \oplus \mathbb{H}_{\alpha}^{\perp}$.

Since α is doubly pure, $\hat{\alpha}$ is also doubly pure; i.e., $\hat{\alpha} \in \{e_0, \tilde{e}_0\}^{\perp}$ and if $\hat{\alpha} \perp \alpha$ and $\hat{\alpha} \perp \tilde{\alpha}$, then $\hat{\alpha} \in \mathbb{H}^{\perp}_{\alpha}$. Now $S^{\sqrt{2}}(\tilde{\mathbb{A}}_{n+1}) = S^{2^{n+1}-3}$ denotes the sphere of radius $\sqrt{2}$ inside of $\tilde{\mathbb{A}}_{n+1}$.

Thus, we have a description of the real Stiefel manifold of 2- orthonormal frames in $\mathbb{R}^{2^{n-1}}$ as follows:

$$V_{2^{n}-1,2} = \{(a,b) \in {}_{o}\mathbb{A}_{n} \times {}_{o}\mathbb{A}_{n} = \widetilde{\mathbb{A}}_{n+1} : ||a|| = ||b|| = 1, a \perp b\}$$

and

$$V_{2^n-1,2} = \{ \alpha \in S^{\sqrt{2}}(\widetilde{\mathbb{A}}_{n+1}) : \hat{\alpha} \in \mathbb{H}_{\alpha}^{\perp} \}.$$

Lemma 2.2. If r and s are in \mathbb{R} with $r^2 + s^2 = 1$ and $(a, b) \in V_{2^n-1,2}$ then $(ra - sb, sa + rb) \in V_{2^n-1,2}$.

Proof: Suppose that ||a|| = ||b|| = 1 and $a \perp b$ in \mathbb{A}_n . Then $||ra - sb||^2 = r^2 ||a||^2 + s^2 ||b||^2 - 2rs\langle a, b \rangle_n$ and $||sa + rb||^2 = s^2 ||a||^2 + r^2 ||b||^2 + 2rs\langle a, b \rangle_n$, so $||ra - sb||^2 = ||sa + rb||^2 = r^2 + s^2 = 1$.

$$\langle ra - sb, sa + rb \rangle_n = rs \langle a, a \rangle_n - sr \langle b, b \rangle_n - s^2 \langle b, a \rangle_n + r^2 \langle a, b \rangle_n$$

= $rs ||a||^2 - rs ||b||^2 + 0 = rs - sr = 0.$

Corollary 2.3. $S^1 \times V_{2^n-1,2} \xrightarrow{a} V_{2^n-1,2}$ given by

$$((r, s), \alpha) \mapsto r\alpha + s\widetilde{\alpha} = (ra - sb, sa + rb)$$

defines a smooth, free S^1 -action on $V_{2^n-1,2}$.

Proof: Clearly $(1,0) \cdot \alpha = \alpha$ and

 $(r,s)[(q,t) \cdot \alpha] = ((r,s)(q,t)) \cdot \alpha = (rq - st, rt + sq) \cdot \alpha$ so a defines an action. It is a smooth action because it is a restriction of a linear action of $GL_2(\mathbb{R})$ on $\widetilde{\mathbb{A}}_{n+1} = \mathbb{R}^{2^{n+1}-2}$.

Finally, a is a free action: if $r\alpha + s\widetilde{\alpha} = \alpha$ then r = 1 and s = 0, because $\alpha \perp \widetilde{\alpha}$ in $\widetilde{\mathbb{A}}_{n+1}$.

Q.E.D.

Now, we identify $V_{2^n-2,2}$, the real Stiefel manifold of 2-orthonormal frames on \mathbb{R}^{2^n-2} , as a submanifold of $V_{2^n-1,2}$ as follows:

$$V_{2^n-2,2} = \{(a,b) \in V_{2^n-1,2} | (a,b) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n\};$$

i.e., $(a, b) \in V_{2^n-1,2}$ belongs to $V_{2^n-2,2}$ whenever a and b are doubly pure elements in \mathbb{A}_n and we have the known fibration [Wh]

thus, $V_{2^n-2,2}$ has dimension $2^n - 3 + 2^n - 4 = 2^{n+1} - 7$.

Since (ra - sb) and (sa + rb) are doubly pure elements in \mathbb{A}_n when a and b are doubly pure elements, we have that

 $V_{2^n-2,2}$ is a S^1 -invariant submanifold of $V_{2^n-1,2}$;

i.e., if $\alpha \in V_{2^n-2,2}$ then $(r,s) \cdot \alpha \in V_{2^n-2,2}$ for all $(r,s) \in S^1$.

We note that \mathbb{A}_{n+1} becomes a complex vector space by defining $i\dot{\alpha} = \tilde{\alpha}$ thus as a complex vector space

$$\widetilde{\mathbb{A}}_{n+1} = {}_{0}\mathbb{A}_{n} \times {}_{0}\mathbb{A}_{n} \cong \mathbb{C} \otimes_{\mathbb{R}} {}_{0}\mathbb{A}_{n}$$

The isomorphism takes $1 \otimes x$ to (x, 0) and $i \otimes y$ to (0, y) and S^1 (the set of modulo 1 complex numbers) acts naturally by multiplication on \mathbb{C} , hence on $\widetilde{\mathbb{A}}_{n+1}$.

Now, we identify the complex Stiefel manifold $W_{2^{n-1}-1,2}$ of 2-orthonormal frames in $\mathbb{C}^{2^{n-1}-1}$ as a submanifold of $V_{2^n-2,2}$ in terms of the Cayley-Dickson algebra \mathbb{A}_{n+1} for $n \geq 3$.

It is known that for a, b and x in \mathbb{A}_n , $\langle ax, b \rangle_n = \langle a, b\overline{x} \rangle_n$.(See [A]). Thus if x is a pure element, i.e., $\overline{x} = -x$ then $\langle ax, b \rangle_n = -\langle a, bx \rangle_n$. That is, right multiplication by a pure non-zero element is a skew-symmetric linear map. In particular $\langle \tilde{a}, b \rangle_n = -\langle a, \tilde{b} \rangle_n$.

Proposition 2.4. For $n \geq 3$, the map $\mathcal{H}_n : \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n \to \mathbb{C}$ given by

$$\mathcal{H}_n(a,b) = 2\langle a,b\rangle_n - 2i\langle \widetilde{a},b\rangle_n$$

defines a Hermitian inner product in $\widetilde{\mathbb{A}}_n$.

Proof: Clearly \mathcal{H}_n is \mathbb{R} -linear and

$$\overline{\mathcal{H}_n(a,b)} = 2\langle a,b\rangle_n + 2i\langle \widetilde{a},b\rangle_n$$
$$= 2\langle a,b\rangle_n - 2i\langle a,\widetilde{b}\rangle_n$$
$$= \mathcal{H}_n(b,a).$$

On the other hand,

$$\begin{aligned} \mathcal{H}_n(\widetilde{a}, b) &= 2\langle \widetilde{a}, b \rangle_n - 2i \langle \widetilde{a}, b \rangle_n \\ &= 2\langle \widetilde{a}, b \rangle_n + 2i \langle a, b \rangle_n \\ &= 2i \langle a, b \rangle_n - 2i^2 \langle \widetilde{a}, b \rangle_n \\ &= i \mathcal{H}_n(a, b). \end{aligned}$$

Q.E.D.

Proposition 2.5. For $n \ge 3$

$$W_{2^{n-1}-1,2} = \{(a,b) \in V_{2^n-2,2} | b \in \mathbb{H}_a^{\perp}\}.$$

Proof: First of all, we observe that $b \in \mathbb{H}_a^{\perp}$ for a and b in $\widetilde{\mathbb{A}}_n$ if and only if $b \perp a$ and $b \perp \widetilde{a}$, i.e. $\mathcal{H}_n(a,b) = 0$. If ||a|| = ||b|| = 1 and $\mathcal{H}_n(a,b) = 0$ then $(a,b) \in W_{m,2}$, where $m = \frac{1}{2}(2^n - 2) = 2^{n-1} - 1$.

Q.E.D.

Proposition 2.6. $W_{2^{n-1}-1,2}$ is S¹-invariant.

Proof: Suppose $(a, b) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$ with ||a|| = ||b|| = 1 and $b \in \mathbb{H}_a^{\perp}$. From this, we have $b \perp a$, and $\widetilde{b} \perp a$ (equivalently $\widetilde{a} \perp b$).

Now r(a, b) + s(-b, a) = (ra - sb, rb + sa) and we know that $(ra - sb) \perp (rb + sa)$.

To finish, we need to show that $(ra - sb) \perp (rb + sa)$.

$$\begin{aligned} \langle ra - sb, rb + sa \rangle_n &= \langle r\widetilde{a} - s\widetilde{b}, rb + sa \rangle_n \\ &= r^2 \langle \widetilde{a}, b \rangle_n - s^2 \langle \widetilde{b}, a \rangle_n + rs \langle \widetilde{a}, a \rangle_n - rs \langle \widetilde{b}, a \rangle_n \\ &= 0, \end{aligned}$$

therefore $(ra - sb) \in \mathbb{H}_{rb+sa}$.

Q.E.D.

Note that we have a fibration

$$S^{2^{n}-5} \rightarrow W_{2^{n-1}-1,2} \xrightarrow{\pi} S(\widetilde{\mathbb{A}}_{n}) = S^{2^{n}-3}$$
$$(a,b) \qquad \mapsto \qquad b$$

 $\pi^{-1}(b) = S(\mathbb{H}_b^{\perp}) = S^{2^n - 5}$ since dim $\mathbb{H}_b^{\perp} = 2^n - 4$. Thus dim $W_{2^{n-1} - 1, 2} = 2^n - 5 + 2^n - 3 = 2^{n+1} - 8$.

In [Mo₁] it is shown that for a and b in \mathbb{A}_n with $n \ge 4$ and ||a|| = ||b|| = 1, then

if ab = 0 then i) $(a, b) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$. ii) $b \in \mathbb{H}_a^{\perp}$ (or equivalently $a \in \mathbb{H}_b^{\perp}$). Thus

$$X_n := \{(a, b) \in \mathbb{A}_n \times \mathbb{A}_n : ||a|| = ||b|| = 1 \text{ and } ab = 0\}$$

is a subset of $W_{2^{n-1}-1,2}$.

Thus we have a chain of inclusions for $n \ge 3$,

$$X_n \subset W_{2^{n-1}-1,2} \subset V_{2^n-2,2} \subset V_{2^n-1,2}.$$

Now we show that X_n and $W_{2^{n-1}-1,2}$ admit a $T := S^1 \times S^1$ action.

Lemma 2.7 For $(a,b) \in V_{2^n-2,2}$ and r,s,q,p in \mathbb{R} with $r^2 + s^2 = 1$ and $p^2 + q^2 = 1$ define

$$(a,b) \stackrel{\tau}{\mapsto} (ra + s\tilde{a}, pb + qb).$$

i) If $(a, b) \in W_{2^{n-1}-1,2}$ then $(ra + s\tilde{a}, pb + q\tilde{b}) \in W_{2^{n-1}-1,2}$. ii) If $(a, b) \in X_n$ then $(ra + s\tilde{a}, pb + q\tilde{b}) \in X_n$. iii) τ defines a free *T*-action on $W_{2^{n-1}-1,2}$ and X_n respectively.

Proof: By direct calculations. If $a \perp b$ and $\tilde{a} \perp b$ then

$$\langle ra + s\tilde{a}, pb + q\tilde{b} \rangle_n = rp\langle a, b \rangle_n + sq\langle \tilde{a}, \tilde{b} \rangle_n + rq\langle a, \tilde{b} \rangle_n + sp\langle \tilde{a}, b \rangle_n$$

= 0 + 0 + 0 + 0
= 0

similarly.

$$\langle ra + s\tilde{b}, (pb + q\tilde{b})\tilde{e_0}\rangle_n = \langle ra + s\tilde{a}, p\tilde{b} - qb\rangle_n = rp\langle a, \tilde{b}\rangle_n + sp\langle \tilde{a}, \tilde{b}\rangle_n - rq\langle a, b\rangle_n - sq\langle \tilde{a}, b\rangle_n = 0 + 0 + 0 + 0 = 0.$$

If ab = 0 then

$$(ra + s\tilde{a})(pb + q\tilde{b}) = rp(ab) + sq\tilde{a}\tilde{b} + sp\tilde{a}b + rqa\tilde{b}$$

= 0.

Also

$$\begin{aligned} ||ra + s\tilde{a}||^2 &= r^2 ||a||^2 + s^2 ||\tilde{a}||^2 = (r^2 + s^2) ||a||^2 = 1 \quad \text{and} \\ ||pb + q\tilde{b}||^2 &= p^2 ||b||^2 + q^2 ||\tilde{b}||^2 = (p^2 + q^2) ||b||^2 = 1 \end{aligned}$$

Therefore we proved (i) and (ii).

Finally $(ra + s\tilde{a}, pb + q\tilde{b}) = (a, b)$ if and only if r = 1, s = 0, p = 1 and q = 0.

Clearly this action is smooth and free (see Corollary 2.5).

Q.E.D.

§ 3. X_n , Octonions and a S^3 action.

In this section, we show that we can attach to every element in X_n a copy of $\mathbb{A}_3 = \emptyset$, the octonions inside of \mathbb{A}_{n+1} for n > 3. This allows us to identify X_n with a subset of algebra monomorphisms of \mathbb{A}_3 into \mathbb{A}_{n+1} , which is our main goal in § 4.

We recall some notation from \S 1.

Let $e_0 \in \mathbb{A}_{n-1}$ be the unit so $(e_0, 0) = e_0$ is the unit in \mathbb{A}_n and $\tilde{e}_0 = (0, e_0)$ in \mathbb{A}_n .

For \tilde{e}_0 in \mathbb{A}_n we denote $\varepsilon = (\tilde{e}_0, 0)$ in \mathbb{A}_{n+1} .

For example for n = 4, $\tilde{e}_0 = e_8$ in \mathbb{A}_4 , and $\varepsilon = (e_8, 0)$ in \mathbb{A}_5

In general $\tilde{e}_0 = e_{2^n}$ in \mathbb{A}_{n+1} and $\varepsilon = e_{2^{n-1}}$ in \mathbb{A}_{n+1} . Since ε is a doubly pure element of norm one, we have that $\mathbb{H}_{\varepsilon} = \operatorname{Span}\{e_0, \tilde{\varepsilon}, \varepsilon, \tilde{e}_0\} \subset \mathbb{A}_{n+1}$ is a copy of \mathbb{A}_2 and a direct sum decomposition $\mathbb{A}_{n+1} = \mathbb{H}_{\varepsilon} \oplus \mathbb{H}_{\varepsilon}^{\perp}$.

By definition $\alpha = (a, b) \in \mathbb{A}_n \times \mathbb{A}_n = \mathbb{A}_{n+1}$ is doubly pure in \mathbb{A}_{n+1} with doubly pure entries in \mathbb{A}_n if and only if $\alpha \in \mathbb{H}_{\varepsilon}^{\perp}$.

In section § 2 we constructed the chain

$$X_n \subset W_{2^{n-1}-1,2} \subset V_{2^n-2,2} \subset V_{2^n-1,2} \subset \mathbb{A}_{n+1}$$

for $n \ge 3$ with $X_3 = \Phi$, the empty set.

Therefore, by definition, $V_{2^n-2,2} = V_{2^n-1,2} \cap \mathbb{H}_{\varepsilon}^{\perp}$.

Lemma 3.1. For $\alpha \in \mathbb{H}_{\varepsilon}^{\perp} \subset \mathbb{A}_{n+1}$ with $\alpha = (a, b) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$, 1) $(\alpha \varepsilon) \in \mathbb{H}_{\varepsilon}^{\perp}$ and $\alpha \varepsilon = (\widetilde{a}, -\widetilde{b})$. 2) $\alpha \widetilde{\varepsilon} \in \mathbb{H}_{\varepsilon}^{\perp}$ and $\alpha \widetilde{\varepsilon} = \widetilde{\alpha} \varepsilon = -\widetilde{\alpha} \widetilde{\varepsilon} = (-\widetilde{b}, -\widetilde{a})$.

Proof: By direct calculation

$$\alpha \varepsilon = (a, b)(\tilde{e}_0, 0) = (a\tilde{e}_0, -b\tilde{e}_0) = (\tilde{a}, -\tilde{b}) \in \tilde{\mathbb{A}}_n \times \tilde{\mathbb{A}}_n = \mathbb{H}_{\varepsilon}^{\perp}$$

and

$$\alpha \tilde{\varepsilon} = (a, b)(0, \tilde{e}_0) = (\tilde{e}_0 b, \tilde{e}_0 a) = (-\tilde{b}, -\tilde{a})$$

by Lemma 1.1 (1).

Finally using Lemma 1.1 (6) and (3) respectively $\alpha \tilde{\varepsilon} = \tilde{\alpha} \varepsilon = -\tilde{\alpha} \varepsilon \in \mathbb{H}_{\varepsilon}^{\perp}$.

Q.E.D.

Corollary 3.2. For a non-zero α in $\mathbb{H}_{\varepsilon}^{\perp} \subset \mathbb{A}_{n+1}$ and $n \geq 3$,

 $\phi_{\alpha} := Span\{e_0, \tilde{\varepsilon}, \varepsilon, \tilde{e}_0, \tilde{\alpha}, \alpha \varepsilon, \tilde{\varepsilon} \alpha, \alpha\} \subset \mathbb{A}_{n+1}$

is an 8-dimensional vector subspace of $\mathbb{A}_{n+1} = \mathbb{R}^{2^{n+1}}$.

Proof: By definition $\{e_0, \tilde{\varepsilon}, \varepsilon, \tilde{e}_0\}, \{e_0, \tilde{\alpha}, \alpha, \tilde{e}_0\}, \{\varepsilon, \alpha, \tilde{\varepsilon}, \tilde{\alpha}\}$ are an orthogonal set of vectors and $\alpha \varepsilon \in \mathbb{H}_{\varepsilon}^{\perp} \cap \mathbb{H}_{\alpha}^{\perp}$. Also by Lemma 3.1. $\tilde{\varepsilon}\alpha = -\alpha \tilde{\varepsilon} \in \mathbb{H}_{\varepsilon}^{\perp} \cap \mathbb{H}_{\alpha}^{\perp}$. Thus $\{e_0, \tilde{\varepsilon}, \varepsilon, \tilde{e}_0, \tilde{\alpha}, \alpha \varepsilon, \tilde{\varepsilon} \alpha, \alpha\}$ is an orthogonal set of vectors in \mathbb{A}_{n+1} . Q.E.D.

Remark: In particular for $\alpha \in V_{2^n-2,2}$ we have that $\phi_{\alpha} \cong \mathbb{R}^8 \subset \mathbb{A}_{n+1}$ and $\phi_{\alpha} \oplus \phi_{\alpha}^{\perp} = \mathbb{A}_{n+1}$.

Lemma 3.3. For $\alpha \in V_{2^n-2,2}$. $\alpha \in W_{2^{n-1}-1,2}$ if and only if $\hat{\alpha} \in \phi_{\alpha}^{\perp}$.

Proof: Recall that by definition $\hat{\alpha} = (b, a)$ if $\alpha = (a, b)$ so $\hat{\alpha} \in (\text{Span}((\{e_0, \tilde{\varepsilon}, \varepsilon, \tilde{e}_0, \alpha, \tilde{\alpha}\}))^{\perp}$ (see Lemma 2.1. above). Now

$$\langle \hat{\alpha}, \tilde{\varepsilon}\alpha \rangle_{n+1} = \langle (b,a), (b,\tilde{a}) \rangle_{n+1} = \langle b,b \rangle_n + \langle a,\tilde{a} \rangle_n = 0 \langle \hat{\alpha}, \alpha\varepsilon \rangle_{n+1} = \langle (b,a), (\tilde{a}, -\tilde{b}) \rangle_{n+1} = \langle b,\tilde{a} \rangle_n - \langle a,\tilde{b} \rangle_n = 2\langle b,\tilde{a} \rangle_n,$$

so $\hat{\alpha} \perp (\alpha \varepsilon)$ in \mathbb{A}_{n+1} if and only if $\tilde{a} \perp b$ in \mathbb{A}_n i.e. $b \in \mathbb{H}_a^{\perp}$. Therefore $\hat{\alpha} \in \phi_{\alpha}^{\perp}$ if and only if $b \in \mathbb{H}_a^{\perp}$.

Thus $W_{2^{n-1}-1,2} = \{ \alpha \in V_{2^n-2,2} | \hat{\alpha} \in \phi_{\alpha}^{\perp} \}.$

Theorem 3.4 For $\alpha \in W_{2^{n-1}-1,2}$ and $n \geq 4$, the following statements are equivalent.

i) $\alpha \in X_n$

ii) α alternate with ε i.e., $(\alpha, \alpha, \varepsilon) = 0$

iii) The vector subspace of \mathbb{A}_{n+1}

$$V(\alpha;\varepsilon) := \operatorname{Span}\{e_0, \alpha, \varepsilon, \alpha\varepsilon\}.$$

is multiplicatively closed and isomorphic to $\mathbb{A}_2 = \mathbb{H}$.

iv) ϕ_{α} is multiplicatively closed and isomorphic to $\mathbb{A}_3 = \phi$.

v) $\hat{\alpha} \in \text{Ker}L_{\alpha} \subset \phi_{\alpha}^{\perp}$, where L_{α} is left multiplication by α .

Proof. First of all, we calculate

 $\alpha(\alpha\varepsilon) = (a,b)[(a,b)(\tilde{e},0)] = (a,b)(\tilde{a},-\tilde{b}) = (a\tilde{a}-\tilde{b}b,-\tilde{b}a-b\tilde{a})$

$$= (-||a||^{2}\tilde{e}_{0} - ||b||^{2}\tilde{e}_{0}, -\tilde{b}a - \tilde{b}a) \text{ (by Lemma 1.1 (2) and (5))} \\ = -||\alpha||^{2}\varepsilon + 2(0, \tilde{b}a).$$

Therefore $\alpha(\alpha \varepsilon) = \alpha^2 \varepsilon = -||\alpha||^2 \varepsilon$ if and only if ba = 0, i.e. $\alpha \in X_n$ and we have (i) \Leftrightarrow (ii).

Clearly if $\alpha \in W_{2^{n-1}-1,2}$ then $\{e_0, \alpha, \varepsilon, \alpha\varepsilon\}$ form an orthonormal set of vectors so dim_{\mathbb{R}} $(V(\alpha; \varepsilon)) = 4$ so $-||\alpha||^2 = \alpha^2 = (\alpha\varepsilon)^2$ and $\alpha(\alpha\varepsilon) = -||\alpha||^2\varepsilon$ if and only if $V(\alpha; \varepsilon) = \mathbb{H}$, and we prove (ii) \Leftrightarrow (iii).

To prove (iii) \Leftrightarrow (iv) we stablish the following correspondence between the canonical basis in \mathbb{A}_3 and the orthonormal basis of ϕ_{α} .

$$e_1 \to \tilde{\varepsilon}; e_2 \mapsto \varepsilon; e_3 \to \tilde{e}_0; ||\alpha||e_4 \to \tilde{\alpha}; ||\alpha||e_5 \to \alpha\varepsilon; ||\alpha||e_6 \to \tilde{\varepsilon}\alpha; ||\alpha||e_7 \to \alpha$$

Using ii) it is a routine calculation to see that this correspondence define an algebra isomorphism. (See also Lemma 4.4 (1) below).

Finally by Lemma 3.3. we know $\hat{\alpha} \in \phi_{\alpha}^{\perp}$ and

$$\alpha \hat{\alpha} = (a, b)(b, a) = (ab + ab, a^2 - b^2) = (2ab, ||b||^2 - ||a||^2)$$

is the Hopf construction. So $\alpha \hat{\alpha} = 0$ in \mathbb{A}_{n+1} if and only if $\alpha \in X_n$. Recall that $\mathbb{A}_3 \cong \phi_{\alpha}$ does not admit zero divisors.

Q.E.D.

Theorem 3.5. $\mathbb{H}_{\varepsilon}^{\perp}$ admits a left \mathbb{H}_{ε} -module structure for $n \geq 3$.

Proof: For $\alpha = (a, b)$ in $\mathbb{H}_{\varepsilon}^{\perp} = \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$ and $u = re_0 + s\widetilde{\varepsilon} + q\varepsilon + p\widetilde{e}_0$ with r, s, q and p in \mathbb{R} .

Define

$$u \cdot \alpha = \alpha u = r\alpha + s\alpha \widetilde{\varepsilon} + q\alpha \varepsilon + p\widetilde{\alpha}.$$

Trivially $\widetilde{\alpha} \in \mathbb{H}_{\varepsilon}^{\perp}$ and $(\alpha \widetilde{\varepsilon})$ and $(\alpha \varepsilon)$ are in $\mathbb{H}_{\varepsilon}^{\perp}$ by Lemma 3.1.(2) and (1) respectively.

Since $\tilde{\varepsilon}, \varepsilon$ and \tilde{e}_0 are alternative elements in \mathbb{A}_{n+1} (actually they belong to the canonical basis) we have that $\tilde{\varepsilon} \cdot \alpha = (\alpha \tilde{\varepsilon})\tilde{\varepsilon} = \alpha(\tilde{\varepsilon})^2 = -||\alpha||^2 e_0 = \tilde{\varepsilon} \cdot \alpha$ and similarly $\varepsilon \cdot (\varepsilon \cdot \alpha) = \varepsilon^2 \cdot \alpha$ and $\tilde{e}_0 \cdot (\tilde{e}_0 \cdot \alpha) = \tilde{e}_0^2 \cdot \alpha$.

Now $\varepsilon \cdot (\widetilde{e}_0 \cdot \alpha) = \varepsilon \cdot (\alpha \widetilde{e}_0) = \varepsilon \cdot \widetilde{\alpha} = \widetilde{\alpha} \varepsilon$ and $(\varepsilon \widetilde{e}_0) \cdot \alpha = \widetilde{\varepsilon} \cdot \alpha = \alpha \widetilde{\varepsilon} = \widetilde{\alpha} \varepsilon$ by Lemma 3.1 (2).

Similarly

$$\begin{aligned} \widetilde{\varepsilon} \cdot (\widetilde{e}_0 \cdot \alpha) &= (\widetilde{\varepsilon} \widetilde{e}_0) \cdot \alpha = \varepsilon \alpha \\ \widetilde{e}_0 \cdot (\widetilde{\varepsilon} \cdot \alpha) &= (\widetilde{e}_0 \widetilde{\varepsilon}) \cdot \alpha = \alpha \varepsilon \\ \varepsilon \cdot (\widetilde{e}_0 \cdot \alpha) &= (\varepsilon \widetilde{e}_0) \cdot \alpha = \widetilde{\alpha} \varepsilon \\ \widetilde{e}_0 \cdot (\varepsilon \cdot \alpha) &= (\widetilde{e}_0 \varepsilon) \cdot \alpha = -\alpha \widetilde{\varepsilon}. \end{aligned}$$

Finally $\widetilde{\varepsilon} \cdot (\varepsilon \cdot \alpha) = \widetilde{\varepsilon} \cdot (\alpha \varepsilon) = (\alpha \varepsilon) \widetilde{\varepsilon} = (\widetilde{\alpha \varepsilon}) \varepsilon = -(\widetilde{\alpha \varepsilon}) \varepsilon = -\widetilde{\alpha} = (\widetilde{\varepsilon} \varepsilon) \cdot \alpha$ and $\varepsilon \cdot (\widetilde{\varepsilon} \cdot \alpha) = \varepsilon \cdot (\alpha \widetilde{\varepsilon}) = (\alpha \widetilde{\varepsilon}) \varepsilon = (\widetilde{\alpha} \varepsilon) \varepsilon = \widetilde{\alpha} = \varepsilon \cdot (\widetilde{\varepsilon} \alpha).$

By Lemma 3.1 and Lemma 1.1. and we are done.

Q.E.D.

Now we define a S^3 action on X_n . Consider the unit sphere inside of $\mathbb{H}_{\varepsilon} \subset \mathbb{A}_{n+1}$.

$$S^{3} = S(\mathbb{H}_{\varepsilon}) = \{ re_{0} + s\tilde{\varepsilon} + q\varepsilon + p\tilde{e}_{0} | r^{2} + s^{2} + q^{2} + p^{2} = 1 \}.$$

For $\alpha \in \mathbb{H}_{\varepsilon}^{\perp}$ with $\alpha = (a, b) \in \tilde{\mathbb{A}}_{n} \times \tilde{\mathbb{A}}_{n}$ define $\mathbb{H}_{\varepsilon}^{\perp} \times S^{3} \to \mathbb{H}_{\varepsilon}^{\perp}$ by

$$\begin{aligned} \alpha(re_0 + s\tilde{\varepsilon} + q\varepsilon + p\tilde{e}_0) &= r\alpha + s\alpha\tilde{\varepsilon} + q\alpha\varepsilon + p\alpha\tilde{e}_0 = r\alpha + s\tilde{\alpha}\varepsilon + q\alpha\varepsilon + p\tilde{\alpha}\\ &= r(a,b) + s(-\tilde{b},-\tilde{a}) + q(\tilde{a},-\tilde{b}) + p(-b,a)\\ &= (ra - s\tilde{b} + q\tilde{a} - pb, rb - s\tilde{a} - q\tilde{b} + pa). \end{aligned}$$

By definition this is a group action which is smooth and free of fixed points.

Corollary 3.6 The above action of $S^3 = S(\mathbb{H}_{\varepsilon})$ on $\mathbb{H}_{\varepsilon}^{\perp}$ is a group action which is smooth, orthogonal and free of fixed points.

Proof: By Theorem 3.5 this is a smooth group action because it is a restriction of a linear action. Since right multiplication by \tilde{e}_0 , ε and $\tilde{\varepsilon}$ are orthogonal linear transformations, we have that the action is orthogonal.

Finally this action is free of fixed points because $\{e_0, \tilde{\varepsilon}, \varepsilon, \varepsilon\}$ is an orthonormal basis, so $\alpha(re_0 + s\tilde{\varepsilon} + q\varepsilon + p\tilde{e}_0) = \alpha$ if and only if r = 1, s = q = p = 0. Q.E.D.

Theorem 3.8.

- i) The subsets X_n and $W_{2^{n-1}-1,2}$ of $\mathbb{H}_{\varepsilon}^{\perp}$ are S^3 -equivariant.
- ii) For α and β in $W_{2^{n-1}-1,2,}$

 $\phi_{\alpha} = \phi_{\beta}$, as vector spaces, if and only if α and β lie in the same S³-orbit.

Proof. For $\alpha \in \mathbb{H}_{\varepsilon}^{\perp}$ with $\alpha = (a, b) \in \tilde{\mathbb{A}}_n \times \tilde{\mathbb{A}}_n$ and r, s, q and p in \mathbb{R} with $r^2 + s^2 + q^2 + p^2 = 1$ we have that

$$\alpha(re_0 + s\tilde{\varepsilon} + q\varepsilon + s\tilde{e}_0) = (ra - s\tilde{b} + q\tilde{a} - pb, rb - s\tilde{a} - q\tilde{b} + pa).$$

Suppose that $\alpha \in W_{2^{n-1}-1,2}$ then $\langle \tilde{a}, b \rangle_n = -\langle a, \tilde{b} \rangle_n = 0$; $\langle a, b \rangle_n = \langle \tilde{a}, \tilde{b} \rangle_n = 0$ and by definition $\langle a, \tilde{a} \rangle_n = \langle b, \tilde{b} \rangle_n = 0$ with $||a|| = ||\tilde{a}|| = ||\tilde{b}|| = ||b|| = 1$ so $\langle ra - s\tilde{b} + q\tilde{a} - pb, rb - s\tilde{a} - q\tilde{b} + pa \rangle = p||a||^2 + sq||b||^2 - qs||\tilde{a}||^2 - pr||b||^2 = 0$ and $(\alpha(re_0 + s\tilde{\varepsilon} + q\varepsilon + s\tilde{e}_0)) \in V_{2^n_2-2,2}.$

Similarly $\langle ra - s\tilde{b} + q\tilde{a} - pb, r\tilde{b} + sa + qb + p\tilde{a} \rangle = 0$ and

$$\alpha(re_0 + s\tilde{\varepsilon} + q\varepsilon + s\tilde{e}_0) \in W_{2^{n-1}-1,2}.$$

A direct calculation shows that if ab = 0 then

$$ab = \tilde{a}b = a\tilde{b} = \tilde{a}\tilde{b} = 0$$

$$(ra - s\tilde{b} + q\tilde{a} - pb)(rb - s\tilde{a} - q\tilde{b} + pa) =$$
$$-rsa\tilde{a} + rpa^2 - sr\tilde{b}b + sq\tilde{b}^2 - qs\tilde{a}^2 + qpaa - prb^2 + pqb\tilde{b} =$$

$$-rs(-||a||^{2}\tilde{e}_{0}+||b||^{2}\tilde{e}_{0})+pq(-||a||^{2}\tilde{e}_{0}+||b||^{2}\tilde{e}_{0})+rp(a^{2}-b^{2})+sq(\tilde{b}^{2}-\tilde{a}^{2})=0$$

because $||a||^2 = ||b||^2 = 1$ and $a^2 = b^2 = -e_0$, so we have (i).

To prove ii) we notice that $\phi_{\alpha} = \mathbb{H}_{\varepsilon} \oplus \operatorname{Span}\{\tilde{\alpha}, \alpha \varepsilon, \tilde{\varepsilon} \alpha, \alpha\}$. Then if $\beta = r\alpha + s\tilde{\alpha}\varepsilon + q\alpha\varepsilon + p\tilde{\alpha}$ (recall that $\tilde{\alpha}\varepsilon = -\tilde{\varepsilon}\alpha$ by Lemma 4.1) and $||\beta|| = 1$ then $r^2 + s^2 + q^2 + p^2 = 1$ and $\alpha \equiv \beta \mod S^3$ if and only if $\phi_{\beta} \subset \phi_{\alpha}$ but

$$\dim \phi_{\beta} = \dim \phi_{\alpha} = 8 \text{ and } \phi_{\beta} = \phi_{\alpha}.$$

Q.E.D. **Remark:** Notice that $T = S^1 \times S^1$, as in Lemma 2.7. and $S^3 = S(\mathbb{H}_{\varepsilon})$ intersect on a copy of S^1 .

Suppose that $r^2 + s^2 + p^2 + q^2 = 1$ and $u^2 + v^2 = 1$ $t^2 + m^2 = 1$ in \mathbb{R} . If $(ra - s\tilde{b} + q\tilde{a} - pb, rb - s\tilde{a} - q\tilde{b} + pa) = (ua + v\tilde{a}, tb + m\tilde{b})$ then r = u, q = v, s = 0, p = 0, r = t, -q = m so

$$S(\mathbb{H}_{\varepsilon}) \cap T = S^1 = \{(r, -q) | r^2 + q^2 = 1\}.$$

§4. X_n and monomorphisms from \mathbb{A}_3 to \mathbb{A}_{n+1} .

In this chapter $1 \leq m \leq n$.

Definition. An algebra monomorphism from \mathbb{A}_m to \mathbb{A}_n is a linear monomorphism $\varphi : \mathbb{A}_m \to \mathbb{A}_n$ such that

i) $\varphi(e_0) = e_0$ (the first e_0 is in \mathbb{A}_m and the second e_0 in \mathbb{A}_n)

ii) $\varphi(xy) = \varphi(x)\varphi(y)$ for all x and y in \mathbb{A}_m .

By definition we have that $\varphi(re_0) = r\varphi(e_0)$ for all r in \mathbb{R} so $\varphi({}_0\mathbb{A}_m) \subset \varphi({}_0\mathbb{A}_n)$ and $\varphi(\overline{x}) = \overline{\varphi(x)}$ therefore $||\varphi(x)||^2 = \varphi(x)\overline{\varphi(x)} = \varphi(x)\varphi(\overline{x}) = \varphi(x\overline{x}) = \varphi(||x||^2) = ||x||^2$ for all $x \in \mathbb{A}_m$ and $||\varphi(x)|| = ||x||$ and φ is an orthogonal linear transformation from $\mathbb{R}^{2^{m-1}}$ to $\mathbb{R}^{2^{n-1}}$.

The *trivial* monomorphism is the one given by $\varphi(x) = (x, 0, 0, ..., 0)$ for $x \in \mathbb{A}_m$ and 0 in \mathbb{A}_m

 $\mathcal{M}(\mathbb{A}_m;\mathbb{A}_n)$ denotes the set of algebra monomorphisms from \mathbb{A}_m to \mathbb{A}_n .

For m = n, $\mathcal{M}(\mathbb{A}_m; \mathbb{A}_n) = \operatorname{Aut}(\mathbb{A}_n)$ the group of algebra automorphisms of \mathbb{A}_n

Proposition 4.1. $\mathcal{M}(\mathbb{A}_1;\mathbb{A}_n) = S({}_{o}\mathbb{A}_n) = S^{2^n-2}.$

Proof: $\mathbb{A}_1 = \mathbb{C} = Span\{e_0, e_1\}.$

If $x \in \mathbb{A}_1$ then $x = re_0 + se_1$ and for $w \in \mathbb{A}_n$ with ||w|| = 1 we have that $\varphi_w(x) = re_0 + sw$ define an algebra monomorphism from \mathbb{A}_1 to \mathbb{A}_n . This can be seen by direct calculations, recalling that, Center $(\mathbb{A}_n) = \mathbb{R}$ for all n and that every associator with one real entries vanish.

Conversely, for $\varphi \in \mathcal{M}(\mathbb{A}_1; \mathbb{A}_n)$, set $w = \varphi(e_1)$ so ||w|| = 1 and $\varphi_w = \varphi$. Q.E.D.

Remark: In particular, we have that

$$Aut(\mathbb{A}_1) = S^0 = \mathbb{Z}/2 = \{ \text{Identity, conjugation} \} = \{ \varphi_{e_1}, \varphi_{-e_1} \}.$$

To calculate $\mathcal{M}(\mathbb{A}_2; \mathbb{A}_n)$ for $n \geq 2$ we need to recall (see [Mo₂]).

Definition: For a and b in \mathbb{A}_n . We say that a alternate with b, we denote it by $a \rightsquigarrow b$, if (a, a, b) = 0.

We say that a alternate strongly with b, we denote it by $a \iff b$, if (a, a, b) = 0 and (a, b, b) = 0.

Clearly a alternate strongly with e_0 for all a in \mathbb{A}_n and if a and b are linearly dependent then $a \leftrightarrow b$ (by flexibility).

Also, by definition, a is an alternative element if and only if $a \rightsquigarrow x$ for all x in \mathbb{A}_n .

By Lemma 1.1 (1) and (2) we have that for any doubly pure element a in $\mathbb{A}_n(a, a, \tilde{e}_0) = 0$ and (by the above remarks) \tilde{e}_0 alternate strongly with any a in \mathbb{A}_n .

For a and b pure elements in \mathbb{A}_n , we define the vector subspace of \mathbb{A}_n

$$V(a;b) = Span\{e_0, a, b, ab\}.$$

Lemma 4.2. If $(a,b) \in V_{2^n-1,2}$ and $a \iff b$ then $V(a;b) = \mathbb{A}_2 = \mathbb{H}$ the quaternions.

Proof: Suppose that $(a, b) \in V_{2^n-1,2}$ and that (a, a, b) = 0 then we have

$$\begin{array}{lll} \langle ab,a\rangle &=& \langle b,\overline{a}a\rangle = \langle b,||a||^2e_0\rangle = ||a||^2\langle b,e_0\rangle = 0\\ \langle ab,a\rangle &=& \langle a,b\overline{b}\rangle = \langle a,||b||^2e_0\rangle = ||b||^2\langle a,e_0\rangle = 0\\ ||ab||^2 &=& \langle ab,ab\rangle &=& \langle \overline{a}(ab),b\rangle = \langle -a(ab),b\rangle = \langle -a^2b,b\rangle\\ &=& -a^2\langle b,b\rangle = ||a||^2||b||^2 = 1 \end{array}$$

so $\{e_0, a, b, ab\}$ is an orthonormal set of vectors in \mathbb{A}_n .

Finally using also that (a, b, b) = 0 and ab = -ba we may check by direct calculation that the multiplication table of $\{e_0, a, b, ab\}$ coincides with the one of the quaternions and by the identification $e_0 \mapsto e_0, a \mapsto e_1, b \mapsto e_2$ and $ab \mapsto e_3$ we have an algebra isomorphism between $\mathbb{A}_2 = \mathbb{H}$ and V(a; b). Q.E.D.

Proposition 4.3. $\mathcal{M}(\mathbb{A}_2; \mathbb{A}_n) = \{(a, b) \in V_{2^n-1,2} | a \iff b\}$ for $n \ge 2$. In particular

$$Aut(\mathbb{A}_2) = \mathcal{M}(\mathbb{A}_2; \mathbb{A}_2) = V_{3,2} = SO(3)$$

and

$$\mathcal{M}(\mathbb{A}_2,\mathbb{A}_3)=V_{7,2}$$

Proof. The inclusion " \supset " follows from Lemma 4.2. Conversely suppose that $\varphi \in \mathcal{M}(\mathbb{A}_2, \mathbb{A}_n)$ then $\varphi(e_0) = e_0, (\varphi(e_1), \varphi(e_2)) \in V_{2^n-1,2}$ and $V(\varphi(e_1), \varphi(e_2)) =$ Im $\varphi = \mathbb{H} \subset \mathbb{A}_n$.

Since \mathbb{A}_2 is an associative algebra and \mathbb{A}_3 is an alternative algebra we have that $a \leftrightarrow b$ for any two elements in \mathbb{A}_n for n = 2 or n = 3.

Q.E.D.

Remark. Recall that $\tilde{\mathbb{A}}_n = \{e_0, \tilde{e}_0\}^{\perp} = \mathbb{R}^{2^n-2}$ denotes the vector subspace of doubly pure elements. Since $a \iff \tilde{e}_0$ for any element in $\tilde{\mathbb{A}}_n$, we have that, if $a \in S(\tilde{\mathbb{A}}_n)$ i.e., ||a|| = 1 then $(a, \tilde{e}_0) \in V_{2^n-1,2}$ and the assignment $a \mapsto (a, \tilde{e}_0)$ defines an inclusion from $S(\tilde{\mathbb{A}}_n) = S^{2^n-3} \hookrightarrow \mathcal{M}(\mathbb{A}_2; \mathbb{A}_n) \subset V_{2^n-1,2}$ which resembles "the bottom cell" inclusion in $V_{2^n-1,2}$.

Now we show that X_n can be identified with a subset of $\mathcal{M}(\mathbb{A}_3; \mathbb{A}_{n+1})$ for $n \geq 4$.

Lemma 4.4. For $\alpha \in \mathbb{H}_{\varepsilon}^{\perp} \subset \mathbb{A}_{n+1}$ and $n \geq 4$. (1) If $||\alpha|| = 1$ then $\phi_{\alpha} = \text{Span}\{e_0, \tilde{\varepsilon}, \varepsilon, \tilde{e}_0, \tilde{\alpha}, \alpha\varepsilon, \tilde{\varepsilon}\alpha, \alpha\}$ is isomorphic as algebra to \mathbb{A}_3 if and only if $(\alpha, \alpha, \varepsilon) = 0$. (2) If $\alpha = (a, b) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$ then $(\alpha, \alpha, \varepsilon) = (0, -(a, \tilde{e}_0, b)) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$

Proof. (1) By definition $\tilde{e}_0 = e_{2^n}$ and $\varepsilon = e_{2^{n-1}}$ are elements in the canonical basis so they are alternative elements (See [Sch]). Since \mathbb{H}_{α} is associative for all α then $(\alpha, \alpha, \tilde{e}_0) = 0$.

Clearly if $\phi_{\alpha} \cong \mathbb{A}_3$ then $(\alpha, \alpha, \varepsilon)$ = because ϕ_{α} is an alternative algebra. Conversely, assume that $(\alpha, \alpha, \varepsilon) = 0$.

We have the following multiplication table that under the mapping $e_0 \mapsto e_0; e_1 \mapsto \tilde{\varepsilon}; e_2 \mapsto \varepsilon; e_3 \mapsto \tilde{e}_0; e_4 \mapsto \tilde{\alpha}; e_5 \mapsto \alpha \varepsilon; e_0 \mapsto \tilde{\varepsilon} \alpha \text{ and } e_7 \mapsto \alpha$

becomes an algebra monomorphism from \mathbb{A}_3 into \mathbb{A}_{n+1}

e_0	$\widetilde{\varepsilon}$	ε	\widetilde{e}_0	$\widetilde{\alpha}$	$\alpha \varepsilon$	$\widetilde{\varepsilon}\alpha$	α
ω	$-e_0$	\widetilde{e}_0	-ω	$\alpha \varepsilon$	$-\widetilde{\alpha}$	$-\alpha$	$\widetilde{\varepsilon}\alpha$
ε	$-\widetilde{e}_0$	$-e_0$	ìω	$\widetilde{\varepsilon}\alpha$	α	$-\widetilde{\alpha}$	$-\alpha\varepsilon$
\widetilde{e}_0	ε	$-\widetilde{\varepsilon}$	$-e_0$	α	$-\widetilde{\varepsilon}\alpha$	$+\alpha\varepsilon$	$-\widetilde{\alpha}$
$\widetilde{\alpha}$	$-\alpha\varepsilon$	$-\widetilde{\varepsilon}\alpha$	$-\alpha$	$-e_0$	ω	ε	$-\widetilde{e}_0$
$\alpha \varepsilon$	$\widetilde{\alpha}$	$-\alpha$	$\widetilde{\varepsilon}\alpha$	$-\widetilde{\varepsilon}$	$-e_0$	$-\widetilde{e}_0$	ε
$\widetilde{\varepsilon}\alpha$	α	$\widetilde{\alpha}$	$-\alpha\varepsilon$	$-\varepsilon$	\widetilde{e}_0	$-e_0$	$-\widetilde{\varepsilon}$
α	$-\widetilde{\varepsilon}\alpha$	$\alpha\varepsilon$	$\widetilde{\alpha}$	\widetilde{e}_0	$-\varepsilon$	$\widetilde{\varepsilon}$	$-e_0$

Notice that this table is skew-symmetric with $-e_0$'s along the diagonal.

The nontrivial calculations are: $\widetilde{\epsilon\alpha} = \alpha \varepsilon$ (by Lemma 1.1 (5)). $\widetilde{\epsilon}(\alpha \varepsilon) = -\widetilde{\epsilon\alpha} = \widetilde{\epsilon(\epsilon\alpha)} = \widetilde{\epsilon^2 \alpha} = -\widetilde{e_0 \alpha} = -\widetilde{\alpha}$ (by Lemma 1.1 (3)). $\varepsilon(\widetilde{\epsilon\alpha}) = -\varepsilon(\widetilde{\epsilon\alpha}) = -\widetilde{\epsilon(\epsilon\alpha)} = -\widetilde{\epsilon(\epsilon\alpha)} = -\widetilde{\alpha}$ (by Lemma 1.1 (1) and (6). $\widetilde{\alpha}(\alpha \varepsilon) = -\alpha(\alpha \varepsilon) = -\alpha^2 \varepsilon = \widetilde{\epsilon}$ because $(\alpha, \alpha, \varepsilon) = 0$ and $||\alpha|| = 1$. $\widetilde{\alpha}(\widetilde{\epsilon\alpha}) = -\widetilde{\alpha}(\widetilde{\epsilon\alpha}) = +\widetilde{\widetilde{\alpha}}(\alpha \varepsilon) = -\alpha(\alpha \varepsilon) = \varepsilon$, so we are done with (1).

To prove (2) we perform similar calculation as in Theorem 3.4.

$$\begin{aligned} \alpha(\alpha\varepsilon) &= (a,b)[(a,b)(\widetilde{e}_{0},0)] = (a,b)(\widetilde{a},-\widetilde{b}) = (a\widetilde{a}-\widetilde{b}b,-\widetilde{b}a-b\widetilde{a}) \\ &= (-||a||^{2}\widetilde{e}_{0}-||b||^{2}e_{0},-(b\widetilde{e}_{0})a+b(\widetilde{e}_{0}a)) \\ &= -||\alpha||^{2}\varepsilon - (0,(b,\widetilde{e}_{0},a)) \end{aligned}$$

Therefore

$$\begin{aligned} (\alpha, \alpha, \varepsilon) &= \alpha^2 \varepsilon - \alpha(\alpha \varepsilon) = -||\alpha||^2 - \alpha(\alpha \varepsilon) \\ &= (0, (b, \widetilde{e}_0, a)) = -(0, (a, \widetilde{e}_0, b)) & \text{by flexibility.} \end{aligned}$$
Q.E.D.

Notation: For $n \ge 4$ consider the following subsets of \mathbb{A}_{n+1}

$$\mathbb{E}_{n} = \{ \alpha \in \mathbb{H}_{\varepsilon}^{\perp} | (\alpha, \alpha, \varepsilon) = 0 \},\$$

$$S(\mathbb{E}_{n}) = \{ \alpha \in \mathbb{E}_{n} : ||\alpha|| = 1 \},\$$

$$P(n) = \{ (a, b) \in \widetilde{\mathbb{A}}_{n} \times \widetilde{\mathbb{A}}_{n} | a \text{ and } b \text{ are } \mathbb{C} - \text{collinear} \}$$

$$\overline{X}_{n} = \{ (a, b) \in \widetilde{\mathbb{A}}_{n} \times \widetilde{\mathbb{A}}_{n} | a \neq 0, b \neq 0 \text{ and } ab = 0 \}$$

and also the following subset of monomorphisms

$$\mathcal{M}_2(\mathbb{A}_3;\mathbb{A}_{n+1}) = \{\varphi \in \mathcal{M}(\mathbb{A}_3;\mathbb{A}_{n+1}) | \mathbb{H}_{\varepsilon} \subset \mathrm{Im}\varphi\}.$$

Remark: By Lemma 4.4 (1) we may identify $S(\mathbb{E}_n)$ and $\mathcal{M}_2(\mathbb{A}_3; \mathbb{A}_{n+1})$, that is, there is a one to one correspondence between this two sets.

Theorem 4.5. For $n \ge 4$

- (i) P(n) and \overline{X}_n are subsets of \mathbb{E}_n with $P(n) \cap \overline{X}_n = \Phi$.
- (ii) There is a continuous retract

$$R: \mathbb{E}_n \backslash P(n) \to \overline{X}_n$$

Proof: If $(a, b) \in P(n)$ then $b \in \mathbb{H}_a$ or $a \in \mathbb{H}_b$ and $(a, \tilde{e}_0, b) = 0$ (recall that \mathbb{H}_a and \mathbb{H}_b are associative), so by Lemma 4.4 (2) $(\alpha, \alpha, \varepsilon) = 0$ in \mathbb{A}_{n+1} when $\alpha = (a, b)$ so $P(n) \subset \mathbb{E}_n$.

On the other hand if $(a, b) \in \overline{X}_n$ then ab = 0 and $b \in \mathbb{H}_a^{\perp} \subset \mathbb{A}_n$ and by Lemma 1.1 (1), (6) and (3)

$$(a, \tilde{e}_0, b) = (a\tilde{e}_0)b - a(\tilde{e}_0b) = \tilde{a}b + a\tilde{b} = \tilde{a}b + \tilde{a}b = 2\tilde{a}b = -2\tilde{a}b.$$

Therefore if ab = 0 then $(a, \tilde{e}_0, b) = 0$ and $(\alpha, \alpha, \varepsilon) = 0$ in \mathbb{A}_{n+1} for $\alpha = (a, b)$ by Lemma 4.4 (2) so $\overline{X}_n \subset \mathbb{E}_n$.

Now if $(a, b) \in P(n) \cap \overline{X}_n$ then $b \in \mathbb{H}_a$ and ab = 0, but \mathbb{H}_a is associative and a = 0 or b = 0 which is a contradiction, therefore $P(n) \cap \overline{X}_n$ is the empty set and we are done with (i).

To prove (ii) suppose that $\alpha = (a, b) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$ with $a \neq 0$. Since

$$\mathbb{A}_n = \mathbb{H}_a \oplus \mathbb{H}_a^{\perp}$$

then there are unique elements c and d in \mathbb{H}_a and \mathbb{H}_a^{\perp} respectively such that b = c + d. Now

$$(a, \tilde{e}_0, b) = (a, \tilde{e}_0, c+d) = (a, \tilde{e}_0, c) + (a, \tilde{e}_0, d) = 0 + (a, \tilde{e}_0, d)$$

because \mathbb{H}_a is associative.

But by Lemma 1.1 (1), (6) and (3)

$$(a, \widetilde{e}_0, d) = (a\widetilde{e}_0)d - a(\widetilde{e}_0d) = \widetilde{a}d + a\widetilde{d} = \widetilde{ad} + \widetilde{a}d = -2\widetilde{ad}.$$

Therefore $(a, \tilde{e}_0, b) = -2\widetilde{ad}$.

Suppose that $\alpha = (a, b)$ is in $\mathbb{E}_n \setminus P(n)$ then $a \neq 0, b \neq 0$,

$$b = c + d \in \mathbb{H}_a \oplus \mathbb{H}_a^{\perp}$$

with $d \neq 0$ and $(a, \tilde{e}_0, b) = 0$ by Lemma 4.6 (2). Thus we have that ad = 0.

Let us define $R : \mathbb{E}_n \setminus P(n) \to \overline{X}_n$ as R(a, b) = (a, d).

Then R(a,b) = (a,b) if $(a,b) \in \overline{X}_n$ and R is continuous, because it is the restriction of the projection map

$$\widetilde{\mathbb{A}}_n \times \mathbb{A}_n \to \widetilde{\mathbb{A}}_n \times (\mathbb{H}_a \oplus \mathbb{H}_a^{\perp}) \to \widetilde{\mathbb{A}}_n \times \mathbb{H}_a^{\perp} (a, b) \to (a, c+d) \to (a, d)$$

which is obviously continuous.

Q.E.D.

Remarks: (1) Recall that $\widetilde{\mathbb{A}}_n$ is a complex vector space by making $ia = \widetilde{a}$. By definition $(a, b) \in P(n)$ if and only if a and b are \mathbb{C} -collinear for $(a, b) \in \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n$. So

$$P(n) \cong (((\widetilde{\mathbb{A}}_n \setminus \{0\}) \times \mathbb{C}) \cup \widetilde{\mathbb{A}}_n).$$

(2) Consider the map $w_n : \widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n \to \widetilde{\mathbb{A}}_n$ given by

$$w_n(a,b) = (a, \widetilde{e}_0, b).$$

Since every associator is a pure element $(a, \tilde{e}_0, b) \perp \tilde{e}_0$ because (a, -, b) is a skew-symmetric linear transformation (see [Mo₁]) then $(a, \tilde{e}_0, b) \in \tilde{\mathbb{A}}_n$.

Now w_n is a polynomia map, actually is a quadraic map, and $\mathbb{E}_n = w_n^{-1}(0)$ is a real algebraic set in $\widetilde{\mathbb{A}}_n \times \widetilde{\mathbb{A}}_n = \mathbb{H}_{\varepsilon}^{\perp} = \mathbb{R}^{2^{n+1}-4}$ with 0 in $\widetilde{\mathbb{A}}_n$ a singular value, by Lemma 4.4 (2).

(3) X_n is a contraction of $\overline{X_n}$ via normalization on each coordinate.

References

- [A] J. Adem. Construction of some normed maps. Bol. Soc. Mat. Mexicana, (2)20 1975, 59-79.
- [C₁] F. Cohen. A course in some aspects of classical homotopy theory. Lecture Notes in Mathematics 1139, 1989, Springer-Verlag, p. 11-120.
- [C₂] F. Cohen. On Whitehead Squares, Cayley-Dickson algebras and rational functions. Bol. Soc. Mat. Mexicana. (2)37 1992, 55-62.
- [D] L.E. Dickson. On quaternions and their generalization and the history of the 8 square theorem. Annals of Math; 20 155-171, 1919.
- [E-K] Eakin-Sathaye. On automorphisms and derivations of Cayley-Dickson algebras. J. Pure Appl. Algebra 129, 263-280, 1990.
 - [L] K. Lam. Construction of Non-singular bilinear maps. Topology 6. 1967, 423-426.
- [H-Y] S.H. Khalil and P. Yiu. The Cayley-Dickson algebras: A theorem of Hurwitz and quaternions. Bol. Soc. Sci. Lett. Lódz Vol. XLVIII 117-169. 1997.
- [Mo₁] G. Moreno. The zero divisors of the Cayley-Dickson algebras over the real numbers. Bol. Soc. Mat. Mexicana (2)4 13-27. 1998
- [Mo₂] G. Moreno. Alternative elements in the Cayley-Dickson algebras. hopf.math.purdue.edu/pub/Moreno.2001
- [Sch] R.D. Schafer. On the algebras formed by the Cayley-Dickson process. Amer. J. Math. 76 (1954) 435-445.
- [Wh] G. Whitehead. Elements of Homotopy theory. Graduate Texts in Math. 61 Springer-Verlag.

Departamento de Matemáticas. CINVESTAV DEL IPN. Apartado Postal 14-740. México D.F. MEXICO. gmoreno@math.cinvestav.mx