

## REAL VALUED DIMENSION FOR MORSE-SARD THEOREM

AZAT AINOULINE

ABSTRACT. A dimension allowing in particular to state necessary and sufficient conditions of the Morse-Sard theorem for real valued function is introduced.

## 1. INTRODUCTION.

The aim of this paper is to find a real valued dimension that is a good extension of the standard integer valued dimension on subsets of Euclidean spaces, in a sense that it would have more "good" properties, as for instance can be used to state necessary and sufficient conditions for the Morse-Sard Theorem.

A number of real valued dimensions have been intensively studied in geometric measure theory (see for example Mattila [6]), and some of the dimensions have desired properties such as :  $\dim \mathbb{R}^n = n$ ,  $A \subseteq B \implies \dim A \leq \dim B$ ,  $\dim \bigcup_{i \in \mathbb{N}} A_i = \sup_{i \in \mathbb{N}} \dim A_i$  etc. One of the central dimensions - the Hausdorff dimension - was used by many authors for sharpening the necessary part of the Morse-Sard theorem , for example:

**Theorem (Federer [4])**

Let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a  $C^k$  function,  $k \in \mathbb{N}$ ,  $C_p(F) = \{x \in \mathbb{R}^n | \text{rank}(DF) \leq p\}$ , then the Hausdorff measure of dimension  $p + \frac{n-p}{k}$  of  $F(C_p(F))$  is zero.

But it was shown that the Hausdorff dimension has lack of the property that  $\dim_H(A \times B) = \dim_H A + \dim_H B$  (see Federer [4] 2.10.29), and as we will see later, it does not give "right" dimension for set of critical values in the Morse-Sard theorem in sense that one cannot state that:

if Hausdorff measure of dimension  $p + \frac{n-p}{k}$  of a set  $A$  is zero then  $A \subseteq F(C_p(F))$  for some  $C^k$  function  $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$  (see theorems 4,8,9 here).

In this article the author introduces a promising candidate for the "right" real valued dimension that can fix at least that weakness of the Hausdorff dimension, regarding to a case of real valued function in the Morse-Sard theorem (theorems 8,9).

**Definition 1.1.** For  $m, n \in \mathbb{N}$ ,  $k \in \mathbb{R}$  a function  $\psi : B \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  is  **$D^k$ -function** if  $\exists K > 0 : \forall b, b' \in B \quad |\psi(b) - \psi(b')|^k \leq K|b - b'|$ .

**Definition 1.2.** For  $n \in \mathbb{N}$  **Outer dimension** of the set  $A \subseteq \mathbb{R}^n$  is a number  $Dm(A) = \inf\{k \in \mathbb{R}; \exists \psi : B \subseteq \mathbb{R}^1 \xrightarrow{D^k} \mathbb{R}^n, k \geq 0, A \subseteq \psi(B)\}$ .

**Theorem 1.** Let  $n \in \mathbb{N}$  then:

1991 *Mathematics Subject Classification.* Primary: 28A35; 28A75; 28A12; Secondary: 58C25, 58K05.

*Key words and phrases.* Dimension, measure, critical values, differentiability, Morse-Sard.

- (1)  $Dm(\emptyset) = 0$
- (2)  $Dm(A) \leq Dm(B)$  whenever  $A \subseteq B \subseteq \mathbb{R}^n$
- (3)  $Dm(\bigcup_{i \in \mathbb{N}} A_i) = \sup_{i \in \mathbb{N}} Dm(A_i)$  whenever  $A_1, A_2, \dots \subseteq \mathbb{R}^n$ .

**Theorem 2.** Every  $m$ -dimensional  $C^1$  manifold  $A \subseteq \mathbb{R}^n$ , ( $m, n \in \mathbb{N}$ ,  $m \leq n$ ) has  $Dm$ -dimension  $m$ .

**Theorem 3.**  $Dm(\psi(B)) \leq k \cdot Dm(B)$  for any  $D^k$ -function  $\psi : B \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$ ,  $k \in \mathbb{R}$ .

**Definition 1.3.** For  $A \subseteq \mathbb{R}^n$  **Hausdorff dimension** of  $A$  is a number  $\dim_H A = \inf \left\{ k; \liminf_{\sigma \rightarrow 0} \left\{ \sum_{i=1}^{\infty} (\text{diam}(B_i))^k; B_i \subseteq \mathbb{R}^n, \text{diam}(B_i) < \sigma, A \subseteq \bigcup_{i=1}^{\infty} B_i \right\} < \infty \right\}$ .

**Theorem 4.** (1)  $\dim_H \leq Dm$   
(2)  $\exists A \subseteq \mathbb{R}^1$ -compact, such that  $\dim_H A < Dm(A)$

**Theorem 5.** If  $A, B \subseteq \mathbb{R}^n$ , ( $n \in \mathbb{N}$ ), then  $Dm(A \times B) \leq Dm(A) + Dm(B)$ .

**Corollary 1** (from Theorem 5 and Marstrand's Theorem [5]). If  $A, B \subseteq \mathbb{R}^n$ , ( $n \in \mathbb{N}$ ), then  $\dim_H A + \dim_H B \leq Dm(A \times B) \leq Dm(A) + Dm(B)$ .

**Theorem 6.** If  $Dm(A) \neq Dm(B)$ , then sets  $A$  and  $B$  are not diffeomorphic.

**Definition 1.4.** For  $n, m \in \mathbb{N}, k \in [0, \infty)$  a function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$  satisfies a  $k$ -**Hölder condition** (or  $f \in H^k$ ) if for every compact neighborhood  $U$  there exists  $M > 0$  such that  $|f(x) - f(y)| \leq M \cdot |x - y|^k$  for all  $x, y \in U$ .

The next theorem is another definition of the  $Dm$  dimension.

**Theorem 7.** Let  $A \subseteq \mathbb{R}^n$ ,  $n \in \mathbb{N}$  then:

$$Dm(A) = \inf \{ 1/k; \exists \psi : B \subseteq \mathbb{R}^1 \xrightarrow{H^k, \text{onto}} A \} = \inf \{ k; \exists \psi : B \subseteq \mathbb{R}^1 \xrightarrow{H^{1/k}, \text{onto}} A \}.$$

**Definition 1.5.** For  $k = 0, 1, 2, \dots$ ,  $\lambda \in [0, 1)$  a function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$  is a  $C^{k, \lambda}$ -function (or  $f \in C^{k, \lambda}$ ), if  $f \in C^k$  and every  $k^{\text{th}}$  partial derivative of  $f$  is a  $H^\lambda$ -function.

Next three theorems show relationship among dimensions,  $C^k$  functions, critical points and critical values.

**Theorem 8.** Let  $F : \mathbb{R}^n \xrightarrow{C^{k, \lambda}} \mathbb{R}^m$ ,  $k \in \mathbb{N}$ ,  $\lambda \in [0, 1)$ ,  $p < \min\{m, n\}$  an integer. Then  $Dm(F(C_p(F))) \leq \min \left\{ p + \frac{n-p}{k+\lambda}, m \right\}$ .

**Theorem 9.** For any  $A \subseteq \mathbb{R}^1$ , ( $Dm(A) < 1$ ), natural  $m$  and real  $k$  ( $1 \leq k < \frac{m}{Dm(A)}$ ),  $A$  is subset of the critical values set of some  $C^{[k], k-[k]}$  function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^1$ , where  $[k]$  is integer part of  $k$ .

Note: In the special case when  $Dm(A) = 1$ , the existence of appropriate function for Theorem 9 is shown in [3].

**Theorem 10.**  $Dm(f(\sum_{k+\lambda} f)) \leq \frac{Dm(B)}{k+\lambda}$  for any  $C^{k, \lambda}$  map  $f : B \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  ( $m, n \in \mathbb{N}$ ,  $k+\lambda \neq 0$ ). where  $\sum_{k+\lambda} f = \{x \in B; \text{every partial derivative of } f \text{ of order } < k + \lambda \text{ vanishes at } x\}$

Next we define **Outer  $Dm$ -measure**  $m_a$ ,  $a \in [0, +\infty)$ .

**Definition 1.6.** Let  $A \subseteq \mathbb{R}^n$ ,  $n \in \mathbb{N}$ ,  $a \in [0, +\infty)$ . Then if  $a \geq Dm(A)$ ,  $m_a(A) = \inf\{meas(B); \exists f : B \subseteq \mathbb{R} \rightarrow \mathbb{R}^n, A \subseteq f(B), \forall b, b' \in B \ |\psi(b) - \psi(b')|^a \leq |b - b'|^a\}$ , where  $meas(B)$  is the Outer Lebesgue measure of  $B$ ; otherwise  $m_a(A) = \infty$ .

To show correctness of the definition 1.6 with regard to Dm-dimensions we will prove the following two theorems:

**Theorem 11.** Let  $n \in \mathbb{N}, a \in [0, +\infty)$  then:

- (1)  $m_a(\emptyset) = 0$
- (2)  $m_a(A) \leq m_a(B)$  whenever  $A \subseteq B \subseteq \mathbb{R}^n$
- (3)  $m_a(\bigcup_{i \in \mathbb{N}} A_i) \leq \sum_{i \in \mathbb{N}} m_a(A_i)$  whenever  $A_1, A_2, \dots \subseteq \mathbb{R}^n$ .

**Theorem 12.** Let  $A \subseteq \mathbb{R}^n$  ( $n \in \mathbb{N}$ ), then for any  $a > Dm(A)$ ,  $m_a(A) = 0$ .

## 2. PROOFS.

**Proof of Theorem 3.** Let  $t \in \mathbb{R}$  be such that  $Dm(B) < t$ , then there exists a  $D^t$  function  $f : A \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$ , that  $B \subseteq f(A)$ . By the "Composition property" [1]  $\psi \circ f : A \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  is a  $D^{tk}$  function and also  $\psi(f(A)) \supseteq \psi(B)$ . It means that  $Dm(\psi(B)) \leq k \cdot t$ , and because it is true for every  $t > Dm(B)$ , then  $Dm(\psi(B)) \leq k \cdot Dm(B)$ .  $\square$

**Lemma 2.1.** Let  $A = \bigcup_{i \in \mathbb{N}} A_i \subseteq \mathbb{R}^n$ ,  $n \in \mathbb{N}$ ,  $k \geq 0$ ,  $k \in \mathbb{R}$  and  $\{f_i : B_i \subseteq [a_i, b_i] \xrightarrow{D^k} \mathbb{R}^n, i \in \mathbb{N}\}$  be such that  $A_i \subseteq f_i(B_i)$ . Then there exists  $f : B \subseteq \mathbb{R}^1 \xrightarrow{D^k} \mathbb{R}^n$  with  $A \subseteq f(B)$ .

**Proof.** By the definition of  $D^k$ -function there exist real positive numbers  $K_i$  ( $i \in \mathbb{N}$ ), such that for each  $i$   $|f_i(x) - f_i(y)|^k \leq K_i|x - y|$ . Also from the "Extension on closure" property [1] every  $f_i(B_i)$  is bounded set in  $\mathbb{R}^n$ . Let us take sets  $\{[x_i, y_i] \subseteq \mathbb{R}^1, i \in \mathbb{N}\}$  and  $\{l_i : [x_i, y_i] \xrightarrow{ont^0} [a_i, b_i], i \in \mathbb{N}\}$  such that:

- (1)  $x_1 = 0$
- (2)  $\forall i \in \mathbb{N}$ 
  - $x_i < y_i < x_{i+1}$
  - $y_i - x_i \geq K_i(b_i - a_i)$
  - $x_{i+1} - y_i > \sup_{j \leq i, a \in [a_j, b_j], b \in [a_{i+1}, b_{i+1}]} |f_j(a) - f_{i+1}(b)|^k$
- (3)  $\forall i \in \mathbb{N}, \forall x \in [x_i, y_i] \quad l_i(x) = \frac{x - x_i}{y_i - x_i}(b_i - a_i) + a_i$ .

Now by defining  $B = \bigcup l_i^{-1}(B_i)$  and  $f : B \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  such that  $f = f_i \circ l_i \upharpoonright (B \cap [x_i, y_i])$  we obtain the required  $D^k$  function  $f$  such that  $|f(x) - f(y)|^k \leq 1 \cdot |x - y| \quad \forall x, y \in B, \quad A \subseteq f(B)$ .  $\square$

**Proof of Theorem 1.** The parts 1),2) of this theorem are obvious. Let's prove the part 3).  $Dm(\bigcup_{i \in \mathbb{N}} A_i) \geq \sup_{i \in \mathbb{N}} Dm(A_i)$  follows from the part 2). Then suffice it to prove that  $Dm(\bigcup_{i \in \mathbb{N}} A_i) \leq \sup_{i \in \mathbb{N}} Dm(A_i)$ . Let  $k \in \mathbb{R}$  be such that  $\sup_{i \in \mathbb{N}} Dm(A_i) < k$ , then for each  $i$  there exists a  $D^k$  function  $g_i : M_i \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  such that  $A_i \subseteq g_i(M_i)$ .

Let us define a denumerable set of  $D^k$  functions:  $F = \{f_{ij} = g_i \upharpoonright B_{ij}; B_{ij} = [j, j+1] \cap M_i, j \in \mathbb{Z}, i \in \mathbb{N}\}$ . Then we have a denumerable set  $P = \{f_{ij}(B_{ij}); i \in \mathbb{N}, j \in \mathbb{Z}\}$  such that  $\bigcup_{i \in \mathbb{N}} A_i = \bigcup_{i \in \mathbb{N}, j \in \mathbb{Z}} f_{ij}(B_{ij})$ . And the sets  $F, P$  satisfy

the condition of Lemma 2.1. So that there exists a function  $f : B \subseteq \mathbb{R}^1 \xrightarrow{D^k} \mathbb{R}^n$  with  $\bigcup_{i \in \mathbb{N}} A_i \subseteq f(B)$ . But it means that  $Dm(\bigcup_{i \in \mathbb{N}} A_i) \leq k$ . And because  $Dm(\bigcup_{i \in \mathbb{N}} A_i) \leq k$  for all  $k$  such that  $\sup_{i \in \mathbb{N}} Dm(A_i) < k$ , it follows that  $Dm(\bigcup_{i \in \mathbb{N}} A_i) \leq \sup_{i \in \mathbb{N}} Dm(A_i)$ .  $\square$

**Proof of Theorem 7.** The proof based on the facts that every  $D^k$ ,  $k > 0$  is a  $H^{1/k}$  function, and on the other hand if  $f \in H^{1/k}$  then  $\forall [a_i, b_i] \subseteq \mathbb{R}$  a function  $f_i = f \upharpoonright [a_i, b_i]$  is a  $D^k$  function. Then by applying the lemma 2.1 we finish the proof of Theorem 7.  $\square$

**Proof of Theorem 5.** We set  $K_0^n = \{Q_{i_0}^n, i_0 \in \mathbb{N}\}$ , where  $Q_{i_0}^n$  is a closed cube in  $\mathbb{R}^n$  with side length 1, and every coordinate of any vertex of  $Q_{i_0}^n$  is an integer. In general, having constructed the cubes of  $K_{s-1}^n$ , divide each  $Q_{i_0, i_1, i_2, \dots, i_{s-1}}^n \in K_{s-1}^n$  into  $2^n$  closed cubes of side  $\frac{1}{2^s}$ , and let  $K_s^n$  be the set of all these cubes. More precisely we will write for  $s \in \mathbb{N}$ :

$$(2.1) \quad K_s^n = \{Q_{i_0, i_1, i_2, \dots, i_{s-1}, i_s}^n ; Q_{i_0, i_1, i_2, \dots, i_{s-1}, i_s}^n \subseteq Q_{i_0, i_1, i_2, \dots, i_{s-1}}^n \in K_{s-1}^n, 1 \leq i_s \leq 2^n\}.$$

And we apply here:

**Lemma 2.2** ([2]). *Let  $n, p \in \mathbb{N}$ ,  $p \leq n$ ,  $k \in \mathbb{R}, k \geq 1$ . Then there exists a continuous space-filling function  $\pi_{k,p}^n = (\pi_1, \pi_2) : [0, 1] \xrightarrow{ont\ o} [0, 1]^n$  such that  $\pi_1 : [0, 1] \xrightarrow{D^{\frac{pk+n-p}{k}}} [0, 1]^p$  and  $\pi_2 : [0, 1] \xrightarrow{D^{pk+n-p}} [0, 1]^{n-p}$  are the component functions of  $\pi_{k,p}^n$ .*

Now let  $k, t$  be real numbers such that  $Dm(A) < k$ ,  $Dm(B) < t$ , then there exists a  $D^k$  function  $\psi^* : A^* \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  with  $A \subseteq \psi^*(A^*)$  and a  $D^t$  function  $\psi^{**} : B^* \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  with  $B \subseteq \psi^{**}(B^*)$ . Let us consider the following:

- a) A function  $\psi : A^* \times B^* \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$  such that  $\psi(x, y) = (\psi^*(x), \psi^{**}(y)) \in \mathbb{R}^{2n} \quad \forall (x, y) \in A^* \times B^*$ .
- b) The set  $K_0^2$  (see (2.1)).
- c) A family of functions  $\psi_\delta = \psi \upharpoonright \delta \cap (A^* \times B^*)$ ,  $\delta \in K_0^2$ .
- d) The function  $\pi_{k/t,1}^2 = (\pi_1, \pi_2) : [0, 1] \xrightarrow{ont\ o} [0, 1]^2$  (see Lemma 2.2).
- e) A family of the identity maps  $I_\delta : [0, 1]^2 \xrightarrow{ont\ o} \delta$  ( $\delta \in K_0^2$ ) such that  $I_\delta = (I_\delta^*, I_\delta^{**})$  and  $I_\delta^*, I_\delta^{**} : [0, 1] \xrightarrow{ont\ o} \delta^*, \delta^{**} \in K_0^1$  are identity maps.
- f) A family of sets  $C_\delta = (\pi_{k/t,1}^2)^{-1}(I_\delta^{-1}(\delta \cap (A^* \times B^*)))$ ,  $\delta \in K_0^2$ .
- g) A family of functions  $f_\delta : C_\delta \subseteq [0, 1] \rightarrow \mathbb{R}^2$  ( $\delta \in K_0^2$ ), such that  $\forall \delta \in K_0^2$ ,  $x \in C_\delta$   $f_\delta(x) = \psi_\delta(I_\delta(\pi_{k/t,1}^2(x)))$ .

Let us prove that

- 1)  $f_\delta$  is  $D^{k+t}$  function for each  $\delta \in K_0^2$
- 2)  $A \times B \subseteq \bigcup_{\delta \in K_0^2} f_\delta(C_\delta)$ .

- 1)  $\forall \delta \in K_0^2$  by Lemma 2.2 and g)  $f_\delta = \psi_\delta \circ I_\delta \circ \pi_{k/t,1}^2 = \psi_\delta \circ I_\delta \circ (\pi_1, \pi_2) = \psi_\delta \circ (I_\delta^* \circ \pi_1, I_\delta^{**} \circ \pi_2) = (\psi^* \circ I_\delta^* \circ \pi_1, \psi^{**} \circ I_\delta^{**} \circ \pi_2)$ , where  $\psi_\delta^*$ ,  $\psi_\delta^{**}$  are the component functions of  $\psi_\delta$ .

Recalling that  $\psi^* \in D^k$ ,  $\psi^{**} \in D^t$ , we can see that  $\psi_\delta^* \in D^k$ ,  $\psi_\delta^{**} \in D^t$ ,  $\pi_1 \in D^{\frac{k/t+2-1}{k/t}}$ ,  $\pi_2 \in D^{k/t+2-1}$  by Lemma 2.2, and  $I_\delta^*, I_\delta^{**} \in D^1$  as

identity maps.

Thus by "Composition property" [1]  $\psi^* \circ I_\delta^* \circ \pi_1 \in D^{k+t}$  and  $\psi^{**} \circ I_\delta^* \circ \pi_2 \in D^{k+t}$ . Consequently  $f_\delta \in D^{k+t}$  as both its component functions are  $D^{k+t}$ .

- 2) Recall that  $A \subseteq \psi^*(A^*)$ ,  $B \subseteq \psi^{**}(B^*)$ , then
- $A \times B \subseteq \text{rang } \psi$  by **a**);
  - $\text{rang } \psi \subseteq \bigcup_{\delta \in K_0^2} \text{rang } \psi_\delta$  by **c**).

Also  $I_\delta : [0, 1]^2 \rightarrow \delta$  is "onto" as identity map and  $\pi_{k/t, 1}^2$  is "onto" by Lemma 2.2. So that by **g**)  $A \times B \subseteq \bigcup_{\delta \in K_0^2} \text{rang } f_\delta = \bigcup_{\delta \in K_0^2} f_\delta(C_\delta)$ .

Now from 1) and 2) we can see that the set  $A \times B$  and the denumerable family of functions  $f_\delta$ ,  $\delta \in K_0^2$  satisfy the conditions of Lemma 2.1, so that there exists  $f : M \subseteq \mathbb{R}^1 \xrightarrow{D^{k+t}} \mathbb{R}^{2n}$  with  $A \times B \subseteq f(M)$ . And because it is true for all  $k, t$  such that  $Dm(A) < k$ ,  $Dm(B) < t$ , then  $Dm(A \times B) \leq Dm(A) + Dm(B)$ .  $\square$

#### Proof of Theorem 4.

Part 1. Let  $k \in \mathbb{R}$  be a number such that  $Dm(A) < k$ , then there exists a  $D^k$  function  $f : B \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  with  $A \subseteq f(B)$ , and consequently there exists  $M > 0$  that  $\forall x, y \in B \quad |f(x) - f(y)|^k < M|x - y|$ . For every  $\sigma > 0$  let us take a natural number  $s_0$  such that  $(\frac{M}{2^{s_0}})^{1/k} < \sigma$ . Now considering the set  $K_{s_0}^1$  (see (2.1)), we can see:

- (1)  $A \subseteq \bigcup_{i \in \mathbb{N}} A_i$ , where  $A_i = f(B \cap [-i, i])$
- (2)  $A_i \subseteq \bigcup_{\delta \in K_{s_0}^1, \delta \subseteq [-i, i]} f(B \cap \delta) \quad \forall i \in \mathbb{N}$
- (3)  $\forall \delta \in K_{s_0}^1 \quad \text{diam } f(B \cap \delta) \leq \sigma$
- (4)  $\forall i \quad \sum_{\delta \in K_{s_0}^1, \delta \subseteq [-i, i]} (\text{diam } f(B \cap \delta))^k < 2Mi$ .

Thus for each  $i \quad \dim_H A_i \leq k$  and therefore  $\dim_H A \leq k$  because  $\dim_H A \leq \sup_i \dim_H A_i$  (see 4.8 in [6]). And since  $\dim_H A \leq k$  for every  $k > Dm(A)$ , it follows that  $\dim_H A \leq Dm(A)$ .

Part 2. There exist compact sets  $A, B \subseteq \mathbb{R}^1$  such that  $\dim_H A + \dim_H B < \dim_H(A \times B)$  (see [4]), then by the Part 1 and theorem 5  $\dim_H A + \dim_H B < Dm(A) + Dm(B)$ , so either  $\dim_H A < Dm(A)$  or  $\dim_H B < Dm(B)$  is true.  $\square$

**Proof of Theorem 2.** As follows from the Theorem 1, it only remains to show that  $Dm([0, 1]^m) = m$ . In [1] it was proven that  $\forall n \in \mathbb{N} \quad \exists f_n : [0, 1] \xrightarrow{D^n, \text{onto}} [0, 1]^n$  so that  $Dm([0, 1]^m) \leq m$ . Now by Theorem 4  $Dm \geq \dim_H$ , and  $\dim_H [0, 1]^m = m$  (see p.59 in [6]), then it follows that  $Dm([0, 1]^m) = m$ .  $\square$

**Lemma 2.3.** Every  $C^{k, \lambda}$  function ( $k + \lambda \neq 0$ )  $f : U \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$ , with  $U$  bounded, is  $D^{\frac{1}{k+\lambda}}$  function on a set  $\sum_k^* f = \{x \in U ; \text{ every partial derivative of } f \text{ of order } \leq k^* \text{ vanishes at } x, k^* = k \text{ if } \lambda \neq 0; k^* = k - 1 \text{ if } \lambda = 0\}$ .

**Proof.** If  $k = 0$ , then conclusion of this theorem follows from the definition of  $C^{0, \lambda}$  function for all  $x \in U$ .

If  $k \in \mathbb{N}$ , we apply Taylor formulae for the points  $x \in \sum_k^* f$  and use the fact that  $D_{k^*} f \in C^{0, \lambda}$  if  $\lambda \neq 0$  or  $D_{k^*} f \in C^1$  if  $\lambda = 0$ .  $\square$

**Proof of Theorem 6.** We can suppose without loss of generality that  $A, B$  are bounded and  $Dm(A) < Dm(B)$ , then there exist  $k \in \mathbb{R} : Dm(A) < k < Dm(B)$  and a  $D^k$  function  $f : E \subseteq \mathbb{R} \xrightarrow{onto} A$ . By contradiction, if  $A, B$  are diffeomorphic then there is a  $C^1$  function  $g : A \xrightarrow{onto} B$ , and by Lemma 2.3  $g \in D^1$ . Now by the "Composition property" ([1]) we have a  $D^k$  function  $g \circ f : E \subseteq \mathbb{R} \xrightarrow{onto} B$  that contradicts  $k < Dm(B)$ .  $\square$

**Proof of Theorem 8.** It follows from Main Theorem [2], Theorem 1 and Lemma 2.3.  $\square$

**Proof of Theorem 10.** It follows from Theorems 3, 1 and Lemma 2.3.  $\square$

**Proof of Theorem 9.** It is possible to choose a real number  $p \in \mathbb{R}$  such that  $m < p$  and  $k < p < \frac{m}{Dm(A)}$ , then  $Dm(A) < \frac{m}{p}$  and by the Definition 1.2 there exists a  $D^{m/p}$  function  $g : B \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^1$  such that  $A \subseteq g(B)$ . By the " $C^{<k}$ -extension property" ([2]) there exists a  $C^{<[\frac{p}{m}] \cdot \frac{p}{m} - [\frac{p}{m}]}$ -function  $q : \mathbb{R}^1 \rightarrow \mathbb{R}^1$  such that  $A \subseteq q(\sum_{p/m} q)$ , ((Designation: We write  $F \in C^{<k, \lambda}$  if  $F \in C^{p, \beta}$  for all  $p + \beta < k + \lambda$ )) then by the Main Theorem [1] there exists a  $C^{<[p] \cdot p - [p]}$ -function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^1$  such that  $q(\sum_{p/m} q)$  is the critical values set of the function  $f$  (Note that  $C^{k+\lambda} \subseteq C^{k, \lambda}$ ). From  $A \subseteq q(\sum_{p/m} q)$  and  $k < p$ , we conclude that  $f \in C^{[k] \cdot k - [k]}$  and  $A$  is subset of the critical values set of  $f$ .  $\square$

**Lemma 2.4.**  $\forall k > 1$  there exists a  $D^k$ -function  $g : B \subseteq \mathbb{R}^1 \xrightarrow{onto} \mathbb{R}^1$  such that  $\text{meas}(B)=0$ .

**Proof.** We define a family of sets  $P = \bigcup_{s \in \mathbb{N}} P_s$  on  $\mathbb{R}^1$ . For  $s = 1$   $P_1 = \{G_{i_0, i_1} : i_0 \in \mathbb{N}, i_1 = 1, 2\}$ , where each  $G_{i_0, i_1}$  is a closed interval of length  $\frac{1}{2^k}$  located in the middle of  $Q_{i_0, i_1}^1 \in K_1^1$  defined in (2.1). And if  $P_{s-1} = \{G_{i_0, \dots, i_j, \dots, i_{s-1}} : i_j = 1, 2\}$  have already been constructed, then  $P_s = \{G_{i_0, \dots, i_j, \dots, i_s} : i_j = 1, 2\}$ , where  $G_{i_0, \dots, i_{s-1}, 1}, G_{i_0, \dots, i_{s-1}, 2}$  are two closed intervals of length  $\frac{1}{2^{ks}}$ , first located at the beginning, second at the end of  $G_{i_0, \dots, i_{s-1}}$ . We can notice that distance between them is

$$(2.2) \quad \frac{1}{2^{k(s-1)}} - \frac{2}{2^{ks}} = (2^k - 2) \frac{1}{2^{ks}}.$$

Let  $B \subseteq \mathbb{R}^1$  be the Cantor set defined by the family of sets  $P$ . Let us define a function  $g : B \subseteq \mathbb{R}^1 \xrightarrow{onto} \mathbb{R}^1$  with the condition  $g(G_{i_0, \dots, i_s} \cap B) = Q_{i_0, \dots, i_s}^1 \in K_s^1$  (see (2.1)). To check that  $g \in D^k$ , let us take arbitrary points  $x, y \in B$ , ( $x < y$ ), then there exists  $s \in \mathbb{N}$  such that  $x \in G_{i_0, \dots, i_{s-1}, 1}$ ,  $y \in G_{i_0, \dots, i_{s-1}, 2}$ .

$$(2.3) \quad |g(x) - g(y)|^k \leq \left( \frac{1}{2^{s-1}} \right)^k \leq K|x - y|$$

where  $K = \frac{2^k}{2^k - 2}$  depends only on  $k$ .

To prove that  $\text{meas}(B) = 0$ , suffice it to show that  $\text{meas}(B \cap [a, a+1]) = 0 \quad \forall a \in \mathbb{Z}$ . And the  $\text{meas}(B \cap [a, a+1]) \leq \lim_{s \rightarrow \infty} 2^s \cdot \frac{1}{2^{ks}} = 0$  because  $k > 1$ .  $\square$

**Proof of Theorem 12.** Let  $a, b \in \mathbb{R}$  be such that  $Dm(A) < b < a$ , then by the definition 1.2 there exists a  $D^b$ -function  $f_a : F \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  such that  $A \subseteq f_a(F)$ . We can suppose that the coefficient  $K$  required for  $f_a$  by the definition 1.1 is equal to 1 (stretching  $F$  if necessary). By Lemma 2.4 and because  $\frac{a}{b} > 1$ , there exists  $g : G \subseteq \mathbb{R}^1 \xrightarrow{onto} \mathbb{R}^1$  such that  $\text{meas}(G) = 0$ ,  $\exists K^* > 0 : \forall x, y \in G \quad |g(x) - g(y)|^{a/b} \leq K^* |x - y|$ .

Let us define a function  $q : Q \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^1$  such that  $Q = \{K^*x ; x \in G\} \quad \forall z \in Q \quad q(z) = g(\frac{z}{K^*})$ , and a function  $h = f_a \circ q \upharpoonright H$ , where  $H = Q \cap q^{-1}(F)$ .

Now we have the following:

- (1)  $A \subseteq h(H)$
- (2)  $\text{meas}(H) = 0$
- (3)  $\forall x, y \in H \quad |h(x) - h(y)|^a = |f_a(q(x)) - f_a(q(y))|^{ab/b} \leq |q(x) - q(y)|^{a/b} = |g(\frac{x}{K^*}) - g(\frac{y}{K^*})|^{a/b} \leq K^* |\frac{x}{K^*} - \frac{y}{K^*}| = 1 \cdot |x - y|$ .

And by the definition 1.6 and from 1,2,3 it follows that  $m_a(A) = 0$ .  $\square$

**Lemma 2.5.** Let  $A = \bigcup_{i \in \mathbb{N}} A_i \subseteq \mathbb{R}^n$ ,  $n \in \mathbb{N}$ ,  $k \geq 0$ ,  $k \in \mathbb{R}$  and  $\{f_i : B_i \subseteq [a_i, b_i] \rightarrow \mathbb{R}^n, i \in \mathbb{N}\}$  be such that  $\forall i \in \mathbb{N} \quad A_i \subseteq f_i(B_i)$ , and  $\forall b, b' \in B_i \quad |f_i(b) - f_i(b')|^k \leq |b - b'|$ . Then there exists  $f : B \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  with  $A \subseteq f(B)$ , such that  $\forall b, b' \in B \quad |f(b) - f(b')|^k \leq |b - b'|$  and  $\text{meas}(B) \leq \sum_{i \in \mathbb{N}} \text{meas}(B_i)$ .

**Proof:** Similar to the prove of lemma 2.1.  $\square$

**Proof of Theorem 11.** 1) and 2) are obvious. 3) Let's consider that  $\forall i \in \mathbb{N} \quad m_a(A_i) < +\infty$  (otherwise it is obvious). Now for an arbitrary fixed  $\epsilon > 0$  take a set  $\{\epsilon_i > 0 : i \in \mathbb{N}\}$  such that  $\sum_{i \in \mathbb{N}} \epsilon_i \leq \epsilon$ . Then by the definition 1.6 there exists a family of function  $\{F_i : B_i \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n, i \in \mathbb{N}\}$  such that  $\forall i \in \mathbb{N} \quad A_i \subseteq F_i(B_i)$ ,  $\forall b, b' \in B_i \quad |F_i(b) - F_i(b')|^a \leq |b - b'|$ , and  $\text{meas}(B_i) \leq m_a(A_i) + \epsilon_i$ . Let's consider a denumerable family of function  $\{f_{ij} = F_i \upharpoonright B_{ij}; B_{ij} = B_i \cap [j, j + 1], i \in \mathbb{N}, j \in \mathbb{Z}\}$ . Then from the lemma 2.5 there exists  $f : B \subseteq \mathbb{R}^1 \rightarrow \mathbb{R}^n$  with  $\bigcup_{i \in \mathbb{N}} A_i = \bigcup_{i \in \mathbb{N}} \bigcup_{j \in \mathbb{Z}} f_{ij}(B_{ij}) \subseteq f(B)$  such that  $\forall b, b' \in B \quad |f(b) - f(b')|^a \leq |b - b'|$  and  $\text{meas}(B) \leq \sum_{i \in \mathbb{N}} \sum_{j \in \mathbb{Z}} \text{meas}(B_{ij}) \leq \sum_{i \in \mathbb{N}} \text{meas}(B_i) \leq \sum_{i \in \mathbb{N}} m_a(A_i) + \epsilon$ . And because it is true for any  $\epsilon > 0$ , by the definition 1.6 we have that  $m_a(\bigcup_{i \in \mathbb{N}} A_i) \leq \sum_{i \in \mathbb{N}} m_a(A_i)$ .  $\square$

## REFERENCES

1. Ainouline, A., Necessary and sufficient condition of the Morse-Sard Theorem for real valued functions, 2003, preprint.
2. Ainouline, A., Critical values lie on a line, 2003, preprint.
3. Bates, S.M., On the image size of singular maps II, Duke Math.J. Vol.68, No.3, December, 1992, 463-476.
4. Federer, H., Geometric Measure Theory, Springer-Verlag, 1969.
5. Marstrand, J.M., The dimension of Cartesian product sets, Proc. Cambridge Philos. Soc. (3) 50 (1954), 198-202.
6. Mattila, P., Geometry of sets and measures in Euclidean spaces,.