

# EQUIDISTRIBUTION OF SMALL SUBVARIETIES OF AN ABELIAN VARIETY

MATTHEW BAKER AND SU-ION IH

ABSTRACT. We prove an equidistribution result for small subvarieties of an abelian variety which generalizes the Szpiro-Ullmo-Zhang theorem on equidistribution of small points.

## 1. INTRODUCTION

1.1. **Notation.** The following notation and conventions will be used throughout this paper:

- $K$  a number field.
- $\mathcal{O}_K$  the ring of integers of  $K$ .
- $A$  an abelian variety defined over  $K$ .

We fix, for future use, a choice of an algebraic closure  $\overline{K}$  of  $K$  and an embedding of  $\overline{K}$  into  $\mathbb{C}$ .

1.2. **Heights of cycles.** Let  $X$  be a smooth projective variety over  $K$  of dimension  $N \geq 1$ , and let  $\mathcal{X}$  be a model for  $X$  over  $\mathcal{O}_K$ , i.e., an integral scheme projective and flat over  $\text{Spec } \mathcal{O}_K$  whose generic fiber is  $X$ .

Let  $\overline{\mathcal{L}}$  be a hermitian line bundle on  $\mathcal{X}$ . A hermitian metric is always assumed to be smooth and invariant under complex conjugation. We assume furthermore that  $\mathcal{L}_K$  is ample, and that the curvature form  $c_1(\overline{\mathcal{L}})$  satisfies  $c_1(\overline{\mathcal{L}}) > 0$ . (See [3] for a discussion of the curvature form associated to a hermitian line bundle).

By using arithmetic intersection theory, one defines the height of a cycle in Arakelov geometry as follows (see e.g. [5]):

**Definition.** The *height* of a nonzero effective cycle  $Y$  (of pure dimension) of  $X$  with respect to  $\overline{\mathcal{L}}$  is

$$h_{\overline{\mathcal{L}}}(Y) := \frac{\widehat{c}_1(\overline{\mathcal{L}}|_{\overline{Y}})^{\dim Y + 1}}{(\dim Y + 1)c_1(\mathcal{L}|_Y)^{\dim Y}},$$

where  $\overline{Y}$  is the (scheme-theoretic) Zariski closure of  $Y$  in  $\mathcal{X}$ .

For a detailed overview of all the properties of curvature forms, arithmetic Chern classes, and heights of arithmetic cycles which we will need, see [1] or [5]. Proofs of the relevant facts can be found in [3].

---

1991 *Mathematics Subject Classification.* 11G10, 11G35, 11G50, 14G05, 14G40.

*Key words and phrases.* Equidistribution, Bogomolov conjecture, height, torsion subvariety. The first author's research was supported in part by NSF Research Grant DMS-0300784.

**1.3. Canonical heights on abelian varieties.** Let  $A/K$  be an abelian variety. Using the choice of an embedding of  $\overline{K}$  into  $\mathbb{C}$ , we view  $A(\overline{K})$  as a subset of  $A(\mathbb{C})$ . Let  $\mathcal{A}$  be a model for  $A$  over  $\text{Spec } \mathcal{O}_K$ . Let  $\overline{\mathcal{L}}$  be a hermitian line bundle on  $\mathcal{A}$  such that  $L := \mathcal{L}_K$  is symmetric and ample, and such that  $\mathcal{L}$  is equipped with the *cubical metric* (see [4]).

Fix a nontrivial multiplication map on  $A$  (e.g. multiplication by 2). One can then construct from  $(\mathcal{A}, \overline{\mathcal{L}})$  a sequence  $(\mathcal{A}_n, \overline{\mathcal{L}}_n)_{n \geq 1}$  of models of  $(A, L)$ , where each  $\mathcal{L}_n$  is equipped with the cubical metric, in such a way that the sequence

$$h_{\overline{\mathcal{L}}_n}(Y) = \frac{\widehat{c}_1(\overline{\mathcal{L}}_n|_{\overline{Y}_n})^{\dim Y + 1}}{(\dim Y + 1)c_1(L|_Y)^{\dim Y}}$$

converges (uniformly in  $Y$ ) to a nonnegative real number  $\hat{h}_L(Y)$ . (Here  $Y$  is a nonzero effective cycle of pure dimension on  $A$ , and  $\overline{Y}_n$  is the Zariski closure of  $Y$  in  $\mathcal{A}_n$ . See [5] or [8] for details.) Note that  $\hat{h}_L$  does not depend on the choice of  $(\mathcal{A}, \mathcal{L})$  or a sequence of models  $(\mathcal{A}_n, \mathcal{L}_n)_{n \geq 1}$ . For  $x \in A(\overline{K})$ ,  $\hat{h}_L(x)$  is the Néron-Tate canonical height of  $x$  with respect to  $L$ .

**1.4. Statement of the main theorem.** We need several definitions in order to state our main result. By a *variety*  $X$  over a field  $k$ , we mean an integral separated scheme of finite type over  $k$ . By a *subvariety* of  $X$ , we mean an integral closed subscheme of  $X$ .

**Definition.** A *torsion subvariety* of  $A$  is a translate of an abelian subvariety of  $A$  by a torsion point.

**Definition.** A sequence  $(X_n)_{n \geq 1}$  of closed subvarieties of  $A$  is *small* if  $\hat{h}_L(X_n) \rightarrow 0$  (as  $n \rightarrow \infty$ ).

**Definition.** A sequence  $(X_n)_{n \geq 1}$  of closed subvarieties of  $X$  is *generic* if it has no subsequence contained in a proper Zariski closed subset of  $X$ .

**Definition.** A sequence  $(X_n)_{n \geq 1}$  of closed subvarieties of  $A$  is *strict* if it has no subsequence contained in a proper torsion subvariety of  $A$ .

Note that the subvarieties  $X_n$  are required to be defined over  $\overline{K}$ , but not necessarily over  $K$ .

The following result is a generalization of the Szpiro-Ullmo-Zhang/Ullmo/Zhang equidistribution theorem to sequences of small subvarieties of an abelian variety.

**Theorem 1.1 (Strict Equidistribution).** *Let  $A/K$  be an abelian variety, let  $L$  be a symmetric ample line bundle on  $A$ , and let  $\overline{L}$  denote  $L$  with the cubical metric. Let  $(X_n)_{n \geq 1}$  be a small strict sequence of closed subvarieties of  $A$ . Then for every real-valued continuous function  $f$  on  $A(\mathbb{C})$ , we have*

$$\int_{A(\mathbb{C})} f \mu_n \longrightarrow \int_{A(\mathbb{C})} f \mu,$$

as  $n \rightarrow \infty$ , where setting  $d_n = \dim X_n$  and  $g = \dim A$ , we have  $\mu_n = \frac{1}{c_1(L|_{X_n})^{d_n}} c_1(\overline{L})^{d_n} \delta_{X_n}$  and  $\mu = \frac{c_1(\overline{L})^g}{c_1(L)^g}$ .

**Remarks.**

1. The first integral is the integral of  $f$  against the restriction of  $c_1(\overline{\mathcal{L}})^{d_n}/\deg_L(X_n)$  to  $X_n(\mathbb{C})$ . The second integral is the integral of  $f$  with respect to the Haar measure  $\mu$  on  $A(\mathbb{C})$ , normalized to have total mass 1.

2. If  $X_n = x_n$  is a point, i.e., if  $d_n = 0$ , note that

$$\int_{A(\mathbb{C})} f(x)\mu_n = \frac{1}{\#O(x_n)} \sum_{x \in O(x_n)} f(x),$$

where  $O(x_n)$  is the orbit of  $x_n$  under the action of  $\text{Gal}(\overline{K}/K)$ .

3. For notational convenience, we write

$$\mu_n \xrightarrow{w} \mu \text{ as } n \rightarrow \infty,$$

and say the sequence  $(\mu_n)_{n \geq 1}$  of measures *weakly converges* to  $\mu$ , if

$$\int_{A(\mathbb{C})} f \mu_n \rightarrow \int_{A(\mathbb{C})} f \mu$$

for every continuous function  $f : A(\mathbb{C}) \rightarrow \mathbb{R}$ . In this case, we say that the  $X_n$ 's are *equidistributed* with respect to  $\mu$ .

4. To get a feeling for what Theorem 1.1 says, consider the following simple example. Let  $E$  be an elliptic curve defined over  $\mathbb{Q}$  and let  $A = E \times E$ . For each  $n \geq 1$ , let  $E_n \subset A$  be the graph of the multiplication-by- $n$  map on  $E$ . Then each  $E_n$  is a torsion subvariety of  $A$  defined over  $\mathbb{Q}$  (in fact,  $E_n$  is  $\mathbb{Q}$ -isogenous to  $E$ ).

It is easy to see that  $\deg(E_n) \rightarrow \infty$  as  $n \rightarrow \infty$ , and that  $\bigcup_{n \geq 1} E_n$  is Zariski dense in  $A$ . Theorem 1.1 says something stronger than this, namely that as  $n \rightarrow \infty$ , the normalized Haar measure on  $E_n$  approximates the normalized Haar measure on  $A$  arbitrarily closely.

5. For related equidistribution results, see Theorem 1.1 of [2], Theorem 4.1 of [5], Theorem 2.3 of [6], and Theorem 1.1 of [9]. In addition, Pascal Autissier has recently obtained a proof of Theorem 2.2 of the present paper independently of the authors.

## 2. GENERIC EQUIDISTRIBUTION

Let  $X$  be a closed subvariety of dimension  $N \geq 1$  of  $A$ . The following result is a special case of Zhang's "Theorem of the successive minima" (see [9] for details):

**Theorem 2.1.** *Define*

$$\lambda_1(X) := \sup_Z \inf_{x \in X-Z} \hat{h}_L(x),$$

where  $Z$  runs over the set of all proper closed subsets of  $X$ . Then

$$\lambda_1(X) \geq \hat{h}_L(X) \geq \frac{1}{N+1} \lambda_1(X).$$

**Definition.** Let  $\overline{\mathcal{L}}$  be a hermitian line bundle on  $\mathcal{A}$ . If  $f$  is a real-valued  $C^\infty$ -function on  $A(\mathbb{C})$ , define

$$\overline{\mathcal{L}}(f) := \overline{\mathcal{L}} \otimes (\mathcal{O}_{\mathcal{A}}, e^{-f})$$

to be the tensor product of  $\overline{L}$  with the trivial bundle, endowed with the metric given by  $\|1\|(P) = e^{-f(P)}$ .

**Theorem 2.2 (Generic Equidistribution).** *Let  $A/K$  be an abelian variety, and let  $L$  be a symmetric ample line bundle on  $A$ . Let  $(X_n)_{n \geq 1}$  be a small generic sequence of closed subvarieties of  $A$ . Then, for every real-valued continuous function  $f$  on  $A(\mathbb{C})$ , we have*

$$\int_{A(\mathbb{C})} f(x) \mu_n \longrightarrow \int_{A(\mathbb{C})} f(x) \mu$$

as  $n \rightarrow \infty$ , where  $d_n = \dim X_n$ ,  $\mu_n = \frac{1}{c_1(L|_{X_n})^{d_n}} c_1(\overline{L})^{d_n} \delta_{X_n}$ ,  $g = \dim A$ ,  $\mu = \frac{c_1(\overline{L})^g}{c_1(L)^g}$ , and  $\overline{L}$  is  $L$  with the cubical metric.

*Proof.* Enumerate the countably many subvarieties  $(Z_n)_{n \geq 1}$  of  $A$  defined over  $\overline{K}$ . Since  $(X_n)_{n \geq 1}$  is generic, we may assume, without loss of generality,  $X_n \not\subset Z_1 \cup \dots \cup Z_n$ . By the definition of  $\lambda_1(X_n)$ , we can find (for each  $n \geq 1$ ) an infinite sequence  $(x_{n,m})_{m \geq 1}$  in  $X_n$  such that:

- (i) for each  $m \geq 1$ ,  $x_{n,m} \notin \bigcup_{1 \leq i \leq n} Z_i$ ;
- (ii)  $|\hat{h}_L(x_{n,m}) - \lambda_1(X_n)| < \frac{1}{n}$  for all  $m \geq 1$ ; and
- (iii) for each  $n \geq 1$ ,  $\lim_{m \rightarrow \infty} \hat{h}_L(x_{n,m}) = \lambda_1(X_n)$ .

By choosing a bijection between  $\mathbf{N}^2$  and  $\mathbf{N}$ , we may consider the doubly-indexed sequence  $(x_{n,m})$  as a sequence indexed by the natural numbers. Property (i) guarantees that the resulting sequence  $(x_{n,m})$  is generic. Furthermore, since  $\hat{h}_L(X_n) \rightarrow 0$  by assumption, it follows from the theorem of the successive minima that  $\lambda_1(X_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Using this observation, properties (ii) and (iii) easily imply that  $\lim_{n,m \rightarrow \infty} \hat{h}_L(x_{n,m}) = 0$ , i.e., that the sequence  $(x_{n,m})$  is small.

Define

$$\alpha_{n,m} := \frac{1}{\#O(x_{n,m})} \sum_{x \in O(x_{n,m})} f(x),$$

where  $O(x_{n,m})$  is the orbit of  $x_{n,m}$  under the action of  $\text{Gal}(\overline{K}/K)$ . By choosing a subsequence of  $(x_{n,m})_{m \geq 1}$  if necessary, we may assume that  $\lim_{m \rightarrow \infty} \alpha_{n,m}$  exists for all  $n \geq 1$ . Note that every subsequence of a small (resp. generic) sequence is small (resp. generic).

Approximating  $f$  by  $C^\infty$ -functions if necessary, we may assume that  $f$  is a  $C^\infty$ -function. Let  $\lambda > 0$  be a real number. Note that  $c_1(\overline{L}_l(\lambda f)) > 0$  if  $\lambda > 0$  is small enough. We then note, for  $l \geq 1$ , that

$$\begin{aligned} h_{\overline{L}_l(\lambda f)}(x_{n,m}) &= h_{\overline{L}_l}(x_{n,m}) + \lambda \alpha_{n,m}; \quad \text{and} \\ \liminf_{m \rightarrow \infty} h_{\overline{L}_l(\lambda f)}(x_{n,m}) &\geq h_{\overline{L}_l(\lambda f)}(X_n) \quad [5], \text{ Proposition 2.1} \\ &= h_{\overline{L}_l}(X_n) + \lambda \int_{A(\mathbb{C})} f(x) \mu_n + O(\lambda^2), \end{aligned}$$

where the last equality follows from [1, Proof of Proposition 2.9]. Here the  $O$ -constant is independent of  $l$  and  $n$ , and  $\lambda > 0$  is sufficiently small.

Fix  $n \geq 1$  and  $\varepsilon > 0$ . Then for  $m$  sufficiently large, we have:

$$h_{\overline{\mathcal{L}}_l}(x_{n,m}) + \lambda \alpha_{n,m} \geq h_{\overline{\mathcal{L}}_l}(X_n) + \lambda \int_{A(\mathbb{C})} f(x) \mu_n + O(\lambda^2) - \varepsilon.$$

Letting  $l \rightarrow \infty$ , we have:

$$\hat{h}_L(x_{n,m}) - \hat{h}_L(X_n) + \lambda \alpha_{n,m} \geq \lambda \int_{A(\mathbb{C})} f(x) \mu_n + O(\lambda^2) - \varepsilon.$$

Now let  $m \rightarrow \infty$ , and we obtain (since  $\varepsilon > 0$  is arbitrary):

$$(1) \quad \lambda_1(X_n) - \hat{h}_L(X_n) + \lambda \lim_{m \rightarrow \infty} \alpha_{n,m} \geq \lambda \int_{A(\mathbb{C})} f(x) \mu_n + O(\lambda^2).$$

On the other hand, the Szpiro-Ullmo-Zhang/Ullmo/Zhang equidistribution theorem ([5], [6], and [9]), applied to the small generic sequence  $(x_{n,m})$ , implies that  $\lim_{n,m \rightarrow \infty} \alpha_{n,m}$  exists, and that

$$(2) \quad \lim_{n,m \rightarrow \infty} \alpha_{n,m} = \int_{A(\mathbb{C})} f(x) \mu.$$

Taking  $\limsup_{n \rightarrow \infty}$  in (1), we have:

$$\lambda \lim_{n,m \rightarrow \infty} \alpha_{n,m} \geq \lambda \limsup_{n \rightarrow \infty} \int_{A(\mathbb{C})} f(x) \mu_n + O(\lambda^2).$$

Now divide both sides by  $\lambda > 0$ , and let  $\lambda \rightarrow 0^+$ . We obtain:

$$(3) \quad \lim_{n,m \rightarrow \infty} \alpha_{n,m} \geq \limsup_{n \rightarrow \infty} \int_{A(\mathbb{C})} f(x) \mu_n.$$

Replacing  $f$  by  $-f$ , we see that:

$$(4) \quad \lim_{n,m \rightarrow \infty} \alpha_{n,m} \leq \liminf_{n \rightarrow \infty} \int_{A(\mathbb{C})} f(x) \mu_n.$$

It then follows from (3) and (4) that  $\lim_{n \rightarrow \infty} \int_{A(\mathbb{C})} f(x) \mu_n$  exists and that

$$\lim_{n \rightarrow \infty} \int_{A(\mathbb{C})} f(x) \mu_n = \lim_{n,m \rightarrow \infty} \alpha_{n,m}.$$

We conclude from (2) that

$$\lim_{n \rightarrow \infty} \int_{A(\mathbb{C})} f(x) \mu_n = \int_{A(\mathbb{C})} f(x) \mu,$$

as desired. □

## 3. STRICT EQUIDISTRIBUTION

The following result is a consequence of two results of Zhang: the generalized Bogomolov conjecture (see [9]) and the theorem of the successive minima. The proof is similar to the proof of Theorem 2.2.

**Theorem 3.1.** *Let  $X$  be a nontorsion subvariety of  $A$ . Then there is an  $\epsilon > 0$  such that the set*

$$\bigcup \{Y : Y \text{ is a closed subvariety of } X \text{ such that } \hat{h}_L(Y) \leq \epsilon\}$$

*is not Zariski dense in  $X$ .*

*Proof.* Suppose, for the sake of contradiction, that  $(Y_n)_{n \geq 1}$  is a sequence of distinct closed subvarieties of  $X$  which is small (i.e.,  $\hat{h}_L(Y_n) \rightarrow 0$ ) and generic in  $X$  (i.e., no subsequence is contained in a proper Zariski closed subset of  $X$ ). Then, proceeding as in the proof of Theorem 2.2, we can construct an infinite sequence  $(y_k)_{k \geq 1}$  of points in  $X$  such that  $\{y_k \in X : k \geq 1\}$  is Zariski dense in  $X$  and  $\hat{h}_L(y_k) \rightarrow 0$ . But Corollary 3 of [9] then implies that  $X$  is a torsion subvariety of  $A$ , a contradiction.  $\square$

Now we are ready to prove Theorem 1.1 (Strict Equidistribution Theorem).

*Proof* (of Theorem 1.1). By Theorem 2.2, it suffices to show that the small and strict sequence  $(X_n)_{n \geq 1}$  is generic. Let  $X'$  be the Zariski closure of  $\bigcup_k X_{n_k}$  for any subsequence  $(X_{n_k})_{k \geq 1}$  of  $(X_n)_{n \geq 1}$ . By Theorem 3.1,  $X'$  must be a torsion subvariety of  $A$ . Since  $(X_n)_{n \geq 1}$  is strict, it follows that  $X' = A$ , so that  $(X_n)_{n \geq 1}$  is generic as desired.  $\square$

## REFERENCES

1. A. Abbes, *Hauteurs et discrétude (d'après L. Szpiro, E. Ullmo et S. Zhang)*, (French) [Heights and discreteness (after L. Szpiro, E. Ullmo and S. Zhang)] Séminaire Bourbaki, Vol. 1996/97. Astérisque No. **245** (1997), Exp. No. **825**, 4, 141–166.
2. Y. Bilu, *Limit distribution of small points on algebraic tori*, Duke Math. J. **89** (1997), no. 3, 465–476.
3. H. Gillet, and C. Soulé, *Arithmetic intersection theory*, Inst. Hautes Études Sci. Publ. Math. No. **72** (1990), 93–174.
4. L. Moret-Bailly, *Métriques permises* (French) [Admissible metrics] Séminaire sur les pinceaux de courbes elliptiques, L. Szpiro (ed.), Astérisque No. **127** (1985), 29–87.
5. L. Szpiro, E. Ullmo, and S. Zhang, *Équirépartition des petits points* (French), [Uniform distribution of small points] Invent. Math. **127** (1997), no. 2, 337–347.
6. E. Ullmo, *Positivité et discrétion des points algébriques des courbes* (French), [Positivity and discreteness of algebraic points of curves] Ann. of Math. (2) **147** (1998), no. 1, 167–179.
7. S. Zhang, *Positive line bundles on arithmetic varieties*, J. Amer. Math. Soc. **8** (1995), no. 1, 187–221.
8. ———, *Small points and adelic metrics*, J. Algebraic Geom. **4** (1995), no. 2, 281–300.
9. ———, *Equidistribution of small points on abelian varieties*, Ann. of Math. (2) **147** (1998), no. 1, 159–165.