

Resolutions of Segre embeddings of projective spaces of any dimension

Elena Rubei

Abstract. This paper deals with syzygies of the ideals of the Segre embeddings. Let $d \geq 3$ and $n_1, \dots, n_d \in \mathbf{N} - \{0\}$. We prove that $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies Green-Lazarsfeld's Property N_p if and only if $p \leq 3$.

1 Introduction

Let L be a very ample line bundle on a smooth complex projective variety Y and let $\varphi_L : Y \rightarrow \mathbf{P}(H^0(Y, L)^*)$ be the map associated to L . We recall the definition of Property N_p of Green-Lazarsfeld, studied for the first time by Green in [8] (see also [10], [9]):

let Y be a smooth complex projective variety and let L be a very ample line bundle on Y defining an embedding $\varphi_L : Y \hookrightarrow \mathbf{P} = \mathbf{P}(H^0(Y, L)^)$; set $S = S(L) = \bigoplus_n \text{Sym}^n H^0(L)$, the homogeneous coordinate ring of the projective space \mathbf{P} , and consider the graded S -module $G = G(L) = \bigoplus_n H^0(Y, L^n)$; let E_**

$$0 \longrightarrow E_l \longrightarrow E_{l-1} \longrightarrow \dots \longrightarrow E_0 \longrightarrow G \longrightarrow 0$$

be a minimal graded free resolution of G ; the line bundle L satisfies Property N_p ($p \in \mathbf{N}$) iff

$$\begin{aligned} E_0 &= S \\ E_i &= \bigoplus S(-i-1) \quad \text{for } 1 \leq i \leq p. \end{aligned}$$

(Thus L satisfies Property N_0 iff $Y \subset \mathbf{P}(H^0(L)^*)$ is projectively normal; L satisfies Property N_1 iff L satisfies N_0 and the homogeneous ideal I of $Y \subset \mathbf{P}(H^0(L)^*)$ is generated by quadrics; L satisfies N_2 iff L satisfies N_1 and the module of syzygies among quadratic generators $Q_i \in I$ is spanned by relations of the form $\sum L_i Q_i = 0$, where L_i are linear polynomials; and so on.)

In this paper we will consider the case of Segre embedding i.e. the case $Y = \mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}$, $L = \mathcal{O}(1, \dots, 1)$.

Among the papers on syzygies of Segre-Verones embeddings we quote [1], [8], [13], [11], [7], [12], [14], [15], [16], [17].

In the first one, the authors examined the cases in which the resolution is “pure” i.e. the minimal generators of each module of syzygies have the same degree. We quote the following results from the other papers:

Case $d = 1$, i.e. the case of the Veronese embedding:

Theorem 1 (Green) [8]. *Let a be a positive integer. The line bundle $\mathcal{O}_{\mathbf{P}^n}(a)$ satisfies Property N_{a+1} .*

Theorem 2 (Schenck-Smith) [17]. *Let a be a positive integer. The line bundle $\mathcal{O}_{\mathbf{P}^n}(a)$ satisfies Property N_{a+1} .*

The case $\mathcal{O}(3)$ satisfies N_4 is also proved by **Rubei** in [16].

Theorem 3 (Ottaviani-Paoletti) [13]. *If $n \geq 2$, $a \geq 3$ and the bundle $\mathcal{O}_{\mathbf{P}^n}(a)$ satisfies Property N_p , then $p \leq 3a - 3$.*

Theorem 4 (Josefiak-Pragacz-Weyman) [11]. *The bundle $\mathcal{O}_{\mathbf{P}^n}(2)$ satisfies Property N_p if and only if $p \leq 5$ when $n \geq 3$ and for all p when $n = 2$.*

(See [13] for a more complete bibliography.)

Case $d = 2$:

Theorem 5 (Gallego-Purnapranja) [7]. *Let $a, b \geq 2$. The line bundle $\mathcal{O}_{\mathbf{P}^1 \times \mathbf{P}^1}(a, b)$ satisfies Property N_p if and only if $p \leq 2a + 2b - 3$.*

Theorem 6 (Lascoux-Pragacz-Weymann) [12], [14]. *Let $n_1, n_2 \geq 2$. The line bundle $\mathcal{O}_{\mathbf{P}^{n_1} \times \mathbf{P}^{n_2}}(1, 1)$ satisfies Property N_p if and only if $p \leq 3$.*

As to the case $d \geq 3$, in [15] we proved

Theorem 7 (Rubei) . *Let $d \geq 3$. The line bundle $\mathcal{O}_{\mathbf{P}^1 \times \dots \times \mathbf{P}^1}(1, \dots, 1)$ (d times) satisfies Property N_p if and only if $p \leq 3$.*

Here we prove

Theorem 8 . *Let $d \geq 3$ and $n_1, \dots, n_d \in \mathbf{N} - \{0\}$. The line bundle $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies Property N_p if and only if $p \leq 3$.*

To prove it we use Theorem 7. As in the proof of it, we use a topological approach, but the problems that arise in proving Theorem 8 are more similar to the problems that arise if we try to solve the open problem of the syzygies of the Veronese embeddings.

2 A preliminary proposition

Proposition 9 . *Let $d \geq 3$ and $n_1, \dots, n_d \in \mathbf{N}$ with $n_i \geq p$ for some i . The line bundle $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_p if and only if $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_i-1} \times \mathbf{P}^p \times \mathbf{P}^{n_i+1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_p .*

Proof. Let L be a very ample line bundle on a smooth complex projective variety Y . We recall from [9] that L satisfies N_p iff

$$(\mathrm{Tor}_p^{S(L)}(G(L), \mathbf{C}))_{p+q} = 0 \quad \forall q \geq 2$$

(see Introduction for the notation) and $(\mathrm{Tor}_p^{S(L)}(G(L), \mathbf{C}))_{p+q}$ is equal to the homology of the Koszul complex

$$\wedge^{p+1} H^0(L) \otimes H^0(L^{q-1}) \rightarrow \wedge^p H^0(L) \otimes H^0(L^q) \rightarrow \wedge^{p-1} H^0(L) \otimes H^0(L^{q+1})$$

Now let $Y = \mathbf{P}(V_1) \times \dots \times \mathbf{P}(V_d)$ and $L = \mathcal{O}(1, \dots, 1)$. In our case $(\text{Tor}_p^{S(L)}(G(L), \mathbf{C}))_{p+q}$ is equal to the homology of

$$\begin{array}{l}
\wedge^{p+1}(V_1 \otimes \dots \otimes V_d) \otimes \text{Sym}^{(q-1)}V_1 \otimes \dots \otimes \text{Sym}^{(q-1)}V_d; \quad \alpha_{p+1, q-1}^{V_1 \times \dots \times V_d} \\
(*_{V_1, \dots, V_d}) \quad \wedge^p(V_1 \otimes \dots \otimes V_d) \otimes \text{Sym}^qV_1 \otimes \dots \otimes \text{Sym}^qV_d; \quad \alpha_{p, q}^{V_1 \times \dots \times V_d} \\
\wedge^{p-1}(V_1 \otimes \dots \otimes V_d) \otimes \text{Sym}^{(q+1)}V_1 \otimes \dots \otimes \text{Sym}^{(q+1)}V_d; \quad \alpha_{p-1, q+1}^{V_1 \times \dots \times V_d}
\end{array}$$

since the maps are $GL(V_i)$ -invariant, $(\text{Tor}_p^{S(L)}(G(L), \mathbf{C}))_{p+q}$ is a $GL(V_i)$ -module, see Rem. of §2 of [8] and Prop. 1.8 [13]. The Young diagrams of the irreducible subrepresentations of the $GL(V_i)$ -module $\wedge^p(V_1 \otimes \dots \otimes V_d) \otimes \text{Sym}^qV_1 \otimes \dots \otimes \text{Sym}^qV_d$ have at most $p+1$ rows (see for instance p.80 [6]). Thus the Young diagrams of the irreducible subrepresentation of $(\text{Tor}_p^{S(L)}(G(L), \mathbf{C}))_{p+q}$ have at most $p+1$ rows and one can easily prove that these Young diagrams are the same for V_1, \dots, V_d and for $V_1, \dots, V_{i-1}, V'_i, V_{i+1}, \dots, V_d$ with $\dim V'_i = p+1$ (by considering an injective map $V'_i \rightarrow V_i$ and the induced commutative diagram with second row $(*_{V_1, \dots, V_d})$ and first row $(*_{V_1, \dots, V_{i-1}, V'_i, V_{i+1}, \dots, V_d})$ seeing every term of the two rows as $GL(V_i)$ and $GL(V'_i)$ representations respectively).

Thus, since the Young diagrams of the irreducible subrepresentations of $(\text{Tor}_p^{S(L)}(G(L), \mathbf{C}))_{p+q}$ have at most $p+1$ rows, we have that if these representations are zero for $\dim(V_i) = p+1$ they are zero also for $\dim(V_i) \geq p+1$. □

3 Recalls on syzygies of toric ideals and on syzygies on the product of two varieties

We recall some facts on toric ideals from [19]. Let $k \in \mathbf{N}$. Let $A = \{a_1, \dots, a_m\} \subset \mathbf{Z}^k$. The toric ideal \mathcal{I}_A is defined as the ideal in $\mathbf{C}[x_1, \dots, x_m]$ generated as vector space by the binomials

$$x_1^{u_1} \dots x_m^{u_m} - x_1^{v_1} \dots x_m^{v_m}$$

for $(u_1, \dots, u_m), (v_1, \dots, v_m) \in \mathbf{N}^m$, with $\sum_{i=1, \dots, m} u_i a_i = \sum_{i=1, \dots, m} v_i a_i$.

We have that \mathcal{I}_A is homogeneous iff $\exists \omega \in \mathbf{Q}^k$ s.t. $\omega \cdot a_i = 1 \forall i = 1, \dots, m$; the rings $\mathbf{C}[x_1, \dots, x_m]$ and $\mathbf{C}[x_1, \dots, x_m]/\mathcal{I}_A$ are multigraded by $\mathbf{N}A$ via $\deg x_i = a_i$; the element $x_1^{u_1} \dots x_m^{u_m}$ has multidegree $b = \sum_i u_i a_i \in \mathbf{N}A$ and degree $\sum_i u_i = b \cdot \omega$; we define $\deg b = b \cdot \omega$. For each $b \in \mathbf{N}A$, let Δ_b be the simplicial complex (see [18]) on the set A defined as follows:

$$\Delta_b = \{\langle F \rangle \mid F \subset A : b - \sum_{a \in F} a \in \mathbf{N}A\}$$

The following theorem studies the syzygies of the ideal \mathcal{I}_A ; it was proved by Campillo and Marijuan for $k = 1$ in [3] and by Campillo and Pison for general k and $j = 0$ in [4]; the following more general statement is due to Sturmfels (Theorem 12.12 p.120 in [19]).

Theorem 10 (see [19], [3], [4]). *Let $A = \{a_1, \dots, a_m\} \subset \mathbf{N}^k$ and \mathcal{I}_A be the associated toric ideal. Let $0 \rightarrow E_n \rightarrow \dots \rightarrow E_1 \rightarrow E_0 \rightarrow G \rightarrow 0$ be a minimal free resolution of $G = \mathbf{C}[x_1, \dots, x_m]/\mathcal{I}_A$ on $\mathbf{C}[x_1, \dots, x_m]$. Each of the generators of E_j has a unique*

multidegree. The number of the generators of multidegree $b \in \mathbf{NA}$ of E_{j+1} equals the rank of the j -th reduced homology group $\tilde{H}_j(\Delta_b, \mathbf{C})$.

Finally we recall the following proposition proved in [15]

Proposition 11 (Rubei) . Let X and Y be two projective varieties and let L be a line bundle on X and M a line bundle on Y . Let $\pi_X : X \times Y \rightarrow X$ and $\pi_Y : X \times Y \rightarrow Y$ be the canonical projections. Suppose L and M satisfy Property N_1 . Let $p \geq 2$. If L does not satisfy Property N_p , then $\pi_X^* L \otimes \pi_Y^* M$ does not satisfy Property N_p , either.

4 Proof of Theorem 8

Notation 12 • homologous means homologous in the reduced homology.

- \sim_A means homologous in A , i.e. $\gamma \sim_A \gamma'$ means that $\exists \beta$ chain in A s.t. $\partial\beta = \gamma - \gamma'$.
- e_i denotes the i -th element of the canonical basis of \mathbf{R}^n .
- The symbol $*$ denotes the joining.
- For any $v \in \mathbf{R}^n$ v_i denotes the i -th coordinate, that is the lower index denotes the coordinate.

If we take $A = A_{n_1, \dots, n_d} = \{(x_1^0, \dots, x_1^{n_1}, \dots, x_d^0, \dots, x_d^{n_d}) \mid \sum_{j=1, \dots, n_i} x_i^j = 1, x_i^j \in \mathbf{N}\}$, we have that $\mathcal{I}_{A_{n_1, \dots, n_d}}$ is the ideal of the embedding of $\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}$ by $\mathcal{O}(1, \dots, 1)$. In this case $\omega = \omega_d = \frac{1}{d}(1, \dots, 1)$.

Let $b \in \mathbf{NA}_{n_1, \dots, n_d}$; we have that $\deg b = (= b \cdot \omega) = k$ iff b is the sum of k (not necessarily distinct) elements of A_{n_1, \dots, n_d} . Observe that a simplex S with vertices in A_{n_1, \dots, n_d} is a simplex of Δ_b iff the sum s of the vertices of S is s.t. $s_i \leq b_i \forall i = 1, \dots, n+1$. We generalize the definition of the simplicial complex Δ_b in §3 in the following way:

Notation 13 Let $v \in \mathbf{N}^{n+1}$. Let Δ_v be the following simplicial complex: a simplex S with vertices in A_{n_1, \dots, n_d} is a simplex of Δ_v iff the sum s of the vertices of S is s.t. $s_i \leq v_i \forall i = 1, \dots, n+1$.

The main points of the proof of the Thm. 8 are Propositions 15 and 16.

Notation 14 Let $d, n_1, \dots, n_d \in \mathbf{N} - \{0\}$ and $b \in \mathbf{NA}_{n_1, \dots, n_d}$. Let X_b be the following simplicial complex on A_{n_1, \dots, n_d} :

$$X_b := \Delta_b \cup \Delta \begin{pmatrix} b_1 - 1 \\ b_2 + 1 \\ b_3 \\ \vdots \\ \vdots \end{pmatrix} \cup \dots \cup \Delta \begin{pmatrix} 0 \\ b_2 + b_1 \\ b_3 \\ \vdots \\ \vdots \end{pmatrix}$$

(in the obvious sense that a simplex with vertices in A_{n_1, \dots, n_d} is a simplex of X_b iff it is a simplex of $\Delta_{b - ke_1 + ke_2}$ for some $k \in \{0, \dots, b_1\}$)

Proposition 15 Let $d, n_1, \dots, n_d, p \in \mathbf{N} - \{0\}$. Let $b \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg(b) \geq p+2$. Let γ be a $(p-1)$ -cycle in Δ_b . If the following conditions hold

- a) $\tilde{H}_{p-3}(\Delta_{c-e_1}) = 0 \forall c \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg(c) \geq p+1$ if $p \geq 3$
- b) $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies Property N_{p-1} ,

then $\exists \gamma'$ cycle in $\Delta \begin{pmatrix} 0 \\ b_1 + b_2 \\ b_3 \\ \vdots \end{pmatrix}$ s.t. $\gamma \sim_{X_b} \gamma'$.

Proposition 16 Let $d, n_1, \dots, n_d \in \mathbf{N} - \{0\}$ and $p \in \{2, 3\}$. Let $b \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg(b) \geq p + 2$. Let γ be a $(p - 1)$ -cycle in Δ_b . If $\gamma \sim_{X_b} 0$ then $\gamma \sim_{\Delta_b} 0$.

Lemma 17 Let $d, n_1, \dots, n_d \in \mathbf{N} - \{0\}$ and $c \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg(c) \geq 4$. Then $\tilde{H}_0(\Delta_{c-e_i}) = 0 \forall i$.

We show now how Thm. 8 follows from Propositions 15 and 16 and Lemma 17.

Proof of Thm. 8. By Prop. 9, to prove $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_3 , it is sufficient to prove our statement when $n_i \leq 3 \forall i$. We prove the statement by induction on $\sum_i n_i$; the first step of induction is given by Thm. 7. By Thm. 10, the bundle $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_p iff $\tilde{H}_{q-1}(\Delta_b) = 0 \forall b \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg b \geq q + 2 \forall q \leq p$; in particular, in order to prove that it satisfies N_3 , we have to prove that $\tilde{H}_{p-1}(\Delta_b) = 0 \forall b \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg b \geq p + 2$ for $p = 2, 3$.

We show that (\star) if $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_{p-1} then $\tilde{H}_{p-1}(\Delta_b) = 0 \forall b \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg b \geq p + 2$ for $p \in \{2, 3\}$. Let $b \in \mathbf{NA}_{n_1, \dots, n_d}$ with $\deg(b) \geq p + 2$, $p \in \{2, 3\}$. Let γ be a $(p - 1)$ -cycle in Δ_b . We want to prove $\gamma \sim_{\Delta_b} 0$. By Prop. 15, $\exists \gamma'$ cycle in $\Delta \begin{pmatrix} 0 \\ b_1 + b_2 \\ b_3 \\ \vdots \end{pmatrix}$ s.t. $\gamma \sim_{X_b} \gamma'$ (the assumption of Prop. 15 in our case is true by Lemma 17 and

by our assumption in (\star)). We have $\tilde{H}_{p-1}(\Delta \begin{pmatrix} 0 \\ b_1 + b_2 \\ b_3 \\ \vdots \end{pmatrix}) = \tilde{H}_{p-1}(\Delta \begin{pmatrix} b_1 + b_2 \\ b_3 \\ \vdots \end{pmatrix}) = 0$, where the

last equality holds since $\mathcal{O}_{\mathbf{P}^{n_1-1} \times \mathbf{P}^{n_2} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_p by induction hypothesis. Thus $\gamma' \sim 0$ in $\Delta \begin{pmatrix} 0 \\ b_1 + b_2 \\ b_3 \\ \vdots \end{pmatrix}$. Thus $\gamma \sim_{X_b} 0$ and then $\gamma \sim_{\Delta_b} 0$ by Prop. 16.

Since $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_1 then, by (\star) , it satisfies N_2 and by applying again (\star) we get that it satisfies also N_3 .

Finally by Theorem 6 and Proposition 11 we know that $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ does not satisfy N_4 if there is $n_i \geq 2$. If $n_i = 1 \forall i$ then we already know the result from Theorem 7. \square

Now we will prove Propositions 15 and 16 and Lemma 17.

Notation 18 Let $b \in \mathbf{NA}_{n_1, \dots, n_d}$ Let γ be a $(p - 1)$ -cycle in X_b .

For every vertex a in γ , let $\mathcal{S}_{a, \gamma}$ be the set of simplexes of γ with vertex a and $\mu_{a, \gamma}$ be the $(p - 2)$ -cycle s.t. $a * \mu_{a, \gamma} = \sum_{\tau \in \mathcal{S}_{a, \gamma}} \tau$. For $\tilde{a} \in A_{n_1, \dots, n_d}$, let

$$\alpha_{a, \tilde{a}, \gamma} = (a - \tilde{a}) * \mu_{a, \gamma}$$

Proof of Prop. 15. We order in some way the (finite) vertices of γ with first coordinate

$$\neq 0: a^1, \dots, a^r. \text{ Let } \tilde{a}^j = \begin{pmatrix} 0 \\ a_2^j + a_1^j \\ a_3^j \\ a_4^j \\ \vdots \end{pmatrix} \text{ for } j = 1, \dots, r.$$

Obviously $\alpha_{a^1, \tilde{a}^1, \gamma} \sim_{X_b} 0$, because $\mu_{a^1, \gamma}$ is in Δ_{b-a^1} and $\tilde{H}_{p-1}((a^1 - \tilde{a}^1) * \Delta_{b-a^1}) = \tilde{H}_{p-2}(\Delta_{b-a^1}) = 0$ (since $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies Property N_{p-1}). Thus $\gamma_1 := \gamma + \alpha_{a^1, \tilde{a}^1, \gamma}$ is homologous to γ in X_b .

We define by induction $\gamma_j := \gamma_{j-1} + \alpha_{a^j, \tilde{a}^j, \gamma_{j-1}}$ for $j = 2, \dots, r$. We want to prove $\gamma_r \sim_{X_b} 0$; to prove this, we prove $\alpha_{a^j, \tilde{a}^j, \gamma_{j-1}} \sim_{X_b} 0$ for $j = 2, \dots, r$.

Observe that $\mu_{a^j, \gamma_{j-1}}$ is in

$$\Delta_{b-a^j} \cup \Delta \begin{pmatrix} (b-a^j)_1 - 1 \\ (b-a^j)_2 + 1 \\ (b-a^j)_3 \\ \vdots \end{pmatrix} \cup \dots \cup \Delta \begin{pmatrix} 0 \\ (b-a^j)_1 + (b-a^j)_2 \\ (b-a^j)_3 \\ \vdots \end{pmatrix}$$

We can find some cycles θ_ε in $\Delta_{b-a^j - \varepsilon e_1 + \varepsilon e_2}$ for $\varepsilon \in \{0, \dots, (b-a^j)_1\}$ s.t. $\mu_{a^j, \gamma_{j-1}} = \sum_{\varepsilon \in \{0, \dots, (b-a^j)_1\}} \theta_\varepsilon$, in fact: if $p = 2$ the statement is obvious; let $p \geq 3$; let σ_0 be the sum of the simplexes of $\mu_{a^j, \gamma_{j-1}}$ in Δ_{b-a^j} and not in $\Delta_{b-a^j - e_1}$; $\partial \sigma_0$ is in $\Delta_{b-a^j - e_1}$ and since $\tilde{H}_{p-3}(\Delta_{c-e_1}) = 0 \forall c$ with $\deg(c) \geq p+1$ then $\exists \sigma'_0$ in $\Delta_{b-a^j - e_1}$ s.t. $\partial \sigma'_0 = \partial \sigma_0$; let $\theta_0 = \sigma_0 - \sigma'_0$; now $\mu_{a^j, \gamma_{j-1}} - \theta_0$ is in

$$\Delta \begin{pmatrix} (b-a^j)_1 - 1 \\ (b-a^j)_2 + 1 \\ (b-a^j)_3 \\ \vdots \end{pmatrix} \cup \dots \cup \Delta \begin{pmatrix} 0 \\ (b-a^j)_1 + (b-a^j)_2 \\ (b-a^j)_3 \\ \vdots \end{pmatrix}$$

and we can go on analogously: let σ_1 be the sum of the simplexes of $\mu_{a^j, \gamma_{j-1}} - \theta_0$ in $\Delta_{b-a^j - e_1 + e_2}$ and not in $\Delta_{b-a^j - 2e_1} \dots$

Since $\deg(b-a^j - \varepsilon e_1 + \varepsilon e_2) \geq p+1$ and $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies N_{p-1} we have $\tilde{H}_{p-2}(\Delta_{b-a^j - \varepsilon e_1 + \varepsilon e_2}) = 0$ thus $\theta_\varepsilon \sim 0$ in $\Delta_{b-a^j - \varepsilon e_1 + \varepsilon e_2}$ and then $(a^j - \tilde{a}^j) * \theta_\varepsilon \sim_{X_b} 0$; therefore $\alpha_{a^j, \tilde{a}^j, \gamma_{j-1}} \sim_{X_b} 0$.

Thus we can define $\gamma' = \gamma_r$: γ' is in $\Delta \begin{pmatrix} 0 \\ b_1 + b_2 \\ b_3 \\ \vdots \end{pmatrix}$ and $\gamma' \sim_{X_b} \gamma$. □

Proof of Lemma 17. Let $P, Q \in \Delta_{c-e_i}$. Then we can find $s \in \mathbf{N}A_{n_1, \dots, n_d}$ with $\deg(s) \geq 3$ s.t. $s_i \leq c_i \forall i$ and $P, Q \in \Delta_s$; since $\mathcal{O}_{\mathbf{P}^{n_1} \times \dots \times \mathbf{P}^{n_d}}(1, \dots, 1)$ satisfies Property N_1 we have that $\tilde{H}_0(\Delta_s)$; thus there is a 1-cycle η in Δ_s and then in Δ_{c-e_i} s.t. $\partial \eta = P - Q$. □

Now we will prove Prop. 16.

Definition 19 Let $d, n_1, \dots, n_d, k \in \mathbf{N} - \{0\}$ and $\beta \in \mathbf{N}A_{n_1, \dots, n_d}$. We say that a $(k-1)$ -chain η in Δ_β is a UFO with axis $\langle a^1, \dots, a^t \rangle$ for the coordinate i (for short we will write $\text{UFO}_{t,k}^i(a^1, \dots, a^t, \Delta_\beta)$) if

$$\eta = \langle a^1, \dots, a^t \rangle * C_\eta$$

for some $(k - t - 1)$ -cycle C_η and a^1, \dots, a^t are distinct vertices in Δ_β with

$$(a^1 + \dots + a^t)_i = \beta_i \quad (a^j)_i > 0 \quad \forall j = 1, \dots, t$$

We will denote the axis $\langle a^1, \dots, a^t \rangle$ by χ_η . Observe $\partial\eta \subset \Delta_{\beta - e_i}$.
(Sometimes we will omit some index when it will be obvious.)




Fig.2 How a $UFO_{2,4}(a_r, a)$ looks like. C_η is in boldface

Lemma 20 “ $UFO_{p+1,p+1}$, $UFO_{p,p+1}$, $UFO_{1,p+1}$ ” Let $d, n_1, \dots, n_d, p \in \mathbf{N} - \{0\}$ and $r, l \in \{1, \dots, \sum_{i=1, \dots, d} (n_i + 1)\}$ with $r \neq l$. Let $\beta \in \mathbf{N}A_{n_1, \dots, n_d}$ with $\deg \beta \geq p + 2$. Let η be a $UFO_{t,p+1}^r(a^1, \dots, a^t, \Delta_\beta)$.

If $t \in \{p + 1, p, 1\}$ then $\exists \tilde{\eta}$ p -chain in $\Delta_{\beta + e_l - e_r}$ with $\partial \tilde{\eta} = \partial \eta$.

Proof. Case $t = p + 1$ Since $\deg \beta \geq p + 2$ and η is a simplex with $p + 1$ vertices (the simplex $\langle a^1, \dots, a^{p+1} \rangle$), then $\exists x \in A_{n_1, \dots, n_d}$ s.t. $x * \eta \subset \Delta_\beta$. Since $(a^1 + \dots + a^{p+1})_r = \beta_r$ then $x_r = 0$. Take $\tilde{\eta} := x * \partial \eta$.

Case $t = p$ If $\deg \beta \geq p + 3$ then $\deg(\beta - a^1 \dots - a^p) \geq 3$, therefore $\tilde{H}_0(\Delta_{\beta - a^1 \dots - a^p}) = 0$, thus $\exists \gamma$ s.t. $\partial \gamma = C_\eta$ and we can take $\tilde{\eta} = \gamma * \partial \chi_\eta$.

Thus we can suppose $\deg \beta = p + 2$. Since C_η is a 0-cycle it is sufficient to prove the statement when $C_\eta = P - Q$ for some $P, Q \in A_{n_1, \dots, n_d}$ with $P = Q + e_i - e_j$ for some i and j (in fact we can write C_η as $\sum_s (-1)^s P_s$ with P_{s+1} obtained from P_s by adding 1 to a coordinate and subtracting 1 to another coordinate). Let $x = \beta - a^1 \dots - a^p - P$ and $y = \beta - a^1 \dots - a^p - Q$. Since $(a^1 + \dots + a^p)_r = \beta_r$ we have $x_r = y_r = 0$.

Suppose first $i, j \notin \{l, r\}$. Let $z = x + e_l - e_j$. The chain $\tilde{\eta} = z * \partial \eta$ is in $\Delta_{\beta + e_l - e_r}$ and $\partial \tilde{\eta} = \partial \eta$.

Suppose now $j \notin \{l, r\}$ and $i = l$. Then $\tilde{\eta} := x * \partial \eta$ is in $\Delta_{\beta + e_l - e_r}$ and $\partial \tilde{\eta} = \partial \eta$.

Case $t = 1$ Let $\tilde{a}^1 = a^1 + e_l - e_r$; take $\tilde{\eta} = \tilde{a}^1 * C_\eta$. □

Lemma 21 Let $d, n_1, \dots, n_d, p \in \mathbf{N} - \{0\}$ and $r \in \{1, \dots, \sum_{i=1, \dots, d} (n_i + 1)\}$ Let $\beta \in \mathbf{N}A_{n_1, \dots, n_d}$ and $\deg \beta \geq p + 2$. Let η be a $UFO_{t,p+1}^r(a^1, \dots, a^t, \Delta_\beta)$. If $C_\eta = \partial \sigma$ where σ is a simplex (with $p + 2 - t$ vertices) in $\Delta_{\beta - a^1 - \dots - a^t}$, then $\exists \tilde{\eta}$ p -chain in $\Delta_{\beta - e_r}$ with $\partial \tilde{\eta} = \partial \eta$.

Proof. Since $\eta = \chi_\eta * C_\eta = \chi_\eta * \partial \sigma$, then $\partial \eta = \partial \chi_\eta * \partial \sigma = \partial(\partial \chi_\eta * \sigma)$. Take $\tilde{\eta} = \partial \chi_\eta * \sigma$. □

Lemma 22 “ $UFO_{2,4}$ ” Let $d, n_1, \dots, n_d, p \in \mathbf{N} - \{0\}$. Let $\beta \in \mathbf{N}A_{n_1, \dots, n_d}$ with $\deg \beta \geq 5$. Let η be a $UFO_{2,4}^2(a^1, a^2, \Delta_\beta)$. If $n_1 \leq 3$ then $\exists \tilde{\eta}$ 3-chain in $\Delta_{\beta + e_1 - e_2}$ with $\partial \tilde{\eta} = \partial \eta$.

Proof. Observe C_η is a 1-cycle and obviously $\beta_2 = 2$.

• *Case where at least one of the first $n_1 + 1$ coordinates of β except the second one is ≥ 2 , say $\beta_i \geq 2$.*

By Lemma 21 we can substitute a simplex of C_η , $\langle T_1, T_2 \rangle$ with $\langle T_1, P \rangle + \langle P, T_1 \rangle$ where P is s.t. $\langle P, T_1, T_2 \rangle * \langle a^1, a^2 \rangle \subset \Delta_\beta$ and then we can suppose C_η is in $\Delta_{\beta - a^1 - a^2 - e_i}$ (if $(T_1 + T_2)_i = (\beta - a^1 - a^2)_i$ we can replace $\langle T_1, T_2 \rangle$ with $\langle T_1, P \rangle + \langle P, T_1 \rangle$ where P is s.t. $\langle P, T_1, T_2 \rangle * \langle a^1, a^2 \rangle \subset \Delta_\beta$). Thus η is $\Delta_{\beta - e_i}$.

Let $\lambda := \langle a^2, \tilde{a}^1 \rangle * C_\eta$, where $\tilde{a}^1 = a^1 - e_2 + e_1$; λ is in $\Delta_{\beta - e_2 + e_1}$; thus $\partial\eta$ is homologous to $\partial(\eta + \lambda)$ in $\Delta_{\beta - e_2 + e_1}$; but $\partial(\eta + \lambda) = (a^1 - \tilde{a}^1) * C_\eta = (a^1 - \tilde{a}^2 + \tilde{a}^2 - \tilde{a}^1) * C_\eta = \partial(\langle a^1, \tilde{a}^2 \rangle + \langle \tilde{a}^2, \tilde{a}^1 \rangle) * C_\eta$ where $\tilde{a}^2 = a^2 - e_2 + e_1$; since $(\langle a^1, \tilde{a}^2 \rangle + \langle \tilde{a}^2, \tilde{a}^1 \rangle) * C_\eta \subset \Delta_{\beta - e_2 + e_1}$, we conclude.

• *Case where the first $n_1 + 1$ coordinates of β except the second one are ≤ 1 .*

In this case, since $n_1 \leq 3$ and $\deg(\beta) \geq 5$, we have that all the first $n_1 + 1$ coordinates except the second one are equal to 1 (and $n_1 = 3$ and $\deg(\beta) = 5$).

Let $\lambda := \langle a^2, \tilde{a}^1 \rangle * C_\eta$, where $\tilde{a}^1 = a^1 - e_2 + e_1$; λ is in $\Delta_{\beta - e_2 + e_1}$; thus $\partial\eta$ is homologous to $\partial(\eta + \lambda)$ in $\Delta_{\beta - e_2 + e_1}$; but $\partial(\eta + \lambda) = (a^1 - \tilde{a}^1) * C_\eta = (a^1 - \tilde{a}^2 + \tilde{a}^2 - \tilde{a}^1) * C_\eta = \partial(\langle a^1, \tilde{a}^2 \rangle + \langle \tilde{a}^2, \tilde{a}^1 \rangle) * C_\eta$ where $\tilde{a}^2 = a^2 - e_2 + e_1$.

If every vertex of C_η has first coordinate 0, then $\eta \subset \Delta_{\beta - e_1}$ and $(\langle a^1, \tilde{a}^2 \rangle + \langle \tilde{a}^2, \tilde{a}^1 \rangle) * C_\eta \subset \Delta_{\beta - e_2 + e_1}$ and then we conclude.

Suppose there is a vertex V of C_η with first coordinate equal to 1 (observe that then, if $\langle P, V \rangle$ is a simplex in C_η , then $P_1 = 0$); then the simplexes of $\langle a^1, \tilde{a}^2 \rangle + \langle \tilde{a}^2, \tilde{a}^1 \rangle * C_\eta$ not in $\Delta_{\beta - e_2 + e_1}$ (which are the simplexes of $\langle \tilde{a}^2, \tilde{a}^1 \rangle * C_\eta$ not in $\Delta_{\beta - e_2 + e_1}$) form $UFO_{3,4}^1(\Delta_{\beta - 2e_2 + 2e_1})$; thus by Lemma 20 Case $t = p$ we can prove that $\partial(\langle a^1, \tilde{a}^2 \rangle + \langle \tilde{a}^2, \tilde{a}^1 \rangle) * C_\eta \sim 0$ in $\Delta_{\beta - e_2 + e_1}$ and this finishes the proof. \square

Corollary 23 *Let $d, n_1, \dots, n_d, p \in \mathbf{N} - \{0\}$ with $n_1 \leq 3$ and $p \in \{2, 3\}$. Let $\beta \in \mathbf{N}A_{n_1, \dots, n_d}$ and $\deg \beta \geq p + 2$. If η is a p -chain in Δ_β with $\partial\eta$ in $\Delta_{\beta - e_2}$, then $\exists \tilde{\eta}$ p -chain in $\Delta_{\beta + e_1 - e_2}$ with $\partial\tilde{\eta} = \partial\eta$.*

Proof. To prove the statement, is sufficient to prove it when η is a $UFO_{t,p+1}^2$ for $t = 1, \dots, p + 1$, $p \in \{2, 3\}$ since η is a sum of $UFO_{t,p+1}^2$ for $t = 1, \dots, p + 1$ $p \in \{2, 3\}$. Thus our statement follows from Lemmas 20 and 22. \square

Proof of Prop. 16. We will show that if γ is a $(p - 1)$ -cycle in Δ_b with $\gamma = \partial\eta$ with η in $\Delta_b \cup \Delta_{b - e_1 + e_2} \cup \dots \cup \Delta_{b - ke_1 + ke_2}$ for some $k \leq b_1$, then we can construct η' in $\Delta_b \cup \dots \cup \Delta_{b - (k-1)e_1 + (k-1)e_2}$ s.t. $\partial\eta' = \gamma$ (this, by induction on b_1 , implies obviously Prop. 16): let ν be the sum of the simplexes of η in $\Delta_{b - ke_1 + ke_2}$ and not in $\Delta_{b - ke_1 + (k-1)e_2}$; $\partial\nu$ is in $\Delta_{b - ke_1 + (k-1)e_2}$; by Corollary 23 $\partial\nu = \partial\nu'$ for some ν' in $\Delta_{b - (k-1)e_1 + (k-1)e_2}$ let $\eta' = \eta - \nu + \nu'$; η' is in $\Delta_b \cup \dots \cup \Delta_{b - (k-1)e_1 + (k-1)e_2}$ and $\partial\eta' = \partial\eta = \gamma$. \square

References

- [1] S. Barcanescu, N. Manolache *Betti numbers of Segre-Veronese singularities* Rev. Roumaine Math. Pures Appl. **26** no. 4, 549-565 (1981)
- [2] Ch. Birkenhake *Linear systems on projective spaces* Manuscripta Math. **88** 177-184 (1995)

- [3] A. Campillo, C. Marijuan *Higher relations for a numerical semigroup* Sem. Theorie Nombres Bordeaux **3**, 249-260 (1991)
- [4] A. Campillo, P. Pison *L'ideal d'un semigroup de type fini* C.R. Acad. Sci. Paris Ser. I, **316** 1303-1306 (1993)
- [5] L. Ein, R. Lazarsfeld *Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension*, Invent. Math., **111** no. 1, 51-67 (1993)
- [6] W. Fulton, J. Harris *Representation Theory* Springer Verlag
- [7] F.J. Gallego, B.P. Purnaprajna *Some results on rational surfaces and Fano varieties* J. Reine Angew. Math. **538** (2001), 25-55.
- [8] M. Green *Koszul cohomology and the geometry of projective varieties I,II* J. Differ. Geom. **20**, 125-171, 279-289 (1984)
- [9] M. Green *Koszul cohomology and geometry*, in: M. Cornalba et al. (eds), Lectures on Riemann Surfaces, World Scientific Press (1989)
- [10] M. Green, R. Lazarsfeld *On the projective normality of complete linear series on an algebraic curve* Invent. Math. **83**, 73-90 (1986)
- [11] T. Josefiak, P. Pragacz, J.Weyman *Resolutions of determinantal varieties and tensor complexes associated with symmetric and antiymmetric matrices* Asterisque **87-88**, 109-189 (1981)
- [12] A. Lascoux *Syzygies des variétés déterminantales* Adv. in Math. **30**, 202-237 (1978)
- [13] G. Ottaviani, R. Paoletti *Syzygies of Veronese embeddings* Compositio Math. **125**, 31-37 (2001)
- [14] P. Pragacz, J.Weyman *Complexes associated with trace and evaluation. Another approach to Lascoux's resolution* Adv. Math. **57**, 163-207 (1985)
- [15] E. Rubei *On syzygies of Segre embeddings* Proc. A.M.S. **130** 12 3483-3493 (2002)
- [16] E. Rubei *A result on resolutions of Veronese embeddings* preprint Dip. Matematica "U. Dini" Firenze, n. 17, October 2002 math.AG/0309102
- [17] H. Schenck, G. Smith *Syzygies of projective toric varieties* math.AG/0308205
- [18] E. Spanier *Algebraic Topology* Springer Verlag
- [19] B. Sturmfels *Gröbner bases and convex polytopes* University Lecture Series A.M.S. **8** (1996)

Address: Elena Rubei, Dipartimento di Matematica "U. Dini", via Morgagni 67/A, 50134 Firenze, Italia. **E-mail address:** rubei@math.unifi.it

2000 Mathematical Subject Classification: 14M25, 13D02.

Key words: syzygies, Segre embeddings.