

NON-MINIMAL SCALAR-FLAT KÄHLER SURFACES AND PARABOLIC STABILITY

YANN ROLLIN AND MICHAEL SINGER

ABSTRACT. A new construction is presented of scalar-flat Kähler metrics on non-minimal ruled surfaces. The method is based on the resolution of singularities of orbifold ruled surfaces which are closely related to rank-2 parabolically stable holomorphic bundles. This rather general construction is shown also to give new examples of low genus: in particular, it is shown that $\mathbb{C}\mathbb{P}^2$ blown up at 10 suitably chosen points, admits a scalar-flat Kähler metric; this answers a question raised by Claude LeBrun in 1986 in connection with the classification of compact self-dual 4-manifolds.

1. INTRODUCTION

The theory of Kähler metrics with constant scalar curvature (CSC) has seen significant progress in the last ten years: some of the obvious highlights are:

- The work of LeBrun and his co-workers [L3, L4, KLP, LS] which gives many compact examples in complex dimension 2;
- The work of Donaldson [D] which shows that for projective varieties with no non-trivial holomorphic vector fields there is at most one metric of constant scalar curvature in any given Kähler class; and that if such a metric *does* exist, then the underlying polarized variety must be *stable* in a suitable algebro-geometric sense;
- The work of X.X. Chen and Tian [CT] which extends Donaldson's uniqueness result, by different methods, to arbitrary compact Kähler manifolds (and to *extremal* Kähler metrics).

The conjecture that the existence of CSC Kähler metrics should be related to the algebro-geometric notion of stability seems to go back to Yau, but, despite the work of Donaldson, Chen and Tian, a proof of the 'obvious conjecture' (stability \Rightarrow existence) is still lacking. In the absence of a general theorem of this kind, special constructions still have an important role to play.

In this paper we give a new construction of compact *scalar-flat Kähler* (SFK) surfaces, in other words Kähler metrics on compact complex surfaces having scalar curvature equal to zero. If M is such a surface and $[\omega]$ is the Kähler class, then $c_1 \cdot [\omega] = 0$, for this is just the integral over M of the scalar curvature. It follows (cf. [L1, LS]) that $c_1^2(M) \leq 0$, with equality if and only if $c_1(M) = 0$ and the metric is Kähler-Einstein. It follows from surface

Date: April 2004.

First author partly supported by NSF grant # DMS-0305130.

Second author partly supported by a Leverhulme research fellowship and an EPSRC small grant.

classification that if $c_1^2(M) \leq 0$ but $c_1(M) \neq 0$, then M is rational or ruled, though not necessarily *minimal*¹.

The simplest examples are therefore blow-ups of the complex projective plane $\mathbb{C}\mathbb{P}^2$. Since $c_1^2(\mathbb{C}\mathbb{P}^2) = 9$ and every blow-up reduces c_1^2 by 1, we see that a 10-point blow-up of $\mathbb{C}\mathbb{P}^2$ is the first such surface that could possibly admit a SFK metric. In [KLP] it was shown that a 14-point blow-up of $\mathbb{C}\mathbb{P}^2$ does admit a SFK metric. Our first result is a sharp improvement of this result, answering a long-standing question [L1] of LeBrun's:

Theorem A. *The complex projective plane $\mathbb{C}\mathbb{P}^2$, blown up at 10 suitably chosen points, admits a scalar flat Kähler metric. Any further blow-up of the resulting complex surface admits a scalar-flat Kähler metric.*

We also obtain new constructions of SFK metrics on ruled surfaces with base an elliptic curve:

Theorem B. *Let \mathbb{T} be a compact Riemann surface of genus 1.*

(i) *Let L_1 and L_2 be two non-isomorphic holomorphic line bundles of same degree over \mathbb{T} . Then there is a 2-point blow-up of $\mathbb{P}(L_1 \oplus L_2)$ which admits a scalar-flat Kähler metric.*

(ii) *There is a 4-point blow-up of $\mathbb{T} \times \mathbb{C}\mathbb{P}^1$ which admits a scalar-flat Kähler metric.*

Any further blow-up of the resulting complex surfaces admit a scalar-flat Kähler metric.

In addition to these specific examples, our construction provides some support for the slogan “stability \Rightarrow existence”. For *minimal* ruled surfaces (i.e., no blow-ups) the relation between stability and existence was noticed in [BB]: a ruled surface of the form $\mathbb{P}(E) \rightarrow \Sigma$, where Σ is a Riemann surface of genus ≥ 2 and $E \rightarrow \Sigma$ is a rank-2 holomorphic vector bundle, admits a SFK metric if and only if E is *polystable*. This result depends on the celebrated theorem of Narasimhan and Seshadri [NS], which allows to construct the metric on $\mathbb{P}(E)$ as a quotient of the Riemannian product metric $\mathbb{C}\mathbb{P}^1 \times \mathbb{H}^2$, where the two factors are equipped with the standard metrics of constant curvature $+1$ and -1 respectively.

For non-minimal ruled surfaces, the following result was proved by LeBrun and the second author:

Proposition C ([LS], Corollary 3.9). *Let M be some blow-up of a compact geometrically ruled surface $\pi : \mathbb{P}(E) \rightarrow \Sigma$. Suppose that M admits a non-zero, periodic holomorphic vector field. Then a Kähler class $[\omega]$ on M with $c_1(M) \cdot [\omega] = 0$ contains a representative of zero scalar curvature if and only if the parabolic bundle E is quasi-stable.*

In this statement, the blow-up and Kähler class are encoded by a parabolic $GL_2(\mathbb{C})$ -structure on E as follows. If the centres of the blow-ups are the points Q_1, \dots, Q_k , then we obtain k marked points $P_j = \pi(Q_j)$ in Σ and flags $0 \subset L_j \subset \pi^{-1}(P_j)$ in the corresponding fibres. The corresponding parabolic weight (β_j, γ_j) is not uniquely defined, but is chosen to satisfy

$$\gamma_j - \beta_j = \frac{\int_{S_j} \omega}{\int_F \omega}, \quad \beta_j, \gamma_j \in]0, 1[$$

¹I.e. M could contain divisors that can be blown down without introducing singularities

where S_j is the exceptional divisor introduced by blowing up P_j .

At the end of [LS] it was conjectured that Proposition C should continue to hold if there is no periodic holomorphic vector field, with “quasi-stable” replaced by “stable”. The methods of this article do not prove this conjecture; instead, we use the parabolic structure to encode an iterated blow-up of $\mathbb{P}(E)$ and define a “map” of the following kind:

$$\boxed{\text{Parabolically stable bundles over a punctured hyperbolic Riemann surface } E \rightarrow \widehat{\Sigma}} \rightarrow \boxed{\text{SFK metrics on a certain blow-up } \widehat{M} \text{ of } \mathbb{P}(E), \text{ encoded by the parabolic structure}}$$

The reason for the quotation marks is that this “map” is only defined for rational values of the parabolic weights, and will not be smooth in any obvious sense. *We state again that although the blow-up \widehat{M} on the right-hand side is encoded by the parabolic structure, its construction is completely different from the one involved in Proposition C.*

By the theorem of Mehta–Seshadri [MS], there is a correspondence between parabolically stable bundles and representations of the fundamental group of the punctured Riemann surface. We use these representations in the statement of our main theorem:

Theorem D. *Let $\widehat{\Sigma}$ be a compact Riemann surface of genus g with a finite set of marked points $\{P_1, P_2, \dots, P_k\}$ and $\rho : \pi_1(\widehat{\Sigma} \setminus \{P_j\}) \rightarrow \text{SU}(2)/\mathbb{Z}_2$ be a homomorphism. Assume in addition that*

- (i) *if l_j is the homotopy class of a small loop around P_j , then $\rho(l_j)$ has finite order q_j ;*
- (ii) $2 - 2g - \sum_{j=1}^k (1 - \frac{1}{q_j}) < 0$;
- (iii) *ρ defines an irreducible representation in the sense that the induced action of $\pi_1(\widehat{\Sigma} \setminus \{P_j\})$ fixes no point of \mathbb{CP}^1 .*

Then there is non-minimal ruled surface $\widehat{M}_\rho \rightarrow \widehat{\Sigma}$ associated canonically to ρ , which admits a scalar-flat Kähler metric.

The construction of \widehat{M}_ρ will be sketched later in the Introduction and is given in detail in §§2–3.

Remarks. (i) We shall see in §2 that the blow-ups made in the construction of our SFK ruled surfaces \widehat{M} are rather non-generic: they all involve iterated blow-ups (a sequence of blow-ups where each centre lies on the exceptional divisor introduced by the previous blow-up). Thus we get SFK metrics on a rather “thin” set in the moduli space of complex structures on \widehat{M} . The general problem of existence of a SFK metric in a Kähler class satisfying $c_1 \cdot [\omega] = 0$ remains mysterious.

- (ii) It is not clear whether Theorem B is sharp: for example, do there exist SFK metrics on a 1-point blow-up of $\mathbb{P}(L_1 \oplus L_2) \rightarrow \mathbb{T}$, if L_1 and L_2 are not isomorphic? Do there exist such metrics on a 3-point blow-up of $\mathbb{T} \times \mathbb{CP}^1$? In this direction, we remark that [LS, Prop. 3.1] shows that there do not exist SFK metrics in the

first case if the centre of the blow-up lies on L_1 or L_2 and in the second case if 2 or fewer points are blown up. (The obstructions come from the non-trivial holomorphic vector field on these spaces.)

In the light of the recent work of Donaldson and Chen–Tian, the following seems a reasonable

Conjecture E. *Let $E \rightarrow \Sigma$ be a parabolic holomorphic bundle of rank 2 (with rational weights) over a Riemann surface, such that the ruled surface $\mathbb{P}(E)$ has no non-trivial holomorphic vector field. If the corresponding iterated blow-up \widehat{M} of $\mathbb{P}(E)$ (cf. Section 2.1 for a precise definition) admits a scalar-flat Kähler metric, then E must be parabolically stable.*

It is ironic that the above conjecture runs in the “easier” direction (existence \Rightarrow stability) and yet we are unable to prove it; the new numerical criterion for stability due to Ross and Thomas [RT] should be useful here, but so far we have not succeeded in applying it.

Another possible approach might be to apply the Tian–Viaclovsky compactness theorem [TV] which shows that under certain conditions, a sequence of SFK metrics can only degenerate to a SFK orbifold metric. (Our gluing theorem, Theorem 4.1.1, gives explicit examples of this degeneration process.)

Outline. This work began from the observation that in the Burns–de Bartolomeis construction, the smooth base Σ can be replaced by an orbifold Riemann surface $\overline{\Sigma}$. More precisely, with the notation of Theorem D, $\overline{\Sigma}$ is the smooth Riemann surface Σ with a finite set of marked points P_j and corresponding integer weights q_j . By a theorem of Troyanov [Tr], if the orbifold Euler characteristic is negative (this is condition (ii) of Theorem D), then $\overline{\Sigma}$ carries a (Kähler) orbifold metric \overline{g} of constant curvature -1 . In particular, \overline{g} is smooth on $\overline{\Sigma} \setminus \{P_j\}$ and has a conical singularity at P_j , with cone angle $2\pi/q_j$. The riemannian product $\overline{\Sigma} \times \mathbb{C}\mathbb{P}^1$ is obviously scalar-flat Kähler, with non-isolated orbifold singularities around the fibres $F_j = \{P_j\} \times \mathbb{C}\mathbb{P}^1$.

In order to replace these by isolated singularities, we twist by a representation ρ of the orbifold fundamental group of $\overline{\Sigma}$, just as was done in the smooth case by Burns and de Bartolomeis. This ρ must be as in condition (i) of Theorem D: using it gives an orbifold SFK metric on an orbifold ruled surface, \overline{M} , say. It turns out that \overline{M} has precisely two isolated cyclic singularities in each fibre F_j . Denote by \widehat{M} the minimal resolution of singularities of \overline{M} .

In §§4–5, it will be shown that \widehat{M} has a SFK metric, by an analytical gluing theorem. On the other hand, we shall see in §§2–3 that \widehat{M} is also a multiple blow-up of a smooth minimal ruled surface $\tilde{M} = \mathbb{P}(E) \rightarrow \widehat{\Sigma}$, say. In fact, \tilde{M} and \overline{M} can be viewed as two different compactifications (one smooth, the other an orbifold) of a non-compact ruled surface $M^* \rightarrow \Sigma \setminus \{P_j\}$, each canonically associated to the representation ρ . Such representations are related by the Mehta–Seshadri Theorem [MS] to parabolic stability of the underlying holomorphic vector bundle. In §2 we shall start from this point, defining a notion of parabolically stable ruled surface \tilde{M} and the corresponding multiple blow-up \widehat{M} . In §3, we shall compare this with the orbifold \overline{M} .

In §2, we shall restate Theorem D in the language of stable ruled surfaces and show how Theorems A and B follow. The advantage of working with parabolically stable bundles is that it is often quite easy to verify stability, whereas it can be rather difficult to find explicit representations of the fundamental group of a punctured Riemann surface.

Acknowledgments. We thank Olivier Biquard for useful conversations and for pointing out a mistake in the proof of the gluing theorem for scalar-flat Kähler metrics in [KS]. This problem is now fixed in section 4 where a self-contained exposition of the result can be found. We also thank Claude LeBrun and Rafe Mazzeo for encouragement and several useful discussions. Finally, we thank the anonymous referee for several suggestions which improved the original manuscript. This work was carried out while the second author was visiting the Mathematics Department at MIT; he thanks MIT for its hospitality and financial support during this visit.

2. PARABOLICALLY STABLE RULED SURFACES

A *geometrically ruled surface* \check{M} is by definition a minimal complex surface obtained as $\check{M} = \mathbb{P}(E)$, where $E \rightarrow \widehat{\Sigma}$ is a holomorphic vector bundle of rank 2 over a Riemann surface $\widehat{\Sigma}$. The induced map $\pi : \check{M} \rightarrow \widehat{\Sigma}$ is called the *ruling*.

A parabolic structure on \check{M} consists of the following data:

- A finite set of distinct points P_1, P_2, \dots, P_n in $\widehat{\Sigma}$;
- for each j , a choice of point $Q_j \in F_j = \pi^{-1}(P_j)$;
- for each j , a choice of weight $\alpha_j \in]0, 1[\cap \mathbb{Q}$.

A geometrically ruled surface with a parabolic structure will be called a *parabolic ruled surface*.

If $S \subset \check{M}$ is a holomorphic section of π , we define its slope

$$\mu(S) = S^2 + \sum_{Q_j \notin S} \alpha_j - \sum_{Q_j \in S} \alpha_j;$$

we say that a parabolic ruled surface is *stable* if for every holomorphic section S , we have $\mu(S) > 0$.

If we return to the vector bundle E , then Q_j defines a line L_j in the fibre of E over P_j . For each j , select $0 \leq \beta_j < \gamma_j < 1$ with $\alpha_j = \gamma_j - \beta_j$. In this way E is endowed with (a family of) parabolic structures. Our notion of stability of a parabolic ruled surface corresponds with the Mehta–Seshadri notion of parabolic stability for E . Indeed, holomorphic cross-sections $S \subset \check{M}$ correspond exactly to holomorphic sub-bundles $L \subset E$. We know that $H^2(\mathbb{P}(E), \mathbb{Z})$ is generated by the class of a fibre F and $H = c_1(\mathcal{O}(1))$ (the fibrewise hyperplane section bundle) on each fibre (cf. for instance [Bea]). They verify

$$H^2 = \deg(E), \quad F^2 = 0, \quad F \cdot H = 1.$$

Moreover, we have

$$S = H - \deg(L)F$$

hence

$$S^2 = \deg(E) - 2 \deg(L).$$

It follows that $\mu(S)$ is equal to twice the difference of the parabolic slopes of E and L in the sense of Mehta–Seshadri [MS], so \check{M} is stable if and only if E is parabolically stable.

2.1. Iterated blow-up of a parabolic ruled surface. Let \check{M} be a parabolic ruled surface. We shall now define a multiple blow-up $\Phi : \widehat{M} \rightarrow \check{M}$ which is canonically determined by the parabolic structure of \check{M} .

In order to simplify the notation, suppose that the parabolic structure on \check{M} is reduced to a single point $P \in \widehat{\Sigma}$; let Q be the corresponding point in $F = \pi^{-1}(P)$ and let $\alpha = \frac{p}{q}$ be the weight, where p and q are two coprime integers, $0 < p < q$. Denote the Hirzebruch–Jung continued fraction expansion of α by

$$\frac{p}{q} = \frac{1}{e_1 - \frac{1}{e_2 - \cdots \frac{1}{e_k}}}; \quad (2.1)$$

define also

$$\frac{q-p}{q} = \frac{1}{e'_1 - \frac{1}{e'_2 - \cdots \frac{1}{e'_l}}}. \quad (2.2)$$

These expansions are unique if, as we shall assume, the e_j and e'_j are all ≥ 2 .

Proposition 2.1.1. *There exists a unique iterated blow-up $\Phi : \widehat{M} \rightarrow \check{M}$ with $\Phi^{-1}(F)$ equal to the following chain of curves:*

$$\begin{array}{cccccccccccccccc} & -e_1 & & -e_2 & & \cdots & & -e_{k-1} & & -e_k & & -1 & & -e'_1 & & -e'_{l-1} & & \cdots & & -e'_2 & & -e'_1 & . \end{array} \quad (2.3)$$

Here the edges represent rational curves, the number above each edge is the self-intersection of the curve, the hollow dots represent transverse intersections with intersection number +1, and the curve of self-intersection $-e_1$ is the proper transform of the exceptional divisor of the first blow-up.

Proof. We give an iterative construction. The first step is to blow up Q , to get a diagram of the form

$$\begin{array}{ccc} & -1 & & -1 & \\ & \text{---} & \circ & \text{---} & \\ & & & & \end{array} \quad (2.4)$$

By blowing up the intersection point of these two curves we get the diagram

$$\begin{array}{ccccccc} & -2 & & -1 & & -2 & \\ & \text{---} & \circ & \text{---} & \circ & \text{---} & \\ & & & & & & \end{array} \quad (2.5)$$

in which we see two (-2) -curves separated by a -1 curve. Suppose by induction that we have used a sequence of blow-ups so that the following chain of curves sits over F :

$$\begin{array}{cccccccccccccccc} & -a_1 & & -a_2 & & \cdots & & -a_{j-1} & & -a_j & & -1 & & -b_r & & -b_{r-1} & & \cdots & & -b_2 & & -b_1 \end{array} \quad (2.6)$$

for integers a_i and $b_i \geq 2$. Then we can increase a_j by one unit by blowing up the point marked A and we can introduce a new curve of self-intersection -2 by blowing up B :

$$\begin{array}{cccccccccccccccc} & -a_1 & & -a_2 & & & -a_{j-1} & -a_{j-1} & -1 & -2 & & -b_r & -b_{r-1} & & & -b_2 & -b_1 \\ & \circ & & \circ & & \circ & \circ & \circ & \circ & \circ & & \circ & \circ & & \circ & \circ & \circ \end{array} \quad (2.7)$$

or

$$\begin{array}{cccccccccccccccc} & -a_1 & & -a_2 & & & -a_{j-1} & -a_j & -2 & -1 & & -b_{r-1} & -b_{r-1} & & & -b_2 & -b_1 \\ & \circ & & \circ & & \circ & \circ & \circ & \circ & \circ & & \circ & \circ & & \circ & \circ & \circ \end{array} \quad (2.8)$$

It is now clear that a suitable sequence of $\sum(e_j - 1)$ blow-ups, starting from the configuration (2.4) yields the required configuration of curves. All that remains is the proof that the integers e'_j in (2.4) do satisfy (2.2). For this we use the fact if

$$\frac{\lambda}{\mu} = \frac{1}{a_1 - \frac{1}{a_2 - \cdots \frac{1}{a_j}}} \quad (2.9)$$

then the reversed continued fraction

$$\frac{1}{a_j - \frac{1}{a_{j-1} - \cdots \frac{1}{a_1}}} = \frac{\lambda'}{\mu} \quad (2.10)$$

where $\lambda\lambda' \equiv 1 \pmod{\mu}$, $0 < \lambda' < \mu$. Suppose by induction that in (2.6) we have

$$\frac{1}{a_1 - \frac{1}{a_2 - \cdots \frac{1}{a_j}}} + \frac{1}{b_1 - \frac{1}{b_2 - \cdots \frac{1}{b_r}}} = 1 \quad (2.11)$$

Since $(\mu - \lambda')(\mu - \lambda)$ is congruent to 1 mod μ , (2.11) implies that

$$\frac{1}{a_j - \frac{1}{a_{j-1} - \cdots \frac{1}{a_1}}} + \frac{1}{b_r - \frac{1}{b_{r-1} - \cdots \frac{1}{b_1}}} = 1 \quad (2.12)$$

Now consider how these fractions change under the blow-up. We have that a_j is replaced by $a_j + 1$, and a new b_{r+1} is introduced, equal to 2. So the inductive step is completed by the trivial identity

$$\frac{1}{1 + \mu/\lambda'} + \frac{1}{2 - (1 - \lambda'/\mu)} = 1.$$

Since the induction clearly starts (consider the diagram (2.5)), the proof is now complete. \square

Our main theorem may now be stated as follows:

Theorem 2.1.2. *Let $\check{M} \rightarrow \widehat{\Sigma}$ be a parabolically stable ruled surface. Suppose that*

$$\chi(\widehat{\Sigma}) - \sum_{j=1}^k \left(1 - \frac{1}{q_j}\right) < 0, \quad (2.13)$$

where $\alpha_j = p_j/q_j$, with p_j and q_j coprime. Then the iterated blow-up \widehat{M} deduced from the parabolic structure carries a SFK metric. Furthermore, any blow-up of \widehat{M} also carries a SFK metric.

The relationship between this statement and Theorem D corresponds to the relationship between representations and parabolically stable bundles, given by the theorem of Mehta and Seshadri [MS]. According to this theorem, given a parabolically stable \check{M} , there is a corresponding irreducible representation

$$\rho' : \pi_1(\widehat{\Sigma} \setminus \{P_j\}) \rightarrow U_2 \quad (2.14)$$

with

$$\rho'(l_j) \text{ conjugate to } \pm \text{diag}(e^{2\pi i\beta_j}, e^{2\pi i\gamma_j}) \quad (2.15)$$

where l_j is the homotopy class of a small loop around P_j and $\gamma_j - \beta_j = \alpha_j$ as before. This representation defines a parabolically stable bundle E with $\mathbb{P}(E) = \check{M}$. We note that E is determined only up to tensoring with a line-bundle; we make use of this freedom (and the freedom of choice in β_j, γ_j) to ensure that the parabolic degree of E is zero, as is necessary for the existence of ρ' .

If $A \in U_2$, write $A = z(A)A_0$, where $z(A)$ is a scalar multiple of the identity and $A_0 \in SU_2/\mathbb{Z}_2$. Letting $\rho(\gamma) = \rho'(\gamma)_0$, we obtain an irreducible representation

$$\rho : \pi_1(\widehat{\Sigma} \setminus \{P_j\}) \rightarrow SU_2/\mathbb{Z}_2 \quad (2.16)$$

with

$$\rho(l_j) \text{ conjugate to } \pm \text{diag}(e^{\pi i\alpha_j}, e^{-\pi i\alpha_j}) \quad (2.17)$$

as required for Theorem D.

Conversely, given a representation with properties (2.16) and (2.17), we clearly get a ruled surface $M^* \rightarrow \widehat{\Sigma} \setminus \{P_j\}$. However, M^* can now be compactified to give \check{M} in a standard way: let P stand for one of the punctures, and identify a neighbourhood of P in $\widehat{\Sigma}$ with the unit disc Δ . Write $L^* = \pi^{-1}(\Delta \setminus \{0\})$ and note that

$$L^* = \mathbb{H}^2 \times \mathbb{C}\mathbb{P}^1/\mathbb{Z} \quad (2.18)$$

where \mathbb{H}^2 is the hyperbolic plane $\{\text{Im}(\xi) > 0\}$ and the \mathbb{Z} -action is generated by

$$(\xi, [w_0 : w_1]) \mapsto (\xi + 2\pi, [e^{\pi i\alpha} w_0, e^{-\pi i\alpha} w_1]). \quad (2.19)$$

Let $\check{L} = \Delta \times \mathbb{C}\mathbb{P}^1$ be covered by two standard holomorphic coordinate charts, $(x_1, y_1) \in \mathbb{C}^2$, $(x_2, y_2) \in \mathbb{C}^2$ glued together by $x_2 = x_1$, $y_2 = y_1^{-1}$. Then the natural embedding of L^* in \check{L} is given by the following map

$$x_1 = x_2 = e^{i\xi}, \quad y_1 = e^{-i\xi\alpha} w_0/w_1, \quad y_2 = e^{i\xi\alpha} w_1/w_0. \quad (2.20)$$

Notice that the pre-image of Q , which corresponds to $(x_1, y_1) = (0, 0)$ is given by $\xi = +i\infty$, $w_0 = 0$, but every other point corresponding $\xi = i\infty$ is mapped to $(x_2, y_2) = (0, 0)$.

In general, we obtain \check{M} as a smooth compactification of M^* by following this recipe at each puncture.

Let us show how Theorem A and Theorem B follow from Theorem 2.1.2.

Corollary 2.1.3. *There is a 9-point iterated blow-up of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ which admits a SFK metric. As a consequence $\mathbb{C}\mathbb{P}^2$ has a 10-point iterated blow-up which admits a scalar-flat Kähler metric.*

Proof. Let $\check{M} = \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$, and let $\pi : \check{M} \rightarrow \mathbb{C}\mathbb{P}^1$ denote projection on the first factor. Pick any 4 points P_1, P_2, P_3, P_4 in $\mathbb{C}\mathbb{P}^1$, with weights $\alpha_1 = \alpha_2 = \alpha_3 = 1/2$ and $\alpha_4 = 1/3$, so that (2.13) is satisfied. Pick $Q_j \in \pi^{-1}(P_j)$. To check when this parabolic structure is stable, note first that any section S is in this case the graph of a meromorphic function f , and $S^2 = 2 \deg f$. Provided that no two of the Q_j lie on the graph of a function of degree 0, we have a parabolically stable ruled surface. In particular, the set of stable configurations of the Q_j is a Zariski open set in the set of all such configurations.

The multiple blow-up \widehat{M} of \check{M} involves a total of 9 blow-ups, 2 each at Q_1, Q_2 and Q_3 and 3 at Q_4 . Hence \widehat{M} is indeed a 9-point blow-up of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$. However, if $n \geq 1$, then any n -point blow-up of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ is isomorphic to an $(n+1)$ -point blow-up of $\mathbb{C}\mathbb{P}^2$. \square

Now we turn to ruled surfaces with base an elliptic curve.

Corollary 2.1.4. *Let \mathbb{T} be an elliptic curve (Riemann surface of genus 1) and let L_1 and L_2 be two non-isomorphic line-bundles of degree 0 over \mathbb{T} . Then, any double blow-up \widehat{M} of $\check{M} = \mathbb{P}(L_1 \oplus L_2)$ at a point which is not on $\mathbb{P}(L_1)$ or $\mathbb{P}(L_2)$ admits a SFK metric. Any further blow-up of \widehat{M} also admits a SFK metric.*

Proof. Endow \check{M} with a parabolic structure by picking an arbitrary point $P \in \mathbb{T}$ and a point $Q \in \pi^{-1}(P)$ so that Q does not lie on $S_1 = \mathbb{P}(L_1 \oplus 0)$ or $S_2 = \mathbb{P}(0 \oplus L_2)$. Set the weight of Q equal to $1/2$. Then condition (2.13) is satisfied and it remains to check that this parabolic structure is stable.

To see this, note first that every section S of π satisfies $S^2 \geq 0$; this follows because at the level of cohomology, $[S] = [S_1] + r[F]$ for some integer r . Then $S^2 = 0$ if $r = 0$; otherwise $S \neq S_1$, so $r = S \cdot S_1 \geq 0$. Hence $S^2 = 2r \geq 0$.

Therefore, $\mu(S) > 0$ unless $S^2 = 0$ and $Q \in S$. Suppose now that $S^2 = 0$, $S \neq S_1$ and $S \neq S_2$. It follows that $S \cdot S_1 = S \cdot S_2 = 0$, and S meets neither S_1 nor S_2 . Such an S defines an isomorphism $L_1 \simeq L_2$ contradicting the hypothesis of the corollary.

The corresponding 2-point blow-up \widehat{M} of \check{M} carries a scalar-flat Kähler metric. \square

In the next corollary, we recover a result of Kim, LeBrun and Pontecorvo [KLP] by a completely different method.

Corollary 2.1.5. *Let \mathbb{T} be an elliptic Riemann surface. There is a 6-point blow-up of $\mathbb{T} \times \mathbb{C}\mathbb{P}^1$ which admits a SFK metric.*

Proof. Let $\check{M} = \mathbb{T} \times \mathbb{C}\mathbb{P}^1$, let π be the projection on the first factor. Endow \check{M} with a parabolic structure by choosing any 3 points P_j in \mathbb{T} , points $Q_j \in \pi^{-1}(P_j)$ each of weight $1/2$. We note that the condition (2.13) is satisfied as before: it remains to analyze the stability of this parabolic structure.

Any holomorphic section S of π corresponds to the graph of a meromorphic function s on \mathbb{T} . Let H be the section corresponding to the constant function 0. Then $S \cdot F = 1$, and $S \cdot H = \sum a_n$, where the $a_n \geq 0$ are the multiplicities of the zeroes of s ; hence

$$S = H + \left(\sum a_n \right) F.$$


If $S^2 = 2 \sum a_n = 0$, then $a_n = 0$ and s must be constant. If we choose the points Q_j such that no two of them lie on the graph of a constant meromorphic function, we must have $\mu(S) \geq 0 + 2/2 > 0$. If $S^2 > 0$, we have $S^2 \geq 2$ for S^2 is even and then $\mu(S) \geq 2 - 3/2 > 0$ whatever the positions of the Q_j .

Applying Theorem 2.1.2, there exists a SFK metric on \widehat{M} , which is in this case a 6-point blow-up of \check{M} ; more precisely \check{M} is obtained by performing a double blow-up at each of Q_1, Q_2 and Q_3 . \square

We now improve the last result.


Corollary 2.1.6. *Let \mathbb{T} be an elliptic curve. There is a 4-point blow-up of $\mathbb{T} \times \mathbb{CP}^1$ which admits a scalar-flat Kähler metric.*

Proof. Let P_1 and P_2 be two points in \mathbb{T} . Let S be a constant section which is not the zero section S_0 or the section at infinity S_∞ , and M' be the blow-up of $\mathbb{T} \times \mathbb{CP}^1$ at $\{P_1\} \times \{0\}$ and $\{P_2\} \times \{\infty\}$. In M' , the proper transform of $\{P_j\} \times \mathbb{CP}^1$ is a (-1) -curve; we blow each of these curves down, getting a new minimal ruled surface \check{M} , giving the following diagram



Here ϕ is a blow-up map with centres at C_1 and C_2 , say, and $\phi^{-1}(C_j)$ is the ψ -proper transform of $\{P_j\} \times \mathbb{CP}^1$.

We shall show that $\check{M} = \mathbb{P}(L_1 \oplus L_2)$ satisfies the conditions of Corollary 2.1.4. Then the double blow-up $\Phi : \widehat{M} \rightarrow \check{M}$ at some point Q is scalar-flat Kähler, and so is any further blow-up of \widehat{M} . In particular, the blow-up $\widehat{\phi} : \widehat{M}' \rightarrow \widehat{M}$ of \widehat{M} at $\Phi^{-1}(C_1)$ and $\Phi^{-1}(C_2)$ is scalar-flat Kähler. Taking Q to be different from C_1 and C_2 , we see that \widehat{M}' is also the double blow-up at $\phi^{-1}(Q)$ of M' , as in the following diagram:



In particular, \widehat{M}' is a 4-point blow-up of $\mathbb{T} \times \mathbb{CP}^1$.

Let us check that \check{M} satisfies the hypotheses of Corollary 2.1.4. Suppose not. Let S', S'_0, S'_∞ be the images by ϕ in \check{M} of the proper transforms of the corresponding curves

S , S_1 and S_∞ in $\mathbb{T} \times \mathbb{CP}^1$. We have

$$(S'_0)^2 = (S'_\infty)^2 = S'_0 \cdot S'_\infty = 0, \quad S'_0 \cdot S' = S'_\infty \cdot S' = 1, \quad (S')^2 = 2.$$

This shows that $M = \mathbb{P}(L_0 \oplus L_\infty)$, where L_0 and L_∞ are two line bundles of degree 0 over \mathbb{T} .

Suppose that $M \simeq \mathbb{T} \times \mathbb{CP}^1$. Then S'_0 and S'_∞ must be two distinct constant sections of $\mathbb{T} \times \mathbb{CP}^1 \rightarrow \mathbb{T}$ as seen in the proof of Corollary 2.1.5. Up to an isomorphism of \mathbb{CP}^1 , we may assume that they are the 0 section and the section at infinity of $\mathbb{T} \times \mathbb{CP}^1$. Now S' is the graph of a meromorphic function s' on \mathbb{T} . Let z_j be the zeroes of s' and w_j its poles. The divisor of s' is

$$(s') = \sum a_j z_j - \sum b_j w_j,$$

where $a_j, b_j \geq 0$ are the multiplicities. Then, $S' \cdot S'_0 = \sum a_j = 1$ hence there is a unique pole with multiplicity 1. Therefore

$$(s') = z - w,$$

which is impossible by Abel's theorem. \square

Remark. — It is a general principle in algebraic geometry that “stability is an open condition”. We have seen in these examples that the set of stable parabolic structures is Zariski dense in the set of all such structures. This will be true in general: if stable parabolic structures exist on a given ruled surface, then they will form a Zariski-open subset in the set of all such structures. Furthermore, stability will be preserved under perturbation of the parabolic weights, keeping the Q_j fixed. However the iterated blow-up \widehat{M} will behave rather wildly under such perturbations: for example, $b_2(\widehat{M})$ will not remain constant.

3. SCALAR-FLAT KÄHLER ORBIFOLDS

The goal of this section is to introduce an orbifold \overline{M} associated to a parabolic ruled surface \check{M} . We shall see that \check{M} and \overline{M} are bimeromorphic to each other, and indeed both are compactifications of the punctured ruled surface M^* . We shall also show that \overline{M} carries an orbifold SFK metric if \check{M} is stable, and that \overline{M} carries no non-trivial holomorphic vector fields.

3.1. Generalities. A complex orbifold X of complex dimension n may be defined as a complex variety having only quotient singularities. More explicitly, for every point $P \in X$, one requires that there is a finite group $G = G_P$ (called the *local isotropy group*) and a *local uniformizing chart*

$$\tilde{U} \rightarrow \tilde{U}/G \xrightarrow{\phi} U. \quad (3.1)$$

Here, \tilde{U} is a neighbourhood of 0 in \mathbb{C}^n , with a given biholomorphic action of G ,

$$G \times \tilde{U} \rightarrow \tilde{U}, \quad (3.2)$$

U is a neighbourhood of P in X and ϕ is a homeomorphism with $\phi(0) = P$. It is a fact that one can always choose ϕ so as to linearize the action (3.2); that is, one can assume $G \subset GL_n(\mathbb{C})$.

The standard notions of differential geometry extend to orbifolds by working G -equivariantly in a local uniformizing chart. For example a smooth orbifold Riemannian metric g on X is defined as usual away from the singular points, and is given by a G_P -invariant smooth metric on \tilde{U} near the singular point P .

3.2. Orbifold Riemann surfaces. A compact orbifold Riemann surface $\bar{\Sigma}$ can be identified with a smooth compact Riemann surface $\hat{\Sigma}$ together with a finite set of marked points P_j , each with a given weight $q_j \in \mathbb{Z}_{\geq 2}$. It is important to note that a smooth orbifold metric on $\bar{\Sigma}$ is not the same as a smooth metric on $\hat{\Sigma}$: a smooth orbifold metric is smooth on $\Sigma = \bar{\Sigma} \setminus \{P_j\}$ but with respect to such a metric, the length of a small circle of radius r centred at P_j , will be approximately $2\pi r/q_j$.

The Euler characteristic of $\bar{\Sigma}$ is defined as follows

$$\chi(\bar{\Sigma}) = \chi(\hat{\Sigma}) - \sum \left(1 - \frac{1}{q_j}\right), \quad (3.3)$$

where $\chi(\hat{\Sigma})$ is the Euler characteristic of the underlying smooth surface $\hat{\Sigma}$.

Just as for smooth Riemann surfaces with negative Euler characteristic, we have the following result of Troyanov [Tr] (see also [McO]).

Theorem 3.2.1. *The orbifold Riemann surface $\bar{\Sigma}$ admits an orbifold metric of constant curvature -1 compatible with the given complex structure if and only if $\chi(\bar{\Sigma}) < 0$.*

In view of this theorem we shall refer to such orbifold Riemann surfaces as *hyperbolic*. In the previous section we saw several examples of orbifold hyperbolic Riemann surfaces.

Next we come to the fundamental group of an orbifold Riemann surface. Recall first the description of the fundamental group of the punctured Riemann surface Σ :

$$\pi_1(\Sigma) = \langle a_1, b_1, \dots, a_g, b_g, l_1, \dots, l_k : [a_1, b_1][a_2, b_2] \dots [a_g, b_g] l_1 \dots l_k = 1 \rangle \quad (3.4)$$

Here the a_j and b_j are standard generators of $\pi_1(\hat{\Sigma})$ and l_j is (the homotopy class of) a small loop around P_j . The orbifold fundamental group is defined by imposing the additional conditions

$$\pi_1^{orb}(\bar{\Sigma}) = \langle a_1, b_1, \dots, a_g, b_g, l_1, \dots, l_k : [a_1, b_1][a_2, b_2] \dots [a_g, b_g] l_1 \dots l_k = l_1^{q_1} = \dots = l_k^{q_k} = 1 \rangle \quad (3.5)$$

From Theorem 3.2.1 it follows that for any orbifold hyperbolic Riemann surface $\bar{\Sigma}$, we have $\bar{\Sigma} = \mathbb{H}^2/\Gamma$, where Γ is the image of the uniformizing representation of $\pi_1^{orb}(\bar{\Sigma})$ in $SL_2(\mathbb{R}) = \text{Isom}(\mathbb{H}^2)$. The only difference from the smooth case is that Γ will not act freely on \mathbb{H}^2 .

3.3. Orbifold ruled surfaces. Let $\bar{\Sigma}$ be a hyperbolic orbifold Riemann surface as in the last section, and let $\rho : \pi_1^{orb}(\bar{\Sigma}) \rightarrow PSL_2(\mathbb{C})$ be a representation. We suppose that ρ is faithful on the loops l_j , so that $\rho(l_j)$ has order precisely q_j . We can form the quotient

$$\bar{M} = \mathbb{H}^2 \times \mathbb{C}\mathbb{P}^1 / \pi_1^{orb}(\bar{\Sigma}) \quad (3.6)$$

by letting $\pi_1^{orb}(\bar{\Sigma})$ act by the uniformizing representation on the upper half-space \mathbb{H}^2 and by ρ on $\mathbb{C}\mathbb{P}^1$. It is clear that \bar{M} is a complex orbifold ruled surface equipped with a ruling

$$\bar{\pi} : \bar{M} \rightarrow \bar{\Sigma}$$

and singularities only in the fibres $\overline{F}_j = \overline{\pi}^{-1}(P_j)$.

In order to analyze these singularities, choose a complex disc Δ with centre at one of the P_j , let \overline{L} denote $\overline{\pi}^{-1}(\Delta)$ and let $L^* = \overline{L} \setminus \overline{\pi}^{-1}(0)$. Complex analytically, we can identify L^* with a quotient as before (see (2.18) and (2.19)). (We do not use the cusp-metric structure on $\Delta \setminus \{0\}$ which comes from this identification.)

Then the q -fold cover \tilde{L}^* of L^* is given by

$$\tilde{L}^* = \mathbb{H}^2 \times \mathbb{C}\mathbb{P}^1/q\mathbb{Z}. \quad (3.7)$$

Since $\alpha = p/q$, the action here is trivial on $\mathbb{C}\mathbb{P}^1$, and so we can introduce coordinates

$$(u_1, v_1) = (e^{i\xi/q}, w_0/w_1), \quad (u_2, v_2) = (e^{i\xi/q}, w_1/w_0), \quad u_1 = u_2, \quad v_1 = v_2^{-1}. \quad (3.8)$$

If $\omega = e^{2\pi i/q}$, then the action of $\mathbb{Z}/q\mathbb{Z}$ on \tilde{L}^* is given by the standard action of $\Gamma_{p,q}$ on the coordinates (u_1, v_1) and by $\Gamma_{q-p,q}$ on (u_2, v_2) . Here we have written $\Gamma_{r,s}$ for the cyclic subgroup

$$\Gamma_{r,s} = \left\{ \begin{pmatrix} e^{2\pi i n/s} & 0 \\ 0 & e^{2\pi i r n/s} \end{pmatrix} : n = 0, 1, \dots, s-1 \right\}. \quad (3.9)$$

of U_2 . Thus in the orbifold \overline{L} , there are two singularities in the fibre $u_1 = u_2 = 0$.

Denote by $\Psi : \widehat{M} \rightarrow \overline{M}$ the minimal resolution of singularities. This is obtained by replacing each $\Gamma_{p,q}$ -singularity of \overline{M} by the corresponding Hirzebruch–Jung string (cf. [BPV, Fu]). In this resolution, $\Psi^{-1}(F_j)$ is exactly the configuration of curves constructed in Proposition 2.1.1. Indeed, as the notation anticipates, this resolution of singularities is isomorphic to the iterated blow-up constructed there:

Theorem 3.3.1. *Let \check{M} be a parabolically ruled surface and let $M^* = \check{M} \setminus \cup F_j$. Let \overline{M} be the corresponding orbifold. Then we have the diagram*

$$\begin{array}{ccc} & \widehat{M} & \\ \Psi \swarrow & & \searrow \Phi \\ \overline{M} & & \check{M} \\ & \swarrow & \searrow \\ & M^* & \end{array} \quad (3.10)$$

where the lower arrows are the natural inclusions of M^* .

Proof. The problem is local to the base, so we use the local models and coordinates introduced above. In terms of these coordinates, the identity map on L^* becomes the singular map $\overline{L} \rightarrow \tilde{L}$

$$F : (u_1, v_1) \mapsto (x_1, y_1) = (u_1^q, u_1^{-p}v_1), \quad F : (u_2, v_2) \mapsto (x_2, y_2) = (u_1^q, u_1^p v_1), \quad (3.11)$$

which is smooth away from the central fibre $u_1 = u_2 = 0$.

We claim that this map has a very simple description in the language of toric geometry, which makes clear the existence of the diagram (3.10).

Fix the standard lattice $\mathbb{Z}^2 \subset \mathbb{R}^2$. The toric description² of \check{L} is in terms of the fan with 2-dimensional cones

$$\sigma_1 = \{(\mu, \nu) : \mu \geq 0, \nu \geq 0\}, \quad \sigma_2 = \{(\mu, \nu) : \mu \geq 0, \nu \leq 0\}.$$

The dual cones are

$$\sigma_1^* = \{(m, n) : m \geq 0, n \geq 0\}, \quad \sigma_2^* = \{(m, n) : m \geq 0, n \leq 0\}.$$

The corresponding coordinate rings are just $\mathbb{C}[X, Y]$ and $\mathbb{C}[X, Y^{-1}]$. Identifying (X, Y) with (x_1, y_1) in the first case and with (x_2, y_2^{-1}) in the second, we arrive at \check{L} , coordinatized as in (2.20). Similarly, \bar{L} is the toric variety corresponding to the fan with two-dimensional cones

$$\tau_1 = \{(\mu, \nu) : \mu \geq 0, p\mu + q\nu \geq 0\}, \quad \tau_2 = \{(\mu, \nu) : \mu \geq 0, p\mu + q\nu \leq 0\}.$$

The dual cones are

$$\tau_1^* = \{(m, n) : qm - pn \geq 0, n \geq 0\}, \quad \tau_2^* = \{(m, n) : qm - pn \geq 0, n \leq 0\}.$$

Using the same indeterminates as before, we have two affine varieties with coordinate rings

$$A_1 = \bigoplus_{qm \geq pn, n \geq 0} \mathbb{C}X^m Y^n, \quad A_2 = \bigoplus_{qm \geq pn, n \leq 0} \mathbb{C}X^m Y^n.$$

Following Fulton [Fu], the first of these is identified with the coordinate ring of $\mathbb{C}^2/\Gamma_{p,q}$ by introducing variables (u_1, v_1) with $u_1^q = X$, $Y = u_1^{-p}v_1$. Then A_1 is precisely the $\Gamma_{p,q}$ -invariant part of $\mathbb{C}[u_1, v_1]$. Similarly A_2 is identified with the coordinate ring of $\mathbb{C}^2/\Gamma_{q-p,q}$ by setting $X = u_2^q$, $Y = u_2^{-p}v_2^{-1}$. Then it is clear that the identity map of \mathbb{Z}^2 gives rise to our singular holomorphic map.

The map F is singular because the identity map does not map the cone τ_1 into either of the cones σ_1, σ_2 . The Hirzebruch–Jung resolution of the two singularities in \bar{L} corresponds to a subdivision of the fan $\{\tau_1, \tau_2\}$. Recall the continued-fraction expansions (2.1), (2.2) and set

$$v_0 = (0, 1), v_1 = (1, 0), v_j = (m_j, -n_j),$$

where

$$\frac{n_j}{m_j} = \frac{1}{e_1 - \frac{1}{e_2 - \cdots - \frac{1}{e_{j-1}}}},$$

so that $v_{k+1} = (q, -p)$. Define similarly

$$v'_0 = (0, -1), v'_1 = (1, -1), v'_j = (m'_j, n'_j - m'_j)$$

where

$$\frac{n'_j}{m'_j} = \frac{1}{e'_1 - \frac{1}{e'_2 - \cdots - \frac{1}{e'_{j-1}}}}$$

²Strictly, of $\mathbb{C} \times \mathbb{CP}^1$ rather than $\Delta \times \mathbb{CP}^1$, but this is not important in this discussion.

are the approximants to $(q - p)/q$. Again $v'_{l+1} = (q, -p)$. Put

$$\rho_j = \mathbb{R}_{\geq 0}v_j \oplus \mathbb{R}_{\geq 0}v_{j+1}, \rho'_j = \mathbb{R}_{\geq 0}v'_j \oplus \mathbb{R}_{\geq 0}v'_{j+1}$$

for every j . Then the fan consisting of all the ρ_j and ρ'_j corresponds precisely to the minimal resolution $\Psi : \widehat{L} \rightarrow \overline{L}$ and the identity map of \mathbb{Z}^2 now induces the (restriction of the) holomorphic map $\Phi : \widehat{L} \rightarrow \check{L}$. \square

Remark. — It is a pleasant exercise to verify from this point of view that $\widehat{L} \rightarrow \check{L}$ is a multiple blow-up. Combinatorially, a (-1) -curve corresponds to the occurrence of adjacent cones of the form

$$\mathbb{R}_{\geq 0}u \oplus \mathbb{R}_{\geq 0}(u + v), \quad \mathbb{R}_{\geq 0}(u + v) \oplus \mathbb{R}_{\geq 0}v$$

in a fan. The blow-down operation corresponds to the replacement of these two cones by the single cone

$$\mathbb{R}_{\geq 0}u \oplus \mathbb{R}_{\geq 0}v.$$

In the above fan, one has to show that there is a sequence of such deletions that always results in a non-singular fan. In fact, at each stage, one deletes the ray generated by the vector v_j or v'_j with the *largest* x -coordinate. The details are left to the interested reader.

3.4. Scalar-flat Kähler orbifold metrics and holomorphic vector fields.

Theorem 3.4.1. *Let \check{M} , \widehat{M} and \overline{M} be as before, and assume that \check{M} is a stable parabolic ruled surface. Then \overline{M} carries a scalar-flat Kähler orbifold metric \overline{g} ; and the algebra $\mathfrak{a}(\overline{M})$ of holomorphic vector fields is 0.*

Proof. Recall from (3.6) that \overline{M} is a quotient of $\mathbb{H}^2 \times \mathbb{C}\mathbb{P}^1$; if the parabolic structure is stable, then $\pi_1^{orb}(\overline{\Sigma})$ acts by isometries of this product, so its SFK structure descends to define an orbifold SFK structure on \overline{M} .

The proof that $\mathfrak{a}(\overline{M}) = 0$ is very close to the proof of [LS, Prop. 3.1]. Suppose that ξ is a holomorphic vector field on \overline{M} . We claim first that ξ must be vertical. To see this, consider the exact sequence

$$0 \longrightarrow \mathcal{F} \longrightarrow T\overline{M} \xrightarrow{h} \mathcal{N} \longrightarrow 0,$$

on \overline{M} , where \mathcal{F} is the tangent space to the fibres of $\overline{\pi}$, while \mathcal{N} is the normal bundle to the fibres. If $F = \overline{\pi}^{-1}(P)$ is a smooth fibre, then $\mathcal{N}|_F$ is the trivial line-bundle on F and so $h(\xi)|_F$ is constant on F and is given by the pull-back of a vector in $T_P\overline{\Sigma}$. These vectors clearly patch together to give a holomorphic vector field V on the punctured Riemann surface Σ . The same argument works near the singular fibres of $\overline{\pi}$, so V extends to define a smooth (orbifold) vector field on $\overline{\Sigma}$. But the base is an orbifold Riemann surface with negative scalar curvature, and it follows as in the smooth case that such a surface carries no non-trivial holomorphic vector fields. So $V = 0$ and ξ is vertical.

The vector field ξ must have zeros, for it is tangent to the smooth fibre F which is a 2-sphere. In particular, it must lie in the subalgebra $\mathfrak{a}_0(\overline{X})$ of non-parallel holomorphic vector fields, so $\mathfrak{a}(\overline{X}) = \mathfrak{a}_0(\overline{X})$.

Because \overline{M} has a SFK metric \overline{g} , $\mathfrak{a}_0(X)$ is the complexification of the Lie algebra of non-parallel infinitesimal isometries of \overline{g} (cf. [Be, Chapter 2]). So we may assume that ξ

is such an infinitesimal isometry. But then ξ lifts to an infinitesimal isometry of $\mathbb{H}^2 \times \mathbb{C}\mathbb{P}^1$, i.e. an element of the Lie algebra

$$\mathfrak{sl}_2(\mathbb{R}) \times \mathfrak{su}_2 \quad (3.12)$$

which is invariant under the induced action of $\pi_1^{orb}(\overline{\Sigma})$. This action is given by composing the representation

$$\pi_1^{orb}(\overline{\Sigma}) \rightarrow SL_2(\mathbb{R}) \times SU_2/\mathbb{Z}_2$$

with the adjoint action of $SL_2(\mathbb{R}) \times SU_2/\mathbb{Z}_2$ on its Lie algebra. Now the adjoint action of SU_2 on its lie algebra is precisely the action of $SU_2/\mathbb{Z}_2 = SO_3$ on \mathbb{R}^3 , so a non-zero $\pi_1^{orb}(\overline{\Sigma})$ -invariant element in (3.12) will determine an invariant line in \mathbb{R}^3 . An intersection of this line with the unit sphere $S^2 \subset \mathbb{R}^3$ defines an invariant point of $\mathbb{C}\mathbb{P}^1 \simeq S^2$, and we obtain an invariant complex line in \mathbb{C}^2 . This contradicts the irreducibility of the representation ρ . \square

3.5. A remark on uniqueness. Before we proceed to the proof of Theorem D (or Theorem 2.1.2), it is worth mentioning that the SFK metrics which are produced, are in fact unique in their Kähler class. More generally, we have the following result which is a direct consequence of [CT, Theorem 1.1].

Proposition 3.5.1. *Let \widehat{M} be a complex surface as in Theorem D. Then, there is at most one SFK metric in each Kähler class.*

Proof. We give here an alternative proof of the proposition, based on the work of Donaldson. If the proposition is false, there are two distinct SFK metrics ω_1, ω_2 with the same Kähler class Ω . The deformation theory for SFK metrics is unobstructed [LS, Theorem 2.8]. By density, we can find arbitrarily small SFK deformations ω'_1 and ω'_2 with Kähler class $\Omega' \in H^2(\widehat{M}, \mathbb{Q})$. We may assume that ω'_1 and ω'_2 are distinct, which is the case for deformations small enough, and that Ω' is an integral class after multiplication by a suitable constant factor. By Kodaira's embedding theorem (\widehat{M}, Ω') is a projective variety.

We remark now that \widehat{M} cannot have any non trivial holomorphic vector field. Let Ξ be a holomorphic vector fields on \widehat{M} . The complex surface \widehat{M} is by definition the resolution $\pi : \widehat{M} \rightarrow \overline{M}$ of the orbifold \overline{M} . The exceptional fibres π have negative self-intersection hence Ξ must be tangent to them. Therefore the vector field Ξ is projectible. By Theorem (3.4.1) that \overline{M} does not admit any non trivial holomorphic vector field. Thus $\pi_*\Xi = 0$ which forces $\Xi = 0$.

According to [D, Corollary 5], there is at most one Kähler metric of constant scalar curvature on a projective manifold with no non-trivial holomorphic vector field. Thus, ω'_1 and ω'_2 must be equal. This is a contradiction, and the proposition holds. \square

4. SCALAR-FLAT METRICS ON \widehat{M}

In this section we prove a gluing theorem that implies Theorem D (or equivalently Theorem 2.1.2). We begin with a statement of the result and an outline of the gluing argument. We then give a rapid description of the needed perturbation theory of SFK and hermitian-ASD metrics and the necessary linear theory, before application of the implicit function theorem to prove the theorem.

4.1. Statement of gluing theorem and outline of proof.

Theorem 4.1.1. *Let $(\overline{M}, \overline{\omega})$ be a compact SFK orbifold of complex dimension 2, with all singularities isolated and cyclic. Suppose further that the algebra $\mathfrak{a}_0(\overline{M})$ of non-parallel holomorphic vector fields on \overline{M} is zero. Then the minimal resolution \widehat{M} of \overline{M} admits a family of SFK metrics $\widehat{\omega}^\varepsilon$ such that $\widehat{\omega}^\varepsilon \rightarrow \overline{\omega}$ on any compact subset $K \subset \widehat{M} \setminus E$, where E is the exceptional divisor. Moreover, any blow-up of \widehat{M} carries a similar family of SFK metrics.*

We have just seen that the orbifold \overline{M} associated to a stable parabolic ruled surface \check{M} carries a SFK metric and satisfies $\mathfrak{a}_0(\overline{M}) = 0$. Our main theorems D and 2.1.2 therefore follow at once from Theorem 4.1.1.

This theorem would follow from the gluing theorems on gluing hermitian–ASD conformal structures stated in [KS]. Unfortunately, the argument given there is not globally consistent; we therefore give a complete proof here. While many of the ingredients are the same, the new argument given here is perhaps a little more direct. We shall now outline the main points; the technical details follow in the rest of this section.

The first step in the proof of Theorem 4.1.1 is to realize the resolution \widehat{M} as a “generalized connected sum”. For this, denote by $X_{p,q}$ the minimal resolution of $\mathbb{C}^2/\Gamma_{p,q}$. If $E \subset X_{p,q}$ is the exceptional divisor, then $X_{p,q} \setminus E$ is canonically biholomorphic to $(\mathbb{C}^2 \setminus 0)/\Gamma_{p,q}$, and E is the Hirzebruch–Jung string constructed from the continued fraction expansion of p/q as before. The resolution of a $\Gamma_{p,q}$ -singularity at a point $0 \in \overline{M}$ can be realized explicitly as follows. Choose two small positive numbers a and b ; cut off $X_{p,q}$ at a large radius a^{-1} and remove a ball $B(0, b)$ from \overline{M} . If z is an asymptotic uniformizing holomorphic coordinate for $X_{p,q}$ and u is a local uniformizing holomorphic system near $0 \in \overline{M}$, then we can perform the resolution by making the identification $u = abz$.

If this model of \widehat{M} is to carry an “approximately” SFK metric, then it is essential that $X_{p,q}$ should itself carry an asymptotically locally euclidean (ALE) SFK metric. The existence of such metrics is guaranteed by the following:

Proposition 4.1.2. *For all relatively prime positive integers $p < q$, $X_{p,q}$ carries an asymptotically locally euclidean SFK metric g . More precisely, there exists a compact subset $K \subset X_{p,q}$ and holomorphic coordinates z on the universal cover of $X_{p,q} \setminus K$ such that*

$$|g - |dz|^2| = O(|z|^{-1})$$

for all sufficiently large $|z|$. Moreover,

$$|\partial^m g| = O(|z|^{-m-1})$$

for all positive integers m .

Such metrics were constructed in [CS] for general p and q . Note that $X_{q-1,q}$ is the complex manifold underlying the A_{q-1} gravitational instanton of Gibbons–Hawking, Hitchin and Kronheimer [K]. On the other hand, $X_{1,q}$ is the total space of $\mathcal{O}(-q) \rightarrow \mathbb{C}\mathbb{P}^1$, and the SFK metric in this case is due to LeBrun [L2]. For general p and q , these metrics were

known to Joyce, though they do not seem to have appeared explicitly in his published work: they are implicit in [J1] and [J2].

The proof that the metrics of [CS] have the correct asymptotic properties is given in §5.

Returning to our outline of the gluing theorem, we use cut-off functions to define a sequence of approximately SFK metrics on \widehat{M} , by gluing the orbifold metric to the ALE metric of Proposition 4.1.2.

Finally we use the implicit function theorem to find a genuine SFK metric close to the approximate one. The successful completion of this step requires in particular that a certain linear operator be surjective in a controlled way as a and b go to 0. This is why we pay a lot of attention to the linear theory in §4.3. The output of the implicit function theorem is only a $C^{2,\alpha}$ metric. However, any such solution will be smooth by elliptic regularity.

For technical reasons, we shall use the above methods to find hermitian-ASD metrics on \widehat{M} . By a result of Boyer [Bo], any such conformal class must have a Kähler representative, and this will automatically be scalar-flat. Indeed, if ω_0 is any representative 2-form in the conformal class, there is a 1-form β defined by

$$d\omega_0 + \beta \wedge \omega_0 = 0. \quad (4.1)$$

Boyer proves (using the compactness of \widehat{M} , and the fact that $b_1(\widehat{M})$ is even) that $\beta = df$, for some f . But then $d(e^{-f}\omega_0) = 0$ and we have found the desired SFK representative.

We start our analysis now by discussing the perturbation theory of SFK metrics and hermitian-ASD conformal structures.

4.2. Perturbation theory of SFK metrics. Let M be a smooth compact complex surface, with complex structure J . To any hermitian conformal structure c on M we associate the (conformal) fundamental 2-form ω as follows:

$$c(\xi, \eta) \leftrightarrow \omega(\xi, \eta) = c(\xi, J\eta)$$

Here we think of c as a positive-definite section of $\Omega^{-1/2}S^2T^*M$, where Ω is the bundle of densities on M ; accordingly, ω is a weightless $(1,1)$ -form, in other words, a section of the bundle $\Omega^{-1/2}\Lambda^{1,1}$.) Fix a background conformal structure c with corresponding $(1,1)$ -form ω . Then the set of J -hermitian conformal structures near c is identified with a neighbourhood U of 0 in $C^\infty(X, \Omega^{-1/2}\Lambda_0^{1,1})$; if $A \in U$, we have the new weightless $(1,1)$ -form $\omega + A$. A is taken point-wise orthogonal to ω to avoid replacing c by a multiple of c . On a complex surface, $\Lambda_0^{1,1} = \Lambda^-$, so we have parameterized the J -hermitian conformal structures near c by a neighbourhood of 0 in $C^\infty(X, \Omega^{-1/2}\Lambda^-)$.

Denote by L_A the operator $\theta \mapsto (\omega + A) \wedge \theta$, and by Λ_A the adjoint “trace” map. It is shown by Boyer [Bo] that

$$\Lambda_A W^+[\omega + A] = 0 \text{ iff } W^+[\omega + A] = 0.$$

On the LHS we have a section of $\Lambda_{\omega+A}^+ \subset \Lambda^2$. Denote by \mathcal{P} the projection $\Lambda^2 \rightarrow \Lambda^+$ to the *fixed* subspace of 2-forms self-dual with respect to the fixed background c and set

$$\mathcal{F}(A) = \mathcal{P}[\Lambda_A W^+[\omega + A]].$$

(Where necessary, we shall denote the dependence of \mathcal{F} on the background conformal structure c by a sub- or super-script.) The necessary facts about \mathcal{F} are summarized as follows:

Proposition 4.2.1. *Given a fixed J -hermitian conformal structure c , there is a map*

$$\mathcal{F} : U \rightarrow C^\infty(X, \Lambda^+)$$

where U is a neighbourhood of $0 \in C^\infty(\Omega^{-1/2}\Lambda^-)$ with the property that $\mathcal{F}^{-1}(0)$ is the set of J -hermitian ASD conformal structures near c on X . Furthermore, there is an expansion

$$\mathcal{F}(A) = \mathcal{F}(0) + S[A] + \varepsilon_1(A, A \otimes \nabla \nabla A) + \varepsilon_2(A, \nabla A \otimes \nabla A) \quad (4.2)$$

where

$$\mathcal{F}(0) = \Lambda W^+[c], \quad S : C^\infty(\Omega^{-1/2}\Lambda^-) \rightarrow C^\infty(\Lambda^+)$$

is a conformally invariant linear elliptic operator, and the nonlinear terms $\varepsilon_j(A, f)$ are real-analytic in the 0-jet of A and linear in the 0-jet of f .

We note further that if g is a SFK metric in the conformal class c then, trivializing $\Omega^{-1/2}$, the operator S gets identified with the operator

$$S : \alpha \mapsto d^+ \delta \alpha + \langle \rho, \alpha \rangle \omega. \quad (4.3)$$

where ρ is the Ricci form [LS].

4.3. Linear theory. From now on, in order to streamline the discussion, denote by (X_1, g_1) the ALE SFK space from Proposition 4.1.2 and denote by (X_2, g_2) the non-compact space $\overline{M} \setminus \overline{M}_{sing}$. In order to save on notation, we assume that \overline{M} has just one singular point, giving a conical singularity that matches the infinity of X_1 . We know that there exist asymptotic (uniformizing) holomorphic coordinates z , defined, say, for $|z| \geq 1/2$ and such that

$$g_1 = |dz|^2 + \eta_1(z)$$

where

$$|\nabla^m \eta_1|_{g_1} = O(|z|^{-m-1}).$$

Similarly, on X_2 , we have (uniformizing) holomorphic coordinates u , say, again defined for $|u| \leq 2$ with respect to which

$$g_2 = |du|^2 + \eta_2(u), \quad |\eta_2(u)| = O(|u|^2), \quad |\nabla \eta_2(u)| = O(|u|), \quad |\nabla^{m+2} \eta_2(u)| = O(1)$$

for $m = 0, 1, \dots$

Remark. — From now on we shall write, for example, $\{|z| \leq 1\}$ as short-hand for $X_1 \setminus \{|z| > 1\}$; and similarly for subsets of X_2 such as $\{|u| \geq 1\}$.

Denote by S_1 and S_2 the linear operators from Proposition 4.2.1 determined by the metrics g_1 and g_2 . These operators are defined over the non-compact spaces X_1 and X_2 , so some care is needed in arranging for them to be Fredholm. The needed results are now standard, having been worked out in various forms by a number of different authors. The most refined results can be found in Melrose's book [Me]; another useful account is [LM]. This does not cover the Hölder spaces that we shall use, however: for that one can consult Mazzeo's paper on edge operators [Ma].

It turns out that S_1 and S_2 are Fredholm when made to operate between suitably defined weighted Hölder (or Sobolev) spaces. In order to define these Hölder spaces, put

$$r_1 = |z| \text{ for } |z| \geq 2, r_1 = 1 \text{ for } |z| \leq 1/2$$

and $r_1 \geq 1$ everywhere. Define r_2 similarly by continuing

$$r_2 = |u| \text{ for } |u| \leq 1/2$$

to 1 for $|u| \geq 2$, so $r_2 \geq 1/2$ for $|u| \geq 1/2$. Define the norms

$$\|f\|_\alpha = \sup |f| + \sup_{z \neq z'} \left((r_1(z) + r_1(z'))^\alpha \frac{|f(z) - f(z')|}{|z - z'|^\alpha} \right)$$

and

$$\begin{aligned} \|f\|_{2,\alpha} &= \sup |f| + \sup |r_1 \nabla f| + \sup |r_1^2 \nabla \nabla f| \\ &\quad + \sup_{z \neq z'} \left((r_1(z) + r_1(z'))^{2+\alpha} \frac{|\nabla \nabla f(z) - \nabla \nabla f(z')|}{|z - z'|^\alpha} \right) \end{aligned}$$

The completion of $C_0^\infty(X_1)$ in $\|\cdot\|_\alpha$ will be denoted by $B^\alpha(X_1)$; its completion in $\|\cdot\|_{2,\alpha}$ will be denoted by $B^{2,\alpha}$. The weighted versions of these spaces are

$$r_1^\delta B^{n,\alpha} = \{f : r_1^{-\delta} f \in B^{n,\alpha}(X_1)\}, \quad \|f\|_{n,\alpha,\delta} = \|r_1^{-\delta} f\|_{n,\alpha}, \quad (n = 0, 2) \quad (4.4)$$

We define $r_2^\delta B^{n,\alpha}(X_2)$ in exactly the same way, using u and $r_2(u)$ in place of z and $r_1(z)$. Similar norms can be introduced in bundles by patching the local definitions.

The basic facts are then that

$$S_j : r_j^\delta B^{2,\alpha}(X_j, \Omega^{-1/2} \Lambda^-) \rightarrow r_j^{\delta-2} B^{0,\alpha}(X_j, \Lambda^+) \quad (4.5)$$

is a bounded linear operator, Fredholm for all but a discrete set of values of δ . Moreover for any one of these "good" weights, S_j is surjective iff the formal adjoint has no null space acting on $r_j^{-2-\delta} B^{0,\alpha}$.

Let

$$S_0 = d^+ d^* : C^\infty(\mathbb{C}^2 \setminus 0, \Omega^{-1/2} \Lambda^-) \rightarrow C^\infty(\mathbb{C}^2 \setminus 0, \Lambda^+) \quad (4.6)$$

(the linearized operator at the euclidean metric). Denote by x a standard system of euclidean coordinates on \mathbb{C}^2 . The bad weights λ correspond to homogeneous solutions ϕ , $S_0 \phi = 0$, $|\phi| = |x|^\lambda$; more precisely, they correspond to the Γ -invariant such solutions, i.e. the solutions that descend to $(\mathbb{C}^2 \setminus 0)/\Gamma$.

Lemma 4.3.1. *The set of bad weights is contained in $\mathbb{Z} \setminus \{-1\}$.*

Proof. Suppose ϕ satisfies $S_0\phi = 0$ and is homogeneous of degree $\lambda \in \mathbb{C}$. By this we mean that if $\phi = \sum \phi_j e_j$, where e_j is the standard parallel orthonormal basis of $\Omega^{-1/2}\Lambda^-$, then each ϕ_j is homogeneous of degree λ . If $\lambda \in \mathbb{C} \setminus \{-4, -5, -6, \dots\}$, then by [H1, Thm. 3.2.3], ϕ has a unique extension $\dot{\phi}$ to \mathbb{C}^2 as a homogeneous distribution. We have

$$S_0\dot{\phi} = f$$

where f is a distribution supported at 0 homogeneous of degree $\lambda - 2$, because S_0 is of second order. Now any distribution supported at 0 is a finite linear combination of derivatives of the δ -function δ_0 . Since a k -th order derivative of δ_0 is homogeneous of degree $-4 - k$, it follows that

$$\text{If } f \neq 0, \lambda = -2 - k, \text{ where } k = 0, 1, 2, \dots$$

On the other hand, if $f = 0$, then $\dot{\phi}$ must be smooth because S_0 is elliptic. In particular, the degree of homogeneity of ϕ must be a non-negative integer. \square

The other essential fact about the operator S_j in (4.5) is that any ϕ_j with $S_j\phi_j = 0$, must have an asymptotic expansion for $|z| \rightarrow \infty$ or $|u| \rightarrow 0$; moreover, the leading term must behave exactly like $|z|^\lambda$ or $|u|^\lambda$, where λ is one of the ‘‘bad weights’’ found in Lemma 4.3.1.

Proposition 4.3.2. *If $0 < \delta < 2$ and $0 < \alpha < 1$ then*

$$S_1 : r_1^{-\delta} B^{2,\alpha}(X_1, \Omega^{-1/2}\Lambda^-) \rightarrow r_1^{-\delta-2} B^{0,\alpha}(X_1, \Lambda^+)$$

has a bounded right-inverse G_1 . If the orbifold \overline{M} satisfies $\mathfrak{a}_0(\overline{M}) = 0$, then

$$S_2 : r_2^{-\delta} B^{2,\alpha}(X_2, \Omega^{-1/2}\Lambda^-) \rightarrow r_2^{-\delta-2} B^{0,\alpha}(X_2, \Lambda^+)$$

has a bounded right-inverse G_2 .

Proof. For δ in the given range, there are no bad weights, so S_j is Fredholm, and is surjective if and only if

$$S_j^* \psi = 0, |\psi| = O(r_j^{\delta-2}) \Rightarrow \psi = 0.$$

If $j = 1$, then we note that $\delta - 2 < 0$ and so $|\psi| = O(|z|^{-2})$. This is enough to force vanishing of ψ by [KS, Theorem 8.4]. If $j = 2$, we use that $\delta - 2 > -2$, from which it follows that $\psi = O(1)$. By elliptic regularity, it follows that ψ extends to a solution on the whole of the orbifold. To show that $\psi = 0$, we follow the analysis of S that was given in [LS, Thm. 2.7]—the argument goes through without change for compact orbifolds. The essential point is that the component of ψ in the direction of the Kähler form ω is a function f that satisfies Lichnerowicz’s equation

$$\Delta^2 f + 2\langle \rho, dd^c f \rangle = 0.$$

Because \overline{M} has constant scalar curvature, $\nabla^{1,0} f \in \mathfrak{a}_0(\overline{M})$. Thus $f = 0$. One shows further that $f = 0$ implies that the component of ψ orthogonal to ω also vanishes. \square

4.4. Gluing construction. We continue with the notation of the previous section, now picking two small numbers a and b . We begin by gluing X_1 and X_2 to produce the complex manifold $\widehat{M} = X_{a,b}$ which is the resolution of singularities of \overline{M} . This is easy: we just identify an annular region $\{a^{-1} \leq |z| \leq 4a^{-1}\}$ in X_1 with a similar region $\{b \leq |u| \leq 4b\}$ in X_2 by the holomorphic map

$$u = abz. \quad (4.7)$$

4.4.1. Gluing metrics. The next step is to glue the metrics. Pick a standard cut-off function θ_1 , $0 \leq \theta_1 \leq 1$, with $\theta_1 = 1$ for $t \leq 1$, $\theta_1 = 0$ for $t \geq 2$. Set $\theta_2 = 1 - \theta_1$. Define new metrics g_1^a on X_1 and g_2^b on X_2 by “flattening” g_1 near infinity and g_2 near 0; thus $g_1^a = g_1$ for $|z| \leq a^{-1}$ and

$$g_1^a = |dz|^2 + \theta_1(a|z|)\eta_1(z) \text{ for } |z| \geq a^{-1}.$$

Similarly, $g_2^b = g_2$ for $|u| \geq 4b$ and

$$g_2^b = |du|^2 + \theta_2((2b)^{-1}|u|)\eta_2(u) \text{ for } |u| \leq 4b.$$

One computes that any curvature quantity R satisfies

$$|R(g_1^a)| = \begin{cases} |R(g_1)| & \text{for } |z| \leq a^{-1} \\ O(a^3) & \text{for } a^{-1} \leq |z| \leq 2a^{-1} \\ 0 & \text{for } |z| \geq 2a^{-1}. \end{cases} \quad (4.8)$$

Similarly,

$$|R(g_2^b)| = \begin{cases} |R(g_2)| & \text{for } |u| \geq 4b \\ O(1) & \text{for } 2b \leq |u| \leq 4b \\ 0 & \text{for } |u| \leq 2b. \end{cases} \quad (4.9)$$

We now define a metric on $X_{a,b}$ by matching g_1^a with $a^{-2}b^{-2}g_2^b$ by (4.7) along the spheres $\{|u| = 2a^{-1}\}$ and $\{|z| = 2b\}$:

$$g^{a,b} = \begin{cases} g_1^a & \text{for } |z| \leq 2a^{-1} \\ a^{-2}b^{-2}g_2^b & \text{for } |u| \geq 2b. \end{cases} \quad (4.10)$$

It is clear that $g^{a,b}$ is *hermitian* with respect to the complex structure of $X_{a,b}$.

Consider now the map $\mathcal{F}^{a,b}$ of Proposition 4.2.1, associated to $g^{a,b}$ (or more accurately to the underlying conformal structure $c^{a,b}$). The main technical theorem can be stated as follows.

Theorem 4.4.2. *There exist Banach spaces \mathbb{E} and \mathbb{F} (depending upon a and b) such that*

- (i) $\mathcal{F} := \mathcal{F}^{a,b}$ extends to a smooth map from a neighbourhood U of $0 \in \mathbb{E}$ to \mathbb{F} ;
- (ii) for $x \in U$,

$$\mathcal{F}(x) = \mathcal{F}(0) + S[x] + Q(x),$$

where S is a Fredholm linear operator and Q satisfies

$$\|Q(x) - Q(y)\| \leq C_1(\|x\| + \|y\|)\|x - y\|.$$

- (iii) $\|\mathcal{F}(0)\| \rightarrow 0$ as $a, b \rightarrow 0$.
- (iv) If a and b are sufficiently small, then S has a right inverse G , with norm uniformly bounded by C_2 , say.

Once one has this result, one obtains a parameterization of $\mathcal{F}^{-1}(0)$ as a graph of a map f from a small ball in \mathbb{E}_0 into \mathbb{E}_1 , where \mathbb{E}_0 is the finite-dimensional null-space of S and \mathbb{E}_1 is the range of G . Indeed, suppose that

$$\|\mathcal{F}(0)\| \leq \frac{\lambda_0}{C_1 C_2^2}, \quad k_1 = \frac{\lambda_1}{C_1 C_2}, \quad k_2 = \frac{\lambda_2}{C_1 C_2^2}.$$

If we replace x by $x + Gy$, where $x \in \mathbb{E}_0$, then the equation $\mathcal{F}(x + Gy) = 0$ becomes

$$y = T_x(y) := -\mathcal{F}(0) - Q(x + Gy)$$

If x is fixed in $\mathbb{E}_0 \cap \{\|x\| \leq k_1\}$ and $\|y\| \leq k_2$, then

$$\|T_x(y)\| \leq \|\mathcal{F}(0)\| + C_1 \|x + Gy\|^2 \leq \frac{\lambda_0 + \lambda_1^2 + \lambda_2^2}{C_1 C_2^2}$$

Therefore, T_x maps the k_2 -ball in \mathbb{F} into itself if

$$\lambda_0 + \lambda_1^2 + \lambda_2^2 \leq \lambda_2 \tag{4.11}$$

Furthermore,

$$\|T_x(y) - T_x(y')\| = \|Q(x + Gy) - Q(x + Gy')\| \leq 2(\lambda_1 + \lambda_2)\|y - y'\|$$

so that T_x is a contraction mapping if

$$\lambda_1 + \lambda_2 < \frac{1}{2}. \tag{4.12}$$

It is easy to find positive numbers λ_0 , λ_1 and λ_2 that simultaneously satisfy (4.11) and (4.12). It follows from the contraction mapping theorem that there is a unique fixed point $y = T_x(y)$ for any given x in the k_1 -ball of \mathbb{E}_0 , and that

$$\|y\| \leq \frac{1}{1 - 2(\lambda_1 + \lambda_2)} \|\mathcal{F}(0)\|.$$

The norm of the corresponding zero $x + Gy$ of \mathcal{F} satisfies

$$\|x + Gy\| \leq \|x\| + \frac{C_2}{1 - 2(\lambda_1 + \lambda_2)} \|\mathcal{F}(0)\|.$$

4.4.3. *Gluing function spaces.* The functions r_1 and r_2 agree in the gluing region $b \leq |u| \leq 4b$, up to a factor of ab . We therefore define

$$w = w^{a,b} = \begin{cases} r_1(z) & \text{for } |z| \leq 2a^{-1} \\ a^{-1}b^{-1}r_2(u) & \text{for } |u| \geq 2b. \end{cases} \tag{4.13}$$

Note that

$$1 \leq w \leq a^{-1}b^{-1} \text{ on } X^{a,b}. \tag{4.14}$$

Set

$$\begin{aligned} {}^w\|f\|_{n,\alpha} &= \sup |f| + \sup |w\nabla f| + \cdots + \sup |w^n \nabla^n f| + \\ &\quad \sup_{P \neq Q} (w(P) + w(Q))^{n+\alpha} \frac{|\nabla^n f(P) - \nabla^n f(Q)|}{d(P, Q)^\alpha}. \end{aligned} \tag{4.15}$$

Here all lengths are measured by the metric $g^{a,b}$. Denote the completion of $C^\infty(X)$ in this norm by $B^{n,\alpha}(X)$. We now define

$$\mathbb{E} = w^{-\delta} B^{2,\alpha}(X, \Omega^{-1/2} \Lambda^-), \quad \mathbb{F} = w^{-\delta-2} B^{0,\alpha}(X, \Lambda^+).$$

The norms in \mathbb{E} and \mathbb{F} will be denoted by subscripts \mathbb{E} and \mathbb{F} or by the notation

$${}^w \|\cdot\|_{2,\alpha,-\delta}, \quad {}^w \|\cdot\|_{0,\alpha,-2-\delta}.$$

By design \mathbb{E} and \mathbb{F} are closely related to the function spaces that were introduced in Proposition 4.3.2. We make a couple of observations concerning \mathbb{E} and \mathbb{F} . Let $A \in C^\infty(X, \Omega^{1/2} \Lambda^-)$, $W \in C^\infty(X, \Lambda^+)$. If A and W have support contained in $|z| \leq 2/a$, then clearly

$$\|A\|_{\mathbb{E}} = \|A\|_{2,\alpha,-\delta}, \quad \|W\|_{\mathbb{F}} = \|W\|_{0,\alpha,-2-\delta}. \quad (4.16)$$

On the other hand, if A and W are supported in the region $|u| \geq 2b$, then

$$\|A\|_{\mathbb{E}} = a^{-\delta} b^{-\delta} \|A\|_{2,\alpha,-\delta}, \quad \|W\|_{\mathbb{F}} = a^{-\delta} b^{-\delta} \|W\|_{0,\alpha,-2-\delta}. \quad (4.17)$$

Proof of Theorem 4.4.2. Parts (i)–(iii) will follow from the expansion in Proposition 4.2.1. Let us begin by computing the \mathbb{F} -norm of $\mathcal{F}(0)$. This is zero away from the gluing region $b \leq |u| \leq 4b$, and can be estimated there by combining (4.8) and (4.9). The result is

$$\|\mathcal{F}(0)\|_{\mathbb{F}} = O(a^{1-\delta} + a^{-\delta} b^2). \quad (4.18)$$

Thus to guarantee (iii), we shall need to choose $0 < \delta < 1$ and make a suitable choice of the relative sizes of a and b , for example $a = b^2$.

In order to show that S extends to a bounded linear map from \mathbb{E} to \mathbb{F} , use a partition of unity to split $S(A)$ into two pieces, the relations (4.16) and (4.17), and the fact that the operators S_1 and S_2 in Proposition 4.3.2 are bounded; here it is important that the same scale factor appears in comparing the norms of A and W in (4.17).

Finally we turn to the non-linear terms in (4.2). These are only defined if $\sup |A|$ is sufficiently small. But

$$\sup |A| \leq \sup w^\delta |A| \leq \|A\|_{\mathbb{E}} \quad (4.19)$$

by (4.14). Assuming, then, that $\|A\|_{\mathbb{E}}$ is sufficiently small, we have

$$\sup |w^{2+\delta} \varepsilon_1(A, A \otimes \nabla^2 A)| \leq \sup |\varepsilon_1(A, w^\delta A \otimes w^{2+\delta} \nabla^2 A)| \leq C \|A\|_{\mathbb{E}}^2,$$

again using (4.14). Since ε_1 is linear in its second variable, we have

$$\begin{aligned} \varepsilon_1(A, A \otimes \nabla^2 A) - \varepsilon_1(B, B \otimes \nabla^2 B) &= \varepsilon_1(A, A \otimes \nabla^2(A - B)) + \varepsilon_1(A, (A - B) \otimes \nabla^2 B) + \\ &\quad + \varepsilon_1(A, B \otimes \nabla^2 B) - \varepsilon_1(B, B \otimes \nabla^2 B) \end{aligned}$$

and it is straightforward to use this to prove that the Hölder quotient in the definition of \mathbb{F} is controlled by $\|A\|_{\mathbb{E}}$ and also the quadratic estimate of Q in (ii). The term $\varepsilon_2(A, \nabla A \otimes \nabla A)$ is estimated in the same way.

It remains to prove that S has a uniformly bounded right inverse. Note that to save on notation, we have not indicated explicitly that S depends upon a and b . One should not lose sight of this dependence, however, in what follows.

First note that if a and b are small, then the operator norms of $S_1 - S_1^a$ and $S_2 - S_2^b$ are small, and so there are operators G_1^a, G_2^b such that

$$S_1^a G_1^a = 1, \quad S_2^b G_2^b = 1, \quad \|G_1 - G_1^a\| \leq 1/2, \quad \|G_2 - G_2^b\| \leq 1/2.$$

(Here the operator norms are defined by viewing S_j as operators between the Banach spaces of Proposition 4.3.2.) We shall now follow the approach of [DK, Chapter 7] to splice these right-inverses to give first an approximate and then an exact right-inverse for S . Recall the partition of unity $\theta_1 + \theta_2 = 1$ on \mathbb{R} that was introduced at the beginning of §4.4.1. For any small positive number λ , which will be fixed later, define

$$\beta_1(z) = \theta_1 \left((a|z|/4)^\lambda \right), \quad \beta_2(u) = \theta_2 \left(2(|u|/2b)^\lambda \right). \quad (4.20)$$

Define also

$$\gamma_1(z) = \theta_1(a|z|/2), \quad \gamma_2(u) = \theta_2(|u|/2b). \quad (4.21)$$

Then $\gamma_1 + \gamma_2 = 1$ on $X^{a,b}$, and

$$\beta_1 \gamma_1 = \gamma_1, \quad \beta_2 \gamma_2 = \gamma_2. \quad (4.22)$$

Moreover, we have estimates of the form

$$\sup |r_1^k \nabla^k \beta_1| = O(\lambda^k), \quad \sup |r_2^k \nabla^k \beta_2| = O(\lambda^k) \quad (4.23)$$

for each positive integer k , with the O 's uniform in a and b .

Now form the operator on $X^{a,b}$,

$$G_0 = \beta_1 G_1^a \gamma_1 + \beta_2 G_2^b \gamma_2; \quad (4.24)$$

we claim first that $\|G_0\|$, regarded as an operator $\mathbb{F} \rightarrow \mathbb{E}$, is bounded independent of a and b . Indeed, we have

$$\begin{aligned} \|G_0 W\|_{\mathbb{E}} &\leq \|\beta_1 G_1^a (\gamma_1 W)\|_{\mathbb{E}} + \|\beta_2 G_2^b (\gamma_2 W)\|_{\mathbb{E}} \\ &= \|\beta_1 G_1^a (\gamma_1 W)\|_{2,\alpha,-\delta} + (ab)^{-\delta} \|\beta_2 G_2^b (\gamma_2 W)\|_{2,\alpha,-\delta} \end{aligned}$$

using the scaling formulae (4.16) and (4.17). Since the operator norms of G_1^a and G_2^b are uniformly bounded, we obtain

$$\|G_0 W\|_{\mathbb{E}} \leq C[\|\gamma_1 W\|_{\mathbb{F}} + \|\gamma_2 W\|_{\mathbb{F}}] \leq C' \|W\|_{\mathbb{F}}$$

since $\gamma_1 + \gamma_2 = 1$.

On the other hand, G_0 is an approximate right-inverse for S :

$$S G_0 = \beta_1 S G_1^a \gamma_1 + \beta_2 S^{a,b} G_2^a \gamma_2 + [S^{a,b}, \beta_1] G_1^a \gamma_1 + [S^{a,b}, \beta_2] G_2^a \gamma_2.$$

On the support of β_1 , S is close to S_1^a , if b is small. Therefore,

$$\beta_1 S G_1^a \gamma_1 = \beta_1 S_1^a G_1^a \gamma_1 + o(1) = \beta_1 \gamma_1 + o(1) = \gamma_1 + o(1).$$

(Here $o(1)$ indicates an operator whose norm tends to zero with a and b .) Similarly,

$$\beta_2 S G_2^b \gamma_2 = \gamma_2 + o(1).$$

We show now that the commutator terms are $O(\lambda)$. Clearly $[S, \beta_1]$ is a first-order differential operator supported where $\nabla \beta_1 \neq 0$, i.e. the interval $[(4/a), (4/a) \cdot 2^{1/\lambda}]$. For fixed λ and small a , S is very close to S_2^b over this interval, and we can estimate norms as before by passing from \mathbb{E} and \mathbb{F} to the corresponding fixed weighted Hölder spaces on

X_2 . The coefficients of $[S, \beta_1]$ are therefore bounded functions on X_2 , multiplied by $\nabla\beta_1$ and $\nabla^2\beta_1$. Hence by (4.23) the operator norm of $[S, \beta_1]$ is $O(\lambda)$, as claimed. The same is true of the other commutator term. In sum, we have found that if λ is chosen small enough, then as a and b tend to zero,

$$SG_0 = 1 + R, \quad \|R\| \leq 1/2$$

Hence $G = G_0(1 + R)^{-1}$ is a controlled right-inverse of S , and the proof of Theorem 4.4.2 is complete.

Applying the implicit function theorem as described above, we obtain solutions in \mathbb{E} of our equation for all sufficiently small a, b , subject to the constraints imposed by (4.18), and by elliptic regularity these solutions will be smooth.

Finally we claim that the corresponding SFK metric is close to the original one. More precisely, it is close to g_1 on any subset of the form $\{|z| \leq C_1\}$ of X_1 , and it is close to $a^{-2}b^{-2}g_2$ on any subset of the form $\{|u| \geq C_2\}$ of X_2 . From the implicit function theorem, our solution has fundamental 2-form

$$\omega^{a,b} + A$$

where $\omega^{a,b}$ is the fundamental 2-form of $g^{a,b}$ and

$$\|A\|_{\mathbb{E}} = O(a^{1-\delta} + a^{-\delta}b^2).$$

Choose $b^2 = a$, $\delta = 1/2$, so that

$$\sup |w^\delta A| + \sup |w^{\delta+1} \nabla A| + \sup |w^{\delta+2} \nabla^2 A| = O(b)$$

Then the equation (4.1) becomes

$$d(\omega^{a,b} + A) + \beta \wedge (\omega^{a,b} + A) = 0.$$

and so

$$\sup w^{\delta+1} |\beta| = O(b).$$

If we define a function f with $df = \beta$ by integrating along curves starting from any base-point fixed in the region $|z| \leq 1/2$ in X_1 , it is not hard to see that such f satisfies

$$\sup |f| = O(b)$$

so that the Kähler form ω_{SFk} satisfies

$$\omega_{SFk} = \omega^{a,b} + O(b)$$

as claimed.

4.5. Summary: the proof of Theorem 2.1.2. Let us explain first how our work applies to prove that \widehat{M} carries a SFK metric. By Theorem 3.3.1, \widehat{M} is the minimal resolution of singularities of the orbifold \overline{M} . By Theorem 3.4.1, \overline{M} is SFK, and $\mathfrak{a}_0(\overline{M}) = 0$. Hence by Theorem 4.1.1, \widehat{M} carries SFK metrics.

Next let us note how to handle blow-ups of \widehat{M} . The whole argument of this section goes through to handle ordinary blow-ups: there is a SFK metric on blow-up of the origin of \mathbb{C}^2 ; this is the ‘‘Burns metric’’, but it also arises by taking $p = 0$, $q = 1$ in

Proposition 4.1.2. The only thing to check is that $\mathfrak{a}_0(\widehat{M}) = 0$, but this follows from the fact that $\mathfrak{a}_0(\overline{M}) = 0$.

5. ASYMPTOTICS OF THE ALE SCALAR-FLAT KÄHLER METRICS

This section is devoted to a proof of Proposition 4.1.2. We shall make extensive use of the notation of [CS].

Recall that the metric on $X_{p,q}$ is determined by a choice of real numbers $y_0 > y_1 > y_2 > \cdots > y_k > y_{k+1} = 0$, and the pairs (m_j, n_j) coming from the continued fraction expansion of p/q . Here $(m_{k+1}, n_{k+1}) = (q, p)$ and we define $(m_{k+2}, n_{k+2}) = (0, 1)$. For comparison with [CS], we note that p and q have been reversed and we normalize y_{k+1} to be 0. We shall also use half-space coordinates $x > 0$ and y , (rather than (ρ, η)) so the hyperbolic metric becomes $x^{-2}(dx^2 + dy^2)$.

If

$$(a_j, b_j) = (m_j - m_{j+1}, n_j - n_{j+1}) \text{ for } j = 0, 1, \dots, k+1,$$

then we can define

$$v_1 = \frac{x}{2} \sum_{j=0}^{k+1} \frac{(a_j, b_j)}{\sqrt{x^2 + (y - y_j)^2}} = \frac{x}{2\sqrt{x^2 + y^2}}(q, p - 1) + \frac{x}{2} \sum_{j=0}^k \frac{(a_j, b_j)}{\sqrt{x^2 + (y - y_j)^2}}, \quad (5.1)$$

and

$$v_2 = \frac{1}{2} \sum_{j=0}^{k+1} \frac{(y - y_j)(a_j, b_j)}{\sqrt{x^2 + (y - y_j)^2}} = \frac{1}{2\sqrt{x^2 + y^2}}(q, p - 1) + \frac{1}{2} \sum_{j=0}^k \frac{(y - y_j)(a_j, b_j)}{\sqrt{x^2 + (y - y_j)^2}}, \quad (5.2)$$

giving an ALE scalar-flat Kähler metric

$$g = \frac{x|\langle v_1, v_2 \rangle|}{x^2 + y^2} \left(\frac{dx^2 + dy^2}{x^2} + \frac{\langle v_1, dt \rangle^2 + \langle v_2, dt \rangle^2}{\langle v_1, v_2 \rangle^2} \right) \quad (5.3)$$

where t is an T^2 -valued flat coordinate and $\langle \cdot, \cdot \rangle$ denotes the standard symplectic form on \mathbb{R}^2 .

The asymptotic region of this metric corresponds to the point $(x, y) = (0, 0)$ in the boundary of the hyperbolic plane. We shall analyze this metric first by introducing coordinates

$$R^2 e^{2i\theta} = \frac{1}{y - ix} \quad (5.4)$$

so that

$$x = R^{-2} \sin 2\theta, \quad y = R^{-2} \cos 2\theta, \quad R > 0, 0 \leq \theta \leq \pi/2. \quad (5.5)$$

Before attempting the computation of this metric in these coordinates, it is worth writing the standard metric on \mathbb{C}^2 in analogous coordinates. Namely, if $(\tilde{z}_1, \tilde{z}_2)$ are standard complex coordinates, let

$$\tilde{z}_1 = R e^{i\phi} \cos \theta, \quad \tilde{z}_2 = R e^{i\psi} \sin \theta, \quad R > 0, 0 \leq \theta \leq \pi/2. \quad (5.6)$$

Then the standard euclidean metric becomes

$$|d\tilde{z}_1|^2 + |d\tilde{z}_2|^2 = dR^2 + R^2 d\theta^2 + R^2 (\cos^2 \theta d\phi^2 + \sin^2 \theta d\psi^2).$$

Now return to the metric (5.3). We shall try to understand it for $R \gg 0$. Referring first to (5.1) we note that each term in the sum from 0 to k is $O(R^{-2})$, so

$$v_1 = \frac{\sin 2\theta}{2}(q, p - 1) + O(R^{-2}). \quad (5.7)$$

The j -th term in the sum in (5.2) is $-(a_j, b_j) + O(R^{-2})$, so

$$v_2 = \frac{1}{2}(1 + \cos 2\theta)(q, p - 1) + (0, 1) + O(R^{-2}). \quad (5.8)$$

Now define new angular variables

$$\psi = t_1/q, \quad \phi = (p/q)t_1 - t_2. \quad (5.9)$$

The fact that the determinant is q^{-1} means that (ϕ, ψ) really live on a q -fold cover.

$$\langle v_1, dt \rangle = q \sin \theta \cos \theta (d\psi - d\phi) + O(R^{-2}) \quad (5.10)$$

and

$$\langle v_2, dt \rangle = -q \sin^2 \theta d\psi - q \cos^2 \theta d\phi + O(R^{-2}) \quad (5.11)$$

Hence the ‘‘angular part’’ of the metric is given by

$$\langle v_1, dt \rangle^2 + \langle v_2, dt \rangle^2 = q^2 \sin^2 \theta d\psi^2 + q^2 \cos^2 \theta d\phi^2 + O(R^{-2}) \quad (5.12)$$

On the other hand, from (5.5),

$$y - ix = R^{-2} e^{-2i\theta}, \quad dx^2 + dy^2 = 4R^{-4}(dR^2 + R^2 d\theta^2). \quad (5.13)$$

Since

$$\langle v_1, v_2 \rangle = \frac{q}{2} \sin 2\theta + O(R^{-2}) \quad (5.14)$$

we obtain for the metric

$$2q (dR^2 + R^2 d\theta^2 + R^2[\alpha^2 \sin^2 \theta + \beta^2 \cos^2 \theta]) + \text{lower order terms}. \quad (5.15)$$

Here the ‘‘lower order terms’’ are just terms that are of order 2 less than in the given expression for the metric (i.e. there are terms like $O(R^{-2})dR^2$ and $O(1)\alpha^2$ etc.)

Thus the metric, pulled back to a uniformizing chart, differs from a constant multiple of the standard flat metric by tensors of order R^{-2} , as measured by the flat metric.

5.1. The complex structure. The metric (5.3) is Kähler with respect to the complex structure J ,

$$Jdt = \frac{1}{\sqrt{x^2 + y^2}} \left((xv_1 - yv_2) \frac{dx}{x} + (yv_1 + xv_2) \frac{dy}{x} \right). \quad (5.16)$$

It follows that the two components of $dt + iJdt$ are $(1, 0)$ -forms. In fact, a simple calculation shows that these components are closed, hence holomorphic.

With respect to the coordinates introduced above, we can rewrite (5.16) as follows

$$Jdt = -\frac{v_1}{\sin \theta \cos \theta} d \log R - \frac{v_2}{\sin \theta \cos \theta} d\theta. \quad (5.17)$$

(To check this, collect the terms in v_1 and v_2 before changing variables.) Now with ψ and ϕ as before, one obtains from (5.7) and (5.8) that

$$\omega_1 := d[i\phi + \log R + \cos \theta] + O(1/R^2) \quad \text{and} \quad \omega_2 := d[i\psi + \log R + \sin \theta] + O(1/R^2);$$

are closed holomorphic 1-forms. Now shift to the complex coordinates $(\tilde{z}_1, \tilde{z}_2)$, which are *not* J -holomorphic. We have

$$\omega_j = \frac{d\tilde{z}_j}{\tilde{z}_j} + F_j(\tilde{z}_1, \tilde{z}_2)$$

where the 1-forms F_j are $O(1/R^2)$. Now set

$$f_j(\tilde{z}_1, \tilde{z}_2) = - \int_{(\tilde{z}_1, \tilde{z}_2)}^{\infty} F_j$$

where the path of integral is $t \mapsto (t\tilde{z}_1, t\tilde{z}_2)$, t from 1 to ∞ . Then $df_j = F_j$ and $z_j := \tilde{z}_j \exp(f_j)$ are J -holomorphic coordinates near infinity close to the standard ones:

$$|z_j - \tilde{z}_j| = O(1/R) \text{ for } R \gg 0,$$

since $f_j = O(1/R)$. Re-expanding the metric in the coordinates (z_1, z_2) will add some $1/R$ -terms to the expansion to (5.15), but this is sufficient to complete the proof of Proposition 4.1.2.

REFERENCES

- [BB] D. Burns, P. de Bartolomeis, *Stability of vector bundles and extremal metrics*, Invent. Math. **92**, 403–407 (1988)
- [Be] A.L. Besse, *Einstein manifolds*, Ergebn. Math. (3) **10**, Springer–Verlag (1987)
- [Bea] A. Beauville, *Complex algebraic surfaces*
- [BPV] W. Barth, C. Peters, A. Van de Ven, *Compact complex surfaces*, Springer (1984).
- [Bi] O. Biquard, *Fibrés paraboliques stables et connexions singulières plates*, Bull. Soc. Math. Fr. **119**, 231–257 (1991)
- [Bo] C. Boyer, *Conformal duality and complex compact surfaces*, Math. Ann., **274** (1986), 517–526
- [CS] D. Calderbank, M. Singer, *Einstein metrics and complex singularities*, Invent. Math. 156 (2004), no. 2, 405–443
- [CT] X.X. Chen, G. Tian, *Geometry of Kähler metrics and holomorphic foliation by discs*, arXiv math.DG/0409433
- [D] S. Donaldson, *Scalar curvature and projective embeddings I*, J. of Diff. Geom. **59** (2001), 479–522.
- [DK] S. Donaldson, P. Kronheimer, *Geometry of 4-manifolds*, Oxford.
- [Fu] W. Fulton, *Toric Varieties*, Princeton University Press.
- [H1] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, Second Edition, Springer (1990).
- [J1] D.D. Joyce, *The hypercomplex quotient and the quaternionic quotient*, Math. Ann. **290** (1991), 323–340.
- [J2] D.D. Joyce, *Explicit construction of self-dual 4-manifolds*, Duke Math. J. **77** (1995), 519–552.
- [KLP] J. Kim, C. LeBrun, M. Pontecorvo, *Scalar-flat Kähler surfaces of all genera*, J. Reine Angew. Math. **486** (1997), 69–95
- [KS] A. Kovalev, M. Singer, *Gluing theorems for complete anti-self-dual spaces*, Geom. Funct. Anal. **11** (2001), no. 6, 1229–1281
- [K] P.B. Kronheimer, *The construction of ALE spaces as hyper-Kähler quotients*, J. Diff. Geom. **29** (1989), 665–683.
- [L1] C. LeBrun, *On the topology of self-dual 4-manifolds*. Proc. Amer. Math. Soc. **98** (1986), no. 4, 637–640.
- [L2] C. LeBrun, *Counter-examples to the generalized positive action conjecture*, Comm. Math. Phys., **118** (1988), 591–596.
- [L3] C. LeBrun, *Scalar-flat Kähler metrics on blown-up ruled surfaces*, J. Reine Angew. Math. **420** (1991), 161–177.

- [L4] C. LeBrun, Anti-self-dual metrics and Kähler geometry, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), 498–507, Birkhäuser, (1995).
- [LS] C. LeBrun, M. Singer, *Existence and deformation theory for scalar-flat Kähler metrics on compact complex surfaces*, Invent. Math. **112** (1993), no. 2, 273–313
- [LM] R.B. Lockhart, R.C. McOwen, *Elliptic differential operators on non-compact manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **12** (1985), 409–447.
- [Me] R.B. Melrose, The Atiyah–Patodi–Singer index theorem. A K Peters Ltd, Wellesley, MA, 1993.
- [Ma] R. Mazzeo, *Elliptic theory of differential edge operators. I.*, Comm. Partial Differential Equations, **16** (1991), 1615–1664.
- [McO] R. McOwen, *Prescribed curvature and singularities of conformal metrics on Riemann surfaces*, J. Math. Anal. Appl. **177** (1993), 287–298.
- [MS] V.B. Mehta, C.S. Seshadri, *Moduli of vector bundles of curves with parabolic structures*, Math. Ann. **248**, 205–239 (1980)
- [NS] M.S. Narasimhan, C.S. Seshadri, *Stable and unitary vector bundles on a compact Riemann surface*, Ann. Math. (2) **82**, 540–564 (1965)
- [RT] J. Ross, R. Thomas, *Slope stability of projective varieties*, Preprint (2004)
- [Tr] M. Troyanov, *Prescribing curvature on compact surfaces with conical singularities*, Trans. Amer. Math. Soc. **324** (1991), no. 2, 793–821
- [TV] G. Tian, J. Viaclovsky, *Moduli spaces of critical Riemannian metrics in dimension four*, Arxiv math.DG/0312318 (2003)

YANN ROLLIN, MIT, 77 MASSACHUSETTS AVENUE, CAMBRIDGE MA 02139, USA
E-mail address: rollin@math.mit.edu

MICHAEL SINGER, SCHOOL OF MATHEMATICS, KING'S BUILDINGS, EDINBURGH SCOTLAND
E-mail address: msinger@math.mit.edu