SPECTRAL COMPUTATIONS ON LAMPLIGHTER GROUPS AND DIESTEL-LEADER GRAPHS

LAURENT BARTHOLDI AND WOLFGANG WOESS

ABSTRACT. The Diestel-Leader graph DL(q,r) is the horocyclic product of the homogeneous trees with respective degrees q+1 and r+1. When q=r, it is the Cayley graph of the lamplighter group (wreath product) $\mathbb{Z}_q \wr \mathbb{Z}$ with respect to a natural generating set. For the "Simple random walk" (SRW) operator on the latter group, GRIGORCHUK AND ŻUK and DICKS AND SCHICK have determined the spectrum and the (on-diagonal) spectral measure (Plancherel measure). Here, we show that thanks to the geometric realization, these results can be obtained for all DL-graphs by directly computing an ℓ^2 -complete orthonormal system of finitely supported eigenfunctions of the SRW. This allows computation of all matrix elements of the spectral resolution, including the Plancherel measure. As one application, we determine the sharp asymptotic behaviour of the N-step return probabilities of SRW. The spectral computations involve a natural approximating sequence of finite subgraphs, and we study the question whether the cumulative spectral distributions of the latter converge weakly to the Plancherel measure. To this end, we provide a general result regarding $F\emptyset$ liner approximations; in the specific case of DL(q,r), the answer is positive only when r=q.

1. Introduction

Let X be a locally finite connected graph. Simple random walk (SRW) on X is the Markov chain on X with transition probabilities

$$p(x,y) = \begin{cases} 1/\deg(x) & \text{if } y \sim x, \\ 0 & \text{otherwise.} \end{cases}$$

Here, \sim denotes neighbourhood, and $\deg(x)$ is the degree (number of neighbours) of vertex $x \in X$. The transition operator associated with SRW acts on real or complex functions f on X by

$$Pf(x) = \sum_{y} p(x, y) f(y).$$

In particular, P acts as a self-adjoint operator on the weighted ℓ^2 -space $\ell^2(X, \deg)$ and has norm = spectral radius ≤ 1 . Here, we shall only consider regular graphs, i.e., deg is constant, and we prefer to use the ordinary space $\ell^2(X)$, where the reference measure is the counting measure (instead of $m(x) = \deg(x)$).

Date: May 11, 2004.

 $^{2000\} Mathematics\ Subject\ Classification.\ 05C50,\ 20E22,\ 47A10,\ 60B15.$

Key words and phrases. lamplighter group, wreath product, Diestel-Leader graph, random walk, spectrum, spectral measures.

Supported by FWF (Austrian Science Fund) project P15577.

Associated with P there is the resolution of the identity. This is an operator-valued measure E defined on $\operatorname{spec}(P)$ such that for all $N \geq 0$,

$$P^N = \int_{\operatorname{spec}(P)} \lambda^n E(d\lambda)$$

It is characterized by the matrix elements

(1.1)
$$\mu_{x,y}(d\lambda) = \langle \delta_x, E(d\lambda)\delta_y \rangle,$$

which in turn are characterized by their moments, which are the N-step transition probabilities (matrix elements of P^N),

$$(1.2) p^{(N)}(x,y) = \int_{\operatorname{spec}(P)} \lambda^N \, \mu_{x,y}(d\lambda) \quad \forall \ N \ge 0 \, .$$

Each $\mu_{x,x}$ is a probability measure, while the off-diagonal $\mu_{x,y}$ are signed measures with total mass 0. When X is a transitive graph (i.e., its group of isometries acts transitively on the vertex set), then all $\mu_{x,x}$ coincide, and we shall just write μ for this measure, whose support is the whole spectrum. This holds, in particular, for Cayley graphs of finitely generated groups. In the spirit of Harmonic Analysis, we call μ the Plancherel measure; more recently, it has also been called the Kesten spectral measure by some authors.

Basic references for the general theory of spectra of *infinite* graphs and groups are the papers of Mohar and Woess [20], de la Harpe, Robertson and Valette [12] and Grigorchuk and Żuk [13]. In [20] and [12], one can also find many examples of specific graphs and groups where spectra and (less frequently) spectral measures are computed: in basic cases $\operatorname{spec}(P)$ is an interval, and the Plancherel measure has a continuous density with respect to Lebesgue measure. This occurs for integer lattices – a classical result from Fourier analysis, see e.g. Pólya [22] – and for free groups, resp. homogeneous trees – see Kesten [15] and Cartier [5]. For other tree-like cases (infinite distance-regular graphs), the situation is almost the same, with a possible additional isolated eigenvalue, as was shown by Kuhn and Soardi [17]; see also Faraut and Picardello [11]. For \tilde{A}_d -buildings with arbitrary d, the situation is similar to that of homogeneous trees, see Cartwright and Młotkowski [7] and Cartwright [6].

The situation is different on typical fractal graphs such as the one associated with the Sierpiński gasket, where the spectrum is *pure point*, i.e., the closure of the set of eigenvalues of P, see Malozemov and Teplyaev [18], Teplyaev [27], Sabot [24] and, for a generalization, Krön [16]. These graphs are regular, but far from being transitive. However, it turned out in recent research that a similar situation may also occur in certain classes of *fractal groups* related with the construction of Grigorchuk and Gupta and Sidki of finitely generated infinite torsion groups with intermediate growth. For a comprehensive survey of these groups and their properties, see Bartholdi, Grigorchuk and Nekrashevych [2] and the references given there, and for the specific computation of a pure point spectrum on such a group, see Bartholdi and Grigorchuk [1].

Coming finally to the types of structures considered in the present paper, a pure point spectrum with the associated Plancherel (Kesten) measure was recently detected for a

¹If Γ is a group and $S = S^{-1} \subset \Gamma$ a finite set of generators, then the Cayley graph $X(\Gamma, S)$ has vertex set Γ , and $x \sim y$ if $x^{-1}y \in S$.

different class of groups, namely the lamplighter groups (wreath products) $F \wr \mathbb{Z}$, where F is a finite group; see GRIGORCHUK AND \dot{Z} UK [14] for $\mathbb{Z}_2 \wr \mathbb{Z}$, and DICKS AND SCHICK [8] for the general case². While [14] uses approximation of the considered Cayley graph by an increasing sequence of finite graphs, [8] applies von-Neumann-algebraic methods.

Note that the present study of random walks on $F \wr \mathbb{Z}$ depends only on the cardinality |F|, whence it suffices to consider $F = \mathbb{Z}_q$. Now, the Cayley graphs of $\mathbb{Z}_q \wr \mathbb{Z}$ considered in [14] and [8] turn out to have geometric realizations as specific examples DL(q,q) in the family of *Diestel-Leader graphs* introduced by DIESTEL AND LEADER [9], i.e., the horocyclic products DL(q,r) of two homogeneous trees with degree q + 1 and r + 1, respectively; see WOESS [29] and §2 below.

In the present paper, we exploit this geometric model to provide in §3 a completely explicit and elementary construction of an ℓ^2 -complete orthonormal system of finitely supported eigenfunctions of the SRW-operator on DL(q,r). This comprises the lamplighter groups, but holds more generally for all Diestel-Leader graphs. When $q \neq r$, the graph DL(q,r) is transitive, but not a Cayley graph, whence the group-specific methods of [8] do not apply here (they rely on identifying eigenfunctions as projections in the von Neumann algebra of a group acting on the graph; here that group is non-discrete and non-unimodular), nor can one use an approximating sequence of Schreier graphs as in [14].

We recover the spectral radius of SRW on DL(q,r), computed previously by SALOFF-COSTE AND WOESS [25],

(1.3)
$$\rho(P) = \frac{2\sqrt{qr}}{q+r}.$$

The spectrum $\operatorname{\mathsf{spec}}(P) = [-\rho(P)\,,\,\rho(P)]$ is the closure of the set of all eigenvalues

$$\left\{\lambda_{m,n} = \rho(P)\cos\frac{m}{n}\pi : n \ge 2, \ 1 \le m \le n-1\right\},\,$$

see Theorem 3.15. This theorem can be used to find an expression for each of the spectral measures, see §4. In particular, the Plancherel measure can be computed explicitly (Corollary 4.6).

We then use the latter in §5 to determine the sharp asymptotic behaviour of the 2N-step return probabilities $p^{(2N)}(x,x)$, as $N \to \infty$, see Theorem 5.4 (note that $p^{(2N+1)}(x,x) = 0$ since DL(q,r) is bipartite). For the lamplighter groups, i.e., on DL(q,q), these asymptotics have been determined for almost the same random walk by Revelle [23]; one has

$$p^{(2N)}(x,x) \sim \bar{A}_1 \exp(-B_1 N^{1/3}) N^{1/6} \qquad (r=q).$$

It is interesting to note that in the case $r \neq q$, the last (polynomial) term changes by a factor of N:

$$p^{(2N)}(x,x) \sim A_1 \rho(P)^{2N} \exp(-B_1 N^{1/3}) N^{-5/6} \qquad (r \neq q).$$

Furthermore, the constants $\bar{A}_1, A_1, B_1 > 0$ are determined explicitly as functions of q and r.

²We denote by \mathbb{Z}_q the cyclic group $\mathbb{Z}/q\mathbb{Z}$ of order q

Next, in §6, we discuss for general vertex transitive graphs under which conditions the cumulative spectral measures of an approximating sequence of finite graphs converge (resp. do not converge) weakly to the Plancherel measure. A positive criterion is given in terms of $F \not older sequences$ of approximating subgraphs, and more generally, $F \not older approximations$ – see Theorem 6.3 and Remark 6.4. In our setting, this applies to DL(q,q) with the natural subgraphs (tetrahedra) used in §3 for computing the spectrum, since they constitute a $F \not older olde$

At the end, in $\S 7$, we add several observations, including random walks with drift, the corresponding spectra, their return probabilities, and also their projections on the two subtrees and on \mathbb{Z} .

2. Lamplighter groups and Diestel-Leader graphs

This section is a short version of §2 in [29]. We explain the structure of the DL-graphs and their relation with the groups $\mathbb{Z}_q \wr \mathbb{Z}$.

Let $\mathbb{T} = \mathbb{T}_q$ be the homogeneous tree with degree q+1, $q \geq 2$. A geodesic path, respectively geodesic ray, respectively infinite geodesic in \mathbb{T} is a finite, respectively one-sided infinite, respectively doubly infinite sequence (x_n) of vertices of \mathbb{T} such that $d(x_i, x_j) = |i-j|$ for all i, j, where $d(\cdot, \cdot)$ denotes the graph distance.

Two rays are equivalent if the symmetric difference of their supports is finite. An end of \mathbb{T} is an equivalence class of rays. The space of ends is denoted $\partial \mathbb{T}$, and we write $\widehat{\mathbb{T}} = \mathbb{T} \cup \partial \mathbb{T}$. For all $w, z \in \widehat{\mathbb{T}}$ there is a unique geodesic \overline{wz} that connects the two. In particular, if $x \in \mathbb{T}$ and $\xi \in \partial \mathbb{T}$ then $\overline{x\xi}$ is the ray that starts at x and represents ξ .

For $x, y \in \mathbb{T}$, $x \neq y$, we define the cone $\widehat{\mathbb{T}}(x, y) = \{w \in \widehat{\mathbb{T}} : y \in \overline{xw}\}$. The collection of all cones is the basis of a topology which makes $\widehat{\mathbb{T}}$ a compact, totally disconnected Hausdorff space with \mathbb{T} as a dense, discrete subset.

We fix a root $o \in \mathbb{T}$. If $w, z \in \widehat{\mathbb{T}}$, then their confluent $c = w \wedge z$ with respect to the root vertex o is defined by $\overline{ow} \cap \overline{oz} = \overline{oc}$. Similarly, we choose and fix a reference end $\omega \in \partial \mathbb{T}$. For $z, v \in \widehat{\mathbb{T}} \setminus \{\omega\}$, their confluent $b = v \wedge z$ with respect to ω is defined by $\overline{v\omega} \cap \overline{z\omega} = \overline{b\omega}$. We write

$$z \leq v$$
 if $z \wedge v = z$.

For $x \in \mathbb{T}$, we describe its relative position with respect to o by the two numbers

$$\mathfrak{u}(x) = d(o, x \curlywedge o)$$
 and $\mathfrak{d}(x) = d(x, x \curlywedge o)$.

In Figure 1, $\mathfrak{u}(x)$ and $\mathfrak{d}(x)$ correspond to the numbers of steps one has to take upwards (in direction of ω), respectively downwards, on the geodesic path from o to x. Thus, $d(x,o) = \mathfrak{u}(x) + \mathfrak{d}(x)$.

The Busemann function $\mathfrak{h}: \mathbb{T} \to \mathbb{Z}$ and the horocycles H_k with respect to ω are defined as

$$\mathfrak{h}(x) = \mathfrak{d}(x) - \mathfrak{u}(x)$$
 and $H_k = \{x \in \mathbb{T} : \mathfrak{h}(x) = k\}$.

Every horocycle is infinite. Every vertex x in H_k has one neighbour x^- (its predecessor) in H_{k-1} and q neighbours (its successors) in H_{k+1} . We set $\partial^* \mathbb{T} = \partial \mathbb{T} \setminus \{\omega\}$.

We label each edge of \mathbb{T} by an element of \mathbb{Z}_q such that for each vertex, the "downward" edges to its q successors carry labels $0, \ldots, q-1$ from left to right (say), see Figure 1. Thus, for each $x \in \mathbb{T}$, the sequence $(\sigma(n))_{n \leq 0}$ of labels on the geodesic $\overline{x} \overline{\omega}$ has finite support $\{n : \sigma(n) \neq 0\}$. We write Σ_q for the set of all those sequences. On every horocycle, there is exactly one vertex corresponding to each $\sigma \in \Sigma_q$. Thus, \mathbb{T}_q is in one-to-one correspondence with the set $\Sigma_q \times \mathbb{Z}$, and the k-th horocycle is $H_k = \Sigma_q \times \{k\}$.




Figure 1

Now consider two trees $\mathbb{T}^1 = \mathbb{T}_q$ and $\mathbb{T}^2 = \mathbb{T}_r$ with roots o_1 and o_2 and reference ends ω_1 and ω_2 , respectively.

(2.1) **Definition.** The Diestel-Leader graph DL(q, r) is

$$DL(q,r) = \{x_1x_2 \in \mathbb{T}_q \times \mathbb{T}_r : \mathfrak{h}(x_1) + \mathfrak{h}(x_2) = 0\},$$

and neighbourhood is given by $x_1x_2 \sim y_1y_2 \iff x_1 \sim y_1$ and $x_2 \sim y_2$.

To visualize DL(q,r), draw \mathbb{T}_q in horocyclic layers with ω_1 at the top and $\partial^*\mathbb{T}_q$ at the bottom, and right to it \mathbb{T}_r in the same way, but upside down, with the respective

horocycles $H_k(\mathbb{T}_q)$ and $H_{-k}(\mathbb{T}_r)$ aligned. Connect the two origins o_1 , o_2 by an elastic spring. It is allowed to move along each of the two trees, may expand infinitely, but must always remain in horizontal position. The vertex set of $DL_{q,r}$ consists of all admissible positions of the spring. From a position x_1x_2 with $\mathfrak{h}(x_1) + \mathfrak{h}(x_2) = 0$ the spring may move downwards to one of the r successors of x_2 in \mathbb{T}_r , and at the same time to the predecessor of x_1 in \mathbb{T}_q , or it may move upwards in the analogous way. Such a move corresponds to going to a neighbour of x_1x_2 . We see that DL(q,r) is regular with degree q + r. As the reference point in DL(q,r), we choose $o = o_1o_2$. Figure 2 illustrates DL(2,2).

The relative position of $x = x_1x_2 \in DL(q,r)$ with respect to o is given by the four numbers $\mathfrak{u}(x_1), \mathfrak{d}(x_1), \mathfrak{u}(x_2), \mathfrak{d}(x_2)$, which satisfy the relation

$$\mathfrak{u}(x_1) + \mathfrak{u}(x_2) = \mathfrak{d}(x_1) + \mathfrak{d}(x_2).$$




Figure 2

The lamplighter group $\mathbb{Z}_q \wr \mathbb{Z}$ is defined as follows: Consider the group of all finitely supported configurations

$$C = \{ \eta : \mathbb{Z} \to \mathbb{Z}_q , \quad |\operatorname{supp}(\eta)| < \infty \}$$

with pointwise addition modulo q. Then the group \mathbb{Z} acts on \mathcal{C} by translations $k \mapsto T_k$: $\mathcal{C} \to \mathcal{C}$ with $T_k \eta(m) = \eta(m-k)$. The resulting semidirect product $\mathbb{Z} \times \mathcal{C}$ is

$$\mathbb{Z}_q \wr \mathbb{Z} = \{(\eta, k) : \eta \in \mathcal{C}, k \in \mathbb{Z}\}$$
 with group operation $(\eta, k)(\eta', k') = (\eta + T_k \eta', k + k')$

We identify each $(\eta, k) \in \mathbb{Z}_q \wr \mathbb{Z}$ with the vertex $x_1 x_2 \in DL(q, q)$, where according to the identification $\mathbb{T}_q \leftrightarrow \Sigma_q \times \mathbb{Z}$, the vertices x_i are given by

both written as sequences over the non-positive integers.

This is clearly a one-to-one correspondence between $\mathbb{Z}_q \wr \mathbb{Z}$ and DL(q,q), and it is also straightforward that this group acts transitively and fixed-point-freely on the graph. The action of $m \in \mathbb{Z}$ is given by $x_1x_2 = (\sigma_1, k)(\sigma_2, -k) \mapsto y_1y_2 = (\sigma_1, k+m)(\sigma_2, -k+m)$, and the action of the group of configurations is pointwise addition modulo q. Write δ_k^{ℓ} for the

configuration in \mathcal{C} with value ℓ at k and 0 elsewhere. Then DL(q,q) is the (right) Cayley graph of $\mathbb{Z}_q \wr \mathbb{Z}$ with respect to the symmetric set of generators

$$\{(\delta_1^{\ell}, 1), (\delta_0^{\ell}, -1) : \ell \in \mathbb{Z}_q\},\$$

i.e., an edge corresponds to multiplying with a generator on the right. This is precisely the set of generators considered in [14] and [8] when computing the spectrum of the associated SRW-operator.

3. Tetrahedra and horizontal functions

In the sequel, we shall often write DL for DL(q, r). We say that a function $f: DL \to \mathbb{C}$ is horizontal if it is finitely supported and

$$(3.1) \quad \sum_{y_2 \in \mathbb{T}^2: \mathfrak{h}(y_2) = -\mathfrak{h}(x_1)} f(x_1 y_2) = \sum_{y_1 \in \mathbb{T}^1: \mathfrak{h}(y_1) = -\mathfrak{h}(x_2)} f(y_1 x_2) = 0 \qquad \forall \ x_1 \in \mathbb{T}^1 \,, \ x_2 \in \mathbb{T}^2 \,.$$

(3.2) Lemma. The linear space of horizontal functions is dense in $\ell^2(DL)$.

Proof. It is sufficient to show that every point mass can be approximated in the ℓ^2 -norm by horizontal functions. Furthermore, by vertex-transitivity, it is sufficient to show this for δ_o where $o = o_1 o_2$. Let $b_1 = b_1^n \in \mathbb{T}^1$ be a vertex on H_{-n}^1 (horocycle in \mathbb{T}^1) for which $\mathfrak{u}(b_1) = n + 1$ and $\mathfrak{d}(b_1) = 1$. Define a function $f_1 = f_1^n$ on \mathbb{T}^1 by

$$f_1(x_1) = \begin{cases} 1 & \text{if } x_1 = o_1, \\ -1/q^n & \text{if } b_1 \leq x_1 \in H_0^1, \\ 0 & \text{otherwise.} \end{cases}$$

In the same way, but replacing q with r, we define a function $f_2 = f_2^n$ on \mathbb{T}^2 . Then the function $f = f_n$, given by $f(x_1x_2) = f_1(x_1)f_2(x_2)$, is horizontal, and

$$||f_n - \delta_o||^2 = \frac{1}{q^n r^n} + \frac{1}{q^n} + \frac{1}{r^n} \to 0 \quad \text{as} \quad n \to \infty$$

(3.3) **Definition.** Let $a_1 \in \mathbb{T}^1$ and $a_2 \in \mathbb{T}^2$ be two vertices with $-\mathfrak{h}(a_2) = \mathfrak{h}(a_1) + n$, where $n \geq 0$. Then the (induced) subgraph of DL given by

$$S = S(a_1, a_2) = \{x_1 x_2 \in DL : a_1 \leq x_1, \ a_2 \leq x_2\}$$

is called a tetrahedron in DL with height n(S) = n.




Figure 3

We shall only be interested in computations on tetrahedra with height $n \geq 2$.

If for i = 1, 2 we write $S^i = S_n^i(a_i) = \{x_i \in \mathbb{T}^i : a_i \leq x_i, d(x_i, a_i) \leq n\}$, then $S = \{x_1x_2 \in S_1 \times S_2 : \mathfrak{h}(x_1) + \mathfrak{h}(x_2) = 0\}$. The boundary of S^i in \mathbb{T}^i is $\{a_i\} \cup \partial^*S^i$, where $\partial^*S^1 = \{b_1 \in S^1 : b_1a_2 \in S\}$, respectively $\partial^*S^2 = \{b_2 \in S^2 : a_1b_2 \in S\}$, and the boundary of S in DL is

$$\partial S = (\{a_1\} \times \partial^* S^2) \cup (\partial^* S^1 \times \{a_2\}).$$

Imagining S as a tetrahedron, two of its faces are copies of S^1 that meet at the common bottom side $\partial^* S_1 \times \{a_2\}$, and the other two faces are copies of S^2 that meet at the common top side $\{a_1\} \times \partial^* S_2$. For $k = 0, \ldots, n$, the k-th level of $S = S(a_1, a_2)$ is the set

$$L_k = L_k(a_1, a_2) = \{x_1 x_2 \in S : \mathfrak{h}(x_1) = \mathfrak{h}(a_1) + k\} = \left(S^1 \cap H^1_{\mathfrak{h}(a_1) + k}\right) \times \left(S^2 \cap H^2_{\mathfrak{h}(a_2) + n - k}\right).$$

It has $q^k r^{n-k}$ elements. Furthermore, we write $v_{1,s}$ for the successor of a_1 where the edge $[a_1, v_{1,s}]$ of \mathbb{T}^1 has label $s \in \mathbb{Z}_q$, and analogously $v_{2,t}$ for the successor of a_2 where the edge $[a_2, v_{2,t}]$ of \mathbb{T}^2 has label $t \in \mathbb{Z}_r$.

We identify functions on S with their extensions to DL, where the latter have value 0 on $DL \setminus S$. In particular, every horizontal function on S must be 0 on ∂S . We now choose non-zero functions φ^1 on \mathbb{Z}_q and φ^2 on \mathbb{Z}_r such that

(3.4)
$$\sum_{s=0}^{q-1} \varphi^1(s) = \sum_{t=0}^{r-1} \varphi^2(t) = 0 \quad \text{and} \quad \sum_{s=0}^{q-1} (\varphi^1(s))^2 = \sum_{t=0}^{r-1} (\varphi^2(t))^2 = 1.$$

(Later on, we shall make specific choices for φ^1 and φ^2 .) Using these two functions, we define functions $f_k^1 = f_k^1[\varphi^1]$ on $\mathbb{T}^1(a_1) = \{x_1 \in \mathbb{T}^1 : a_1 \preccurlyeq x_1\}$ and $f_k^2 = f_k^2[\varphi^2]$ on $\mathbb{T}^2(a_2) = \{x_2 \in \mathbb{T}^1 : a_2 \preccurlyeq x_2\}$ by $f_0^1 \equiv 0$, respectively $f_0^2 \equiv 0$, and for $k \geq 1$,

(3.5)
$$f_k^1(x_1) = \begin{cases} \varphi^1(s)q^{(1-k)/2} & \text{if } v_{1,s} \leq x_1 \in H^1_{\mathfrak{h}(a_1)+k}, \\ 0 & \text{otherwise}, \end{cases} \quad \text{and}$$

$$f_k^2(x_2) = \begin{cases} \varphi^2(t)r^{(1-k)/2} & \text{if } v_{2,t} \leq x_2 \in H^2_{\mathfrak{h}(a_2)+k}, \\ 0 & \text{otherwise}. \end{cases}$$

For $k \geq 1$, these functions have ℓ^2 -norm 1. Now we define for $0 \leq k \leq n$

(3.6)
$$f_{k,n}(x) = f_k[S, \varphi^1, \varphi^2](x) = f_k^1(x_1) f_{n-k}^2(x_2), \quad x = x_1 x_2 \in S = S(a_1, a_2).$$

Recall that $n = -\mathfrak{h}(a_2) - \mathfrak{h}(a_1) \geq 2$ is the height of S. The following is a straightforward exercise.

(3.7) **Lemma.** The functions $f_{k,n}$, k = 1, ..., n-1, are horizontal and orthonormal in $\ell^2(S)$. The SRW-operator P satisfies

$$Pf_{k,n} = \frac{\sqrt{qr}}{q+r} (f_{k-1,n} + f_{k+1,n}).$$

Thus, since $f_{k,0} = f_{k,n} = 0$, the action of P on the linear space spanned by $f_{k,n}$, $k = 1, \ldots, n-1$, is described by the $(n-1) \times (n-1)$ tridiagonal matrix

(3.8)
$$M_{n-1} = \frac{\sqrt{qr}}{q+r} \begin{pmatrix} 0 & 1 & & \\ 1 & \ddots & \ddots & \\ & \ddots & \ddots & 1 \\ & & 1 & 0 \end{pmatrix}$$

Its eigenvalues $\lambda_{m,n}$ and associated orthonormal eigenvectors $\psi_{m,n}$ (the latter written as functions on $\{1,\ldots,n-1\}$) are

(3.9)
$$\lambda_{m,n} = \frac{2\sqrt{qr}}{q+r}\cos\frac{m}{n}\pi \quad \text{and}$$

$$\psi_{m,n}(k) = \sqrt{\frac{2}{n}}\sin\frac{km}{n}\pi, \quad m, k = 1, \dots, n-1$$

(3.10) Corollary. The functions $g_{m,n}$ on S, m = 1, ..., n-1, defined by

$$g_{m,n} = \sum_{k=1}^{n-1} \psi_{m,n}(k) f_{k,n}$$

are horizontal and orthonormal in $\ell^2(S)$ as well as in $\ell^2(DL)$. They satisfy

$$Pg_{m,n}=\lambda_{m,n}\cdot g_{m,n}\qquad and$$

$$\mathrm{span}\{g_{m,n}:m=1,\ldots,n-1\}=\mathrm{span}\{f_{k,n}:k=1,\ldots,n-1\}\,.$$

Once more, recall that besides depending on the height n of S, each $g_{m,n}$ depends on a_1, a_2, φ^1 and φ^2 ,

$$g_{m,n} = g_m[S, \varphi^1, \varphi^2].$$

- (3.11) Lemma. Let $S(a_1, a_2)$ and $S(\tilde{a}_1, \tilde{a}_2)$ be two tetrahedra of heights n and $\tilde{n} \geq 2$, respectively. Let φ^1 and φ^2 , respectively $\tilde{\varphi}^1$ and $\tilde{\varphi}^2$ be as in (3.4). Write $f_{k,n} = f_k[a_1, a_2, \varphi^1, \varphi^2]$ and $\tilde{f}_{l,\tilde{n}} = f_l[\tilde{a}_1, \tilde{a}_2, \tilde{\varphi}^1, \tilde{\varphi}^2]$. If one of
 - (i) $(a_1, a_2) \neq (\tilde{a}_1, \tilde{a}_2)$, or
 - (ii) $(a_1, a_2) = (\tilde{a}_1, \tilde{a}_2)$ and $\varphi^1 \perp \tilde{\varphi}^1$ or $\varphi^2 \perp \tilde{\varphi}^2$

holds, then

$$\operatorname{\mathsf{span}}\{f_{k,n}: k=1,\ldots,n-1\} \perp \operatorname{\mathsf{span}}\{\tilde{f}_{l,\tilde{n}}: l=1,\ldots,\tilde{n}-1\}$$
 .

Proof. (i) If $S(a_1, a_2) \cap S(\tilde{a}_1, \tilde{a}_2) = \emptyset$ then perpendicularity is obvious.

If $S(a_1, a_2) \cap S(\tilde{a}_1, \tilde{a}_2) \neq \emptyset$ then both a_1, \tilde{a}_1 and a_2, \tilde{a}_2 must be comparable with respect to the partial order \preceq . Assume that $a_1 \preceq \tilde{a}_1$ and $a_1 \neq \tilde{a}_1$. (The other three cases are treated analogously). Let $\kappa = \mathfrak{h}(\tilde{a}_1) - \mathfrak{h}(a_1)$.

If $k \neq \kappa + l$ then we certainly have $f_{k,n} \perp \tilde{f}_{l,\tilde{n}}$ (since the two functions have disjoint support).

If $k = \kappa + l$ then by construction, the function f_k^1 on $\mathbb{T}^1(a_1)$, as defined in (3.5), is constant on supp \tilde{f}_l^1 , where the latter is given as in (3.5), but on $\mathbb{T}^1(\tilde{a}_1)$. Since $\sum \tilde{f}_l^1 = 0$, we

have $f_k^1 \perp \tilde{f}^1 l$, Therefore, with f_k^2 and \tilde{f}_l^2 defined in the same way on $\mathbb{T}^2(a_2)$, respectively $\mathbb{T}^2(\tilde{a}_2)$.

$$\sum_{x} f_{k,n}(x)\tilde{f}_{l,\tilde{n}}(x) = \sum_{x_2} f_{n-k}^2(x_2)\tilde{f}_{\tilde{n}-l}^2(x_2) \underbrace{\sum_{x_1} f_k^1(x_1)\tilde{f}_l^2(x_1)}_{-0} = 0.$$

(ii) If $(a_1, a_2) = (\tilde{a}_1, \tilde{a}_2)$ then $\tilde{n} = n$, and $f_{k,n} \perp \tilde{f}_{l,n}$ if $l \neq k$, since the functions have disjoint support. Finally,

$$\langle f_{k,n}, \tilde{f}_{k,n} \rangle = \langle \varphi^1, \tilde{\varphi}^1 \rangle \langle \varphi^2, \tilde{\varphi}^2 \rangle = 0$$

We now specify our choices for the functions φ^1 and φ^2 . For $i = 1, \ldots, q-1$ and $s \in \mathbb{Z}_q$, let

(3.12)
$$\varphi_i^1(s) = \begin{cases} \frac{q-i}{\sqrt{(q-i)(q+1-i)}}, & s=i-1, \\ -\frac{1}{\sqrt{(q-i)(q+1-i)}}, & s=i,\dots,q-1, \\ 0, & \text{otherwise.} \end{cases}$$

These functions are orthogonal and satisfy (3.4). Analogously, replacing i with j and q with r, we define the orthogonal functions $\varphi_j^2(t)$, $j = 1, \ldots, r-1$ $(t \in \mathbb{Z}_r)$. We shall write

(3.13)
$$f_k^{[S,i,j]} = f_k[S, \varphi_i^1, \varphi_i^2] \text{ and } g_m^{[S,i,j]} = g_m[S, \varphi_i^1, \varphi_i^2]$$

(3.14) Proposition. The set

$$\mathfrak{B}_{S} = \left\{ g_{m}^{[\tilde{S},i,j]} : \tilde{S} \subseteq S; m \in \{1,\ldots,n(\tilde{S})-1\}; i \in \{1,\ldots,q-1\}; j \in \{1,\ldots,r-1\} \right\}$$

constitutes an orthonormal basis of the linear space of all horizontal functions on the tetrahedron S. Here, \tilde{S} runs through all tetrahedra in DL(q,r) that are contained in S and have height $n(\tilde{S}) \geq 2$.

Proof. Instead of the functions $g_m^{[\tilde{S},i,j]}$, we may equivalently work with the functions $f_m^{[\tilde{S},i,j]}$, since they are also linearly independent and span the same space as \mathfrak{B}_S .

If L_k is the k-th level of S (where S is assumed to have height n) and f is any horizontal function with support in L_k , then f must satisfy each of the following $q^k + r^{n-k}$ equations

$$\sum_{x_2:x_1x_2\in L_k} f(x_1x_2) = 0 \quad \text{and} \quad \sum_{x_1:x_1x_2\in L_k} f(x_1x_2) = 0.$$

Thus, the dimension of the linear space of all horizontal functions with support in L_k is $(q^k - 1)(r^{n-k} - 1)$.

On the other hand, we can count all $f_m^{[\tilde{S},i,j]}$ where $\tilde{S} \subseteq S$ and $\operatorname{supp} f_m^{[\tilde{S},i,j]} \subseteq L_k$. We find (q-1)(r-1) functions of this type (one for each pair (i,j)) associated with every tetrahedron $\tilde{S} = S(\tilde{a}_1, \tilde{a}_2)$, where $a_1 \preceq \tilde{a}_1$ and $d(\tilde{a}_1, a_1) \leq k-1$, and at the same time $a_2 \preceq \tilde{a}_2$ and $d(\tilde{a}_2, a_2) \leq n-k-1$. There are precisely

$$\sum_{\kappa=0}^{k-1} q^{\kappa} \sum_{\nu=0}^{n-k-1} r^{\nu} = \frac{(q^k - 1)(r^{n-k} - 1)}{(q - 1)(r - 1)}$$

choices for $(\tilde{a}_1, \tilde{a}_2)$. Thus, the number of all functions $f_m^{[\tilde{S},i,j]}$ with support in L_k (which are linearly independent) coincides with the dimension of the space of all horizontal functions on L_k , and we have a basis of that space. Putting together the different levels of S, we obtain the proposed result.

Thus, we obtain the following spectral decomposition of $\ell^2(DL)$.

(3.15) **Theorem.** The spectrum of of the SRW-operator P on DL(q,r) is the interval $[-\rho(P), \rho(P)]$, where $\rho(P) = 2\sqrt{qr}/(q+r)$.

It is a pure point spectrum, being the closure of the set

$$\left\{\lambda_{m,n} = \rho(P)\cos\frac{m}{n}\pi : n \ge 2, \ 1 \le m \le n-1\right\}.$$

Furthermore, the set of all functions

$$\mathfrak{B} = \left\{ g_m^{[S,i,j]} : \begin{array}{c} S \text{ tetrahedron in DL}; \ m \in \{1, \dots, n(\tilde{S}) - 1\}; \\ i \in \{1, \dots, q - 1\}; \ j \in \{1, \dots, r - 1\} \end{array} \right\},$$

constructed in Corollary 3.10, respectively (3.13), is a complete orthonormal system in $\ell^2(DL)$ consisting of finitely supported functions; we have

$$Pg_m^{[S,i,j]} = \lambda_{m,n} \cdot g_m^{[S,i,j]},$$

where n = n(S) is the height of S.

Proof. If f is any horizontal function on DL, then there is some tetrahedron S containing its support. By Proposition 3.14, f is a linear combination of elements of \mathfrak{B}_S . Thus, $\mathfrak{B} = \bigcup_S \mathfrak{B}_S$ is an orthonormal system that spans the space of all horizontal functions. Now Lemma 3.2 completes the proof.

4. The spectral measures

Using Theorem 3.15, we can compute the spectral measures (1.1). Indeed, if $x \in DL(q, r)$, then the Fourier expansion of δ_x with respect to the orthonormal system of (3.15) is

$$\delta_x = \sum_{S} \sum_{m=1}^{n(S)-1} \sum_{i=1}^{q-1} \sum_{j=1}^{r-1} g_m^{[S,i,j]}(x) g_m^{[S,i,j]}(\cdot).$$

Therefore, for $x, y \in DL$,

$$p^{(N)}(x,y) = \langle \delta_x, P^N \delta_y \rangle = \sum_{S} \sum_{m=1}^{n(S)-1} \sum_{i=1}^{q-1} \sum_{j=1}^{r-1} \lambda_{m,n(S)}^N g_m^{[S,i,j]}(x) g_m^{[S,i,j]}(y),$$

and comparing this with (1.2), we find that for any continuous function \mathbf{f} on $\operatorname{spec}(P)$, its integral with respect to $\mu_{x,y}$ is

(4.1)
$$\int_{\operatorname{spec}(P)} \mathbf{f}(\lambda) \, \mu_{x,y}(d\lambda) = \sum_{S} \sum_{m=1}^{n(S)-1} \sum_{i=1}^{q-1} \sum_{j=1}^{r-1} \mathbf{f}(\lambda_{m,n(S)}) \, g_m^{[S,i,j]}(x) \, g_m^{[S,i,j]}(y) \,,$$

a countable sum of point masses. Since DL is transitive, we only need the measures $\mu_{o,x}$, where $x \in DL$. Let $x = x_1x_2$ with $\mathfrak{u}_i = \mathfrak{u}(x_i)$ and $\mathfrak{d}_i = \mathfrak{d}(x_i)$. Furthermore, for i = 1, 2 let $c_i = x_i \curlywedge o_i$, so that $S(c_1, c_2)$ has height $\mathfrak{s} = \mathfrak{u}_1 + \mathfrak{u}_2 = \mathfrak{d}_1 + \mathfrak{d}_2 = (d(x_1, o_1) + d(x_2, o_2))/2$, see Figure 4.




Figure 4

Since $\lambda_{m,n}$ depends only on m/n, we choose m and n relatively prime $(1 \leq m < n)$. In order to compute the mass $\mu_{o,x}(\lambda_{m,n})$, we have to consider all tetrahedra $S = S(a_1, a_2)$ with height $n(S) = \ell n$ that contain $S(c_1, c_2)$, that is, $a_i \in \overline{c_i, \omega_i}$ for i = 1, 2. For those S, note that $g_m^{[S,i,j]}(o) = 0$ when $(i,j) \neq (1,1)$. Therefore, denoting by $\lceil \cdot \rceil$ the next larger integer, (4.1) now yields

(4.2)
$$\mu_{o,x}(\lambda_{m,n}) = \sum_{\ell=\lceil \mathfrak{s}/n \rceil}^{\infty} \sum_{\substack{S \supseteq S(c_1, c_2) \\ n(S) = \ell n}} g_{\ell m}^{[S,1,1]}(o) g_{\ell m}^{[S,1,1]}(x),$$

and if we set

$$k_i = k_i(S) = d(a_i, c_i) = \mathfrak{h}(c_i) - \mathfrak{h}(a_i), \quad i = 1, 2,$$

then by Corollary 3.10

(4.3)
$$g_{\ell m}^{[S,1,1]}(x) g_{\ell m}^{[S,1,1]}(o) = C_{k_1,k_2} q^{-k_1 - (\mathfrak{u}_1 + \mathfrak{d}_1)/2} r^{-k_2 - (\mathfrak{u}_2 + \mathfrak{d}_2)/2} \times \frac{2}{\ell n} \sin\left((k_1 + \mathfrak{u}_1)\frac{m}{n}\pi\right) \sin\left((k_1 + \mathfrak{d}_1)\frac{m}{n}\pi\right).$$

where $n(S) = \ell n$ and

(4.4)
$$C_{k_1,k_2} = \begin{cases} (q-1)(r-1), & \text{if } k_1 > 0 \text{ and } k_2 > 0, \\ -(r-1), & \text{if } k_1 = 0 \text{ and } k_2 > 0, \\ -(q-1), & \text{if } k_1 > 0 \text{ and } k_2 = 0, \\ 1, & \text{if } k_1 = k_2 = 0. \end{cases}$$

The last case occurs only when $a_1 = c_1$ and $a_2 = c_2$. Also, note that $k_1 + k_2 = n(S) - \mathfrak{s}$ and that $\mathfrak{u}_i + \mathfrak{d}_i = d(x_i, o_i)$, i = 1, 2. We obtain

(4.5) Proposition. If $1 \le m < n$ and m and n are relatively prime, then with constants as in (4.4),

$$\mu_{o,x}(\lambda_{m,n}) = \sum_{\ell=\lceil \mathfrak{s}/n \rceil}^{\infty} \frac{2r^{-\ell n}}{\ell n} \sum_{k=0}^{\ell n-\mathfrak{s}} C_{k,\ell n-\mathfrak{s}-k} (r/q)^{k+d(x_1,o_1)/2} \sin\left((k+\mathfrak{u}_1)\frac{m}{n}\pi\right) \sin\left((k+\mathfrak{d}_1)\frac{m}{n}\pi\right).$$

Elementary computations yield the following.

(4.6) Corollary. The Plancherel measure $\mu = \mu_{o,o}$ is given by

$$\mu(\lambda_{m,n}) = \begin{cases} \frac{\log(1-r^{-n}) - \log(1-q^{-n})}{n(r-q)} & \frac{2qr(q+r)(q-1)(r-1)\sin^2\frac{m}{n}\pi}{(r-q)^2 + 4qr\sin^2\frac{m}{n}\pi}, & \text{if } r \neq q, \\ \frac{(q-1)^2}{q^n - 1}, & \text{if } r = q, \end{cases}$$

where $\lambda_{m,n} = \frac{2\sqrt{qr}}{q+r}\cos\frac{m}{n}\pi$ and m and n are relatively prime $(1 \le m < n)$.

Note that $\lambda_{n-m,n} = -\lambda_{m,n}$ and $\mu(\lambda_{n-m,n}) = \mu(-\lambda_{m,n})$, that is, the Plancherel measure is symmetric, as it has to be, since DL(q,r) is a bipartite graph (it has no odd cycles). The formula for μ in the case r = q (lamplighter group) was obtained previously in [14] and [8]. For $x \neq o$, the inner sum in Proposition 4.5 can be computed in (lengthy) closed form, but in general not the outer one.

5. Asymptotic behaviour of the return probabilities

Combining (1.2) with Corollary 4.6, we can determine the exact asymptotic behaviour of the return probabilities $p^{(N)}(o, o)$ as $N \to \infty$. For odd N, these probabilities are = 0. For dealing with $p^{(2N)}(o, o)$, the following standard technical lemma will be useful.

(5.1) Lemma. For $k \in \mathbb{N}$ and $\gamma \in \mathbb{R}$ and any sequence ε_n tending to 0, let

$$\Sigma(N) = \Sigma(N; k, \gamma) = \sum_{n=2k+1}^{\infty} (1 + \varepsilon_n) n^{\gamma} q^{-n} \cos^{2N} \frac{k}{n} \pi.$$

Then

$$\Sigma(N) \sim \xi_k^{\gamma} (2\pi/C_k)^{1/2} \exp(-B_k N^{1/3}) N^{(1+2\gamma)/6}$$
 as $N \to \infty$,

where

$$\Phi_k(\xi) = \xi \log q + \frac{(k\pi)^2}{\xi^2}, \quad \xi_k = \left(\frac{2(k\pi)^2}{\log q}\right)^{1/3};$$

$$B_k = \Phi_k(\xi_k) = 3\left(\frac{k\pi \log q}{2}\right)^{2/3}, \quad \Phi'_k(\xi_k) = 0 \quad and \quad C_k = \Phi''_k(\xi_k) = 6\left(\frac{(\log q)^2}{4k\pi}\right)^{2/3}.$$

(Here, as usual, \sim denotes asymptotic equivalence, i.e., quotients tending to 1. It will always be clear from the context whether \sim means neighbourhood in a graph or asymptotic equivalence.)

Proof. We decompose $\Sigma(N) = \Sigma_1(N) + \Sigma_2(N)$, where the sum $\Sigma_1(N)$ ranges over all n with $2k < n < \kappa N^{1/3}$, and $\Sigma_2(N)$ ranges over all $n \ge \kappa N^{1/3}$, and where $\kappa = \xi_k/2$.

We start with $\Sigma_2(N)$. For $n \to \infty$,

$$\cos^{2N} \frac{k}{n} \pi = \exp\left(-(k\pi)^2 N(n^{-2} + O(n^{-4}))\right).$$

Therefore, we have as $N \to \infty$

(5.2)
$$\Sigma_2(N) \sim N^{\gamma/3} \sum_{n \ge \kappa N^{1/3}} \xi_{N,n}^{\gamma} \exp\left(-N^{1/3} \Phi_k(\xi_{N,n})\right)$$
, where $\xi_{N,n} = \frac{n}{N^{1/3}}$.

The point where Φ_k attains its minimum is ξ_k , and we compute the values $\Phi_k(\xi_k) = B_k$ and $\Phi''_k(\xi_k) = C_k$, as given above. Therefore

$$\Phi_k(\xi) = B_k + \frac{C_k}{2}(\xi - \xi_k)^2 + R(\xi)(\xi - \xi_k)^3$$

with $R(\xi)$ continuous for $\xi > 0$. We now "substitute"

$$\tau_{N,n} = N^{1/6}(\xi_{N,n} - \xi_k)$$
, with $\Delta \tau_{N,n} = \tau_N(n+1) - \tau_{N,n} = N^{-1/6} \to 0$.

Then we can rewrite (5.2) as

(5.3)
$$\Sigma_{2}(N) \sim \exp\left(-B_{k} N^{1/3}\right) N^{(1+2\gamma)/6} \times \left\{ \sum_{\substack{n:\\ \tau_{N,n} \geq -\kappa N^{1/6}}} (\xi_{k} + N^{-1/6} \tau_{N,n})^{\gamma} \exp\left\{-\frac{C}{2} \tau_{N,n}^{2} - R_{N}^{*}(\tau_{N,n})\right\} \Delta \tau_{N,n} \right\}$$

where $R_N^*(\tau) = N^{-1/6} R(\xi_0 + N^{-1/6} \tau) \tau^3$.

It is now standard that the sum in (5.3) converges to $\xi_k^{\gamma} \int_{-\infty}^{\infty} e^{-C_k \tau^2/2} d\tau = \xi_k^{\gamma} \sqrt{2\pi/C}$. (One has to use dominated convergence in a suitable central piece of the sum and control the two tails.) Thus, $\Sigma_2(N)$ has the asymptotic behaviour that we have proposed for $\Sigma(N)$.

Let us now look at $\Sigma_1(N)$. Set $M = \sup_n |1 + \varepsilon_n| n^{\gamma} q^{-n}$. Then

$$\Sigma_1(N) \le M \, \kappa \, N^{1/3} \, \cos^{2N} \left(\frac{k\pi}{\kappa N^{1/3}} \right) \sim M \, \kappa \, N^{1/3} \, \exp \left(-\frac{(k\pi)^2}{\kappa^2} N^{1/3} \right) \, .$$

With our choice $\kappa = \xi_k/2$, one checks that $(k\pi)^2/\kappa^2 > B_k$. Therefore $\Sigma_1(N)/\Sigma_2(N) \to 0$ as $N \to \infty$.

For the following, recall from Theorem 3.15 that $\rho(P) = 1$ when r = q. When $r \neq q$, it is enough to consider only r > q.

- (5.4) **Theorem.** Let ξ_1, B_1 and C_1 be as defined in Lemma 5.1.
- (i) If r > q then

$$p^{(2N)}(o,o) \sim A_1 \, \rho(P)^{2N} \, \exp(-B_1 \, N^{1/3}) \, N^{-5/6} \qquad as \ N \to \infty \,,$$

where

$$A_1 = 4\pi^2 \xi_1^{-3} (2\pi/C_1)^{1/2} qr(q+r)(q-1)(r-1)/(r-q)^3.$$

(ii) If r = q then

$$p^{(2N)}(o,o) \sim \bar{A}_1 \exp(-B_1 N^{1/3}) N^{1/6}$$
 as $N \to \infty$,

where

$$\bar{A}_1 = 2(q-1)^2 (2\pi/C_1)^{1/2}$$
.

Proof. We decompose (using $\lambda_{n-m,n} = -\lambda_{m,n}$ and $\lambda_{1,2} = 0$)

$$p^{(2N)}(o,o) = \sum_{n=3}^{\infty} \sum_{\substack{m=1 \\ \gcd(m,n)=1}}^{n-1} \mu(\lambda_{m,n}) \lambda_{m,n}^{2N} = S_1(N) + S_2(N),$$

where

$$S_1(N) = 2 \sum_{n=3}^{\infty} \mu(\lambda_{1,n}) \, \lambda_{1,n}^{2N}$$
 and $S_2(N) = \sum_{n=4}^{\infty} \sum_{\substack{m=2 \ \gcd(m,n)=1}}^{n-2} \mu(\lambda_{m,n}) \, \lambda_{m,n}^{2N}$.

Case r > q. Then, for $n \to \infty$,

$$2 \mu(\lambda_{1,n}) = 2 \pi^2 A_0 n^{-3} q^{-n} (1 + O(n^{-2}))$$
 where $A_0 = 2qr(q+r)(q-1)(r-1)/(r-q)^3$.

Therefore, using Lemma 5.1,

(5.5)
$$S_1(N) = A_0 \rho(P)^{2N} \Sigma(N; 1, -3)$$

has the asymptotic behaviour that we have proposed for $p^{(2N)}(o, o)$. Thus, it remains to show that $S_2(N)$ is dominated by $S_1(N)$. Note that for $2 \le m \le n-2$, we have

$$\lambda_{m,n}^2 \le \rho(P)^2 \cos^2 \frac{2}{n} \pi \,,$$

and

$$\sum_{\substack{m=2\\\gcd(m,n)=1}}^{n-2} \mu(\lambda_{m,n}) \le A_0 \left(\log(1-r^{-n}) - \log(1-q^{-n})\right) = A_0 \left(1+\varepsilon_n\right) q^{-n},$$

where $\varepsilon_n \to 0$ as $n \to \infty$. Therefore,

(5.6)
$$S_2(N) \le A_0 \, \rho(P)^{2N} \, \Sigma(N; 2, 0) \, .$$

Since $B_2 > B_1$, Lemma 5.1 shows that $\Sigma(N; 2, 0)/\Sigma(N; 1, -3) \to 0$, and comparing (5.5) with (5.6), we see that $S_2(N)/S_1(N) \to 0$ as $N \to \infty$.

Case r = q. The proof is basically the same. The only difference is that in this case, Theorem 3.15 yields

$$\mu(\lambda_{1,n}) \sim (q-1)^2 q^{-n}$$

while in case r > q, we had the additional factor n^{-3} . Therefore

$$S_1(N) = 2(q-1)^2 \Sigma(N; 1, 0)$$
 and $S_2(N) \le (q-1)^2 \Sigma(N; 2, 1)$,

whence Lemma 5.1 implies the result.

It is quite surprising that the polynomial terms $N^{-5/6}$ versus $N^{1/6}$ are different when $r \neq q$, respectively r = q. The asymptotics in case r = q were computed previously by Revelle [23] for a very similar random walk on the lamplighter group: in terms of DL(q,r) this is SRW on the graph obtained by adding edges in the first tree (\mathbb{T}_q) , so that each vertex is connected to each of the siblings of its predecessor and then taking the horocyclic product as before. It turns out that for q = r, the N-step return probabilities of that random walk are just $2 p^{(N)}(o, o)$, see [29], §5 for details. The specific computations of [23] are very similar to ours, although [23] does not use the Plancherel measure.

6. Plancherel measure and cumulative spectral measure

Let (X, o) and (X', o') be locally finite, infinite graphs with respective roots o and o'. Let R(X, X') be the largest radius R for which there is a root-preserving isomorphism between the balls $B_X(o, R)$ and $B_{X'}(o', R)$ in the respective graph metrics. A sequence of rooted graphs (X_n, o_n) is said to converge to the graph (X, o), if $R(X, X_n) \to \infty$. In this situation, given vertices $x, y \in X$, we can consider them via the respective isomorphisms as elements of X_n for all $n \ge n(x, y)$, and we have the corresponding spectral measures $\mu_{x,y}^{(n)}$ associated with the SRW operator P_n on X_n , as well as the measure $\mu_{x,y}$ associated with P on X. (In particular, we consider o as the common root of all graphs in the sequence.)

In this setting, it is a well known fact regarding operator convergence that

(6.1)
$$\mu_{x,y}^{(n)} \to \mu_{x,y}$$
 weakly, as $n \to \infty$;

see e.g. Grigorchuk and Żuk [13]. The main interest here is in the diagonal elements $\mu_{x,x}^{(n)} \to \mu_{x,x}$, in particular in the case when X is vertex-transitive, and $\mu_{x,x} = \mu_{o,o} \ \forall \ x \in X$. When the X_n are finite graphs, another type of spectral measure is of interest in the place of $\mu_{o,o}^{(n)}$, namely the cumulative spectral measure

$$\tilde{\mu}_{X_n} = \tilde{\mu}_n = \frac{1}{|X_n|} \sum_{x \in X_n} \mu_{x,x}^{(n)} = \frac{1}{|X_n|} \sum_{\lambda \in \operatorname{spec}(P_n)} \operatorname{mult}(\lambda) \, \delta_\lambda \,,$$

where $\operatorname{\mathsf{mult}}(\lambda)$ is the multiplicity of λ as an eigenvalue of P_n . If X_n is vertex-transitive then $\tilde{\mu}_n = \mu_{o,o}^{(n)}$. In general, the following two questions are of interest.

(6.2) (a) Does the sequence
$$(\tilde{\mu}_n)$$
 converge weakly to some probability measure $\tilde{\mu}$? (b) Is $\tilde{\mu} = \mu_{o,o}$?

In case of a positive answer to question (6.2.a), we call $\tilde{\mu}$ the *cumulative spectral measure* of (X, o) with respect to the sequence (X_n, o) . Recently, the names Von Neumann and Serre have been associated with that measure.

All this applies, in particular, when each X_n is an induced subgraph of X. ("Induced" means that when $x, y \in X_n$ are neighbours in X, then also $x \sim y$ in X_n .) In this situation, it may have advantages to use instead of P_n the restriction (truncation) $P_{|n} = P|_{X_n}$ of P_n on P_n on P_n and P_n coincide in the interior of P_n , while they differ in the points of the boundary ∂X_n of P_n (i.e., the points of P_n having a neighbour in P_n while P_n uses different weights (vertex degrees) at the boundary points. The spectral measures of

 $P_{|n}$ also converge weakly to the respective spectral measures of X, and one can as well study the cumulative spectral measures $\tilde{\mu}_{|n}$ associated with $P_{|n}$ and their possible limit, again denoted $\tilde{\mu}$.

We remark that in the literature, the distinction between Plancherel (Kesten) and cumulative spectral measures has not always been very clear.

Question (6.2) has first been dealt with explicitly by MCKAY [19], who showed that when the X_n are (q+1)-regular graphs with (asymptotically) few cycles, then the sequence $\tilde{\mu}_n$ converges weakly to the Plancherel measure of the tree \mathbb{T}_q . A systematic answer to question (6.2.a) is given by SERRE [26]. In [14], the spectrum of SRW on the lamplighter group $\mathbb{Z}_2 \wr \mathbb{Z}$ (i.e., DL(2,2)) is computed via an approximating sequence of Schreier graphs, and it is shown that the corresponding cumulative spectral measure coincides with $\mu_{o,o}$.

The sequence (X_n) of subsets of X is called a $F\emptyset$ lner sequence, if $\inf_n |\partial X_n|/|X_n| = 0$. It is called expanding, if that infimum is positive. Recall that a graph is called amenable if it has a $F\emptyset$ lner sequence. By Dodziuk [10], this holds if and only if the spectral radius of SRW satisfies $\rho(P) = 1$. This notion comes from group theory; a group is called amenable if it carries a finitely additive, left-invariant probability measure, and a finitely generated group is amenable if and only if one (equivalently, each) of its Cayley graphs with respect to a finite, symmetric generating set is amenable. For more details in the context of random walks, see e.g. [28], §10 and §12.

- (6.3) **Theorem.** Let X be an infinite, connected locally finite vertex-transitive graph, and (X_n) an increasing subsequence of finite subgraphs whose union is X.
- (a) If (X_n) is a Følner sequence then both $\tilde{\mu}_n$ and $\tilde{\mu}_{|n}$ converge weakly to the Plancherel measure of X.
- (b) If (X_n) is an expanding sequence then $\tilde{\mu}_{|n|}$ does not converge to the Plancherel measure of X.

Proof. All involved measures are probability measures on \mathbb{R} with compact support. Therefore, weak convergence holds if and only if for each $N \in \mathbb{N}$, the N-th moment of $\tilde{\mu}_n$ (resp. $\tilde{\mu}_{|n}$) converges to the N-th moment of the Plancherel measure. The latter is $M_N(\mu_{o,o}) = p^{(N)}(o,o) = p^{(N)}(x,x)$ for all $x \in X$ (by transitivity), while

$$M_N(\tilde{\mu}_n) = \frac{1}{|X_n|} \sum_{x \in X_n} p_{X_n}^{(N)}(x, x)$$

(analogously for $\tilde{\mu}_{|n}$). Now consider

$$\partial_N X_n = \{ x \in X_n : d(x, \partial X_n) \le N/2 \}$$

If $x \in X_n \setminus \partial_N X_n$ then $p_{X_n}^{(N)}(x,x) = p_{|X_n}^{(N)}(x,x) = p^{(N)}(x,x)$. Therefore

$$M_N(\mu_{o,o}) - M_N(\tilde{\mu}_{|n}) = \frac{1}{|X_n|} \sum_{x \in X_n} \left(p^{(N)}(x, x) - p_{|X_n}^{(N)}(x, x) \right)$$
$$= \frac{1}{|X_n|} \sum_{x \in \partial_N X_n} \left(p^{(N)}(x, x) - p_{|X_n}^{(N)}(x, x) \right)$$

Now, if (X_n) is a Følner sequence then

$$|M_N(\mu_{o,o}) - M_N(\tilde{\mu}_{|n})| \le 2 |\partial_N X_n|/|X_n| \to 0 \quad (n \to \infty),$$

and the same holds for $\tilde{\mu}_n$. This proves (a).

To see (b), first note that $p^{(N)}(x,x) - p_{|X_n}^{(N)}(x,x) > 0$ for every $x \in \partial X_n$. The involved transition probabilities regard only what happens in the ball B(x, N/2), and the restriction to X_n means that only a part of that ball is admitted for the walker, while the rest is taboo. By transitivity, all these balls are isomorphic, and up to isometry, there are only finitely many ways to subdivide a ball into the admitted and taboo parts. Hence, as x varies, while N is fixed, there are only finitely many different values of $p_{|X_n}^{(N)}(x,x)$. Consequently, there is $\varepsilon_N > 0$ such that

$$p^{(N)}(x,x) - p_{|X_n}^{(N)}(x,x) \ge \varepsilon_N \quad \forall n \in \mathbb{N}, x \in \partial X_n.$$

Therefore

$$M_N(\mu_{o,o}) - M_N(\tilde{\mu}_{|n}) \ge \varepsilon_N |\partial X_n|/|X_n|,$$

which does not tend to zero as $n \to \infty$.

- **(6.4) Remark.** (1) Theorem 6.3 is of course also valid for the adjacency matrix of X and its restriction to X_n , acting on $\ell^2(X)$, resp. $\ell^2(X_n) \subset \ell^2(X)$.
- (2) Part (a) can also be formulated for a sequence of finite graphs X_n converging to X that are not necessarily subgraphs of X. In that case, define

$$\partial^N X_n = \{x \in X_n : B_{X_n}(x, N/2) \text{ is } not \text{ isomorphic with } B_X(o, N/2)\}.$$

Then we call (X_n) a $F \emptyset lner approximation$ of X if $|\partial^N X_n|/|X_n| \to 0$ as $n \to \infty$ for every $N \in \mathbb{N}$.

This condition requires that X is vertex-transitive. If (X_n) is a Følner approximation of X, then the above argument shows that the cumulative spectral measures of X_n converge weakly to the Plancherel measure of X.

(3) As mentioned above, GRIGORCHUK AND \dot{Z} UK [14] consider $\mathbb{Z}_2 \wr \mathbb{Z}$, i.e., DL(2,2), and use an approximation by Schreier graphs and the associated cumulative spectral measures. They show that the latter converge to the Plancherel measure. This can also be interpreted in terms of a Følner approximation. Indeed, the graphs X_n defined by the action of the automaton considered in [14] can be defined alternatively as follows: let m be the smallest power of 2 that is $\geq n$, and consider the group $\Gamma_n = \mathbb{Z}_2^n \rtimes \mathbb{Z}_m$, with \mathbb{Z}_m acting as a matrix with 1's on the diagonal and just above, 0's elsewhere. This is the permutation group acting on level n. The generator a generates \mathbb{Z}_m , and $b = \sigma a$ where σ is any non-trivial element of \mathbb{Z}_2^m . The Schreier graph is the homogeneous space $X_n = \mathbb{Z}_m \backslash \Gamma_n$.

The elements in X_n can be naturally identified with vectors in \mathbb{Z}_2^n . Fix an integer N, and let Y_n denote those vectors that are not periodic of period less than N. On one hand, $|Y_n|/|X_n| \to 1$ as $n \to \infty$, and on the other hand, any ball of radius N around $y \in Y_n$ embeds in the lamplighter group $\mathbb{Z}_2^{\infty} \rtimes \mathbb{Z}$; therefore the graphs X_n have the Følner approximation property.

Our computation of the spectrum of SRW on DL(q,r) in §3 is linked with tetrahedra. In this context, it is natural to take for X_n an increasing family of tetrahedra S_n with height $n \to \infty$, whose union is DL, and consider the associated cumulative measures $\tilde{\mu}_{|n}$.

If r = q, i.e., for the lamplighter group, we have $|S_n| = (n+1)q^n$ and $|\partial S_n| = 2q^n$. Therefore (S_n) is a very natural Følner sequence in DL(q,q), which is indeed a Cayley graph of an amenable group.

On the other hand, if r > q then $|S_n| \sim r^{n+1}/(r-q)$ and $|\partial S_n| \sim r^n$, as $n \to \infty$, whence (S_n) is an expanding sequence. We know that in this case DL(q,r) is a nonamenable graph [25].

(6.5) Corollary. If r = q then for S_n , both sequences $\tilde{\mu}_n$ and $\tilde{\mu}_{|n}$ converge weakly to the Plancherel measure of DL(q,q).

If $r \neq q$ then $\tilde{\mu}_{|n}$ does not converge to the Plancherel measure of DL(q,r).

We remark that we did not compute the actual limit of $\tilde{\mu}_{|n}$, when $r \neq q$. This can be done along the lines of the following computations.

Let us now consider the sequence $\tilde{\mu}_n$ corresponding to SRW on the graphs S_n for $r \neq q$. We shall always suppose that r > q.

Let $S = S(a_1, a_2)$ be any tetrahedron with height n. We want to compute the cumulative spectral measure of SRW P_S on S. We define

$$\partial^1 S = \{b_1 a_2 : b_1 \in \mathbb{T}_q, \ b_1 a_2 \in DL\} \text{ and } \partial^2 S = \{a_1 b_2 : b_2 \in \mathbb{T}_r, \ a_1 b_2 \in DL\}$$

(the lower and upper parts of ∂S). Then P_S coincides with P in the interior $S \setminus \partial S$ of S, while

$$p_S(b_1a_2, x_1x_2) = 1/r$$
 if $b_1a_2 \in \partial^1 S$, $x_1 = b_1^-$, $x_2^- = a_2$, and $p_S(a_1b_2, x_1x_2) = 1/q$ if $a_1b_2 \in \partial^2 S$, $x_1^- = a_1$, $x_2 = b_2^-$.

In order to compare with P acting on $\ell^2(X)$, it is more natural to consider P_S as a self-adjoint operator on $\ell^2(S, \mathbf{m}_S)$, where

$$\mathsf{m}_{S}(x_{1}x_{2}) = \begin{cases} 1, & \text{if } x_{1}x_{2} \in S \setminus \partial S, \\ r/(r+q), & \text{if } x_{1}x_{2} \in \partial^{1}S, \\ q/(r+q), & \text{if } x_{1}x_{2} \in \partial^{2}S, \end{cases}$$

instead of using the reference measure $\deg_S = (q+r)\mathsf{m}_S$.

We already know part of the spectrum of P_S , namely, the spectrum of P_S acting on the space of all horizontal functions with support in S. We need further eigenfunctions besides the horizontal ones. Recall the functions $f_k^{[S,i,j]}$ and $g_m^{[S,i,j]}$ constructed in (3.13), $i=1,\ldots,q-1,\ j=1,\ldots,r-1$, with $k,m=1,\ldots,n-1$. We can also include k=0,n with $f_0^{[S,i,j]}=f_n^{[S,i,j]}=0$. We shall now extend the range of i and j, adding also the values i=0 and j=0. Namely, in analogy with (3.5) and (3.6), and using (3.12), we define for

 $k = 0, \dots, n \text{ and } x = x_1 x_2 \in S$

$$f_k^{[S,0,0]}(x_1x_2) = \begin{cases} q^{-k/2} r^{-(n-k)/2} & \text{if } x_1 \in H^1_{\mathfrak{h}(a_1)+k}, \\ 0 & \text{otherwise}, \end{cases}$$

$$(6.6) \qquad f_k^{[S,0,j]}(x_1x_2) = \begin{cases} q^{-k/2} f_{n-k}^2 [\varphi_j^2](x_2) & \text{if } x_1 \in H^1_{\mathfrak{h}(a_1)+k}, \\ 0 & \text{otherwise} \end{cases} \qquad (j = 1, \dots, r-1),$$

$$f_k^{[S,i,0]}(x_1x_2) = \begin{cases} f_k^1 [\varphi_i^1](x_1) r^{-(n-k)/2} & \text{if } x_1 \in H^1_{\mathfrak{h}(a_1)+k}, \\ 0 & \text{otherwise} \end{cases} \qquad (i = 1, \dots, q-1).$$

The different functions $f_k^{[S,i,j]}$ are all orthogonal, and when $k \in \{1,\ldots,n-1\}$, then they have norm one. On the other hand,

(6.7)
$$||f_0^{[S,0,j]}||^2 = \frac{q}{r+q} \quad \text{and} \quad ||f_0^{[S,i,0]}||^2 = \frac{r}{r+q} \quad \text{in} \quad \ell^2(S, \mathsf{m}_S) \,.$$

As usual, we also think of all functions of (6.6) as being extended to the whole of DL, with value 0 outside of S. It is also convenient to set $f_k^{[S,i,j]} \equiv 0$ when k < 0 or k > n. Let $\rho = \rho(P) = \frac{2\sqrt{qr}}{q+r}$. One has for all pairs (i,j)

$$P_{S}f_{k}^{[S,i,j]} = \frac{\rho}{2} \left(f_{k-1}^{[S,i,j]} + f_{k+1}^{[S,i,j]} \right) \begin{cases} \forall \ k \in \mathbb{Z} \,, & \text{if } i, j \neq 0 \,, \\ \forall \ k \in \mathbb{Z} \setminus \{\pm 1\} \,, & \text{if } i = 0, j \neq 0 \,, \\ \forall \ k \in \mathbb{Z} \setminus \{n \pm 1\} \,, & \text{if } i \neq 0, j = 0 \,, \\ \forall \ k \in \mathbb{Z} \setminus \{\pm 1, n \pm 1\} \,, & \text{if } i \neq 0, j = 0 \,, \end{cases}$$

$$P_{S}f_{1}^{[S,0,j]} = \frac{\sqrt{r}}{\sqrt{q}} f_{0}^{[S,0,j]} + \frac{\rho}{2} f_{2}^{[S,0,j]} \,, \quad \text{and}$$

$$P_{S}f_{n-1}^{[S,i,0]} = \frac{\rho}{2} f_{n-2}^{[S,i,0]} + \frac{\sqrt{q}}{\sqrt{r}} f_{n}^{[S,i,0]} \,.$$

<u>Case 1.</u> $i, j \neq 0$. In this case, (6.8) was stated in Lemma 3.7, and we find the corresponding eigenvalues and eigenfunctions as in §3.

Case 2. i=j=0. None of $f_0^{[S,0,0]}$ and $f_n^{[S,0,0]}$ vanish. The action of P_S on the space spanned by $f_k^{[S,0,0]}$ $(k=0,\ldots,n)$ is described by the $(n+1)\times(n+1)$ -matrix

$$M_{n+1}^{o} = \frac{\rho}{2} \begin{pmatrix} 0 & 1 & & & & \\ \frac{q+r}{q} & \ddots & \ddots & & & \\ & 1 & \ddots & \ddots & & \\ & & \ddots & \ddots & 1 & \\ & & & \ddots & \ddots & \frac{q+r}{r} \\ & & & 1 & 0 \end{pmatrix}$$

over the indices k = 0, ..., n. In general, if we have an eigenvalue λ of M_{n+1}^o and an associated *left* eigenvector ψ , written as a function on $\{0, ..., n\}$, then we obtain a normalized

eigenfunction g of P_S and its norm in $\ell^2(S, \mathsf{m}_S)$ by setting

(6.9)
$$g = C \sum_{k=0}^{n} \psi(k) f_k^{[S,0,0]}, \text{ where } C^2 = \frac{q}{r+q} \psi(0)^2 + \sum_{k=1}^{n-1} \psi(k)^2 + \frac{r}{r+q} \psi(n)^2$$

(using (6.7)), as in §3. Applying this recipe, we start with the following eigenvalues and eigenfunctions of M_{n+1}^o , resp. P_S .

$$(6.10) \lambda_{0,n} = 1, \quad \psi_{0,n}(k) = \sqrt{q/r}^{k}, \qquad g_0^{[S,0,0]}(x) = \left(2qr\frac{r^n - q^n}{r^2 - q^2}\right)^{-1/2}, \text{ and}$$

$$\lambda_{n,n} = -1, \quad \psi_{n,n}(k) = \left(-\sqrt{q/r}\right)^{k}, \quad g_n^{[S,0,0]}(x) = (-1)^{\mathfrak{h}(x_1)} \left(2qr\frac{r^n - q^n}{r^2 - q^2}\right)^{-1/2},$$

where $x \in S$. Next, for m = 1, ..., n - 1, set

$$\beta_{m,n}^o = \arctan\left(\frac{q+r}{q-r}\tan\frac{m}{n}\pi\right).$$

Some computations yield the same eigenvalue as in (3.9) and the following corresponding eigenfunction of P_S :

(6.11)
$$\lambda_{m,n} = \rho \cos \frac{m}{n} \pi, \qquad g_m^{[S,0,0]} = \sum_{k=0}^n \frac{\sqrt{2}}{\sqrt{n}} \sin \left(\frac{km}{n} \pi + \beta_{m,n}^o\right) f_k^{[S,0,0]}.$$

Case 3. i=0, $j\neq 0$. Then $f_0^{[S,0,j]}$ does not vanish, while $f_n^{[S,0,j]}\equiv 0$. The action of P_S on the space spanned by $f_k^{[S,0,j0]}$ $(k=0,\ldots,n-1)$ is described by the $(n\times n)$ -matrix

$$M'_{n} = \frac{\rho}{2} \begin{pmatrix} 0 & 1 & & & \\ \frac{q+r}{q} & \ddots & \ddots & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & 0 \end{pmatrix}$$

over the indices k = 0, ..., n - 1. Computations are slightly more involved in this case; we present the results.

Case 3.A $n \ge \frac{r+q}{r-q}$.

<u>A.1</u> If $n > \frac{r+q}{r-q}$ then there is precisely one $\alpha = \alpha_{0,n} > 0$ that solves the equation

(6.12)
$$e^{2n\alpha} = \frac{q - re^{2\alpha}}{qe^{2\alpha} - r}.$$

(If $n \leq \frac{r+q}{r-q}$ then there is no such solution.) We get

$$\lambda'_{0,n} = \rho \cosh \alpha_{0,n}, \qquad g_0^{[S,0,j]} = C'_{0,n} \sum_{k=0}^{n-1} \sinh((n-k)\alpha_{0,n}) f_k^{[S,0,j]}$$

$$(6.13)$$

$$\lambda'_{n-1,n} = -\rho \cosh \alpha_{0,n}, \qquad g_{n-1}^{[S,0,j]} = C'_{n-1,n} \sum_{k=0}^{n-1} (-1)^k \sinh((n-k)\alpha_{0,n}) f_k^{[S,0,j]}.$$

<u>A.2</u> If $n = \frac{r+q}{r-q}$ then we find, setting $\alpha_{0,n} = 0$ and $\alpha_{n-1,n} = \pi$,

$$\lambda'_{0,n} = \rho = \rho \cos \alpha_{0,n}, \qquad g_0^{[S,0,j]} = C'_{0,n} \sum_{k=0}^{n-1} \left(1 - \frac{k}{n}\right) f_k^{[S,0,j]}$$

$$(6.14)$$

$$\lambda'_{n-1,n} = -\rho = \rho \cos \alpha_{n-1,n}, \qquad g_{n-1}^{[S,0,j]} = C'_{n-1,n} \sum_{k=0}^{n-1} (-1)^k \left(1 - \frac{k}{n}\right) f_k^{[S,0,j]}.$$

In both subcases A.1 and A.2, the equation

(6.15)
$$\cot(n\alpha) = \frac{r-q}{r+q}\cot\alpha, \qquad \alpha \in (0, \pi)$$

has exactly n-2 distinct solutions $\alpha_{m,m}$, $m=1,\ldots,n-2$. For each $\alpha_{m,n}$, we get

(6.16)
$$\lambda'_{m,n} = \rho \cos \alpha_{m,n}, \qquad g_m^{[S,0,j]} = C'_{m,n} \sum_{k=0}^{n-1} \sin((n-k)\alpha_{m,n}) f_k^{[S,0,j]}.$$

In (6.13), (6.14) and (6.16), the normalizing constants $C'_{m,n}$ are computed as C in (6.9). $\underline{\text{Case 3.B}}$ $n < \frac{r+q}{r-q}$.

In this case, the equation (6.15) has exactly n distinct solutions $\alpha_{m,n}$, $m = 0, \ldots, n-1$. Associated with each of those $\alpha_{m,n}$ there is a solution of the form (6.16).

Case 4. $i \neq 0$, j = 0. Then $f_n^{[S,i,0]}$ does not vanish, while $f_0^{[S,i,0]} \equiv 0$. The action of P_S on the space spanned by $f_k^{[S,i,0]}$ (k = 1, ..., n) is described by the $(n \times n)$ -matrix

$$M_n'' = \frac{\rho}{2} \begin{pmatrix} 0 & 1 & & & \\ 1 & \ddots & \ddots & & \\ & \ddots & \ddots & 1 & \\ & & \ddots & \ddots & \frac{q+r}{r} \\ & & & 1 & 0 \end{pmatrix}$$

over the indices k = 1, ..., n. This is analogous to Case 3.B, exchanging q with r and k, m with n - k, m - k. The equation

(6.17)
$$\cot(n\gamma) = -\frac{r-q}{r+q}\cot\gamma, \qquad \gamma \in (0, \pi)$$

has exactly n distinct solutions $\gamma_{m,n}$, $m=1,\ldots,n$. Associated with each of them, we find

(6.18)
$$\lambda''_{m,n} = \rho \cos \gamma_{m,n}, \qquad g_m^{[S,i,0]} = C''_{m,n} \sum_{k=1}^n \sin(k\gamma_{m,n}) f_k^{[S,i,0]},$$

with normalizing constant $C''_{m,n}$ according to (6.9).

We set
$$\mathcal{A}_S^o = \{g_m^{[S,0,0]} : m = 0, \dots, n\},$$

$$\mathcal{A}_S' = \{g_m^{[S,0,j]} : m = 0, \dots, n-1, \ j = 1, \dots, r-1\},$$

$$\mathcal{A}_S'' = \{g_m^{[S,i,0]} : m = 1, \dots, n, \ i = 1, \dots, q-1\}.$$

(6.19) Lemma. The orthogonal complement of the subspace of horizontal functions in $\ell^2(S)$, where $S = S(a_1, a_2)$, is spanned by

$$\mathcal{A}_{S} = \mathcal{A}_{S}^{o} \cup \bigcup \left\{ \mathcal{A}_{S(a_{1},c_{2})}' : S(a_{1},c_{2}) \subseteq S \right\} \cup \bigcup \left\{ \mathcal{A}_{S(c_{1},a_{2})}'' : S(c_{1},a_{2}) \subseteq S \right\},\,$$

where c_1 and c_2 vary. The functions in A_S are all orthonormal with respect to each other.

Proof. Orthonormality follows from the straightforward verification that this is true for the corresponding functions f_m in the place of the g_m . To prove that \mathcal{A}_S spans the orthogonal complement of horizontal functions, we proceed as in the proof of Proposition 3.14 and consider the k-th level L_k of S. The space of functions supported in L_k has dimension $q^k r^{n-k}$, whence the codimension of the space of horizontal functions supported in L^k is $q^k + r^{n-k} - 1$. Direct counting shows that this is precisely the number of functions $f_m^{[S(c_1,c_2),i,j]}$ with $S(c_1,c_2) \subseteq S$, $c_1 = a_1$ or $c_2 = a_2$, and i = 0 or j = 0, that do not vanish on L_k .

Thus, our long computations lead to the following result, which also shows that $\tilde{\mu}_n$ does not converge to the Plancherel measure when $r \neq q$.

(6.20) Proposition. If r > q, then the cumulative spectral measures $\tilde{\mu}_n$ associated with (non-truncated) SRW on S_n converge weakly to the measure $\tilde{\mu} + \tilde{\nu}$, given as follows.

$$\tilde{\mu} = (q-1)(r-1)\sum_{N=2}^{\infty} r^{-N} \sum_{m=1}^{N-1} \delta_{\lambda_{m,N}} + (r-q)(r-1) \sum_{2 \le N \le \frac{r+q}{r-q}} r^{-N-1} \sum_{m=0}^{N-1} \delta_{\lambda'_{m,N}}$$

+
$$(r-q)(r-1)\sum_{N>\frac{r+q}{r-q}}r^{-N-1}\sum_{m=1}^{N-2}\delta_{\lambda'_{m,N}}$$

with $\lambda_{m,N} = \rho(P)\cos(\frac{m}{M}\pi)$ and $\lambda'_{m,N} = \rho(P)\cos\alpha_{m,N}$ given by (6.15).

$$\tilde{\nu} = (r - q)(r - 1) \sum_{N > \frac{r+q}{r-q}} r^{-N-1} \left(\delta_{\lambda'_{0,N}} + \delta_{-\lambda'_{0,N}} \right),$$

with $\lambda'_{0,N} = \rho(P) \cosh \alpha_{0,N}$ given by (6.12).

The support of the measure $\tilde{\mu}$ is the interval $[-\rho(P), \rho(P)]$.

The support $\{\pm \lambda'_{0,N}: N > \frac{r+q}{r-q}\}$ of $\tilde{\nu}$ is contained in $[-1, -\rho(P)] \cup [\rho(P), 1]$. The sequence $(\lambda'_{0,N})$ is strictly increasing with limit 1.

Proof. First of all, $|S_n| = (r^{n+1} - q^{n+1})/(r - q)$.

If we fix n and consider $N \in \{2, ..., n\}$, then $S_n = S(a_1, a_2)$ contains $(r^{n-N+1} - q^{n-N+1})/(r-q)$ different tetrahedra $S(c_1, c_2)$ with height N. With each of those, and each $i \in \{1, ..., q-1\}$ and $j \in \{1, ..., r-1\}$, we associate each of the eigenvalues $\lambda_{m,N}$, m = 1, ..., N-1. Thus, taking into account all those tetrahedra of height N, we count each $\lambda_{m,N}$ precisely $(q-1)(r-1)(r^{n-N+1}-q^{n-N+1})/(r-q)$ times. If we divide by $|S_n|$ and let $n \to \infty$, we get the first of the three parts of $\tilde{\mu}$ (with an implicit use of dominated convergence).

Also, S_n contains r^{n-N} different tetrahedra $S(a_1, c_2)$ with height N, where only c_2 is allowed to vary. Associated with each of them, and with each $j \in \{1, \ldots, r-1\}$, we have

each of the eigenvalues $\lambda'_{m,N}$, $m=1,\ldots,N-1$. Again, taking into account all those tetrahedra, dividing by $|S_n|$, and letting $n\to\infty$, we obtain the second and third parts of $\tilde{\mu}$, plus $\tilde{\nu}$. The subdivision is according to whether $|\lambda'_{m,N}| \leq \rho(P)$ or $> \rho(P)$, respectively.

The contributions to the spectrum of SRW on S_n that come from $\mathcal{A}^o_{S(a_1,a_2)}$ and $\mathcal{A}''_{S(c_1,a_2)}$ (where $S(c_1,a_2) \subseteq S(a_1,a_2)$ and c_1 varies) vanish as $n \to \infty$, because $q^n/r^n \to 0$.

7. Final observations

A. Random walks with drift. Besides SRW on DL(q,r), it may also be instructive to consider the following variant. If the actual position is $x = x_1x_2 \in DL(q,r)$, then we first toss a coin, where "head" comes up with probability $\alpha \in (0, 1)$. If head comes up, then we step at random to one among the q neighbours $y = y_1y_2$ of x with $y_1^- = x_1$ (i.e., downwards in Figure 2). Otherwise, we step at random to one among the r neighbours r0 of r2 with r3 of simple random walk on r4 with r5 of simple random walk on r6 of simple random walk on r6 of simple random walk on r7 of simple random walk on r8 of simple random walk on r9 of

(7.1)
$$p_{\alpha}(x,y) = \begin{cases} \alpha/q & \text{if } y_{1}^{-} = x_{1} \text{ and } y_{2} = x_{2}^{-} \\ (1-\alpha)/r & \text{if } y_{1} = x_{1}^{-} \text{ and } y_{2}^{-} = x_{2} \\ 0 & \text{otherwise.} \end{cases}$$

In order to interpret this in terms of a lamplighter when r=q, it is best to think of the lamps not placed at each vertex of the two-way-infinite path \mathbb{Z} , but at the middle of each edge. Each lamp may have q different intensities or states (\equiv elements of \mathbb{Z}_q), the state "off" corresponding to $0 \in \mathbb{Z}_q$. Only finitely many lamps may be switched on. At each step, the lamplighter tosses his α -coin. If "head" comes up, he moves "down" (from k to k+1) and switches the lamp on the transversed edge to a random state. Otherwise, he moves "up" (to k-1) and again switches the lamp on the transversed edge to a random state.

We remark that for all values q, r, the random walk P_{α} on DL(q, r) may be interpreted as a lamplighter walk in an extended sense. Imagine that on each edge of \mathbb{Z} , there is a green lamp with q possible intensities (including "off") plus a red lamp with r possible intensities (including "off"). The rule is that only finitely many lamps may be switched on, and in addition, if the lamplighter stands at k, then all lamps between k and $-\infty$ have to be in a green state, while all lamps between k and $+\infty$ must be in a red state. The lamplighter tosses his α -coin. If "head" comes up, he moves "down" (from k to k+1) and switches the green lamp on the transversed edge to a random state, while switching off the red lamp on that edge. Otherwise, he moves "up" (to k-1) and switches the red lamp on the transversed edge to a random state, while switching off the green lamp on that edge.

SRW on DL(q,r) is P_{α} with $\alpha = \frac{q}{q+r}$. In order to compare DL(q,r) with DL(q,q), it may be more natural to consider the same α in each case.

For arbitrary α , we have

(7.2)
$$m_{\alpha}(x)p_{\alpha}(x,y) = m_{\alpha}(y)p_{\alpha}(y,x)$$
, where $m_{\alpha}(x) = \left(\frac{\alpha r}{(1-\alpha)q}\right)^{\mathfrak{h}(x_1)}$,

i.e., P_{α} is m_{α} -reversible. Therefore, P_{α} acts as a self-adjoint operator on the weighted space $\ell^{2}(DL, m_{\alpha})$ with inner product

$$\langle f, g \rangle_{\alpha} = \sum_{x} f(x)g(x) m_{\alpha}(x).$$

A quick computation shows that

(7.3)
$$\frac{\sqrt{m_{\alpha}(x)}}{\sqrt{m_{\alpha}(y)}} p_{\alpha}(x,y) = t_{\alpha} p(x,y) \quad \forall \ x,y \in DL, \quad \text{where} \quad t_{\alpha} = \frac{\sqrt{4\alpha(1-\alpha)}}{\rho(P)}.$$

Here, p(x,y) refers to simple random walk, and $\rho(P) = 2\sqrt{qr}/(q+r)$ is the spectral radius of the latter. This can be interpreted in terms of the Hilbert space isomorphism $T_{\alpha}: \ell^2(DL, m_{\alpha}) \to \ell^2(DL)$ (the latter with respect to the counting measure), where $T_{\alpha}f(x) = \sqrt{m_{\alpha}(x)} f(x)$. We find that

$$T_{\alpha} P_{\alpha} f = t_{\alpha} \cdot P T_{\alpha} f \quad \forall f \in \ell^{2}(DL, m_{\alpha}).$$

(7.4) Corollary. The spectrum of the operator P_{α} acting on $\ell^2(DL, m_{\alpha})$ is the the interval $\left[-\sqrt{4\alpha(1-\alpha)}, \sqrt{4\alpha(1-\alpha)}\right]$. It is obtained by dilating the spectrum of P acting on $\ell^2(DL)$ (computed in Theorem 3.15) by the factor t_{α} . The Plancherel measure associated with P_{α} is the image of the Plancherel measure associated with P (computed in Corollary 4.6) under this dilation.

(7.5) Corollary. The return probabilities of P_{α} behave asymptotically as follows.

(i) If r > q then

$$p^{(2N)}(o,o) \sim A_1 \left(4\alpha (1-\alpha) \right)^N \exp\left(-B_1 N^{1/3} \right) N^{-5/6} \quad as \quad N \to \infty.$$

(ii) If r = q then

$$p^{(2N)}(o,o) \sim \bar{A}_1 \left(4\alpha (1-\alpha) \right)^N \exp(-B_1 N^{1/3}) N^{1/6}$$
 as $N \to \infty$.

(The constants A_1 , \bar{A}_1 and B_1 are as in Theorem 5.4).

This direct comparison of the P_{α} underlines how surprising it is that for r > q (red and green lamps with different numbers of states) the asymptotics scale down by a factor of N with respect to the case r = q (only one type of lamps, or equivalently, red and green lamps with the same number of states).

We remark here that BERTACCHI [3] has proved several basic results for general random walks on DL(q,r), without being aware that they apply, in particular, to random walks on $\mathbb{Z}_q \wr \mathbb{Z}$. For example, when applied to P_{α} , one gets a rate of escape theorem and a central limit theorem, as follows.

In Z_n is the random vertex at time n according to P_α , with starting point $Z_0 = o$, then

$$\frac{d(Z_n, Z_0)}{n} \to |2\alpha - 1|$$
 almost surely.

If $\alpha \neq 1/2$ then

$$\frac{d(Z_n, Z_0) - |2\alpha - 1|n}{\sqrt{4\alpha(1 - \alpha)n}} \to \mathcal{N}(0, 1) \quad \text{in law},$$

where $\mathcal{N}(0,1)$ is the standard normal distribution. If $\alpha = 1/2$ then

$$\frac{d(Z_n, Z_0)}{\sqrt{n}} \to \mathcal{M}$$
 in law,

where \mathcal{M} is the probability distribution on \mathbb{R}^+ with density

$$\frac{d\mathcal{M}}{dt} = \frac{1}{\sqrt{2\pi}} \left(e^{-t^2/4} - e^{-t^2} \right) \mathbf{1}_{[0,\infty)}(t) .$$

Note that $2\alpha - 1$ and $4\alpha(1 - \alpha)$ are mean and variance (respectively) of the projected random variable $\widetilde{\Pi}(Z_1)$ on \mathbb{Z} (see definition of $\widetilde{\Pi}$ a few lines below). Also note that these results, contrary to Corollary 7.5, do *not* differ when r = q, resp. $r \neq q$.

B. Projections. We have the natural projections Π_1 , Π_2 and $\widetilde{\Pi}$ of DL(q,r) onto \mathbb{T}_q and \mathbb{T}_r respectively, namely, $\Pi_i(x_1x_2) = x_i$ and $\widetilde{\Pi}(x_1x_2) = \mathfrak{h}(x_1) = \mathfrak{h} \circ \Pi_1(x)$. Associated with them, we have the projected random walks $P_{\alpha,q}$, $P_{1-\alpha,r}$ and \widetilde{P}_{α} , where

$$p_{\alpha,q}(x_1,y_1) = \begin{cases} \alpha/q \,, & \text{if } y_1^- = x_1 \,, \\ 1-\alpha \,, & \text{if } y_1 = x_1^- \,, \\ 0 \,, & \text{otherwise.} \end{cases} \quad \text{and} \quad \widetilde{P}_{\alpha}(k,l) = \begin{cases} \alpha \,, & \text{if } l = k+1 \,, \\ 1-\alpha \,, & \text{if } l = k-1 \,, \\ 0 \,, & \text{otherwise.} \end{cases}$$

 $(P_{1-\alpha,r})$ is analogous to $P_{\alpha,q}$.) The projections are compatible with these transition operators in the sense of factorization of Markov chains, i.e.,

(7.6)
$$p_{\alpha,q}(x_1, y_1) = \sum_{\Pi_1(y) = y_1} p_{\alpha}(x, y) \quad \text{for every } x \text{ with } \Pi_1(x) = x_1, \quad \text{and} \quad \widetilde{p}(k, l) = \sum_{\widetilde{\Pi}_1(y) = l} p_{\alpha}(x, y) \quad \text{for every } x \text{ with } \widetilde{\Pi}(x) = k.$$

Now, $P_{\alpha,q}$ is reversible with respect to $m_{\alpha,q}(x_1) = \left(\frac{\alpha}{q(1-\alpha)}\right)^{\mathfrak{h}(x_1)}$, where $x_1 \in \mathbb{T}_q$. Also, \widetilde{P}_{α} is reversible with respect to $\widetilde{m}_{\alpha}(k) = \left(\frac{\alpha}{(1-\alpha)}\right)^k$. These operators are self-adjoint on the respective spaces $\ell^2(\mathbb{T}_q, m_{\alpha,q})$ and $\ell^2(\mathbb{Z}, \widetilde{m}_{\alpha})$. Their spectra are both known to be the interval $\left[-\sqrt{4\alpha(1-\alpha)}, \sqrt{4\alpha(1-\alpha)}\right]$. The respective Plancherel measures are continuous with respect to Lebesgue measure on that interval; they are

(7.7)
$$\frac{q+1}{2\pi} \frac{\sqrt{4\alpha(1-\alpha)-\lambda^2}}{\tau_{\alpha}^2 - \lambda^2} d\lambda \quad \text{with } \tau_{\alpha} = \frac{\sqrt{4\alpha(1-\alpha)}}{2\sqrt{q}/(q+1)} \quad \text{for } P_{\alpha,q} \text{ on } \mathbb{T}_q, \\
\frac{1}{\pi} \frac{1}{\sqrt{4\alpha(1-\alpha)-\lambda^2}} d\lambda \quad \text{for } \widetilde{P}_{\alpha} \text{ on } \mathbb{Z}.$$

This follows from the well known results for SRW on \mathbb{T}_q and \mathbb{Z} (see e.g. [28], p. 192 and (19.27) on p.214). To adapt the latter methods to the α -walks, one can use the same method as used above for Corollary 7.4, conjugating as in (7.3) with the square root of the reversing measure. The corresponding dilation factors are τ_{α} when passing to $P_{\alpha,q}$ from SRW on \mathbb{T}_q (whose spectral radius is the denominator of τ_{α} above), and $\sqrt{4\alpha(1-\alpha)}$ when passing to \widetilde{P}_{α} from SRW on \mathbb{Z} .

The interesting fact to observe here is that the projections preserve the spectrum, while the respective Plancherel measures differ drastically, starting with a sum of point masses over a dense, countable subset, and ending up with measures having continuous densities with respect to Lebesgue measure.

C. Green kernel estimates. Using the spectral measures, one may also undertake a computation of the asymptotic behaviour of the *Green kernel*

$$G_{\alpha}(o,x) = \sum_{N=0}^{\infty} p^{N}(x,y) = \int_{\operatorname{spec}(P_{\alpha})} \frac{1}{1-\lambda} \, d\mu_{o,x}(\lambda) \,, \quad x,y \in DL(q,r) \,.$$

in space, i.e., as $d(o, x) \to \infty$. Due to the many oscillating terms that occur in these integrals, the spectral method becomes quite tedious, while a more probabilistic reasoning is more efficient and likely to admit extensions to more general random walks, see Brofferio And Woess [4]. The asymptotics depend on the way (direction) in which x tends to ∞ in DL, and they also differ according to whether the drift is non-zero ($\alpha \neq 1/2$) or zero ($\alpha = 1/2$).

References

- [1] Bartholdi, L., and Grigorchuk, R. I.: On the spectrum of Hecke type operators related to some fractal groups, Proc. Steklov Inst. Math. 231 (2000) 1–41.
- [2] Bartholdi, L., Grigorchuk, R. I., and Nekrashevych, V.: From fractal groups to fractal sets, in: Fractals in Graz 2001 (eds. P. Grabner and W. Woess), pp. 25–118, Birkhäuser, Basel, 2003.
- [3] Bertacchi, D.: Random walks on Diestel-Leader graphs, Abh. Math. Sem. Univ. Hamburg **71** (2001) 205–224.
- [4] Brofferio, S., and Woess, W.: Green kernel estimates and the full Martin boundary for random walks on lamplighter groups and Diestel-Leader graphs, TU Graz, in preparation.
- [5] Cartier, P.: Harmonic analysis on trees, Proc. Sympos. Pure Math. 26 (1973) 419–424.
- [6] Cartwright, D. I.: Spherical harmonic analysis on buildings of type \tilde{A}_n , Monatsh. Math. 133 (2001) 93–109.
- [7] Cartwright, D. I., and Młotkowski, W.: Harmonic analysis for groups acting on triangle buildings, J. Austral. Math. Soc. Ser. A **56** (1994) 345–383.
- [8] Dicks, W., and Schick, Th.: The spectral measure of certain elements of the complex group ring of a wreath product, Geom. Dedicata **93** (2002) 121–137.
- [9] Diestel, R., and Leader, I.: A conjecture concerning a limit of non-Cayley graphs, J. Algebraic Combin. 14 (2001) 17–25.
- [10] Dodziuk, J.: Difference equations, isoperimetric inequality, and transience of certain random walks, Trans. Amer. Math. Soc. 284 (1984) 787–794.
- [11] Faraut, J., and Picardello, M. A.: The Plancherel measure for symmetric graphs, Ann. Mat. Pura Appl. 138 (1984) 151–155.
- [12] de la Harpe, P., Robertson, A. G., and Valette, A.: On the spectrum of the sum of generators for a finitely generated group, Israel J. Math. 81 (1993) 65–96.
- [13] Grigorchuk, R. I., and Żuk, A.: On the asymptotic spectrum of random walks on infinite families of graphs, Symposia Math. 39 (1999), 188–204.
- [14] Grigorchuk, R. I., and Żuk, A.: The lamplighter group as a group generated by a 2-state automaton, and its spectrum, Geom. Dedicata 87 (2001) 209–244.
- [15] Kesten, H.: Symmetric random walks on groups, Trans. Amer. Math. Soc. 92 (1959) 336–354.
- [16] Krön, B.: Green functions on self-similar graphs and bounds for the spectrum of the Laplacian, Ann. Inst. Fourier (Grenoble) **52** (2002), 1875–1900.
- [17] Kuhn, G., and Soardi, P. M.: The Plancherel measure for polygonal graphs, Ann. Mat. Pura Appl. 134 (1983) 393–401.

- [18] Malozemov, L., and Teplyaev, A.: Pure point spectrum of the Laplacians on fractal graphs, J. Funct. Anal. 129 (1995) 390–405.
- [19] McKay, B.: The expected eigenvalue distribution of a large regular graph, Linear Algebra and its Appl. 40 (1981) 203–216.
- [20] Mohar, B., and Woess, W.: A survey on spectra of infinite graphs, Bull. London Math. Soc. 21 (1989) 209–234.
- [21] Pittet, C., and Saloff-Coste, L.: On random walks on wreath products, Ann. Probab. **30** (2002) 948–977.
- [22] Pólya, G.: Über eine Aufgabe der Wahrscheinlichkeitsrechnung betreffend die Irrfahrt im Strasennetz, Mat. Ann. 84 (1921) 149–160.
- [23] Revelle, D.: Heat kernel asymptotics on the lamplighter group, preprint, Cornell Univ. (2002).
- [24] Sabot, Ch.: Pure point spectrum for the Laplacian on unbounded nested fractals, J. Funct. Anal. 173 (2000) 497–524.
- [25] Saloff-Coste, L., and Woess, W.: Transition operators, groups, norms, and spectral radii, Pacific J. Math. 180 (1997) 333–367.
- [26] Serre, J.-P.: Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p , Jour. Amer. Math. Soc. 10 (1997), 75–102.
- [27] Teplyaev, A.: Spectral analysis on infinite Sierpiński gaskets, J. Funct. Anal. 159 (1998) 537–567.
- [28] Woess, W.: Random Walks on Infinite Graphs and Groups, Cambridge Tracts in Mathematics 138, Cambridge University Press, Cambridge, 2000.
- [29] Woess, W.: Lamplighters, Diestel-Leader graphs, random walks, and harmonic functions, Combinatorics, Probability & Computing, to appear.

IGAT, BÂTIMENT BCH, ÉCOLE POLYTECHNIQUE FÉDÉRALE, CH-1015 LAUSANNE, SWITZERLAND

 $E ext{-}mail\ address: laurent.bartholdi@epfl.ch}$

INSTITUT FÜR MATHEMATIK C, TECHNISCHE UNIVERSITÄT GRAZ, STEYRERGASSE 30, A-8010 GRAZ, AUSTRIA

E-mail address: woess@TUGraz.at