

SOME EXTREMELY AMENABLE GROUPS RELATED TO OPERATOR ALGEBRAS AND ERGODIC THEORY

THIERRY GIORDANO AND VLADIMIR PESTOV

ABSTRACT. A topological group G is called extremely amenable if every continuous action of G on a compact space has a fixed point. This concept is linked with geometry of high dimensions (concentration of measure). We show that a von Neumann algebra is approximately finite-dimensional if and only if its unitary group with the strong topology is the product of an extremely amenable group with a compact group, which strengthens a result by de la Harpe. As a consequence, we describe nuclear C^* -algebras A as those whose unitary group $U(A)$ with the relative weak topology is strongly amenable in the sense of Glasner. We prove that the group of automorphisms of a Lebesgue space with a non-atomic measure is extremely amenable with the weak topology. We establish a similar result for groups of non-singular transformations, using concentration of measure for groups of automorphisms of measure spaces with finitely many points. As a consequence, we prove extreme amenability of the groups of isometries of $L^p(0, 1)$, $1 \leq p < \infty$, extending the classical result of Gromov and Milman ($p = 2$). We show that a measure class preserving equivalence relation $\mathcal{R}$ on a standard Borel space is hyperfinite if and only if the full group $[\mathcal{R}]$, equipped with the uniform topology, is extremely amenable. Finally, we give natural examples of concentration to a nontrivial space in the sense of Gromov occurring in the automorphism groups of injective factors of type *III*. Some of our results have been announced in C.R. Acad. Sci. Paris Sér. Math. **334** (2002), 273–278.

1. INTRODUCTION

Extreme amenability [19] (or the fixed point on compacta property [38]) is a relatively recent concept, useful for study of ‘large’ (non locally compact) topological groups. Recall that a topological group G is *amenable* [27] if every continuous action of G by affine transformations on a convex compact subset of a topological vector space has a fixed point. A topological group G is *extremely amenable* if every continuous action of G on a compact space has a fixed point.

This is a strong nonlinear fixed point property, never observed in locally compact groups [20, 51]. Beginning with the discovery by Gromov and Milman [23] that the unitary group of an infinite dimensional Hilbert space with the strong operator topology is extremely amenable, a large number of concrete large groups of importance are now known to have this property.

2000 Mathematical Subject Classification. 22A05, 22F50, 37A15, 43A07, 46L05, 46L10.

The concept is interesting for several reasons. Extreme amenability is closely linked to asymptotic geometric analysis, and in many examples it is a manifestation of the phenomenon of concentration of measure on high-dimensional structures, captured in the concept of a Lévy group [23, 15, 13]. Extreme amenability of groups of automorphisms of various combinatorial structures is intimately related to Ramsey-type theorems for those structures [40, 32]. While surely not all infinite dimensional groups are extremely amenable, the fixed point on compacta property of subgroups can be used to make conclusions about the dynamical properties of larger groups, for instance, to describe their universal minimal compact flows [40, 16]. And the recent work by Glasner, Tsirelson and Weiss [18] links Lévy groups to the existence of spatial models for near actions.

No non-trivial locally compact group has the fixed point on compacta property, which is thus exclusively a property of ‘large’ topological groups. This was shown by Granirer and Lau [20]. Later Veech [51] has proved his well-known theorem stating that every locally compact group acts freely on a compact space. (Now there are at least four known proofs of this result, see [32] and references therein.)

The first example of a non-trivial extremely amenable group has been constructed by Herer and Christensen [29]. Gromov and Milman [23] have shown that the unitary group of an infinite dimensional Hilbert space, equipped with the strong operator topology, is extremely amenable. The idea of the proof, using concentration of measure on high-dimensional structures, has turned out to be applicable in a great many situations. For example, Glasner, and also (independently, unpublished) Furstenberg and Weiss, have shown that the group $L_0(\mathbb{I}, U(1))$ of measurable maps from the standard Lebesgue space to the circle rotation group $U(1)$, equipped with the topology of convergence in measure, is extremely amenable, see [15]. The group $U(1)$ here can be replaced by any amenable locally compact group [42].

A direct link between extreme amenability and Ramsey theory has been established by one of the present authors, who has proved in [40] extreme amenability of the group $\text{Aut}(\mathbb{Q}, \leq)$ of order-preserving bijections of the rationals, equipped with the topology of pointwise convergence on $\mathbb{Q}$ viewed as discrete. This statement is a reformulation of the classical Finite Ramsey Theorem. The recent paper by Kechris, Pestov and Todorcevic [32] explores this trend extensively, by establishing extreme amenability of groups of automorphisms of numerous countable Fraïssé structures. It should be noted that extreme amenability of a topological group G can be restated in terms of the so-called Ramsey-Dvoretzky-Milman property (Section 9.3 in [21]) of transitive isometric actions of G on metric spaces, which is in turn linked to both discrete Ramsey theory and Ramsey-type properties in geometric functional analysis [43].

The universal minimal flow $M(G)$ of a topological group G is, in the case where G is locally compact and non-compact, a highly non-constructive object (for instance, non-metrizable, [32], Appendix 2). Of course, if G is an extremely amenable group, then $M(G) = \{*\}$ is a singleton. It appears that the first instance where the flow $M(G)$, different from a point, was described explicitly, was the case where $G = \text{Homeo}_+(\mathbb{S}^1)$

is the group of orientation-preserving homeomorphisms of the circle with the compact-open topology. In this situation, $M(G)$ is the circle $\mathbb{S}^1$ itself, equipped with the canonical action of G [40]. The proof was using the existence of a large extremely amenable subgroup, namely $\text{Homeo}_+(\mathbb{I})$. Glasner and Weiss have subsequently described in [16] the universal minimal flow of the infinite symmetric group with its (unique) Polish topology as the space of linear orders on the natural numbers, and then in [17] the universal minimal flow of the group of homeomorphisms of the Cantor set, C , as the space of maximal chains of closed subsets of C (a construction proposed by Uspenskij [50]). Numerous other examples of explicit computations of universal minimal flows of groups of automorphisms of countable structures can be found in the above cited paper [32].

Overall, the papers [43] and [32] will together provide an introduction to, a survey of, and bibliographical references to most of what is known to date about extremely amenable groups.

In this article we will establish extreme amenability of some concrete topological groups of importance using the concentration of measure, as well as connect the property with known concepts from operator algebras and ergodic theory. We investigate extreme amenability of topological groups of two types: unitary groups of von Neumann algebras and group of transformations of spaces with measure.

It was proved by de la Harpe [28] that a von Neumann algebra M is approximately finite dimensional if and only if its unitary group $U(M)$, equipped with the relative $s(M, M_*)$ -topology, is amenable. We strengthen this result by showing that this is the case if and only if $U(M)$ is the product of a compact group and an extremely amenable group. Paterson [39] has deduced from de la Harpe's result the following: a C^* -algebra A is nuclear if and only if the unitary group $U(A)$, equipped with the relative weak (that is, $s(A, A^*)$) topology, is amenable. We describe nuclear C^* -algebras as those C^* -algebras A whose unitary group $U(A)$ with the relative weak topology is strongly amenable in the sense of Glasner [14].

Let now (X, ν) be a standard non-atomic Borel probability space. We show that the group $\text{Aut}(X, \nu)$ of all measure-preserving automorphisms of (X, ν) , equipped with the weak topology, is extremely amenable. This result is a consequence of the Rokhlin–Kakutani lemma and the concentration of measure on finite symmetric groups discovered by Maurey [35]. The group $\text{Aut}(X, \nu)$, equipped with the uniform topology, is no longer extremely amenable. We also establish the extreme amenability of the group $\text{Aut}^*(X, \nu)$ of measure class preserving transformations of (X, ν) , equipped with the weak topology. As a consequence of this result and a description of groups of isometries of spaces $L^p(X, \mu)$ belonging to Banach [1] and Lamperti [33], we prove that those groups, equipped with the strong operator topology, are extremely amenable for all $1 \leq p < \infty$, extending Gromov and Milman's classical result [23] (obtained for $p = 2$).

We consider measure class preserving equivalence relations $\mathcal{R}$ on the standard Lebesgue measure space and prove that such a relation is amenable in the sense of Zimmer [52],

[53] if and only if the full group $[\mathcal{R}]$, equipped with the uniform topology, is extremely amenable. In order to obtain this and the previous result, we generalize Maurey's theorem to automorphism groups of measure spaces with finitely many points.

The concentration of measure phenomenon can be interpreted in terms of convergence of a family of spaces with metric and measure (*mm*-spaces) to a one-point space with respect to a suitable metric, as was shown by Gromov [22]. This leads to a more general concept of concentration to a non-trivial space. In the concluding part of the article, we show some natural examples of concentration of compact subgroups and other subobjects of the groups of automorphisms of injective factors of type *III* to non-trivial spaces.

Some of the above results have been announced by the present authors in the C.r. note [13].

Acknowledgements. The investigation has been partially supported by NSERC operating grants and by the Swiss Fund for Scientific Research (both authors), as well as the Marsden Fund of the Royal Society of New Zealand and a University of Ottawa start-up grant (V.P. only). Both authors are grateful to Alexander Kechris for useful comments, and to Pierre de la Harpe for his hospitality at the University of Geneva where the present research collaboration started in April 1999.

2. CONCENTRATION AND EXTREME AMENABILITY

In this section we will outline a general scheme of deducing fixed point theorems from the phenomenon of concentration of measure, introduced by Gromov and Milman [23]. We work in a slightly more general context and prove some new results, in particular a concentration of measure result for automorphism groups of finite measure spaces.

2.1. Concentration of measure and Lévy families. For a subset $A \subseteq X$ of a separated uniform space $X = (X, \mathcal{U})$ let $V[A] = \{x \in X : \exists a \in A, (x, a) \in V\}$ be the V -neighbourhood of A . We say that a net (μ_α) of probability measures on X has the *Lévy concentration property*, or simply *concentrates*, if for every family of Borel subsets $A_\alpha \subseteq X$ satisfying $\liminf_\alpha \mu_\alpha(A_\alpha) > 0$ and every entourage $V \in \mathcal{U}_X$ one has $\mu_\alpha(V[A_\alpha]) \rightarrow_\alpha 1$.

A triple (X, d, μ) , where d is a metric on a set X and μ is a probability Borel measure on the metric space (X, d) , is called a *metric space with measure*, or else an *mm-space* [22]. An infinite family (X_n, d_n, μ_n) of *mm*-spaces is a *Lévy family* if, whenever $A_n \subseteq X_n$ are Borel subsets satisfying $\liminf_n \mu_n(A_n) > 0$, one has for each $\varepsilon > 0$ $\lim_{n \rightarrow \infty} \mu_n((A_n)_\varepsilon) = 1$, where A_ε denotes the ε -neighbourhood of a set A .

The above property is equivalent to concentration of the family (μ_n) considered as probability measures supported on the disjoint union $\bigoplus_{n=1}^\infty X_n$, equipped with a metric d inducing the metrics d_n on each X_n and making X_n into an open and closed subset.

The next Lemma first appears in [23], 2.2. For a proof, see Proposition 3.8 in [34]. (The case of uniform spaces clearly reduces to that of *mm*-spaces.)

Lemma 2.1. *Let a net of probability measures (μ_α) (resp. (ν_α)) on an uniform space X (resp. Y) concentrate. Then the net of product measures $(\mu_\alpha \otimes \nu_\alpha)$ concentrates on the product space $X \times Y$. $\square$*

The concentration function, α_X , of an mm-space [23] is defined for $\varepsilon \geq 0$ by

$$\alpha_X(\varepsilon) = \begin{cases} \frac{1}{2}, & \text{if } \varepsilon = 0, \\ 1 - \inf\{\mu(A_\varepsilon) : A \subseteq X \text{ is Borel}, \mu(A) \geq \frac{1}{2}\}, & \text{if } \varepsilon > 0. \end{cases}$$

The following is simple and well-known.

Lemma 2.2. *Let (Y, ν) be a probability measure space, and let d, ρ be two Lipschitz equivalent measurable metrics on Y . Denote by α_d the concentration function of the mm-space (Y, d, ν) , and similarly by α_ρ the concentration function of the mm-space (Y, ρ, ν) . Denote by L the Lipschitz constant of the identity map*

$$(Y, d, \nu) \rightarrow (Y, \rho, \nu)$$

Then for every $\varepsilon > 0$

$$\alpha_\rho(\varepsilon) \leq \alpha_d(L^{-1}\varepsilon).$$

Proof. Indeed, the $L^{-1}\varepsilon$ -neighbourhood (w.r.t. d) of an arbitrary measurable subset $A \subseteq Y$ having measure $\geq \frac{1}{2}$ is contained in the ε -neighbourhood (w.r.t. ρ) of A , and therefore the measure of the latter is at least as large as the measure of the former. $\square$

A family (X_n, d_n, μ_n) is a Lévy family if and only if the concentration functions α_n of X_n converge to zero pointwise as $n \rightarrow \infty$ on the interval $(0, +\infty)$.

A well-known manifestation of concentration of measure (see [36, 37, 34]) is that on a highly-concentrated space every uniformly continuous function is “nearly constant nearly everywhere.” More precisely:

Lemma 2.3. *Let a family (μ_β) of probability measures on a uniform space X be Lévy. Then for every bounded uniformly continuous function f on X there exists a net of constants (c_β) such that for every $\varepsilon > 0$,*

$$(2.1) \quad \mu_\beta\{x \in X : |f_\beta(x) - c_\beta| > \varepsilon\} \rightarrow_\beta 0.$$

The convergence is uniform in f , with the left hand side bounded above by $2\alpha_{X_\beta}(\varepsilon)$.

Proof, sketch. One can choose as c_α median values for f_α , that is, numbers with the property that sets $M_\alpha^-(f) = \{x \in X : f(x) \leq c_\alpha\}$ and $M_\alpha^+(f) = \{x \in X : f(x) \geq c_\alpha\}$ have both μ_α -measure at least $1/2$. Now the upper bound is obtained by applying to those two sets the definition of a Lévy family of measures. $\square$

Here are a few examples of Lévy families.

Example 2.4. The family of special unitary groups $SU(n)$, $n \in \mathbb{N}$, equipped with the normalized Haar measure and the Hilbert-Schmidt metric is a Lévy family [23].

Example 2.5. Let $X = (X, \mu)$ be a probability measure space. Introduce on X^n the product measure $\mu^{\otimes n}$ and the Hamming distance $d(x, y) = |\{i: x_i \neq y_i\}|$. The family X^n , $n \in \mathbb{N}$, is then a Lévy family, with the concentration function α_n of X_n satisfying

$$(2.2) \quad \alpha_n(\varepsilon) \leq 2 \exp(-\varepsilon^2 N)$$

(See e.g. [48], Prop. 2.1.1.)

Here is an immediate consequence. Let $X_1, X_2, \dots$ be a sequence of probability measure spaces. Equip the n -fold product $Y_n = X_1 \times \dots \times X_n$ with the product measure and the Hamming distance. Then the family $(Y_n)_{n=1}^\infty$ is Lévy, with the same concentration functions as in Eq. (2.2). (Apply Example 2.5 and Lemma 2.2 to the Lévy family $\mathbb{I}^n$ with Lebesgue measures and the Hamming distance.)

Example 2.6. Symmetric groups $\mathfrak{S}_n$ of rank n , equipped with the normalized Hamming distance

$$d_{ham}(\sigma, \tau) := \frac{1}{n} |\{i: \sigma(i) \neq \tau(i)\}|$$

and the normalized counting measure

$$\mu(A) := \frac{|A|}{n!}$$

form a normal Lévy family, with the concentration functions satisfying the estimate

$$(2.3) \quad \alpha_{\mathfrak{S}_n}(\varepsilon) \leq \exp(-\varepsilon^2 n / 64).$$

(Maurey [35]; see also [37], Theorem 7.5, and [34], Corollary 4.3.)

There are many known techniques for establishing concentration inequalities. The following approach, based on martingales, allows one to prove Examples 2.5 and 2.6, as well as to obtain some useful variations and generalizations.

Following Milman and Schechtman [37], say that a finite metric space (Ω, d) is of *length at most ℓ* if there exist positive numbers $a_1, a_2, \dots, a_n$ with $\ell = (\sum_{i=1}^n a_i^2)^{1/2}$ and a refining sequence $(\Omega^k)_{k=0}^n$, $\Omega^k = \{A_i^k\}_{i=1}^{m_k}$ of partitions of Ω such that $\Omega^0 = \{\Omega\}$, $\Omega^n = \{\{x\}: x \in \Omega\}$, and for every $k = 1, \dots, n$ and every two elements A, B of the partition Ω^k that are both contained in some element of Ω^{k-1} there exists a bijection $\varphi: A \rightarrow B$ with $d(x, \varphi(x)) \leq a_k$ for all $x \in A$.

The following is Theorem 7.8 in [37], and the present improved constants are taken from Theorem 4.2 in [34].

Theorem 2.7. *Let (Ω, d) be a finite metric space of length at most ℓ , equipped with the normalized counting measure. Then*

$$\alpha_\Omega(\varepsilon) \leq \exp\left(-\frac{\varepsilon^2}{8\ell^2}\right).$$

□

The following is Theorem 7.12 in [37], and (with the present, better constants) Theorem 4.4 in [34].

Theorem 2.8. *Let G be a compact group equipped with a bi-invariant metric d , and let $\{e\} = H_0 < H_1 < \dots < H_n = G$ be a sequence of closed subgroups. Equip every factor-space H_{i+1}/H_i with the factor-distance of d , and let d_i denote the diameter of H_{i+1}/H_i . Then the concentration function of the mm-space (G, d, μ) , where μ is the normalized Haar measure, satisfies*

$$\alpha_G(\varepsilon) \leq \exp \left(-\frac{\varepsilon^2}{8 \sum_{i=0}^{n-1} d_i^2} \right).$$

□

2.2. Two new examples of Lévy families. Here we apply Theorems 2.7 and 2.8 to deduce two new natural examples of Lévy families, which we will use in the sequel.

If (X, μ) be a probability measure space and (G, d) is a metric group of finite diameter, we denote by $L^1(X, \mu; G)$ the group of all measurable maps from (X, μ) to G , equipped with the $L^1(\mu)$ -metric $d^\dagger$ by

$$d^\dagger(f, g) = \int_X d(f(x), g(x)) \, d\mu(x).$$

Corollary 2.9. *Let (X, μ) be a probability measure space with finitely many points, and let (K, d) be a compact metric group of diameter 1. The concentration function of the compact metric group $L^1(X, \mu; K)$, equipped with the normalized Haar measure, satisfies*

$$\alpha(\varepsilon) \leq \exp \left(-\frac{\varepsilon^2}{8 \sum_{x \in X} \mu(\{x\})^2} \right).$$

Proof. If $X = \{x_1, x_2, \dots, x_n\}$, then let H_i consist of all functions vanishing on $x_{i+1}, x_{i+2}, \dots, x_n$. The group H_{i+1}/H_i is isometrically isomorphic to the group $(K, \mu(\{x_{i+1}\})d)$, whose diameter equals $\mu(\{x_{i+1}\})$, and the result follows by Theorem 2.8. □

Corollary 2.10. *Let (X_n, μ_n) be a sequence of probability spaces with finitely many points in each, having the property that the mass of the largest atom in X_n goes to zero as $n \rightarrow \infty$. Let (K, d) be a compact metric group of diameter 1. Then the family of compact groups $L^1(X_n, \mu_n; K)$, equipped with the normalized Haar measure, is Lévy, with*

$$\alpha_{X_n}(\varepsilon) \leq \exp \left(-\frac{\varepsilon^2}{8 \max_{x \in X_n} \mu_n(\{x\})} \right).$$

Proof. Let $v_n = (\mu_n(\{x\}))_{x \in X_n}$. By Hölder's inequality,

$$\sum_{x \in X_n} \mu_n(\{x\})^2 = \langle v_n, v_n \rangle \leq \|v_n\|_1 \cdot \|v_n\|_\infty = \max_{x \in X_n} \mu_n(\{x\}) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Corollary 2.9 now applies. $\square$

Let $X = (X, \mu)$ be a measure space. Denote by $\text{Aut}^*(X, \mu)$ the group of all invertible measurable and non-singular transformations of X . In particular, if X has finitely many points and the measure has full support, then $\text{Aut}^*(X)$ is, as an abstract group, just the symmetric group of rank $n = |X|$.

Define a left-invariant metric on $\text{Aut}^*(X, \mu)$ (the uniform metric) as follows:

$$(2.4) \quad d_{\text{unif}}(\tau, \sigma) = \mu\{x \in X : \tau(x) \neq \sigma(x)\}.$$

For instance, if a finite set X is equipped with a uniform measure, the uniform metric on $\text{Aut}^*(X) \cong \mathfrak{S}_n$ coincides with the Hamming distance. For every measure space, the uniform metric generates a group topology, known as the *uniform topology*, which will play an important role later (cf. Subsection 4.1).

Theorem 2.11. *Let $X = (X, \mu)$ be a probability space with finitely many points. The concentration function α of the automorphism group $\text{Aut}^*(X, \mu)$, equipped with the uniform metric (2.4) and the normalized counting measure, satisfies*

$$(2.5) \quad \alpha(\epsilon) \leq \exp\left(-\frac{\epsilon^2}{32 \sum_{x \in X} \mu(\{x\})^2}\right).$$

Proof. Let $X = \{x_1, x_2, \dots, x_n\}$, where

$$\mu(\{x_1\}) \geq \mu(\{x_2\}) \geq \dots \geq \mu(\{x_{n-1}\}) \geq \mu(\{x_n\}).$$

For $k = 0, 1, \dots, n$, let H_k be the subgroup stabilizing each element $x_1, \dots, x_k$. Thus,

$$\text{Aut}^*(X, \mu) = H_0 > H_1 > H_2 > \dots > H_n = \{e\}.$$

Let Ω^k be the partition of $G = \text{Aut}^*(X, \mu)$ into left H_k -cosets σH_k , $\sigma \in G$.

Suppose $A = \sigma H_k$ and $B = \tau H_k$ are contained in the same left H_{k-1} -coset. Then $\sigma(x_i) = \tau(x_i) = a_i$ for $i = 1, 2, \dots, k-1$, while $a = \sigma(x_k)$ and $b = \tau(x_k)$ need not coincide. Thus, elements π of A are defined by the conditions

$$\pi(x_1) = a_1, \pi(x_2) = a_2, \dots, \pi(x_{k-1}) = a_{k-1}, \pi(x_k) = a,$$

while elements $\pi \in B$ are defined by the conditions

$$\pi(x_1) = a_1, \pi(x_2) = a_2, \dots, \pi(x_{k-1}) = a_{k-1}, \pi(x_k) = b.$$

Let $t_{a,b}$ denote the transposition of a and b . Consider the map

$$(2.6) \quad \varphi: \sigma H_k \ni j \mapsto t_{a,b} \circ j \in \tau H_k.$$

Clearly, φ is a bijection between $A = \sigma H_k$ and $B = \tau H_k$. The values of j and $t_{a,b} \circ j$ differ in at most two inputs, $x_k = j^{-1}(a)$ and $j^{-1}(b)$. Since $a, b \notin \{a_1, a_2, \dots, a_{k-1}\}$, it

follows that $j^{-1}(b) \notin \{x_1, x_2, \dots, x_{k-1}\}$ and $\mu(\{j^{-1}(b)\}) \leq \mu(\{x_k\})$. We conclude: for every $j \in A = \sigma H_k$

$$d_{unif}(j, \varphi(j)) \leq 2\mu(\{x_k\}).$$

Consequently, the metric space $\text{Aut}^*(X, \mu)$ has length at most $\ell = 2(\sum_{i=1}^n \mu(\{x_i\}^2))^{1/2}$, and Theorem 2.7 accomplishes the proof. $\square$

Corollary 2.12. *Let (X_n, μ_n) be a sequence of probability spaces with finitely many points in each, having the property that the mass of the largest atom in X_n goes to zero as $n \rightarrow \infty$. Then the family of automorphism groups $\text{Aut}^*(X_n, \mu_n)$, $n \in \mathbb{N}$, equipped with the uniform metric and the normalized counting measure, is Lévy, with*

$$\alpha_{X_n}(\varepsilon) \leq \exp\left(-\frac{\varepsilon^2}{32 \max_{x \in X_n} \mu_n(\{x\})}\right).$$

Proof. Same as the proof of Corollary 2.10. $\square$

2.3. Fixed points. If $X = (X, \mathcal{U})$ is a uniform space, then σX will denote the *Samuel*, or *universal, compactification* of X , that is, the space of maximal ideals of the C^* -algebra $\text{UCB}(X)$ of all uniformly continuous bounded complex-valued functions on X . (See [12].)

If a group G acts on X by uniform isomorphisms, this action uniquely extends to an action of G on σX by homeomorphisms.

The following result is essentially Th. 2.12 in [42], to which we give a different proof.

Theorem 2.13. *Let a group G act by uniform isomorphisms on a uniform space X . Suppose a net (ν_α) of probability measures on X concentrates and has the property: (*) for every $g \in G$, every $A \subseteq X$ and every $V \in \mathcal{U}_X$,*

$$\limsup_{\alpha} |\nu_\alpha(V[A]) - \nu_\alpha(g \cdot V[A])| < 1.$$

Then the Samuel compactification σX has a G -fixed point.

Proof. The net (ϕ_α) of means on $C(\sigma X)$, defined by $\phi_\alpha(f) = \int_X f(x) d\nu_\alpha(x)$, has a weak* cluster point, say ϕ . By selecting a subnet, assume that $\phi_\alpha \rightarrow \phi$ in weak* topology. As a consequence of Lemma 2.3, for any bounded uniformly continuous function f on X , the $L_1(\nu_\alpha)$ -distance between f and a suitable constant function (for instance, $M_\alpha(f)$, a ν_α -median of f) goes to zero as $\alpha \rightarrow \infty$. Consequently, the $L_1(\nu_\alpha)$ -distance between $f \cdot g$ and $M_\alpha(f) \cdot M_\alpha(g)$ tends to zero as well, and

$$\left| \int f g d\nu_\alpha - \left(\int f d\nu_\alpha \right) \cdot \left(\int g d\nu_\alpha \right) \right| \rightarrow_\alpha 0,$$

for all $f, g \in C(\sigma X)$, implying that the mean ϕ is multiplicative.

The condition $(*)$, jointly with concentration of (ν_α) , easily implies that whenever $\liminf_\alpha \nu_\alpha(A) > 0$, $g \in G$, and $V \in \mathcal{U}_X$, one has $\limsup_\alpha \nu_\alpha(VA \cap gVA) = 1$. Consequently, if $f \in C(\sigma X)$, there is a subnet (β) with $M_\beta(f) - M_\beta(gf) \rightarrow 0$, where ${}^g f(x) = f(g^{-1}x)$. We conclude: $\phi_\beta(f) - \phi_\beta(gf) \rightarrow_\beta 0$, and as $\phi_\beta(f) \rightarrow_\beta \phi(f)$, it follows that ϕ is a G -invariant mean. $\square$

Let X be a topological group G equipped with the right uniform structure, which has as a basis the entourages of the diagonal of the form $V_\uparrow = \{(x, y) \in G \times G \mid xy^{-1} \in V\}$, where V is a neighbourhood of e in G . The corresponding Samuel compactification, denoted by $S(G)$, is the *greatest ambit* of G , that is, the maximal compact G -space containing a distinguished point with an everywhere dense orbit. The existence of a G -fixed point in $S(G)$ is therefore equivalent to extreme amenability of G .

The condition $(*)$ is a weak form of invariance of the measures ν_α . In many examples, one can assume that the measures ν_α are virtually invariant, that is, for every $g \in G$, one has $g*\nu_\alpha = \nu_\alpha$ for all $\alpha \geq \alpha_0$. A topological group G is called a *Lévy group* [23, 15] if it contains a family $\mathcal{K}$ of compact subgroups, directed by inclusion, having everywhere dense union in G , and such that the corresponding normalized Haar measures, μ_K , on the groups $K \in \mathcal{K}$ concentrate in $G_\uparrow$.

We expand the concept of a Lévy group as follows.

Definition 2.14. Let us say that a topological group G is a *generalized Lévy group* if there is a net $(K_\alpha)_{\alpha \in A}$ of compact subgroups of G with the following properties:

1. The family of normalized Haar measures μ_α on K_α concentrates in G ;
2. For every non-empty open subset $V \subseteq G$ there is an $\alpha \in A$ such that $V \cap K_\beta \neq \emptyset$ for all $\beta \geq \alpha$.

Remark 2.15. The second condition is equivalent to the following, formally stronger, condition: for every finite collection $g_1, g_2, \dots, g_N \in G$, $N \in \mathbb{N}$, and for every neighbourhood V of the identity in G there is an $\alpha \in A$ such that for all $\beta \geq \alpha$ and $i = 1, 2, \dots, N$, one has $Vg_i \cap K_\beta \neq \emptyset$.

Theorem 2.16. *Every generalized Lévy group is extremely amenable.* $\square$

Proof. In view of Theorem 2.13, it is enough to verify the condition $(*)$. Assume it fails. Then for some $g \in G$, some (non-empty) $A \subset G$, and some symmetric neighbourhood of identity, V , one can construct a subnet (K_β) of groups with the properties $\mu_\beta(V^2A) \rightarrow_\beta 1$ and $\mu_\beta(gV^2A) \rightarrow_\beta 0$.

Because of condition (2) in the definition of a generalized Lévy group, one can assume without loss in generality that for all β , $gV \cap K_\beta \neq \emptyset$. Choose a net $g'_\beta \in K_\beta$ with $g = g'_\beta v$, where $v \in V$. Since $vV^2A \supseteq VA$, one has $\nu_\beta(gV^2A) \geq \nu_\beta(g'_\beta VA) = \nu_\beta(VA) \rightarrow_\beta 1$, meaning that $\nu_\beta(gV^2A) \rightarrow_\beta 1$ as well, a contradiction. $\square$

Corollary 2.17 (Gromov and Milman). *Every Lévy group is extremely amenable.* $\square$

To deduce Corollary 2.17 directly from Theorem 2.13, it is enough to note the extreme amenability of the everywhere dense subgroup $\cup \mathcal{K}$ of G , since both topological groups share the same greatest ambit.

There are still situations where one needs the condition $(*)$ in its full generality. For instance, this is how one proves the following result.

Theorem 2.18. *Let G be an amenable locally compact group and let X be a non-atomic Lebesgue measure space. Then the group $L_0(X, G)$ of all measurable maps from X to G , equipped with the topology of convergence in measure, is extremely amenable.*

Proof, sketch. By force of amenability of G , there is an invariant probability measure, ν , on the greatest ambit $\mathcal{S}(G)$ of G . Choose a net (ν_α) of probability measures on G (for instance, with finite support), weak* converging to ν . For every measurable partition $\mathcal{W}$ of X define a measure $\nu_{\alpha, \mathcal{W}}$ on $L_0(X, G)$, supported on the subgroup of all $\mathcal{W}$ -simple maps $X \rightarrow G$ (identified with the product $G^{|\mathcal{W}|}$), as the product measure, $\nu_{\alpha, \mathcal{W}} = \nu_\alpha^{\otimes |\mathcal{W}|}$. The net $(\nu_{\alpha, \mathcal{W}})$ has the desired properties: concentration follows e.g. from [48], p. 76 and Prop. 2.1.1, while the condition $(*)$ is checked directly. $\square$

In the case where G is a compact group, the theorem was established by Glasner [15] and (independently, unpublished) Furstenberg and Weiss. The above version is Thm. 2.2 in [42].

Remark 2.19. Notice that, for $G = \mathbb{R}$, the topological group $L_0(X)$ is extremely amenable, but not a Lévy group, nor indeed a generalized Lévy group in the sense of our Definition 2.14, because it contains no non-trivial compact subgroups.

Let us finally state a simple and useful fact that belongs to the folklore.

Lemma 2.20. *Let G be a topological group, and let $\mathcal{G}$ be a family of topological subgroups of G with the property: if $H, K \in \mathcal{G}$, then $H \cup K \subseteq L$ for some $L \in \mathcal{G}$. Suppose that $\cup \{G : G \in \mathcal{G}\}$ is everywhere dense in G and that every $H \in \mathcal{G}$ is contained in a suitable extremely amenable subgroup of G . Then G is extremely amenable.*

Proof. Let G act continuously on a compact space X . For every $H \in \mathcal{G}$ denote by F_H the set of all H -fixed points in X . As H is contained in a suitable extremely amenable subgroup of G , F_H is non-empty and compact. As $\mathcal{G}$ is upward directed, the family $\{F_H : H \in \mathcal{G}\}$ is centered, that is, for every finite subset $H_1, \dots, H_k \in \mathcal{G}$, one has $\cap_{i=1}^k F_{H_i} \neq \emptyset$. Hence, $\cap_{H \in \mathcal{G}} F_H \neq \emptyset$, and every element of this intersection is a fixed point for G . $\square$

Corollary 2.21. *Let G be a topological group containing a family of compact subgroups directed by inclusion and having an everywhere dense union. Let X be a non-atomic Lebesgue measure space. Then the group $L_0(X, G)$ of all measurable maps from X to G , equipped with the topology of convergence in measure, is extremely amenable.*

Proof. Let $\mathcal{K}$ be a family of compact subgroups of G as in the assumptions of the theorem. The group $L_0(X, \cup \mathcal{K})$ is everywhere dense in $L_0(X, G)$, and it is extremely amenable by Lemma 2.20, because each of the topological subgroups of the form $L_0(X, K)$, $K \in \mathcal{K}$ is extremely amenable by Theorem 2.18. $\square$

3. UNITARY GROUPS OF OPERATOR ALGEBRAS

3.1. A characterization of approximately finite dimensional von Neumann algebras. Let $\mathcal{H}$ be a Hilbert space, let $M \subset \mathcal{B}(\mathcal{H})$ be a von Neumann algebra, and let M_* be the predual of M . Recall that the *strong topology* on M , or the $s(M, M_*)$ -topology is the topology on M induced by the family of semi-norms

$$x \in M \mapsto \|x\|_\varphi = \varphi(x^*x)^{1/2}, \quad \varphi \in M_*^+$$

The $s(M, M_*)$ -topology on M is equal to the restriction of the $s(\mathcal{B}(\mathcal{H}), \mathcal{B}(\mathcal{H})_*)$ -topology to M .

Let us mention the following technical result we will need.

Lemma 3.1. *Let A be an index set and let $(M_\alpha)_{\alpha \in A}$ be a family of von Neumann algebras. If M denotes the direct sum $\oplus_{\alpha \in A} M_\alpha$, then $U(M)$, endowed with the $s(M, M_*)$ -topology, is isomorphic (as a topological group) to $\prod_{\alpha \in A} U(M_\alpha)$, where each $U(M_\alpha)$ is endowed with the $s(M_\alpha, (M_\alpha)_*)$ -topology and $\prod_{\alpha \in A} U(M_\alpha)$ with the product topology.*

Proof. As

$$M_* = \{(\varphi_\alpha)_{\alpha \in A} : \varphi_\alpha \in (M_\alpha)_*, \sum_{\alpha \in A} \|\varphi_\alpha\| < \infty\},$$

the proof is clear. $\square$

The main result of this section is the following.

Theorem 3.2. *If M is an injective von Neumann algebra, then its unitary group $U(M)$ with the $s(M, M_*)$ -topology is the direct product of a compact group and an extremely amenable group.*

Let M be an injective von Neumann algebra. By [11], Thm. 4, M is generated by an upwards directed collection $(M_\alpha)_{\alpha \in A}$ of countably generated injective von Neumann subalgebras. By [28], Lemma 4, $U(M)$ is the $s(M, M_*)$ -closure of the upward directed collection of groups $U(M_\alpha)$. (The separability condition in [28] is not essential, as remarked by Haagerup in [25].) A countably generated injective von Neumann algebra is a direct sum of injective von Neumann algebras with separable predual. By Lemma 3.1, we can therefore assume in the proof of Theorem 3.2 that M_* is separable.

Moreover, M can be decomposed as a direct sum of finite von Neumann algebras of type I and type II, and of a properly infinite von Neumann algebra. Therefore Theorem 3.2 will follow directly from Proposition 3.5, Corollary 3.7 and Proposition 3.8.

Lemma 3.3. *Let $\mathbb{S}$ denote the unit sphere of a Hilbert space $\mathcal{H}$, and let (X, μ) be a finite measure space. The restriction of the $s(L^\infty(X, \mu; \mathcal{H}), L^1(X, \mu; \mathcal{H}))$ -topology to $L^\infty(X, \mu; \mathbb{S})$ coincides with the restriction of the topology of convergence in measure.*

Proof. An easy geometric property of the unit ball $\mathbb{B}$ of $\mathcal{H}$ (uniform convexity) implies that for every $a, b \in \mathbb{S}$ with $d(a, b) > \varepsilon$ one has $\langle a, a_0 \rangle < \sqrt{1 - \varepsilon^2}$. Consequently, for every function $f: X \rightarrow \mathbb{S}$ the condition $\int_X (f(x) - a) d\mu(x) < 1 - \sqrt{1 - \varepsilon^2}$ implies $\int_X |f(x) - a| d\mu(x) < \varepsilon$.

It remains to notice that the L^1 -distance induces the topology of convergence in measure on every $\|\cdot\|_\infty$ -bounded subset of $L^\infty(X, \mu; \mathcal{H})$, such as $L^\infty(X, \mu; \mathbb{S})$. $\square$

Corollary 3.4. *Let $\mathcal{H}$ be a finite-dimensional Hilbert space, and let (X, μ) be a finite measure space. The $s(L^\infty(X, \mu; \mathcal{B}(\mathcal{H})), L^1(X, \mu; \mathcal{B}(\mathcal{H})_*))$ -topology restricted to $L^\infty(X, \mu; U(\mathcal{H}))$ coincides with the restriction of the topology of convergence in measure on $L^\infty(X, \mu; U(\mathcal{H}))$.*

Proof. The unitary group $U(\mathcal{H})$ is a subset of the unit sphere of $\mathcal{B}(\mathcal{H})$ with the Hilbert-Schmidt norm. $\square$

Proposition 3.5. *If M is a finite von Neumann algebra of type I, then $U(M)$, with the $s(M, M_*)$ -topology, is the direct product of a family of compact groups and extremely amenable groups.*

Proof. By the structure theorem for finite type I von Neumann algebras (see [46], Theorem V.1.27), M is isomorphic to a direct sum of algebras of the form $L^\infty(\Gamma, \mu) \otimes \mathcal{B}(\mathcal{H})$, where $\dim \mathcal{H} < \infty$ and Γ is a locally compact space, with a positive (finite) Radon measure μ .

If μ is atomic, then

$$U(L^\infty(\Gamma, \mu) \otimes \mathcal{B}(\mathcal{H})) \cong \prod_{\text{supp } \mu} U(\mathcal{H})$$

and therefore is a compact group.

If μ is non-atomic, then the unitary group $U(L^\infty(\Gamma, \mu) \otimes \mathcal{B}(\mathcal{H}))$ is canonically isomorphic to $L^\infty(\Gamma, \mu; U(\mathcal{H}))$. It follows that $U(L^\infty(\Gamma, \mu) \otimes \mathcal{B}(\mathcal{H}))$ is extremely amenable by Corollaries 3.4 and 2.21. By Lemma 3.1, the Proposition is proved. $\square$

Proposition 3.6. *Let M be a von Neumann algebra and let $(N_k)_{k \geq 1}$ be an increasing sequence of finite-dimensional subfactors of M such that $N_\infty = \cup_{k \geq 1} N_k$ is $s(M, M_*)$ -dense in M . Then $U(M)$, with the $s(M, M_*)$ -topology, is a Lévy group.*

Proof. For $k \geq 1$, let $(e_{i,j}^k)_{1 \leq i,j \leq n_k}$ be a system of matrix units of N_k . If $u \in U(N_k)$, let $(u_{ij}) \in U(n_k, \mathbb{C})$ denote the matrix defined by $u = \sum_{i,j=1}^{n_k} u_{ij} e_{ij}^k$. Let $SU(N_k) = \{u \in U(N_k) : \det(u_{ij}) = 1\}$.

Let $\varphi \in M_*$ be a faithful state and let $\varepsilon > 0$ be given. As $\varphi(1) = \sum_{i=1}^{n_k} \varphi(e_{ii}^k)$, there exists K such that for $k \geq K$, $\varphi(e_{ii}^k) \leq \varepsilon/4$ for some $1 \leq i \leq n_k$.

Let $u \in U(N)$. By [28], Lemma 4, there exist $k \geq K$ and $u_k \in U(N_k)$ such that $\|u - u_k\|_\varphi \leq \varepsilon/2$. Let $1 \leq i \leq n_k$ be such that $\varphi(e_{ii}^k) \leq \varepsilon/4$ and $v_k = 1 - e_{ii}^k + \overline{\det(u_{ij}^k)}e_{ii}^k \in N_k$. Then $u_k v_k \in SU(N_k)$ and

$$\begin{aligned} \|u - u_k v_k\| &\leq \|u - u_k\|_\varphi + \|u_k(1 - v_k)\|_\varphi \\ &\leq \frac{\varepsilon}{2} + \|1 - v_k\|_\varphi \\ &\leq \frac{\varepsilon}{2} + |1 - \det(u_{ij}^k)|\varphi(e_{ii}^k) \\ &\leq \frac{\varepsilon}{2} + 2\frac{\varepsilon}{4} = \varepsilon. \end{aligned}$$

Hence, $\cup_{k \geq 1} SU(N_k)$ is $s(M, M_*)$ -dense in $U(M)$.

As $N_k \cong M_{n_k}(\mathbb{C})$, there exists $h_\varphi \in M_{n_k}(\mathbb{C})$, $h_\varphi > 0$ such that for $x = \sum_{i,j=1}^{n_k} x_{i,j} e_{i,j}^k \in N_k$, $\varphi(x) = \text{Tr}(h_\varphi(x_{i,j}))$, where Tr denotes the trace on $M_{n_k}(\mathbb{C})$. As $\varphi(1) = \text{Tr}(h_\varphi) = 1$, $\|h_\varphi\| \leq 1$ and for $x \in N_k$,

$$\|x\|_\varphi \leq \|(x_{ij})\|_2,$$

where $\|\cdot\|_2$ denotes the Hilbert–Schmidt norm. By [23], $(SU(N_k))_{k \geq 1}$ forms a Lévy family with regard to the Hilbert-Schmidt metric, and consequently the $\|\cdot\|_\varphi$ -metric, which ends the proof of Proposition. $\square$

Let M be a properly infinite injective von Neumann algebra with separable predual. By [11], there exists an increasing sequence $(N_k)_{k \geq 1}$ of finite dimensional subfactors whose union is $s(M, M_*)$ -dense in M . By Proposition 3.6, we get

Corollary 3.7. *Let M be a properly infinite injective von Neumann algebra with separable predual. Then the unitary group $U(M)$, with the $s(M, M_*)$ -topology, is extremely amenable.* $\square$

Let M be a finite continuous injective von Neumann algebra with separable predual. By considering the standard representation of M , we can assume that M acts on a separable Hilbert space. Then by reduction theory and uniqueness of the injective factor of type II_1 , M is isomorphic to $L^\infty(X, \mu) \otimes R$, where X is a standard Borel space, with a finite measure μ and R is the hyperfinite factor of type II_1 .

By Proposition 3.6 and Corollary 2.17, $U(R)$, endowed with the $s(R, R_*)$ -topology, is extremely amenable. By the same argument as in the proof of 3.5, if μ is an atomic measure, then $U(M)$ is isomorphic to $\prod_{\text{supp} \mu} U(R)$, equipped with the product $s(R, R_*)$ -topology. If μ is a non-atomic measure, then $U(M)$, with the $s(M, M_*)$ -topology, is isomorphic to the group of all measurable maps from X to $U(R)$, equipped with the topology of convergence in measure. Then by Lemma 3.1, Corollary 2.21, and Corollary 3.7 we get

Proposition 3.8. *If M is an injective finite and continuous von Neumann algebra with separable predual, then $U(M)$, endowed with the $s(M, M_*)$ -topology, is extremely amenable.* $\square$

3.2. A characterization of nuclear C^* -algebras. Recall that a continuous action of a topological group G on a compact space X is *proximal* if for every two points x, y and every entourage of the diagonal $V \subseteq X \times X$ there is a $g \in G$ with $(gx, gy) \in V$. A topological group G is called *strongly amenable* if every continuous proximal action of G on a compact space has a fixed point [14]. For instance, every compact group is strongly amenable simply because it has no non-trivial proximal actions on compacta. Every extremely amenable group is strongly amenable for obvious reasons. The class of strongly amenable groups is closed under direct products.

By the *completion* $\hat{G}$ of a topological group G we mean, as usual, the completion of G with regard to the two-sided uniform structure, that is, the supremum of the left and the right uniform structures on G . This is again a topological group. (See, e.g., Chapter 10 in [45].)

It is a standard fact, easily proved, that the completion, $\hat{G}$, of a topological group G is amenable (strongly amenable, extremely amenable) if and only if G is amenable (resp. strongly amenable, extremely amenable).

The following corollary of our results strengthens a result due to Paterson [39] (Thm. 2).

Theorem 3.9. *A C^* -algebra A is nuclear if and only if its unitary group $U(A)$, equipped with the topology $s(A, A^*)$, is strongly amenable.*

Proof. Denote by M the universal von Neumann envelope of A . As is well known, M can be obtained as the completion of the C^* -algebra A with regard to the additive uniform structure determined by the relative weak topology. ([10], Corollaire 12.1.3.) As a consequence, one can prove that the completion, $\widehat{U(A)}$, of the unitary group $U(A)$, equipped with the relative weak topology, is isomorphic to the unitary group of the von Neumann envelope M of A , equipped with the $s(M, M_*)$ -topology.

Necessity ($\Rightarrow$): if A is nuclear, then the von Neumann envelope M is approximately finite dimensional, and therefore, by Theorem 3.2, the unitary group $U(M)$ with the topology $\sigma(A, A^*)$ is the product of a compact group and an extremely amenable group. In particular, $U(M)$ is strongly amenable, and so is its everywhere dense topological subgroup $U(A)$.

Sufficiency ($\Leftarrow$): follows from de la Harpe's result in [28], as noted by Paterson [39]. If the topological group $U(A)$ is strongly amenable, then it is in particular amenable, and so is the topological group completion, $U(M)$. It follows by de la Harpe's result that M is approximately finite-dimensional. $\square$

3.3. Automorphism groups of von Neumann algebras. If M is a von Neumann algebra, let $\text{Aut}(M)$ denote, as usual, the group of all $*$ -automorphisms of M . We

consider on $\text{Aut}(M)$ the topology of norm pointwise convergence for the action of $\text{Aut}(M)$ on M_* , given by

$$\alpha \in \text{Aut}(M), \varphi \in M_* \mapsto \varphi \circ \alpha^{-1} \in M_*.$$

With this topology, called the *u-topology*, $\text{Aut}(M)$ is a topological group and it is a Polish group if M_* is separable.

If $u \in U(M)$, let $\text{Ad } u$ denote the inner automorphism of M given by $\text{Ad } u(x) = uxu^*$, $x \in M$, and let $\text{Inn}(M)$ be the subgroup of all inner automorphisms of M .

If $U(M)$ is endowed with the $s(M, M_*)$ -topology and $\text{Inn}(M)$ with the *u-topology*, the canonical surjection $U(M) \rightarrow \text{Inn}(M)$ is continuous, as for $\varphi \in M_*$, $u \in U(M)$

$$\|\varphi \circ \text{Ad } u - \varphi\| \leq 2 \|\varphi\| \|u - 1\|_\varphi.$$

By Theorem 3.2, we therefore get:

Proposition 3.10. *If M is an injective continuous von Neumann algebra, then $\text{Inn}(M)$, with the topology of pointwise convergence in norm on M_* , is extremely amenable. $\square$*

4. GROUPS OF TRANSFORMATIONS OF SPACES WITH MEASURE

4.1. Weak and uniform topologies. For a standard non-atomic Lebesgue measure space $X = (X, \mu)$ denote by $\text{Aut}(X, \mu)$ the group of (equivalence classes of) invertible measure-preserving transformations of X , and by $\text{Aut}^*(X, \mu)$ the group of all invertible measurable and non-singular transformations of X .

The *weak topology* on $\text{Aut}^*(X, \mu)$ is the weakest topology making continuous every function

$$\text{Aut}^* X \ni \tau \mapsto \mu(A \cap \tau(A)) \in \mathbb{R},$$

where $A \subseteq X$ is a measurable subset. It is enough to consider elements of a generating family of measurable sets. Equivalently, the weak topology is induced by the strong operator topology on the isometry group of $L^p(X)$ under the quasi-regular representation of the group $\text{Aut}^*(X, \mu)$ in the Banach space $L^p(X)$, where $1 \leq p < \infty$ is any. (See [6], Theorem 8.)

The weak topology makes $\text{Aut}^*(X, \mu)$ into a Polish topological group, while $\text{Aut } X$ is a closed (therefore also Polish) subgroup.

Define a left-invariant metric on $\text{Aut}^*(X, \mu)$ as follows:

$$d_{\text{unif}}(\tau, \sigma) = \mu\{x \in X : \tau(x) \neq \sigma(x)\}.$$

The topology induced on $\text{Aut}^*(X)$ by the metric d is a group topology, known as the *uniform topology*. It is strictly finer than the strong topology ([49], Prop. 3). For example, the uniform topology makes $\text{Aut}^*(X, \mu)$ into a (path-connected) non-separable group.

Both the weak topology and the uniform topology only depend on the equivalence class of the measure μ , rather than on μ itself.

Sometimes we will indicate the uniform topology by the subscript u , as in $\text{Aut}(X, \mu)_u$. Similarly, for the weak topology the subscript w will be used.

Note that the restriction of d to $\text{Aut}(X, \mu)$ is bi-invariant, and so the topological group $\text{Aut}(X, \mu)_u$ is SIN (= has small invariant neighbourhoods), that is, the left and the right uniform structures coincide.

4.2. Extreme amenability of $\text{Aut}(X, \mu)$ with the weak topology. The following theorem (cf. e.g. [26], pp. 65–68), belonging to the Kakutani–Rokhlin circle of results, is well-known. Identify the symmetric group $\mathfrak{S}_n$ with the subgroup of all measure-preserving automorphisms of $\mathbb{I} = [0, 1]$ with the Lebesgue measure mapping each dyadic interval of rank n onto a dyadic interval of rank n via a translation (the interval exchange transformations).

Theorem 4.1 (Weak Approximation Theorem). *The union of the subgroups $\mathfrak{S}_n$, $n \in \mathbb{N}$ is everywhere dense in $\text{Aut}(\mathbb{I})$ with respect to the weak (=strong) topology.* $\square$

The *normalized Hamming distance* on the finite permutation group $\mathfrak{S}_n$ of rank $n \geq 1$ is given by

$$(4.1) \quad d_n(\sigma, \tau) = \frac{1}{n} |\{i: \sigma(i) \neq \tau(i)\}|.$$

Under the above embedding $\mathfrak{S}_n \hookrightarrow \text{Aut}(\mathbb{I})$, the normalized Hamming distance on $\mathfrak{S}_n$ coincides with the restriction of the metric d .

Theorem 4.2. *The group $\text{Aut}(X, \mu)_w$ of all measure-preserving automorphisms of a standard non-atomic finite or sigma-finite measure space, equipped with the weak topology, is a Lévy group and therefore extremely amenable.*

Proof. For X finite, the group $\text{Aut}(X, \mu)$ is topologically isomorphic with the group $\text{Aut}(\mathbb{I}, \lambda)$, where $\mathbb{I} = [0, 1]$ is the unit interval with the Lebesgue measure. According to Maurey’s theorem [35], the permutation groups $\mathfrak{S}_n$ of finite rank n , equipped with the normalized counting measure and the normalized Hamming distance, form a Lévy family (even with regard to the uniform topology). By Theorem 4.1, $\text{Aut}(X, \mu)_w$ forms a Lévy group, and Corollary 2.17 applies.

For μ sigma-finite it is enough to prove the result for the real line $\mathbb{R}$, equipped with the Lebesgue measure λ . For every $n \in \mathbb{N}$, let G_n denote the group of interval exchange transformations of the system of n^2 intervals of length $1/n$ each, covering the interval $[-n, n]$. A natural group isomorphism from the permutation group S_{n^2} of rank n^2 equipped with the normalized Hamming distance onto G_n , equipped with the uniform metric, is Lipschitz with $L = n$. By Lemma 2.2 and Maurey’s result (Example 2.6), the family (G_n) is a Lévy family, with

$$\alpha_{G_n}(\varepsilon) \leq \exp(-\varepsilon^2 n^2 / 64n) = \exp(-\varepsilon^2 n / 64).$$

Since the groups (G_n) form an increasing chain with everywhere dense union in $\text{Aut}(\mathbb{R}, \lambda)$, we are done. $\square$

4.3. Non-amenability of $\text{Aut}(X, \mu)$ with uniform topology. Let π be a unitary representation of a group G in a Hilbert space $\mathcal{H}$, that is, $\mathcal{H}$ is a unitary G -module. One says that π (or the G -module $\mathcal{H}$) is *amenable in the sense of Bekka* [2] if there exists a state, ϕ , on the von Neumann algebra $\mathcal{B}(\mathcal{H})$ that is invariant under the action of G by conjugations: for all $T \in \mathcal{B}(\mathcal{H})$ and all $g \in G$,

$$\phi(T) = \phi(\pi_g^* T \pi_g).$$

Denote by $\mathbb{S}$ the unit sphere in the space $\mathcal{H}$. Notice that G acts on the space $\mathbb{C}^{\mathbb{S}}$ of all functions $\mathbb{S} \rightarrow \mathbb{C}$ by

$${}^g f(\xi) := f(\pi_g^*(\xi)) \equiv f(\pi_{g^{-1}}(\xi)),$$

where $g \in G$, $f \in \mathbb{C}^{\mathbb{S}}$, and $\xi \in \mathbb{S}$. This action leaves invariant the space $\text{UCB}(\mathbb{S})$ of all uniformly continuous bounded complex-valued functions on the sphere. The following was proved by one of the present authors [41].

Theorem 4.3. *A unitary representation π of a group G is amenable if and only if there exists a G -invariant mean on the space $\text{UCB}(\mathbb{S})$. $\square$*

Let $\xi \in \mathbb{S}$. For every (complex-valued) function f on $\mathbb{S}$ denote by $\tilde{f}$ a function on G defined as follows:

$$\tilde{f}(g) := f(\pi_g \xi),$$

where $g \in G$. If $f \in \ell_{\infty}(\mathbb{S})$, then $\tilde{f} \in \ell_{\infty}(G)$. The latter is a Banach G -module with respect to the left regular representation of G :

$${}^g \varphi(x) := \varphi(g^{-1}x).$$

Lemma 4.4. *The mapping $f \mapsto \tilde{f}$ from $\ell_{\infty}(\mathbb{S})$ to $\ell_{\infty}(G)$ is G -equivariant, that is, commutes with the action of G .*

Proof. $\tilde{{}^g f}(h) = {}^g f(\pi_h(\xi)) = f(\pi_g^* \pi_h(\xi)) = f(\pi_{g^{-1}h}(\xi)) = \tilde{f}(g^{-1}h) = {}^g(\tilde{f})(h)$. $\square$

Now assume that G is a topological group and the representation π is strongly continuous.

Lemma 4.5. *If $f \in \text{UCB}(\mathbb{S})$, then $\tilde{f}$ is left uniformly continuous on G .*

Proof. Let $\varepsilon > 0$. Find a $\delta > 0$ such that, whenever $\|\xi - \zeta\| < \delta$, one has $|f(\xi) - f(\zeta)| < \varepsilon$. Due to the strong continuity of π , there is a neighbourhood of the identity, V , in G such that, whenever $g \in V$, one has $\|\xi - \pi_g \xi\| < \delta$. If now $g, h \in G$ are such that $g^{-1}h \in V$, then $|\pi_{g^{-1}h}(\xi) - \xi| < \delta$, that is, $|\pi_h(\xi) - \pi_g(\xi)| < \delta$, and $|\tilde{f}(g) - \tilde{f}(h)| = |f(\pi_g(\xi)) - f(\pi_h(\xi))| < \varepsilon$. $\square$

Denote by $\text{LUCB}(G)$ the C^* -algebra of all bounded left uniformly continuous complex-valued functions on G . It is now easy to verify that the mapping $f \mapsto \tilde{f}$ is a positive linear operator from $\text{UCB}(\mathbb{S})$ to $\text{LUCB}(G)$, sending 1 to 1 and commuting with the

action of G . Therefore, if there is an invariant mean on $\text{LUCB}(G)$, by composing it with the mapping $f \mapsto \tilde{f}$, one gets an invariant mean on $\text{UCB}(\mathbb{S})$. We obtain:

Proposition 4.6. *If a topological group G admits a left-invariant mean on the space $\text{LUCB}(G)$ of all left uniformly continuous bounded functions on G , then every strongly continuous unitary representation of G is amenable.* $\square$

In particular, if G is a SIN group, that is, the left and the right uniformities on G coincide, we obtain the following.

Corollary 4.7. *Every strongly continuous unitary representation of an amenable SIN group is amenable.* $\square$

Remark 4.8. In general, the condition that G is SIN cannot be dropped here, unless we presume that G is locally compact. For instance, the infinite unitary group $U(\ell_2)$ with the strong operator topology is (extremely) amenable, while the standard unitary representation of $U(\ell_2)$ is non-amenable (because this would clearly imply, for instance, amenability of every unitary representation of every countable group).

Theorem 4.9. *The topological group $\text{Aut}(X, \mu)_u$ is non-amenable.*

Proof. Take as $V = L_0^2(X)$ the closed unitary G -submodule of $L^2(X)$ consisting of all functions $f \in L^2(X)$ with

$$\int_X f(x) d\mu(x) = 0.$$

According to Proposition 4.6, if $\text{Aut}(X, \mu)_u$ were amenable, then the regular representation γ of $\text{Aut}(X, \mu)_u$ in $L_0^2(X)$ would be amenable. In particular, so would be the restriction of γ to any subgroup G of $\text{Aut}(X, \mu)_u$. It is, therefore, enough to discover a group G acting on a standard non-atomic Lebesgue space X by measure-preserving transformations in such a manner that the associated regular representation of G in $L_0^2(X)$ is not amenable in the sense of Bekka.

The following is inspired by Bekka's work [3]. Let G be a semisimple real Lie group with finite centre and without compact factors, having property (T). (Cf. [3]; e.g. $SL_3(\mathbb{R})$.) Let Γ denote a lattice in G , that is, a discrete subgroup such that the Haar measure induces an invariant finite measure on G/Γ . Set $X = G/\Gamma$.

Assume that the standard representation π of G in $L_0^2(X) = L_0^2(G/\Gamma)$ is amenable. Since G is a Kazhdan group, according to a result by Bekka [2], π must have a finite-dimensional subrepresentation. (Later it was shown [4] that this property characterizes groups with property (T).)

Note that $L_0^2(X)$ has nontrivial G -invariant vectors: since the action of G on X is transitive, such vectors must be constant functions. Therefore, a finite-dimensional subrepresentation of π must be non-trivial. But this would result in a non-trivial continuous group homomorphism from G to a finite-dimensional unitary group, which is impossible by assumption. $\square$

4.4. Groups of non-singular transformations. Let (X, μ) be a standard Lebesgue measure space. We will prove that $\text{Aut}^*(X, \mu)$ with the weak topology forms a generalized Lévy group.

Following Tulcea [49], call a measurable mapping $f: X \rightarrow Y$ between two finite measure spaces a *Lebesgue mapping* if for every measurable $A \subseteq X$

$$\mu_Y(f(A)) = \frac{\mu_Y(Y)}{\mu_X(X)} \mu_X(A).$$

If X and Y are measurable subsets of a standard Lebesgue probability space, there is always a Lebesgue mapping from X onto Y .

Definition 4.10. Say that a non-singular transformation $g \in \text{Aut}^*(X, \mu)$ is a *Lebesgue permutation* with respect to a finite measurable partition $\mathcal{P}$ of X , if

1. for each element $A \in \mathcal{P}$ the image $g(A)$ also belongs to $\mathcal{P}$, and the restriction $g|_A$ is a Lebesgue mapping, and
2. if $A \in \mathcal{P}$ and $g(A) = A$, then $g|_A = \text{Id}_A$.

The following was deduced by Tulcea ([49], Theorem 1) from a theorem, attributed to Linderholm.

Theorem 4.11. *Lebesgue permutations are everywhere dense in the uniform topology on the group $\text{Aut}^*(X, \mu)$. Moreover, given a transformation $\tau \in \text{Aut}^*(X, \mu)$, an $\varepsilon > 0$, and a finite measurable partition $\mathcal{Q}$ of X , there exists a finite partition $\mathcal{Q}_1 \prec \mathcal{Q}$ and a Lebesgue permutation σ with respect to $\mathcal{Q}_1$ such that $d_{\text{unif}}(\tau, \sigma) < \varepsilon$. $\square$*

Lemma 4.12. *Let (X, μ) be the standard Lebesgue probability space, let $\tau_1, \tau_2, \dots, \tau_n$ be an arbitrary finite collection of transformations from $\text{Aut}^*(X, \mu)$, and let V be a strong neighbourhood of the identity in $\text{Aut}^*(X, \mu)$. Then there exist*

1. a finite partition $\mathcal{W}$ of X ,
2. a subgroup $\mathfrak{S}_w$ of $\text{Aut}^*(X, \mu)$ isomorphic to the permutation group of rank $w = |\mathcal{W}|$ and consisting of Lebesgue permutations with respect to $\mathcal{W}$, and
3. transformations $\sigma_1, \sigma_2, \dots, \sigma_n \in \mathfrak{S}_w$ such that $\tau_i \sigma_i^{-1} \in V$ for all $i = 1, 2, \dots, n$.

Proof. Let V be determined by an $\varepsilon > 0$ and a finite measurable partition $\mathcal{Q}$ of X , so that

$$V = \{\tau \in \text{Aut}^*(X, \mu) : \|\chi_A - \tau\chi_A\|_1 < \varepsilon \text{ for each } A \in \mathcal{Q}\}.$$

Using Theorem 4.11, choose for every $i = 1, 2, \dots, n$ a finite partition $\mathcal{Q}_i \prec \mathcal{Q}$ refining $\mathcal{Q}$ and a Lebesgue permutation σ_i with respect to $\mathcal{Q}_i$ such that $d_{\text{unif}}(\tau_i, \sigma_i) < \varepsilon$. Form a new partition $\mathcal{R} = \bigvee_{i=1}^n \mathcal{Q}_i$. The transformations σ_i need not be Lebesgue permutations with respect to $\mathcal{R}$.

Let $r = |\mathcal{R}|$. Denote by $\Sigma(\mathcal{R})$ the algebra of sets generated by $\mathcal{R}$. Let $0 < r_1 < r_2 < \dots < r_s$ be the collection of all real numbers $\mu(A)/\mu(B)$, $A, B \in \Sigma(\mathcal{R})$, $\mu(B) > 0$.

Choose a $\delta > 0$ so small that each of the numbers r_i , $i = 1, 2, \dots, s$, is approximated to within $\varepsilon/2$ by a suitable integer power of $(1 + \delta)$, say $m_1, m_2, \dots, m_s$:

$$(4.2) \quad |(1 + \delta)^{m_i} - r_i| < \frac{\varepsilon}{2}.$$

Let N, M be so large that

$$(4.3) \quad \frac{r}{N} < \frac{\varepsilon}{16},$$

$$(4.4) \quad \frac{\max_{i=1}^s |m_i|}{M} < \frac{\varepsilon}{8},$$

and

$$(4.5) \quad \frac{r(1 + \delta)}{M\delta} < 1.$$

For every $i = 1, 2, \dots, M$ denote

$$N_i = \lfloor (1 + \delta)^{i-1} N \rfloor.$$

Denote by $\mathcal{W} = \mathcal{P}_X(M, N, \delta)$ a partition of Y into the sets X_{ij} , $i = 1, 2, \dots, M$, $j = 1, 2, \dots, N_i + 1$, where for every $i = 1, 2, \dots, M$ the sets X_{ij} , $j = 1, 2, \dots, N_i$ have equal measure $\mu(X)/(MN(1 + \delta)^{i-1})$, while the remainder set X_{i, N_i+1} has measure

$$\frac{\mu(Y) \langle (1 + \delta)^{i-1} N \rangle}{MN(1 + \delta)^{i-1}}.$$

(The angular brackets denote the fractional part of a number.) Thus, the elements of our partition are arranged in M rows with $N_i + 1$ entries in each of them, so that the total measure of pieces making up each row is $\frac{1}{M}$, and all the bits in each row save perhaps the last one are of equal measure.

The total measure of all the M remainder sets does not exceed $\mu(X)(1 + \delta)/(\delta MN) < \mu(X)/N$. (Sum of a geometric progression and the condition (4.5).)

We can construct $\mathcal{W}$ so as to satisfy two additional conditions:

(i) each element $A \in \mathcal{R}$ and each row of $\mathcal{W}$ and are ‘independent events,’ that is, for each $i = 1, \dots, M$, $\mu(A) = \mu(A \cap \cup_{j=1}^{N_i} X_{i,j}) / \mu(\cup_{j=1}^{N_i} X_{i,j})$, and

(ii) for each $A \in \mathcal{R}$, in every row of $\mathcal{W}$ at most one element has non-null intersection with both A and $X \setminus A$.

Such a partition can be chosen by first forming the rows and then subdividing them into bits X_{ij} .

Thus, if we denote for an $A \in \mathcal{R}$

$$(4.6) \quad \tilde{A} = \bigcup_{X_{i,j} \subseteq A} X_{i,j},$$

then $\mu(A \setminus \tilde{A}) < N^{-1}$.

Fix a group of Lebesgue permutations, $\mathfrak{S}_w$, with respect to the partition $\mathcal{W}$, where $w = |\mathcal{W}|$, in the sense of Definition 4.10. Let σ be an arbitrary Lebesgue permutation

σ formed with respect to an *arbitrary* finite partition $\mathcal{Q}' \prec \mathcal{R}$. We claim that $\sigma \in V\mathfrak{S}_w$, which will finish the proof.

Let $A \in \mathcal{Q}'$. Denote $B = \sigma(A) \in \mathcal{Q}' \prec \mathcal{R}$. Let $A = \cup_i A_i$, $B = \cup_j B_j$, where $A_i, B_j \in \mathcal{R}$. Denote $\tilde{A} = \cup_i \tilde{A}_i$, $\tilde{B} = \cup_j \tilde{B}_j$.

The number of elements in the i -th row of $\mathcal{W}$ entirely contained in A is $\sum_i [\mu(A_i)N_i]$. This number differs from $[N_i\mu(A)]$ by no more than $r = |\mathcal{R}|$. Computing the sum of geometric progression, we conclude that $\mu(A \setminus \tilde{A}) < r(1 + \delta)/(MN\delta) + 1/N < 2/N < \varepsilon/8$.

A similar estimate is available for B . One conclusion is that $\mu(B \setminus \tilde{B}) < \varepsilon/4$. Also, it follows that the number of ‘unbroken’ elements contained in the i -th row in A differs from the number of ‘unbroken’ elements in the $(i + m)$ -th row in B by no more than r .

Without loss in generality, we may assume that $\mu(B) \geq \mu(A)$. Choose an $m = m_i$ such that $(1 + \delta)^m$ approximates the ratio $\mu(B)/\mu(A) \geq 1$ to within $\varepsilon/2$.

Construct a maximal partial bijection, $f = f^A$, between finite subsets of $\mathcal{W}$, with the following property. Suppose X_{ij} belongs to the domain of f^A . Then:

1. $X_{ij} \subseteq A$,
2. $X_{i',j'} = f^A(X_{ij}) \subseteq B$, and
3. $i' = i + m$.

In each row $i \leq M - m$, the number of elements of $\mathcal{W}$ that intersect A and do not belong to the domain of f^A , is at most r . The same applies to the elements of $\mathcal{W}$ contained in B that are not in the range of f^A , provided the row number is $\leq N + m$.

The total measure of the union of m rows in A outside of the domain of f^A is less than $m\mu(A)/M < \varepsilon\mu(A)/8$, and the total measure of the union of elements of m rows in B not in the range of f^A is $< m\mu(B)/M < \varepsilon\mu(B)/8$.

Denote by $\hat{A} = \cup \text{dom } f^A \subseteq \tilde{A}$, and $\hat{B} = \cup \text{range } f^A \subseteq \tilde{B}$. We conclude:

$$(4.7) \quad \mu(A \setminus \hat{A}) < \mu(A \setminus \tilde{A}) + \frac{\varepsilon}{8} < \frac{\varepsilon}{4}.$$

Similarly,

$$(4.8) \quad \mu(B \setminus \hat{B}) < \frac{\varepsilon}{4}.$$

Select such a partial bijection f^A for each $A \in \mathcal{Q}'$. Now coalesce the partial bijections f^A to obtain a partial bijection f between two subsets of $\mathcal{W}$. Extend f to a full permutation f' of $\mathcal{W}$. Let ς be the Lebesgue permutation from $\mathfrak{S}_w$ corresponding to f' , that is, $\varsigma(Y) = f'(Y)$ for each $Y \in \mathcal{W}$.

Clearly, for every $A \in \mathcal{Q}'$, the Radon-Nikodym derivative of $\varsigma_*(\mu)$ assumes at all elements $x \in \tilde{B}$ the constant value $(1 + \delta)^m$ and thus differs from $(d\sigma_*(\mu)/d\mu)(x)$ (which equals $\mu(B)/\mu(A)$) by less than $\varepsilon/2$, by virtue of our choice of m .

Finally, one has for every $A \in \mathcal{Q}$:

$$\begin{aligned} \|\sigma\chi_A - \varsigma\chi_A\|_1 &= \int_X \left| \chi_{\sigma(A)}(x) \frac{d\sigma_*\mu}{d\mu}(x) - \chi_{\varsigma(A)}(x) \frac{d\varsigma_*\mu}{d\mu}(x) \right| d\mu(x) \\ &\leq \int_{\tilde{B}} \left| \frac{d\sigma_*\mu}{d\mu}(x) - \frac{d\varsigma_*\mu}{d\mu}(x) \right| d\mu(x) + 2\mu(A \setminus \tilde{A}) \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

□

Theorem 4.13. $\text{Aut}^*(X, \mu)_w$ with the weak topology is a generalized Lévy group and therefore extremely amenable.

Proof. Fix a countable basis $V_1, V_2, \dots$ of the weak topology at the identity of $\text{Aut}^*(X)$ and a countable everywhere dense subset $\tau_1, \tau_2, \dots$ of $\text{Aut}^*(X)$. For each n choose, using Lemma 4.12, a finite measurable partition $\mathcal{W}_n$ and a subgroup G_n of $\text{Aut}^*(X, \mu)$ isomorphic to the permutation group of rank m_n and consisting of Lebesgue permutations with respect to $\mathcal{W}_n$, such that $G_n \cap V_n \tau_i \neq \emptyset$, $i = 1, 2, \dots, n$.

It is clear that as $n \rightarrow \infty$, the size of the largest element of the partition $\mathcal{W}_n$ goes to zero. According to Corollary 2.12, (G_n) form a Lévy family. This verifies the condition (1) in the definition 2.14. The condition (2) evidently follows from the choice of groups G_n . □

Remark 4.14. We do not know if the group $\text{Aut}^*(X, \mu)_u$ with the uniform topology is (extremely) amenable.

4.5. Groups of isometries of L^p . In this subsection we will prove the following result.

Theorem 4.15. *Let (X, μ) be a standard Borel space with a non-atomic measure. The group of isometries $\text{Iso}(L^p(X, \mu))$ with the strong operator topology is a generalized Lévy group (and therefore extremely amenable).*

In case $p = 2$, the result by Gromov and Milman [23] that the unitary group $U(\ell^2)$ with the strong operator topology is a Lévy group has started the current problematics. For $p \neq 2$, however, the proof is different.

A description of the isometries of $L^p(X, \mu)$ was obtained by S. Banach ([1], théorème 11.5.I, p. 178), but the proof was apparently never published. The first published proof (of a more general result) belongs to Lamperti [33], whose statement of the main result (Theorem 3.1) was not quite correct, as noticed in [5].

Let $(X, \mathcal{F}, \mu)$ be a finite measure space. Recall that a set map $\phi: \mathcal{F} \rightarrow \mathcal{F}$, defined modulo zero measure sets, is called a *regular set isomorphism* if

1. $\phi(X \setminus A) = \phi(X) \setminus \phi(A)$, $A \in \mathcal{F}$,
2. $\phi(\cup_{n=1}^{\infty} A_n) = \cup_{n=1}^{\infty} \phi(A_n)$ for disjoint $A_n \in \mathcal{F}$,

3. $\mu(\phi(A)) = 0$ if and only if $\mu(A) = 0$, $A \in \mathcal{F}$.

A regular set isomorphism ϕ determines a positive linear operator on $L^p(X, \mu)$, denoted by the same letter ϕ , by the condition $\phi(\chi_A) = \chi_{\phi(A)}$.

Here is a corrected form of the Lamperti theorem, cf. [5],[24].

Theorem 4.16. *Let $1 \leq p < \infty$, $p \neq 2$. Then every isometry, T , of $L^p(X, \mathcal{F}, \mu)$ (not necessarily onto) is of the form*

$$(4.9) \quad (Tf)(x) = h(x)(\phi(f))(x),$$

where ϕ is a regular set isomorphism and $E\{|h|^p: \phi(\mathcal{F})\} = d(\mu \circ \phi^{-1})/d\mu$, that is,

$$\int_{\phi(A)} |h|^p d\mu = \int_{\phi(A)} \frac{d(\mu \circ \phi^{-1})}{d\mu} d\mu \text{ for all } A \in \mathcal{F}.$$

□

Recall that $\text{Aut}^*(X, \mu)_w$ can be identified with a (closed) topological subgroup of the group $\text{Iso}(L^p(X, \mu))$, equipped with the strong operator topology, through the left quasi-regular representation:

$$\sigma f(x) = f(\sigma^{-1}x) \frac{d(\mu \circ \sigma^{-1})}{d\mu}, \quad \sigma \in \text{Aut}^*(X, \mu).$$

Invertible (the same as onto) isometries of $L^p(X, \mu)$ correspond to the invertible regular set isomorphisms, that is, invertible measure class preserving transformations of (X, μ) , in which case $h\phi(f) = \phi^{-1}f$. We obtain:

Corollary 4.17. *Let $1 \leq p < \infty$, $p \neq 2$, let (X, μ) be a finite measure space, and let T be a surjective isometry of $L^p(0, 1)$. Then there is an invertible measure class preserving transformation σ of X and a measurable function h with $|h| = 1$ such that*

$$(4.10) \quad Tf = h \sigma f.$$

□

Corollary 4.18. *Let $1 \leq p < \infty$, $p \neq 2$, let (X, μ) be a finite measure space. The group $\text{Iso}(L^p(X, \mu))$ is the semidirect product of the subgroup $\text{Aut}^*(X, \mu)$ and the normal subgroup $L(X, \mu; U(1))$ of all measurable maps from X to the circle rotation group. Moreover, the group $\text{Iso}(L^p(X, \mu))$ with the strong operator topology is the semidirect product of $\text{Aut}^*(X, \mu)_w$ and the group $L^0(X, \mu; U(1))$ equipped with the topology of convergence in measure.*

Proof. According to Corollary 4.17 of Theorem 4.16, every element $T \in \text{Iso}(L^p(X, \mu))$ admits a (clearly unique) decomposition of the form $T = h \circ \sigma$, where $h \in L(X, \mu; U(1))$ and $\sigma \in \text{Aut}^*(X, \mu)$. To establish the first claim, it is therefore enough to prove that $L(X, \mu; U(1))$ is normal in $\text{Iso}(L^p(X, \mu))$ or, which is the same, invariant under inner

automorphisms generated by the elements of $\text{Aut}^*(X, \mu)$. Let $h \in L(X, \mu; U(1))$ and $\sigma \in \text{Aut}^*(X, \mu)$. One has

$$\begin{aligned}
 (\sigma^{-1}h\sigma)(f)(x) &\equiv \sigma^{-1}(h \cdot {}^\sigma f)(x) \\
 &= (h \cdot {}^\sigma f)(\sigma x) \frac{d(\mu \circ \sigma)}{d\mu}(x) \\
 &= h(\sigma x) f(\sigma^{-1}\sigma x) \frac{d(\mu \circ \sigma^{-1})}{d\mu}(\sigma x) \frac{d(\mu \circ \sigma)}{d\mu}(x) \\
 &= \sigma^{-1}(h)(x) f(x),
 \end{aligned}$$

and therefore $\sigma^{-1}h\sigma = \sigma^{-1}(h) \in L(X, \mu; U(1))$.

We have already noted that $\text{Aut}^*(X, \mu)_w$ is a closed topological subgroup of $\text{Iso}(L^p(X, \mu))$. The same is true of $L^0(X, \mu; U(1))$. Indeed, every L^p -metric, $1 \leq p < \infty$, induces the topology of convergence in measure on $L(X, \mu; U(1))$, and using this observation, it is easy to verify that, first, for every $f \in L^p(X, \mu)$ the orbit map of the multiplication action

$$L^0(X, \mu; U(1)) \ni h \mapsto h \cdot f \in L^p(X, \mu)$$

is continuous, while for $f(x) \equiv 1$ it is a homeomorphic embedding.

As a corollary (which can be also checked directly), the action of $\text{Aut}^*(X, \mu)_w$ on $L^0(X, \mu; U(1))$, defined by $(\sigma, h) \mapsto \sigma^{-1}(h)$, is continuous as a map $\text{Aut}^*(X, \mu)_w \times L^0(X, \mu; U(1)) \rightarrow L^0(X, \mu; U(1))$. Therefore, the semi-direct product $\text{Aut}^*(X, \mu)_w \ltimes L^0(X, \mu; U(1))$ is a (Polish) topological group.

The algebraic automorphism given by the multiplication map,

$$\text{Aut}^*(X, \mu)_w \ltimes L^0(X, \mu; U(1)) \rightarrow \text{Iso}(L^p(X, \mu)),$$

is continuous, and since all groups involved are Polish, it is a topological isomorphism by the standard Open Mapping Theorem for Polish groups, cf. [31], Corollary 3, p. 98 and Theorem 3, p. 90. $\square$

Proof of Theorem 4.15. Choose a countable cofinal family $(\mathcal{W}_n)$ of refining partitions of X . For every n select a group $S(n)$ of Lebesgue transformations with regard to $\mathcal{W}_n$.

Let L_n denote the subgroup of $L(X, \mu; U(1))$ consisting of functions constant on elements of $\mathcal{W}_n$. From the proof of Corollary 4.18 it is clear that the group L_n is invariant under all inner automorphisms generated by elements of $S(n)$.

The normalized Haar measures on $S(n)$ concentrate in $\text{Aut}^*(X, \mu)_w$ by Corollary 2.12. Since concentration depends on the uniform structure, one can assume without loss in generality that the groups L_n are equipped with the $L^1(\mu)$ -metric induced from $L^0(X, \mu; U(1))$ (because it induces the topology of convergence in measure on the former group and is translation invariant, therefore induces the additive uniformity). By Corollary 2.10, the normalized Haar measures on L_n , $n \in \mathbb{N}$, concentrate in $L^0(X, \mu; U(1))$.

The semidirect product groups $S(n) \ltimes L_n$ are compact, and the normalized Haar measures on them are products of Haar measures, respectively, on $S(n)$ and on L_n . Therefore, those measures concentrate in $\text{Iso}(L^p(X, \mu))$ by Lemma 2.1.

Finally, condition (2) from Definition 2.14 of a generalized Lévy group is satisfied by force of Theorem 4.11 and the Weak Approximation Theorem 4.1. $\square$

Theorem 4.15 was conjectured by one of the present authors in [44].

5. FULL GROUPS OF AMENABLE EQUIVALENCE RELATIONS

5.1. The full group. Let $\mathcal{R}$ be a discrete measured equivalence relation, i.e. $\mathcal{R}$ is an equivalence relation with countable equivalence classes on a standard Borel space S , the graph $\mathcal{R} \subset S \times S$ is Borel, and μ is a quasi-invariant probability measure on S .

The full group $[\mathcal{R}]$ is defined as the group of all bimeasurable transformations σ of (S, μ) with

$$(s, \sigma(s)) \in \mathcal{R}, \quad \mu\text{-a.e.}$$

We will consider on $[\mathcal{R}]$ the uniform topology, determined by the measure μ . With this topology, $[\mathcal{R}]$ is a Polish group [30],[9].

If G is a countable discrete group and (S, μ) is a (right) free G -space, then an element σ of the *full group* $[G]$ of G is determined by a measurable partition with the property

$$S = \coprod_{g \in G} S_g = \coprod_{g \in G} S_g g,$$

in such a way that $s\sigma = sg$ if $s \in S_g$. In particular, G is a subgroup of $[G]$, every element $h \in G$ being determined by the partition

$$S_g = \begin{cases} S, & \text{if } g = h, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Remark 5.1. To the measured dynamical system (S, μ, G) is associated a von Neumann algebra, the so-called Murray–von Neumann group measure space construction. One has the following unitary representations U, V of G and isometric $*$ -representations M and N of $L^\infty(S)$ on $L^2(S \times G)$, given for $g \in G$, $f \in L^\infty(S, \mu)$, by

$$U_g \xi(x, h) = r(x, g)^{1/2} \xi(xg, hg), \quad V_g \xi(x, h) = \xi(x, g^{-1}h),$$

$$M_f \xi(x, h) = f(x) \xi(x, h), \quad N_f \xi(x, h) = f(xh^{-1}) \xi(x, h),$$

where $r(x, g) = d\mu(xg)/d\mu(x)$ is the Radon–Nikodym cocycle of the action of G on (S, μ) .

The von Neumann algebra generated by the operators V_g and N_f is denoted by L , while that generated by the operators U_g and M_f is denoted by R .

If $J \in U(L^2(S \times G))$ is given by

$$J\xi(x, g) = f(xg^{-1}, g^{-1}) r(x, g^{-1})^{1/2}, \quad (x, g) \in S \times G,$$

then $J^2 = 1$, and $JLJ = R$.

To any $\sigma \in [G]$, defined by the partitions $(S_g)_{g \in G}$ and $(S_g g)_{g \in G}$ of S , there are naturally associated unitaries $U_\sigma \in R$ and $V_\sigma \in L$ given by

$$U_\sigma = \sum_{g \in G} M_{f_g} U_g \in R \text{ and } V_\sigma = \sum_{g \in G} N_{f_g} V_g \in L,$$

where $f_g = \chi_{S_g}$.

If $U([G]) \subset U(R)$ (resp. $V([G]) \subset U(L)$) is endowed with the strong operator topology and $[G]$ with the uniform topology, it is easy to check that U (resp. V) is a topological isomorphism from $[G]$ onto its image.

5.2. In this subsection, we will show that if $\mathcal{R}$ is an amenable equivalence relation, then the full group $[\mathcal{R}]$ with the uniform topology is extremely amenable. For a definition of an amenable equivalence relation, see [52], Sect. 3.

By [8], we can assume that S is the compact abelian group $\prod\{0, 1\}$, the equivalence relation $\mathcal{R}$ is induced by the action of the subgroup $\oplus \mathbb{Z}/2$, acting by addition, and μ is a G -ergodic non-atomic measure on S .

For $n \geq 1$, let $S_n = \prod_{k=1}^n \{0, 1\}$, and for $x \in S_n$ let $C(x)$ denote the cylinder set $\{(y_k)_{k \geq 1} \in S : y_k = x_k, 1 \leq k \leq n\}$ of S . For every bijection $\sigma \in B_{ij}(S_n)$, let σ be the element of $[G]$ defined for $y = (x, y^n) \in C(x)$, $x \in S_n$, by $\sigma(y) = (\sigma(x), y^n)$, and let $B_{ij}(n)$ denote the finite subgroup of $[G]$ such transformations form.

For $n \geq 1$, if $G_n = \{g = (g_k)_{k \geq 1} \in G : g_k = 0 \ \forall k \geq n+1\}$, then $[G_n] = B_{ij}(n)$.

Lemma 5.2. *Keeping the above notation, $\cup_n B_{ij}(n)$ is uniformly dense in $[G]$, and therefore the group $[G]$ with the uniform topology is amenable.*

Proof. Let $(S_g)_{g \in G}$ and $(S_g g)_{g \in G}$ be a measurable partition of S and let $\alpha \in [G]$ be given by $\alpha(x) = xg$, $x \in S_g$. Let $0 < \varepsilon < 1$ be given and let N_1 be such that $\mu\left(\coprod_{g \in G_{N_1}} S_g\right) \geq 1 - \varepsilon$. Let us fix $0 < \delta < \frac{\varepsilon}{3 \cdot 2^{2N_1}}$.

The probability measure $\nu = \frac{1}{2^{N_1}} \sum_{g \in G_{N_1}} \mu \circ g$ is G_{N_1} -invariant and equivalent to μ .

For $n \geq 1$, let E_n denote the conditional expectation with respect to ν , i.e. for a measurable subset $B \subset S$,

$$E_n(\chi_B) = \sum_{x \in S_n} r_x^n(B) \chi_{C(x)},$$

where

$$r_x^n(B) = \frac{\nu(C(x) \cap B)}{\nu(C(x))}.$$

Let $N_2 \geq N_1$ be such that for all $B = S_g$ or $B = S_g g$, for $g \in G_{N_1}$,

$$(5.1) \quad \|E_{N_2}(\chi_B) - \chi_B\|_{1,\nu} = 2 \sum_{x \in S_{N_2}} r_x^{N_2}(B) (1 - r_x^{N_2}(B)) \nu(C(x)) < \delta^2$$

If A is a measurable subset of S , let $S_{N_2}^A$ denote $\{x \in S_{N_2} : r_x^{N_2}(A) \geq 1 - \delta\}$ and let $\tilde{A}$ denote the disjoint union $\coprod_{x \in S_{N_2}^A} C(x)$. Then the sets $(\tilde{S}_g)_{g \in G_{N_1}}$ satisfy for $g, h \in G_{N_1}$ the following properties:

- i) $\|\chi_{S_g} - \chi_{\tilde{S}_g}\|_{1,\nu} \leq 3\delta$,
- ii) $\tilde{S}_g \cap \tilde{S}_h = \emptyset$, for $g \neq h$,
- iii) $\tilde{S}_g g \cap \tilde{S}_h h = \emptyset$, for $g \neq h$.

Indeed, for $g \in G_{N_1}$ we have by definition of $\tilde{S}_g$:

$$\begin{aligned} \|E_{N_2}(\chi_{S_g}) - \chi_{\tilde{S}_g}\|_{1,\nu} &\leq \delta \sum \{\nu(C(x)) : x \in S_{N_2}, r_x^{N_2}(S_g) \leq \delta \text{ or } r_x^{N_2}(S_g) \geq 1 - \delta\} + \\ &\quad \sum \{\nu(C(x)) : x \in S_{N_2}, r_x^{N_2}(S_g)(1 - r_x^{N_2}(S_g)) > \delta(1 - \delta)\}. \end{aligned}$$

By (5.1) we have

$$\begin{aligned} \|E_{N_2}(\chi_{S_g}) - \chi_{\tilde{S}_g}\|_{1,\nu} &\leq \delta + \frac{1}{2\delta(1 - \delta)} \|E_{N_2}(\chi_{S_g}) - \chi_{S_g}\| \\ &\leq \delta + \frac{\delta}{2(1 - \delta)} \leq \frac{5\delta}{2}. \end{aligned}$$

Then,

$$\begin{aligned} \|\chi_{S_g} - \chi_{\tilde{S}_g}\|_{1,\nu} &\leq \|E_{N_2}(\chi_B) - \chi_B\|_{1,\nu} + \|E_{N_2}(\chi_{S_g}) - \chi_{\tilde{S}_g}\|_{1,\nu} \\ &\leq \delta^2 + \frac{5\delta}{2} \leq 3\delta, \end{aligned}$$

which proves (i).

For $x \in S_{N_2}$,

$$1 \geq \sum_{g \in G_{N_1}} \frac{\nu(C(x) \cap S_g)}{\nu(C(x))} = \sum_{g \in G_{N_1}} r_x^{N_2}(S_g) \geq 0.$$

Hence $S_{N_2}^{S_g} \cap S_{N_2}^{S_h} = \emptyset$ and $\tilde{S}_g \cap \tilde{S}_h = \emptyset$, which proves (ii).

For (iii), notice that for $x \in S_{N_2}$, $g \in G_{N_1}$,

$$r_x^{N_2}(S_g) = \frac{\nu(S_g \cap C(xg))}{\nu(C(xg))} = \frac{\nu((S_g g \cap C(x))g)}{\nu(C(xg))} = r_x^{N_2}(S_g g),$$

as ν is G_{N_1} -invariant. Hence $S_{N_2}^{S_g}g = S_{N_2}^{S_g g}$ and

$$\tilde{S}_g = \coprod_{x \in S_{N_2}^{S_g}} C(xg) = \coprod_{x \in S_{N_2}^{S_g g}} C(x) = \coprod_{x \in S_{N_2}^{S_g g}} C(x).$$

If $g, h \in G_{N_1}$, $g \neq h$, then $S_g g \cap S_h h = \emptyset$ and (iii) holds.

We now define $\sigma \in B_{ij}(N_2)$ such that $d_\mu(\alpha, \sigma) < \varepsilon$. As $S_{N_2}^{S_g}g = S_{N_2}^{S_g g}$, for $g \in G_{N_1}$, $A = \left\{x \in S_{N_2} : x \notin \coprod_{g \in G_{N_1}} S_{N_2}^{S_g g}\right\}$ and $B = \left\{x \in S_{N_2} : x \notin \coprod_{g \in G_{N_1}} \left(S_{N_2}^{S_g}g\right)\right\}$ have the same cardinality. Let $\tilde{\sigma}$ be an arbitrary bijection from A to B , and let σ denote the bijection of S_{N_2} defined by

$$\sigma(x) = \begin{cases} xg, & \text{if } x \in S_{N_2}^{S_g}, g \in G_{N_1}, \\ \tilde{\sigma}(x), & \text{if } x \in A. \end{cases}$$

We shall also denote by σ the corresponding element of $B_{ij}(N_2)$. Then

$$\{x \in S : \sigma(x) \neq \alpha(x)\} \subseteq \left(S \setminus \coprod_{g \in G_{N_1}} S_g\right) \coprod \left(\coprod_{g \in G_{N_1}} (S_g \setminus \tilde{S}_g)\right)$$

and

$$d_\mu(\alpha, \sigma) \leq \mu\left(S \setminus \coprod_{g \in G_{N_1}} S_g\right) + \sum_{g \in G_{N_1}} \mu(S_g \Delta \tilde{S}_g).$$

But $\mu(S_g \Delta \tilde{S}_g) \leq 2^{N_1} \nu(S_g \Delta \tilde{S}_g) = 2^{N_1} \left\| \chi_{S_g} - \chi_{\tilde{S}_g} \right\|_{1, \nu} \leq 2^{N_1+1} 3\delta$ and $d_\mu(\alpha, \sigma) \leq \varepsilon + 2^{2N_1} 3\delta \leq 2\varepsilon$. $\square$

As the size of the largest cylinder set determined by the first n coordinates clearly goes to zero as $n \rightarrow \infty$, Corollary 2.12 implies that the family of groups $B_{ij}(n)$ equipped with the uniform metric and the normalized counting measure is a Lévy family. Applying Lemma 5.2, we deduce:

Proposition 5.3. *Let $\mathcal{R}$ be a discrete measured equivalence relation, acting ergodically on a Lebesgue space (S, μ) . If $\mathcal{R}$ is an amenable equivalence relation, then the full group $[\mathcal{R}]$, endowed with the uniform topology, is a Lévy group (and in particular extremely amenable).* $\square$

Remark 5.4. As pointed out to us by A.S. Kechris, extreme amenability of the full group $[G]$ in Proposition 5.3 can be alternatively deduced by applying an argument similar to that employed by one of the present authors to give a new proof of extreme amenability of the unitary group $U(\ell^2)_s$ in [42], subsection 4.5.

Namely, for $n \geq 1$, let $S^n = \prod_{k \geq n+1} \{0, 1\}$. To any $f \in L(S^n, S_{2^n})$ associate the element $\tilde{f} \in [G]$, defined for $x = (x_k)_{k \geq 1} \in S$, by $\tilde{f}(x) = (f(x^n)(x_1, \dots, x_n), x^n)$,

where $x^n = (x_{n+1}, x_{n+1}, \dots) \in S^n$. Let L_n denote the subgroup of $[G]$ formed by $\{\tilde{f} \in [G]: f: S^n \rightarrow \mathfrak{S}_{2^n} \text{ is measurable}\}$. Let $\tilde{\mu}$ be the probability measure on S^n , given for $A \subset S^n$, A measurable, by

$$\tilde{\mu}(A) = \sum_{\bar{x} \in S_n} \mu(\{(\bar{x}, y): y \in A\}).$$

For $f \in L(S^n, \mathfrak{S}_{2^n})$ we have

$$\begin{aligned} \mu(\{x \in S: \tilde{f}(x) \neq x\}) &= \sum_{\bar{x} \in S_n} \mu(\{(\bar{x}, y): f(y)(\bar{x}) \neq \bar{x}\}) \\ &\subset \sum_{\bar{x} \in S_n} \mu(\{(\bar{x}, y): f(y) \neq \text{Id}\}) \\ &= \tilde{\mu}(\{y \in S^n: f(y) \neq \text{Id}\}). \end{aligned}$$

Therefore, if $L(S^n, \mathfrak{S}_{2^n})$ is endowed with the topology of convergence in measure with respect to $\tilde{\mu}$ and $L_n \subset [G]$ with the uniform topology, then the group isomorphism $L(S^n, \mathfrak{S}_{2^n}) \ni f \mapsto \tilde{f} \in L_n$ is continuous.

By Theorem 2.18, $(L_n)_{n \geq 1}$ forms a sequence of extremely amenable subgroups of $[G]$, with $[G_k] \subset L_n$, $\forall n \geq 1$. As $([G_n])_{n \geq 1}$ forms an increasing sequence of subgroups of $[G]$, whose union is uniformly dense, we conclude by Lemma 2.20 that if G and X are as in Prop. 5.3, then the full group $[G]$, endowed with the uniform topology, is extremely amenable.

Notice, however, that the actual conclusion of our Proposition 5.3 is stronger, because there exist extremely amenable groups that are not Lévy groups, cf. e.g. Remark 2.19.

5.3. In this subsection, we show that if $[\mathcal{R}]$ endowed with the uniform topology is amenable as a topological group, then $\mathcal{R}$ is an amenable equivalence relation.

If E is a separable Banach space, let $\text{Iso}(E)$ denote the group of isometric isomorphisms of E , endowed with the Borel structure induced by the strong operator topology. As is well-known (see e.g. [53], Lemma 1.3), the map from $\text{Iso}(E)$ to $\text{Homeo}(E_1^*)$, given by duality, is continuous, hence Borel, if we consider the topology of uniform convergence on $\text{Homeo}(E_1^*)$.

Let $\alpha: \mathcal{R} \rightarrow \text{Iso}(E)$ be a cocycle, that is, a measurable map such that

$$\alpha(x, y)\alpha(y, z) = \alpha(x, z) \quad \text{if } (x, y), (y, z) \in \mathcal{R}.$$

Let $\alpha^*: \mathcal{R} \rightarrow \text{Homeo}(E_1^*)$ denote the cocycle given by

$$\alpha^*(x, y) = (\alpha(x, y)^{-1})^* = \alpha(y, x)^*, \quad (x, y) \in \mathcal{R}.$$

For $f \in L^1(S, E)$ and $\sigma \in [\mathcal{R}]$, let $T(\sigma)f$ denote the element of $L^1(S, E)$ given by

$$T(\sigma)f(s) = r(s, \sigma)\alpha(s, s\sigma)f(s\sigma), \quad s \in S,$$

where $r(s, \sigma)$ denotes the Radon-Nikodym cocycle given for $s \in S$ and $\sigma \in [\mathcal{R}]$ by $d\mu(s\sigma) = r(s, \sigma)d\mu(s)$.

One checks easily that T defines a representation of $[\mathcal{R}]$ into $\text{Iso}(L^1(S, E))$.

Lemma 5.5. *If $[\mathcal{R}]$ is endowed with the uniform topology, then T is strongly continuous.*

Proof. For $\sigma \in [\mathcal{R}]$ one has

$$\begin{aligned} \|T(\sigma)\tilde{f} - \tilde{f}\| &= \int_S \|r(s, \sigma)\alpha(s, s\sigma)\tilde{f}(s\sigma) - \tilde{f}(s)\| d\mu(s) \\ &= \int_{\text{supp}(\sigma)} \|r(s, \sigma)f(s\sigma)\alpha(s, s\sigma)\xi - f(s)\xi\| d\mu(s) \\ &\leq \|\xi\| \int_{\text{supp}(\sigma)} (r(s, \sigma)|f(s\sigma)| + |f(s)|) d\mu(s) \\ &\leq 2\|\xi\| \|f\|_\infty \mu(\text{supp}(\sigma)) = 2\|\xi\| \|f\|_\infty d_\mu(\sigma, \text{Id}), \end{aligned}$$

where $\text{supp}(\sigma) = \{s \in S : s\sigma \neq s\}$. □

5.4. Let $\mathfrak{A} = \{A_s : s \in S\}$ be a Borel field of (non-empty) compact convex subsets of E^* such that for all $(x, y) \in \mathcal{R}$,

$$\alpha^*(x, y)(A_y) = A_x \quad \mu\text{-a.e.}$$

By [53], Prop. 2.2,

$$A = \{\lambda \in L^\infty(S, E^*) : \lambda(s) \in A_s \quad \mu\text{-a.e.}\}$$

is a weak* compact convex non-empty subset of $L^\infty(S, E^*)_1$.

Moreover, if T^* denotes the dual representation of T in $L^\infty(S, E^*)$, given for $\sigma \in [\mathcal{R}]$ and $\lambda \in L^\infty(S, E^*)$ by

$$T^*(\sigma)\lambda(s) = \alpha^*(s, s\sigma)f(s\sigma), \quad s \in S,$$

then A is invariant under the (affine) action T^* of $[\mathcal{R}]$.

We have already noted that this action is continuous if A is equipped with the weak* topology. Therefore, if the full group $[\mathcal{R}]$, equipped with the uniform topology, is amenable as a topological group, then there exists a $[\mathcal{R}]$ -invariant section of the Borel field $\mathfrak{A}$. In particular, this section is $\mathcal{R}$ -invariant. Therefore, we have proved the following proposition.

Proposition 5.6. *Let $\mathcal{R}$ be a discrete measured equivalence relation, acting ergodically on a Lebesgue space (S, μ) . If the full group $[\mathcal{R}]$, equipped with the uniform topology, is amenable as a topological group, then the equivalence relation $\mathcal{R}$ is amenable.*

We summarize the results of this section in the following theorem.

Theorem 5.7. *Let $\mathcal{R}$ be a discrete measured equivalence relation, acting ergodically on a Lebesgue space (S, μ) . Then the following are equivalent.*

1. $\mathcal{R}$ is an amenable equivalence relation.
2. $[\mathcal{R}]$ with the uniform topology is amenable.

- 3. $[\mathcal{R}]$ with the uniform topology is extremely amenable.
- 4. $[\mathcal{R}]$ with the uniform topology is a Lévy group.

Proof. The implications (4) $\implies$ (3) $\implies$ (2) are trivial, (2) $\implies$ (1) is Proposition 5.6, and (1) $\implies$ (4) is Proposition 5.3. $\square$

Remark 5.8. Let G be a discrete countable group acting freely and ergodically on a Lebesgue space (S, μ) . Let $N[G]$ denote the normalizer of the full group. Any element of $N[G]$ (resp. $[G]$) induces an automorphism (resp. an inner automorphism) of the von Neumann factor R , normalizing $M(L^\infty(S, \mu))$ (we keep the notation of Subs. 5.1). Then the u -topology (see Section 3.3) on $N[G]$ coincides with the normal topology defined in Section 3.1 of [9] and is coarser than the uniform topology. By Proposition 5.3, if (S, μ) is an amenable G -space, then $[G]$, endowed with the u -topology, is a Lévy group and in particular extremely amenable.

6. CONCENTRATION TO A NONTRIVIAL SPACE

M. Gromov [22], section 3 $\frac{1}{2}$.45 (p. 200) has defined a metric on the space of equivalence classes of Polish mm -spaces in such a way that a sequence $X_n = (X_n, d_n, \mu_n)$ of mm -spaces forms a Lévy family if and only if it converges to the trivial mm -space $\{*\}$.

This approach allows one to talk of *concentration to a nontrivial mm -space*. According to Gromov, this type of concentration commonly occurs in statistical physics. At the same time, there are very few known non-trivial examples of this kind in the context of transformation groups. The aim of this Section is to give some natural examples of concentration of this type.

We will recall Gromov's construction. Let $\mu^{(1)}$ denote the Lebesgue measure on the unit interval $\mathbb{I} = [0, 1]$. On the space $L(0, 1)$ of all (equivalence classes) of measurable real-valued functions define the metric me_1 , generating the topology of convergence in measure, as follows: $\text{me}_1(h_1, h_2)$ is the infimum of all $\lambda > 0$ with the property

$$\mu^{(1)}\{|h_1(x) - h_2(x)| > \lambda\} < \lambda.$$

Let $X = (X, d_X, \mu_X)$ and $Y = (Y, d_Y, \mu_Y)$ be two Polish mm -spaces. There exist measurable maps $f: \mathbb{I} \rightarrow X$, $g: \mathbb{I} \rightarrow Y$ such that $\mu_X = f_* \mu^{(1)}$ and $\mu_Y = g_* \mu^{(1)}$. Denote by L_f the set of all functions of the form $h = h_1 \circ f$, where $h_1: X \rightarrow \mathbb{R}$ is 1-Lipschitz, having the additional property $h(0) = 0$. Similarly, define the set L_g . Both L_f and L_g are compact subsets of $L^0(0, 1)$.

Let $\underline{H}_1 \mathcal{L} \iota(X, Y)$ be the infimum of Hausdorff distances between L_f and L_g (with regard to the metric me_1), taken over all parametrizations f and g as above. One can prove that $\underline{H}_1 \mathcal{L} \iota$ is a metric on the space of (isomorphism classes of) all Polish mm -spaces. (The difficult part of the proof is verifying the first axiom of a metric.)

A sequence of mm -spaces $X_n = (X_n, d_n, \mu_n)$ forms a Lévy family if and only if it converges to the trivial (one-point) mm -space in the metric $\underline{H}_1 \mathcal{L} \iota$:

$$X_n \xrightarrow{\underline{H}_1 \mathcal{L} \iota} \{*\}.$$

(This is a reformulation of Lemma 2.3.)

Theorem 6.1 (Gromov). *Let $Y = (Y, d_Y, \nu)$ and $Z_n = (Z_n, d_n, \mu_n)$, $n \in \mathbb{N}$ be mm-spaces, where (Z_n) forms a Lévy family. Let $d_Y \oplus^{(2)} d_n$ denote the ℓ_2 -type sum of the metrics d_Y and d_n on the product space $Y \times Z_n$. Then the family of mm-spaces $(Y \times Z_n, d_Y \oplus^{(2)} d_n, \nu \otimes \mu_n)$ concentrates to the mm-space (Y, d_Y, ν) :*

$$(Y \times Z_n, d_Y \oplus^{(2)} d_n, \nu \otimes \mu_n) \xrightarrow{H_1 \mathcal{L}_t} (Y, d_Y, \nu).$$

□

We are going to show a situation where Gromov's theorem applies naturally.

Lemma 6.2. *Let G be a metrizable topological group which is, as a topological group, a semidirect product of a compact subgroup H and a closed normal subgroup N :*

$$G = H \ltimes N.$$

Assume that N is a Lévy group, that is, there is a directed family of compact subgroups (K_α) of N , having an everywhere dense union and such that (K_α, μ_α) form a Lévy family with regard to the right uniform structure on G . Let ν denote the normalized Haar measure on H . Then there exist a right-invariant compatible metric ρ on G such that the family of mm-spaces $H \cdot K_\alpha$, equipped with the metric ρ and the convolution of normalized Haar measures, concentrates to the mm-space $(H, \rho|_H, \nu)$:

$$(H \cdot K_\alpha, \rho, \nu * \mu_\alpha) \xrightarrow{H_1 \mathcal{L}_t} (H, \rho|_H, \nu).$$

If in addition each of the subgroups (K_α) is invariant under the action of H by conjugations, then every $H \cdot K_\alpha$ is a compact subgroup of G .

Proof. Denote by τ the continuous action of H on N by topological group isomorphisms, so that $G = H \ltimes_\tau N$. Fix any compatible right-invariant metric, d , on N , and define a new metric, ς , as follows: for $x, y \in N$, set

$$\varsigma_N(x, y) = \int_H d(\tau_h x, \tau_h y) \, d\nu(h).$$

This ς_N is also a right-invariant compatible metric on N , which in addition is invariant under the action τ .

Now choose a bi-invariant compatible metric, ς_H , on the compact group H . Define a metric ρ on G as the ℓ_2 -type sum of ς_H and ς_N :

$$\rho((h, x), (h', x')) = \sqrt{\varsigma_H(h, h')^2 + \varsigma_N(x, x')^2}.$$

It is easily verified that ρ is a right-invariant compatible metric on G . Besides, ρ is invariant under left multiplication by elements of H :

$$\begin{aligned}
\rho((h, e) \cdot (g, x), (h, e) \cdot (g', x')) &= \rho((hg, \tau_h x), (hg', \tau_h x')) \\
&= (\varsigma_H(hg, hg')^2 + \varsigma_V(\tau_h x, \tau_h x')^2)^{\frac{1}{2}} \\
&= (\varsigma_H(g, g')^2 + \varsigma_V(x, x')^2)^{\frac{1}{2}} \\
&= \rho((g, x), (g', x')).
\end{aligned}$$

The restriction of ρ to each $H \cdot K_\alpha$ is just the ℓ_2 -type sum of ς_H with the restriction of ς_N to K_α .

Since the multiplication mapping $H \times K_\alpha \rightarrow H \cdot K_\alpha$ is one-to-one (by assumption) and also of course continuous, it is a homeomorphic embedding of $H \times K_\alpha$ into our topological group G . In particular, the convolution of measures $\nu * \mu_\alpha$ is in fact the product measure, $\nu \otimes \mu_\alpha$. Now Gromov's theorem 6.1 applies.

The last statement of the Lemma is obvious. $\square$

Remark 6.3. As a direct consequence from [44], Lemma 2.3, under assumptions of Lemma 6.2, H forms the universal minimal flow of the topological group G . It would be interesting to find more general situations where concentration of subobjects of G to a nontrivial compact space X would mean that X is a universal minimal flow of G .

Using the description of the automorphism groups of injective factors of type *III*, we now present concrete examples of concentration to a non-trivial space. In particular, in the last Example 6.6 we get a sequence of compact subgroups concentrating to a non-trivial space.

Example 6.4. For $0 < \lambda < 1$, let us consider $G = \oplus(\mathbb{Z}/2)$ acting freely and ergodically on $(X, \mu_\lambda) = \prod(\{0, 1\}, \{\frac{1}{1+\lambda}, \frac{\lambda}{1+\lambda}\})$, and let R_λ denote the von Neumann factor of type *III* $_\lambda$, obtained by the group measure space construction. Recall that R_λ is approximately finite dimensional and by [7] is up to isomorphism the unique injective factor of type *III* $_\lambda$. To any automorphism α of R_λ is associated its modulus, $\text{mod } \alpha$, which belongs to $\mathbb{R}/(\log \lambda)\mathbb{Z}$ (see for example [47], Chap. XII). If $\text{Aut}(R_\lambda)$ is endowed with the u -topology, then the mod map $\text{mod} : \text{Aut}(R_\lambda) \rightarrow \mathbb{R}/(\log \lambda)\mathbb{Z}$ is a continuous homomorphism. Moreover (see [47], Thm. XVIII.1.11) there is a natural short exact sequence

$$1 \rightarrow \overline{\text{Inn}(R_\lambda)} \rightarrow \text{Aut}(R_\lambda) \xrightarrow{\text{mod}} \mathbb{R}/(\log \lambda)\mathbb{Z} \rightarrow 1,$$

and a continuous action $\tilde{\theta} : \mathbb{R}/(\log \lambda)\mathbb{Z} \rightarrow \text{Aut}(R_\lambda)$ with

$$\text{mod } (\tilde{\theta}(s)) = s, \quad s \in \mathbb{R}/(\log \lambda)\mathbb{Z}.$$

By Proposition 3.10, $\text{Inn}(R_\lambda)$ and consequently $\overline{\text{Inn}(R_\lambda)}$ are Lévy groups. Hence by Lemma 6.2 the group $\text{Aut}(R_\lambda)$ contains a chain of compact mm -spaces with everywhere dense union which concentrates to $\mathbb{R}/(\log \lambda)\mathbb{Z}$.

Example 6.5. Let G be a countable discrete group acting freely and ergodically on the Lebesgue space (X, μ) . By Proposition 5.3, if (X, μ) is an amenable G -space, then the full group $[G]$, endowed with the u -topology, is extremely amenable.

If the dynamical system (X, μ, G) is of type III_λ , $0 < \lambda < 1$, then there is a split exact sequence

$$1 \rightarrow \overline{[G]}^{u\text{-top}} \rightarrow N[G] \xrightarrow{\text{mod}} \mathbb{R}/(\log \lambda)\mathbb{Z} \rightarrow 1$$

(See [9], 3.3.) As $\overline{[G]}^{u\text{-top}}$ is extremely amenable and moreover a Lévy group, $N[G]$ contains a chain of compact mm -spaces with everywhere dense union which concentrates to $\mathbb{R}/(\log \lambda)\mathbb{Z}$.

Example 6.6. Let R_λ be the injective factor of type III_λ (see Example 6.4), realized as an infinite tensor product of factors of type I_2 , i.e. $R_\lambda = \otimes (M_2(\mathbb{C}), \varphi_\lambda)$, where $\varphi_\lambda = \text{Tr}(h_\lambda, \cdot)$ and h_λ is the density matrix $\begin{pmatrix} \frac{1}{1+\lambda} & \\ & \frac{\lambda}{1+\lambda} \end{pmatrix}$. If φ denotes the state $\otimes \varphi_\lambda$, then the modular group of automorphisms σ^φ is given by

$$\sigma_t^\varphi = \otimes \text{Ad} \begin{pmatrix} 1 & \\ & \lambda^{it} \end{pmatrix}, \quad t \in \mathbb{R}.$$

For $m \geq 1$, let N_m be the type I_{2^m} subfactor of M generated by

$$\{x_1 \otimes x_2 \otimes \dots \otimes x_m \otimes 1 \otimes 1 \otimes \dots : x_k \in M_2(\mathbb{C})\}.$$

Let K_m denote the compact subgroup of $\text{Inn}(R_\alpha)$ given by $\{\text{Ad}(u) : u \in U(N_m)\}$. Let $\text{Cnt}(R_\lambda) \triangleleft \text{Aut}(R_\lambda)$ be the normal subgroup of centrally trivial automorphisms of R_λ . Let $T = \frac{-2\pi}{\log \lambda}$. By [47], Thm. XVIII.1.11, $\text{Cnt}(R_\lambda) = \text{Inn}(R_\lambda) \rtimes_{\sigma^\varphi} \mathbb{R}/T\mathbb{Z}$. By Propositions 3.6 and 3.10, $\{K_m\}_{m \geq 1}$ forms a Lévy family of $\text{Inn}(R_\lambda)$. As for $t \in \mathbb{R}$, σ_t^φ leaves N_m globally fixed,

$$K_m \cdot \mathbb{R}/T\mathbb{Z} = \{\text{Ad } u \circ \sigma_s^\varphi : u \in U(N_m), s \in \mathbb{R}\}$$

is a compact subgroup of $\text{Cnt}(R_\lambda)$. Therefore, $\text{Cnt}(R_\lambda)$ contains a chain of compact subgroups with everywhere dense union which concentrates to $\mathbb{R}/T\mathbb{Z}$.

Notice that the compact space $\mathbb{R}/T\mathbb{Z}$ forms the universal minimal flow of the topological group $\text{Cnt}(R_\lambda)$, cf. Remark 6.3.

REFERENCES

- [1] S. Banach, *Théorie des opérations linéaires*, Second edition, Chelsea Publishing Co., New York. (Reprint of the first edition, 1932, with corrections and an addendum.)
- [2] M.E.B. Bekka, *Amenable unitary representations of locally compact groups*, Invent. Math. **100** (1990), 383–401.

- [3] M.B. Bekka, *On uniqueness of invariant means*, Proc. Amer. Math. Soc. **126** (1998), 507–514.
- [4] M.E.B. Bekka and A. Valette, *Kazhdan's property (T) and amenable representations*, Math. Z. **212** (1993), 293–299.
- [5] S.L. Campbell, G.D. Faulkner, and M.L. Gardner, *Isometries on L^p spaces and copies of l^p shifts*, Proc. Amer. Math. Soc. **77** (1979), 198–200.
- [6] J.R. Choksi and S. Kakutani, *Residuality of ergodic measurable transformations and of ergodic transformations which preserve an infinite measure*, Indiana Univ. Math. J. **28** (1979), 453–469.
- [7] A. Connes, *Classification of injective factors*, Ann. of Math. **104** (1976), 73–115.
- [8] A. Connes, J. Feldman, and B. Weiss, *An amenable equivalence relation is generated by a single transformation*, Ergodic Theory Dynamical Systems **1** (1981), 431–450.
- [9] A. Danilenko, *The topological structure of Polish groups and groupoids of measure space transformations*, Publ. Res. Inst. Math. Sci. **31** (1995), 913–940.
- [10] J. Dixmier, *Les C^* -algèbres et leurs représentations*, Gauthier-Villars, Paris, 1964.
- [11] G.A. Elliott, *On approximately finite-dimensional von Neumann algebras. II*, Canad. Math. Bull. **21** (1978), 415–418.
- [12] R. Engelking, *General Topology*, Math. Monographs, **60**, PWN - Polish Scient. Publishers, Warsaw, 1977.
- [13] T. Giordano and V. Pestov, *Some extremely amenable groups*. – C.R. Acad. Sci. Paris, Sér. I **334** (2002), No. 4, 273–278.
- [14] S. Glasner, *Proximal Flows*, Lecture Notes Math. **517**, Springer-Verlag, Berlin-New York, 1976.
- [15] S. Glasner, *On minimal actions of Polish groups*, Top. Appl. **85** (1998), 119–125.
- [16] E. Glasner and B. Weiss, *Minimal actions of the group $S(\mathbb{Z})$ of permutations of the integers*, Geom. and Funct. Anal. **12** (2002), 964–988.
- [17] E. Glasner and B. Weiss, *The universal minimal system for the group of homeomorphisms of the Cantor set*, Fund. Math. **176** (2003), no. 3, 277–289.
- [18] E. Glasner, B. Tsirelson, and B. Weiss, *The automorphism group of the Gaussian measure cannot act pointwise*, ArXiv e-print math.DS/0311450, v. 2 (March 2004), 18 pp. To appear in Israel J. Math.
- [19] E. Granirer, *Extremely amenable semigroups I,II*, Math. Scand. **17** (1965), 177–179; **20** (1967), 93–113.
- [20] E. Granirer and A.T. Lau, *Invariant means on locally compact groups*, Ill. J. Math. **15** (1971), 249–257.
- [21] M. Gromov, *Filling Riemannian manifolds*, J. Diff. Geom. **18** (1983), 1–147.
- [22] M. Gromov, *Metric Structures for Riemannian and Non-Riemannian Spaces*, Progress in Mathematics **152**, Birkhauser Verlag, 1999.
- [23] M. Gromov and V.D. Milman, *A topological application of the isoperimetric inequality*, Amer. J. Math. **105** (1983), 843–854.
- [24] R. Grzaślewicz, *Isometries of $L^1 \cap L^p$* , Proc. Amer. Math. Soc. **93** (1985), 493–496.
- [25] U. Haagerup, *All nuclear C^* -algebras are amenable*, Invent. Math. **74** (1983), 305–319.
- [26] P.R. Halmos, *Lectures on Ergodic Theory*, Chelsea Publ. Co., New York, 1960.
- [27] P. de la Harpe, *Moyennabilité de quelques groupes topologiques de dimension infinie*, C.R. Acad. Sci. Paris, Sér. A **277** (1973), 1037–1040.
- [28] P. de la Harpe, *Moyennabilité du groupe unitaire et propriété P de Schwartz des algèbres de von Neumann*, in: Algèbres d'opérateurs (Sém., Les Plaines-sur-Bex, 1978), pp. 220–227, Lecture Notes in Math. **725**, Springer-Verlag, Berlin, 1979.
- [29] W. Herer and J.P.R. Christensen, *On the existence of pathological submeasures and the construction of exotic topological groups*, Math. Ann. **213** (1975), 203–210.
- [30] T. Hamachi and M. Osikawa, *Ergodic groups of automorphisms and Krieger's theorems*, Seminar on Mathematical Sciences, **3**, Keio University, Dept. Math., Yokohama, 1981. 113 pp.

- [31] T. Husain, *Introduction to Topological Groups*, W. B. Saunders Co., Philadelphia, Pa.-London, 1966,
- [32] A.S. Kechris, V.G. Pestov and S. Todorcevic, *Fraïssé limits, Ramsey theory, and topological dynamics of automorphism groups*. – ArXiv e-print math.LO/0305241, May 2003, 68 pp. To appear in Geometric and Functional Analysis.
- [33] J. Lamperti, *On the isometries of certain function-spaces*, Pacific J. Math. **8** (1958), 459–466.
- [34] M. Ledoux, *The concentration of measure phenomenon*. Math. Surveys and Monographs, **89**, Amer. Math. Soc., 2001.
- [35] B. Maurey, *Constructions de suites symétriques*, C.R. Acad. Sc. Paris, Sér. **A-B** **288** (1979), 679–681.
- [36] V.D. Milman, The heritage of P. Lévy in geometrical functional analysis, *Astérisque* **157–158** (1988), 273–301.
- [37] V.D. Milman and G. Schechtman, *Asymptotic theory of finite-dimensional normed spaces (with an Appendix by M. Gromov)*, Lecture Notes in Math., **1200**, Springer, 1986.
- [38] T. Mitchell, Fixed points and multiplicative left invariant means, *Trans. Am. Math. Soc.* **122** (1966), 195–202.
- [39] A.L.T. Paterson, *Nuclear C^* -algebras have amenable unitary groups*, Proc. Amer. Math. Soc. **114** (1992), 719–721.
- [40] V.G. Pestov, *On free actions, minimal flows, and a problem by Ellis*, Trans. Amer. Math. Soc. **350** (1998), 4149–4165.
- [41] V.G. Pestov, *Amenable representations and dynamics of the unit sphere in an infinite-dimensional Hilbert space*, Geometric and Functional Analysis **10** (2000), 1171–1201.
- [42] V. Pestov, *Ramsey–Milman phenomenon, Urysohn metric spaces, and extremely amenable groups*. – Israel Journal of Mathematics **127** (2002), 317–358.
- [43] V. Pestov, *mm-Spaces and group actions*. – L’Enseignement Mathématique **48** (2002), 209–236.
- [44] V. Pestov, *Remarks on actions on compacta by some infinite-dimensional groups*, *Infinite dimensional Lie groups in geometry and representation theory, (Washington, DC 2000)*, World Sci., 145–163, 2002.
- [45] W. Roelcke and S. Dierolf, *Uniform Structures on Topological Groups and Their Quotients*, McGraw-Hill, 1981.
- [46] M. Takesaki, *Theory of Operator Algebras. I*, Springer-Verlag, New York-Heidelberg, 1979.
- [47] M. Takesaki, *Theory of Operator Algebras. III*, Springer-Verlag, Berlin, 2003.
- [48] M. Talagrand, *Concentration of measure and isoperimetric inequalities in product spaces*, Publ. Math. IHES **81** (1995), 73–205.
- [49] A.I. Tulcea, *On the category of certain classes of transformations in ergodic theory*, Trans. Amer. Math. Soc. **114** (1965), 261–279.
- [50] V. Uspenskij, *On universal minimal compact G -spaces*, *Topology Proc.* **25** (2000), 301–308.
- [51] W.A. Veech, *Topological dynamics*, Bull. Amer. Math. Soc. **83** (1977), 775–830.
- [52] R.J. Zimmer, *Hyperfiniteness factors and amenable ergodic actions*, Invent. Math. **41** (1977), 23–31.
- [53] R.J. Zimmer, *Amenable ergodic group actions and an application to Poisson boundaries of random walks*, J. Functional Analysis **27** (1978), 350–372.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, 585 KING EDWARD AVE., OTTAWA, ONTARIO, CANADA K1N 6N5.

E-mail address: giordano@uottawa.ca, vpest283@uottawa.ca