

**ALGEBRAS ASSOCIATED TO PSEUDO-ROOTS  
OF NONCOMMUTATIVE POLYNOMIALS  
ARE KOSZUL**

DMITRI PIONTKOVSKI

ABSTRACT. Quadratic algebras associated to pseudo-roots of noncommutative polynomials have been introduced by I. Gelfand, Retakh, and Wilson in connection with studying the decompositions of noncommutative polynomials. Later they (with S. Gelfand and Serconek) shown that the Hilbert series of these algebras and their quadratic duals satisfy the necessary condition for Koszulity. It is proved in this note that these algebras are Koszul.

1. INTRODUCTION

In this note we investigate the quadratic algebras  $Q_n$  associated with pseudo-roots of noncommutative polynomials introduced in [GRW]. This is a homogeneous quadratic algebra over a commutative field  $k$  (the results do not depend on the choice of it) with  $2^n - 1$  generators ( $n \geq 2$ ). It is shown in [GRW, GGRSW] how this algebra encodes an information about factorizations of noncommutative polynomials of degree  $n$  and about noncommutative symmetric functions.

Our purpose is to show that  $Q_n$  is Koszul. It was partially done in [GGRSW]. Namely, the Hilbert series of  $Q_n$  and of its quadratic dual algebra  $Q_n^!$  have been calculated, and it occurs that these formal power series satisfy the Froberg relation  $Q_n(z)Q_n^!(-z) = 1$ , like Hilbert series of dual Koszul algebras. However, this equality do not imply the Koszulity [Po, Ro]. Moreover, for general quadratic algebra  $A$  it may happens that there is a Koszul quadratic algebra  $S$  such that  $A(z) = S(z)$  and  $A^!(z) = S^!(z)$  while  $A$  is not Koszul [Pi].

Fortunately, for some particular Koszul algebras  $S$  the reverse implication is true. Assume that  $A$  is filtered by a  $\mathbb{N}$ -graded ordered semigroup  $\Gamma$  such that the filtration on the graded component  $A_n$  is induced by the filtration on the space of generators  $V = A_1$ . Then  $A$  has also an  $\mathbb{N}$ -filtration induced by the grading of  $\Gamma$ . Then the associated graded algebra  $S = \mathbf{gr} A$  has the same Hilbert series as  $A$ , and  $A$  is Koszul provided that  $S$  is (see [PP]; idea of the proof: the induced filtration on the bar-complex leads to a spectral sequence  $E_1 = \text{Ext}_S(k, k) \implies \text{Ext}_A(k, k)$ ). We use this fact to establish

**Theorem 1.1.** *The algebra  $Q_n$  is Koszul for every  $n \geq 2$ .*

Since the Hilbert function of its dual algebra  $Q_n^!$  is a polynomial of degree  $n$  (see [GGRSW]), we have

**Corollary 1.2.** *The algebra  $Q_n$  has global dimension  $n$ .*

---

2000 *Mathematics Subject Classification.* 16S37 (16W30).

*Key words and phrases.* Koszul algebra, noncommutative polynomial.

Partially supported by the grant 02-01-00468 of the Russian Basic Research Foundation.

## 2. KOSZULITY

Recall that a graded connected  $k$ -algebra  $A$  is called Koszul if the trivial module  $k_A$  admits a linear free minimal resolution, that is, every graded vector space  $\mathrm{Tor}_i^A(k, k) \simeq \mathrm{Ext}_A^i(k, k)$  is concentrated in degree  $i$ . The standard references on Koszulity are [Pr, B, L]; a survey could be found in [PP].

**2.1. Associated graded algebra  $\mathbf{gr} Q_n$ .** There are several ways to choose the generators of  $Q_n$  [GRW, GGRSW]; let us fix one choice. Following [GGRSW], the algebra  $Q_n$  is generated by the variables  $r_A$  where  $A$  runs all nonempty subsets  $A \subset I_n = \{1, \dots, n\}$ . These generators are connected by the relations

$$f_{A;i,j} := r_A(r_{A \setminus i} - r_{A \setminus j}) + (r_{A \setminus i} - r_{A \setminus j})r_{A \setminus \{i,j\}} - r_{A \setminus i}^2 + r_{A \setminus j}^2 = 0$$

for all  $i, j \in A \subset I_n$  (where we simply write  $A \setminus i$  instead of  $A \setminus \{i\}$  etc). These relations are not linearly independent, because, e. g.,  $f_{A;i,j} = -f_{A;j,i}$ .

Recall [GGRSW] that  $Q_n$  is filtered by the ordered semigroup  $\Gamma$  of monomials on the generators (the order is degree-lexicographical with  $r_A < r_B$  iff either  $|A| < |B|$  or, if  $|A| = |B| = k$  for some  $k$ , then  $A = \{a_1, \dots, a_k\}$  and  $B = \{b_1, \dots, b_k\}$  with  $a_1 < \dots < a_k, b_1 < \dots < b_k$  and for some  $t < k$  we have  $a_i = b_i$  for all  $i < t$  but  $a_t > b_t$ ). The additional  $\mathbb{N}$ -grading on  $\Gamma$  is induced by the grading  $\deg r_A = |A|$  on generators. It induces a  $\mathbb{N}$ -filtration  $\mathcal{N}$  on  $Q_n$ . By [GGRSW], the associated graded algebra  $\mathbf{gr} Q_n$  has the same generators  $r_A, A \subset I_n$ , and relations

$$h_{A;i,j} := r_A(r_{A \setminus i} - r_{A \setminus j}) = 0.$$

Our next purpose is to prove

**Theorem 2.1.** *Algebra  $\mathbf{gr} Q_n$  is Koszul for every  $n \geq 2$ .*

By the consideration above, we have

**Corollary 2.2** (Theorem 1.1). *Algebra  $Q_n$  is Koszul for every  $n \geq 2$ .*

**2.2. Groebner basis of  $\mathbf{gr} Q_n$ .** To be familiar with canonical forms of elements in  $\mathbf{gr} Q_n$ , we first describe the Groebner basis of its relations. It was essentially done in [GGRSW], but in another language. Recall [U] that a set  $f = \{f_i\}$  of elements of a free algebra  $\mathcal{F} = k\langle x_1, \dots, x_n \rangle$  with Noetherian order “ $<$ ” on monomials is called a *Groebner basis* of an ideal  $I \subset \mathcal{F}$  (= Groebner basis of the factor algebra  $\mathcal{F}/I$ ) iff  $f \subset I$  and for every element  $a \in I$  its leading monomial  $\hat{a}$  has a subword equal to a leading monomial  $\hat{f}_i$  of some  $f_i$ . A Groebner basis is called *reduced*, if every monomial of the decomposition of  $f_i$  has no subword equal to  $\hat{f}_j$  for  $j \neq i$ , and the coefficients of  $\hat{f}_i$  in  $f_i$  are unit.

For every nonempty  $A \subset I_n$  and every  $0 \leq t \leq |A|$ , let us define a monomial  $S_A^t$  in the free algebra  $F = k\langle r_B | \emptyset \neq B \subset I_n \rangle$  as follows : if  $A = \{a_1, \dots, a_m\}$  with  $a_1 < \dots < a_m$ , then  $S_A^t = r_A r_{A \setminus \{a_1\}} \dots r_{A \setminus \{a_1, \dots, a_t\}}$ .

**Proposition 2.3.** *The reduced Groebner basis of  $\mathbf{gr} Q_n$  consists of all the elements  $g_{A,B}^t = S_A^t r_{A \setminus B} - S_A^{t+1}$  for  $t \geq 0$ , where  $|B| = t + 1, a_1 \notin B \subset A$ .*

It follows that both algebras  $\mathbf{gr} Q_n$  and  $Q_n$  has the same linear basis consisting of monomials (*normal monomials* w. r. t. the Groebner basis above)

$$S_{A^1}^{j_1} \dots S_{A^m}^{j_m},$$

where for every  $1 \leq i < m$  either  $A^{i+1} \not\subset A^i$  or  $|A^{i+1}| \neq |A^i| - j_i - 1$ . Notice that similar bases has been constructed in [GGRSW].

*Proof.* Let  $I$  be an ideal in  $F$  generated by the relations  $h_{A;i,j}$  of  $\mathbf{gr} Q_n$ . Let us show that all the elements  $g_{A,B}^t$  lie in this ideal.

First, every element  $g_{A,B}^t$  with  $a_1 \in B, t \geq 2$  lies in the ideal  $I^t$  generated by  $\{g_{A,B}^j | j < t, A, B \text{ as above}\}$ , because  $g_{A,B}^t = r_A g_{A \setminus a_1, B \setminus a_1}^{t-1}$ . By induction, assume that  $I^t \subset I$ . Let  $b \in B$ , let  $B' = B \setminus \{b\}$ , and let  $A' = A \setminus \{a_1\}$ . If  $a_1 \notin B$ , we have

$$g_{A,B}^t = r_A g_{A',B'}^{t-1} + g_{A,B'}^{t-1} (r_{A' \setminus B'} - r_{A \setminus B}) + S_A^{t-1} h_{A \setminus B'; b, a_1} \in I.$$

Obviously, the basis  $g = \{g_{A,B}^t\}$  is reduced. Since  $h_{A;i,j} = g_{A,\{i\}}^0 - g_{A,\{j\}}^0$ , it generates  $I$ . By Diamond lemma (see [U]), it is sufficient to show that every  $s$ -polynomial of  $g$  is reducible to zero. Since  $\widehat{g_{A,B}^t} = S_A^t r_{A \setminus B}$ , every such  $s$ -polynomial has the form

$$s_{A,B,C,p,q} = g_{A,B}^p S_{A \setminus B \setminus a_1}^{q-1} r_{A \setminus B \setminus C} - S_A^p g_{A \setminus B, C}^q = S_A^p S_{A \setminus B}^{q+1} - S_A^{p+1} S_{A \setminus B \setminus a_1}^{q-1} r_{A \setminus B \setminus C},$$

where  $a_1 \notin B \subset A, a_1 \notin C \subset A \setminus B, |B| = p+1, |C| = q+1$ , and  $S_*^{-1} = 1$ . It is clear that every monomial of the form  $r_A r_{A \setminus B_1} \dots r_{A \setminus B_t}$ , where  $B_i \subset A, |B_i| = i$ , is reduced to  $S_A^t$  by a sequence of reductions w. r. t. the elements of type  $g_{A \setminus \{a_1, \dots, a_j\}, B_i}^i$ . Therefore, both monomials in the representation of  $s_{A,B,C,p,q}$  are reduced to the same monomial  $S_A^{p+q+1}$ .  $\square$

The following consequence will be used without notification.

**Corollary 2.4.** (a) *The ideals  $r_C \mathbf{gr} Q_n, \emptyset \neq C \subset I_n$  have zero pairwise intersections.*

(b) *Let  $G = \{u_1, \dots, u_s\}$  be a subset of the generators of  $\mathbf{gr} Q_n$  such that if  $B \subset C \in G$ , then  $B \in G$ , and let  $U$  be a subalgebra (with unit) in  $\mathbf{gr} Q_n$  generated by  $G$ . Then there is a natural isomorphism of left  $U$ -modules*

$$\mathbf{gr} Q_n \simeq U \oplus U \otimes \bigoplus_{C \not\subset U} r_C \mathbf{gr} Q_n.$$

*Proof.* Obviously, a normal form of an element  $r_A a \in F$  is  $r_A a'$  for some linear combination  $a'$  of normal monomials. This proves (a).

To prove (b), it is sufficient to notice that every monomial  $ar_C b$  is normal provided that  $a \in U$  is normal,  $C \notin G$ , and  $r_C b$  is normal.  $\square$

### 2.3. Algebra $\mathbf{gr} Q_n$ is Koszul.

*Proof of Theorem 2.1.* Now, we are ready to prove that the algebra  $\mathbf{gr} Q_n$  is Koszul. Consider the following complex  $K^n$  of right  $\mathbf{gr} Q_n$ -modules:

$$0 \rightarrow K_n^n \rightarrow K_{n-1}^n \rightarrow \dots \rightarrow K_0^n \rightarrow 0,$$

where  $K_0^n = \overline{\mathbf{gr} Q_n} = \sum_{r_C} r_C \mathbf{gr} Q_n$  is the augmentation ideal of  $\mathbf{gr} Q_n$ , and every  $K_{t+1}^n, t \geq 0$  is a free module with generators  $S(A : B)$ , where  $A \neq \emptyset, \min A \notin B \subset A$ , and  $|B| = t$ . The differential  $d : K_1^n \rightarrow K_0^n$  sends  $S(A : \emptyset)$  to  $r_A$ , while in higher degrees it is defined on the symbol  $S(A : B)$ , where  $A = \{a_1, \dots, a_m\}$  and  $B = \{b_1, \dots, b_t\}$  with  $a_i < a_{i+1}, b_j < b_{j+1}$ , by the rule

$$d(S(A : B)) = \sum_{i=1}^t (-1)^{t-1} S(A : B) \otimes (r_{A \setminus B} - r_{(A \setminus a_i) \setminus (B \setminus b_i)}).$$

In fact, it is a standard Koszul complex [Pr] for the quadratic algebra  $\mathbf{gr} Q_n$ , but with non-standard generators.

We claim that the complex  $K^n$  is acyclic. Then the complex  $K^n \oplus k$  (the addition is in the term  $K_0^n$ ) forms a linear free resolution of the trivial  $\mathbf{gr} Q_n$ -module  $k$ , hence, it follows that the algebra  $\mathbf{gr} Q_n$  is Koszul.

For completeness, let us also define the algebra  $Q_1 = \mathbf{gr} Q_1$  as a free algebra generated by  $r_{\{1\}}$  and the complex  $K^1 : 0 \rightarrow S(\{1\} : \emptyset) \rightarrow r_{\{1\}} \mathbf{gr} Q_1 \rightarrow 0$ . Obviously, it is acyclic.

We will show that  $K^n$  is acyclic by the induction on  $n \geq 2$ . First, notice that  $K^n$  is a direct sum of the complexes  $K_A^n$  for  $\emptyset \neq A \subset I_n$ , where the differential submodule  $K_A^n$  is spanned by the generators  $S(A : B)$ , where  $B \subset A$ , and  $r_A \in K_0^n = \mathbf{gr} Q_n$ . We have to show that every complex  $K_A^n$  is acyclic.

If  $|A| = k > 0$ , then the complex  $K_A^n$  is obviously isomorphic to  $K_{I_k}^n$ . Consider a subalgebra  $P_k \subset \mathbf{gr} Q_n$  generated by  $\{r_A | A \subset I_k\}$ . It is canonically isomorphic to  $\mathbf{gr} Q_k$  by [GRW]. Therefore, the subcomplex  $\widetilde{K}_{I_k}^n = r_{I_k} P_k \oplus \sum_{B \subset I_k} S(I_k : B) \otimes P_k$  is isomorphic to  $K_{I_k}^k$ . (It is indeed a subcomplex, because  $d(S(I_k : B)) \in \widetilde{K}_{I_k}^n$ .) If  $k < n$ , it is acyclic by the induction. Because of the decomposition

$$\mathbf{gr} Q_n = P_k \oplus P_k \otimes \bigoplus_{C \not\subset I_k} r_C \mathbf{gr} Q_n,$$

we have  $K_{I_k}^n \simeq \widetilde{K}_{I_k}^n \otimes (k + \bigoplus_{C \not\subset I_k} r_C \mathbf{gr} Q_n)$ . Therefore, all complexes  $K_A^n$  are acyclic whenever  $|A| < n$ .

It remains to show that the complex  $K = K_{I_n}^n$  is acyclic. Consider the following filtration on the complex  $K$ :

$$0 = J_0^n \subset \dots \subset J_n^n = K,$$

where for  $i \geq 2$  the differential submodule  $J_i^n$  is spanned by the set  $T_i^n = \{S(I_n : B) | \max B \leq i\}$ . We are going to show that all of them are acyclic. Because of the exact triples

$$0 \rightarrow J_i^n \rightarrow J_{i+1}^n \rightarrow J_{i+1}^n / J_i^n \rightarrow 0,$$

it equivalent to say that all quotients  $J_{i+1}^n / J_i^n$  are acyclic. By the induction arguments, we will assume that all the complexes  $J_t^n$  are acyclic for  $t < n$ .

First, the complex  $J_n^n / J_{n-1}^n = K / J_{n-1}^n$  is isomorphic to  $K_{I_{n-1}}^n$ ; the isomorphism is given by the map  $S(I_n : B) \rightarrow (-1)^{|B|} S(I_{n-1} : B \setminus n)$ . By the above, it is acyclic.

Now, consider the quotient complex  $J_t^n / J_{t-1}^n$  for  $t \leq n-1$ . It is spanned by the set  $U_t^n = \{S(I_n : B) | \max B = t\}$  with the differential

$$dS(I_n : B) = \sum_{i=1}^k (-1)^i S(I_n : B \setminus b_i) \otimes (r_{I_n \setminus B} - r_{I_n \setminus 1 \setminus (B \setminus b_i)})$$

(where  $B = \{b_1, \dots, b_k, t\}$  with  $b_1 < \dots < b_k < t$ ). We have  $dU_t^n \subset U_t^n \otimes N_t$ , where  $N_t$  is the subalgebra in  $\mathbf{gr} Q_n$  generated by  $\{r_C | t \notin C\}$ . Let us show that the subcomplex  $U_t^n \otimes N_t$  is isomorphic to the complex  $J_{t-1}^{n-1}$ . Consider the injective map  $U_t^n \otimes N_t \rightarrow K_{I_n \setminus t}^n$  given by  $\phi : S(I_n : B) \mapsto S(I_n \setminus t : B \setminus t)$ . The cyclic permutation  $\sigma := (n \rightarrow (n-1) \rightarrow \dots \rightarrow t \rightarrow n)$  maps the generators of  $K_{I_n \setminus t}^n$  to the set  $\{S(I_{n-1} : B) | n \notin B\}$  and maps  $N_t$  to  $N_n$ . Then the image of the composition  $\sigma \phi(U_t^n \otimes N_t)$  is exactly  $J_{t-1}^{n-1}$ . By the induction, the last complex is acyclic, and so is the complex  $U_t^n \otimes N_t$ . Because

$$J_t^n / J_{t-1}^n = U_t^n \otimes \mathbf{gr} Q_n \simeq U_t^n \otimes N_t \otimes (k \oplus \bigoplus_{C: t \in C} r_C \mathbf{gr} Q_n),$$

the complex  $J_t^n/J_{t-1}^n$  is acyclic as well.

Therefore, all the complexes  $J_t^n$  are acyclic, thus, the algebra  $\mathbf{gr} Q_n$  is Koszul.  $\square$

### 3. ACKNOWLEDGMENT

I am grateful to Mittag–Leffler Institute for hospitality during preparation this note.

### 4. ADDITION

Another proof of Theorem 1.1, due to S. Serconek and R. L. Wilson, has been announced in [GGRW]. According to [Re], this proof is based on another approach and seems to be more complicated.

### REFERENCES

- [B] J. Backelin, *A distributiveness property of augmented algebras, and some related homological results*, Ph. D. thesis, Stockholm, 1982
- [GRW] I. Gelfand, V. Retakh, R. L. Wilson, *Quadratic linear algebras associated with factorizations of noncommutative polynomials and noncommutative differential polynomials*, *Selecta Math. (N.S.)*, **7** (2001), 4, p. 493–523
- [GGRW] I. Gelfand, S. Gelfand, V. Retakh, R. Wilson, *Quasideterminants*, preprint math.QA/0208146
- [GGRSW] I. Gelfand, S. Gelfand, V. Retakh, S. Serconek, R. L. Wilson, *Hilbert series of quadratic algebras associated with pseudo-roots of noncommutative polynomials*, *J. Algebra*, **254** (2002), 2, p. 279–299
- [L] C. Lofwall, *On the subalgebra generated by the one-dimensional elements in the Yoneda Extalgebra*, *Lect. Notes in Math.*, **1352** (1988), pp. 291–338
- [Pi] D. I. Piontkovski, *On the Hilbert series of Koszul algebras*, *Functional Anal. and its Appl.*, **35** (2001), 2
- [PP] A. Polishchuk and L. Positselski, *Quadratic algebras*, preprint (1994–2000)
- [Po] L. E. Positselski, *Relation between the Hilbert series of dual quadratic algebras does not imply Koszulity*, *Functional Anal. and its Appl.*, **29**, 3, 83–87 (1995)
- [Pr] S. B. Priddy, *Koszul resolutions*, *Trans. AMS*, **152** (1970), 1, p. 39–60
- [Re] V. S. Retakh, private communication
- [Ro] J.–E. Roos, *On the characterization of Koszul algebras. Four counter-examples*, *Comptes Rendus Acad. Sci. Paris, Sér. I*, **321**, 1, 15–20 (1995)
- [U] V. A. Ufnarovsky, *Combinatorial and asymptotical methods in algebra*, *Sovr. probl. mat., Fund. napr.*, **57** (1990), p. 5–177 [Russian] Engl. transl.: *Algebra VI*, *Encycl. Math. Sci.*, Springer, Berlin 1995, p. 1–196

CENTRAL INSTITUTE OF ECONOMICS AND MATHEMATICS, NAKHIMOVSKY PROSP. 47, MOSCOW 117418, RUSSIA

*E-mail address:* piont@mccme.ru