

Fibonomial Cumulative Connection Constants

A.K.Kwaśniewski

Higher School of Mathematic and Applied Informatics
 PL-15-021 Białystok, ul.Kamienna 17, POLAND
 e-mail: kwandr@uwb.edu.pl

May 21, 2019

Abstract

In this note we present examples of cumulative connection constants - new Fibonomial ones included. All examples posses combinatorial interpretation.

vol 44 may 2005 Bulletin of the Institute of Combinatorics and its Applications ; to appear

1 Introduction

The cumulative connection constants (**ccc**) C_n were introduced by the Authors of [1]:

$$C_n = \sum_{k \geq 0} c_{n,k}, \quad n \geq 0,$$

where

$$p_n(x) = \sum_{0 \leq k \leq n} c_{n,k} q_k(x)$$

and Pascal-like array $\{c_{n,k}\}_{n,k \geq 0}$ of connection constants $c_{n,k}$, $n, k \geq 0$ "connects" two polynomial sequences (Note: $\{p_k(x)\}_{k \geq 0}$ is a polynomial sequence if $\deg p_k = k$).

Motivation 1

"The connection constants problem" i.e. combinatorial interpretations, algorithms of calculation, recurrences and other properties - is one of the central issues of the binomial enumeration in Finite Operator Calculus (FOC) of Rota-Roman and Others and in its afterwards extensions (see: abundant references in [8]-[10],[13]). Important knowledge on recurrences for $\{c_{n,k}\}_{n,k \geq 0}$ one draws from [2] - consult also the NAVIMA group program: <http://webs.uvigo.es/t10/navima/> .

Motivation 2

The cumulative connection constants (**ccc**) as defined above appear to represent combinatorial quantities of primary importance - as shown by examples below .

A special case of interest considered in [2, 1] is the case of monic persistent root polynomial sequences [3] characterized by the following conditions:

$$p_0(x) = q_0(x) = 1 ; \quad q_n(x) = q_{n-1}(x)(x-r_n) , \quad p_n(x) = p_{n-1}(x)(x-s_n) \quad n \geq 1.$$

The pair of persistent root polynomial sequences and hence corresponding connection constants array are then bijectively labeled by the pair of root sequences :

$$[r] = \{r_k\}_{k \geq 1} , \quad [s] = \{s_k\}_{k \geq 1} .$$

In this particular case of persistent root polynomial sequences the authors of [2] gave a particular name to connection constants : "generalized Lah numbers" - now denoted as follows $c_{n,k} = L_{n,k}$. The generalized Lah numbers do satisfy the following recurrence [2, 1] :

$$L_{n+1,k} = L_{n,k-1} + (r_{k+1} - s_{n+1})L_{n,k} \tag{1}$$

$$L_{0,0} = 1 , \quad L_{0,-1} = 0 \quad n, k \geq 0$$

from which the recurrence for cumulative connection constants (**ccc**) follows

$$C_{n+1} = (1 - s_{n+1})C_n + \sum_{k=0}^n c_{n,k}r_{k+1} \tag{2}$$

where $c_{n,k} = L_{n,k}$.

The clue examples of [1] are given by the following choice of a pair of root sequences :

the Fibonacci sequence

$$C_n = F_n; \quad n > 0 \longleftrightarrow [r] = \{r_k\}_{k \geq 1} = \{0, 0, 0, 0, 0, 0, \dots\}, \quad [s] = \{s_k\}_{k \geq 1} = \{0, 0, \dots, 0, \dots\} = [0]$$

the Lucas sequence

$$C_n = L_n; \quad n > 0 \longleftrightarrow [r] = \{r_k\}_{k \geq 1}; \quad r_1 = 0, r_2 = 2, r_k = 1 - (r_{k-1})^2 \text{ for } k > 2 \text{ and } [s] = [0].$$

2 Examples of cumulative connection constants - old and new

We shall supply now several examples of connection constants arrays and corresponding ccc's - Fibonomial case included - altogether with their combinatorial interpretation. (HELP! - For notation-help - see: Appendix).

Example 1. $c_{n,k} = \binom{n}{k}$, $x^n = \sum_{k \geq 0} \binom{n}{k} (x-1)^k \quad n \geq 0$;
 $E^{-1}(D)x^k = (x-1)^k \quad n \geq 0$.

$$C_n = \sum_{k \geq 0} \binom{n}{k} = 2^n = \text{number of all subsets of } S; \quad n = |S|; n \geq 0.$$

Here $E^a(D) = \sum_{n \geq 0} \frac{a^n}{n!} D^n$ is the translation operator ($D = \frac{d}{dx}$) and the recurrence (2) is obvious.

Example 2. $c_{n,k} = \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ denote Stirling numbers of the second type.
Then $x^n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} x^k$. Let $B_n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$, then
 $B_n = \text{number of all partitions of } S; \quad n = |S|; B_n \equiv \text{Bell numbers}.$

As the recurrence for Bell numbers reads $B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k$; $n \geq 0$
we easily get from (2) the inspiring identity

$$\sum_{k=0}^n \binom{n}{k} B_k = B_n + \sum_{k=1}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} k.$$

Example 3. $c_{n,k} = \left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]$ are Stirling numbers of the first kind. Then

$x^{\bar{n}} = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} x^k$, $n \geq 0$. $C_n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} = n!$ and the obvious recurrence $C_{n+1} = (n+1)C_n$ coincides with (2) .

Example 4. $c_{n,k} = \begin{pmatrix} n \\ k \end{pmatrix}_q$, $x^n = \sum_{k \geq 0} \begin{pmatrix} n \\ k \end{pmatrix}_q \Phi_k(x)$, $\Phi_k(x) = \prod_{s=0}^{k-1} (x - q^s)$,

$$C_n = \sum_{k \geq 0} \begin{pmatrix} n \\ k \end{pmatrix}_q = \text{number of all subspaces of } V(n, q) \text{ i.e.}$$

$$C_n = \sum_{k=0}^n \begin{pmatrix} n \\ k \end{pmatrix}_q = |L(n, q)| \equiv (1 +_q 1)^n,$$

where $L(n, q)$ denotes the lattice of all subspaces of $V(n, q)$ - (the n -th dimensional space $V(n, q)$ over Galois field $GF(q)$) - see: [4] .

According to Konvalina

$$C_n = \sum_{k=0}^n \begin{pmatrix} n \\ k \end{pmatrix}_q \equiv (1 +_q 1)^n$$

is at the same time the number of all possible choices with repetition of objects from exponentially weighted boxes [5, 6].

As q -Gaussian polynomials $\Phi_k(x) = \prod_{s=0}^{k-1} (x - q^s)$ constitute a persistent root polynomial sequence - the generalized Lah numbers - which are now are $\begin{pmatrix} n \\ k \end{pmatrix}_q = c_{n,k} = L_{n,k}$ must satisfy the recurrence equation

$$L_{n+1,k} = L_{n,k-1} + r_{k+1} L_{n,k}, \quad L_{0,0} = 1, \quad L_{0,-1} = 0 \quad n, k \geq 0,$$

where $[r] = \{q^{k-1}\}_{k \geq 1}$ is the root sequence determining $\{\Phi_n\}_{n \geq 0}$. See Appendix 2. Hence the recurrence (2) for the number C_n of all possible choices with repetition of objects from exponentially weighted boxes [5, 6] is of the form

$$C_{n+1} = C_n + \sum_{k=0}^n \begin{pmatrix} n \\ k \end{pmatrix}_q q^k$$

because

$$C_{n+1} = C_n + \sum_{k=0}^n c_{n,k} r_{k+1}.$$

In another q -umbral notation (see: Appendix 2)

$$(1 +_q 1)^{n+1} = (1 +_q 1)^n + \sum_{k=0}^n \begin{pmatrix} n \\ k \end{pmatrix}_q q^k.$$

Occasionally note that using the Möbius inversion formula (see: [4] - the q -binomial theorem Corollaries in section 5 - included) we have

$$x^n = \sum_{k \geq 0} \binom{n}{k}_q \Phi_k(x), \quad \Phi_k(x) = \sum_{l \geq 0} \binom{k}{l}_q (-1)^l q^{\binom{l}{2}} x^{k-l}, \quad n \geq 0$$

i.e. the identity

$$\prod_{s=0}^{k-1} (x - q^s) \equiv \sum_{l \geq 0} \binom{k}{l}_q (1)_q^l q^{\binom{l}{2}} x^{k-l}.$$

"The Möbius q -inverse example" is then the following.

Example 4'. $c_{n,k} = \binom{n}{k}_q (-1)^{n-k} q^{\binom{n-k}{2}}$, $x^n = \sum_{k \geq 0} c_{n,k} H_k(x)$, where (written with help of generalized shift operator [7, 8, 9, 10] - analogously to the use of $E^{-1}(D)x^k = (x-1)^k$ above in Example 1.) we have (see Appendix 2):

$$H_k(x) = E(\partial_q)x^k \equiv (x +_q 1)^k = \sum_{l \geq 0} \binom{k}{l}_q x^l.$$

Compare with $H_k(x, y)|_{y=1} = H_k(x)$ in [11] (pp. 240-241) and note that

$$C_n = \sum_{k=0}^n |c_{n,k}| = \sum_{k=0}^n \binom{n}{k}_q q^{\binom{k}{2}}$$

is the number of all possible choices without repetition of objects from exponentially weighted boxes [5, 6]. It is then combinatorial - natural to consider also another triad .

Example 4''. $c_{n,k} = \binom{n}{k}_q q^{\binom{k}{2}}$, $x^n = \sum_{k \geq 0} \binom{n}{k}_q q^{\binom{k}{2}} \Gamma_k(x)$,
 $\Gamma_k(x) = \sum_{l \geq 0} \binom{k}{l}_q (-1)^{k-l} x^l$, $n \geq 0$.

Example 5. This is The Fibonomial Example: $c_{n,k} = \binom{n}{k}_F$,
 $x^n = \sum_{k \geq 0} \binom{n}{k}_F \Xi_k(x)$ where $\binom{n}{k}_F$ are called Fibonomial coefficients (see A3).
Using the Möbius inversion formula - analogously to the Example 4 case - we get

$$\Xi_k(x) = \sum_{l \geq 0} \binom{k}{l}_f \mu(l, k) x^{k-1}, \quad n \geq 0,$$

where Möbius matrix μ is the unique inverse of the incidence matrix ζ defined for the "Fibonacci cobweb" poset introduced in [12] (see: A4 for explicit

formula of ζ from [16]). For combinatorial interpretation of these Fibonomial coefficients see: Appendix 4. Interpretation of C_n stated below results from combinatorial interpretation of Fibonomial coefficients. Combinatorial interpretation of ccc C_n in this case is then the following:

$C_n = \sum_{k \geq 0} \binom{n}{k}_F = \text{number of \underline{all} such subposets of the Fibonacci cobweb poset } P \text{ which are cobweb subposets starting from the level } F_k \text{ and ending at the level labeled by } F_n.$

As in the q -umbral case of the Example 4 we have " F -umbral" representation of the Fibonomial ccc C_n (see A3) . Namely

$$C_n = \sum_{k=0}^n \binom{n}{k}_F \equiv (1 +_F 1)^n$$

3 Remark on ccc - recurrences "exercise"

ccc-recurrences from examples 1-4 were an easy game to play. All these four cases of ccc might be interpreted on equal footing due to the ingenious scheme of one unified combinatorial interpretation by Konvalina [5, 6].

As for the other cases the "exercise" of finding the recurrences for ccc - up to the Fibonomial case - might be the slightly harder task. In this last case with Fibonomial coefficients one may derive the well known recurrence

$$\binom{n+1}{k}_F = F_{k-1} \binom{n}{k}_F + F_{n-k+2} \binom{n}{k-1}_F \quad (3)$$

$$\binom{n}{0}_F = 1, \quad \binom{0}{k}_F = 0 \text{ for } k > 0$$

or equivalently

$$\binom{n+1}{k}_F = F_{k+1} \binom{n}{k}_F + F_{n-k} \binom{n}{k-1}_F$$

$$\binom{n}{0}_F = 1, \quad \binom{0}{k}_F = 0 \text{ for } k > 0$$

also by combinatorial reasoning [16] (it might be a hard exercise - not to check it but to prove it another way - a use of [2] - being recommended). Derivation of the recurrence for Fibonomial ccc C_n - we leave as an exercise.

4 Appendix

4.1 On the notation used through out in this note

q -binomial and Fibonomial cases are specifications of the ψ -sequence choice in the so called ψ -calculus notation [8]-[10] in conformity with [14] referring back to works of Brenke and Boas on generalized Appell polynomials (see abundant references in [9])

4.2 q -Gaussian ccc

$$\begin{aligned} \binom{n+1}{k}_q &= q^k \binom{n}{k}_q + \binom{n}{k-1}_q \\ \binom{n}{0}_q &= 1, \quad n \geq 0, k \geq 1, \end{aligned} \quad (4)$$

where $n_q! = n_q(n-1)_q!; \quad 1_q! = 1, \quad n_q^k = n_q(n-1)_q \dots (n-k+1)_q,$
 $\binom{n}{k}_q \equiv \frac{n_q^k}{k_q!}.$

Remark 4.1. (see [15]) The dual to (4) recurrence is then given by

$$x\Phi_n(x) = q^n\Phi_n(x) + \Phi_{n+1}(x) \quad (5)$$

$$\Phi_0(x) = 1, \quad \Phi_{-1}(x) = 0; \quad n \geq 0,$$

in accordance with a well known fact (see: [4]) that

$$x^n = \sum_{k \geq 0} \binom{n}{k}_q \Phi_k(x), \quad (6)$$

where $\Phi_k(x) = \prod_{s=0}^{k-1} (x - q^s)$ are the q -Gaussian polynomials.

Recall now that the Jackson ∂_q -derivative is defined as follows [7]-[11]:
 $(\partial_q \varphi)(x) = \frac{\varphi(x) - \varphi(qx)}{(1-q)x}$ - and is also called the q -derivative. The consequent notation for the generalized shift operator [7]-[10],[13] is:

$$E^a(\partial_q)x^n = \exp_q\{a\partial_q\} = \sum_{k=0}^{\infty} \frac{a^k}{k_q!} \partial_q^k.$$

Then we identify $E^y(\partial_q)x^n \equiv (x +_q y)^n$. (Ward in [7] does not use any subscript like q or F or ψ . Naturally his " + " means not + but a "plus with a subscript" in our notation).

4.3 Fibonomial notation

In straightforward analogy consider now the *Fibonomial coefficients* $c_{n,k}$,

$$c_{n,k} = \binom{n}{k}_F = \frac{F_n!}{F_k! F_{n-k}!} = \binom{n}{n-k}_F,$$

where $n_F \equiv F_n, n_F! \equiv n_F(n-1)_F(n-2)_F(n-3)_F \dots 2_F 1_F$; $0_F! = 1; n_F^{\frac{k}{F}} = n_F(n-1)_F \dots (n-k+1)_F$; $\binom{n}{k}_F \equiv \frac{n_F^{\frac{k}{F}}}{k_F!}$ and linear difference operator ∂_F acting as follows: $\partial_F; \partial_F x^n = n_F x^{n-1}$; $n \geq 0$ - we shall call the F -derivative. Then in conformity with [7] and with notation as in [10, 9, 8, 13] one writes:

- (1) $(x +_F a)^n \equiv \sum_{k \geq 0} \binom{n}{k}_F a^k x^{n-k}$ where $\binom{n}{k}_F \equiv \frac{n_F^{\frac{k}{F}}}{k_F!}$
and $n_F^{\frac{k}{F}} = n_F(n-1)_F \dots (n-k+1)_F$;
- (2) $(x +_F a)^n \equiv E^a(\partial_F x^n)$; $E^a(\partial_F) = \sum_{n \geq 0} \frac{a^n}{n_F!} \partial_F^n$;
 $E^a(\partial_F) f(x) = f(x +_F a)$, $E^a(\partial_F)$ is corresponding generalized translation operator.

4.4 Combinatorial interpretation of Fibonomial Coefficients [12]

Let us depict a partially ordered infinite set P via its finite part Hasse diagram to be continued ad infinitum in an obvious way as seen from the figure below.

This Fibonacci cobweb partially ordered infinite set P is defined as in [12] via its finite part - cobweb subposet $P_m(\text{rooted at } F_1 \text{ level subposet})$ to be continued ad infinitum in an obvious way as seen from the figure of P_5 below. It looks like the Fibonacci tree with a specific "cobweb". It is identified with Hasse diagram of the partial order set P .




Fig. 1. The construction of the Fibonacci "cobweb" poset

If one defines this poset P with help of its incidence matrix ζ representing P uniquely then one arrives at ζ with easily recognizable staircase-like structure - of zeros in the upper part of this upper triangle matrix ζ . This structure is depicted by the Figure 2 where: empty places mean zero values (under diagonal) and filled with $-$ places mean values one (above the diagonal).




Fig. 2. The staircase structure of incidence matrix ζ

The characteristic function ζ of the partial order relation in the Fibonacci cobweb poset P might be expressed by Kronecker delta as follows [16] :

$$\zeta = \zeta_1 - \zeta_0$$

where for $x, y \in \mathbf{N}$:

$$\begin{aligned} \zeta_1(x, y) &= \sum_{k=0}^{\infty} \delta(x + k, y) \\ \zeta_0(x, y) &= \sum_{k \geq 0} \sum_{s \geq 0} \delta(x, F_{s+1} + k) \sum_{r=1}^{F_s - k - 1} \delta(k + F_{s+1} + r, y) \end{aligned}$$

and naturally

$$\delta(x, y) = \begin{cases} 1 & x = y \\ 0 & x \neq y \end{cases}.$$

Right from the Hasse diagram of P here now obvious observations follow

Observation 4.1. *The number of maximal chains starting from the root (level F_1) to reach any point at the n -th level labeled by F_n is equal to $n_F!$.*

Observation 4.2. *The number of maximal chains starting from any fixed point at the level labeled by F_k to reach any point at the n -th level labeled by F_n is equal to n_F^m ($n = k + m$).*

Let us denote by P_m a subposet of P consisting of points up to m -th level points

$$\bigcup_{s=1}^m \Phi_s ; \Phi_s \text{ is the set of elements of the } s\text{-th level}$$

Consider now the following behavior of a "sub-cob" moving from any given point of the F_k level of the poset up. It behaves as it has been born right there and can reach at first F_2 points up then F_3 points up, F_4 and so on - thus climbing up to the level $F_{k+m} = F_n$ of the poset P . It can see - as its Great Ancestor at the root F_1 -th level and potentially follow one of its own accessible finite subposet P_m . One of many P_m 's rooted at the k -th level might be found. How many?

The answer is :

Observation 4.3. *Let $n = k + m$. The number of subposets P_m rooted at any fixed point at the level labeled by F_k and ending at the n -th level labeled by F_n is equal to*

$$\binom{n}{m}_F = \binom{n}{k}_F = \frac{n_F^k}{k_F!}.$$

References

- [1] L. Colucci, O. D’Antona, C. Mereghetti: *Fibonacci and Lucas Numbers as Cumulative Connection Constants*, The Fibonacci Quarterly 38.2 (2000), pp.157-164
- [2] E.Damiani,O. D’Antona, G. Naldi *On the Connection Constants*,Studies in Appl. Math. 85(4)(1991), pp. 289-302
- [3] A. Di Bucchianico ,D. Loeb: *Sequences of Binomial type with Persistent Roots*, J. Math. Anal. Appl. 199(1996),pp. 39-58
- [4] Goldman J. Rota G-C.: *On the Foundations of Combinatorial Theory IV; Finite-vector spaces and Eulerian generating functions* Studies in Appl. Math. vol. 49 1970, pp. 239-258
- [5] J.Konvalina: *Generalized binomial coefficients and the subset-subspace problem*, Adv. in Appl. Math. Vol. 21 (1998), pp. 228-240
- [6] J. Konvalina: *A Unified Interpretation of the Binomial Coefficients, the Stirling Numbers and the Gaussian Coefficients*, The American Mathematical Monthly vol. 107, No 10 , (2000), pp.901-910
- [7] Ward M.: *A calculus of sequences*, Amer.J. Math. 58 (1936) pp. 255-266
- [8] A. K. Kwaśniewski: *Main theorems of extended finite operator calculus*, Integral Transforms and Special Functions Vol 14 , No 6, (2003),pp. 499-516
- [9] A.K.Kwaśniewski: *Towards ψ -extension of Finite Operator Calculus of Rota*, Rep. Math. Phys.48 No 3 (2001)pp. 305-342

- [10] A.K.Kwaśniewski: *On Simple Characterisations of Sheffer ψ -polynomials and Related Propositions of the Calculus of Sequences*, Bulletin de la Soc. des Sciences et de Lettres de Lodz ; 52, Ser. Rech. Deform. 36 (2002): 45-65 see :ArXiv : math.CO/0312397
- [11] S.R.Roman: *More on the umbral calculus with emphasis on the q -umbral calculus*, J. Math. Anal. Appl. 107(1985), pp. 222-254
- [12] A.K.Kwaśniewski: *Information on combinatorial interpretation of Fibonacci coefficients*, Bull. Soc. Sci. Lett. Lodz Ser. Rech. Deform. 53, Ser. Rech.Deform. 42 (2003) pp. 39-41 ArXiv: math.CO/0402291 v 1 22 Feb 2004
- [13] A.K.Kwaśniewski: *On extended finite operator calculus of Rota and quantum groups*, Integral Transforms and Special Functions, Vol. 2, No.4(2001) pp. 333-340.
- [14] O.V. Viskov: *Operator characterization of generalized Appel polynomials*, Soviet Math. Dokl. 16 (1975),pp. 1521-1524
- [15] A. K. Kwaśniewski: *On duality triads*, Bull. Soc. Sci. Lett. Lodz Ser. Rech. Deform. 53, Ser. Rech.Deform. 42 (2003) pp.11-25 ArXiv: math.GM/0402260 v 1 Feb. 2004 ; A. K. Kwaśniewski: *The second part of on duality triads' paper* Bull. Soc. Sci. Lett. Lodz Ser. Rech. Deform. 53, Ser. Rech.Deform. 42 (2003) pp.27 -37 ArXiv: math.GM/0402288 v 1 Feb. 2004
- [16] A.K.Kwaśniewski: *Combinatorial iterpretation of the recurrence relation for fibonomial coefficients*, ArXiv: math.CO/0403017 v1 1 March 2004