

MORE ON: THE REVISED GCH AND MIDDLE DIAMOND

SAHARON SHELAH

The Hebrew University of Jerusalem
Einstein Institute of Mathematics
Edmond J. Safra Campus, Givat Ram
Jerusalem 91904, Israel

Department of Mathematics
Hill Center-Busch Campus
Rutgers, The State University of New Jersey
110 Frelinghuysen Road
Piscataway, NJ 08854-8019 USA

ABSTRACT. We strengthen the revised GCH theorem by showing, e.g., that for $\lambda = \text{cf}(\lambda) > \beth_\omega$, for all but finitely many regular $\kappa < \beth_\omega$, λ is accessible on cofinality κ in a weak version of it holds. In particular, $\lambda = 2^\mu = \mu^+ > \beth_\omega$ implies the diamond on λ is restricted to cofinality κ for all but finitely many $\kappa \in \text{Reg} \cap \beth_\omega$ and we strengthen the results on the middle diamond. Moreover, we get stronger results on the middle diamond.

The author would like to thank the United States-Israel Binational Science Foundation for partial support of this research and Alice Leonhardt for the beautiful typing. Publication 829

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - TEX

§0 INTRODUCTION

The main result of this paper is to define for any cardinal λ a set $\mathfrak{d}_0(\lambda)$ of regular cardinals $< \lambda$ such that for strong limit $\theta < \lambda$ we prove that $\theta \cap \mathfrak{d}_0(\lambda)$ is finite, and for every $\kappa \in \text{Reg} \cap \theta \setminus \mathfrak{d}_0(\lambda)$, in some sense λ has not too many subsets of cardinality κ . This serves as our main aim here to use this to show: if $\text{cf}(\lambda) > \mu$ and $\kappa \in \text{Reg} \cap \mu$ satisfies $\lambda = \sup\{\alpha : \kappa \notin \mathfrak{d}_0(\alpha)\}$ then λ has a “good” sequence $\langle \mathcal{P}_\alpha : \alpha < \lambda \rangle$, $\mathcal{P}_\alpha \subseteq [\alpha]^{\leq \kappa}$ and if $\lambda = \lambda^\mu$ more (see 3.5, 3.8).

This gives as a main consequence that: if $\mu \geq \theta$, $\lambda = \text{cf}(2^\mu)$ then (λ, κ) has middle diamond for all but finitely many $\kappa \in \text{Reg}$ satisfying $\beth_\omega(\kappa) \leq \mu$. Also $\lambda = 2^\mu = \mu^+ > \theta \Rightarrow \lambda$ has diamond on cofinality κ for all regular κ for which $\beth_\omega(\kappa) < \lambda$ except finitely many. We also strengthen the results on the middle diamond (in ZFC, see below).

So this is part of pcf theory ([Sh:g]) continuing in particular [Sh 460]. As I have been very happy about [Sh 460] and the proofs here give a shorter proof of the main theorem there, we have here (§1) gives a self-contained proof of the revised GCH, the main theorem of [Sh 460] and discuss it.

By pcf theory ([Sh:g],[Sh 460]) a worthwhile choice of power is (for $\kappa < \lambda$ regular) $\lambda^{[\kappa]}$ (or $\lambda^{<\kappa>}$), the minimal cardinality of a family of subsets of λ each of cardinality $\leq \kappa$ such that any other subset of λ of cardinality κ is equal to (or is contained in) the union of $< \kappa$ members of the family (see Definition 1.2). Let $\mathfrak{d}^+(\lambda) = \{\kappa : \kappa \text{ is regular } < \lambda \text{ and } \lambda < \lambda^{<\kappa>}\}$.

This gives a good partition of the exponentiation as $\lambda^\kappa = \lambda \Leftrightarrow 2^\kappa \leq \lambda$ & $(\forall \sigma)(\sigma = \text{cf}(\sigma) \leq \kappa \Rightarrow \lambda^{<\sigma>} = \lambda)$. So G.C.H. is equivalent to: κ regular $\Rightarrow 2^\kappa = \kappa^+$ and $[\kappa < \lambda \text{ are regular} \Rightarrow \lambda^{<\kappa>} = \lambda]$. See more in [Sh 460] where it is proved that (the revised G.C.H.):

- ⊗ if $\lambda > \beth_\omega$ then $\mathfrak{d}^+(\lambda) \cap \beth_\omega$ is bounded, i.e., $\lambda = \lambda^{<\kappa>}$ for every regular $\kappa < \beth_\omega$ large enough.

We can replace $\beth_\omega$ by any strong limit singular cardinal θ .

The advances in pcf theory reveal several natural hypotheses. The Strongest Hypothesis ($\text{pp}(\mu) = \mu^+$ for every singular μ) is very nice but it implies the SCH hence it does not follow from ZFC. The status of the Weak Hypothesis (somewhat more than $\{\mu : \text{cf}(\mu) < \mu < \lambda \leq \text{pp}(\mu)\}$ is at most countable) is not known but I am sure is consistent though it has large consistency strength. I am not sure about $(\forall \mathfrak{a})(|\mathfrak{a}| \geq |\text{pcf}(\mathfrak{a})|)$. Still better than ⊗ would be (we believe but do not know it, particularly (2)).

- 0.1 Conjecture.* 1) For every λ , $\mathfrak{d}^+(\lambda)$ is finite, or at least
2) For every strong limit μ , $\lambda \geq \mu \Rightarrow \mathfrak{d}^+(\lambda) \cap \mu$ is finite.

Now $\mathfrak{d}_0(\lambda) \cap \theta$ being finite is a step in the right direction and is enough to improve the results of [Sh 775]. In particular being able to use $\kappa = \aleph_n$ for some n rather than κ regular $< \beth_\omega$ seems crucial in abelian group theory (as there are non-free almost κ -free abelian group of cardinality κ when $\kappa = \aleph_n$).

So we can hope to get the right objects in each cardinality $\aleph_n$ whereas consistently they may not exist for arbitrary $\kappa = \text{cf}(\kappa) < \beth_\omega$, this is the case in abelian group theory.

This work also continues ones on $I[\lambda]$. By [Sh 108] if θ is strong limit singular, for some $A \in I[\lambda]$ for some $\mathbf{c} : [\mu^+]^2 \rightarrow \text{cf}(\mu)$, if $B \subseteq \mu, \mathbf{c} \upharpoonright [B]^2$ constant (or just has bounded range), $\delta = \sup(B)$, $\text{cf}(\delta) \neq \text{cf}(\mu)$ then $\delta \in A$.

By Dzamanja and Shelah [DjSh 562], using [Sh 460], if $\lambda = \text{cf}(\lambda) > \theta$, a strong limit singular, for some $\kappa < \theta$, for some $A \in I[\lambda]$, if for every $A' \subseteq A, |A'| < \theta$ for some $\mathbf{c} : [A'] \rightarrow \kappa$, we have: if $B \subseteq A', \mathbf{c} \upharpoonright [B]$ is constant, $\delta = \sup[B]$, $\text{cf}(\delta) > \kappa$ then $\delta \in A$. By [Sh 589, 5.20], T_D helps.

We try to make this paper as self-contained as is reasonably possible.

0.2 Definition. 1) For an ideal J on a set X :

- (a) $J^+ = \mathcal{P}(X) \setminus J$; we agree that J determines X so $X = \text{Dom}(J)$, this is an abuse of notation when $\cup\{A : A \in J\} \subset X$ but usually clear in the context
- (b) for a two-place relation R on Y and for $f, g \in {}^XY$, let fR_Jg means $\{t \in X : \neg f(t)Rg(t)\} \in J$; the cases we shall use are $=, \neq, <, \leq$.

2) If D is a filter on X , J the dual ideal on X (i.e., $J = \{X \setminus A : A \in D\}$) we may replace J by D .

3) Let $(\forall^J t)\varphi(t)$ mean $\{t : \neg\varphi(t)\} \in J$ similarly $\exists^J, \forall^D, \exists^D$.

4) Let $S_\kappa^\lambda = \{\delta < \lambda : \text{cf}(\delta) = \kappa\}$ and $S_{<\kappa}^\lambda = \{\delta < \lambda : \text{cf}(\delta) < \kappa\}$.

We quote (see [Sh 71], [Sh 506, §3], in 0.3, 0.4, 0.5)).

0.3 Definition. 1) Let $\bar{A} = \langle A_i : i \in X \rangle, D$ a filter on X and for simplicity first assume $i \in X \Rightarrow A_i \neq \emptyset$. We let

- (a) $T_D^0(\bar{A}) = \sup\{|\mathcal{F}| : \mathcal{F} \subseteq \Pi(\bar{A}) \text{ and } f_1 \neq f_2 \in \mathcal{F} \Rightarrow f_1 \neq_D f_2\}$
- (b)

$$T_D^1(\bar{A}) = \text{Min}\{|\mathcal{F}| : (i) \quad \mathcal{F} \subseteq \Pi(\bar{A})$$

$$(ii) \quad f_1 \neq f_2 \in \mathcal{F} \Rightarrow f_1 \neq_D f_2$$

$$(iii) \quad \mathcal{F} \text{ maximal under } (i) + (ii)\}$$

- (c) $T_D^2(\bar{A}) = \text{Min}\{|\mathcal{F}| : \mathcal{F} \subseteq \Pi\bar{A} \text{ and for every } f_1 \in \Pi\bar{A}, \text{ for some } f_2 \in \mathcal{F} \text{ we have } \neg(f_1 \neq_D f_2)\}$.

- 2) If $\{i : A_i = \emptyset\} \in J$ then we let $T_D^\ell(\bar{A}) = T_{D \upharpoonright Y}^\ell(\bar{A} \upharpoonright Y)$ where $Y = \{i : A_i \neq \emptyset\}$; if $\{i : A_i \neq \emptyset\} \in J$ then $T_D^\ell(\bar{A}) = 0$.
- 3) For $f \in {}^\kappa \text{Ord}$ and $\ell < 3$ let $T_D^\ell(f)$ means $T_D^\ell(\langle f(\alpha) : \alpha < \kappa \rangle)$.
- 4) If $T_D^0(\bar{A}) = T_D^1(\bar{A}) = T_D^2(\bar{A})$ then let $T_D(\bar{A}) = T_D^l(\bar{A})$ for $l < 3$; similarly $T_D(\bar{A})$; we say that $\mathcal{F}$ witness $T_D(\bar{A}) = \lambda$ if it is as in the definition of $T_D^1(\bar{A}) = \lambda$; similarly $T_D(f)$.

Remark. Actually the case $\bar{A} = \langle \lambda_\alpha : \alpha < \lambda \rangle$ is enough so we concentrate on it.

0.4 Claim. 0) If $D_0 \subseteq D_1$ are filters on κ then $T_{D_0}^\ell(\bar{\lambda}) \leq T_{D_1}^\ell(\bar{\lambda})$ for $\ell = 0, 2$.

- 1) $T_D^2(\bar{\lambda}) \leq T_D^1(\bar{\lambda}) \leq T_D^0(\bar{\lambda})$, in particular $T_D^1(\bar{\lambda})$ is well defined.
- 2) If $(\forall i) \lambda_i > 2^\kappa$ then $T_D^0(\bar{\lambda}) = T_D^1(\bar{\lambda}) = T_D^2(\bar{\lambda})$ so the supremum in 0.3(a) is obtained (so, e.g., $T_D^0(\bar{\lambda}) > 2^\kappa$ suffice; also $(\forall i) \lambda_i \geq 2^\kappa$ suffice).

Proof. 0) Check.

- 1) If $\mathcal{F}$ exemplifies the value of $T_D^1(\bar{\lambda})$, it also exemplifies $T_D^2(\bar{\lambda}) \leq |\mathcal{F}|$ hence but easily $T_D^2(\bar{\lambda}) \leq T_D^1(\bar{\lambda})$ is as in the definition it can be extended. In the definition of $T_D^0(\bar{\lambda})$ the Min is taken over a non-empty set (as maximal such $\mathcal{F}$ exists), so $T_D^0(\bar{\lambda})$ is well defined as well as $T_D^1(\bar{\lambda})$.

Lastly, if $\mathcal{F}$ exemplifies the value of $T_D^1(\bar{\lambda})$ it also exemplifies $T_D^0(\bar{\lambda}) \geq |\mathcal{F}|$, so $T_D^1(\bar{\lambda}) \leq T_D^0(\bar{\lambda})$.

- 2) Let μ be 2^κ . Assume that the desired conclusion fails so $T_D^2(\bar{\lambda}) < T_D^0(\bar{\lambda})$, so there is $\mathcal{F}_0 \subseteq \Pi \bar{\lambda}$, such that $[f_1 \neq f_2 \in \mathcal{F}_0 \Rightarrow f_1 \neq_D f_2]$, and $|\mathcal{F}_0| > T_D^2(\bar{\lambda}) + \mu$ (by the definition of $T_D^0(\bar{\lambda})$). Also there is $\mathcal{F}_2 \subseteq \Pi \bar{\lambda}$ exemplifying the value of $T_D^2(\bar{\lambda})$. For every $f \in \mathcal{F}_0$ there is $g_f \in \mathcal{F}_2$ such that $\neg(f \neq_D g_f)$ (by the choice of $\mathcal{F}_2$). As $|\mathcal{F}_0| > T_D^2(\bar{\lambda}) + \mu$ for some $g \in \mathcal{F}_2$, $\mathcal{F}^* =: \{f \in \mathcal{F}_0 : g_f = g\}$ has cardinality $> T_D^2(\bar{\lambda}) + \mu$. Now for each $f \in \mathcal{F}^*$ let $A_f = \{i < \kappa : f(i) = g(i)\}$ clearly $A_f \in D^+$. Now $f \mapsto A_f/D$ is a function from $\mathcal{F}^*$ into $\mathcal{P}(\kappa)/D$, hence, if as $\mu \geq |\mathcal{P}(\kappa)/D|$, it is not one to one (by cardinality consideration) so for some $f' \neq f''$ from $\mathcal{F}^*$ (hence from $\mathcal{F}_0$) we have $A_{f'}/D = A_{f''}/D$; but so

$$\{i < \kappa : f'(i) = f''(i)\} \supseteq \{i < \kappa : f'(i) = g(i)\} \cap \{i < \kappa : f''(i) = g(i)\} = A_{f'}/D$$

hence is $\neq \emptyset \pmod D$, so $\neg(f' \neq_D f'')$, contradiction the choice of $\mathcal{F}_0$. $\square_{0.4}$

0.5 Claim. *Let J be a σ -complete ideal on κ .*

- 1) *If $\bar{A} = \langle A_i : i < \kappa \rangle, \bar{\lambda} = \langle \lambda_i : i < \kappa \rangle, \lambda_i = |A_i|$ then $T_J^\ell(\bar{A}) = T_J^\ell(\bar{\lambda})$ and if $A \in J, B = \kappa \setminus A$ then $T_J^\ell(\bar{\lambda}) = T_{J \upharpoonright B}^\ell(\bar{\lambda} \upharpoonright B)$.*
- 2) *$T_J(\bar{\lambda}) > 2^\kappa$ iff $(\forall_J t)(\lambda_t > 2^\kappa)$.*
- 3) *$T_J^\ell(\bar{\lambda}^1) \leq T_J^\ell(\bar{\lambda}^2)$ if $(\forall_J t)(\lambda_t^1 \leq \lambda_t^2)$.*
- 4) *If $\text{Dom}(J) = \cup\{A_\varepsilon : \varepsilon < \zeta\}, \zeta < \sigma$ and $\lambda_i \geq 2^\kappa$ for $i < \kappa$ then $T_J^0(\bar{\lambda}) = \text{Min}\{T_{J \upharpoonright A_\varepsilon}^0(\bar{\lambda} \upharpoonright A_\varepsilon) : \varepsilon < \zeta \text{ and } A_\varepsilon \in J^+\}$.*

Proof. E.g. (and the one we use):

- 4) Let $A'_\varepsilon = A_\varepsilon \setminus \cup\{A_\xi : \xi < \varepsilon\}$ for $\varepsilon < \zeta$.

First assume that $\mathcal{F} \subseteq \Pi \bar{\lambda}$ and $f_1 \neq f_2 \in \mathcal{F}_2 \Rightarrow f_1 \neq_J f_2$. Then for each $\varepsilon < \zeta$ satisfying $A_\varepsilon \in J^+$ clearly $\mathcal{F}^{[\varepsilon]} = \{f \upharpoonright A_\varepsilon : f \in \mathcal{F}\}$ satisfies $|\mathcal{F}^{[\varepsilon]}| = |\mathcal{F}|$ as $f \mapsto f \upharpoonright A_\varepsilon$ is one to one by the assumption on $\mathcal{F}$ and $\mathcal{F}^{[\varepsilon]} \subseteq \prod_{i \in A_\varepsilon} \lambda_i$; so

$|\mathcal{F}| = |\mathcal{F}^{[\varepsilon]}| \leq T_{J \upharpoonright A_\varepsilon}(\bar{\lambda} \upharpoonright A_\varepsilon)$. As this holds for every $\varepsilon < \zeta$ for which $A_\varepsilon \in J^+$ we get $|\mathcal{F}| \leq \text{Min}\{T_{J \upharpoonright A_\varepsilon}(\bar{\lambda} \upharpoonright A_\varepsilon) : \varepsilon < \zeta, A_\varepsilon \in J^+\}$. By the demand on $\mathcal{F}$ we get the inequality $\leq$ in part (4). Second, assume $\mu < \text{Min}\{T_{J \upharpoonright A_\varepsilon}^0(\bar{\lambda} \upharpoonright A_\varepsilon) : \varepsilon < \zeta, A_\varepsilon \in J^+\}$. So for each such ε there is $\mathcal{F}_\varepsilon \subseteq \prod_{i \in A_\varepsilon} \lambda_i$ such that $f \neq g \in \mathcal{F}_\varepsilon \Rightarrow f \neq_{J \upharpoonright A_\varepsilon} g$, $|\mathcal{F}_\varepsilon| \geq \mu^+$. Let $f_\alpha^\varepsilon \in \mathcal{F}_\varepsilon$ be pairwise distinct, and define $f_\alpha \in \Pi \bar{\lambda}$ for $\alpha < \lambda^+$ as follows $f_\alpha \upharpoonright A'_\varepsilon = f_\alpha^\varepsilon$ when $A_\varepsilon \in J^+, f_\alpha \upharpoonright A'_\xi$ is zero otherwise.

Now check.

0.6 Definition. For regular λ and stationary $S \subseteq \lambda$ let $(D\ell)_{\lambda, S}$ mean that we can find $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha \in S \rangle, \mathcal{P}_\alpha \subseteq \mathcal{P}(\alpha)$ of cardinality $< \lambda$ such that for every $A \subseteq \lambda$ the set $\{\alpha \in S : A \cap \alpha \in \mathcal{P}_\alpha\}$ is stationary.

0.7 Definition. For λ regular uncountable let $I[\lambda]$ be the family of sets $S \subseteq \lambda$ which have a witness $(E, \bar{\mathcal{P}})$ for $S \in I[\lambda]$, which means

- (*) E is a club of $\lambda, \bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha < \lambda \rangle, \mathcal{P}_\alpha \subseteq \mathcal{P}(\alpha), |\mathcal{P}_\alpha| < \lambda$, and for every $\delta \in E \cap S$ there is an unbounded subset C of δ of order type $< \delta$ satisfying $\alpha \in C \Rightarrow C \cap \alpha \in \mathcal{P}_\alpha$.

0.8 Claim. ([Sh 420]): 1) For λ regular uncountable, $S \in I[\lambda]$ iff there is a pair $(E, \bar{\mathcal{P}}), E$ a club of $\lambda, \bar{a} = \langle a_\alpha : \alpha < \lambda \rangle, a_\alpha \subseteq \alpha$ such that $\beta \in a_\alpha \Rightarrow a_\beta = a_\alpha \cap \beta$ and $\delta \in E \cap S \Rightarrow \delta = \sup(a_\delta) > \text{otp}(a_\delta)$ (or even $\delta \in E \cap S \Rightarrow \delta = \sup(a_\delta)$), $\text{otp}(a_\delta) = \text{cf}(\delta) < \delta$.

2) If $\kappa^+ < \lambda$ and λ, κ are regular, then for some stationary $S \in I[\lambda]$ we have $\delta \in S \Rightarrow \text{cf}(\delta) = \kappa$.

0.9 Claim. 1) Assume that $f_\alpha \in {}^\kappa\text{Ord}$ for $\alpha < \lambda$, $\lambda = (2^\kappa)^+$ or just $\lambda = \text{cf}(\lambda)$ and $(\forall \alpha < \lambda)(|\alpha|^\kappa < \lambda)$ and $S_1 \subseteq \{\delta < \lambda: \text{cf}(\delta) > \kappa\}$ is stationary. Then for some stationary $S_2 \subseteq S_1$ we have: for each $i < \kappa$ the sequence $\langle f_\alpha(i) : \alpha \in S_2 \rangle$ is increasing or is constant.

2) If D is a filter on κ and $f_\alpha \in {}^\kappa\text{Ord}$ for $\alpha < \delta$ is $<_D$ -increasing and $\text{cf}(\delta) > 2^\kappa$ then $\langle f_\alpha : \alpha < \delta \rangle$ has a $<_D$ -eub, i.e.,

$$(i) \quad \alpha < \delta \Rightarrow f_\alpha \leq_D f_\delta$$

$$(ii) \quad f' \in {}^\kappa\text{Ord} \ \& \ f' <_D \text{Max}\{f, 1_\kappa\} \text{ then } (\exists \alpha < \delta)(f' <_D f_\alpha).$$

Proof. See [Sh 110], [Sh:g].

§1 THE REVISED G.C.H. REVISITED

Here we give a proof of the RGCH which requires little knowledge; this is the main theorem of [Sh 460], see also [Sh 513, §1]. The presentation is self-contained; in particular, do not use the pcf-theorem (hence repeat some proofs, in weak forms).

- 1.1 Definition.** 1) For $\lambda > \theta \geq \sigma = \text{cf}(\sigma)$ let $\lambda^{[\sigma, \theta]} = \text{Min}\{|\mathcal{P}| : \mathcal{P} \subseteq [\lambda]^{\leq \theta} \text{ and every } u \in [\lambda]^{\leq \theta} \text{ is the union of } < \sigma \text{ members of } \mathcal{P}\}$.
 2) Let $\lambda^{[\sigma]} = \lambda^{[\sigma, \sigma]}$.
 3) For $\lambda \geq \theta^{[\sigma, \kappa]}$ let $\lambda^{[\sigma, \kappa, \theta]} = \text{Min}\{|\mathcal{P}| : \mathcal{P} \subseteq [\lambda]^{\leq \kappa} \text{ such that for every } u \subseteq \lambda \text{ of cardinality } \leq \theta \text{ we can find } i^* < \sigma \text{ and } u_i \subseteq u \text{ for } i < i^* \text{ such that } u = \cup\{u_i : i < i^*\} \text{ and } [u_i]^\kappa \subseteq \mathcal{P}\}$.
 4) We may replace θ by $< \theta$ with the obvious meaning (also $< \kappa$).

- 1.2 Definition.** 1) For $\lambda > \theta \geq \text{cf}(\sigma) = \sigma$ let $\lambda^{< \sigma, \theta >} = \text{Min}\{|\mathcal{P}| : \mathcal{P} \subseteq [\lambda]^\theta \text{ and every } u \in [\lambda]^{\leq \theta} \text{ is included in the union of } < \sigma \text{ members of } \mathcal{P}\}$.
 2) Let $\lambda^{< \sigma >} = \lambda^{< \sigma, \sigma >}$.
 3) For $\lambda \geq \theta^{< \sigma, \kappa >}$ let $\lambda^{< \sigma, \kappa, \theta >} = \text{Min}\{|\mathcal{P}| : \mathcal{P} \subseteq [\lambda]^{\leq \kappa} \text{ such that for every } u \subseteq \lambda \text{ of cardinality } \leq \theta \text{ we can find } i^* < \sigma \text{ and } u_i \subseteq u \text{ for } i < i^* \text{ such that } u \subseteq \cup\{u_i : i < i^*\} \text{ satisfying } (\forall v \in [u_i]^{\leq \kappa})(\exists w \in \mathcal{P})(v \subseteq w)\}$.
 4) We may replace θ by $< \theta$ with the obvious meaning (also $< \kappa$).

1.3 Observation. Let $\lambda > \theta \geq \kappa \geq \sigma = \text{cf}(\sigma)$.

- 1) $\lambda^{< \kappa >} \leq \lambda^{[\kappa]} \leq \lambda^{< \kappa >} + 2^\kappa$.
 2) $\lambda^{< \sigma, \theta >} \leq \lambda^{[\sigma, \theta]} \leq \lambda^{< \sigma, \theta >} + 2^\theta$ (but see (3)).
 3) If $\text{cf}(\theta) < \sigma$ then $\lambda^{< \sigma, \theta >} = \Sigma\{\lambda^{< \sigma, \theta' >} : \sigma \leq \theta' < \theta\}$ and $\lambda^{[\sigma, \theta]} = \Sigma\{\lambda^{[\sigma, \theta']} : \sigma \leq \theta' < \theta\}$.
 4) $\lambda^{< \sigma, \kappa, \theta >} \leq \lambda^{[\sigma, \kappa, \theta]} \leq \lambda^{< \sigma, \kappa, \theta >} + 2^\kappa$.

Proof. Easy.

The main claim of this section is

1.4 Claim. Assume

- (a) $\aleph_0 < \sigma = \text{cf}(\sigma) \leq \kappa < \partial \leq \theta$
 (b) J is a σ -complete ideal on κ
 (c) $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$

- (d) $T_J(\bar{\lambda}) = \lambda$
- (e) $\lambda_i^{[\partial, \theta]} = \lambda_i$ for $i < \kappa$ (yes ∂ not ∂_i !)
- (f) if $\partial_i < \partial$ for $i < \kappa$ then $\prod_{i < \kappa} \partial_i < \partial$
- (g) $\theta = \theta^\kappa$ and $2^\theta \leq \lambda$.

Then $\lambda^{[\partial, \theta]} = \lambda$.

Remark. 1) We may consider using a μ^+ -free family $\bar{f}$ (see §2).
 2) Actually we use less than $T_J^1(\bar{\lambda}) = \lambda$, we just use

- (a) there are $f_\alpha \in \prod_{i < \kappa} \lambda_i$ for $\alpha < \lambda$ such that $\alpha < \beta \Rightarrow f_\alpha \neq_J f_\beta$
- (b) there are $f_\alpha \in \prod_{i < \kappa} \lambda_i$ for $\alpha < \lambda$ such that for every $f \in \prod_{i < \kappa} \lambda_i$ for some α , $\neg(f \neq_J f_\alpha)$.

Proof. Let $\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle$ be pairwise J -different, $f_\alpha \in \prod_{i < \kappa} \lambda_i$ (i.e. $\alpha \neq \beta \Rightarrow \{i : f_\alpha(i) \neq f_\beta(i)\} \in J$).

For each $i < \kappa$ let $\mathcal{P}_i \subseteq [\lambda_i]^{\leq \theta}$ be of cardinality λ_i and witness $\lambda_i^{[\partial, \theta]} = \lambda_i$ that is: every $u \in [\lambda_i]^{\leq \theta}$ is the union of $< \partial$ members of $\mathcal{P}_i$; such family exists by assumption (e). Let $M \prec (\mathcal{H}(\chi), \in)$ be of cardinality λ such that $\lambda + 1 \subseteq M$ and $\bar{f}, \langle \lambda_i : i < \kappa \rangle, \langle \mathcal{P}_i : i < \kappa \rangle, J, \mathcal{P}(\kappa)$ belong to M .

Let $\mathcal{P} = M \cap [\lambda]^{\leq \theta}$. We shall show that $\mathcal{P}$ exemplifies the desired conclusion. Now $\mathcal{P}$ is a family of $\leq \|M\| = \lambda$ subsets of λ each of cardinality $\leq \theta$ hence it is enough to show

- (*) if $u \in [\lambda]^{\leq \theta}$ then u is included in the union of $< \partial$ members of $\mathcal{P}$ (or equal to; equivalent here as $2^\theta \leq \lambda$ hence $u_1 \subseteq u_2 \in \mathcal{P} \Rightarrow u_1 \in \mathcal{P}$).

Proof of ().* Let $u_i = \{f_\alpha(i) : \alpha \in u\}$; so $u_i \in [\lambda_i]^{\leq \theta}$, hence we can find $\langle v_{i,j} : j < j_i \rangle$ such that $v_{i,j} \in \mathcal{P}_i$ and $u_i = \cup \{v_{i,j} : j < j_i\}$ and $0 < j_i < \partial$. For each $\eta \in \prod_{i < \kappa} j_i$

let

$$w_\eta = \{\alpha \in u : i < \kappa \Rightarrow f_\alpha(i) \in v_{i, \eta(i)}\}.$$

Clearly $u = \cup\{w_\eta : \eta \in \prod_{i < \kappa} j_i\}$ as for any $\alpha \in u$ for each $i < \kappa$ we can define $\varepsilon_i(\alpha) < j_i$ such that $f_\alpha(i) \in v_{i, \varepsilon_i(\alpha)}$ and let $\eta_\alpha = \langle \varepsilon_i(\alpha) : i < \kappa \rangle$, clearly $\eta_\alpha \in \prod_{i < \kappa} j_i$ and so $\alpha \in w_{\eta_\alpha}$. By the assumption (f) as $i < \kappa \Rightarrow j_i < \partial$ clearly $|\prod_{i < \kappa} j_i| < \partial$ hence it is enough to prove that $\eta \in \prod_{i < \kappa} j_i \Rightarrow w_\eta \in \mathcal{P}$. As $u \in M \wedge |u| \leq \theta \Rightarrow \mathcal{P}(u) \subseteq M$ it is enough to prove for $\eta \in \prod_{i < \kappa} j_i$, that

⊛ w_η is included in some $w \in M \cap [\lambda]^{\leq \theta}$.

Proof of ⊛: As $i < \kappa \Rightarrow |\mathcal{P}_i| = \lambda_i$ and $T_J(\bar{\lambda}) = \lambda$ by 0.5 there is $\mathcal{G} \subseteq \{g : g \in \prod_{i < \kappa} \mathcal{P}_i\}$ satisfying $|\mathcal{G}| = \lambda$ and $(\forall g \in \prod_{i < \kappa} \mathcal{P}_i)(\exists g' \in \mathcal{G})(\{i : g(i) = g'(i)\} \in J^+)$. As $\langle \mathcal{P}_i : i < \kappa \rangle \in M$ without loss of generality $\mathcal{G} \in M$ and as $\lambda + 1 \subseteq M$ we have $\mathcal{G} \subseteq M$. Apply the choice of $\mathcal{G}$ to $\langle v_{i, \eta(i)} : i < \kappa \rangle \in \prod_{i < \kappa} \mathcal{P}_i$, so for some $g \in \mathcal{G} \subseteq M$ the set $B =: \{i < \kappa : v_{i, \eta(i)} = g(i)\}$ belongs to J^+ . Clearly $B \in M$ (as $B \subseteq \kappa$, $\mathcal{P}(\kappa) \in M$ and $|\mathcal{P}(\kappa)| \leq 2^\kappa \leq \theta^\kappa \leq \lambda \subseteq M$) hence $\langle v_{i, \eta(i)} : i \in B \rangle \in M$ hence $w = \{\alpha < \lambda : \text{for every } i \in B \text{ we have } f_\alpha(i) \in v_{i, \eta(i)}\}$ belongs to M . Now $|w| \leq \prod_{i \in B} |v_{i, \eta(i)}| \leq \theta^\kappa = \theta$ because $\alpha < \beta < \lambda \Rightarrow f_\alpha \neq_J f_\beta \Rightarrow f_\alpha \restriction B \neq f_\beta \restriction B$. Lastly $w_\eta \subseteq w$ as $\alpha \in w_\eta$ & $i < \kappa \Rightarrow f_\alpha(i) \in v_{i, \eta(i)}$, so we are done. $\square_{1.4}$

Remark. We could have used $\mathcal{F} = \{\alpha < \lambda : \{i : f_\alpha(i) \in v_{i, \eta(i)}\} \in J^+\}$.

To make this section free of quoting the pcf theorem we use the following definition.

1.5 Definition/Observation. 1) For a set $\mathfrak{a}$ of regular cardinals and $\sigma = \text{cf}(\sigma) \leq \text{cf}(\lambda)$ let

$$J_\lambda^\sigma[\mathfrak{a}] = \{\mathfrak{b} \subseteq \mathfrak{a} : \text{there is a set } \mathcal{F} \subseteq \pi \mathfrak{b} \text{ of cardinality } < \lambda \\ \text{such that for every } g \in \pi \mathfrak{b} \text{ we can find } j < \sigma \text{ and} \\ f_i \in \mathcal{F} \text{ for } i < j \text{ satisfying } \theta \in \mathfrak{b} \Rightarrow (\exists i < j)(g(\theta) < f_i(\theta))\}.$$

2) Clearly $J_\lambda^\sigma[\mathfrak{a}]$ is a σ -complete ideal on $\mathfrak{a}$ but possibly $\mathfrak{a} \in J_\lambda^\sigma[\mathfrak{a}]$.

Remark. In fact, if $\text{Min}(\mathfrak{a}) > |\mathfrak{a}|$, $J_\lambda^\sigma[\mathfrak{a}] = \{\mathfrak{b} \subseteq \mathfrak{a} : \text{pcf}_{\sigma\text{-complete}}(\mathfrak{b}) \subseteq \lambda\} = \{b \subseteq \mathfrak{a} : \mathfrak{b} \text{ is the union of } < \sigma \text{ members of } J_\lambda[\mathfrak{a}]\}$ can be proved but this is irrelevant here.

For completeness we recall and prove then we know

1.6 Claim. $\lambda = \lambda^{[\sigma, < \theta]}$ when

- (a) $\lambda \geq 2^{< \theta} \geq \sigma = \text{cf}(\sigma) > \aleph_0$ and $\text{cf}(\theta) \notin [\sigma, \theta]$
- (b) for every set $\mathfrak{a} \subseteq \text{Reg} \cap \lambda \setminus \theta$ of cardinality $< \theta$ we have $\mathfrak{a} \in J_\lambda^\sigma[\mathfrak{a}]$.

Proof. Let χ be large enough, choose $M \prec (\mathcal{H}(\chi), \in, <_\chi^*)$ of cardinality λ where $<_\chi^*$ is any well ordering of $\mathcal{H}(\chi)$ such that $\lambda + 1 \subseteq M$ and let $\mathcal{P} = M \cap [\lambda]^{< \theta}$ and we shall prove that $\mathcal{P}$ exemplifies $\lambda = \lambda^{[\sigma, < \theta]}$.

Clearly $\mathcal{P} \subseteq [\lambda]^{< \theta}$ has cardinality λ so let $u \in [\lambda]^{< \theta}$ and as $2^{< \theta} \leq \lambda$ it is enough to show that u is included in a union of $< \sigma$ members of $\mathcal{P}$ thus finishing.

Let f be a one-to-one function from $\kappa =: |u|$ onto u so $\kappa < \theta$. By induction on n we choose $f_n, \bar{v}_n$ such that

- ⊗ (a) f_n is a function from κ to $\lambda + 1$
- (b) $\bar{v}_n = \langle v_{n,\varepsilon} : \varepsilon < \varepsilon_n \rangle$ is a partition of κ which satisfies
 $\varepsilon_n < \sigma$ and $\kappa = \bigcup \{v_{n,\varepsilon} : \varepsilon < \varepsilon_n\}$
- (c) $f_0(i) = \lambda$ for every $i < \kappa$
- (d) $f_{n+1}(i) \leq f_n(i)$ for $i < \kappa$
- (e) $f(i) \leq f_n(i)$ and if $f(i) < f_n(i)$ then $f_{n+1}(i) < f_n(i)$
- (f) $f_n \upharpoonright v_{n,\varepsilon} \in M$.

This is sufficient: $\{\text{Rang}(f_n \upharpoonright v_{n,\varepsilon}) : n < \omega, \varepsilon < \varepsilon_n\}$ is a family of $< \sigma$ sets (as $\sigma = \text{cf}(\sigma) > \aleph_0, \sigma > \varepsilon_n$) each belonging to $\mathcal{P}$ (as $f_n \upharpoonright v_{n,\varepsilon} \in M$) and their union includes u because for every $i < \kappa$, $f_n(i) = f(i)$ for every n large enough (by clauses (d) + (e) of ⊗).

For $n = 0$, f_n is constantly λ . So let $n = m + 1$, f_m be given, let

$$u_{n,0} = \{i < \kappa : f_m(i) = f(i)\}$$

$$u_{n,1} = \{i < \kappa : f_m(i) > f(i) \text{ and } i \text{ is a successor ordinal or just has cofinality } < \theta\}$$

$$u_{n,2} = \kappa \setminus u_{n,0} \setminus u_{n,1}.$$

As $2^\kappa \leq 2^{<\theta} \leq \lambda$, clearly the partition $\langle u_{n,0}, u_{n,1}, u_{n,2} \rangle$ of κ belongs to M , so it is enough to choose $f_{n+1} \upharpoonright u_{n,\ell}$ separately for $\ell = 0, 1, 2$.

Let $f_n \upharpoonright u_{n,0} = f_m \upharpoonright u_{n,0}$.

Let $\bar{C} = \langle C_\alpha : \alpha \leq \lambda \rangle \in M$ be such that $C_0 = \emptyset, C_{\alpha+1} = \{\alpha\}, C_\delta$ is a club of δ of order type $\text{cf}(\delta)$ for limit ordinal $\delta \leq \lambda$. Let $f_n \upharpoonright u_{n,1}$ be defined by $f_n(i) = \text{Min}(C_{f_m(i)} \setminus f(i))$. For each $\varepsilon < \varepsilon_m$ the function $f_n \upharpoonright (u_{n,1} \cap v_{m,\varepsilon})$ belongs to M hence $\langle C_{f_m(i)} : i \in u_{n,1} \cap v_{m,\varepsilon} \rangle$ belongs to M , but the product $\prod_{i \in u_{n,1} \cap v_{m,\varepsilon}} C_{f_m(i)}$

has cardinality $\leq \theta^\kappa \leq 2^{<\theta} \leq \lambda$ if $\text{cf}(\theta) \geq \kappa$ and without loss of generality $\theta_1 = \sup\{|C_{f_m(i)}| : i \in u_{n,1} \cap v_{m,\varepsilon}\} < \theta$ hence has cardinality $\leq 2^{\theta_1+\kappa} \leq 2^{<\theta} \leq \lambda$ if $\text{cf}(\theta) < \sigma$.

Lastly, it is enough to define $f_n \upharpoonright (v_{m,\varepsilon} \cap u_{n,2})$ for each $\varepsilon < \varepsilon_m$. Let $\lambda_{n,i} = \text{cf}(f_m(i))$, so $\langle \lambda_{n,\zeta} : \zeta \in v_{m,\varepsilon} \cap u_{n,2} \rangle \in M$ hence there is a sequence $\langle h_{n,\zeta} : \zeta \in v_{m,\varepsilon} \cap u_{n,2} \rangle \in M$ where $h_{n,\zeta}$ is increasing continuous function from $\lambda_{n,\zeta}$ onto some club of $f_m(i)$.

Let $\mathfrak{a} = \{\lambda_{n,\zeta} : \zeta \in v_{m,\varepsilon} \cap u_{n,2}\}$. Applying assumption (b), Definition 1.5(1) it is easy to finish. $\square_{1.6}$

1.7 Claim. *There is $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ and σ -complete ideal J on κ such that $T_J(\bar{\lambda}) \geq \lambda$ and $i < \kappa \Rightarrow 2^\kappa < \lambda_i < \lambda$ when*

- ⊗ (a) $2^\kappa < \lambda, \aleph_0 < \sigma = \text{cf}(\sigma) \leq \kappa$
- (b) $\mathfrak{a} \subseteq \text{Reg} \cap \lambda \setminus (2^\kappa)^+$ has cardinality $\leq \kappa$ and $\mathfrak{a} \notin J_\lambda^\sigma[\mathfrak{a}]$.

Proof. Let $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ list $\mathfrak{a}$ and let $J = J_\lambda^\sigma[\mathfrak{a}]$ and by induction on $\alpha < \lambda$ a function $f_\alpha \in \pi(\mathfrak{a})$ such that $\beta < \alpha \Rightarrow f_\beta <_J f_\alpha$. Arriving to α for every $\mathfrak{b} \subseteq \mathfrak{a}$ let $\mathcal{F}_\mathfrak{b}^\alpha = \{f_\beta \upharpoonright \mathfrak{b} : \beta < \alpha\}$, so by the definition of $J_\lambda^\sigma[\mathfrak{a}]$, for every $\mathfrak{b} \in J^+ := \mathcal{P}(\mathfrak{a}) \setminus J$, $\mathcal{F}_\mathfrak{b}^\alpha$ does not witness $\mathfrak{b} \in J_\lambda^\sigma[\mathfrak{a}]$ hence there is $g_\mathfrak{b}^\alpha \in \pi(\mathfrak{b})$ witnessing it. Let $f_\alpha \in \pi(\mathfrak{a})$ be defined by $f_\alpha(\theta) = \sup\{g_\mathfrak{b}^\alpha(\theta) : \mathfrak{b} \in J^+ \text{ and } \theta \in \mathfrak{b}\}$. Now $f_\alpha \in \Pi \mathfrak{a}$ as $\theta \in \mathfrak{a} \Rightarrow f_\alpha(\theta) < \theta$ which holds as $|J^+| \leq 2^{|\mathfrak{a}|} \leq 2^\kappa < \theta$. Also if $\beta < \alpha$ and we let $\mathfrak{b}_\beta^\alpha = \{\theta \in \mathfrak{a} : f_\beta(\theta) \geq f_\alpha(\theta)\}$ then $\mathfrak{b}_\beta^\alpha \in J^+$ implies easy contradiction to the choice of $g_{\mathfrak{b}_\beta^\alpha}^\alpha$ (and f_α). So we can carry the induction and so $\langle f_\alpha : \alpha < \lambda \rangle, f_\alpha \in \pi(\bar{\lambda})$ where $f'_\alpha(i) = f_\alpha(\lambda_i)$ exemplify $T_J(\bar{\lambda}) \geq \lambda$ as required. $\square_{1.7}$

Remark. This is the case $\text{Min}(\mathfrak{a}) > 2^{|\mathfrak{a}|}$ from [Sh:b, XIII).

1.8 Claim. *If $\circledast$ from 1.7 then there is $\bar{\lambda}' = \langle \lambda'_i : i < \kappa \rangle$ such that*

- (α) $2^\kappa < \lambda'_i \leq \lambda_i$
- (β) *if $f \in \prod_{i < \kappa} \lambda_i$ then $T_J(f) < \lambda$*
- (γ) $T_J(\bar{\lambda}') = \lambda$

where

- $\circledast$ (a) $2^\kappa < \lambda, \aleph_0 < \sigma = \text{cf}(\sigma) \leq \kappa$
- (b) $2^\kappa < \lambda_i < \lambda$
- (c) J is a σ -complete ideal on κ
- (d) $T_J(\bar{\lambda}) \geq \lambda$.

Proof. Clearly $\{i : \lambda_i \leq (2^\kappa)^{+n}\} \in J$ for $n < \omega$ (as $((2^\kappa)^{+n})^\kappa = (2^\kappa)^{+n}$) so by 0.5(1) without loss of generality $i < \kappa \Rightarrow \lambda_i > (2^\kappa)^{+2}$.

As $(\prod_{i < \kappa} \lambda_i^+, <_J)$ is well founded and there is $f \in \prod_{i < \kappa} (\lambda_i + 1)$ satisfying $T_J(f) \geq \lambda$

(i.e. $\bar{\lambda}$ itself) clearly there is $f \in \prod_{i < \kappa} (\lambda_i + 1)$ for which $T_J(f) \geq \lambda$ satisfying

$g \in \prod_{i < \kappa} (\lambda_i + 1), g <_J f$ implies $T_D(g) < \lambda$. Now easily $\{i < \kappa : f(i) \leq (2^\kappa)^{+2}\} \in J$,

so without loss of generality $i < \kappa \Rightarrow f(i) > (2^\kappa)^{+2}$. Let $\lambda'_i = |f(i)|$, hence $\bar{\lambda}'$ satisfies demands (α) + (β) of the desired conclusion, i.e., and $T_J(\bar{\lambda}') = T_J(f) \geq \lambda$. So assume toward contradiction that it fails clause (γ), so by the last sentence we have $T_J(\bar{\lambda}') > \lambda$ and we shall derive a contradiction thus finishing. So there is $\{f_\alpha : \alpha < \lambda^+\} \subseteq \prod_{i < \kappa} \lambda'_i$ such that $\alpha \neq \beta \Rightarrow f_\alpha \neq_J f_\beta$ and let $u_\alpha = \{\beta : f_\beta <_J f_\alpha\}$.

If for some $\alpha < \lambda, |u_\alpha| \geq \lambda$ then $\{f_\beta : \beta \in u_\alpha\}$ exemplifies that $T_D(f_\alpha) \geq \lambda$ and clearly $f_\alpha <_J \bar{\lambda}' \leq f$, contradiction to the choice of f . So $\alpha < \lambda^+ \Rightarrow |u_\alpha| < \lambda$. Hence there is $S \subseteq \lambda$ such that $(\forall \alpha \neq \beta \in S)(\beta \notin u_\alpha)$ contradicting 0.9. $\square_{1.8}$

1.9 The revised GCH Theorem. *If θ is strong limit singular then for every $\lambda \geq \theta$ for some $\sigma < \theta$ we have $\lambda = \lambda^{[\sigma, \theta]}$.*

Remark. 1) Hence for every $\lambda \geq \theta$ for some $n < \omega$ and $\kappa_\ell < \theta (\ell \leq n), \aleph_0 = \kappa_0 < \kappa_1 < \dots < \kappa_n = \theta$, for each $\ell < n, 2^{\kappa_\ell} \geq \kappa_{\ell+1}$ or $\lambda = \lambda^{[\kappa'_\ell, < \kappa_{\ell+1}]}$ where $\kappa'_\ell = (2^{\kappa_\ell})^+$.
2) If $\sigma \in (\text{cf}(\theta), \theta)$ and $\lambda \geq \theta$ then $\lambda^{[\sigma, \theta]} = \lambda^{[\sigma, < \theta]} = \Sigma\{\lambda^{[\sigma, \theta']} : \theta' \in [\sigma, \theta]\}$.

Proof. We prove by induction on $\lambda \geq \theta$.

Let $\sigma =: (\text{cf}(\theta))^+ < \theta$.

Case 0: $\lambda = \theta$.

Let $\mathcal{P}$ be the family of bounded subsets of θ , so $|\mathcal{P}| = \theta$ and every $u \in [\theta]^{<\theta}$ is the union of $\text{cf}(\theta)$ members of $\mathcal{P}$.

Case 1: For every $\mathfrak{a} \subseteq \text{Reg} \cap \lambda^+ \setminus \theta$ of cardinality $< \theta$ we have $\text{pcf}_{\sigma\text{-comp}}(\mathfrak{a}) \subseteq \lambda^+$. By 1.6, we have $\lambda^{[\sigma, <\theta]} = \lambda$ (recalling $\text{cf}(\theta) < \sigma$ and 1.3).

Case 2: Neither Case 0 nor Case 1.

As not Case 1, the assumption of Claim 1.7 holds for some κ for which $\sigma \leq \kappa < \theta$, hence its conclusion holds for some $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ and J , i.e., we have $2^\kappa < \lambda_i < \lambda$ and $T_J(\bar{\lambda}) \geq \lambda$ where J is a σ -complete ideal on κ . So the assumptions of Claim 1.8 holds, hence its conclusion so by 1.8

- ⊗ (i) J is a σ -complete ideal on κ
- (ii) $\bar{\lambda}' = \langle \lambda'_i : i < \kappa \rangle$
- (iii) $2^\kappa < \lambda'_i < \lambda$
- (iv) $T_J(\bar{\lambda}') = \lambda$
- (v) $T_J(f) < \lambda$ if $f \in \prod_{i < \kappa} \lambda'_i$.

We can find an increasing sequence $\langle \theta_\varepsilon : \varepsilon < \text{cf}(\theta) \rangle$ of regular cardinals from (σ, θ) with limit θ . By the induction hypothesis for each $i < \lambda$ there is $\varepsilon(i)$ such that $\lambda_i = \lambda_i^{[\theta_{\varepsilon(i)}, <\theta]} \geq \theta$ or $\lambda_i \leq \theta_{\varepsilon(i)}$. For $\zeta < \text{cf}(\theta)$ define $A_\zeta = \{i < \kappa : \lambda_i \geq \theta, \varepsilon(i) = \zeta\}$ and $A_{\text{cf}(\theta)+\zeta} = \{i < \kappa : \lambda_i < \theta \text{ and } \varepsilon(i) = \zeta\}$. So $\langle A_\varepsilon : \varepsilon < \text{cf}(\theta) + \text{cf}(\theta) \rangle$ is a partition of κ to $< \sigma$ sets hence by 0.5(4) we know that

$$T_J^1(\bar{\lambda}') = \text{Min}\{T_{J \upharpoonright A_\varepsilon}^1(\bar{\lambda}' \upharpoonright A_\varepsilon) : \varepsilon < \text{cf}(\theta) + \text{cf}(\theta) \text{ and } A_\varepsilon \in J^+\}$$

hence for some $\zeta < \text{cf}(\theta) + \text{cf}(\theta)$, $T_J(\bar{\lambda}') = T_{J \upharpoonright A_\zeta}(\bar{\lambda}' \upharpoonright A_\zeta)$ and $A_\zeta \in J^+$, so by renaming without loss of generality $A_\zeta = \kappa$. If $\zeta \geq \text{cf}(\theta)$ as $\kappa < \theta$, θ strong limit we get $T_J(\bar{\lambda}') \leq \prod_{i < \kappa} \lambda'_i < (\theta_\zeta)^\kappa < \theta$, a contradiction so $\zeta < \text{cf}(\theta)$. Now apply 1.4 with $J, \bar{\lambda}', \sigma, \kappa, (2^\kappa)^+, \theta_\zeta$ here standing for $J, \bar{\lambda}, \sigma, \kappa, \partial, \theta$ and get the desired result.

□_{1.9}

1.10 Concluding Remarks. 1) We can in 1.4 assume less. Instead $\theta = \theta^\kappa$, it is enough (which follows to [Sh 506, §3], see 0.4)

⊗ for every $\lambda' < \lambda$ we can find $\mathcal{F} \subseteq \prod_{i < \kappa} \lambda_i$ of cardinality λ' such that $f \neq g \in \mathcal{F} \Rightarrow f \neq_J g$.

This is seemingly a gain, but in the induction the case $(\forall \mathfrak{a} \subseteq \text{Reg} \cap \lambda^+ \setminus \theta)(|\mathfrak{a}| \leq \kappa \Rightarrow \text{pcf}_{\aleph_1\text{-comp}}(\mathfrak{a}) \subset \lambda^+)$ is problematic.

§2 THE FINITELY MANY EXCEPTIONS

We can prove a relative of 1.4 assuming $i < \kappa \Rightarrow \lambda_i^{[\sigma, \kappa, \theta]} = \lambda_i$, replacing “ $\theta = \theta^\kappa + 2^\theta \leq \lambda$ by $2^\kappa \leq \lambda$ and getting $\lambda^{[\sigma, \kappa, \theta]} = \lambda$. But so far it has no conclusion parallel to 1.9.

In full:

2.1 Claim. *Assume*

- (a) $\aleph_0 < \sigma = \text{cf}(\sigma) \leq \theta$ and $\mu \leq \theta$
- (b) J is a σ -complete ideal on κ
- (c) $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$
- (d) $T_J(\bar{\lambda}) = \lambda$, moreover this is exemplified by a μ^{++} -free family
- (e) $\lambda_i^{<\partial, \mu, \theta>} = \lambda_i$ for $i < \kappa$
- (f) if $\partial_i < \partial$ for $i < \kappa$ then $\prod_{i < \kappa} \partial_i < \partial$
- (g) $\theta = \theta^\kappa$ and $2^\kappa \leq \lambda$.

Then $\lambda^{<\partial, \mu, \theta>} = \lambda$.

Proof. Let $\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle$ be θ^+ -free, $f_\alpha \in \prod_{i < \kappa} \lambda_i$ pairwise J -different, (i.e., $\alpha \neq \beta \Rightarrow \{i : f_\alpha(i) = f_\beta(i)\} \in J$) exists by clause (d) of the assumption.

For each $i < \kappa$ let $\mathcal{P}_i \subseteq [\lambda_i]^{\leq \mu}$ be of cardinality λ_i and witness $\lambda_i^{<\sigma, \mu, \theta>} = \lambda_i$ that is: every $u \in [\lambda_i]^{\leq \theta}$ is the union of $< \sigma$ members of

$\text{set}_{\theta, \mu}(\mathcal{P}_i) =: \{v : v \in [\lambda_i]^{\leq \mu} \text{ and every } w \in [v]^{\leq \mu} \text{ is included in some member of } \mathcal{P}_i\}$;

such family exists by assumption (e). Let $M \prec (\mathcal{H}(\chi), \in)$ be of cardinality λ such that $\lambda + 1 \subseteq M$ and $\langle \lambda_i : i < \kappa \rangle, \langle \mathcal{P}_i : i < \kappa \rangle, J, \mathcal{P}(\kappa)$ belong to M .

Let $\mathcal{P} = M \cap [\lambda]^{\leq \mu}$. We shall show that $\mathcal{P}$ exemplifies the desired conclusion. Now $\mathcal{P}$ is a family of $\leq \|M\| = \lambda$ of subsets of λ each of cardinality $\leq \mu$ hence it is enough to show

- (*) if $u \in [\lambda]^{\leq \mu}$ then u is included in the union of $< \partial$ sets $v \in \text{set}_{\theta, \mu}(\mathcal{P}_i)$.

Proof of ().* Let $u_i = \{f_\alpha(i) : \alpha \in u\}$; so $u_i \in [\lambda_i]^{\leq \theta}$, hence we can find $\langle v_{i,j} : j < j_i \rangle$ such that $[v_{i,j}]^{\leq \mu} \in \text{set}_{\theta,\mu}(\mathcal{P}_i)$ and $u_i = \cup\{u_{i,j} : j < j_i\}$ and $0 < j_i < \partial$. For each $\eta \in \prod_{i < \kappa} j_i$ let

$$w_\eta = \{\alpha \in u : i < \kappa \Rightarrow f_\alpha(i) \in v_{i,\eta(i)}\}.$$

Clearly $u = \cup\{w_\eta : \eta \in \prod_{i < \kappa} j_i\}$ as for any $\alpha \in u$ for each $i < \kappa$ we can choose $\varepsilon_i(\alpha) < j$ such that $f_\alpha(i) \in v_{i,\varepsilon_i(\alpha)}$ and let $\eta_\alpha = \langle \varepsilon_i(\alpha) : i < \kappa \rangle$ clearly $\eta_\alpha \in \prod_{i < \kappa} j_i$ and $\alpha \in w_{\eta_\alpha}$. By the assumption (f) as $i < \kappa \Rightarrow j_i < \partial$ clearly $|\prod_{i < \kappa} j_i| < \sigma$ hence it is enough to prove that $\eta \in \prod_{i < \kappa} j_i \Rightarrow w_\eta \in \text{set}_{\theta,\mu}(\mathcal{P})$. So it is enough to prove for $\eta \in \prod_{i < \kappa} j_i$ and $w \in [w_\eta]^{\leq \mu}$ that

⊗ w is included in some $w' \in M \cap [\lambda]^{\leq \mu}$.

Proof of ⊗: As $i < \kappa \Rightarrow |\mathcal{P}_i| = \lambda_i$ and $T_J(\bar{\lambda}) = \lambda$ there is $\mathcal{G} \subseteq \{g : g \in \prod_{i < \kappa} \mathcal{P}_i\}$ satisfying $|\mathcal{G}| = \lambda$ and $(\forall g \in \prod_{i < \kappa} \mathcal{P}_i)(\exists g' \in \mathcal{G})(\{i : g(i) = g'(i)\} \in J^+)$. As $\langle \mathcal{P}_i : i < \kappa \rangle \in M$ without loss of generality $\mathcal{G} \in M$ and as $\lambda + 1 \subseteq M$ we have $\mathcal{G} \subseteq M$. For each $i < \kappa$ we have $A_i = \{f_\alpha(i) : \alpha \in w\}$ is a subset of some $A'_i, \mathcal{P}_i$. Apply the choice of $\mathcal{G}$ to $\langle A'_i : i < \kappa \rangle \in \prod_{i < \kappa} \mathcal{P}_i$, so for some $g \in \mathcal{G} \subseteq M$ the set $B = \{i : A'_i = g(i)\}$ belongs to J^+ . Clearly $B \in M$ (as $2^\kappa \leq \theta^\kappa \leq \lambda$) hence $\langle A'_i : i \in B \rangle \in M$ hence $w' = \{\alpha < \lambda : \text{for some } Y \in J^+ \text{ for every } i \in Y \text{ we have } f_\alpha(i) \in A'_i\}$ belongs to M . Now $|w| \leq \mu^\kappa$; as $\alpha < \beta < \lambda \Rightarrow f_\alpha \neq_J f_\beta$ but as $\bar{f}$ is μ^{++} -free we even have $|w| \leq \mu$. Lastly $w \subseteq \mathcal{F}$ as $\alpha \in w$ & $i < \kappa \Rightarrow f(i) \in A_i$, so we are done. $\square_{2.1}$

2.2 Claim. If $\theta > \sigma = \text{cf}(\sigma) > \aleph_0$ and $\lambda > \theta_* = 2^{<\theta}$ then there is $\langle (\lambda_\eta, \mathcal{D}_\eta, D_\eta, \kappa_\eta) : \eta \in \mathcal{T} \rangle$ such that

- (a) $\mathcal{T}$ is a subtree of ${}^\omega > \theta$ (i.e. $\langle \rangle \in \mathcal{T} \subseteq {}^\omega > \theta$, $\mathcal{T}$ is closed under initial segments) with no ω -branch, let $\text{max}_{\mathcal{T}}$ be the set of maximal nodes of $\mathcal{T}$
- (b) λ_η is a cardinal $\in (2^{<\theta}, \lambda]$ and $\nu \triangleleft \eta \Rightarrow \lambda_\nu > \lambda_\eta$ and $\lambda_{\langle \rangle} = \lambda$
- (c) κ_η is a regular cardinal $\in [\sigma, \theta)$ if $\eta \in \mathcal{T} \setminus \text{max}_{\mathcal{T}}$ and κ_η is zero or undefined if $\eta \in \text{max}_{\mathcal{T}}$ and $\eta \cap \langle \alpha \rangle \in \mathcal{T} \Leftrightarrow \alpha < \kappa_\eta$

- (d) if $\eta \in \max_{\mathcal{T}}$ then for no $\kappa < \theta$ and σ -complete filter $\mathcal{D}$ on κ and cardinals $\lambda_i \in (2^{<\theta}, \lambda_\eta)$ for $i < \kappa$ do we have $T_{\mathcal{D}}(\langle \lambda_i : i < \kappa \rangle) \geq \lambda$
- (e) $\mathcal{D}_\eta$ is a σ -complete filter on κ_η when $\eta \in \mathcal{T} \setminus \max_{\mathcal{T}}$
- (f) $T_{\mathcal{D}_\eta}(\langle \lambda_{\eta \frown \langle \alpha \rangle} : \alpha < \kappa_\eta \rangle) = \lambda_\eta$ if $\eta \in \mathcal{T} \setminus \max_{\mathcal{T}}$
- (g) if $f \in \prod_{\alpha < \kappa_\eta} \lambda_{\eta \frown \langle \alpha \rangle}$ then $T_{\mathcal{D}_\eta}(f) < \lambda_\eta$
- (h) D_η is the σ -complete filter on $\max_{\mathcal{T}, \eta} = \{\nu \in \max_{\mathcal{T}} : \eta \leq \nu\}$ such that
 - (α) if $\eta \in \max_{\mathcal{T}}$, $D_\eta = \{\{\eta\}\}$
 - (β) if $\eta \in \mathcal{T} \setminus \max_{\mathcal{T}}$ then

$$D_\eta = \{A \subseteq \max_{\mathcal{T}, \eta} : \text{the following set belongs to } \mathcal{D}_\eta \\ \{\alpha < \kappa_\eta : A \cap \max_{\mathcal{T}, \eta \frown \langle \alpha \rangle} \in D_{\eta \frown \langle \alpha \rangle}\}\}$$

- (i) (α) if $\text{cf}(\lambda) > \theta_*$ then $\eta \in \mathcal{T} \Rightarrow \text{cf}(\lambda_\eta) > \theta_*$
- (β) we can replace $\theta_* = 2^{<\theta}$ by any cardinal such that $\text{cf}(\theta_*) \geq \theta \wedge (\forall \sigma < \sigma)(\forall \alpha < \theta_*)|\alpha|^\sigma < \theta_*$
- (γ) if λ is regular then every λ_η is regular [FILL proof].

Proof.

Case 1: Ignoring clause (i).

We prove this by induction $\lambda > 2^{<\theta}$. If λ satisfies the requirement in clause (d) let $\mathcal{T} = \{\langle \rangle\}$, $\lambda_\eta = \lambda$ and $\kappa_{\langle \rangle}, D_{\langle \rangle}$ are trivial. If λ fails that demand use claims 1.7 + 1.8 to find $\mathcal{D}, \kappa, \bar{\lambda}$ such that

$$(*) \quad \kappa \in [\sigma, 2^{<\theta}], \mathcal{D} \text{ is a } \sigma\text{-complete filter on } \kappa, \bar{\lambda} = \langle \lambda_\alpha : \alpha < \kappa \rangle \text{ and } \lambda_\alpha \in [\kappa, 2^{<\theta}], \text{ a cardinal } T_{\mathcal{D}}(\bar{\lambda}) = \lambda \text{ but } f \in \prod_{\alpha < \kappa} \lambda_\alpha \Rightarrow T_{\mathcal{D}}(f) < \lambda.$$

Now for each $\alpha < \kappa$ we can use the induction hypothesis to find $\langle (\lambda_\eta^\alpha : \mathcal{D}_\eta^\alpha, \mathcal{D}_\eta^\alpha) : \eta \in \mathcal{T}_\alpha \rangle$ as required in the claim for λ_α . Now we let:

- ⊗ (a) $\mathcal{T} = \{\langle \rangle\} \cup \{\langle \alpha \rangle \frown \eta : \eta \in \mathcal{T}_\alpha\}$
- (b) $\lambda_{\langle \rangle}^0 = \lambda, \kappa_{\langle \rangle} = \kappa$
- (c) $\lambda_{\langle \alpha \rangle \frown \eta} = \lambda_\eta^\alpha$ for $\alpha < \kappa, \eta \in \mathcal{T}_\alpha$
- (d) $\mathcal{D}_{\langle \rangle} = \mathcal{D}$

- (e) $\mathcal{D}_{\langle \alpha \rangle \frown \eta} = \mathcal{D}_\eta^\alpha$ for $\alpha < \kappa, \eta \in \mathcal{T}_\alpha$
- (f) $D_{<} = \{A : A \subseteq \max \mathcal{T}_{<} \text{ and } \alpha < \kappa : \{\eta : \langle \alpha \rangle \frown \eta \in A \cap \max \mathcal{T}_{\alpha, <}\} \text{ belongs to } \mathcal{D}\}$
- (g) $D_{\langle \alpha \rangle \frown \eta} = \{\langle \alpha \rangle \frown \nu : \nu \in B\} : B \in D_\eta^\alpha\}$.

Easily they are as required.

Case 2: Proving the claim with (i)(α), so dealing with λ satisfying $\text{cf}(\lambda) > \theta_x$.

If λ satisfies the requirement in clause (a) we finish as above. Otherwise, we can find $\kappa \in [\sigma, \theta), D, \bar{\lambda}$ such that

- (*) σ -complete filter $\mathcal{D}$ on $\kappa, \bar{\lambda} = \langle \lambda_\alpha : \alpha < \kappa \rangle$ and $\lambda_\alpha \in (2^{<\theta}, \lambda)$ such that $\lambda = T_D(\langle \lambda_\alpha : \alpha < \kappa \rangle)$ and $f \in \prod_{\alpha < \kappa} \lambda_\alpha \Rightarrow T_D(f) < \lambda$.

Let $B = \{\alpha : \text{cf}(\lambda_\alpha) > \theta_*\}$. If $B \in \mathcal{D}^+$ and $T_{\mathcal{D} \upharpoonright B}(f \upharpoonright B) < \lambda$ for $f \in \Pi \bar{\lambda}$ (hence $T_{\mathcal{D} \upharpoonright B}(\bar{\lambda} \upharpoonright B)(\lambda_{\eta \frown \langle \varepsilon \rangle}) = \lambda$), then we can use $\bar{\lambda} \upharpoonright B, \mathcal{D} \upharpoonright B$ (and renaming), so we are done, so assume that this fails hence $\bar{\lambda}(\kappa \setminus B), \mathcal{D} \upharpoonright (\kappa \setminus B)$ are as required in (*), so by renaming without loss of generality $B = \emptyset$. For each $\alpha < \kappa$ let $\langle \lambda_{\alpha, \varepsilon} : \varepsilon < \text{cf}(\lambda_\alpha) \rangle$ be increasing continuous with limit λ_α and let $\bar{f} = \langle f_\alpha : \alpha < \lambda \rangle$ witness $T_{\mathcal{D}_\eta}(\bar{\lambda}) \geq \lambda$. For each $\alpha < \lambda$ for some $h_\alpha \in \prod_{\alpha < \kappa_\eta} \text{cf}(\lambda_\alpha), f_\alpha < \langle \lambda_{\alpha, h(\alpha)} : \alpha < \kappa \rangle$ so

for some $h, |\{\alpha : h_\alpha = h\}| = \lambda_\eta$, contradiction.

We finish as in case (1).

Case 3: Restricting ourselves to a regular λ . Similar to case to using 2.5(3) below.

$\square_{2.2}$

2.3 Definition. 1) We say that $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ is a D -representation of λ when:

- (a) D is a filter on κ
- (b) $T_D(\bar{\lambda}) = \lambda$
- (c) if $f \in \prod_{i < \kappa} \lambda_i$ then $T_D(f) < \lambda$.

2) We say that $\bar{\lambda}$ is an exact D -representation of λ when

- (a) D is a filter on κ
- (b) $T_{D+A}(\bar{\lambda}) = \lambda$ for $A \in D^+$
- (c) if $f \in \prod_{i < \lambda} \lambda_i$ and $A \in D^+$ then $T_{D+A}(f) < \lambda$.

3) We say that the representation is true when:

$$(d) \text{ cf}(\lambda) = \text{tcf}(\Pi \bar{\lambda}, <_D).$$

4) We can replace the filter by the dual ideal.

2.4 Definition. 1) We say $\langle (\lambda_\eta, \mathcal{D}_\eta, D_\eta, \kappa_\eta) : \eta \in \mathcal{T} \rangle$ is a representation if the conditions in Definition 2.2 holds.

2) We say it is an exact/true representation when each $\langle \lambda_{\eta \smallfrown \alpha} : \alpha < \kappa_\eta \rangle$ is exact/true a D_η -representation of λ_η .

2.5 Claim. 1) Assume

- ⊗ (a) $\bar{\lambda}^* = \langle \lambda_i : i < \kappa \rangle$ is a D_* -representation of λ
- (b) $\bar{\lambda}^i = \langle \lambda_{i,j} : j < \kappa_i \rangle$ is a D_i -representation of λ_i
- (c) D is $\Sigma_{D_*} \langle D_i : i < \kappa \rangle$, i.e., the filter on $u = \{(i, j) : i < \kappa, j < \kappa_i\}$ defined by $D = \{A^* \subseteq S : \{i : \{j < \kappa_i : (i, j) \in A\} \in D_i\} \in D_*\}$
- (d) $\text{cf}(\lambda), \text{cf}(\lambda_i)$ are $> |u|$ and $\lambda, \lambda_i, \lambda_{i,j}$ are $> 2^{|u|}$.

Then $\bar{\lambda} = \langle \lambda_{i,j} : (i, j) \in u \rangle$ is a D -representation of λ .

2) Similarly for exact representations, i.e., if in ⊗(a), (b) we further assume that the representations are exact then also $\bar{\lambda}$ an exact D -representation of λ .

3) Similarly for true representations: if $\lambda_i = \text{tcf}(\prod_{j < \kappa_i} \lambda_{i,j}, <_{D_i})$, $\lambda = \text{tcf}(\prod_{i < \kappa} \lambda_i, <_D)$

then $\lambda = \text{tcf}(\prod_{(i,j) \in u} \lambda_{i,j}, <_D)$. Similarly for min-cf, etc.

4) If D is an $\aleph_1$ -complete filter on κ , $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ and $T_D(\bar{\lambda}) > \lambda > 2^\kappa$ and $i < \kappa \Rightarrow \lambda_i > 2^\kappa$. Then we can find $\bar{\lambda}'$ such that $i < \kappa \Rightarrow 2^\kappa \leq \lambda'_i < \lambda_i$ and $\bar{\lambda}'$ is a D -representation of λ . If we demand only $T_D(\bar{\lambda}) \geq \lambda$ then we know only $\lambda'_i \leq \lambda_i$.

Proof. 1)

- (*)₁ $\lambda = T_D(\langle \lambda_{i,j} : (i, j) \in u \rangle)$
 [why? Let $\mathcal{G}^i = \{g_\alpha^i : \alpha < \lambda_i\}$ witness that $T_{D_i}(\bar{\lambda}^i) = \lambda_i$ and let $\mathcal{G}^* = \{g_\alpha^* : \alpha < \lambda\}$ witness that $T_{D_*}(\bar{\lambda}^*) = \lambda$. We now define $\mathcal{G} = \{g_\alpha : \alpha < \lambda\}$ where $g_\alpha \in \prod_{(i,j) \in u} \lambda_{i,j}$ is defined by $g_\alpha((i, j)) = g_{g_\alpha^*(i)}^i(j)$ and we can easily check that $\alpha < \beta < \lambda \Rightarrow g_\alpha \neq g_\beta \text{ mod } D$. Now if $g \in \prod_{(i,j) \in u} \lambda_{i,j}$ then for each i

the function (i.e. sequence) $\langle g((i, j)) : j < \kappa_{i,j} \rangle$ belongs to $\prod_{j < \kappa_i} \lambda_{i,j}$ so for some $\gamma_i < \lambda_i$ we have $\{j : g((i, j)) = g_{\gamma_i}^i(j)\} \in D_i^+$. Similarly for some $\beta < \lambda$, $\{i < \kappa : \gamma_i = g_\beta^*(i)\} \in D_*^+$. Easily $\{(i, j) \in u : g_\beta(i, j) = g(i, j)\} \in D^+$, so $\mathcal{G}$ witness that $T_D(\bar{\lambda}) = \lambda$ is as required.]

- (*)₂ if $g \in \Pi\{\lambda_{i,j} : (i, j) \in u\}$ then $T_D(g) < \lambda$
 [Why? Without loss of generality $g((i, j)) > 0$ for every $(i, j) \in u$. For each $i < \kappa$, let $g_i \in \prod_{j < \kappa_i} \lambda_{i,j}$ be defined by $g_i(j) = g((i, j))$. So $g_i \in \prod_{j < \kappa_i} \lambda_{i,j}$ hence $\mu_i =: T_{D_i}(g_i) < \lambda_i$ hence there is a sequence $\langle h_\alpha^i : i < \mu_i \rangle$ such that $h_\alpha^i \in \prod_{j < \kappa_i} g_i(j)$ and $(\forall h \in \prod_{j < \kappa_i} g_i(j))(\exists \alpha < \mu_i)(\neg(h \neq_{D_i} h'_\alpha))$. Clearly $\bar{\mu} = \langle \mu_i : i < \kappa \rangle \in \prod_{j < \kappa} \lambda_i$ hence $\mu_* =: T_{D_*}(\bar{\mu}) < \lambda$ and let $\langle g_\alpha^{**} : \alpha < \mu \rangle$ exemplifies this. We now define $f_\alpha^{**} \in \prod_{(i,j) \in u} g((i, j))$ by $f_\alpha^{**}((i, j)) = h_{g_\alpha^{**}(i)}^i(j)$ and it suffices to show that $T_D(g) \leq \mu(< \lambda)$ is exemplified by $\{f_\alpha^{**} : \alpha < \mu\}$ which is proved as in (*)₁, the second half of the proof.]

So we are done.

2) Similarly.

3) By [Sh:e, I].

4) Easy (and proved above). □_{2.5}

2.6 Remark. 1) So if D is defined from $D^*, \langle D_i : i < \kappa \rangle$, as in 2.5 and $\bar{\lambda} = \langle \lambda_{i,j} : (i, j) \in u \rangle, \lambda_i = T_{D_i}(\langle \lambda_{i,j} : j < \kappa_i \rangle), \lambda = T_{D_*}(\langle \lambda_i : i < \kappa \rangle)$ then $\lambda = T_D(\bar{\lambda})$.

We may wonder

2.7 Question: Can we add in Claim 2.2:

- 1) If λ is regular can we add: each λ_η is regular.
- 2) Can add the case of nice filters.

See below.

2.8 Claim. *If $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ is a J -representation of $\lambda, \lambda = \text{cf}(\lambda) > 2^\kappa$ and J is an $\aleph_1$ -complete ideal on κ then for some $J' \in J^+$, the sequence $\bar{\lambda}$ is a J' -representation of λ and $\prod_{i < \kappa} \lambda_i / J'$ has true cofinality λ (hence $\{i : \lambda_i \text{ singular}\} \in J$).*

Proof. By the pcf theorem there is $u^* \subseteq \kappa$ such that $\lambda \notin \text{pcf}\{\lambda_i : i \in \kappa(u^*)\}$ and $\lambda \geq \text{cf}(\prod_{i \in u^*} \lambda_i)$. If $\bar{\lambda}$ is a $(J + u^*)$ -representation of λ then $\lambda = T_{J+u^*}(\bar{\lambda})$ but this

implies that for some $u \in (J + u^*)^+$ we have $\prod_{i \in u} \lambda_i / (J + u^*) \upharpoonright u$ has true cofinality

λ by [Sh 589, 1.1], contradiction. Hence $\bar{\lambda} \upharpoonright u^*$ is a $(J \upharpoonright u^*)$ -representation of λ (by ?), so without loss of generality $u^* = \kappa$, so $\lambda = \max \text{pcf}\{\lambda_i : i < \kappa\}$. Let

→ $\text{scite}\{\text{xxX}\}$ undefined

$J_1 = \{u \subseteq \kappa : u \in J \text{ or } u \notin J \text{ and } \mathcal{P}(u) \cap J_2 \subseteq J_1 \text{ where } J_\chi = \{u \subseteq \kappa : u \in J \text{ or for some } v \in J \text{ we have } \lambda > \max \text{pcf}\{\lambda_i : i \in u \setminus v\}\}$. Clearly J_1, J_2 are ideals on κ extending J and by the pcf theorem we have $J_1 \cap J_2 = J$. So by ? for some

→ $\text{scite}\{\text{xxX}\}$ undefined

$\ell \in \{1, 2\}$, $\bar{\lambda}$ is a J_ℓ -representation of λ .

Case 1: $\ell = 1$.

So $\lambda = T_{J_1}(\bar{\lambda})$ hence by [Sh 589, 1.1] for some $v \in (J_1)^+$, $\prod_{i \in v} \lambda_i (J_1 \upharpoonright v)$ has true cofinality λ . So if $u \in J_2 \setminus J$ then for some $u' \subseteq u$, $\lambda > \max \text{pcf}(\{\lambda_i : i \in u \setminus u'\})$ and $J_2 \upharpoonright u = J \upharpoonright u$ hence $v \cap u \in J$. But this means $v \cap u \in J$ for every $u \in J_2 \setminus J$ (and, of course, for $u \in J$) hence $v \in J_1$, contradiction.

Case 2: $\ell = 2$.

By the pcf theorem $\prod_{i < \kappa} \lambda_i / J_2$ has true cofinality λ . □_{2.8}

2.9 Conclusion. In 2.2 we can add:

- (j) if λ is regular then every λ_η is regular and for $\eta \in \mathcal{T} \setminus \max \mathcal{T}$ we have
- $$\lambda_\eta = \text{tcf}(\prod_{\alpha < \kappa_\eta} \lambda_{\eta \cap \langle \alpha \rangle} / \mathcal{D}_\eta).$$

Now 2.7(2) (and also 2.7(1)) are answered by:

Saharon: Change 2.9 to cover singleton λ as needed in 3.1!

2.10 Observation. 1) Assume that

- (a) J_1, J_2 are ideals on κ with intersection J_0
- (b) $f_i \in {}^\kappa(\text{Ord} \setminus \omega)$.

Then $T_J(f) = \text{Min}\{T_{J_1}(f), T_{J_2}(f)\}$.

2) If (a) above holds and $\bar{\lambda}$ is a J -representation of λ , then for some $\ell \in \{1, 2\}$, $\bar{\lambda}$ is

a J_ℓ -representaiton of λ .

3) See D_1 in the proof of 3.1.

2.11 Claim. Assume¹ that the pair $(\mathbf{K}[S], \mathbf{V})$ fails the covering lemma for every $S \subseteq \beth_2(\kappa)$ (or less). Then in 2.2 we can add:

1) If $\mathfrak{a} \subseteq \text{Reg} \cap \lambda \setminus (2^{<\theta})^+$ and $|\mathfrak{a}| < \theta$ and $\text{pcf}_{\sigma\text{-comp}}(\mathfrak{a}) \not\subseteq \lambda, \lambda > 2^{<\theta}$ then for some $\kappa = \text{cf}(\kappa) \in [\sigma, \theta)$ and κ -complete ideal J on κ and $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ we have:

- (a) $\text{cf}(\lambda) > 2^{<\theta} \Rightarrow \text{cf}(\lambda_i) > 2^{<\theta}$
- (b) $\langle \lambda_i : i < \kappa \rangle$ is an exact true J -representation of λ
- (c) if λ is regular then every λ_i is regular.

2) For any normal filter D on κ we can further demand in part (1) that for some $\iota : \kappa \rightarrow \kappa, (J, \iota)$ is nice and $A \in D \Rightarrow \iota^{-1}(\kappa \setminus A) \in J$.

2A) If $\sigma \geq \partial = \text{cf}(\partial) > \aleph_0$ and D is a normal filter on ∂ we can add in part (1) that (J, ι) is nice and $A \in D = \iota^{-1}(\kappa \setminus A)$. Similarly for normal filters on $[\sigma_1]^{<\partial}$.

3) So in 2.2, we can strengthen clauses (f), (g) to

- (f)⁺ if $A \in \mathcal{D}_\eta^+, \eta \in \mathcal{T} \setminus \max_{\mathcal{T}}$ and $f \in \prod_{\alpha < \kappa_\eta} \lambda_{\eta \frown \langle \alpha \rangle}$ then $T_{\mathcal{D}_\eta + A}(\langle \lambda_{\eta \frown \langle \alpha \rangle} : \alpha < \kappa_\eta \rangle) = \lambda_\eta > T_{\mathcal{D}_\eta + A}(f)$ (hence the parallel result for D_η)
- (g)⁺ if $\eta \in \mathcal{T} \setminus \max_{\mathcal{T}}, A \in \mathcal{D}_\eta^+$ and $f \in \prod_{\alpha < \kappa_\eta} \lambda_{\eta \frown \langle \alpha \rangle}$ then $T_{\mathcal{D}_\eta + A}(f) < \lambda_\eta$ (hence the parallel result for D_η)
- (h)⁺ for each $\eta \in \mathcal{T} \setminus \max_I$ for some $\iota_\eta : \kappa_\eta \rightarrow \kappa_\eta$ the pair $(\mathcal{D}_\eta, \iota_\eta)$ is nice
- (i) (γ) if λ is regular then every λ_η is regular.

Proof. By [Sh 420, §3], better [Sh 835], very close to [Sh 386].

1) There are κ a κ -complete filter on κ and $\lambda_i < \lambda$ such that $T_D(\langle \lambda_i : i < \kappa \rangle) \geq \lambda$. By the results quoted above without loss of generality D is a normal filter on $\kappa \times \kappa$ with $\iota(\alpha, \beta) = \alpha$. Now we can choose $(D, \bar{\lambda})$ such that D is a nice filter on $\kappa \times \kappa, T_D(\bar{\lambda}) \geq \lambda$ and $\text{rk}_D^3(\bar{\lambda})$ is minimal. As $D_1 \subseteq D_2 \Rightarrow T_{D_1}(\bar{\lambda}) \leq T_{D_2}(\bar{\lambda})$ without loss of generality $\text{rk}_D^3(\bar{\lambda}) = \text{rk}_D^2(\bar{\lambda})$ and so $A \in D^+ \Rightarrow \text{rk}_{D+A}^3(\lambda) = \text{rk}_{D+A}^2(\lambda) = \text{rk}_D^3(\bar{\lambda})$ and $T_{D+A}(\bar{\lambda}) \geq T_D(\bar{\lambda})$. If $T_{D+A}(\bar{\lambda})$ then for some $f \in \pi \bar{\lambda}, T_{D+A}(f) \geq \lambda$, let $\bar{\lambda}' = \langle f(i) : i < \kappa \rangle$, so $\bar{\lambda}' <_D \bar{\lambda}$ hence $\text{rk}_{D+A}^3(\bar{\lambda}') < \text{rk}_{D+A}^3(\bar{\lambda})$ and we get a contradiction).

2), 2A), 3) Left to the reader. □_{2.11}

¹without this assumption much more follows, see [Sh:g, V].

2.12 Definition. Assume $\sigma < \theta < \lambda$.

- 1) Let $\mathfrak{d}_0(\lambda) = \mathfrak{d}_{\sigma, \theta}^0(\lambda) = \{\kappa : \kappa = \text{cf}(\kappa) \in \text{Reg} \cap \theta \setminus \sigma \text{ such that we cannot find } \langle (\lambda_\eta, \mathcal{D}_\eta, D_\eta, \kappa_\eta) : \eta \in \mathcal{T} \rangle \text{ satisfying } [\eta \in \mathcal{T} \Rightarrow |\mathcal{T}| < \lambda_\eta] \text{ and } \kappa \notin \{\kappa_\eta : \eta \in \mathcal{T}\}\}$ (so finite!) If $\sigma = \aleph_1$ we may omit it. If $\sigma = \aleph_1, \theta = \lambda$ we may omit both.
- 2) Let $\mathfrak{d}_1(\lambda) = \mathfrak{d}_{\sigma, \theta}^1(\lambda) = \{\kappa : \kappa = \text{cf}(\kappa) < \lambda \text{ and for arbitrarily large } \alpha < \lambda \text{ we have } \kappa \in \mathfrak{d}_0(|\alpha|)\}$; note that if $\text{cf}(\lambda) > \aleph_0$ we can deduce the finiteness of $\mathfrak{d}_1(\lambda)$ from the finiteness of $\mathfrak{d}_0(\lambda)$.
- 3) Let $\mathfrak{d}'_\ell(\lambda) = \mathfrak{d}_\ell(\lambda) \cup \{\aleph_0\}$ for $\ell = 0, 1$; similarly $\mathfrak{d}'_{\ell, \theta, \sigma}$ or $\mathfrak{d}'_{\ell, \theta}$.

§3 THE MAIN RESULTS ($\text{Pr}_\ell, \text{Ps}_\ell$)

3.1 Lemma. Assume that $\mu > \aleph_0$ is strong limit singular, $\lambda \geq \text{cf}(\lambda) > \mu$ and $h : \text{cf}(\lambda) \rightarrow \lambda$ is increasing continuous with $\lambda = \cup\{h(\alpha) : \alpha < \lambda\}$ and satisfies $\text{cf}(\lambda) < \lambda \Rightarrow \text{Rang}(h) \subseteq \text{Card}$ and $\lambda = \text{cf}(\lambda) \Rightarrow h(\alpha) = \alpha$. Then for some $\kappa < \mu$ and finite $\mathfrak{d} \subseteq \text{Reg} \cap \mu$ there is $\bar{\mathcal{P}}$ such that

- (*) $_{\lambda, \bar{\mathcal{P}}}$ $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha < \text{cf}(\lambda) \rangle$, $\mathcal{P}_\alpha \subseteq [h(\alpha)]^{<\mu}$, $|\mathcal{P}_\alpha| < \lambda$, $\mathcal{P}_\alpha$ increasing,
 (*) $_{\lambda, \bar{\mathcal{P}}}^{\mathfrak{d}, \kappa}$ for every A satisfying $A \subseteq \text{cf}(\lambda)$ or (more generally) $A \subseteq \lambda$ & $(\forall \alpha \in A)[\text{Min}(A \setminus (\alpha+1) < \text{Min}(\text{Rang}(h)) \setminus (\alpha+1))]$ and satisfying $|A| < \mu$ there is $\mathbf{c} : [A]^2 \rightarrow \kappa$ such that:
 if $B \subseteq A$ has no last element and $\mathbf{c} \upharpoonright [B]^2$ is constant and $\delta = \cup\{\text{Min}(\alpha+1) : \gamma < h(\alpha)\} : \gamma \in B\}$ has cofinality $\in (\text{Reg} \cap \mu \setminus \mathfrak{d})$ so $B \subseteq h(\delta)$ then $B \in \mathcal{P}_\delta$.

3.2 Remark. 1) The proof of 3.1 is simpler if λ is regular.

2) The conclusion of 3.1 means that for $\lambda > \mu$, for all but finitely many $\kappa = \text{cf}(\kappa) < \mu$, $\text{Pr}(\lambda, \text{cf}(\lambda), \kappa)$ holds (see Definition 3.9(b)).

Similarly

3.3 Claim. In fact in 3.1 we can choose $\mathfrak{d} = \mathfrak{d}'_{0, \mu}(\lambda)$.

Proof. Without loss of generality $\text{cf}(\mu) = \aleph_0$ (this is no loss by Fodor's Lemma; otherwise we may use $\mu > \text{cf}(\mu)$ or replace $\aleph_1$ by $(\text{cf}(\mu))^+$.) We first prove the desired conclusions for cardinal λ such that

$$\boxtimes_\lambda \mathfrak{a} \subseteq \lambda \cap \text{Reg} \setminus \mu \text{ \& } |\mathfrak{a}| < \mu \Rightarrow \text{pcf}_{\aleph_1\text{-comp}}(\mathfrak{a}) \subseteq \lambda.$$

Let $\bar{M} = \langle M_\alpha : \alpha < \text{cf}(\lambda) \rangle$ be such that

- ⊗₁ (a) $M_\alpha \prec (\mathcal{H}(\chi), \in)$ is increasing continuous
 (b) $\lambda \in M_\alpha$, $\|M_\alpha\| < \lambda$, $h(\alpha) \subseteq M_\alpha$
 (c) $\langle M_\alpha : \alpha \leq \beta \rangle \in M_{\beta+1}$
 (d) (α) if λ is regular then $M_\alpha \cap \lambda \in \lambda$
 (β) if λ is singular and $\alpha < \text{cf}(\lambda)$ then $\mu_\alpha + 1 \subseteq M_{\alpha+1}$
 where $\mu_\alpha = \text{Min}\{\mu : \text{if } \mathfrak{a} \subseteq \mu \cap \text{Reg} \setminus \mu \text{ and } |\mathfrak{a}| < \mu \text{ then } \text{pcf}_{\aleph_1\text{-com}}(\mathfrak{a}) \subseteq \mu \text{ and } \mu \geq \|M_\alpha\|\}$.

We let $\mathcal{P}_\alpha =: M_{\alpha+1} \cap [h(\alpha)]^{<\mu}$ and $\mathfrak{d} = \{\aleph_0\}$ and $\kappa = \aleph_0$ and will show that $\langle \mathcal{P}_\alpha : \alpha < \text{cf}(\lambda) \rangle, \mathfrak{d}$ are as required. Now $(*)_{\lambda, \mathcal{P}}$ of the claim holds trivially. To prove $(*)_{\lambda, \mathcal{P}}^{\mathfrak{d}, \kappa}$ let $A \subseteq \lambda$, $\text{otp}(A) \leq \mu$ be as there and let $\{\alpha_\varepsilon : \varepsilon < \varepsilon(*)\}$ list A in increasing order. Hence there is $\langle \beta_\varepsilon : \varepsilon < \varepsilon(*) \rangle$ increasing continuous such that $\beta_\varepsilon < \text{cf}(\lambda)$, $h(\beta_\varepsilon) \leq \alpha_\varepsilon < h(\beta_{\varepsilon+1})$. By the assumption (and 1.6, i.e., [Sh:g, II,5.4]) if λ is regular then for each $\varepsilon < \varepsilon(*)$ there is a set $\mathcal{P}^\varepsilon \subseteq [h(\beta_\varepsilon)]^{<\mu}$ of cardinality $< \lambda$ such that every $a \in [h(\beta_\varepsilon)]^{<\mu}$ is equal to the union of $\leq \kappa$ of them (by the choice of κ) hence without loss of generality $\mathcal{P}^\varepsilon \in M_{\beta_\varepsilon+1}$ hence $\mathcal{P}^\varepsilon \subseteq M_{\beta_\varepsilon+1} \cap [h(\beta_\varepsilon)]^{<\mu} = \mathcal{P}_{\beta_\varepsilon}$. If λ is singular using clause (d)(β) we get the same conclusion. So there is a sequence $\langle A_{\varepsilon,i} : i < \kappa \rangle$ such that $A_{\varepsilon,i} \in \mathcal{P}_{\beta_\varepsilon}$, $A \cap \alpha_\varepsilon = \cup \{A_{\varepsilon,i} : i < \kappa\}$. We defined $\mathbf{c} : [A]^2 \rightarrow \kappa$ by: for $\varepsilon < \zeta < \varepsilon(*)$, $\mathbf{c}\{\alpha_\varepsilon, \alpha_\zeta\} = \text{Min}\{i : \alpha_\varepsilon \in A_{\zeta,i}\}$. So assume $B \subseteq A$ and $\mathbf{c} \upharpoonright [B]^2$ is constantly $i < \kappa$ and $\delta = \sup(B)$ has cofinality $\theta \in \text{Reg} \cap \mu \setminus \mathfrak{d}$. Clearly $\alpha_\varepsilon \in B \Rightarrow \alpha_\varepsilon \cap B \subseteq \{\alpha_\zeta : \zeta < \varepsilon, \mathbf{c}\{\alpha_\zeta, \alpha_\varepsilon\} = i\} \subseteq A_{\varepsilon,i} \in \mathcal{P}_{\beta_\varepsilon}$. But $\mathcal{P}_\alpha = M_{\alpha+1} \cap [h(\alpha)]^{<\mu}$ is closed under subsets hence $\alpha_\varepsilon \in B \Rightarrow \alpha_\varepsilon \cap B \in \mathcal{P}_{\beta_\varepsilon}$.

Now in $M_{\delta+1}$ we can define a tree $\mathcal{T}$; it has $\text{otp}(B)$ levels;

the i -level is $\{a \in M_\delta : a \subseteq \delta \text{ and } \text{otp}(a) = i\}$

the order is $\triangleleft$, being initial segments.

So by the assumptions (and [Sh 589, §2]), as $\aleph_1 \leq \text{cf}(\delta) < \mu$ the number of δ -branches of $\mathcal{T}$ is $< \lambda$, so as $\mathcal{T} \in M_{\delta+1}$, every δ -branch of $\mathcal{T}$ belongs to $M_{\delta+1}$, hence $B \in M_{\delta+1}$, which implies that $B \in \mathcal{P}_\delta$, as required.

Now we prove the statement in general.

We prove this by induction on λ . For $\lambda = \mu^+$ this is trivial by the first part of the proof. So assume $\lambda > \mu^+$ and the conclusion fails, but the first part does not apply.

In particular for some $\mathfrak{a} \subseteq \text{Reg} \cap \lambda^+$, $|\mathfrak{a}| < \mu$ and $\text{pcf}_{\aleph_1\text{-comp}}(\mathfrak{a}) \not\subseteq \lambda$. Hence by 1.7 + 1.8 + 2.2 + 2.8 (alternatively + 2.11 as in the proof of 2.2; if the hypothesis of 2.11 fails that as in [Sh 386] Jensen Dodd covering lemma gives much more), for some $\kappa = \text{cf}(\kappa) \in [\aleph_1, \mu)$,

$(*)_1$ there is a sequence $\langle \lambda_i : i < \kappa \rangle$ and an $\aleph_1$ -complete filter D on κ such that

$$(a) \quad T_D\left(\prod_{i < \kappa} \lambda_i\right) = \lambda$$

$$(b) \quad \text{cf}(\lambda_i) > \mu$$

$$(c) \quad \text{if } \lambda'_i < \lambda_i \text{ for } i < \kappa, \text{ then } T_D(\langle \lambda'_i : i < \kappa \rangle) < \lambda$$

$$(d) \quad \text{tcf}\left(\prod_{i < \kappa} \lambda_i, <_D\right) = \text{cf}(\lambda).$$

Clearly we can find $\langle h_i : i < \kappa \rangle$ such that

$(*)_2$ h_i is an increasing continuous function from $\text{cf}(\lambda_i)$ to λ_i .

Let

$$D_1 = \{A : A \in D \text{ or } A \notin D, A \in D^+ \text{ and } T_{D+(\kappa \setminus A)}(\bar{\lambda}') \geq \lambda \text{ for some } \bar{\lambda}' \in \prod_{i < \kappa} \lambda_i\}.$$

Clearly

- (*)₃ D_1 is a κ -complete filter on κ extending D and we can replace D by $D + A$ whenever $A \in D_1^+$.

By the induction hypothesis applied to λ_i , as $\lambda_i > \mu$ there is a pair $(\kappa_i, \mathfrak{d}_i)$ as in the conclusion. Without loss of generality $\kappa_i^\kappa = \kappa_i$. So for some $m(*) < \omega$ and $\kappa(*) < \mu$ the set $\{i < \kappa : |\mathfrak{d}_i| = m(*), \kappa(i) \leq \kappa(*)\} \in D_1^+$ so without loss of generality

- (*)₄ $i < \kappa \Rightarrow |\mathfrak{d}_i| = m(*) \ \& \ \kappa_i = \kappa(*)$.

By (d) of (*)₁ there is $\bar{f}$ such that

- (*)₅ $\bar{f} = \langle f_\alpha : \alpha < \text{cf}(\lambda) \rangle$ is $<_D$ -increasing and cofinal in $\prod_{i < \kappa} \lambda_i$ and if $\delta < \text{cf}(\lambda), \text{cf}(\delta) < \mu$; and $\bar{f} \upharpoonright \delta$ has a $<_D$ -eub then f_δ is such a $<_D$ -eub and let $f'_\alpha \in \prod_{i < \kappa} \lambda_i$ be $f'_\alpha(i) = \text{Min}(\text{Rang}(h_i) \setminus f_\alpha(i))$ and $f''_\alpha(i) = h_i^{-1}(f'_\alpha(i))$.

For each i let $\bar{\mathcal{P}}^i = \langle \mathcal{P}_\alpha^i : \alpha < \text{cf}(\lambda_i) \rangle$ be such that $(*)_{\lambda_i, \bar{\mathcal{P}}^i, h_i}^{\mathfrak{d}_i, \kappa}$ holds. We now choose M_α for $\alpha < \text{cf}(\lambda)$ such that

- ⊗₂ (a) $M_\alpha \prec (\mathcal{H}(\chi), \in)$
- (b) $\|M_\alpha\| < \lambda$, M_α is increasing continuous, $\beta < \alpha \Rightarrow h(\beta) \subseteq M_{\alpha+1}$ and $\beta < \alpha \Rightarrow \langle M_\gamma : \gamma \leq \beta \rangle \in M_{\alpha+1}$
- (c) the following objects belong to M_α : $\langle \bar{\mathcal{P}}^i : i < \kappa \rangle$, $\langle \lambda_i, h_i : i < \kappa \rangle$, $\bar{f}$, D and μ
- (d) if $A \in D_1^+$, so $T_{D+A}(\langle |\mathcal{P}_{f_\alpha(i)}^i| : i < \kappa \rangle) < \lambda$ then $T_D(f_\alpha) + 1 \subseteq M_{\alpha+1}$ (remember $\text{cf}(\lambda) > \mu > 2^\kappa$).

Let $\mathfrak{d}^* = \{\theta : \theta = \kappa \text{ or } \{i < \kappa : \theta \notin \mathfrak{d}_i\} = \emptyset \text{ mod } D_1\}$; it should be clear that $|\mathfrak{d}^*| \leq m(*) + 1$.

Let $\mathcal{P}_\alpha = M_{\alpha+1} \cap [h(\alpha)]^{<\mu}$ and $\bar{\mathcal{P}} = \langle \mathcal{P}_\alpha : \alpha < \text{cf}(\lambda) \rangle$.

It is enough now to prove that $(*)_{\lambda, \bar{\mathcal{P}}, h}^{\mathfrak{d}^*, \kappa(*)}$ holds.

Let $A \subseteq \lambda, |A| < \mu$ be as in the assumption and we should find $\mathbf{c} : [A] \rightarrow \kappa(*)$

as required. For $i < \kappa$ let $A_i = \{f_\alpha(i) : \alpha \in A\}$, so $A_i \in [\lambda_i]^{<\mu}$ hence there is $\mathbf{c}_i : [A_i]^2 \rightarrow \kappa(*)$ as required. Recalling that $\kappa(*)^\kappa = \kappa(*)$, we can define $\mathbf{c} : [A]^2 \rightarrow \kappa(*)$ such that

- ⊗₃ if $\alpha_1 < \beta_1, \alpha_2 < \beta_2$ are from A and $\mathbf{c}\{\alpha_1, \beta_1\} = \mathbf{c}\{\alpha_2, \beta_2\}$ then
 - (i) if $i < \kappa$ then $f_{\alpha_1}(i) < f_{\beta_1}(i) \equiv f_{\alpha_2}(i) < f_{\beta_2}(i)$
 - (ii) if $i < \kappa$ then $f_{\alpha_1}(i) > f_{\beta_1}(i) \equiv f_{\alpha_2}(i) > f_{\beta_2}(i)$
 - (iii) if $i < \kappa$ and $f_{\alpha_1}(i) < f_{\beta_1}(i)$ then $\mathbf{c}_i\{f_{\alpha_1}(i), f_{\beta_1}(i)\} = \mathbf{c}_i\{f_{\alpha_2}(i), f_{\beta_2}(i)\}$.

Let $\theta \in \text{Reg} \cap \mu \setminus \mathfrak{d}^*$ and let $\delta < \text{cf}(\lambda)$ and $B \subseteq A \cap h(\delta)$ be such that $\mathbf{c} \upharpoonright [B]^2$ is constantly i and $\theta = \text{cf}(\delta)$ and $\delta = \sup(B)$. We can replace D by $D + \{i < \kappa : \theta \notin \mathfrak{d}_i\}$. So for some set $a \subseteq \theta$ we have

- ⊗₄ if $\alpha < \beta$ are from B then $a = \{i < \kappa : f_\alpha(i) < f_\beta(i)\}$.

Clearly $a \in D$. Note that by ⊗₃ for each $i \in a$ the sequence $\langle f_\alpha(i) : \alpha \in B \rangle$ is increasing and let $B_i = \{f_\alpha(i) : \alpha \in B\}$ so $\delta_i =: \sup(B_i)$ has cofinality θ and $\mathbf{c}_i \upharpoonright [B_i]^2$ is constant. Hence by the choice of $\bar{\mathcal{P}}^i$ and ⊗(iii) clearly $B_i \in \mathcal{P}_{\delta_i}^i$. Also by the choice of a and ⊗ above (and [Sh:g, II, §1]) $\bar{f} \upharpoonright \delta$ has a $\leq_D$ -eub $f', f'(i) =: \cup\{f_\alpha(i) : i \in B\}$, hence² $a' = \{i \in a : f_\delta(i) = \delta_i\}$ belongs to D . Now $|\mathcal{P}_{f_\delta(i)}^i| < \lambda_i$, hence $T_D(\langle \mathcal{P}_{f_\delta(i)}^i : i < \kappa \rangle) < \lambda$, so there is $\mathcal{F} \subseteq \prod_{i < \kappa} \mathcal{P}_{f_\delta(i)}^i, |\mathcal{F}| < \lambda$ such that for

every $g \in \prod_{i < \kappa} \mathcal{P}_{f_\delta(i)}^i$ there is $g' \in \mathcal{F}$ such that $\{i : g(i) = g'(i)\} \in D^+$. By the choice of $M_{\delta+1}$, i.e., clause (d) of ⊗₂ without loss of generality $\mathcal{F} \in M_{\delta+1}$, hence $\mathcal{F} \subseteq M_{\delta+1}$. We can define $g \in \prod_{i < \kappa} \mathcal{P}_{f_\delta(i)}^i$ by letting $i \in a' \Rightarrow g(i) = B_i$. So there is $g' \in \mathcal{F} \subseteq M_{\delta+1}$ such that $b = \{i : g(i) = g'(i)\} \in D^+$ hence $b \cap a' \in D^+$. That is $b' =: \{i \in a' : g'(i) = B_i\} \in D^+$. Clearly $b' \in M_{\delta+1}$ (as $\mu \in M_{\delta+1}$ hence $\mathcal{H}(\mu) \subseteq M_{\delta+1}$) and $g' \in M_{\delta+1}$, hence $g' \upharpoonright b' \in M_{\delta+1}$, hence also the set B^* belongs to $M_{\delta+1}$ where

$$B^* =: \{\gamma < \lambda : \{i \in b' : f_\gamma(i) \in g'(i) = g(i) = B_i\} \in D^+\}.$$

Now $|B^*| \leq \prod_{i < \kappa} B_i < \mu$ and $\alpha \in B \Rightarrow \alpha \in B^*$. But as $B^* \in M_{\delta+1}$ every subset of B^* belongs to $M_{\delta+1}$ hence $B \in M_{\delta+1}$ so $B \in \mathcal{P}_\delta$, as required. $\square_{3.1}$

²Note that here we use $\theta \neq \kappa$, in fact the only point that we use it, if we could avoid it, then $\mathfrak{d}$ could be chosen as $\{\aleph_0\}$

3.4 Discussion: 1) Note that in a sense what was done in [Sh 108], i.e., $I[x]$ large for $\lambda = \mu^+$ is done in 3.1 for any λ with $\text{cf}(\lambda) > \mu$.

2) We may consider replacing $\mathfrak{d}_\lambda$ by $\{\aleph_0\}$ in 3.1. The base of the induction is clear ($\text{pcf}_{\aleph_1}$ -inaccessibility). So eventually we have f_δ for it as above $\langle f_\alpha : \alpha \in B \rangle$, the hard case is $\text{cf}(\text{otp}(B)) = \kappa$, we have the induced $h_* \in {}^\kappa \kappa$ such that $\alpha < \kappa \Rightarrow \{i : d < h_*(i)\} \in D$, but $(\forall^D i)[\text{cf}(h_*(i)) = \aleph_0]$ (otherwise using niceness of the filter, etc., we are done).

Note that is problem appear even in the problem “assume μ is strong limit of cofinality $\aleph_1$ (or $\kappa \in [\aleph_1, \mu)$) and $2^\mu = \mu^+$, does it follows that $\diamond_{S_{\text{cf}(\mu)}^{\mu^+}}$ holds?” See [Sh 186], Cummings Dzamonja Shelah [CDSSh 571], Dzamonja Shelah [DjSh 692].

But if $\kappa = \text{cf}(\kappa) > \aleph_0$ and in 2.12 we use $D = D_\kappa + S_{\aleph_1}^\kappa$. For each $\alpha < \kappa$ we should consider $\iota(t)$; if D -positively we have $\iota(t) \leq h_*(t)$ we are done. But if $\iota(t) > h_*(t)$, D -positively then on some $A \in D^+$, $h_* \upharpoonright A$ is constant.

3.5 Conclusion. Assume $\mu < \lambda$, μ is strong limit $> \aleph_0$, λ is regular (or just $\text{cf}(\lambda) > \mu$). Then for some $\kappa < \mu$ and finite $\mathfrak{d} \subseteq \text{Reg} \cap \mu$ to which $\aleph_0$ belongs (in fact $(\mathfrak{d}_\mu^0(\lambda) \cup \{\aleph_0\})$ is O.K.), there is $\bar{\mathcal{F}}$ such that

- $\circledast_{\lambda, \bar{\mathcal{F}}}^{\mu, \mathfrak{d}, \kappa}$ (a) $\bar{\mathcal{F}} = \langle \mathcal{F}_\alpha : \alpha < \lambda \rangle, |\mathcal{F}_\alpha| < \lambda$ for $\alpha < \lambda$
- (b) $\mathcal{F}_\alpha \subseteq \{f : f \text{ a partial function from } \alpha \text{ to } \alpha, |\text{Dom}(f)| < \mu\}$, $\mathcal{F}_\alpha$ closed under restriction
- (c) for every $A \subseteq \lambda, |A| < \mu$ and $f : A \rightarrow \lambda$ for some $\mathbf{c} : [A]^2 \rightarrow \kappa$ we have
 - $\square_1$ if $B \subseteq A, \delta = \sup(B) \in E, \mathbf{c} \upharpoonright [B]^2$ is constant, $[\alpha \in B \Rightarrow f(\alpha) < \delta]$ and $\text{cf}(\delta) \notin \mathfrak{d}$ then $f \upharpoonright B \in \mathcal{F}_\delta$ and $\alpha \in B \Rightarrow f \upharpoonright (B \cap \alpha) \in \mathcal{F}_\alpha$.

Proof. We use the result of 3.1.

For clause (c) we use pairing function pr on λ such that $\text{pr}(\alpha, \beta) < \text{Max}\{\omega, \alpha + |\alpha|, \beta + |\beta|\}$ to replace the function f in clause (c) by the set $\{\text{pr}(\alpha, f(\alpha)) : \alpha \in A\}$ and first we restrict ourselves to δ in some club E of λ (the range of h in 3.1's notation) such that $\delta \in E \Rightarrow |\delta|$ divides δ (hence δ is closed under pr); so if $B \subseteq \lambda, \sup(B) \in E$ we are done. The other cases are easier as without loss of generality if $\alpha < \delta \in E$, then $\alpha + \text{Min}\{\chi : \mu \geq |\alpha| \text{ and if } \mathfrak{a} \subseteq \text{Reg} \cap \chi^+, |\mathfrak{a}| < \mu, \text{pcf}_{\kappa(*)^+ \text{-comp}}(\mathfrak{a}) \subseteq \mu^+\} < \delta$, and easy to finish. $\square_{3.5}$

3.6 Conclusion. Assume that μ is strong limit singular, $\lambda = \lambda^{<\mu}$ (equivalently $\lambda = \lambda^\mu$) and $\text{cl} : [\lambda]^{<\mu} \rightarrow [\lambda]^{<\mu}$ satisfies for notational simplicity $\text{cl}(B) = \cup\{\text{cl}(B \cap (\alpha + 1)) : \alpha \in B\}$ and $B_1 \trianglelefteq B_2 \Rightarrow B_1 \subseteq \text{cl}(B_1) \subseteq \text{cl}(B_2)$.

Then in 3.5 we can add to (a),(b) and (c) also

- (d) $\mathbf{g}$ is a function from $\{f \restriction u : f \in {}^\lambda \lambda \text{ and } u \in [\lambda]^{<\mu}\}$ to λ
- (e) for every $f : \lambda \rightarrow \lambda$ for some $g_f : [\lambda]^{<\mu} \rightarrow \lambda$ (in fact $g_f(u) = \mathbf{g}(f \restriction \text{cl}(u))$) we have
 - ⊠ for every $A \subseteq \lambda$ of cardinality $< \mu$ such that $\alpha \in A \Rightarrow g_f(A \cap \alpha) < \alpha$, for some function $\mathbf{c} : [A]^2 \rightarrow \kappa$ we have
 - ⊗ if $B \subseteq A$, $\mathbf{c} \restriction [B]$ is constant and B has no last element, $\delta = \sup(B)$ has cofinality $\notin \mathfrak{d}$ then $f \restriction \text{cl}(B)$ belong to $\mathcal{F}_\delta$ and $\alpha \in B \Rightarrow f \restriction \text{cl}(B \cap \alpha) \in \cup\{\mathcal{F}_\beta : \beta < \delta\}$
- (f) if λ is regular then there is a sequence $\bar{C} = \langle C_\delta : \delta \in S \rangle$ is
 - (α) $S \subseteq S^* = \{\delta < \lambda : \text{cf}(\delta) \in [\aleph_1, \mu)\}$,
 - (β) C_δ a club of δ of order type $\text{cf}(\delta)$ and in clause (e) we can add:
 - (γ) $f \restriction \text{cl}(C_\delta) \in \mathcal{F}_\delta$ and
 - (δ) $\alpha \in C_\delta \Rightarrow f \restriction \text{cl}(C_\delta \cap \alpha) \in \bigcup_{\beta < \delta} \mathcal{F}_\beta$ and
 - (ε) $\alpha \in \text{nacc}(C_{\delta_1}) \cap \text{nacc}(C_{\delta_2}) \Rightarrow C_{\delta_1} \cap \alpha = C_{\delta_2} \cap \alpha$
 - (ζ) if $\alpha < \text{cf}(\lambda)$ is limit, $\text{cf}(\alpha) \notin \mathfrak{d}$ then $\{C_\delta \cap \alpha : \alpha \in \text{acc}(C_\delta)\}$ has coordinates $< \lambda$
 - (η) if $\beta \leq \lambda, |B| < \mu$ then for some $\mathbf{c} : [B]^2 \rightarrow \kappa$ if $B' \subseteq B$ has no last member and $\mathbf{c} \restriction B'$ is constant and $\text{cf}(\sup(B')) \notin \mathfrak{d}$ then $\sup(B') \in S$.

Proof. We repeat the proof of 3.1.

Choose $h : \text{cf}(\lambda) \rightarrow \lambda$ and $\langle M_\alpha : \alpha < \text{cf}(\lambda) \rangle$ as in the proof of 3.1 but add the requirement that $\text{cl} \in M_0$ and still use $\mathcal{F}_\alpha = M_{\alpha+1} \cap \{f : f \text{ a partial function from } \alpha \text{ to } \alpha \text{ with domain of cardinality } < \mu\}$.

Choose $\mathbf{g}$ such that

- ⊠ (a) $\mathbf{g}$ is a function from $\{f \restriction u : f \in {}^\lambda \lambda \text{ and } u \in [\lambda]^{<\mu}\}$ onto λ
- (b) $f_1 \subseteq f_2 \in \text{Dom}(\mathbf{g}) \Rightarrow \mathbf{g}(f_1) \leq \mathbf{g}(f_2)$ and
- (c) for each $\alpha < \lambda$ for some $f \in \text{Dom}(\mathbf{g})$ we have $\mathbf{g}(f) = \alpha$ & $(\forall f')[\mathbf{g}(f') = \alpha \Rightarrow f' \subseteq f]$
- (d) if $f : B_2 \rightarrow \lambda$ and $B_1 \triangleleft B_2$ then $\mathbf{g}(B_1) < \mathbf{g}(B_2)$
- (e) $\mathbf{g}(f) = \alpha \Rightarrow \text{Dom}(f) \subseteq \alpha$.

Without loss of generality $\mathbf{g} \in M_0$ so clause (d) (of the conclusion of 3.6) holds trivially and let us prove clause (e). So, as $\mathbf{g}$ has already been chosen, we are given $A \subseteq \text{cf}(\lambda)$ of cardinality $< \mu$ and $f : A \rightarrow \lambda$ such that $\alpha \in A \Rightarrow \mathbf{g}(f \upharpoonright (A \cap \alpha)) < \alpha$.

Now $\alpha \mapsto \mathbf{g}(f \upharpoonright (A \cap \alpha))$ is an increasing function from A to λ and let $A' = \{\mathbf{g}(f \upharpoonright (A \cap \alpha)) : \alpha \in A\}$ and let $\mathbf{c}' : [A']^2 \rightarrow \kappa$ be as proved to exist in 3.1 and by $\mathbf{c} : [A]^2 \rightarrow \kappa$ be defined by $\mathbf{c}\{\alpha, \beta\} = \mathbf{c}'\{\mathbf{g}(f \upharpoonright (A \cap \alpha)), \mathbf{g}(f \upharpoonright (A \cap \beta))\}$.

It is easy to check that $\mathbf{c}$ is as required. We turn to proving clause (f) of the claim. Now there is a function $F : {}^\omega \lambda \rightarrow \lambda$ such that for any $\bar{\alpha} \in {}^\omega \lambda$ for every large enough $n < \omega$ there are $m_0 < m_1 < m_2 < \dots < \omega$ which are $> n$ and $\alpha_n = F(\alpha_{m_0}, \alpha_{m_1}, \dots)$; by [EH]. For any $u \in [\lambda]^{<\mu}$ we define $cl_*(u)$ as follows: let $u^{+\mathbf{g}} = u \cup \{\mathbf{g}(1_v) : v \subseteq u \cap \alpha \text{ and } \alpha \in u\}$ and let $cl_*(u)$ be the minimal set v such that $u^{+\mathbf{g}} \subseteq v$ and $[\delta = \sup(v \cap \delta) < \sup(u^{+\mathbf{g}}) \ \& \ \text{cf}(\delta) \leq |u| \Rightarrow \delta \in v]$ and $[\mathbf{g}(1_w) \in v \ \& \ |w| \leq |u| \Rightarrow w \subseteq v]$ and $\bar{\alpha} \in {}^\omega v \Rightarrow F(\alpha) \in v$; so $|cl_*(u)| \leq (|u|^+ + 2)^{\aleph_0}$.

In the proof above we can replace cl by $cl_* \circ cl$. Now if $\delta < \lambda, \aleph_0 < \text{cf}(\delta) < \mu$ for some club C_δ^* of δ of order type $\text{cf}(\delta)$ we have: if $C \subseteq C_\delta^*$ is a club of δ then $cl_* \circ cl(C) = cl_* \circ cl(C_\delta^*)$ (exists by the choice of F). Alternatively, let $C'_\delta = \cap \{cl_*(C) : C \text{ a club of } \delta\}$; however, C'_δ seemingly has order type just $< (\text{cf}(\delta)^{\aleph_0})^+$. Now if C_δ^* satisfies $(\forall \alpha \in C_\delta^*)(\mathbf{g}(1_{C_\delta \cap \alpha}) < \delta)$ then we can find C_δ^{**}, C_δ such that:

- ⊗₁ $C_\delta^{**} \subseteq cl_* \circ cl(C_\delta^*)$ is a club of δ of order type $\text{cf}(\delta)$ such that $\alpha \in \text{nacc}(C_\delta^{**}) \Rightarrow \sup((C_\delta^{**} \cup \{0\}) \cap \alpha) < \mathbf{g}(1_{(C_\delta^{**} \cup \{0\}) \cap \alpha}) < \alpha$
- ⊗₂ C_δ is $\{\mathbf{g}(1_{(C_\delta^{**} \cup \{0\}) \cap \alpha}) : \alpha \in \text{nacc}(C_\delta^{**})\} \cup \text{acc}(C_\delta^{**})$.

Clearly

- ⊗₃ $C_\delta \subseteq cl_*(B)$ whenever $B \subseteq \delta = \sup(B)$
- ⊗₄ if $\alpha \in \text{nacc}(C_{\delta_1}) \cap \text{nacc}(C_{\delta_2})$ then $C_{\delta_1} \cap \alpha = C_{\delta_2} \cap \alpha$.

We are done as we have used $cl_* \circ cl$ and

- (*) if $\delta < \lambda, \aleph_0 < \text{cf}(\delta) < \mu$ and B is an unbounded subset of δ then $C_\delta \subseteq cl_*(B)$.

□_{3.6}

3.7 Remark. 1) In 3.1, 3.5, 3.6 if λ is regular; then

$$A_{\bar{M}} = \{\delta : \delta < \lambda, \text{cf}(\delta) < \delta \text{ and there is } u \subseteq \delta = \sup(u), \text{otp}(u) < \delta, (\forall \alpha < \delta)(u \cap \alpha \in M_\alpha)\}$$

belongs to $I[\lambda]$ and the δ mentioned in $(*)_{\lambda, \mathcal{P}}^{\mathfrak{d}, \kappa}$ of 3.1, (c) of 3.5 necessarily belongs to $A_{\bar{M}}$. So $A_{\bar{M}}$, for ordinals of cofinality $\in \text{Reg} \cap \mu \setminus \mathfrak{d}$ contains “almost all of them” in the appropriate sense.

2) We can use them to upgrade if $\{\delta < \omega_2 : S_{\kappa}^{\bar{\square}^{\delta+1}} \in I(\bar{\square}_{\delta}^+)\}$ then $S_{\kappa}^{\bar{\square}^{\omega+1}} \in I[\bar{\square}_{\omega_1+1}^+]$ when $\kappa = \text{cf}(\kappa) > \aleph_1$, see [Sh 589].

3.8 Main Conclusion. 1) If μ is strong limit and $\lambda = \lambda^{<\mu}$ then for all but finitely many regular $\kappa < \mu$ (actually $\kappa \notin \mathfrak{d}_{\mu}^0(\lambda) \cup \{\aleph_0\}$ is enough) we have $\text{Ps}_1(\lambda, \kappa)$, see Definition 3.9 below.

2) We also get $\text{Ps}_1(\text{cf}(\lambda), \lambda, \kappa)$ when $\kappa > \aleph_0$.

Proof. 1) By 3.5, 3.6.

3.9 Definition. 1) $\text{Ps}_1(\lambda, \kappa)$ means that $\text{Ps}_2(\lambda, S)$ for some stationary $S \subseteq S_{\kappa}^{\lambda}$.

2) $\text{Ps}_2(\lambda, S)$ means that for some $\bar{C} = \langle C_{\delta} : \delta \in S \rangle$ we have $\text{Ps}_3(\lambda, \bar{C})$.

3) $\text{Ps}_3(\lambda, \bar{C})$ means that for some $\bar{\mathcal{F}}$ we have $\text{Ps}_4(\lambda, \bar{C}, \bar{\mathcal{F}})$.

4) $\text{Ps}_4(\lambda, \bar{C}, \bar{\mathcal{F}})$ means that for some S :

- (a) S is a stationary subset of λ
- (b) $\bar{C}$ has the form $\langle C_{\delta} : \delta \in S \rangle$
- (c) $\bar{\mathcal{F}}$ has the form $\bar{\mathcal{F}} = \langle \mathcal{F}_{\alpha} : \alpha \in S \rangle$
- (d) C_{δ} is a club of δ of order type $\text{cf}(\delta)$ and $\alpha \in \text{nacc}(C_{\delta_1}) \cap \text{nacc}(C_{\delta_2}) \Rightarrow \alpha \cap C_{\delta_1} = \alpha \cap C_{\delta_2}$
- (e) $\mathcal{F}_{\delta}$ is a set of functions from C_{δ} to δ of cardinality $< \lambda$
- (f) if $f : \lambda \rightarrow \lambda$ then for stationarily many $\delta \in S$ we have $f \restriction C_{\delta} \in \mathcal{F}_{\delta}$.

5) $\text{Ps}_4(\lambda, \mu, h, \bar{C}, \bar{\mathcal{F}})$ is defined similarly (so λ is regular) except that

- (e)'₁ h is an increasing continuous function from λ to μ with limit μ
- (e)₂ $\mathcal{F}_{\delta}$ is a set of functions from δ to $h(\delta)$ of cardinality $< \mu$
- (f) if $f : \lambda \rightarrow \mu$ then for stationarily many $\delta \in S$ we have $f \restriction C_{\delta} \in \mathcal{F}_{\delta}$.

6) If in (5) we omit h we mean some h .

7) $\text{Ps}_1(\lambda, \mu, \kappa)$, $\text{Ps}_2(\lambda, \mu, S)$, $\text{Ps}_3(\lambda, \mu, \bar{C})$ are defined parallelly.

3.10 Definition. Pr_ℓ are defined similarly except not using $\bar{C}$ and $\mathcal{F}_\delta$ is a set of functions from some unbounded subset of δ into δ (or $h(\delta)$) that is:

- 1) $\text{Pr}_1(\lambda, \kappa)$ means that $\text{Pr}_2(\lambda, S)$ for some stationary $S \subseteq S_\kappa^\lambda$.
- 2) $\text{Pr}_4(\lambda, \mathcal{F})$ means that for some S :

- (a) S is a stationary subset of λ
- (b) $\mathcal{F}$ has the form $\mathcal{F} = \langle \mathcal{F}_\alpha : \alpha \in S \rangle$
- (c) $\mathcal{F}_\delta$ is a set of cardinality $< \lambda$ of functions from some unbounded subset of δ to δ
- (d) if $f : \lambda \rightarrow \lambda$ then for stationarily many $\delta \in S$ we have $f \upharpoonright C \in \mathcal{F}_\delta$ for some $A \subseteq \delta = \sup(C)$.

- 3) $\text{Pr}_4(\lambda, \mu, h, \mathcal{F})$ is defined similarly to except that

- (c)₁' h is an increasing continuous function from λ to μ with limit μ
- (c)₂ $\mathcal{F}_\delta$ is a set of cardinality $< \lambda$ of functions from some unbounded subset of δ to $h(\delta)$
- (d) if $f : \lambda \rightarrow \mu$ then for stationarily many $\delta \in S$ we have $f \upharpoonright A \in \mathcal{F}_\delta$ for some $A \subseteq \delta = \sup(A)$.

- 4) If in (5) we omit h we mean some h .

3.11 Observation.: If $\text{Ps}_4(\lambda, \bar{C}, \mathcal{F})$, $\lambda_1 = \text{cf}(\lambda) < \lambda$, $\bar{C} = \langle C_\delta : \delta \in S \rangle$, $(\forall \delta \in S)[\text{cf}(\delta) > \aleph_0]$, $h : \lambda_1 \rightarrow \lambda$ is increasing continuous with limit λ , $S' = \{\delta < \lambda_1 : h(\delta) \in S\}$, $C'_\delta = \{\alpha < \delta : h(\alpha) \in C_\delta\}$, $\bar{C}' = \langle C'_\delta : \delta \in S' \rangle$, $\mathcal{F}'_\delta = \{h \circ f : f \in \mathcal{F}_\delta\}$ then $\text{Ps}_4(\lambda_1, \lambda, h, \bar{C}', \mathcal{F}')$.

3.12 Conclusion. 1) If $\lambda = \text{cf}(\lambda) > \mu > \aleph_0$, μ strong limit singular then for some $A \in I[\lambda]$, $\kappa < \mu$ and finite $\mathfrak{d} \subseteq \text{Reg} \cap \mu$ (in fact $\mathfrak{d} = \mathfrak{d}'_{0,\mu}(\lambda)$) we have:

- (*) for every $\kappa(2) = \kappa(2)^{\kappa(1)} < \mu$, $\kappa(1) > \kappa$ and increasing continuous sequence $\langle \alpha_\varepsilon : \varepsilon < \kappa(2)^+ \rangle$ we have: there is a club C of $\kappa(2)^+$ such that $\{\alpha \in C : \text{cf}(\alpha) \notin \mathfrak{d} \text{ and } \text{cf}(\alpha) \leq \kappa(1)^+\} \subseteq A$.

- 2) If above $\lambda = \lambda^{<\lambda}$ we can add: $\kappa \in \text{Reg} \cap \mu \setminus \mathfrak{a} \Rightarrow (D\ell)_{S_\kappa^\lambda}$ (and is even $(D\ell)_S$ for any $S \subseteq S_\kappa^\lambda$ which is $\neq \emptyset$ modulo for a suitable filter similarly in (3)).

On diamond from instances of GCH see [Sh 460]. Whereas $\lambda = \mu^+$ successor of regular cardinals has strong partial squares ([Sh 351, §4]) for successor of singular we have much less. If $\lambda = \mu^+$, $\mu^\theta = \mu$ for cofinalities $\leq \theta$ we still have this.

3.13 Conclusion. Assume $\lambda = \text{cf}(\lambda) > \mu > \aleph_0, \mu$ strong limit and $\mathfrak{d} = \mathfrak{d}'_{1,\mu}(\lambda) = \{\kappa : \kappa = \text{cf}(\kappa) \text{ and } \kappa \in \mathfrak{d}'_0(|\alpha|) \text{ for arbitrarily large } \alpha < \lambda\}$ which is finite. If $\lambda = \chi^+ = 2^\chi$ and $\kappa \in \text{Reg} \cap \mu \setminus \mathfrak{d}$ then $\diamond_{S_\kappa}^\lambda$.

§4 MIDDLE DIAMONDS AND BLACK BOXES

We use §3 to improve the main results of [Sh 775] on the middle diamond. The point is that there we use [Sh 460], here we use §3 instead. For our aim we quote some results and definitions. See 4.4 and 4.5. The conclusions are

In full

4.1 The Middle Diamond Claim. *Assume*

- (a) $\lambda = \text{cf}(2^\mu)$, D is a μ^+ -complete filter on λ extending the club filter
- (b) $\kappa = \text{cf}(\kappa) < \lambda$ and $S \subseteq S_\kappa^\lambda$
- (c) $\bar{C} = \langle C_\delta : \delta \in S \rangle$, C_δ a club of δ of order type κ and $\lambda = 2^\mu$ & $\delta \in S \Rightarrow \lambda > |\{C_\delta \cap \alpha : \alpha \in \text{nacc}(C_\delta)\}|$ and $S \in D$
- (d) $2^{<\chi} \leq 2^\mu$ and $\theta \leq \mu$
- (e) $\text{Ps}_1(\lambda, 2^\mu, \bar{C})$ (see Definition 3.9)
- (f) $\text{Sep}(\mu, \theta)$ (see Definition 4.2 below and 4.3 on sufficient conditions).

Then $\bar{C}$ has the $(D, 2^\mu, \theta)$ -Md-property (recall that this means that the number of colours is θ not just 2, see Definition 4.5 below).

Proof. By the proof of [Sh 775, 1.10] or see the more general claim on the squared middle diamond (here or in subsequent work).

4.2 Definition. 1) $\text{Sep}(\mu, \theta)$ means that for some $\bar{f}$ and Υ :

- (a) $\bar{f} = \langle f_\varepsilon : \varepsilon < \mu \rangle$
- (b) f_ε is a function from ${}^\mu\theta$ to θ
- (c) for every $\partial \in {}^\chi\theta$ the set $\{\nu \in {}^\mu\theta : \text{for every } \varepsilon < \chi \text{ we have } f_\varepsilon(\nu) \neq \partial(\varepsilon)\}$ has cardinality $< \Upsilon$
- (d) $\Upsilon = \text{cf}(\Upsilon) \leq 2^\mu$.

2) $\text{Sep}_\sigma(\mu, \theta)$ means that for some $\bar{f}$, R and Υ we have

- (a) $\bar{f} = \langle f_\varepsilon^i : \varepsilon < \mu \text{ and } i < \sigma \rangle$
- (b) f_α^i is a function from ${}^M\theta$ to ${}^\mu\theta$
- (c) $R \subseteq {}^\mu\theta$; $|R| = 2^\mu$ (if $R = {}^\mu\theta$ we may omit it)
- (d) $\bar{\mathcal{I}} = \langle \mathcal{I}_i : i < \sigma \rangle$, $\mathcal{I}_i \subseteq \mathcal{P}(\mu)$ and if $A_j \in \mathcal{I}_j$ for $j < j^* < \sigma$ then $\mu \neq \bigcup \{A_j : j < j^*\}$ (e.g. $\mathcal{I}_i = \mathcal{I}$ a σ -complete ideal on μ)

(e) if $\eta \in {}^\mu\theta$ and $i < \sigma$ then $\Upsilon > |\text{Sol}_\eta|$ where

$$\text{Sol}_\eta = \{\rho \in R : \text{the set } \{\varepsilon < \mu : (f_\alpha^i(\eta))(\varepsilon) \neq \eta(\varepsilon)\} \text{ belong to } I\}.$$

We may wonder if clause (f) of the assumption is reasonable; the following Claim gives some sufficient conditions for clause (f) to hold.

4.3 Claim. *Clause (f) of 4.1 holds, i.e., $\text{Sep}(\mu, \theta)$ holds, if at least one of the following holds:*

- (a) $\mu = \mu^\theta$
- (b) $\mathbf{U}_\theta(\mu) = \mu + 2^\theta \leq \mu$,
- (c) $\mathbf{U}_J(\mu) = \mu$ where for some σ we have $J = [\sigma]^{<\theta}$, $\theta \leq \sigma$, $2^{<\sigma} < \mu$
- (d) μ is a strong limit of cofinality $\neq \theta = \text{cf}(\theta) < \mu$
- (e) $\mu \geq \beth_\omega(\theta)$.

Proof. This is [Sh 775, 1.11].

4.4 Definition. 1) We say that λ has the (κ, θ) -MD⁺ property when there are $\bar{C}^i = \langle C_\delta : \delta \in S_i \rangle$ for $i < \lambda$ such that

- $\square_{\bar{C}}^{\lambda, \kappa}$ (a) S_i are pairwise disjoint stationary subsets of λ
- (b) $\delta \in S_i \Rightarrow \text{cf}(\delta) = \kappa$
- (c) C_δ is a club of δ of order type κ and every $\alpha \in \text{nacc}(C_\delta)$ is a successor ordinal,
- (d) if $\alpha \in \text{nacc}(C_{\delta_1}) \cap \text{nacc}(C_{\delta_2})$ then $C_{\delta_1} \cap \alpha = C_{\delta_2} \cap \alpha$
- (e) $\bar{C}^i$ has the θ -MD property which means that there is $\bar{f} = \langle f_\delta : \delta \in S_i \rangle$ such that $f_\delta : {}^\omega(C_\delta) \rightarrow \theta$ and for every $f \in {}^\omega \lambda \rightarrow \theta$ for stationarily many $\delta \in S_i$ we have $f_\delta = f \upharpoonright C_\delta$.

2) We write MD instead of MD⁺ if we omit clause (d); we write MD[±] if we replace “ C_δ a club of δ ” by “ $C_\delta \subseteq \delta = \sup(C_\delta)$ ” and MD[−] if we do both changes.

4.5 Definition. 1) We say that $\bar{C}$ exemplifies $\text{Md}^+(\lambda, \kappa, \theta, \Upsilon, D)$ when

- (a) $\lambda > \kappa$ are regular cardinals, Υ an ordinal (or a function with domain λ or ${}^\omega > \lambda$ in this case a function f from X to Υ means that f is a function with domain X and $f(x) \in \Upsilon(x)$, so ${}^C \Upsilon = \{f : f \text{ is a function with } \text{Dom}(f) = C \text{ and } \alpha \in C \Rightarrow f(\alpha) \in \Upsilon(\alpha)\}$)
- (b) $C = \langle C_\delta : \delta \in S \rangle$, S a stationary subset of λ such that $\delta \in S \Rightarrow \text{cf}(\delta) = \kappa$
- (c)⁺ C_δ is a club of δ disjoint to S and $\alpha \in \text{nacc}(C_{\delta_1}) \cap \text{nacc}(C_{\delta_2}) \Rightarrow C_{\delta_2} \cap \alpha = C_{\delta_1} \cap \alpha$ so we may define $C_\alpha = C_\delta \cap \alpha$ when $\alpha \in \text{nacc}(C_\delta)$
- (d) if $\mathbf{F}$ is a function from $\bigcup_{\delta \in S} \{f : f \text{ is a function from } {}^\omega > (C_\delta) \text{ to } \Upsilon\}$ to θ then for some $\mathbf{c} \in {}^S \theta$ for every $f \in {}^\lambda \Upsilon$ the set $\{\delta \in S : \mathbf{F}(f \upharpoonright C_\delta) = \mathbf{c}(\delta)\} \in D^+$.

2) We write Md instead Md^+ if we weaken (c)⁺ to

- (c) C_δ is an unbounded subset of δ .

3) We say $\bar{C}$ has the (D, Υ, θ) -Md property when clauses (a),(b),(c),(d) above holds; we say λ has this property if some $\bar{C} = \langle C_\delta : \delta \in S \rangle$ has it, $S \subseteq S_\theta^\lambda$ stationary.

4.6 Remark. 1) How strong is the demand that S can be divided to λ sets S_i with the property? It is hard not to have it.

2) In 4.7 to have more than one exception is a heavy demand on $\mathcal{H}(\mu)$.

3) We can improve 4.7 including the case $\text{cf}(\mu_*) = \aleph_0$, even $\mu_* = \beth_{\alpha+\omega}$. Then probably in part (2) we have to distinguish λ successor of regular (easy), success of singular (harder), rest (hardest).

4.7 Conclusion. 1) If μ_* is strong limit $> \aleph_0$, $\mu \geq \mu_*$, $\lambda = \text{cf}(2^\mu)$ and $\Upsilon = 2^\mu$ then for all but finitely many $\kappa \in \text{Reg} \cap \mu_*$ (even every $\kappa \in \text{Reg} \cap \mu_* \setminus \mathfrak{d}'_{1,\mu_*}(2^\mu)$), there is $\bar{C} = \langle C_\delta : \delta \in S \rangle$ exemplifies $\text{Md}^+(\lambda, \kappa, \theta, \Upsilon)$ hence (κ, θ) -MD⁺.

2) Assume μ_* is strong limit singular of uncountable cofinality and $\lambda = \text{cf}(\lambda) > \mu_*$ is not strongly inaccessible. Then for all but finitely many $\kappa \in \text{Reg} \cap \mu_*$ for every $\theta < \mu_*$, λ has (κ, θ) -MD hence (κ, θ) -MD⁺ (moreover only one of the exceptions depend on λ).

Proof. 1) Let $\mathfrak{d} = \mathfrak{d}'_{\mu_*,0}(\lambda)$. So by §3 we have $\kappa \in \text{Reg} \cap \mu_* \setminus \mathfrak{d} \Rightarrow \text{Ps}_1(\lambda, 2^\mu, \bar{C})$ for some $\bar{C}$ satisfying clause (c) of 4.1, moreover clauses (c) (d) of 4.4(1). So we apply 4.1.

2) Let $\langle \mu_i : i < \text{cf}(\mu_*) \rangle$ be increasing continuous with limit μ_* each μ_i is strong

limit singular. For each $i < \text{cf}(\mu_*)$ let $\mathfrak{d}_i = \mathfrak{d}'_{\mu_i, 0}(\text{cf}(2^{\mu_i}))$, so it is finite and let $\mathfrak{d} = \{\kappa : \kappa = \text{cf}(\kappa) < \mu_* \text{ and } \kappa \in \mathfrak{d}_i \text{ for every } i < \text{cf}(\mu_*) \text{ large enough}\}$.

Case 1: $(\forall \alpha < \lambda)[|\alpha|^{<\mu_*} < \lambda]$.

So we can find $\mu < \lambda \leq 2^\lambda$, let $\mu_1 = ((\mu)^{<\mu_*})^{<\mu_*}$ this cardinal is $< \lambda$ and $\mu_1 = (\mu_1)^{\mu_*}$.

Now use [Sh 775, §2].

Case 2: $(\exists \alpha < \lambda)[|\alpha|^{<\mu_*} \geq \lambda]$.

As λ is regular for some $\kappa < \lambda$, $\mu < \lambda$ we have $\mu^\kappa \geq \lambda$. Let $\mu = \text{Min}\{\mu : \mu^\kappa \geq \lambda \text{ for some } \kappa < \mu_*\}$.

NOTE: Here getting λ pairwise disjoint S_i should be done. Again we use [Sh 775, §2].

Remark. $\aleph_0 \in \mathfrak{d}$ as we need $F : {}^\omega \lambda \rightarrow \lambda$ as in §3!!

□_{4.7}

4.8 Definition. We say that $\bar{C}$ exemplify $\text{BB}_0(\lambda, \kappa, \theta)$ when

- (a) $\lambda > \kappa$ are regular
- (b) $\bar{C} = \langle C_\delta : \delta \in S \rangle$, S a stationary subset of λ such that $\delta \in S \Rightarrow \text{cf}(\delta) = \kappa$
- (c) C_δ is an unbounded subset of δ disjoint to S such that $\alpha \in C_{\delta_1} \cap C_{\delta_2} \Rightarrow C_{\delta_1} \cap \alpha = C_{\delta_2} \cap \alpha$
- (d) assume $\tau_0 \subseteq \tau_1 \subseteq \tau_2$ are vocabularies of cardinality $\leq \theta$, $\tau_1 \setminus \tau_0$ has only predicates, $\tau_2 \setminus \tau_1$ has only function symbols (allowed to be partial) $\mathfrak{B}$ is a τ_0 -model with universe λ (if $\lambda = \lambda^\theta$ also individual constants) then we can find $\langle \mathcal{M}_\delta : \delta \in S \rangle$ such that
 - (α) every $M \in \mathcal{M}_\delta$ is a τ_2 -model of cardinality θ expanding $\mathfrak{B} \upharpoonright |M_\delta|$
 - (β) if $M \in \mathcal{M}_\delta$, $F \in \tau_2 \setminus \tau_1$ then F^M has domain $\subseteq C_\delta$ (i.e., $\text{arity } F(C_\delta)$)
 - (γ) every $M \in \mathcal{M}_\delta$ has universe which includes C_δ and is included in δ and the universe of M is the $\mathfrak{B}$ -closure of $C_\delta \cup \{F(\bar{\alpha}) : F \in \tau_2 \setminus \tau_1 \text{ and } \bar{\alpha} \in {}^{\text{arity}(F)}(C_\delta)\}$
 - (δ) if $M', M'' \in \mathcal{M}_\delta$ then $(M', \gamma)_{\gamma \in C_\delta}, (M'', \gamma)_{\gamma \in C_\delta}$ are isomorphic
 - (ε) if $\mathfrak{B}^+$ is a τ_2 -expansion of $\mathfrak{B}$ then for stationarily many $\delta \in S$ for some $M \in \mathcal{M}_\delta$ we have:
 - (i) $F \in \tau_2 \setminus \tau_2 \Rightarrow F^{\mathfrak{B}^+} \upharpoonright C_\delta = F^M \upharpoonright C_\delta (= F^M)$
 - (ii) $M \upharpoonright \tau_1 \subseteq \mathfrak{B}^+ \upharpoonright \tau_1$.

4.9 Observation. 1) In 4.8 if the order $<$ on λ is a relation of $\mathfrak{B}$ (which is no loss) then the isomorphism is unique as it is necessarily the unique order preserving function from $|M'|$ onto $|M''|$.

2) In 4.8, if the function F_i where $\alpha < \beta \in C_\delta, \alpha \in C_\delta, \text{otp}(C_\delta \cap \alpha) = i \Rightarrow F_i(\beta) = \alpha$, then for any $M \in \cup\{\mathcal{M}_\delta : \delta \in S\}$ and $\delta, M \cap C_\delta$ is an initial segment of C_δ .

4.10 Definition. We say that $\bar{C}$ exemplifies $\text{BB}_1(\lambda, \kappa, \theta)$ when (a),(b),(d),(e) from 4.8 holds + (ε) below. $\text{BB}_2(\lambda, \kappa, \theta)$ holds when we add (ζ) to clause (d) where

- (ε) the isomorphism type of $(M, \gamma)_{\gamma \in C_\delta}$ for $M \in \mathcal{M}_\delta$ depend on τ_0, τ_1, τ_1 but not on $\mathfrak{B}$
- (ζ) if $M', M'' \in \mathcal{M}_\delta$ and π is an isomorphism from M' onto M'' and $\delta', \delta'' \in S, C_{\delta'} \subseteq M', C_{\delta''} \subseteq M''$ and π maps $C_{\delta'}$ onto $C_{\delta''}$, then for any $N' \in \mathcal{M}_{\delta'}, N'' \in \mathcal{M}_{\delta''}$ we have $(N_{\delta'}, \gamma)_{\gamma \in C_{\delta'}} \cong (N_{\delta''}, \gamma)_{\gamma \in C_{\delta''}}$.

4.11 Claim. If $\mu > \aleph_0$ is strong limit and $\lambda = \text{cf}(2^\mu)$ or $\lambda > 2^{2^\mu}$ not strongly inaccessible then for all but finitely many $\kappa \in \text{Reg} \cap \theta$ ($\kappa \in \text{Reg} \cap \mu \setminus \mathfrak{d}'_1(2^\mu)$) for every $\theta < \mu$, $\text{BB}_1(\lambda, \kappa, \theta)$ holds.

Proof. Use also 4.13 below.

4.12 Observation. 1) If $\bar{C}$ exemplifies $\text{BB}_\ell(\lambda, \kappa, \theta)$ then for some pairwise disjoint $\langle S_\varepsilon : \varepsilon < \lambda \rangle$ we have each $\bar{C} \upharpoonright S_\varepsilon$ exemplifies $\text{BB}_\ell(\lambda, \kappa, \theta)$.

2) If $\lambda = \lambda^\theta$ we can allow in $\tau_1 \setminus \tau_0$ individual constant.

We delay their proof as we first use them.

Now we turn to proving 4.11, 4.12.

4.13 Claim. 1) If $\bar{C}$ exemplifies $\text{MD}(\lambda, \kappa, 2^\theta, \lambda)$ then $\bar{C}$ exemplifies $\text{BB}_1(\lambda, \kappa, \theta)$ [Rethink: if we use $C * \chi, \chi = \sqsupset_\kappa$ enough to have many guess.]

2) $\bar{C}$ exemplifies $\text{BB}_1(\lambda, \kappa, \theta)$ when there are $\lambda_1, \bar{C}^1$

- (a) $\bar{C}$ exemplifies $\text{MD}(\lambda, \kappa, 2^\theta, \lambda)$ (hence $\bar{C}^1 = \langle C_\delta^1 : \delta \in S_1 \rangle$ exemplifies $\text{BB}_1(\lambda, \kappa, \theta)$ but apparently we need more
- (b) $\bar{h} = \langle h_\delta : \delta \in S_1 \rangle$ where h_δ is an increasing function from C_δ onto some $\gamma = \gamma(\delta) \in S_1$
- (c) for every club C of λ there is an increasing continuous function g from λ_1 into C such that $\alpha \in S_1 \Rightarrow g(\alpha) \in S$ & $\gamma_{g(\alpha)} = \alpha$.

3) If $\bar{C}$ exemplifies $\text{MD}(\lambda, \kappa, 2^\theta)$ then $\bar{C}$ exemplifies $\text{BB}_2(\lambda, \kappa, \theta)$.

Proof. 1) $\bar{C}$ has the $(D, 2^\mu, \theta)$ -Md-property (which is like the desired conclusion except that we write $F_\delta(\nu \restriction C_\delta)$ instead of $F(\nu \restriction C_\delta, \bar{C} \restriction \bar{C}_\delta)$). But let $\beta = \alpha/\theta$ mean that $\theta\beta \leq \alpha < \theta\beta + 1$. But define $F'_\delta(\nu) = F_\delta(\langle \nu(\alpha)/\theta : \alpha \in C_\delta \rangle, \langle \nu(\alpha) - \theta(\nu(\alpha)/\theta) : \alpha \in C_\delta \rangle)$. So for $\langle F'_\delta : \delta \in S \rangle$ we have $\bar{c}$ as required in the original requirement; the same $\bar{c}$ is as required for our $\bar{F}$.

2), 3) Left to the reader. □_{4.13}

4.14 Conclusion. If $\lambda = \text{cf}(\lambda) > \beth_{\omega+3}$ not strongly inaccessible, then for every regular $\kappa < \beth_\omega$ except possibly finitely many we have:

⊗ for some topological space X and $\bar{C} = \langle C_\delta : \delta \in S \rangle$ we have

(a) X is Hausdorff with a clopen basis set

(b) every $Y \subseteq X$ of cardinality $< \kappa$ is closed

(c) every point has a neighborhood of cardinality $\leq \kappa$

(d) there is $f : X \rightarrow \kappa$ such that:

if $X = \bigcup_{\alpha < \beta} X_\alpha, \beta < \alpha$ then some non-isolated point x has a neighborhood included in $x_{f(x)}$ (so $f(x) < \beta$).

REFERENCES.

- [CDS_{Sh} 571] James Cummings, Mirna Džamonja, and Saharon Shelah. A consistency result on weak reflection. *Fundamenta Mathematicae*, **148**:91–100, 1995. math.LO/9504221.
- [DjSh 562] Mirna Džamonja and Saharon Shelah. On squares, outside guessing of clubs and $I_{<f}[\lambda]$. *Fundamenta Mathematicae*, **148**:165–198, 1995. math.LO/9510216.
- [DjSh 692] Mirna Džamonja and Saharon Shelah. On $\triangleleft^*$ -maximality. *Annals of Pure and Applied Logic*, **125**:119–158, 2004. math.LO/0009087.
- [EH] Paul Erdős and Andras Hajnal. Unsolved Problems in set theory. In *Axiomatic Set Theory*, volume XIII Part I of *Proc. of Symp. in Pure Math.*, pages 17–48, Providence, R.I., 1971. AMS.
- [Sh 775] Saharon Shelah. Middle Diamond. *Archive for Mathematical Logic*, **submitted**. math.LO/0212249.
- [Sh:e] Saharon Shelah. *Non-structure theory*, accepted. Oxford University Press.
- [Sh 835] Saharon Shelah. PCF without choice. *Preprint*.
- [Sh 108] Saharon Shelah. On successors of singular cardinals. In *Logic Colloquium '78 (Mons, 1978)*, volume 97 of *Stud. Logic Foundations Math*, pages 357–380. North-Holland, Amsterdam-New York, 1979.
- [Sh 71] Saharon Shelah. A note on cardinal exponentiation. *The Journal of Symbolic Logic*, **45**:56–66, 1980.
- [Sh 110] Saharon Shelah. Better quasi-orders for uncountable cardinals. *Israel Journal of Mathematics*, **42**:177–226, 1982.
- [Sh:b] Saharon Shelah. *Proper forcing*, volume 940 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, xxix+496 pp, 1982.
- [Sh 186] Saharon Shelah. Diamonds, uniformization. *The Journal of Symbolic Logic*, **49**:1022–1033, 1984.
- [Sh 351] Saharon Shelah. Reflecting stationary sets and successors of singular cardinals. *Archive for Mathematical Logic*, **31**:25–53, 1991.
- [Sh 420] Saharon Shelah. Advances in Cardinal Arithmetic. In *Finite and Infinite Combinatorics in Sets and Logic*, pages 355–383. Kluwer Academic Publishers, 1993. N.W. Sauer et al (eds.).

- [Sh 386] Saharon Shelah. Bounding $pp(\mu)$ when $cf(\mu) > \mu > \aleph_0$ using ranks and normal ideals. In *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*, chapter VI. Oxford University Press, 1994.
- [Sh:g] Saharon Shelah. *Cardinal Arithmetic*, volume 29 of *Oxford Logic Guides*. Oxford University Press, 1994.
- [Sh 506] Saharon Shelah. The pcf-theorem revisited. In *The Mathematics of Paul Erdős, II*, volume 14 of *Algorithms and Combinatorics*, pages 420–459. Springer, 1997. Graham, Nešetřil, eds.. math.LO/9502233.
- [Sh 589] Saharon Shelah. Applications of PCF theory. *Journal of Symbolic Logic*, **65**:1624–1674, 2000.
- [Sh 460] Saharon Shelah. The Generalized Continuum Hypothesis revisited. *Israel Journal of Mathematics*, **116**:285–321, 2000. math.LO/9809200.
- [Sh 513] Saharon Shelah. PCF and infinite free subsets in an algebra. *Archive for Mathematical Logic*, **41**:321–359, 2002. math.LO/9807177.