

# Finite blocking property *vs* pure periodicity

Thierry Monteil \*

## Abstract

A translation surface  $\mathcal{S}$  is said to have the finite blocking property if for every pair  $(O, A)$  of points in  $\mathcal{S}$  there exists a finite number of “blocking” points  $B_1, \dots, B_n$  such that every geodesic from  $O$  to  $A$  meets one of the  $B_i$ ’s.  $\mathcal{S}$  is said to be purely periodic if the directional flow is periodic in each direction whose directional flow contains a periodic trajectory (this implies that  $\mathcal{S}$  admits a cylinder decomposition in such directions). We will prove that finite blocking property implies pure periodicity. We will also classify the surfaces that have the finite blocking property in genus 2: such surfaces are exactly the torus branched coverings. Moreover, we prove that in every stratum, such surfaces form a set of null measure.

*Keywords:* *blocking property, polygonal billiards, translation surfaces, Veech surfaces, torus branched covering, complete periodicity, illumination, quadratic differentials.*

---

\*Institut de Mathématiques de Luminy, CNRS UMR 6206, Case 907, 163 Avenue de Luminy, 13288 Marseille cedex 09, France – E-Mail: [monteil@iml.univ-mrs.fr](mailto:monteil@iml.univ-mrs.fr) – Tel: +33 4 91 26 96 77 – Fax : +33 4 91 26 96 55 –

# Introduction

A *translation surface* is a triple  $(\mathcal{S}, \Sigma, \omega)$  such that  $\mathcal{S}$  is a topological compact connected surface,  $\Sigma$  is a finite subset of  $\mathcal{S}$  (whose elements are called *singularities*) and  $\omega = (U_i, \phi_i)_{i \in I}$  is an atlas of  $\mathcal{S} \setminus \Sigma$  (consistent with the topological structure on  $\mathcal{S}$ ) such that the transition maps (i.e. the  $\phi_j \circ \phi_i^{-1} : \phi_i(U_i \cap U_j) \rightarrow \phi_j(U_i \cap U_j)$  for  $(i, j) \in I^2$ ) are translations. This atlas gives to  $\mathcal{S} \setminus \Sigma$  a Riemannian structure; therefore, we have notions of length, angle, measure, geodesic... Moreover, we assume that  $\mathcal{S}$  is the completion of  $\mathcal{S} \setminus \Sigma$  for this metric.

A translation surface can also be seen as a holomorphic differential  $h$  on a Riemann surface (the singularities correspond to the zeroes of the differential, and in an admissible atlas  $\omega$ ,  $h$  is of the form  $h = dz$ ). This point of view is useful to give coordinates (and therefore a topology and a measure) to the moduli space of all translation surfaces, while the first one allows to make pictures in the plane.

Translation surfaces provide one of the main tool for the study of rational polygonal billiards.

Since the unit tangent bundle of  $\mathcal{S}$  enjoys a canonical global decomposition  $U\mathcal{S} = \mathcal{S} \times \mathbb{S}^1$ , the study of the geodesic flow on  $\mathcal{S}$  can be done through two points of view depending on whether the variable is the first or the second projection:

1. We can fix one particular direction  $\theta \in \mathbb{S}^1$ : this corresponds to the study of the *directional flow*  $\phi_\theta : \mathcal{S} \times \mathbb{R} \rightarrow \mathcal{S}$ . We are usually interested in the ergodic properties of this flow depending on whether  $\theta$  is a saddle connection direction or not, or on the way  $\theta$  can be approximated by such saddle connections (see [KMS], [Ve], [MT], [Vo], [Ma], [MS], [Ch]). This class of problems is called *dynamical problems*. This point of view is the most studied by mathematicians.
2. We can also fix one point in  $x \in \mathcal{S}$ : this corresponds to the study of the *exponential flow*  $\exp_x : \mathbb{S}^1 \times \mathbb{R} \rightarrow \mathcal{S}$ . We are usually interested in looking at which points we can reach when we let the second variable move, or at the statistics of the return to (or near)  $x$  (see [Kl], [To], [Mo1], [Mo2], [BGZ], [CG]). This class of problems is called *illumination problems*. This point of view is connected with computer science (computational geometry, ray tracing, art gallery).

Note that those two flows are defined on a dense  $G_\delta$ -set of full measure but not everywhere because the singularities don't allow some (few) geodesic to be defined on  $\mathbb{R}$ .

We are interested here by the connection between an illumination problem called the finite blocking property, and a dynamical one named the pure periodicity.

A translation surface  $\mathcal{S}$  is said to have the *finite blocking property* if for every pair  $(O, A)$  of points in  $\mathcal{S}$ , there exists a finite number of points  $B_1, \dots, B_n$  (different from  $O$  and  $A$ ) such that every geodesic from  $O$  to  $A$  meets one of the  $B_i$ 's.

A translation surface  $\mathcal{S}$  is said to be *purely periodic* if for any  $\theta \in \mathbb{S}^1$ , the existence of a periodic orbit in the direction  $\theta$  implies that the directional flow  $\phi_\theta$  is periodic (i.e. there exists  $T > 0$  such that  $\phi_\theta^T = Id_{\mathcal{S}}$  a.e.). This notion is a stronger version of the complete periodicity defined in [Ca].

The paper is organized as follows :

In the first section, we recall some background and known results about the finite blocking property. The second one is devoted to the proof of the following result :

**Theorem 1.** *Let  $\mathcal{S}$  be a translation surface with the finite blocking property. Then  $\mathcal{S}$  is purely periodic.*

The third section is devoted to some applications of this theorem. First, we give an improvement of [Mo2], Theorem 3 :

**Theorem 2.** *In genus  $g \geq 2$ , the set of translation surfaces that fail the finite blocking property is of full measure in every stratum.*

Then, we give the complete classification in genus 2 :

**Theorem 3.** *Let  $\mathcal{S}$  be a translation surface of genus 2. Then  $\mathcal{S}$  has the finite blocking property if and only if  $\mathcal{S}$  is a torus branched covering.*

# 1 Background

In this section, we present some results about the finite blocking property that were proven in [Mo2].

## 1.1 Stability

The finite blocking property is stable under some classical operations on translation surfaces:

**Singularities** We have used the convention that a geodesic stops when it reaches a singularity. This convention is not restrictive for the study of the finite blocking property since we can add those singularities in any blocking configuration. Moreover the finite blocking property is not perturbed when we add or remove removable singularities (a singularity  $\sigma \in \Sigma$  is said to be *removable* if there exists an atlas  $\omega' \supset \omega$  such that  $(\mathcal{S}, \Sigma \setminus \{\sigma\}, \omega')$  is a translation surface).

**Branched covering** A *branched translation covering* between two translation surfaces is a map  $\pi : (\mathcal{S}, \Sigma) \rightarrow (\mathcal{S}', \Sigma')$  that is a topological branched covering that locally preserves the translation structure. A *torus branched covering* is a translation surface  $\mathcal{S}$  such that there exists a branched translation covering from  $\mathcal{S}$  to a flat torus.

**Proposition 1.** *Let  $\pi : (\mathcal{S}, \Sigma) \rightarrow (\mathcal{S}', \Sigma')$  be a translation covering branched on a finite set  $\mathcal{R}' \subset \mathcal{S}'$ . Then  $\mathcal{S}$  has the finite blocking property if and only if  $\mathcal{S}'$  has.*

**Rational billiards** Let  $\mathcal{P}$  denote a polygon in  $\mathbb{R}^2$ . Let  $\Gamma \subset O(2, \mathbb{R})$  be the group generated by the linear parts of the reflections in the sides of  $\mathcal{P}$ . When  $\Gamma$  is finite, we say that  $\mathcal{P}$  is a *rational polygonal billiard table*. When  $\mathcal{P}$  is simply connected,  $\mathcal{P}$  is rational if and only if all the angles between edges are rational multiples of  $\pi$ . A classical construction due to Zemljakov and Katok (see [ZK], [MT]) allows us to associate to each rational billiard table  $\mathcal{P}$  a translation surface  $ZK(\mathcal{P})$  (as an example, the translation surface associated to the square billiard table is the torus  $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ ).

**Proposition 2.** *Let  $\mathcal{P}$  be a rational polygonal billiard table. Then  $ZK(\mathcal{P})$  has the finite blocking property if and only if  $\mathcal{P}$  has.*

**Action of  $GL(2, \mathbb{R})$**  If  $A \in GL(2, \mathbb{R})$ , we can define the translation surface

$$A.(\mathcal{S}, \Sigma, (U_i, \phi_i)_{i \in I}) \stackrel{\text{def}}{=} (\mathcal{S}, \Sigma, (U_i, A \circ \phi_i)_{i \in I});$$

hence we have an action of  $GL(2, \mathbb{R})$  on the class of translation surfaces. We classically consider only elements of  $SL(2, \mathbb{R})$  (see [MT]), but area is not assumed to be preserved here.

**Proposition 3.** *Let  $\mathcal{S}$  be a translation surface and  $A$  be in  $GL(2, \mathbb{R})$ . Then  $\mathcal{S}$  has the finite blocking property if and only if  $A.\mathcal{S}$  has.*

## 1.2 Sufficient conditions

**Proposition 4 (Fomin [Fo]).** *The square billiard table has the finite blocking property.*

Hence, combining propositions 1, 2, 3 and 4, we have :

**Proposition 5.** *Any torus branched covering has the finite blocking property.*

## 1.3 Necessary conditions

We first give some definitions and conventions concerning isometries and cylinders. In this paper, “isometry” means “orientation-preserving local isometry”. Since an isometry  $i$  from an open subset  $U$  of  $\mathbb{R}^2$  to a translation surface  $\mathcal{S}$  can be uniquely extended by continuity to  $\overline{U}$  (it is uniformly continuous), we will also denote by  $i$  this extension (that is also an isometry). In this process, we may lose the injectivity of  $i$ .

A *subcylinder*  $\mathcal{C}$  is an isometric copy of  $\mathbb{R}/w\mathbb{Z} \times ]0, h[$  in a translation surface  $\mathcal{S}$  ( $w > 0$ ,  $h > 0$ ).  $w$  and  $h$  are unique and called the *width* and the *height* of  $\mathcal{C}$ . The *direction* of  $\mathcal{C}$  is the direction of the image of  $\mathbb{R}/w\mathbb{Z} \times \{h/2\}$  (which is considered here as an oriented closed geodesic); it is defined modulo  $2\pi$ . The images in  $\mathcal{S}$  of  $\mathbb{R}/w\mathbb{Z} \times \{0\}$  and  $\mathbb{R}/w\mathbb{Z} \times \{h\}$  are called the *sides* of  $\mathcal{C}$ . The side  $\mathbb{R}/w\mathbb{Z} \times \{h\}$  is called the *upper side*.

A *cylinder* is a maximal subcylinder (for the inclusion). Any periodic trajectory for  $\phi_\theta$  can be thickened to obtain a cylinder in the direction  $\theta$ . Whenever  $\mathcal{S}$  is not a torus, each side of a cylinder contains at least one singularity and is a finite union of saddle connections (a *saddle connection* is a geodesic joining two singularities). We say that  $\mathcal{S}$  admits a *cylinder decomposition* in the direction  $\theta$  if the (necessarily finite) union of cylinders whose direction is  $\theta$  is dense in  $\mathcal{S}$  (then, the complement of this union is the union of the sides of those cylinders).

Let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  be two cylinders with width  $w_1$  (resp.  $w_2$ ) and height  $h_1$  (resp.  $h_2$ ) in a translation surface  $\mathcal{S}$ . We suppose that  $\mathcal{C}_1$  and  $\mathcal{C}_2$  are two different cylinders in the same direction whose closures have a nontrivial (i.e. not reduced to a finite set) intersection. The situation is described in Figure 1.




Figure 1: The opposite vertical sides are identified by translation while the dotted horizontal ones are glued with the rest of the surface;  $l > 0$ .

**Lemma 1.** *Let  $\mathcal{S}$  be a translation surface that contains two different cylinders  $\mathcal{C}_1$  and  $\mathcal{C}_2$  in the same direction whose closures have a nontrivial intersection. If their widths are uncommensurable, then  $\mathcal{S}$  fails the finite blocking property.*

This lemma implies all the remaining results of section 1 and will be useful in the proof of Theorem 1.

**Proposition 6.** *The only regular polygons (considered as billiard tables) with the finite blocking property are the equilateral triangle, the square and the regular hexagon.*

**Proposition 7.** *A Veech surface has the finite blocking property if and only if it is a torus covering, branched over only one point.*

In other terms, arithmetic surfaces are exactly the Veech surfaces with the finite blocking property. Note that proposition 7 was proven independently and later by Eugene Gutkin, and is the main result of [Gu].

## 1.4 Global point of view: moduli spaces of holomorphic forms

A singularity  $\sigma \in \Sigma$  has a conical angle of the form  $2(k+1)\pi$ , with  $k \geq 0$ ; we say that  $\sigma$  is of *multiplicity*  $k$ . Removable singularities are the singularities of multiplicity 0. In terms of holomorphic differentials, it is equivalent to say that there is a chart around  $\sigma$  such that  $h = z^k dz$  (i.e.  $\sigma$  is a zero of order  $k$  of  $h$ ).

If  $1 \leq k_1 \leq k_2 \leq \dots \leq k_n$  is a sequence of integers whose sum is even, we denote by  $\mathcal{H}(k_1, k_2, \dots, k_n)$  the *stratum* of translation surfaces with exactly  $n$  singularities whose multiplicities are  $k_1, k_2, \dots, k_n$  (we consider only surfaces without removable singularities). A translation surface in  $\mathcal{H}(k_1, k_2, \dots, k_n)$  has genus  $g = 1 + (k_1 + k_2 + \dots + k_n)/2$ . Each stratum carries a natural topology and an  $SL(2, \mathbb{R})$ -invariant measure that are for example defined in [Ko].

**Proposition 8.** *In genus  $g \geq 2$ , the set of translation surfaces that fail the finite blocking property is dense in every stratum.*

In fact, we proved in [Mo2] that the hypotheses of Lemma 1 are satisfied on a dense subset of each stratum. Unfortunately, those hypotheses are valid only on a set of zero measure. Proposition 8 will be considerably improved by Theorem 2.

## 2 Main result

A translation surface is said to be *completely periodic* if the following property holds (see [Ca]): if there exists a periodic orbit in the direction  $\theta$ , then the surface admits a cylinder decomposition in the direction  $\theta$ .

**Proposition 9.** *Let  $\mathcal{S}$  be a translation surface with the finite blocking property. Then  $\mathcal{S}$  is completely periodic.*

**Proof:** Let  $\theta$  be a direction that contains a periodic orbit. Up to a rotation, we assume that this direction is the horizontal one (defined by  $\theta = 0 \bmod 2\pi$ ). We can thicken all of such horizontal periodic orbits to obtain a finite number of cylinders. Assume by contradiction, that  $\mathcal{S}$  is not fully decomposed into cylinders in the horizontal direction: there exists a cylinder  $\mathcal{C}$  and singularities  $\sigma$  and  $\sigma'$  (possibly equal) that lie on the upper side of  $\mathcal{C}$  such that the horizontal geodesic  $\gamma$  (whose length is denoted by  $l(\gamma)$  and image by  $|\gamma|$ ) from  $\sigma$  to  $\sigma'$  exists (the singularities are consecutive for the cyclic order on the upper side of  $\mathcal{C}$ ) and is not a part of a side of any other cylinder. To be more precise, there exists  $s > 0$  (by compactity of  $|\gamma|$  and by finiteness of the number of horizontal cylinders) such that there is an injective isometry  $i : ]0, l(\gamma)[ \times ]0, s[ \rightarrow \mathcal{S}$  such that  $i([0, l(\gamma)] \times \{0\}) = |\gamma|$  and  $i(]0, l(\gamma)[ \times ]0, s[)$  doesn't meet any horizontal cylinder. Up to a dilatation (that belongs to  $GL^+(2, \mathbb{R})$ ), we assume that the width of  $\mathcal{C}$  is 1 and that its height is 2. The situation is described in Figure 2.




Figure 2: The opposite vertical black sides are identified by translation; the dotted ones are glued with the rest of the surface; the dark zone doesn't meet any horizontal cylinder.

The heart of the proof is organised into three steps. A summary concludes each of them.

**Step 1** *Construction of certain thin rectangles in  $\mathcal{S}$  that contain a common point  $A$ .*

Let  $\varepsilon > 0$  be smaller than  $s$ .

Let  $T_\varepsilon$  be the supremum of all  $t > 0$  such that there exists an injective isometry  $j_t : ]0, t[ \times ]0, \varepsilon[ \rightarrow \mathcal{S}$  that coincides with  $\gamma$  on  $]0, \min(l(\gamma), t)[ \times \{0\}$ .  $T_\varepsilon$  is greater than or equal to  $l(\gamma)$  and therefore strictly positive.  $\mathcal{S}$  has finite area hence  $T_\varepsilon$  is

finite. Moreover  $T_\varepsilon$  is a maximum (indeed, if  $0 < t \leq t' < T_\varepsilon$ , then  $j_{t'}$  extends  $j_t$ ): there exists an injective isometry  $j_{T_\varepsilon} : ]0, T_\varepsilon[ \times ]0, \varepsilon[ \rightarrow \mathcal{S}$  that coincide with  $\gamma$  on  $]0, l(\gamma)[ \times \{0\}$ .

There are only three possibilities depending on what happens on  $j_{T_\varepsilon}(\{T_\varepsilon\} \times ]0, \varepsilon[)$  (if  $j_{T_\varepsilon}(\{T_\varepsilon\} \times ]0, \varepsilon[)$  does not meet a singularity, then  $j_{T_\varepsilon}$  can be extended to an isometry that fails to be injective, hence  $I_\varepsilon = j_{T_\varepsilon}(\{T_\varepsilon\} \times ]0, \varepsilon[)$  and  $J_\varepsilon = j_{T_\varepsilon}(\{0\} \times ]0, \varepsilon[)$  intersect each other):

1.  $I_\varepsilon$  meets a singularity denoted by  $\sigma_\varepsilon$ .
2.  $I_\varepsilon = J_\varepsilon$ : we just have constructed a cylinder where it is forbidden, so this case does not occur.
3. The vertical intervals  $I_\varepsilon$  and  $J_\varepsilon$  intersect each other but we are neither in the first nor in the second case. Since  $\sigma = j_{T_\varepsilon}(0, 0)$  is a singularity,  $I_\varepsilon \cap J_\varepsilon$  is a subinterval of  $J_\varepsilon$  that hits the extremity of  $J_\varepsilon$  that is not  $\sigma$  (i.e.  $j_{T_\varepsilon}(0, \varepsilon)$ ) (see Figure 3).




Figure 3: The dark zone is the image of  $j_{T_\varepsilon}$ .

So, there exists  $0 < \varepsilon' \leq \varepsilon$  such that  $J_\varepsilon \setminus I_\varepsilon = j_{T_\varepsilon}(\{0\} \times ]0, \varepsilon'])$ . Now, the same three possibilities are offered to  $j_{T_{\varepsilon'}}$ , but the third case can't happen. Indeed, otherwise there is  $0 < \varepsilon'' < \varepsilon'$  such that  $j_{T_{\varepsilon'}}(0, \varepsilon') = j_{T_{\varepsilon'}}(T_{\varepsilon'}, 0) = j_{T_{\varepsilon'}}(T_{\varepsilon'}, \varepsilon' - \varepsilon'')$  is not a singularity but has a conical angle greater than  $2\pi$ .

So, up to reducing the size of  $\varepsilon$  (by replacing  $\varepsilon$  by  $\varepsilon'$  whenever we are in the third case), we can consider that we are in the first case, and, since  $\Sigma$  is finite, there exists a sequence  $(\varepsilon_n)_{n \in \mathbb{N}}$  of strictly positive numbers that converges to 0 and a singularity  $\sigma_\infty \in \Sigma$  such that for all  $n$  in  $\mathbb{N}$ ,  $\sigma_{\varepsilon_n} = \sigma_\infty$ . If  $n \in \mathbb{N}$ , there exists  $0 < y_n < \varepsilon_n$  such that  $\sigma_\infty = j_{T_{\varepsilon_n}}(T_{\varepsilon_n}, y_n)$ . Let  $A = j_{T_{\varepsilon_n}}(T_{\varepsilon_n} - l(\gamma)/2, y_n)$  ( $A$  does not depend on  $n \in \mathbb{N}$  and is in the image of every  $j_{T_{\varepsilon_n}}$ ).



To sum up this step and fix notations, we proved that there exists  $A \in \mathcal{S}$  such that for all  $\varepsilon > 0$  there exists  $l_\varepsilon \geq l(\gamma)$  and  $0 < h_\varepsilon \leq \varepsilon$  such that there exists an injective isometry  $i_\varepsilon$  from  $]0, l_\varepsilon[ \times ]0, h_\varepsilon[$  to  $\mathcal{S}$  that coincides with  $\gamma$  on  $]0, l(\gamma)[ \times \{0\}$  and whose image contains  $A$ .

The situation is summarized in Figure 4. Let  $O$  be the point drawn on this Figure. We denote  $A = i_\varepsilon(x_\varepsilon, y_\varepsilon)$ .




Figure 4: Summary of Step 1.

**Step 2** Construction of “teepees” of more and more geodesics between  $O$  and  $A$ .

Let  $\varepsilon > 0$ . The unfolding of  $\mathcal{C}$  allows us to define an extension of  $i_\varepsilon$  as in Figure 5: we get an isometry  $i_\varepsilon : ]0, l_\varepsilon[ \times [0, h_\varepsilon[ \cup \mathbb{R} \times ]-2, 0[ \rightarrow \mathcal{S}$ . In  $\mathbb{R}^2$ , we denote without ambiguity  $\sigma = (0, 0)$ ,  $\sigma' = (l(\gamma), 0)$  and  $A = (x_\varepsilon, y_\varepsilon)$ .




Figure 5: First (topological) application of the theorem of Thales.

Let  $S_\varepsilon$  (resp.  $S'_\varepsilon$ ) be the point of the line that passes through  $A$  and  $\sigma$  (resp.  $\sigma'$ ) whose ordinate is  $-1$ . Since  $(\sigma, \sigma')$  and  $(S_\varepsilon, S'_\varepsilon)$  are parallel, the theorem of Thales

asserts that

$$\frac{d(S_\varepsilon, S'_\varepsilon)}{(1 + y_\varepsilon)} = \frac{l(\gamma)}{y_\varepsilon},$$

where  $d$  stands for the euclidean distance of  $\mathbb{R}^2$ . Let  $\mathcal{O}_\varepsilon = i_\varepsilon^{-1}(\{O\}) \cap ]S_\varepsilon, S'_\varepsilon[$ . We have

$$\text{card } \mathcal{O}_\varepsilon \geq d(S_\varepsilon, S'_\varepsilon) - 1 = l(\gamma)(1 + y_\varepsilon)/y_\varepsilon - 1 \xrightarrow{\varepsilon \rightarrow 0} \infty.$$

Since the open triangle  $(A, S_\varepsilon, S'_\varepsilon)$  is included in the source of  $i_\varepsilon$ , we define a set  $\mathcal{T}_\varepsilon$  (called a *teepee*) of geodesics in  $\mathcal{S}$  from  $O$  to  $A$  by taking the images by  $i_\varepsilon$  of the segments joining an element of  $\mathcal{O}_\varepsilon$  to  $A$ . Of course,

$$\text{card } \mathcal{T}_\varepsilon = \text{card } \mathcal{O}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0} \infty.$$

To sum up this step, we constructed, for all  $\varepsilon > 0$ , a set  $\mathcal{T}_\varepsilon$  of more and more geodesics from  $O$  to  $A$  whose images are included in  $\mathcal{C} \cup |\gamma| \cup i_\varepsilon([0, l_\varepsilon[ \times ]0, h_\varepsilon[)$ .

**Step 3** *The density of the geodesics of a teepee that are blocked by a fixed point vanishes.*

Let  $B$  be a fixed point in  $\mathcal{S}$  that is different from  $O$  and  $A$ . For  $\varepsilon > 0$ , let  $\mathcal{T}_\varepsilon(B)$  be the set of elements of  $\mathcal{T}_\varepsilon$  that are blocked by  $B$  (i.e. that pass through  $B$ ).

- Suppose that  $B$  is not in  $\mathcal{C}$ . Let  $\varepsilon > 0$ . If  $B$  is not in the image of  $i_\varepsilon$ , then  $\mathcal{T}_\varepsilon(B)$  is empty. Otherwise,  $B$  is in  $i_\varepsilon(]0, l_\varepsilon[ \times ]0, h_\varepsilon[)$ , then the cardinal of  $\mathcal{T}_\varepsilon(B)$  is at most 1 (indeed, the elements of  $\mathcal{T}_\varepsilon$  are disjoint on  $i_\varepsilon(]0, l_\varepsilon[ \times ]0, h_\varepsilon[) \setminus \{A\}$ ). Hence,  $\text{card } \mathcal{T}_\varepsilon(B) / \text{card } \mathcal{T}_\varepsilon \xrightarrow{\varepsilon \rightarrow 0} 0$ .
- Suppose that  $B$  is in  $\mathcal{C}$ . Let  $-h$  be the ordinate of any preimage of  $B$  by any  $i_\varepsilon$ . We can assume that  $h \in ]0, 1[$ , otherwise  $\mathcal{T}_\varepsilon(B)$  is empty for any  $\varepsilon > 0$ . Let  $\varepsilon > 0$ . Let  $\beta_1$  and  $\beta_2$  be two distinct elements of  $\mathcal{T}_\varepsilon(B)$ : for  $i \in \{1, 2\}$ , there exists  $O_i$  in  $\mathcal{O}_\varepsilon$  and  $B_i$  in  $i_\varepsilon^{-1}(B)$  such that  $B_i$  lies in the segment  $[O_i, A]$  whose image by  $i_\varepsilon$  is  $\beta_i$  (see Figure 6).




Figure 6: Second (arithmetic) application of the theorem of Thales.

We choose  $\beta_1$  and  $\beta_2$  such that  $d(O_1, O_2)$  is minimal, and set  $p_\varepsilon = d(B_1, B_2)$  and  $q_\varepsilon = d(O_1, O_2)$ . By minimality,

$$\text{card } \mathcal{T}_\varepsilon(B) \leq \frac{\text{card } \mathcal{T}_\varepsilon}{q_\varepsilon} + 1.$$

Since  $i_\varepsilon(B_1) = i_\varepsilon(B_2) = B$  and  $i_\varepsilon(O_1) = i_\varepsilon(O_2) = O$ ,  $p_\varepsilon$  and  $q_\varepsilon$  are integers. Since  $(B_1, B_2)$  and  $(O_1, O_2)$  are parallel, the theorem of Thales asserts that

$$\frac{p_\varepsilon}{q_\varepsilon} = \frac{(h + y_\varepsilon)}{(1 + y_\varepsilon)}.$$

The map  $\left( \begin{array}{ccc} \mathbb{R}_+^* & \longrightarrow & ]h, \infty[ \\ \varepsilon & \longmapsto & (h + y_\varepsilon)/(1 + y_\varepsilon) \end{array} \right)$  converges to  $h$  when  $\varepsilon$  converges to 0 but never takes the value  $h$ , hence  $q_\varepsilon \xrightarrow{\varepsilon \rightarrow 0} \infty$  (indeed, any set of rational numbers whose denominators are bounded is a closed discrete subset of  $\mathbb{R}$ , hence it cannot have an accumulation point).

Hence,

$$\frac{\text{card } \mathcal{T}_\varepsilon(B)}{\text{card } \mathcal{T}_\varepsilon} \leq \frac{1}{q_\varepsilon} + \frac{1}{\text{card } \mathcal{T}_\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} 0.$$

*To sum up this step, we proved that for any point  $B$  in  $\mathcal{S}$  that is different from  $O$  and  $A$ , the density of the geodesics of  $\mathcal{T}_\varepsilon$  that are blocked by  $B$  tends to 0 when  $\varepsilon$  tends to 0.*

Now, we assumed that  $\mathcal{S}$  has the finite blocking property, so there exists a finite set  $\{B_1, \dots, B_n\}$  in  $\mathcal{S} \setminus \{O, A\}$  that blocks any geodesic from  $O$  to  $A$ .

Let  $\varepsilon > 0$  be small enough such that  $\sum_{i=1}^n \text{card } \mathcal{T}_\varepsilon(B_i) / \text{card } \mathcal{T}_\varepsilon < 1$ : there exists a geodesic in  $\mathcal{T}_\varepsilon \setminus \bigcup_{i=1}^n \mathcal{T}_\varepsilon(B_i)$  from  $O$  to  $A$  that doesn't meet any  $B_i$ , leading to a contradiction. So,  $\mathcal{S}$  is completely periodic. □

**Theorem 1.** *Let  $\mathcal{S}$  be a translation surface with the finite blocking property. Then  $\mathcal{S}$  is purely periodic.*

**Proof:** Let  $\theta$  be a direction that contains a periodic trajectory. By virtue of Proposition 9,  $\mathcal{S}$  admits a cylinder decomposition  $(\mathcal{C}_i)_{i=1}^n$  in the direction  $\theta$ . Starting from one of those cylinders and applying Lemma 1 step by step, we show that all the widths of those cylinders are commensurable (remember that  $\mathcal{S}$  is connected). Hence there exists  $w > 0$  such that the width of  $\mathcal{C}_i$  is equal to  $p_i w$  with  $p_i \in \mathbb{N}^*$ . Setting  $T = w p_1 \dots p_n > 0$ , we have  $\phi_\theta^T = Id_{\mathcal{S}}$  a.e. □

### 3 Some applications

**Theorem 2.** *In genus  $g \geq 2$ , the set of translation surfaces that fail the finite blocking property is of full measure in every stratum.*

**Proof:** It is known that completely periodic translation surfaces form a set of null measure in every stratum (in genus  $g \geq 2$ ) ([Zo]). □

**Theorem 3.** *Let  $\mathcal{S}$  be a translation surface of genus 2. Then  $\mathcal{S}$  has the finite blocking property if and only if  $\mathcal{S}$  is a torus branched covering.*

**Proof:**

By Proposition 5, if  $\mathcal{S}$  is a torus branched covering, then  $\mathcal{S}$  has the finite blocking property. Conversely, suppose that  $\mathcal{S}$  has the finite blocking property. By Proposition 9,  $\mathcal{S}$  is completely periodic. Such surfaces of genus 2 were classified in [Ca]. If  $\mathcal{S}$  is in  $\mathcal{H}(2)$ , then  $\mathcal{S}$  is a Veech surface and by Proposition 7,  $\mathcal{S}$  is a torus branched covering. If  $\mathcal{S}$  is in  $\mathcal{H}(1, 1)$ , we assume by contradiction that  $\mathcal{S}$  is not a torus branched covering. A direct consequence of Theorem 1.2 of [Ca] is that after rescaling  $\mathcal{S}$ , there exists a square-free integer  $d > 0$ , four positive numbers  $w_1, w_2, s_1, s_2$  in  $\mathbb{Q}(\sqrt{d})$  and a direction  $\theta \in \mathbb{S}^1$  such that:

- $\mathcal{S}$  is decomposed into cylinders  $\mathcal{C}_1, \mathcal{C}_2$  (and sometimes  $\mathcal{C}_3$ ) in the direction  $\theta$
- $w_1$  (resp.  $w_2$ ) is the width of  $\mathcal{C}_1$  (resp.  $\mathcal{C}_2$ )
- 

$$\frac{w_1}{w_2}\bar{s}_1 + \bar{s}_2 = 0$$

Since  $w_1, w_2, s_1, s_2$  are in  $\mathbb{Q}(\sqrt{d})$ , there exists two rational numbers  $r_1$  and  $r_2$  such that

$$\frac{w_1}{w_2}s_1 + s_2 = r_1 + r_2\sqrt{d}.$$

The addition of those two equalities gives  $\frac{w_1}{w_2}(s_1 + \bar{s}_1) + (s_2 + \bar{s}_2) = r_1 + r_2\sqrt{d}$ . Theorem 1 asserts that  $\mathcal{S}$  is purely periodic, so  $\frac{w_1}{w_2}$  is rational, so  $\frac{w_1}{w_2}(s_1 + \bar{s}_1) + (s_2 + \bar{s}_2)$  is rational, so  $r_2 = 0$ . The subtraction of the two equalities leads to  $r_1 = 0$ .

Hence,  $\frac{w_1}{w_2}s_1 + s_2$  is both null and positive, which is impossible:  $\mathcal{S}$  is a torus branched covering. □

Of course, the notion of finite blocking property and the results presented in this paper can be translated in the vocabulary of quadratic differentials.

# Conclusion

We conclude this paper by the following open question :

Does pure periodicity imply being a torus branched covering?

It was positively answered in [Mo2] for Veech surfaces and in the present paper for surfaces of genus two. Proving that in full generality would provide an equivalence between the following assertions :

1.  $\mathcal{S}$  is a torus branched covering
2. there exists  $A \in GL(2, \mathbb{R})$  such that  $hol(H_1(A.\mathcal{S}, \mathbb{Z})) = \mathbb{Z} + i\mathbb{Z}$
3.  $Vect_{\mathbb{Q}}(hol(H_1(\mathcal{S}, \mathbb{Z})))$  has dimension 2
4.  $\mathcal{S}$  has the finite blocking property
5.  $\mathcal{S}$  has the bounded blocking property: there exists  $n \in \mathbb{N}$  such that for every pair  $(O, A)$  of points in  $\mathcal{S}$ , there exists  $n$  points  $B_1, \dots, B_n$  (different from  $O$  and  $A$ ) such that every geodesic from  $O$  to  $A$  meets one of the  $B_i$ 's
6.  $\mathcal{S}$  has the middle blocking property: For every pair  $(O, A)$  of points in  $\mathcal{S}$ , the set of the midpoints of the geodesics from  $O$  to  $A$  is finite
7.  $\mathcal{S}$  has the bounded middle blocking property: there exists  $n \in \mathbb{N}$  such that the set of the midpoints of the geodesics from  $O$  to  $A$  has cardinal less than  $n$
8.  $\mathcal{S}$  is purely periodic: in every direction  $\theta$  that contains a periodic orbit, the directional flow  $\phi_\theta$  is periodic (i.e. there exists  $T > 0$  such that  $\phi_\theta^T = Id_{\mathcal{S}}$  a.e.)
9. in every direction that contains a periodic orbit,  $\mathcal{S}$  is decomposable into cylinders whose widths are commensurable

Properties 1, 2 and 3 are geometric, whereas properties 4, 5, 6, 7 are exponential and properties 8 and 9 are dynamic.

# References

- [BGZ] F. BOCA, R. GOLOGAN, A. ZAHARESCU, *The statistics of the trajectory of a certain billiard in a flat torus*, Comm. Math. Phys. 240 (2003), p 53–76.
- [Ca] K. CALTA, *Veech surfaces and complete periodicity in genus 2*, J. Amer. Math. Soc., to appear.
- [Ch] Y. CHEUNG, *Hausdorff dimension of the set of nonergodic directions*, Ann. Math., 158 (2003), 661–678.
- [CG] N. CHERNOV, G. GALPERIN, *Search light in billiard tables*, Regul. Chaotic Dyn. 8 (2003), no. 2, 225–241.

- [Fo] D. FOMIN, ЗАДАЧИ ЛЕНИНГРАДСКИХ МАТЕМАТИЧЕСКИХ ОЛИМПИАД, Leningrad (1990), 77.
- [Gu] E. GUTKIN, *Blocking of orbits and the phenomenon of (in)security for the billiard in polygons and flat surfaces*, preprint (2003).
- [HS] P. HIEMER, V. SNURNIKOV, *Polygonal billiards with small obstacles*, J. Statist. Phys. 90 (1998), no. 1-2, 453–466.
- [Kl] V. KLEE, *Is every polygonal region illuminable from some point?*, Amer. Math. Monthly 76(80) (1969).
- [KMS] S. KERCKHOFF, H. MASUR, J. SMILLIE, *Ergodicity of billiard flows and quadratic differentials*, Ann. Math. 124 (1986), no. 2, p 293–311.
- [Ko] M. KONTSEVICH, *Lyapunov exponents and Hodge theory*, The mathematical beauty of physics (Saclay, 1996), p 318–332, Adv. Ser. Math. Phys., 24, World Sci. Publishing, River Edge, NJ, 1997.
- [Ma] H. MASUR, *Hausdorff dimension of the set of nonergodic foliations of a quadratic differential*, Duke Math. J. 66 (1992), no. 3, p 387–442.
- [MS] H. MASUR, J. SMILLIE, *Hausdorff dimension of sets of nonergodic measured foliations*, Ann. Math. 134 (1991), no. 3, 455–543.
- [MT] H. MASUR, S. TABACHNIKOV, *Rational billiards and flat structures*, Handbook on dynamical systems, Vol. 1A, 1015–1089, North-Holland, Amsterdam, (2002).
- [Mo1] T. MONTEIL, *A counter-example to the theorem of Hiemer and Snurnikov*, J. Statist. Phys. 114 (2004), no. 5-6, 1619–1623.
- [Mo2] T. MONTEIL, *On the finite blocking property*, Ann. Inst. Fourier, to appear.
- [To] G. TOKARSKY, *Polygonal rooms not illuminable from every point*, Amer. Math. Monthly 102 (1995), no. 10, 867–879.
- [Ve] W. VEECH, *Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards*, Invent. Math. 97 (1989), no. 3, 553–583.
- [Vo] Y. VOROBETS, *Planar structures and billiards in rational polygons: the Veech alternative*, Russian Math. Surveys 51 (1996), no. 5, 779–817.
- [ZK] A. ZEMLJAKOV, A. KATOK, *Topological transitivity of billiards in polygons*, Mat. Zametki 18 (1975), no. 2, 291–300.
- [Zo] A. ZORICH, *private communication*.