

On the extreme eigenvalues of regular graphs

Sebastian M. Cioabă^{a,1}

^a*Department of Mathematics, Queen's University, Kingston, Ontario K7L 3N6,
Canada*

To the memory of Dom de Caen

Abstract

In this paper, we present an elementary proof of a theorem of Serre concerning the greatest eigenvalues of k -regular graphs. We also prove an analogue of Serre's theorem regarding the least eigenvalues of k -regular graphs: given $\epsilon > 0$, there exist a positive constant $c = c(\epsilon, k)$ and a nonnegative integer $g = g(\epsilon, k)$ such that for any k -regular graph X with no odd cycles of length less than g , the number of eigenvalues μ of X such that $\mu \leq -(2 - \epsilon)\sqrt{k - 1}$ is at least $c|X|$. This implies a result of Winnie Li.

Key words: Alon-Boppana theorem, eigenvalues of graphs

1 Preliminaries

Let X be a graph and let v_0 be a vertex of X . A *closed walk* in X of length $r \geq 0$ starting at v_0 is a sequence v_0, v_1, \dots, v_r of vertices of X such that $v_r = v_0$ and v_{i-1} is adjacent to v_i for $1 \leq i \leq r$. For $r \geq 0$, let $\Phi_r(X)$ denote

¹ Research partially supported by an Ontario Graduate Scholarship

the number of closed walks of length r in X . A *cycle* of length r in X is a subgraph of X whose vertices can be labeled v_0, \dots, v_r such that v_0, \dots, v_r is a closed walk in X and $v_i \neq v_j$ for all i, j with $0 \leq i < j \leq r$. The *girth*, denoted $\text{girth}(X)$, of X is the length of a smallest cycle in X if such a cycle exists and ∞ otherwise; the *oddgirth*, denoted $\text{oddg}(X)$, of X is the length of a smallest odd cycle in X if such a cycle exists and ∞ otherwise. The adjacency matrix of X is the matrix $A = A(X)$ of order $|X|$, where the (u, v) entry is 1 if the vertices u and v are adjacent and 0 otherwise. It is a well known fact that $\Phi_r(X) = \text{Tr}(A^r)$, for any $r \geq 0$. The eigenvalues of X are the eigenvalues of A . If X is k -regular, then it is easy to see that k is an eigenvalue of X with multiplicity equal to the number of components of X and that any eigenvalue λ of X satisfies $|\lambda| \leq k$. For $l \geq 1$, we denote by $\lambda_l(X)$ the l th greatest eigenvalue of X and by $\mu_l(X)$ the l th least eigenvalue of X .

2 An elementary proof of Serre's theorem

Serre has proved the following theorem (see [4,5,7,15]) using Chebyshev polynomials. See also [2] for related results. In this section, we present an elementary proof of Serre's result.

Theorem 1 *For each $\epsilon > 0$, there exists a positive constant $c = c(\epsilon, k)$ such that for any k -regular graph X , the number of eigenvalues λ of X with $\lambda \geq (2 - \epsilon)\sqrt{k - 1}$ is at least $c|X|$.*

For the proof of this theorem we require the next lemma which can be deduced from McKay's work [11], Lemma 2.1. For the sake of completeness, we include a short proof here.

Lemma 2 *Let v_0 be a vertex of a k -regular graph X . Then the number of closed walks of length $2s$ in X starting at v_0 is greater than or equal to $\frac{1}{s+1} \binom{2s}{s} k(k-1)^{s-1}$.*

Proof of Lemma 2 The number of closed walks of length $2s$ in X starting at v_0 is at least the number of closed walks of length $2s$ starting at a vertex u_0 in the infinite k -regular tree. To each closed walk in the infinite k -regular tree, there corresponds a sequence of nonnegative integers $\delta_1, \dots, \delta_{2s}$, where δ_i is the distance from u_0 after i steps. The number of such sequences is the s -th Catalan number $\frac{1}{s+1} \binom{2s}{s}$. For each sequence of distances, there are at least $k(k-1)^{s-1}$ closed walks of length $2s$ since for each step away from u_0 there are $k-1$ choices (k if the walk is at u_0).

□

By Stirling's bound on $s!$ or by a simple induction argument it is easy to see that $\binom{2s}{s} \geq \frac{4^s}{s+1}$, for any $s \geq 1$. Hence, for any k -regular graph X and for any $s \geq 1$, we have by Lemma 2

$$\mathrm{Tr}(A^{2s}) \geq |X| \frac{1}{s+1} \binom{2s}{s} k(k-1)^{s-1} > |X| \frac{1}{(s+1)^2} (2\sqrt{k-1})^{2s} \quad (1)$$

Proof of Theorem 1 Let X be k -regular graph of order n with eigenvalues $k = \lambda_1 \geq \dots \geq \lambda_n \geq -k$. Given $\epsilon > 0$, let m be the number of eigenvalues λ of X with $\lambda \geq (2-\epsilon)\sqrt{k-1}$. Then $n-m$ of the eigenvalues of X are less

than $(2 - \epsilon)\sqrt{k-1}$. Thus

$$\begin{aligned}\mathrm{Tr}(kI + A)^{2s} &= \sum_{i=1}^n (k + \lambda_i)^{2s} \\ &< (n - m)(k + (2 - \epsilon)\sqrt{k-1})^{2s} + m(2k)^{2s} \\ &= m((2k)^{2s} - (k + (2 - \epsilon)\sqrt{k-1})^{2s}) + n(k + (2 - \epsilon)\sqrt{k-1})^{2s}\end{aligned}$$

On the other hand, the binomial expansion and relation (1) give

$$\begin{aligned}\mathrm{Tr}(kI + A)^{2s} &= \sum_{i=0}^{2s} \binom{2s}{i} k^i \mathrm{Tr}(A^{2s-i}) \\ &\geq \sum_{j=0}^s \binom{2s}{2j} k^{2j} \mathrm{Tr}(A^{2s-2j}) \\ &> \frac{n}{(s+1)^2} \sum_{j=0}^s \binom{2s}{2j} k^{2j} (2\sqrt{k-1})^{2s-2j} \\ &= \frac{n}{2(s+1)^2} ((k + 2\sqrt{k-1})^{2s} + (k - 2\sqrt{k-1})^{2s}) \\ &> \frac{n}{2(s+1)^2} (k + 2\sqrt{k-1})^{2s}\end{aligned}$$

Thus,

$$\frac{m}{n} > \frac{\frac{1}{2(s+1)^2} (k + 2\sqrt{k-1})^{2s} - (k + (2 - \epsilon)\sqrt{k-1})^{2s}}{(2k)^{2s} - (k + (2 - \epsilon)\sqrt{k-1})^{2s}}$$

for any $s \geq 1$. Since

$$\begin{aligned}\lim_{s \rightarrow \infty} \left(\frac{(k + 2\sqrt{k-1})^{2s}}{2(s+1)^2} \right)^{\frac{1}{2s}} &= k + 2\sqrt{k-1} \\ &> k + (2 - \epsilon)\sqrt{k-1} = \lim_{s \rightarrow \infty} \left(2(k + (2 - \epsilon)\sqrt{k-1})^{2s} \right)^{\frac{1}{2s}}\end{aligned}$$

it follows that there exists $s_0 = s_0(\epsilon, k)$ such that for all $s \geq s_0$

$$\frac{(k + 2\sqrt{k-1})^{2s}}{2(s+1)^2} - (k + (2 - \epsilon)\sqrt{k-1})^{2s} > (k + (2 - \epsilon)\sqrt{k-1})^{2s}$$

Hence, if

$$c(\epsilon, k) = \frac{(k + (2 - \epsilon)\sqrt{k-1})^{2s_0}}{(2k)^{2s_0} - (k + (2 - \epsilon)\sqrt{k-1})^{2s_0}}$$

then $c(\epsilon, k) > 0$ and $m > c(\epsilon, k)n$.

□

The proofs of Serre's theorem given in [4,5,7] don't allow an easy estimation of the constant $c(\epsilon, k)$ in terms of ϵ and k . We relegate the detailed analysis of the constant obtained by those arguments to a future work [3]. We should mention that Serre's theorem can be also deduced from the work of Friedman [6] or Nilli [13]. Friedman's results imply an estimate of $\left(\frac{1}{2}\right)^{O\left(\frac{\log k}{\sqrt{\epsilon}}\right)}$ for the proportion of the eigenvalues that are at least $(2 - \epsilon)\sqrt{k-1}$. Nilli's work provides a bound of $\left(\frac{1}{2}\right)^{O\left(\frac{\log k}{\arccos(1-\epsilon)}\right)}$. Their methods provide better bounds on $c(\epsilon, k)$ than ours. From our proof of Serre's theorem, we obtain that a proportion of $\left(\frac{1}{2}\right)^{O\left(\frac{\sqrt{k}}{\epsilon} \log\left(\frac{\sqrt{k}}{\epsilon}\right)\right)}$ of the eigenvalues are at least $(2 - \epsilon)\sqrt{k-1}$. This is because in Theorem 1 we pick s_0 such that $\frac{s_0}{\log s_0} = \Theta\left(\frac{\sqrt{k}}{\epsilon}\right)$.

Theorem 1 has the following consequence regarding the asymptotics of the greatest eigenvalues of k -regular graphs.

Corollary 3 *Let $(X_i)_{i \geq 0}$ be a sequence of k -regular graphs such that*

$\lim_{i \rightarrow \infty} |X_i| = \infty$. Then for each $l \geq 1$,

$$\liminf_{i \rightarrow \infty} \lambda_l(X_i) \geq 2\sqrt{k-1}$$

This corollary has also been proved directly by Serre in an appendix to [8] using the eigenvalue distribution theorem in [16]. When $l = 2$, we obtain the asymptotic version of the Alon-Boppana theorem (see [1,10,12,14] for more details).

3 Analogous theorems for the least eigenvalues of regular graphs

The analogous result to Theorem 1 for the least eigenvalues of a k -regular graph is not true. For example, the eigenvalues of line graphs are all at least -2 . However, by adding an extra condition to the hypothesis of Theorem 1, we can prove an analogue of Serre's theorem for the least eigenvalues of a k -regular graph.

Theorem 4 *For any $\epsilon > 0$, there exist a positive constant $c = c(\epsilon, k)$ and a non-negative integer $g = g(\epsilon, k)$ such that for any k -regular graph X with $\text{oddg}(X) > g$, the number of eigenvalues μ of X with $\mu \leq -(2 - \epsilon)\sqrt{k - 1}$ is at least $c|X|$.*

Proof Let X be a k -regular graph of order n with eigenvalues $-k \leq \mu_1 \leq \mu_2 \leq \dots \leq \mu_n = k$. Given $\epsilon > 0$, let m be the number of eigenvalues μ of X with $\mu \leq -(2 - \epsilon)\sqrt{k - 1}$. Then $n - m$ of the eigenvalues of X are greater than $-(2 - \epsilon)\sqrt{k - 1}$. Thus

$$\begin{aligned} \text{Tr}(kI - A)^{2s} &= \sum_{i=1}^n (k - \mu_i)^{2s} < (n - m)(k + (2 - \epsilon)\sqrt{k - 1})^{2s} + m(2k)^{2s} \\ &= m((2k)^{2s} - (k + (2 - \epsilon)\sqrt{k - 1})^{2s}) + n(k + (2 - \epsilon)\sqrt{k - 1})^{2s} \end{aligned}$$

In the previous section, we proved that there exists $s_0 = s_0(\epsilon, k)$ such that for all $s \geq s_0$

$$\frac{(k + 2\sqrt{k - 1})^{2s_0}}{2(s_0 + 1)^2} - (k + (2 - \epsilon)\sqrt{k - 1})^{2s_0} > (k + (2 - \epsilon)\sqrt{k - 1})^{2s_0}$$

Let $g(\epsilon, k) = 2s_0$. If $\text{oddg}(X) > 2s_0$, then for $0 \leq j \leq s_0 - 1$, the number of closed walks of length $2s_0 - 2j - 1$ in X is 0. Hence, $\text{Tr}(A^{2s_0 - 2j - 1}) = 0$, for

$0 \leq j \leq s_0 - 1$. Using also (1), we obtain

$$\begin{aligned} \operatorname{Tr}(kI - A)^{2s_0} &= \sum_{j=0}^{s_0} \binom{2s_0}{2j} k^{2j} \operatorname{Tr}(A^{2s_0-2j}) - \sum_{j=0}^{s_0-1} \binom{2s_0}{2j+1} k^{2j+1} \operatorname{Tr}(A^{2s_0-2j-1}) \\ &= \sum_{j=0}^{s_0} \binom{2s_0}{2j} k^{2j} \operatorname{Tr}(A^{2s_0-2j}) > \frac{n}{(s_0+1)^2} \sum_{j=0}^{2s_0} \binom{2s_0}{2j} k^{2j} (2\sqrt{k-1})^{2s_0-2j} \\ &> \frac{n}{2(s_0+1)^2} (k + 2\sqrt{k-1})^{2s_0}. \end{aligned}$$

From the previous inequalities, it follows that if

$$c(\epsilon, k) = \frac{(k + (2 - \epsilon)\sqrt{k-1})^{2s_0}}{(2k)^{2s_0} - (k + (2 - \epsilon)\sqrt{k-1})^{2s_0}}$$

then $c(\epsilon, k) > 0$ and $m > c(\epsilon, k)n$.

□

The next result is an immediate consequence of Theorem 4.

Corollary 5 *Let $(X_i)_{i \geq 0}$ be a sequence of k -regular graphs such that*

$\lim_{i \rightarrow \infty} \operatorname{oddg}(X_i) = \infty$. Then for each $l \geq 1$

$$\limsup_{i \rightarrow \infty} \mu_l(X_i) \leq -2\sqrt{k-1}$$

When $l = 1$, we get the main result from [8]. Also, Corollary 5 holds when $l = 1$ and $\lim_{i \rightarrow \infty} \operatorname{girth}(X_i) = \infty$. This special case of Corollary 5 was proved directly in [9] using orthogonal polynomials and is also a consequence of the eigenvalue distribution theorem from [11].

A theorem stronger than Corollary 5 has been proved by Serre in [8] using the eigenvalue distribution results from [16]. We now present an elementary proof of this theorem. For $r \geq 0$, let $c_r(X)$ be the number of cycles of length r in a graph X .

Theorem 6 Let $(X_i)_{i \geq 0}$ be a sequence of k -regular graphs such that $\lim_{i \rightarrow \infty} |X_i| = \infty$. If $\lim_{i \rightarrow \infty} \frac{c_{2r+1}(X_i)}{|X_i|} = 0$ for each $r \geq 1$, then for each $l \geq 1$

$$\limsup_{i \rightarrow \infty} \mu_l(X_i) \leq -2\sqrt{k-1}$$

Proof Let $l \geq 1$. For a graph X and $r \geq 1$, let $n_{2r+1}(X)$ denote the number of vertices v_0 in the graph X such that the subgraph of X induced by the vertices at distance at most r from v_0 is bipartite. Thus, $|X| - n_{2r+1}(X)$ is the number of vertices u_0 of X such that the subgraph of X induced by the vertices at distance at most r from u_0 contains at least one odd cycle. Since each such vertex is no further than r from each of the vertices of an odd cycle of length at most $2r + 1$, it follows that

$$|X| - n_{2r+1}(X) \leq \sum_{l=1}^{r-1} \alpha_{l,r} c_{2l+1}(X)$$

where $0 \leq \alpha_{l,r} \leq 3(2l+1)(k-1)^r$. Thus, we have the following inequalities

$$1 - \sum_{l=1}^{r-1} \alpha_{l,r} \frac{c_{2l+1}(X_i)}{|X_i|} \leq \frac{n_{2r+1}(X_i)}{|X_i|} \leq 1$$

for all $r \geq 1, i \geq 0$. Hence, for each $r \geq 1$

$$\lim_{i \rightarrow \infty} \frac{n_{2r+1}(X_i)}{|X_i|} = 1 \tag{2}$$

For $i \geq 0$, let $A_i = A(X_i)$. Then, for $i \geq 0$ and $r \geq 1$, we have

$$\text{Tr}(A_i^{2r+1}) = n_{2r+1}(X_i) \cdot 0 + (|X_i| - n_{2r+1}(X_i))\theta_{2r+1}(X_i) \tag{3}$$

where $0 \leq \theta_{2r+1}(X_i) \leq k^{2r+1}$. From (2) and (3), we obtain that for each $r \geq 1$

$$\lim_{i \rightarrow \infty} \frac{\text{Tr}(A_i^{2r+1})}{|X_i|} = 0 \tag{4}$$

By using relation (1), it follows that for each $r \geq 1$

$$\liminf_{i \rightarrow \infty} \frac{\text{Tr}(A_i^{2r})}{|X_i|} \geq \frac{(2\sqrt{k-1})^{2r}}{(r+1)^2} \quad (5)$$

Now for each $i \geq 0$, we have

$$\text{Tr}(kI - A_i)^{2s} = \sum_{j=1}^{|X_i|} (k - \lambda_j(X_i))^{2s} \leq (|X_i| - l)(k - \mu_l(X_i))^{2s} + l(2k)^{2s}$$

Once again, the binomial expansion gives us

$$\text{Tr}(kI - A_i)^{2s} = \sum_{j=0}^{2s} \binom{2s}{j} k^j (-1)^{2s-j} \text{Tr}(A_i^{2s-j})$$

From the previous two relations, we get that

$$(k - \mu_l(X_i))^{2s} + \frac{4^s l k^{2s}}{|X_i| - l} \geq \sum_{j=0}^{2s} \binom{2s}{j} k^j (-1)^{2s-j} \frac{\text{Tr}(A_i^{2s-j})}{|X_i| - l}$$

Using relations (4) and (5), it follows that

$$\begin{aligned} k - \limsup_{i \rightarrow \infty} \mu_l(X_i) &\geq \left(\sum_{j=0}^s \binom{2s}{2j} k^{2j} \frac{(2\sqrt{k-1})^{2s-2j}}{(s-j+1)^2} \right)^{\frac{1}{2s}} \\ &> \left(\frac{1}{(s+1)^2} \sum_{j=0}^s \binom{2s}{2j} k^{2j} (2\sqrt{k-1})^{2s-2j} \right)^{\frac{1}{2s}} \\ &> \left(\frac{1}{2(s+1)^2} \right)^{\frac{1}{2s}} (k + 2\sqrt{k-1}) \end{aligned}$$

for any $s \geq 1$. By taking the limit as $s \rightarrow \infty$, we get

$$k - \limsup_{i \rightarrow \infty} \mu_l(X_i) \geq k + 2\sqrt{k-1}$$

which implies the inequality stated in the theorem.

□

Acknowledgments

This paper is part of my Ph.D. thesis at Queen's University. I am grateful to my thesis advisors: David Gregory, Ram Murty and David Wehlau for their help and support and to Chris Godsil and Shlomo Hoory for their comments. I thank the referees for many useful suggestions.

References

- [1] N. Alon, Eigenvalues and Expanders, *Combinatorica*, **6** (1986), 83-96.
- [2] S. M. Cioabă, Eigenvalues, Expanders and Gaps between Primes, Ph.D. Thesis, Queen's University at Kingston, submitted (2005).
- [3] S. M. Cioabă and R. Murty, Expander Graphs and Gaps between Primes, in progress.
- [4] G. Davidoff, P. Sarnak and A. Vallete, *Elementary Number Theory, Group Theory and Ramanujan Graphs*, Cambridge University Press, (2003).
- [5] K. Feng and W.-C. Winnie Li, Spectra of hypergraphs and applications, *J. of Number Theory*, **60** (1996), no.1, 1-22.
- [6] J. Friedman, Some geometric aspects of graphs and their eigenfunctions, *Duke Math. J.*, **69** (1993), 487-525.
- [7] W.-C. Winnie Li, Number Theory with Applications, Series of University Mathematics, Vol.7, World Scientific, (1996).
- [8] W.-C. Winnie Li (with an appendix by J.-P. Serre), On negative eigenvalues of regular graphs, *Comptes Rendus de l'Académie des Sciences*, **333** (2001), issue 10, 907-912.

- [9] W.-C. Winnie Li and P. Solé, Spectra of regular graphs and hypergraphs and orthogonal polynomials, *Europ.J.Combin.*, **17** (1996), 461-477.
- [10] A. Lubotzky, R. Phillips and P. Sarnak, Ramanujan Graphs, *Combinatorica*, **8** (1988), no.3, 261-277.
- [11] B. McKay, The expected eigenvalue distribution of a large regular graph, *Linear Algebra and its Applications*, **40** (1981), 203-216.
- [12] A. Nilli, On the second eigenvalue of a graph, *Discrete Mathematics*, **91** (1991), 207-210.
- [13] A. Nilli, Tight estimates for eigenvalues of regular graphs, *Electronic Journal of Combinatorics*, **11** (2004), N9.
- [14] A. Pizer, Ramanujan Graphs, *Computational perspectives on number theory* (Chicago, IL, 1995), 159-178, AMS/IP Stud. Adv. Math., 7, Amer. Math. Soc., Providence, RI, (1998).
- [15] J.-P. Serre, Private letters to W. Li dated October 8, 1990 and November 5, 1990.
- [16] J.-P. Serre, Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p , *J.Amer.Math.Soc.*, **10** (1997), no.1, 75-102.