

# Betti strata of height two ideals

Anthony Iarrobino

*Department of Mathematics, Northeastern University, Boston, MA 02115, USA.*

June 29, 2004

## Abstract

Let  $R = k[x, y]$  denote the polynomial ring in two variables over a field  $k$ . We study the Betti strata of the family  $G(H)$  parametrizing graded Artinian quotients of  $R = k[x, y]$  having given Hilbert function  $H$ . The Betti stratum  $G_\beta(H)$  parametrizes all quotients  $A$  of having the graded Betti numbers determined by  $H$  and the relation degrees  $\beta$ . We recover that the Betti strata are irreducible, and we calculate their codimension in the family  $G(H)$ .

**Theorem.** *The codimension of  $G_\beta(H)$  in  $G(H)$  satisfies, letting  $\nu_i = \#\{\text{generators of degree } i\}$ , and  $\beta_i = \#\{\text{relations of degree } i\}$ ,*

$$\text{cod } G_\beta(H) = \sum_{i > \mu} (\beta_i)(\nu_i).$$

Here  $\mu = \min\{i \mid H_i < i+1\}$ , the initial degree of the ideals. When  $k$  is algebraically closed, we also show that the closure of a Betti stratum is the union of more special strata, and is Cohen-Macaulay (Theorem 2.17). Our method is to identify  $G_\beta(H)$  as a product of determinantal varieties. Ours is a more direct argument than that of [IK, Theorem 5.63], which obtains the codimension of Betti substrata of a postulation stratum for punctual subschemes of  $\mathbb{P}^2$ ; there we relied on a result of M. Boij determining the codimension of Betti strata of height three Gorenstein algebras [Bj]. Key tools throughout include properties of an invariant  $\tau(V)$ , the number of generators of the ancestor ideal  $\overline{V} \subset R$ , and previous results concerning the projective variety  $G(H)$  in [I1]. We also adapt a method of M. Boij that reduces the calculation of the codimension of the Betti strata to showing that the most special stratum has the right codimension.

## 1 Introduction

Let  $R = k[x, y]$  be the polynomial ring in two variables over a field  $k$ ; for our main results we assume that  $\text{char } k = 0$  or  $\text{char } k$  is large enough – greater than  $j = \max\{i \mid H_i \neq 0\}$ . We study the Betti strata  $G_\beta(H)$  of the family  $G(H)$  of standard graded Artinian quotients  $A = R/I$  of  $R$  having given Hilbert function  $H$ ; here  $\beta$  is a sequence specifying the number of relations in each degree for the ideal  $I$ .

For arbitrary graded ideals in  $R$ , the Hilbert function does not determine the graded Betti numbers, in contrast to the situation for the three special algebras determined by a vector space  $V$  of degree- $j$  forms in  $R$  studied in [I3] — the algebra  $R/(V)$ , the level algebra  $LA(V)$  and the ancestor algebra  $\text{Anc}(V)$ . Which graded Betti numbers are possible, given the Hilbert function is well known (see [Cam], [Fr, §5], and [I1, Lemma 4.5]). By the Hilbert-Burch theorem the Betti stratum  $G_\beta(H)$  is determined by the Hilbert function  $H = H(R/I)$ , together with the sequence  $\beta = \beta(I)$  specifying the number of relations in each degree. A key invariant we used in studying

the special algebras, and that we use again here is  $\tau(V)$ , a measure of the complexity of  $V$ : it is the number of generators of the ancestor ideal  $\bar{V}$  (see [I3] and Lemma 2.2 below).

We will show that, given  $H$ , to specify  $\beta(I)$  is the same as to specify the invariant  $\tau(I)$ , which fixes  $\tau_i = \tau(I_i)$  for each  $i$ , and we give sharp bounds on these sequences of integers (Proposition 2.7). We then show that each Betti stratum  $G_\beta(H)$  is irreducible, and we determine the codimension of  $G_\beta(H)$  in the projective variety  $G(H)$  parametrizing all graded ideals of  $R$  such that  $H(R/I) = H$ . Letting  $\beta_i, \nu_i$  denote the number of degree- $i$  relations and generators, respectively of  $I$ , and  $\mu = \mu(H) = \min\{i \mid H_i < i + 1\}$  the *order* of  $H$  (initial degree of the ideals  $I$ ), we show (Theorem 2.17(i), equation (2.38)) that this codimension has the strikingly simple form

$$\text{cod } G_\beta(H) = \sum_{i > \mu} (\beta_i)(\nu_i).$$

We also show that the Zariski closure of a Betti stratum satisfies the *frontier property*,

$$\overline{G_\beta(H)} = \bigcup_{\beta' \geq \beta} G_{\beta'}(H),$$

where  $\beta' \geq \beta$  if for each  $i > \mu(H)$ ,  $\beta'_i \geq \beta_i$ ; also, the closure is Cohen-Macaulay with singularity locus  $\bigcup_{\beta' > \beta} G_{\beta'}(H)$ . (Theorem 2.17(ii), (iii)).

Our method is to identify (an open subset of) the  $G_\beta(H)$  stratum as the product of determinantal varieties, where each determinantal variety is the rank  $\tau_i$  locus of a certain homomorphism  $\theta_i$  (see (2.32)). Here, we mean by determinantal variety a variety defined by suitable minors of a matrix, and each component of which has the right codimension.

As a consequence of our main result, we determine in Section 3 the Hilbert functions possible for algebras of given socle type when  $r = 2$ , a result announced in [I2, Theorem 4.6C], whose complete proof was referenced to the paper [I3], but is instead shown here (Theorem 3.2). We then determine the Hilbert function of the intersection of a set of general enough level ideals  $I(i) \subset R$  each having given invariants ( $\tau$ , order, type) and each having a given Hilbert function  $H(i) = H(R/I(i))$  (Theorem 3.6).

That we can show a codimension formula and frontier property for Betti strata of height two Cohen-Macaulay algebras, is not surprising nowadays, as similar results have been found for the Betti strata of height three graded Gorenstein ideals [Bj, Di]. Also, V. Kanev and the author using the results of S. Diesel and M. Boij derived analogous codimension formulas and showed a frontier property for the Betti substrata of the schemes  $\text{Hilb}^{\Sigma(H)}(\mathbb{P}^2)$ , which are the postulation Hilbert schemes parametrizing those punctual schemes in  $\mathbb{P}^2$  with Hilbert function  $\Sigma(H)$  [IK, Theorem 5.63]. The postulation Hilbert function  $\Sigma(H)$  there is the sum function of  $H$  here.

There, we used as input M. Boij's formula for the codimension of Betti strata for height three Gorenstein ideals of a given Hilbert function  $T$  containing a subsequence  $(t, t, t)$ , and also S. J. Diesel's proof of a frontier property for those strata [Bj, Di]. We also used the connections between certain families of codimension three Gorenstein Artinian algebras and punctual schemes in  $\mathbb{P}^2$ , studied extensively in [IK, Part II]. The codimension result here Theorem 2.17(i) could be derived from [IK, Theorem 5.63]. The argument here is rather more direct. Conversely, Theorem 2.17 below and the argument of V. Kanev and the author in [IK, §5.5] could be used to recover the codimension formula of M. Boij for the Betti strata of height three, socle-degree  $j$  Gorenstein ideals of Hilbert function  $T$ , with  $(\Delta T)_{\leq j/2} = H$ , [Bj]; this recovers the M. Boij result only for the special case that  $T$  contains a subsequence  $(t, t, t)$ , where  $t$  is the length  $|H| = \sum H_i$ .

The irreducibility of the Betti strata here is a natural consequence of the Hilbert-Burch theorem giving a determinantal minimal resolution for height two Cohen-Macaulay ideals. G. Ellingsrud used this approach [Ell, p.427] to show the irreducibility of the union of the Betti strata with  $\beta$  smaller or equal a given  $\beta_0$  (so more general than a given stratum), as part of his proof of the smoothness of the Hilbert scheme of codimension two ACM ideals. The frontier property here

would also admit of a proof analagous to that of S. J. Diesel for the Betti strata of Gorenstein height three ideals, by deforming the relation matrix [Di].

However, we have chosen to base our proof of Theorem 2.17 on the relation matrix for the standard generators found in [I1] for height two graded Artinian quotients of  $R$ . More precisely, we base it on a the study of a certain homomorphism  $\theta_i$ , which for each degree  $i$ ,  $\mu \leq i < j$  has rank  $\tau_i = \tau(I_i)$ , the dimension of the vector space quotient  $R_1 I_i / y I_i$ . We feel that the interpretation of the Betti strata as the product of the rank  $\tau_i$  strata of these homomorphisms illuminates the codimension formulas (2.37), (2.38), and also allows us to apply the theory of determinantal ideals. A key step in the proof uses a method of M. Boij, that determines the codimension of each Betti stratum — a putative product of determinantal varieties — by verifying the codimension of the most special stratum [Bj, Proposition 3.2]. Here the most special stratum satisfies,  $i \geq \mu(H)$  implies  $\tau(I_i) = 1$ : this is equivalent to  $I_i$  being the degree- $i$  homogeneous component of a principal ideal of degree  $H_i$  (that depends on  $i$ ), and is the case  $\beta = \beta_{\max}(H)$  of (2.19) and Corollary 2.9.

In higher dimensions, several authors have studied the postulation of height two irreducible arithmetic Cohen-Macaulay curves in  $\mathbb{P}^3$ , in particular showing that they have “decreasing type” [GruP, MaR, HTV, GM]. In Remark 3.7 we discuss the consequences of our main result for Betti strata of height two ACM schemes of given  $h$ -vector  $H$  and higher embedding dimension than two.

## 2 Betti strata for graded ideals of $R = k[x, y]$

We assume throughout that  $R = k[x, y]$  is the polynomial ring in two variables over an arbitrary field  $k$ , unless  $k$  is otherwise specified. We study standard graded Artinian quotients  $A = R/I$  having given Hilbert function  $H$ . In Section 2.1 we show that, given  $H$ , a Betti stratum is equivalent to fixing  $\tau(I_i)$  for each  $i$  (Proposition 2.7 and Corollary 2.8). In Section 2.2 we study the codimension and Zariski closure of the Betti stratum  $G_\beta(H)$ . Our main result is Theorem 2.17. We now state several preparatory results. First, we define the invariant  $\tau(V)$ .

**Definition 2.1.** For  $a > 0$  and  $V \subset R_j$  we let

$$R_a V = \langle f v \mid f \in R_a, v \in V \rangle \subset R_{a+j} \quad (2.1)$$

$$R_{-a} V = \{f \in R_{j-a} \mid R_a f \subset V\}. \quad (2.2)$$

We define  $\tau(V)$  by

$$\tau(V) = \dim R_1 V - \dim V, \quad (2.3)$$

and the *ancestor ideal*

$$\overline{V} = R_{-j} V + \cdots + R_{-1} V + (V) \quad (2.4)$$

**Lemma 2.2.** [I3] *We have*

$$\tau(V) = \dim V - \dim R_{-1} V \text{ and} \quad (2.5)$$

$$\tau(V) = \#\{\text{generators of } \overline{V}\}. \quad (2.6)$$

*Remark on proof.* The equation 2.5 results from the exact sequence

$$0 \rightarrow R_{-1} V \xrightarrow{\phi} R_1 \otimes V \xrightarrow{\theta} R_1 \cdot V \rightarrow 0, \quad (2.7)$$

where  $\phi : f \rightarrow y \otimes x f - x \otimes y f$ , and  $\theta : \sum_i \ell_i \otimes v_i \rightarrow \sum_i \ell_i v_i$ , where the  $\ell_i$  are elements of  $R_1$  (linear forms). The equation 2.6 results from a repeated application of (2.7) (see [I3, Lemma 2.2]).  $\square$

We recall the characterization of  $O$ -sequences  $H$  — the sequences possible for Hilbert functions  $H(R/I)$  in two variables. We let  $\mu = \min\{i \mid H_i \neq i + 1\}$ .

**Lemma 2.3.** *The sequence  $H$  is an  $O$ -sequence for some quotient of  $R = k[x, y]$  if and only if*

$$H = (1, 2, \dots, \mu, H_\mu, H_{\mu+1}, \dots, H_i, \dots) \text{ where} \\ \mu \geq H_\mu \geq H_{\mu+1} \geq \dots \geq c_H \text{ and } \lim_{i \rightarrow \infty} H_i = c_H \geq 0. \quad (2.8)$$

Recall that the variety  $G(H)$  parametrizes graded ideals in  $R$  of Hilbert function  $H(R/I) = H$ ; it is defined as the closed subvariety

$$G(H) \subset \prod_{\mu \leq i \leq s} \text{Grass}(i+1-H_i, R_i) \quad (2.9)$$

of the product of Grassmanians parametrizing dimension  $(i+1-H_i)$  subspaces  $V_i \subset R_i$ : here  $G(H)$  parametrizes the ideals, satisfying  $R_1 V_i \subset V_{i+1}$ . That this is a closed condition is evident, since it is the intersection of the closed conditions  $xV_i \subset V_{i+1}$  and  $yV_i \subset V_{i+1}$ .

When  $c = c_H > 0$ , we denote by  $H : c$  the sequence  $H : c_i = H_{i+c} - c$ . Then we have the well-known

**Lemma 2.4.** *(See [I3, Corollary 2.14]) Let  $I \subset R = k[x, y]$  be an ideal satisfying  $H(R/I) = H, c = c_H > 0$ . Then  $\exists f \in R_c$  such that  $I = f \cdot I'$  where  $R/I'$  is an Artinian quotient of  $R$  having Hilbert function  $H : c$ . Also,  $G(H) \cong G(H : c) \times \mathbb{P}^c$ .*

Evidently, the generators and relations for  $I$  are simply related to those for  $I'$ , and the Betti stratum  $G_\beta(H)$  of  $G(H)$  is the product of a corresponding Betti stratum for  $G(H')$  by  $\mathbb{P}^c$ . Hence it will be sufficient to treat the case  $c_H = 0$  below.

We will use the following result, quoted from [I3, Theorem 1.10], valid over a field  $k$  of arbitrary characteristic. The proof is essentially from [I1], but the necessary adjustments for an arbitrary field  $k$  are outlined in [I3, Theorem 1.10]. For an integer  $n \in \mathbb{Z}$  we set  $n^+ = \max\{0, n\}$ .

**Theorem 2.5.** *[I1, Theorems 2.9, 2.12, 3.13, 4.3, Proposition 4.4, Equation 4.7], [I3, Theorem 1.10] Let  $r = 2$ , and for (i) let the field  $k$  be algebraically closed. Let  $H$  be an  $O$ -sequence that is eventually constant, so  $H$  is a sequence satisfying (2.8), let  $c = c_H$  and let  $H_s = c_H, H_{s-1} \neq c_H$ .*

- i. *Then  $G(H)$  is a smooth projective variety of dimension  $c + \sum_{i \geq \mu} (e_i + 1)(e_{i+1})$ .  $G(H)$  has a finite cover by opens in an affine space of this dimension. If  $\text{char } k = 0$  or  $\text{char } k > s$  then  $G(H)$  has a finite cover by opens that are affine spaces.*
- ii. *[I1, Theorem 4.3] The number of generators  $\nu(I)$  of a graded ideal  $I$  for which  $H(R/I) = H$ , satisfies  $\nu(I) \geq \nu(H) = 1 + e_\mu + \sum_{i \geq \mu} (e_{i+1} - e_i)^+$ .*
- iii. *[I1, Proposition 4.4] Assume that  $k$  is an infinite field. The family  $U(H)$  of graded ideals  $I$  satisfying  $H(R/I) = H$  and having the minimum possible number  $\nu(H)$  of generators given by equality in (ii) form an open subscheme of  $G(H)$  having the dimension specified in (i). Also  $U(H)$  is dense in  $G(H)$  when  $k$  is algebraically closed.*

## 2.1 The $\tau$ invariant and the Betti strata

The equations (2.13) and (2.14) of Proposition 2.7 below bounding the generator and relation sequences of a height two CM ideal in terms of the Hilbert function are well known: for example, (2.13) follows from [I1, Lemma 4.5]. They are also a consequence of R. Fröberg's result that every set of Betti numbers for a CM height two ideal in  $R$  is attained by a monomial ideal [Fr, §5]. Indeed, the possible Betti sequences for height two CM ideals were determined by G. Campanella in [Cam].

**Definition 2.6.** Let  $H$  be an  $O$ -sequence — a sequence satisfying (2.8). We define  $E(H) = (e_\mu, \dots, e_i, \dots)$  as the first difference sequence  $e_i = H_{i-1} - H_i$ . Let  $s = s(H) = \min\{i \mid H_i = c_H\}$ . We denote by  $\beta(I)$  the sequence of integers

$$\beta(I) = (\beta_{\mu+1}(I), \dots, \beta_i(I), \dots, \beta_s(I)), \quad (2.10)$$

where  $\beta_i(I)$  is the minimal number of relations of  $I$  in degree  $i$  (so  $\beta_i$  is uniquely determined by  $I$ ). We denote by  $\nu(I) = (\nu_\mu, \dots, \nu_s)$  the corresponding sequence where  $\nu_i$  is the number of degree- $i$  generators of  $I$ , and by  $\tau(I)$  the sequence

$$\tau(I) = (\tau(I_\mu), \tau(I_{\mu+1}), \dots, \tau(I_s) = 1) = (\tau_\mu, \tau_{\mu+1}, \dots, \tau_s). \quad (2.11)$$

We will need the well known fact for the socle  $0 : M \subset A$  of a graded quotient  $A = R/I$ ,  $R = k[x, y]$

$$\dim \text{Soc}(A)_i = \beta(I)_{i+2}. \quad (2.12)$$

**Proposition 2.7.** Let  $H = H(R/I)$  where  $I$  is a graded ideal of  $R = k[x, y]$  be a sequence satisfying (2.8), and let  $\mu(H) = \min\{i \mid H_i < i + 1\}$ . Let  $i \geq \mu(H)$ . Then the number of generators  $\nu_{i+1}(I)$  of  $I$  having degree  $i + 1$  satisfies

$$e_{i+1} \geq \nu_{i+1}(I) \geq (e_{i+1} - e_i)^+; \quad (2.13)$$

and  $\nu_\mu(I) = e_\mu + 1$ . For  $i \geq \mu$  the number of degree  $i + 1$  relations  $\beta_{i+1}(I)$  satisfies

$$e_i \geq \beta_{i+1}(I) \geq (e_i - e_{i+1})^+, \quad (2.14)$$

and  $\beta_i = 0$  for  $i \leq \mu$ . The  $\tau$  invariant is zero for  $i < \mu$ ; for  $i \geq \mu$  it satisfies

$$\tau(I_i) = e_{i+1}(H) + 1 - \nu_{i+1}(I) = e_i(H) + 1 - \beta_{i+1}(I) \text{ and} \quad (2.15)$$

$$\tau(I_i) = \tau(I_{i-1}) + \nu_i(I) - \beta_{i+1}(I), \quad (2.16)$$

and also for  $i \geq \mu$  the inequality

$$1 \leq \tau(I_i) \leq 1 + \min\{e_i(H), e_{i+1}(H)\}. \quad (2.17)$$

For an open dense subset of  $G(H)$  the invariants  $\beta(I)$ ,  $\nu(I)$  and  $\tau(I)$ , respectively, satisfy equality on the right sides of (2.13), (2.14) and (2.17).

*Proof.* We have evidently  $\dim I_{i+1} = \dim R_1 I_i + \nu_{i+1}(I)$ , and also as a consequence of (2.12) that  $\dim R_{-1} I = \dim I_{i-1} + \beta_{i+1}(I)$ . Thus by the definition 2.3 of  $\tau$  and Lemma 2.2 we have for  $i \geq \mu$ ,

$$\tau(I_i) = \dim R_1 I_i - \dim I_i = \dim I_{i+1} - \nu_{i+1}(I) - \dim I_i = e_{i+1} + 1 - \nu_{i+1}(I) \text{ and}$$

$$\tau(I_i) = \dim I_i - \dim R_{-1} I_i = \dim I_i - (\dim I_{i-1} + \beta_{i+1}(I)) = e_i + 1 - \beta_{i+1}(I).$$

Here the second equation follows from the first and the usual calculation that the second difference

$$\Delta^2(H)_{i+1} = e_i - e_{i+1} = \beta_{i+1}(I) - \nu_{i+1}(I). \quad (2.18)$$

This shows (2.15) and implies all the equations (2.13) to (2.17) for  $i \geq \mu(H)$ . For example, (2.18) directly implies the right sides of (2.13) and (2.14), while  $\tau(I_i) \geq 1$  from Lemma 2.2 and (2.15) imply the left sides of (2.13) and (2.14); and (2.15) directly implies the right side of (2.17). By Theorem 2.5(iii), an open dense subset of ideals  $I \in G(H)$  consists of ideals having the minimum possible number of generators, given by  $\nu(H) = 1 + e_\mu + \sum_{i \geq \mu} (e_{i+1} - e_i)^+$ . This subset consists of ideals satisfying equality on the right of each of the inequalities (2.13), (2.14), (2.17) for  $\nu(I)$ ,  $\beta(I)$  and  $\tau(I)$ , respectively.  $\square$

**Corollary 2.8.** *Let  $I$  be an arbitrary graded ideal of  $R = k[x, y]$ . Then the pair  $(H(R/I), \tau(I))$  are equivalent to the pair  $(\nu(I), \beta(I))$  of sequences giving the graded Betti numbers of  $I$ . Given  $H = H(R/I)$  each of the sequences  $\nu(I), \beta(I), \tau(I)$  determines the other two.*

*Proof.* We have  $\tau_\mu(I) = e_\mu(H) + 1 = \nu_\mu(I)$ , and  $\beta_\mu = 0$ . Then (2.15) shows that the pair  $(H(R/I), \tau(I))$  determines both the generator and relation degrees, and (2.16) shows that the latter pair determine the former.  $\square$

We define  $\beta_{\max}(H)$  — the most special stratum — by equality in the upper bound (left) of (2.14)

$$(\beta_{\max})_i = \begin{cases} e_{i-1} & \text{if } \mu < i \leq s(H) \\ 0 & \text{otherwise.} \end{cases} \quad (2.19)$$

The corresponding  $\tau_{\max}$  satisfies  $\tau_i = 1$  for  $i \geq \mu$ , and  $\nu_{\max}$  is given by equality on the left of (2.13).

**Corollary 2.9.** *Let  $H$  be a sequence satisfying (2.8). The subscheme  $G_{\beta_{\max}}(H)$  of  $G(H)$  is isomorphic to  $(\prod_{i=\mu+1}^s \mathbb{P}^{e_i}) \cdot \mathbb{P}^{c_H}$ , and has dimension  $H_\mu$ .*

*Proof.* When  $\beta = \beta_{\max}$ ,  $\tau(I) = \tau_{\min} = (1, 1, \dots, 1)$ . By Lemma 2.2 equation (2.6) we have  $\bar{I}_i = (f_i)$  of degree  $H_i$ , and it is easy to see that we have  $f_s \mid f_{s-1} \dots \mid f_\mu$ , whence each such ideal  $I$  determines a point  $(f_\mu/f_{\mu-1}, \dots, f_i/f_{i+1}, \dots, f_{s-1}/f_s; f_s)$  of  $(\prod_{i=\mu+1}^s \mathbb{P}^{e_i}) \cdot \mathbb{P}^{c_H}$ ; and conversely.  $\square$

**Definition 2.10.** Let  $H$  be fixed satisfying (2.8), and  $\beta$  be a sequence satisfying (2.10). We denote by  $G_\beta(H)$  the reduced subscheme of  $G(H)$  parametrizing all ideals  $I \in R$  such that  $H(R/I) = H$  and  $\beta(I) = \beta$ .

We will say for two sequences  $\beta', \beta$  satisfying (2.14) that

$$\beta' \geq \beta \text{ if and only if } \beta'_i \geq \beta_i \ \forall i. \quad (2.20)$$

Evidently,  $G_\beta(H)$  is a locally closed subscheme of  $G(H)$ , as fixing each  $\tau_i$  is to fix the vector space dimension of  $R_1 I_i$ , or, equivalently, the rank of the image of  $R_1 \otimes I_i \rightarrow I_{i+1}$  under the multiplication map.

## 2.2 Codimension and closure of the Betti strata

We now state and prove our main result about the Betti strata  $G_\beta(H)$  of  $G(H)$ , the projective variety parametrizing graded quotients of  $R = k[x, y]$  having Hilbert function  $H$  (Theorem 2.17). We need an adaptation of a result of M. Boij [Bj]. Recall from Proposition 2.7 that for  $i > \mu$  we have  $(\beta_{\max})_{i+1} = e_i$ . Recall that for any ideal  $I \in G(H)$  we have a resolution

$$0 \rightarrow \oplus_{i \geq \mu} R(-i-1)^{e_i} \rightarrow R_\mu \oplus_{i \geq \mu(H)} R(-i)^{e_i} \rightarrow R \rightarrow R/I \rightarrow 0, \quad (2.21)$$

that is minimal only when  $\beta(I) = \beta_{\max}$ . Denote by  $X(H)$  the open subfamily of all maps

$$\oplus_{i \geq \mu(H)} R(-i)^{(\beta_{\max})_i} \rightarrow R_\mu \oplus_{i \geq \mu(H)} R^{e_i} \quad (2.22)$$

that have a rank one cokernel; restricting to an affine subset, we have a morphism  $\pi : X(H) \rightarrow G(H)$ ; we denote by  $X_\beta(H)$  the inverse image of  $G_\beta(H)$ . Then, following M. Boij, we have a commutative fiber diagram

$$\begin{array}{ccc} X_\beta(H) & \longrightarrow & X(H) \\ \downarrow & & \downarrow \\ \overline{G_\beta(H)} & \longrightarrow & G(H) \end{array} \quad (2.23)$$

where the vertical morphisms are dominant and the horizontal morphisms are injective. We can replace all by the associated reduced schemes.

**Lemma 2.11.** (after [Bj, Proposition 3.2]). For  $\beta$  satisfying (2.14) we have that the codimensions of  $G_\beta(H)$  in  $G(H)$  and  $X_\beta(H)$  in  $X(H)$  satisfy

$$\text{cod } G_{\beta_{\max}}(H) - \text{cod } X_{\beta_{\max}}(H) \geq \text{cod } G_\beta(H) - \text{cod } X_\beta(H) \geq 0. \quad (2.24)$$

*Proof.* Since  $\pi : X_\beta(H) \rightarrow \overline{G_\beta(H)}$  is a dominant morphism of integral schemes the dimension of its general fibre is no greater than that of its special fibres; the same is true for  $\pi : X(H) \rightarrow G(H)$ , hence we have

$$\dim X(H) - \dim G(H) \leq \dim X_\beta(H) - \dim G_\beta(H) \leq \dim X_{\beta_{\max}}(H) - \dim G_{\beta_{\max}}(H), \quad (2.25)$$

which is (2.24).  $\square$

We now define an initial monomial ideal  $M(H)$  and an open subset  $U_B$  of  $G(H)$ , the ideals having *normal pattern* or cobasis (see [I1]). Recall that  $s(H) = \min\{i \mid H_i = c_H\}$ . Recall that given a partition  $P$ , the dual partition  $P^*$  is that obtained by exchanging the rows and columns of the Ferrer's graph of  $P$ . We use this to define the “alignment character” of  $H$  below.

**Definition 2.12 (Normal pattern).** Given  $H$  satisfying (2.8) we define a *normal pattern* or *standard cobasis*:

$$B = B_H = \bigoplus_{u=0}^s B_u, \quad B_u = \langle y^{H_u-1} x^{u+1-H_u}, \dots, x^u \rangle. \quad (2.26)$$

We say that the ideal  $I \in G(H)$  has *normal pattern* in the coordinates  $x, y$  for  $R$  and write  $I \in U_B, B = B_H$  if  $I$  satisfies

$$I_u \oplus B_u = R_u \quad \forall u \mid \mu \leq u \leq s(H). \quad (2.27)$$

We denote by  $M(H)$  the unique monomial ideal  $M(H)$  having normal pattern in the basis  $H$ . It satisfies  $M(H) = \bigoplus M(H)_u$ , where

$$M(H)_u = \langle y^u, y^{u-1}x, \dots, y^{H_u}x^{u-H_u} \rangle \quad (2.28)$$

*Preparation for standard generators.* We will need the standard generators for a graded ideal  $I \subset R$  having *normal pattern* or cobasis  $B$ , and a variation of the coordinates for  $U_B$  used in the proofs [I1, Theorem 4.3 and Proposition 4.4]. In [I1, Proposition 3.2] it is shown that given  $\text{char } k = 0$  or  $\text{char } k > s(H)$  then  $G(H)$  has a finite cover by affine open subvarieties, each isomorphic to  $U_B, B = B_H$ . We will now assume that  $I \in U_B$ . Our standard generating set  $F = F(I)$  for  $I$  has  $e_i$  elements of degree  $i$  for  $i > \mu$ , and  $1 + e_\mu$  elements of degree  $\mu$ , and it is not usually a minimal generating set. To present  $F$ , following [I1, Proposition 2.5, Lemmas 2.7, 2.8], we first define a sequence of integers  $K = (k_0 > \dots > k_\mu)$  by adding  $(\mu - 1, \dots, 1)$  to the dual partition  $(h_1^* \geq \dots)$  to  $(H_{\mu-1} = \mu, H_\mu, \dots, H_{s-1})$ : thus,

$$k_0 = h_1^* + \mu - 1, \dots, k_i = h_{i+1}^* + \mu - (i + 1), \dots \quad (2.29)$$

The sequence  $K$  specifies the lengths of the horizontal rows of the bar graph of  $H$  (see Figure 1 and Example 2.19), and has sometimes been called the *alignment character* of  $H$ : see [GPS, Hari], and also [IK, p. 122-124] for a discussion and further references. For  $0 \leq i \leq \mu$ , the monomial  $v_i = y^i x^{k_i}$  is the unique monomial of the ideal  $M_H$  having  $y$ -degree  $i$  and lowest total degree.

**Definition 2.13 (Standard generators).** For each  $I \in U_B$  we let  $F = F(I) = (f_\mu, \dots, f_0)$  be the unique homogeneous elements of  $I$  such that

$$f_i = y^i x^{k_i} + g_i \text{ where } g_i \in B_{i+k_i}. \quad (2.30)$$

We define  $a_{i,u} \in k$  for  $0 \leq u \leq i \leq \mu$  by writing

$$F : f_i = y^i x^{k_i} + \sum_{\deg F_u > \deg F_i} a_{i,u} (f_u : x^{k_u - k_i - (i-u)}). \quad (2.31)$$

Note that  $F_v = (f_{H_{v-1}-1}, \dots, f_{H_v})$ , for  $s(H) > v > \mu$ , and  $F_\mu = (f_\mu, \dots, f_{H_\mu})$ . comprise the elements  $F_v = F \cap R_v$  of  $F$  having degree  $v$  for each  $v \geq \mu$ . We have  $\#\{F_v\} = e_v$  when  $v > \mu$  but  $\#\{F_\mu\} = e_\mu + 1$ , and  $\#\{F\} = \mu + 1$ .

We define for  $v \geq \mu$  the set  $A_v$  of coefficients, in two equivalent ways

- i.  $A_v = \{a_{i,u} \text{ such that } H_v \geq i \geq H_{v+1} \text{ and } H_{v+1} - 1 \geq u \geq H_{v+2}\};$
  - ii.  $A_v = \{a_{i,u} \text{ such that } f_i \in F_v, \text{ or is the next generator } f_{H_v} \text{ of a lower degree, and } f_u \in F_{v+1}\}.$
- We let  $A = \bigcup_{\mu \leq v} A_v$ .

These may also be described as the coefficients of  $f_u$  that land outside the pattern  $B_H$  in  $x^{w_u} f_u$ ,  $w_u = k_{u-1} - k_u$  (see [I1] and also [IY, §3B] for a discussion).

Evidently  $\#\{A_v\} = (e_v + 1)e_{v+1}$ . Our standard generators differ from those in [I1, Proposition 2.7] by using the forms  $f_u : x^a$  here in place of the monomials  $y^u x^{i-u}$  used there to write  $f_i$ . The chief advantage is to simplify the matrices  $N_i$  constructed below, but there is no change in determining which coefficients are parameters for  $U_B$ . According to [I1, Lemma 2.7, Theorem 2.9] these parameters are exactly the set  $A$  of coefficients defined above. Thus we have,

**Theorem 2.14.** [I1, Proposition 2.7, Theorem 2.9] *Let  $k$  be infinite. Fix an  $O$ -sequence  $H$  satisfying (2.8) and  $c_H = 0$ , and let  $B = B_H$ , the normal pattern of (2.26).*

- i. *Each ideal  $I \in U_B$  is generated by  $F(I)$ .*
- ii. *The subfamily  $U_B \subset G(H)$  is isomorphic to an affine space, with coordinates given by the coefficients  $A$ .*

*Proof.* That  $F(I)$  generates  $I$  is evident. Although the coefficients used for  $U_B$  differ slightly from those in [I1, Proposition 2.7, Theorem 2.9] by using the forms  $f_u : x^a$  here in place of the monomials  $y^u x^{i-u}$  used there to write  $f_i$  and hence define  $a_{i,u}$ ; there is no essential change in the result, and in particular, there is no change in the definition of the coefficients used as parameters (those in  $A$ ). Furthermore, each coefficient  $a_{i,u}$  that is not in  $A$  may be written as an explicit polynomial in those parameters in  $A$  corresponding to monomials of equal or higher degrees to  $\deg f_i$ . Evidently  $\#\{A\} = \sum_{i \geq \mu} (e_i + 1)e_{i+1}$ , which is the dimension of  $G(H)$  given in Theorem 2.5 above.  $\square$

We now prepare for our main result, Theorem 2.17, by defining key homomorphisms  $\theta_i$ . Using these we will show that the Betti strata  $G_\beta(H)$  are determinantal varieties having an expected dimension. We assume that  $H$  has been chosen satisfying (2.8), and that  $I \in U_B$ .

*The homomorphism  $\theta_i$  and the vector space  $\langle R_1 I_i / y I_i \rangle$ .* We have evidently from the Definition 2.3 of  $\tau$

$$\tau(I_i) = \dim R_1 I_i - \dim I_i = \dim \langle R_1 I_i / y I_i \rangle.$$

For now we fix  $i$  and let  $t = H_{i-1}$ ,  $t' = H_i$ . We denote by  $F'_i$  the  $k$ -vector space spanned by  $(x^{i-\deg f_t} f_t, F_i)$  and by  $F'_{i+1}$  the span of  $(x^{i+1-\deg f_{t'}} f_{t'}, F_{i+1})$ . Suppose  $I \in U_B$  and consider the homomorphism

$$\theta_i : F'_i \rightarrow F'_{i+1}, \quad (2.32)$$



where  $\theta_i$  is the composite  $\theta_i = \eta_i \circ \theta'_i \circ m_x$  of the homomorphisms

$$F'_i \xrightarrow{m_x} \langle R_1 I_i \rangle \xrightarrow{\theta'_i} \langle R_1 I_i / y I_i \rangle \xrightarrow{\eta_i} F'_{i+1}, \quad (2.33)$$

where  $m_x$  denotes multiplication by  $x$ , the homomorphism  $\theta'_i$  is simply reduction mod  $y I_i$ , and the inclusion  $\eta_i$  arises from the inverse of the isomorphism

$$F'_{i+1} \rightarrow \langle F'_{i+1} + y I_i \rangle / y I_i \rightarrow I_{i+1} / y I_i; \quad (2.34)$$

The composite map of (2.34) is an isomorphism, since  $I_{i+1} \cap B_{i+1} = 0$ , and the normal form of  $I$  assures that  $I_{i+1} = y I_i \oplus F'_{i+1}$ .

**Lemma 2.15.** *The homomorphism  $\theta_i$  of (2.32) is from a vector space of dimension  $e_i + 1$  to another of dimension  $e_{i+1} + 1$ ; and for  $I \in U_{B(H)}$ , we have*

$$\text{rank } \theta_i = \tau(I_i). \quad (2.35)$$

*Proof.* Since  $m_x$  and  $\eta_i$  are monomorphisms (2.35) follows.  $\square$

We denote by  $N_i$  the  $(e_{i+1} + 1) \times (e_i + 1)$  matrix for the homomorphism, where we list both domain and range in descending order of the index of the standard generators involved, and where we write the entries in terms of the parameters  $A$  (see Examples 2.19, 2.20). We denote by  $A'_{u,v}$  the subset of the parameters involving pairs  $f_{u'}, f_{v'}$  with  $u' < u$ , or  $u' = u$  and  $v' < v$ , that is, all those parameters “earlier than” the coefficient of  $f_u$  on  $f_v : x^{k_v - k_u + u - v}$ .

**Lemma 2.16.** *The entries of  $N_i$  are elements of the polynomial ring  $k[A]$ . The entry  $N_i(v, u)$ , corresponding to the component of  $\theta_i(f_u)$  involving  $f_v$ , satisfies*

$$N_i(v, u) = a_{u,v} \pmod{k[A'_{u,v}]}, \quad (2.36)$$

for all pairs  $u, v$  with  $f_v \in F'_{i+1}$  and  $v \neq t'$ . Also  $N_i$  has last  $f_{t'}$  column  $(1, 0, \dots)^T$ . After column reduction (change of basis in the domain of  $\theta_i$ ), the resulting matrix  $N'_i$  has first row  $(0, \dots, 0, 1)$ .

*Proof.* It is easy to see inductively, beginning with  $f_0$ , that for an ideal  $I \in U_B$ , the coefficients of the standard generators  $f_i$  are elements of the polynomial ring  $k[A]$ . From the simple definition of the homomorphism  $\theta_i$ , it follows that the entries of  $N_i$  are polynomials in the coefficients of the  $f_i$  involved, hence are also elements of the polynomial ring  $k[A]$ . Equation (2.36) is straightforward, and follows from a closer examination of the coefficients of the standard generators involved in calculating the matrix entry of  $N_i$  corresponding to the  $\theta_i(f_u)$  component on an  $f_v, v < u$ : this coefficient only involves either “earlier”  $f_{u'}, u' < u$ , or the same  $f_u$  but earlier  $f_{v'}, v' < v$ . However,  $a_{u,v}$  does occur, as one evaluates  $x^{k_v - k_u + u - v} f_u$  on  $f_v$ . We have  $\theta(f_{t'}) = x f_{t'}$ , hence the last column of  $N_i$  is  $(1, 0, \dots, 0)^T$ , as claimed.  $\square$

Recall that we denote by  $G_\beta(H)$  the Betti stratum of  $G(H)$  (see Definition 2.10).

**Theorem 2.17.** *Let  $H = (1, 2, \dots, 0)$  be an  $O$ -sequence for an Artinian quotient of  $R$ : so  $H$  satisfies (2.8) with  $c_H = 0$ . Let  $\beta$  denote a sequence of integers (potential relation degrees) satisfying the condition (2.14), and let  $\tau$  and  $\nu$  be the sequences defined from  $\beta$  by (2.15) and (2.18), respectively. Let  $k$  be an infinite field satisfying either  $\text{char } k = 0$  or  $\text{char } k = p \geq s(H)$ .*

- i. Then  $G_\beta(H)$  is irreducible. The intersection  $G_\beta(H) \cap U_B$  is open dense in  $G_\beta(H)$ , and is the product of the rank  $\tau_i$  loci of the homomorphisms  $\theta_i$  of (2.32), determined by  $(e_{i+1} + 1) \times (e_i + 1)$  matrices,  $\mu \leq i \leq s - 1$ . The codimension in  $G(H)$  of the Betti stratum  $G_\beta(H)$  satisfies*

$$\text{cod } G_\beta(H) = \sum_{i \geq \mu} (e_i + 1 - \tau_i)(e_{i+1} + 1 - \tau_i) \quad (2.37)$$

$$= \sum_{i > \mu} (\beta_i)(\nu_i). \quad (2.38)$$

ii. Assume that  $k$  is algebraically closed. The closure  $\overline{G_\beta(H)}$  satisfies the frontier property.

$$\overline{G_\beta(H)} = \bigcup_{\beta' \geq \beta} G_{\beta'}(H). \quad (2.39)$$

iii. Assume that  $k$  is algebraically closed. Then  $\overline{G_\beta(H)}$  is Cohen-Macaulay, and has singular locus  $\bigcup_{\beta' > \beta} G_{\beta'}(H)$ .

*Proof.* The irreducibility of each Betti stratum follows from the Hilbert-Burch theorem. The frontier property will follow from our characterization of the Betti strata, as an intersection of determinantal varieties, and the closure properties of determinantal varieties. To define the appropriate determinantal varieties we study the vector space  $\langle R_1 I_i / y I_i \rangle$ .

By Proposition 2.7 and Corollary 2.9 the intersection of the rank one loci of  $N_\mu, \dots, N_{s-1}$  is the  $\beta_{\max}$  stratum  $G_{\beta_{\max}}(H)$  of  $G(H)$ , and it has dimension  $\sum_{i > \mu} e_i = H_\mu$ . Since  $\dim G(H) = \sum_{i \geq \mu} (e_i + 1)(e_{i+1})$  by Theorem 2.5(i), the stratum  $G_{\beta_{\max}}(H)$  has the expected codimension in  $G(H)$ , namely

$$\text{cod } G_{\beta_{\max}}(H) = \sum_{i \geq \mu}^{s-1} (e_i)(e_{i+1}). \quad (2.40)$$

Thus, this locus is the intersection of determinantal varieties, and the difference on the left of the inequality (2.24) in M. Boij's Lemma 2.11. satisfies

$$\text{cod } G_{\beta_{\max}}(H) - \text{cod } X_{\beta_{\max}}(H) = 0.$$

This implies by (2.24) that for each  $\beta$  the difference

$$\text{cod } G_\beta(H) - \text{cod } X_\beta(H) = 0. \quad (2.41)$$

Recall that here  $\beta$  is defined by (2.15) where  $\tau$  satisfies (2.17), so  $G_\beta(H) \cap U_B$  is the subfamily parametrizing  $I \in U_B$  satisfying  $\tau(I) = \tau$ . But the codimension  $\text{cod } X_\beta$  is just the expected codimension  $\sum \beta_i \cdot \nu_i$ , as by definition the variety  $X_\beta$  is a product of generic determinantal varieties. It follows from (2.41) that each intermediate stratum  $G_\beta(H) \cap U_B$  has the expected codimension. It follows from (2.35) that each such locus  $G_\beta(H) \cap U_B$  is the product of the determinantal varieties given by the rank- $\tau_i$  locus of  $\overline{\theta_i}$ , as claimed. When  $k$  is algebraically closed, the frontier property and Cohen-Macaulayness of  $\overline{G_\beta(H)} \cap U_B$  and also the remainder of (iii) concerning the singularity locus follow from the analogous properties for a determinantal variety (see [BV], [W, §6.1], [ACGH, II.4.1]), since  $\overline{G_\beta(H)}$  is a product of determinantal varieties. This completes the proof of Theorem 2.17.  $\square$

The following corollary is the analogue of a result of S. Diesel for height three Gorenstein algebras (see [Di, §3] and also [IK, Theorem 5.25v.f]). We will denote by  $[n]$  where  $n$  is an integer, the simply ordered set  $\{0 \leq \dots \leq n\}$ . We denote by  $\mathcal{L}(H)$  the lattice

$$\prod_{\mu(H) \leq i < s(H)} [e_i - (e_i - e_{i+1})^+], \quad (2.42)$$

and we consider the map  $\eta$  on  $\beta$  sequences satisfying (2.14) to  $\mathcal{L}(H)$ ,

$$\eta(\beta) = (\beta_{\mu+1} - (e_\mu - e_{\mu+1})^+, \dots, \beta_{i+1} - (e_i - e_{i+1})^+, \dots, \beta_s - (e_{s-1} - e_s)^+). \quad (2.43)$$

**Corollary 2.18.** *Given  $H$  satisfying 2.8, the relation sequences  $\beta$  that occur, for which  $G_\beta(H)$  is nonempty are exactly those of (2.14). Under the termwise order of Definition 2.10, this set forms a lattice that under the map  $\eta$  is isomorphic to  $\mathcal{L}(H)$ . The Zariski closure  $\overline{G_\beta(H)} \supset G_{\beta'}(H)$  if and only if  $\beta' \geq \beta$  in the lattice, and  $\overline{G_\beta(H)} \cap G_{\beta'}(H) = \emptyset$  unless  $\beta' \geq \beta$ .*

					$y^3$
			$y^2$	$y^2x$	$y^2x^2$
		$y$	$yx$	$yx^2$	$yx^3$
1	$x$	$x^2$	$x^3$	$x^4$	$x^5$

Figure 1: Normal pattern  $B_H$ ,  $H = (1, 2, 3, 3, 1, 0)$ , (see Example 2.19).

*Proof.* This is immediate from Theorem 2.17 and Equation 2.14.  $\square$

**Example 2.19.** Let  $H = (1, 2, 3, 3, 1, 0)$ ; then we have  $\mu = 3$ , and the alignment character

$$K = (3, 3, 1)^* + (2, 1, 0) = (3, 2, 2) + (2, 1, 0) = (5, 3, 2),$$

the lengths of the rows of the “normal pattern  $B_H$  depicted in Figure 1. The standard generators  $F$  are (using  $a, b, c, d, e$  in place of  $a_{3,2}, \dots$ ),

$$f_0 = x^5, f_1 = yx^3 + ex^4, f_2 = y^2x^2 + dx^4, f_3 = y^3 + af_2 : x + bf_1 : x + cx^3.$$

All the coefficients form the parameters  $A$  for  $U_B$ . We have  $E(H) = (e_3, e_4, e_5) = (0, 2, 1)$ , and  $\dim G(H) = (1 \cdot 2 + 3 \cdot 1) = 5$ , as expected. The map  $\theta_4$  from  $\langle xf_3, f_2, f_1 \rangle$  to  $\langle xf_1, f_0 \rangle$ , has matrix

$$N_4 = \begin{pmatrix} -d - ae & -e & 1 \\ c + ed - ad + ae^2 & d - e^2 & 0 \end{pmatrix}.$$

After natural column operations as described in the proof of Theorem 2.17, the matrix becomes

$$N_4' = \begin{pmatrix} 0 & 0 & 1 \\ c + ed & d - e^2 & 0 \end{pmatrix}.$$

The map  $\theta_3$  has matrix  $N_3 = N_3' = (1, 0, 0)^T$ , of rank one. The rank one locus of  $\theta_4$  is given by  $d = -e^2, c = -ed = e^3$ , so has codimension two, and dimension three equal to  $H_\nu = H_3$  (see Corollary 2.9). Also, by equation 2.37 of Theorem 2.17 for  $\tau_4 = 1$ , this a codimension  $(3 - 1) \times (2 - 1) = \beta_4 \times \nu_4 = 2$  locus. The only other stratum is the rank two locus of  $\theta_4$ , which is the generic stratum of dimension five, open dense in  $G(H)$ .

**Example 2.20.** Let  $H = (1, 2, 3, 4, 2, 1, 0)$ . Then  $E(H) = (e_4, e_5, e_6) = (2, 1, 1)$ , and  $G(H)$  has dimension  $(3 \cdot 1 + 2 \cdot 1) = 5$ . The standard generators  $F$  are (using  $a, b, c, d, e$  in place of  $a_{4,2}, \dots$ ),

$$\begin{aligned} f_0 &= x^6, f_1 = yx^4 + ex^5, f_2 = y^2x^2 + cf_1 : x + dx^4, \\ f_3 &= y^3x + bf_1 : x + \alpha_1x^4, f_4 = y^4 + af_1 : x + \alpha_2x^4, \end{aligned}$$

where  $a, b, c, d, e$  are the parameters  $A$ , and  $\alpha_1, \alpha_2$  may be written as polynomials in the parameters. The non-trivial  $\theta$  maps are  $\theta_4$  and  $\theta_5$ . The matrix  $N_4'$  for the map  $\theta_4$  from  $\langle f_4, f_3, f_2 \rangle \rightarrow \langle xf_2, f_1 \rangle$  satisfies

$$N_4' = \begin{pmatrix} 0 & 0 & 1 \\ a & b - d & 0 \end{pmatrix} \pmod{A^2}.$$

The matrix  $N'_5$  for the map  $\theta_5$  from  $\langle xf_2, f_2 \rangle \rightarrow \langle xf_1, f_0 \rangle$  satisfies

$$N'_5 = \begin{pmatrix} 0 & 1 \\ d - ce + e^2 & 0 \end{pmatrix}.$$

The Betti strata of  $G(H)$  are determined by  $\tau(I_4) = \text{rank}(\theta_4)$ , and  $\tau(I_5)$  which can each be one or two. So there are four Betti strata of  $G(H)$ , of codimensions zero,  $(G_{\beta(0)}(H) : \tau_4 = \tau_5 = 2)$ , one  $(G_{\beta(1)}(H) : \tau_4 = 2, \tau_5 = 1)$ , two  $(G_{\beta(2)}(H) : \tau_4 = 1, \tau_5 = 2)$ , and three  $(G_{\beta_{\max}}(H) : \tau_4 = \tau_5 = 1)$ , respectively. The values of  $(\beta_5, \beta_6)$  that correspond are  $\beta(0) : (1, 0)$ ,  $\beta(1) : (1, 1)$ ,  $\beta(2) : (2, 0)$ ,  $\beta_{\max} : (2, 1)$ . By Corollary 2.18 the lattice  $\mathcal{L}(H) = [1] \times [1]$ , so is not simply ordered. In fact,  $\overline{G_{\beta(1)}(H)} = G_{\beta(1)}(H) \cup G_{\beta_{\max}}(H)$  and  $\overline{G_{\beta(2)}(H)} = G_{\beta(2)}(H) \cup G_{\beta_{\max}}(H)$ .

The following Corollary is known [Cam].

**Corollary 2.21.** *Let the sequence  $H$  satisfy (2.8), and  $\lim_{i \rightarrow \infty} H_i = 0$ , and suppose  $k$  is arbitrary. Each sequence of integers  $\beta$  satisfying (2.14) occurs as the relation degrees for an ideal of Hilbert function  $H$ . Likewise, each sequence  $\nu$ , satisfying (2.13), or sequence  $\tau$  satisfying (2.17) occurs.*

*Proof.* This is immediate from the codimension formula in Theorem 2.17. Since it is an existence result, the field  $k$  may be arbitrary.  $\square$

### 3 Algebra quotients of $k[x, y]$ having fixed socle type

In this section we characterize the Hilbert functions consistent with a given socle type (Theorem 3.2), and we also determine the Hilbert function of the intersection of a general enough set of  $t$  level ideals, each having a specified Hilbert function (Theorem 3.6).

**Definition 3.1.** The *socle type* ST of an Artinian algebra  $A = R/I$  is the Hilbert function of the socle  $\text{Soc}(A) = \langle 0 : M \rangle \subset A$ .

The following result was announced as [I2, Theorem 4.6C]. The case ST concentrated in a single degree  $j$  (so  $\text{ST}_i = 0$  unless  $i = j$ ), announced in [I2, Theorem 4.6A], was shown in [I3, Theorems 2.17, 2.19] for  $\text{LA}_N(d, j)$  (see Definition 3.4), where the resulting Hilbert function was related to a partition  $P$ .

**Theorem 3.2.** *Let the sequence  $H$  satisfy (2.8), and  $\lim_{i \rightarrow \infty} H_i = 0$ , and suppose  $k$  is arbitrary. The sequence  $H$  occurs for the Hilbert function of a graded algebra of socle type ST if and only if for all  $i$ ,*

$$e_{i+1} - e_{i+2} \leq \text{ST}_i \leq e_{i+1}. \quad (3.1)$$

*Proof.* By (2.12), that  $H$  and ST must satisfy (3.1) is equivalent to

$$\beta_{i+2} \leq e_{i+1} \leq e_{i+2} + \beta_{i+2},$$

which follows from (2.14) and (2.18). Suppose conversely ST satisfies (3.1). Setting  $\beta_i = \text{ST}_{i-2}$  and letting  $\nu_{i+2} = \text{ST}_i - (e_{i+1} - e_{i+2})$ , we have sequences  $\beta$  and  $\nu$  satisfying the conditions (2.14) or (2.13), so by Corollary 2.21 they occur.  $\square$

**Lemma 3.3.** *Let  $H$  satisfy the conditions of (2.8), with  $\lim_{i \rightarrow \infty} H_i = 0$ , and let  $W = \oplus W_i, W_i \subset R_i$  where  $\mu(H) \leq i \leq s-1$  be given vector spaces of dimension  $\dim_k W_i \leq H_i$ . Then there is an ideal  $I$  satisfying both  $I_i \cap W_i = 0$  and  $\tau(I_i) = 1 \forall i \mid \mu \leq i \leq s-1$ . Given a sequence  $\beta$  satisfying (2.14), there is an open dense subset of ideals  $I \in G_\beta(H)$  satisfying  $I \cap W = 0$ .*

*Proof.* The last assertion is a consequence of the first assertion and Theorem 2.17(ii), as the stratum  $H_{\beta_{\max}}$  is in the closure of  $H_\beta$ , and the condition that  $I$  miss  $W$  is an open condition on Artinian algebras  $A = R/I$  in  $H_\beta$ . For the first assertion we may assume without loss of generality that  $\dim W_i = H_i$ . Given the characterization of the extremal Betti stratum in the proof of Proposition 2.7, it suffices to show that given  $W_i$  of dimension  $c = H_i$ , there is a vector space of the form  $fR_{i-c}$ ,  $\deg f = c$ , missing  $W_i$ . Dualizing, we have

$$W_i \cap fR_{i-c} = 0 \Leftrightarrow W_i^\perp + \langle fR_{i-c} \rangle^\perp = \mathcal{R}_i.$$

Let  $f = \prod_{v=1}^c (x - a_v y)$ , with distinct  $a_v$ . Then we have (see, for example, [EmI] or [IK, Lemma 1.15])

$$\langle fR_{i-c} \rangle^\perp = \langle L_1^i, \dots, L_c^i \rangle, \text{ where } L_v = a_v X + Y. \quad (3.2)$$

It is well known and classical that one can assure that  $W_i^\perp \cap L = 0$ , by choosing for  $L$  a complementary-dimension vector space spanned by  $i$ -th powers of linear forms (this is used extensively in [I2]). Thus, one may build  $I$  as follows, beginning in degree  $i = (s-1)$  taking  $f_s = 1$ , and continuing down to degree  $i = \mu$ . We choose as the first part of the  $i$ -labelled step,  $e_{i+1}$  linear forms such that for each degree  $u \leq i$  the  $u$ -th powers of the  $H_i = e_{i+1} + \dots + e_s$  forms chosen so far are linearly independent from  $W_u^\perp$ . In the second part of each step, we determine  $f_i$  by multiplying the previously chosen  $f_{i+1}$  by the degree  $e_{i+1}$  homogeneous form whose linear factors are the annihilators as in (3.2) of the  $e_{i+1}$  linear forms chosen in the first part of the step.  $\square$

**Definition 3.4.** An algebra  $A = R/I$  is a *level algebra* if its socle  $\text{Soc}(A) = 0 : M$  is concentrated in a single degree  $j$ . The level algebra  $A(V)$  determined by a vector subspace  $V \subset R_j$  is the quotient  $A(V) = R/L(V)$  of  $R$  by the *level ideal*  $L(V) = \overline{V} + M^{j+1}$ , satisfying

$$L(V) = R_{-j}V \oplus \dots \oplus R_{-1}V \oplus V \oplus M^{j+1}. \quad (3.3)$$

We denote by  $\text{LA}_N(d, j)$  the family of level algebras having socle type  $d$  in degree  $j$ , and Hilbert function  $N$ .

It follows from (2.12) that  $\text{LA}_N(d, j)$  is just the open dense subscheme Betti stratum of  $G(N)$ , parametrizing those ideals having the minimum possible Betti numbers. We recall the following characterization of level Hilbert functions  $N$  from [I3, Lemma 2.15]:

**Lemma 3.5.** *The Hilbert function  $N = (1, \dots, N_j, 0)$  of a level algebra  $\text{LA}(V)$  determined by the vector subspace  $V \subset R_j$ ,  $\dim V = d$  satisfies*

$$\begin{aligned} N_j &= j+1-d, N_i = 0 \text{ for } i > j, \text{ and} \\ e_{j+1}(N) &= j+1-d \geq e_j(N) = \tau(V) - 1 \geq e_{j-1}(N) \geq \dots, \end{aligned} \quad (3.4)$$

where  $1 \leq \tau(V) \leq \min\{d, j+2-d\}$ .

The following result characterizes the Hilbert function of the intersection of general enough level algebras. The special case for *compressed algebras* — when each  $N(i)$  is the maximum Hilbert function given  $(d_i, j_i)$  — has an analogue in embedding dimension  $r$  that is a consequence of [I2, Proposition 3.6]: for the statement, one replaces  $u+1$  in (3.5) by  $\dim R_u$ . Likewise, there is an analogue in embedding dimension  $r$  for the slightly more general case of *power sum algebras*, whose Hilbert functions  $N(i)$  are the maximum bounded by  $t_i$ , given  $d_i, j_i$  [I2, Theorem 4.8B]. What is unusual in our result here is that there is no restriction on the level algebra Hilbert functions  $N(i)$ . Usually, it is difficult to build algebras of expected “sum” Hilbert functions from intersecting a set of ideals of given Hilbert functions: it is difficult to show that the intersections are dimensionally proper. Of course, it is the embedding dimension two that helps us.

**Theorem 3.6.** *Let  $R = k[x, y]$ . Fix Hilbert functions  $N(1), \dots, N(t)$ , and sequences of integers  $j_1 \geq \dots \geq j_t$ ,  $d_1, \dots, d_t$  where  $1 \leq d_i \leq j_i + 1$ , such that each  $N(i)$  satisfies (3.4) of Lemma 3.5 for the Hilbert function of level algebras  $L(V(i))$  determined by a vector subspace  $V(i)$  of dimension  $d_i$  and degree  $j_i$  in  $R = k[x, y]$ . Let  $I(1), I(2), \dots, I(t)$ , where  $I_i = L(V_i) = \bar{V}_i + M^{j_i+1}$ ,  $V_i \subset R_{j_i}$ ,  $j_1 \geq j_2 \geq \dots \geq j_t$  be general enough level ideals of Hilbert functions  $H(R/I(i)) = N(i)$ . Then the intersection  $I = I(1) \cap \dots \cap I(t)$  satisfies*

$$H(R/I)_u = \min\{\sum H(R/I(i))_u, u + 1\}. \quad (3.5)$$

For general enough  $I(1), \dots, I(t)$  the  $\tau$  sequence  $\tau(I)$  for the intersection  $I$  satisfies for  $u \geq \mu(I)$

$$\tau(I_u) = \sum_{i \geq u} \tau(V_i). \quad (3.6)$$

*Proof.* The key issue is to satisfy (3.5), which is the condition that the vector spaces  $I(1)_u, \dots, I(t)_u$  intersect properly in  $R_u$  for each  $u$ . This is an open condition on the product  $P = \prod_i \text{LA}_{N(i)}(d_i, j_i)$ , an irreducible variety by [I3, Theorem 2.17]. So it suffices to produce a single algebra having that Hilbert function and the given socle type, and such that successively, the algebras determined by  $N(1)$ , by  $\{N(1), N(2)\}, \dots$  (beginning in the highest socle degree) have the expected Hilbert functions. We accomplish this by successive applications of Theorem 2.17 and Lemma 3.3.  $\square$

**Remark 3.7.** BETTI STRATA FOR ACM CODIMENSION TWO PUNCTUAL SCHEMES IN  $\mathbb{P}^2$ , AND ACM CURVES IN  $\mathbb{P}^3$ . The codimension formula of Theorem 2.17 is also valid for the Betti strata of the postulation stratum  $\text{Hilb}^T(\mathbb{P}^2)$  of the punctual Hilbert scheme  $\text{Hilb}^n(\mathbb{P}^2)$ ,  $n = \sum H_i$ ,  $H = \Delta T$  (here  $H$  is termed the  $h$ -vector of  $T$ ), since a punctual scheme is arithmetically Cohen-Macaulay (ACM). The proof can use Gotzmann's result that  $\text{Hilb}^T(\mathbb{P}^2)$  is fibred over  $G(T)$  by an affine space of fixed dimension [Gotz] (see [IK, §5.5, especially p. 180]). A well known result of L. Gruson and C. Peskine, and R. Maggioni and A. Ragusa, is that a reduced, irreducible ACM curve  $X \subset \mathbb{P}^3$  has a *numerical character* without gaps, or, equivalently, has  $H$ -vector  $H_X = \Delta^2 H(k[x_0, x_1, x_2, x_3]/I_X)$  that is *decreasing*: for  $i \geq \mu(H)$ ,  $t_i < \mu \Rightarrow t_i > t_{i+1}$ : that is the  $h$ -vector  $H$  is strictly decreasing after its first decrease from its maximum value of  $\mu$  [GruP, MaR].

Considering ACM curves in  $\mathbb{P}^3$ , T. Sauer has shown that each Betti stratum possible for an Artinian algebra  $A = k[x, y]/I$  having decreasing Hilbert function  $H$  occurs for a smooth ACM curve  $X$  in  $\mathbb{P}^3$  [Sa]. J. Herzog, N. V. Trung and G. Valla show further that a reduced ACM variety  $Y \subset \mathbb{P}^n$  of codimension two is a hyperplane section of a reduced irreducible CM normal variety  $X \subset \mathbb{P}^{n+1}$  of codimension two if and only if the degree-matrix of the minimal resolution satisfies the condition equivalent to the  $H$ -vector decreasing [HTV, Theorem 1.1]. See [GM] for a careful discussion of these results. It follows from the above mentioned results of T. Sauer et al that the codimension formula of Theorem 2.17 for the special case  $H$  decreasing will also give the codimension of the corresponding Betti substratum of the postulation scheme  $\text{Hilb}_{\text{Sm}}^{d,g,T}(\mathbb{P}^3)$  parametrizing smooth ACM curves in  $\mathbb{P}^3$ , of postulation determined by the  $h$ -vector  $H = \Delta T$ , degree  $d = \sum H_i$  and genus  $g = \sum_{i \geq 2} (i-1)H_i$ . The postulation stratum for smooth ACM curves, as well as its Betti substrata are irreducible. Of course, there can be other irreducible components of  $\text{Hilb}^{d,g}(\mathbb{P}^3)$  corresponding to non-ACM curves [PS].

**Acknowledgment.** *We acknowledge gratefully discussions with Mats Boij, whose method in studying the Betti strata of height three Gorenstein algebras we apply here. Discussions with Jacques Emsalem and Vassil Kanev were helpful background for this article. We are grateful also to a referee for [I3] who suggested making this a separate article. We thank Juan Migliore for some comments.*

# References

- [ACGH] Arbarello E., Cornalba M., Griffiths P.A., Harris J.: *Geometry of Algebraic Curves*, Vol. 1, Grundlehren der math. Wissen. #267, (1985), Springer-Verlag, New York.
- [Bj] Boij M.: : *Betti number strata of the space of codimension three Gorenstein Artin algebras*, preprint, 2001.
- [BV] Bruns W, Vetter U.: *Determinantal rings*, Springer LNM #1327, Springer, Berlin, 1988.
- [Cam] Campanella G.: *Standard bases of perfect homogeneous polynomial ideals of height two*, J. Algebra **101** (1986), 47–60.
- [CaCh] Carlini E., Chipalkatti J.: *On Waring’s problem for several algebraic forms*. Comment. Math. Helv. 78 (2003), no. 3, 494–517.
- [Di] Diesel S. J.: *Some irreducibility and dimension theorems for families of height 3 Gorenstein algebras*, Pacific J. Math. **172** (1996), 365–397.
- [Ell] Ellingsrud G.: *Sur le schéma de Hilbert des variétés de codimension 2 dans  $\mathbb{P}^e$  a cône de Cohen-Macaulay*, Ann. Scient. Ecole Norm. Sup. 4<sup>e</sup> serie, t.8 (1975), 423–432.
- [EmI] Emsalem J., Iarrobino A.: *Inverse system of a symbolic power I*, J. Algebra **174** (1995), 1080–1090.
- [Fr] Fröberg R.: *An inequality for Hilbert series of graded algebras*, Math. Scandinavica **56** (1985), 117–144.
- [GM] Geramita A. , Migliore, J.: *Hyperplane sections of a smooth curve in  $\mathbb{P}^3$* , Comm. in Algebra **17**(12), (1989), 3129–3164.
- [GPS] ———, Pucci M., Shin Y.S.: *Smooth points of  $\text{Gor}(T)$* , J. of Pure and Applied Algebra, 122 (1997), 209–241.
- [Gotz] Gotzmann G.: *A stratification of the Hilbert scheme of points in the projective plane*, Math Z. **199** (1988) no. 4, 539–547.
- [GruP] Gruson L., Peskine C.: *Genre des courbes de l’espace projectif*, in Algebraic Geometry, (Proc. Sympos., Univ. Tromsø, Tromsø,1977), pp. 31–59, Lecture Notes in Math. #687, (1978), Springer, Berlin.
- [Hari] Harima T.: *Some examples of unimodal Gorenstein sequences*, J. Pure and Applied Algebra **103** (1995), 313–324.
- [HTV] Herzog J., Trung N.V., Valla G.: *On hyperplane sections of reduced irreducible varieties of low codimension*, J. Math. Kyoto Univ. **34** (1994), no. 1, 47–72.
- [I1] Iarrobino A.: *Punctual Hilbert schemes*, A.M.S. Memoirs Vol. 10, no. 188, Amer. Math. Soc. 1977.
- [I2] ———: *Compressed algebras: Artin algebras having given socle degrees and maximal length*, Transactions of the A.M.S., vol. 285, no. 1 pp. 337–378, 1984.
- [I3] ———: *Ancestor ideals of vector spaces of forms, and level algebras*, J. Algebra 272 (2004), 530–580.

- [IK] ———, Kanev V.: *Power Sums, Gorenstein Algebras, and Determinantal Loci*, 345+xxvii p. (1999) Springer Lecture Notes in Mathematics # 1721, Springer, Heidelberg.
- [IY] ———, Yaméogo J.: *The family  $G_T$  of graded Artinian quotients of  $k[x, y]$  of given Hilbert function*, Communications in Algebra, Vol. 31 # 8 (2003), 3863–3916.
- [Klp] Kleppe, J. O.: *Deformations of graded algebras* Math. Scand. 45 (1979), 205–231.
- [MaR] Maggione R., Ragusa A.: *The Hilbert function of generic plane sections of curves of  $\mathbb{P}^3$* , Invent. Math. **91** (1988), no. 2, 253–258.
- [PS] Piene R. , Schlessinger M.: *On the Hilbert scheme compactification of the space of twisted cubics*, Amer. J. of Math, 107 (1985), 761–774.
- [Sa] Sauer T.: *Smoothing projectively Cohen-Macaulay space curves*, Math. Ann. **272**, (1985), 83–90.
- [W] Weyman J.: *Cohomology of Vector Bundles and Syzygies*, Camb. Tracts in Math. #149, Cambridge Univ. Press, 2003.