PRINCIPLE SUBSPACE FOR BOSONIC VERTEX OPERATOR $\phi_{\sqrt{2m}}(z)$ AND JACK POLYNOMIALS

B.FEIGIN AND E.FEIGIN

ABSTRACT. Let $\phi_{\sqrt{2m}}(z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-m}$, $m \in \mathbb{N}$ be bosonic vertex operator, L some irreducible representation of the vertex algebra $\mathfrak{A}_{(m)}$, associated with one-dimensional lattice $\mathbb{Z}l$, generated by the vector l, $\langle l, l \rangle = 2m$. Fix some extremal vector $v \in L$. We study the principal subspace $\mathbb{C}[a_i]_{i \in \mathbb{Z}} \cdot v$ and its finitization $\mathbb{C}[a_i]_{i>N} \cdot v$. We construct their bases and find characters. In the case of finitization basis is given in terms of Jack polynomials.

INTRODUCTION

Let $L_{0,1}$ and $L_{1,1}$ be irreducible representations of the Lie algebra \mathfrak{sl}_2 , $v_{2n} \in L_{0,1}, v_{2n+1} \in L_{1,1}$ the set of extremal vectors (for example, v_0 is a vacuum vector). For $x \in \mathfrak{sl}_2$ consider the current $x(z) = \sum_{i \in \mathbb{Z}} x_i z^{-i-1}$ (here we use the notation $x_i = x \otimes t^i$). Let e, h, f be standard basis of \mathfrak{sl}_2 . Then $e_i v_p = 0$ for $i \geq p$. Consider the principle subspace $V_p = \mathbb{C}[e_{p-1}, e_{p-2}, \ldots] \cdot v_p$. Let us list some properties of V_p (see [FL, FF1, FF2, CP]).

1) $V_p \simeq \mathbb{C}[e_{p-1}, e_{p-2}, \ldots]/I_p$, where I_p is an ideal, generated by coefficients of series $(e_{p-1} + ze_{p-2} + z^2e_{p-3} + \ldots)^2$.

2) Elements

$$e_{i_1} \cdots e_{i_k} v, \ i_1 < p, \ i_\alpha - i_{\alpha+1} \ge 2, \ k = 0, 1, \dots$$

form the basis of V_p . Using this basis one can write formula for the character of V_p and construct semi-infinite basis of $L_{0,1}$ and $L_{1,1}$.

3) Consider finitization: the subspace $V_p(n) \hookrightarrow V_p$, $V_p(n) = \mathbb{C}[e_{p-1}, \ldots, e_{p-n}] \cdot v_p$. Then dim $V_p(n) = 2^n$.

Recall that the current e(z) can be realized as bosonic vertex operator $\phi_{\sqrt{2}}(z)$. In this paper we generalize the above results to the case of $\phi_{\sqrt{2m}}(z)$ for an arbitrary $m \in \mathbb{N}$ (see [FJM] for the discussion on this topic).

Let $\mathfrak{A}_{(m)}$ be lattice vertex algebra, associated with a one-dimensional lattice $\mathbb{Z}l$, generated by vector l, $\langle l, l \rangle = 2m$. Let $L_{(m),i}$ be the set of irreducible representations of $\mathfrak{A}_{(m)}$ (see [D]). We have

$$L_{(m),i} = \bigoplus_{n \in \mathbb{Z}} H_{\frac{2nm+i}{\sqrt{2m}}}, \quad 0 \le i \le 2m - 1.$$

Here H_{λ} is highest weight representation of the Heisenberg algebra $H = span\{h_i\}, i \in \mathbb{Z}$ with a highest weight vector $|\lambda\rangle$. Fix an action of the Virasoro algebra on $L_{(m),i}$:

$$L_n = \frac{1}{2} \sum_{i+j=n} : h_i h_j : -\frac{m-1}{\sqrt{2m}} (n+1)h_n.$$

Introduce a notation for the extremal vectors of $L_{(m),i}$: $v_p = |\frac{-p}{\sqrt{2m}}\rangle$, $p \in \mathbb{Z}$. Note that $a_i v_p = 0$ for $i \ge p$. Following [FS] define the principle subspace

$$V_{(m),p} = \mathbb{C}[a_{p-1}, a_{p-2}, \ldots] \cdot v_p.$$

First we describe such ideal $I_{(m),p}$ that $V_{(m),p} \simeq \mathbb{C}[a_{p-1}, a_{p-2}, \ldots]/I_{(m),p}$. Namely, $I_{(m),p}$ is generated by the coefficients of series

$$((a_{p-1} + a_{p-2}z + a_{p-3}z^2 + \cdots)^{(i)})^2, \quad 0 \le i < m,$$

where superscript (i) stands for the *i*-th derivative. Using functional realization (see [FS]) of the dual space $V^*_{(m),p}$ we obtain its character, construct the fermionic realization of $V_{(m),p}$ and find its monomial basis. Namely we prove that elements

$$a_{i_1} \cdots a_{i_k} v_p, \ i_1 \le p-1, \ i_\alpha - i_{\alpha+1} \ge 2m, \ k = 0, 1, \dots$$

form the basis of $V_{(m),p}$. We also show that this basis is compatible with the operation of taking coinvariants. It means that the images of vectors $a_{i_1} \cdots a_{i_k} v_p$ with

$$i_{\alpha} - i_{\alpha+1} \ge 2m, \ i_1 \le p - 1, \ i_k > N$$

form basis of the quotient $V_{(m),p}/span\{a_i V_{(m),p}, i \leq N\}$.

Now consider the subspace

$$V_{(m),p}(n) \hookrightarrow V_{(m),p}, \ V_{(m),p}(n) = \mathbb{C}[a_{p-1}, a_{p-2}, \dots, a_{p-n}] \cdot v_p.$$

Define numbers $F_i^{(m)}$ as follows:

$$F_i^{(m)} = i+1, \ i = 0, 1, \dots, m-1; \ F_{i+m}^{(m)} = F_i^{(m)} + F_{i+m-1}^{(m)}, \ i \ge 0.$$

For example, $F_i^{(1)} = 2^i$ and $F_i^{(2)}$ are Fibonacci numbers. We prove that the dimension of $V_{(m),p}(n)$ equals to $F_n^{(m)}$. The proof is based on the construction of basis of $V_{(m),p}(n)$ in terms of Jack polynomials $J_{\lambda}(m; \mathbf{x})$. (Which reduce to Schur polynomials in $\widehat{\mathfrak{sl}}_2$ case).

Namely, let $h_i, i \in \mathbb{Z}$ be the basis of Heisenberg Lie algebra H. Then each space H_{λ} can be regarded as the space of polynomials in infinite number of variables via the identification (h_i with i < 0 are generating operators)

$$h_{i_1}\cdots h_{i_k}|\lambda\rangle \mapsto \left(\sqrt{\frac{m}{2}}\right)^k p_{-i_1}\cdots p_{-i_k}, \ i_\alpha < 0,$$

where p_j is the *j*-th power sum. Combining results from [TY, MY] (see also [AMOS]) we obtain that

$$a_0^k v_p = c \cdot J_{((p-(k-1)m-1)^k)}(m; \mathbf{x}),$$

where c is some non-vanishing constant and $((p - (k - 1)m - 1)^k)$ is the partition

$$\underbrace{(p-(k-1)m-1,\ldots,p-(k-1)m-1)}_k.$$

For $p \in \mathbb{Z}$ let $V_{(m),p}^{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}(n) = V_{(m),p} \cap H_{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}$. Note that because of the independence (up to a shift of variables) of $V_{(m),p}$ on p spaces $V_{(m),n}(p)$ and $V_{(m),p}(p)$ are isomorphic. Denote the latter space simply by $V_{(m)}(p)$. We show that

$$V_{(m)}^{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}(p) = \mathbb{C}[h_1, h_2, \ldots] \cdot a_0^k v_p$$

and in addition $\mathbb{C}[h_1, h_2, \ldots] \cdot J_{((p-m(k-1)-1)^k)}(m; \mathbf{x})$ is a linear span of Jack polynomials $J_{\lambda}(m; \mathbf{x})$ with Young diagram λ being the subdiagram of $((p-m(k-1)-1)^k)$. This gives us the dimension and character of $V_{(m)}(p)$.

Recall that in [FF1] spaces $V_{(1)}(n)$ were obtained via the fusion procedure. Let $Z = (z_1, \ldots, z_n) \in \mathbb{C}^n, z_i \neq z_j$. Consider the ring $R = \mathbb{C}[y_1, \ldots, y_n]/\langle y_i^2 \rangle_{1 \leq i \leq n}$ and pick its generators $a_i = \sum_{j=1}^n y_j z_j^i$, $i = 0, \ldots, n-1$. We have

$$R \simeq \mathbb{C}[a_0, \ldots, a_{n-1}]/I_Z(n),$$

where $I_Z(n)$ is some ideal. Then the following is true:

$$V_{(1)}(n) \simeq \mathbb{C}[a_0, \dots, a_{n-1}] / \lim_{Z \to 0} I_Z(n).$$

Conjecturally, the analogous construction exists for the general m. Namely one must put R to be $\mathbb{C}[y_1, \ldots, y_n]/\langle y_i y_j \rangle_{|i-j| < m}$. Note that the dimension of this ring obviously equals $F_n^{(m)}$. We discuss this conjecture in the last section.

As it was mentioned above $L_{(m),i}$ generalize irreducible representations $L_{0,1}$ and $L_{1,1}$ of $\widehat{\mathfrak{sl}_2}$ on the level 1. Using the semi-infinite construction of $L_{(m),i}$ (see [FS] for the case m = 1) we obtain bases of the latter spaces.

The paper is organized as follows.

In the first section we consider the quotient of $\mathbb{C}[\xi_0, \xi_1, \ldots]$ by the ideal, generated by coefficients of series $((\xi_0 + \xi_1 z + \ldots)^{(k)})^2$, $k = 0, 1, \ldots, m-1$. We find basis of this quotient and basis of the space of coinvariants.

In the second section we study the principle subspaces $V_{(m),p}$ and finitizations $V_{(m)}(p)$. We find relations between Fourier coefficients of $\phi_{\sqrt{2m}}(z)$ and construct bases of $V_{(m),p}$ and $V_{(m)}(p)$.

Third section is devoted to the computation of character and dimension of $V_{(m)}(p)$.

Fourth section contains construction of semi-infinite basis for $L_{(m),i}$.

And in the last section we discuss some possible generalizations and open questions.

Acknowledgment. EF wants to thank V.Dotsenko for helpful information about Jack polynomials. First named author was partially supported by grants RFBR 02.01.01015, SS 2044.2003.2 and INTAS 03-51-3350. Second named author was partially supported by grant RFBR 03-01-00167.

1. Commutative Algebras

Let ξ_0, ξ_1, \ldots be some commuting variables. Consider an algebra

(1)
$$A_{(m)} = \mathbb{C}[\xi_0, \xi_1, \ldots] / \langle (\xi(z)^{(i)})^2 \rangle_{0 \le i \le m-1}.$$

Here $\xi(z) = \sum_{j=0}^{\infty} \xi_j z^j$, $\xi(z)^{(i)}$ its *i*-th derivative, and the ideal of relations (the right hand side of (1)) is generated by the coefficients of $(\xi(z)^{(i)})^2$, $i = 0, 1, \ldots, m-1$. Surely, $A_{(m)}$ is bi-graded:

$$A_{(m)} = \bigoplus_{k,s \ge 0} A_{(m)}^{k,s}, \ A_{(m)}^{k,s} = span\{\xi_{i_1} \cdots \xi_{i_k}, \ \sum_{\alpha=1}^k i_\alpha = s\}.$$

Define $\operatorname{ch}(A_{(m)}) = \sum_{k,s \ge 0} z^k q^s \dim A_{(m)}^{k,s}$.

Lemma 1.1. $ch(A_{(m)}) = \sum_{k \ge 0} z^k \frac{q^{mk(k-1)}}{(1-q)\cdots(1-q^k)}$

Proof. The dual space $(A_{(m)}^{k,s})^*$ can be realized as a subspace of symmetric polynomials $f(z_1, \ldots, z_k)$ of degree s. Namely for $\theta \in (A_{(m)}^{k,s})^*$ define

$$f_{\theta}(z_1,\ldots,z_k) = \sum_{i_1,\ldots,i_k \ge 0} z_1^{i_1} \cdots z_k^{i_k} \theta(\xi_{i_1} \cdots \xi_{i_k}).$$

One can show that $f = f_{\theta}$ for some $\theta \in (A_{(m)}^{k,s})^*$ if and only if

(2)
$$f(z_1, \dots, z_k) = \prod_{1 \le i < j \le k} (z_i - z_j)^{2m} g(z_1, \dots, z_k)$$

with some symmetric g. (This is because of the condition $(\xi(z)^{(i)})^2 = 0$ for $i = 0, \ldots, m-1$). Formula (2) gives us the character of $A_{(m)}$.

Now we want to construct the monomial basis of $A_{(m)}$. In order to do that we embed $A_{(m)}$ into the algebra, generated by fermions. Consider algebra F, generated by variables $\psi_s(i)$, $s = 1, \ldots, 2m$, $i \in \mathbb{N} \cup \{0\}$, subject to the relations $\psi_s(i)\psi_t(j) = -\psi_t(j)\psi_s(i)$. Define the currents $\psi_s(z) = \sum_{i\geq 0} \psi_s(i)z^i$. Let $\tilde{\xi}_i \in F$, $i \geq 0$, be elements defined by

$$\tilde{\xi}(z) = \sum_{i \ge 0} \tilde{\xi}_i z^i = \psi_1(z) \cdots \psi_{2m}(z).$$

Lemma 1.2. Algebra, generated by $\tilde{\xi}_i$, $i \ge 0$ is isomorphic to $A_{(m)}$ via the identification $\tilde{\xi}_i \mapsto \xi_i$.

Proof. First it is easy to check that $(\tilde{\xi}(z)^{(i)})^2 = 0$ for $i = 0, \ldots, m-1$. We need to prove that these are defining relations. To do that it is enough to find the set of linearly independent monomials in variables $\tilde{\xi}_0, \tilde{\xi}_1, \ldots$, whose character (with respect to the grading $\deg_z \tilde{\xi}_i = 1$, $\deg_q \tilde{\xi}_i = i$) is equal to the character of $A_{(m)}$.

Consider the set of admissible monomials

$$\xi_{i_1}\cdots\xi_{i_k}, \ 0 \le i_1, \ i_{\alpha+1}-i_{\alpha} \ge 2m, \ \alpha=1,\ldots,k-1.$$

We prove that they are linearly independent. In fact, let some linear combination of admissible monomials vanish:

(3)
$$\sum \alpha_{i_1,\dots,i_k} \tilde{\xi}_{i_1} \cdots \tilde{\xi}_{i_k} = 0.$$

Pick monomial $\tilde{\xi}_{i_1^0} \cdots \tilde{\xi}_{i_k^0}$ with a nonzero $\alpha_{i_1^0, \dots, i_k^0}$ such that for any other k-tuple (i_1, \dots, i_k) with $\alpha_{i_1, \dots, i_k} \neq 0$ the following is true: there exists such l < k that

$$i_1^0 = i_1, \dots, \ i_l^0 = i_l, \ i_{l+1}^0 > i_{l+1}.$$

We state that $\tilde{\xi}_{i_1^0} \cdots \tilde{\xi}_{i_k^0}$ can not be written as a sum of other monomials from (3). In fact,

(4)
$$\tilde{\xi}_{i_1^0} \cdots \tilde{\xi}_{i_k^0} = \prod_{s=1}^{\kappa} \sum_{\beta_1^s + \dots + \beta_{2m}^s = i_s^0} \psi_1(\beta_1^s) \cdots \psi_{2m}(\beta_{2m}^s).$$

Right hand side of (4) contains nonzero (because of the condition $i_{\alpha+1} - i_{\alpha} \ge 2m$) term

(5)
$$\prod_{s=1}^{k} \psi_1\left(\left[\frac{i_s^0}{2m}\right]\right) \psi_2\left(\left[\frac{i_s^0+1}{2m}\right]\right) \cdots \psi_{2m}\left(\left[\frac{i_s^0+2m-1}{2m}\right]\right)$$

([x] is a maximal integer not exceeding x). By definition of (i_1^0, \ldots, i_k^0) the term (5) appears in the sum (3) only from the product $\tilde{\xi}_{i_1^0} \cdots \tilde{\xi}_{i_k^0}$. Thus (3) is impossible.

To finish the proof note that the character of admissible monomials coincide with the character of $A_{(m)}$.

Corollary 1.1. $\xi_{i_1} \cdots \xi_{i_k}$, $i_{\alpha+1} - i_{\alpha} \ge 2m$, form basis of $A_{(m)}$.

Proposition 1.1. Images of the monomials $\xi_{i_1} \cdots \xi_{i_k}$, $i_{\alpha+1} - i_{\alpha} \ge 2m$, $i_k \le N$ in the quotient $A_{(m)}/span\{\xi_i A_{(m)}, i > N\}$ form the basis of the latter quotient.

Proof. Note that the term (5) with $i_k^0 \leq N$ can not appear in the sum of the form

(6)
$$\xi_{N+1}x_1 + \xi_{N+2}x_2 + \ldots + \xi_{N+p}x_p, \ x_i \in A_{(m)}.$$

In fact, substituting ξ_i instead of ξ_i we obtain that (6) is a linear combination of the products of the fermions

$$\prod_{s=1}^{2m} \psi_s(\beta_1^s) \cdots \psi_s(\beta_k^s)$$

with a property that there exists such map $\sigma : \{1, \ldots, 2m\} \to \{1, \ldots, k\}$ that $\beta_{\sigma(1)}^1 + \ldots + \beta_{\sigma(2m)}^{2m} > N$. But the latter is not true for (5). Thus the sum (6) does not contain monomial (5). Proposition is proved.

2. VERTEX OPERATORS AND FINITIZATION

Consider the lattice vertex operator algebra $\mathfrak{A}_{(m)}$, associated with one-dimensional lattice $\mathbb{Z}l$, generated by vector l, $\langle l, l \rangle = 2m$. Let $\phi_{\sqrt{2m}}(z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-m}$ be corresponding bosonic vertex operator. Consider the set $L_{(m),i}$ of irreducible representations of $\mathfrak{A}_{(m)}$. Then

$$L_{(m),i} = \bigoplus_{n \in \mathbb{Z}} H_{\frac{2nm+i}{\sqrt{2m}}}, \ 0 \le i \le 2m-1.$$

Here H_{λ} is highest weight representation of the Heisenberg algebra H with a basis $h_i, i \in \mathbb{Z}$. We denote the highest weight vector by $|\lambda\rangle$. Consider the action of Virasoro algebra on $L_{(m),i}$:

$$L_n = \frac{1}{2} \sum_{i+j=n} : h_i h_j : -\frac{m-1}{\sqrt{2m}} (n+1)h_n,$$

where : : is a normal ordering sign.

Spaces $L_{(m),i}$ are bi-graded by operators h_0 (z-degree) and L_0 (q-degree). For any subspace $V \hookrightarrow L_{(m),i}$ define

$$\operatorname{ch}(z,q,V) = \operatorname{Tr}(z^{h_0}q^{L_0}|_V).$$

Denote

$$v_p = \left|\frac{-p}{\sqrt{2m}}\right\rangle \in \bigoplus_{i=0}^{2m-1} L_{(m),i}$$

For the eigenvector v of the operators h_0 and L_0 we denote the corresponding eigenvalues by $\deg_z v$ and $\deg_q v$. For example

$$\deg_z v_p = \frac{-p}{\sqrt{2m}}, \ \deg_q v_p = \frac{p^2}{4m} + \frac{p(m-1)}{2m}.$$

B.FEIGIN AND E.FEIGIN

Introduce the principle subspace $V_{(m),p} = \mathbb{C}[a_{p-1}, a_{p-2}, \ldots] \cdot v_p$ (note that $a_i v_p = 0$ for $i \ge p$ because the difference of q-degrees of v_p and v_{p-2m} equals p).

In what follows we use the connection between vertex operators and Jack polynomials from [MY] (see also [AMOS]). Recall the definition of Jack polynomials (see [Mac]). Let Λ be the algebra of symmetric polynomials in infinite number of variables, $p_k \in \Lambda$ – power-sums, $p_k = \sum_{i \in \mathbb{N}} x_i^k$. Define the scalar product on $\Lambda = \mathbb{C}[p_1, p_2, \ldots]$, depending on the coupling constant α :

$$\langle p_1^{i_1}\cdots p_k^{i_k}, p_1^{i_1}\cdots p_k^{i_k}\rangle_{\alpha} = \alpha^{-\sum_{s=1}^k i_s} \prod_{s=1}^k s^{i_s} i_s!$$

and the products of power sums form the orthogonal basis of Λ with respect to $\langle \cdot, \cdot \rangle_{\alpha}$.

Now let $\lambda = (\lambda_1 \geq \lambda_2 \ldots \geq \lambda_s \geq 0)$ be partition. Let $l(\lambda)$ be the length of λ , i.e. the number of nonzero λ_i . Young diagram, attached to λ , is the following subset of \mathbb{Z}^2 : $\{(i, j) : 1 \leq i \leq l(\lambda), 1 \leq j \leq \lambda_i\}$. Jack polynomials $J_{\lambda}(\alpha; \mathbf{x}) = J_{\lambda}(\alpha; x_1, x_2, \ldots) \in \Lambda$ depend on the partition λ and coupling constant α . They are uniquely determined by following properties $(m_{\lambda} \text{ is symmetrization of the monomial } x_1^{\lambda_1} \cdots x_s^{\lambda_s})$:

$$J_{\lambda}(\alpha, \mathbf{x}) = m_{\lambda} + \sum_{\mu < \lambda} v_{\lambda,\mu}(\alpha) m_{\mu}$$
$$\langle J_{\lambda}(\alpha, \mathbf{x}), J_{\mu}(\alpha, \mathbf{x}) \rangle_{\alpha} = 0 \text{ if } \lambda \neq \mu,$$

where for two partitions λ, μ we write $\mu \leq \lambda$ if $\mu_1 + \ldots + \mu_i \leq \lambda_1 + \ldots + \lambda_i$ for any *i*. Note that in the similar way one can define Jack polynomials in finite number of variables. In this case $J_{\lambda}(\alpha; x_1, \ldots, x_N) = 0$ if $N < l(\lambda)$.

Now fix $m \in \mathbb{N}$ and consider Fock space $H_s = \mathbb{C}[h_{-1}, h_{-2}, \ldots] \cdot |s\rangle$. Identify H_s with Λ in the following way:

$$I: h_{-i_1} \cdots h_{-i_k} | s \rangle \mapsto \left(\sqrt{\frac{m}{2}}\right)^k p_{i_1} \dots p_{i_k}$$

Then $h_i, i > 0$ acts on Λ by $i\sqrt{\frac{2}{m}}\frac{\partial}{\partial p_i}$. Denote by $\widehat{J}_{\lambda}(m; \mathbf{x})$ vector $I^{-1}(J_{\lambda}(m, \mathbf{x}))$. Following theorem was proved in [MY]:

Theorem 2.1. For any p > 0 the following equality holds in $H_{\frac{-p}{2}+k\sqrt{2m}}$:

$$a_0^k v_p = C \cdot \widehat{J}_{((p-m(k-1)-1)^k)}(m; \mathbf{x}),$$

where C is some nonzero constant and (N^k) is a partition $(\underbrace{N,\ldots,N}_k)$.

Recall the principle subspace $V_{(m),p}$. First we study

$$V_{(m),-m+1} = \mathbb{C}[a_{-m}, a_{-m-1}, \ldots] \cdot v_{1-m}.$$

We denote it simply by $V_{(m)}$.

Lemma 2.1. Let
$$\phi^+_{\sqrt{2m}}(z) = a_{-m} + za_{-m-1} + z^2 a_{-m-2} + \cdots$$
. Then
 $V_{(m)} \simeq \mathbb{C}[a_{-m}, a_{-m-1}, \ldots] / \langle (\phi^+_{\sqrt{2m}}(z)^{(i)})^2 \rangle_{0 \le i \le m-1},$

where $\phi^+_{\sqrt{2m}}(z)^{(i)}$ is the *i*-th derivative. It means that the ideal of relations in $V_{(m)}$ is generated by the coefficients of series $(\phi_{\sqrt{2m}}^+(z)^{(i)})^2$, namely by

$$\sum_{\substack{\alpha_1 + \alpha_2 = s \\ \alpha_1, \alpha_2 \ge 0}} a_{-m-\alpha_1} a_{-m-\alpha_2} \alpha_1(\alpha_1 - 1) \cdots (\alpha_1 - i + 1) \alpha_2(\alpha_2 - 1) \cdots (\alpha_2 - i + 1),$$

$$s \ge 0, \quad i = 0, \dots, m - 1.$$

Proof. Recall that

$$\phi_{\sqrt{2m}}(z)\phi_{\sqrt{2m}}(w) = (z-w)^{2m} : \phi_{\sqrt{2m}}(z)\phi_{\sqrt{2m}}(w) : .$$

Thus because of $a_i v_{1-m} = 0$ for i > -m we obtain $(\phi_{\sqrt{2m}}^+(z)^{(i)})^2 v_{1-m} = 0, 0 \le 0$ $i \leq m-1$. Because of the lemma (1.1) it is enough to prove that the character of $V_{(m)}$ is greater or equal (in each weight component) then

$$z^{\frac{m-1}{\sqrt{2m}}}q^{\frac{-(m-1)^2}{4m}}\sum_{k=0}^{\infty}\frac{z^kq^{mk^2}}{(1-q)(1-q^2)\cdots(1-q^k)}.$$

(Note that $\deg_q v_{1-m} = \frac{-(m-1)^2}{4m} v_{1-m}$, $\deg_z v_{1-m} = \frac{m-1}{\sqrt{2m}} v_{1-m}$.) Consider the decomposition $V_{(m)} = \bigoplus_{k,s \ge 0} \tilde{V}_{(m)}^{k,s}$,

$$\tilde{V}_{(m)}^{k,s} = span\{a_{i_1}\dots a_{i_k}v_{1-m}, \ \sum_{\alpha=1}^k i_\alpha = -s\}.$$

Denote $\tilde{V}_{(m)}^k = \bigoplus_s \tilde{V}_{(m)}^{k,s}$. Identify $(\tilde{V}_{(m)}^k)^*$ with the subspace of symmetric polynomials in k variables in the following way:

$$(\tilde{V}_{(m)}^k)^* \ni \theta \mapsto f_\theta(z_1, \dots, z_k) = \sum_{i_1, \dots, i_k \ge 0} z_1^{i_1} \cdots z_k^{i_k} \theta(a_{-i_1} \cdots a_{-i_k} v_{1-m}).$$

Because of the relations on $\phi^+_{\sqrt{2m}}(z)$ we obtain that f_{θ} is of the following form:

(7)
$$f_{\theta}(z_1, \dots, z_k) = \prod_{i=1}^k z_i^m \prod_{1 \le i < j \le k} (z_i - z_j)^{2m} g(z_1, \dots, z_k),$$

where q is some symmetric polynomial. Note that character of the right hand side of (7) with g running over all symmetric polynomials in k variables coincides with $\frac{q^{mk^2}}{(1-q)(1-q^2)\cdots(1-q^k)}$. Thus we only need to prove that any symmetric g appears in

the right hand side of (7) for some θ . First note that $\tilde{V}_{(m)}^{k,mk^2}$ is a subspace of one-dimensional space, spanned by vector v_{1-m-2k} . At the same time one can show that $\tilde{V}_{(m)}^{k,mk^2}$ is not trivial. Thus for some nontrivial element $\theta \in (\tilde{V}_{(m)}^{k,mk^2})^*$

$$f_{\theta}(z_1, \dots, z_k) = \prod_{i=1}^k z_i^m \prod_{1 \le i < j \le k} (z_i - z_j)^{2m}.$$

Because of the relations $[h_i, a_j] = \sqrt{2m} a_{i+j}$ operators h_i with i > 0 acts on $\tilde{V}^k_{(m)}$. Let h^*_i be the dual operators, acting on $(\tilde{V}^k_{(m)})^*$. One can see that

$$f_{h_i^*\theta}(z_1,\ldots,z_k) = \sqrt{2m} \left(z_1^i + \ldots + z_k^i \right) f_{\theta}(z_1,\ldots,z_k).$$

To finish the proof it is enough to mention that power sums generates the algebra of symmetric polynomials. $\hfill \Box$

Following lemma can be easily verified.

Lemma 2.2. The map
$$\mathbb{C}[a_{-m}, a_{-m-1}, \ldots] \cdot v_{1-m} \to \mathbb{C}[a_{p-1}, a_{p-2}, \ldots] \cdot v_p$$
,

 $a_{i_1}\cdots a_{i_k}v_{1-m}\mapsto a_{i_1+m+p-1}\cdots a_{i_k+m+p-1}v_p$

is well-defined isomorphism.

Now we will study the finitization of $V_{(m),p}$. Namely for p > 0 define the subspace

$$V_{(m),p} \leftarrow V_{(m)}(p) = \mathbb{C}[a_{p-1}, a_{p-2}, \dots, a_0] \cdot v_p.$$

Introduce a notation

$$V_{(m)}^{k,s}(p) = \{ v \in V_{(m)}(p) : h_0 v = kv, L_0 v = sv \}, \quad V_{(m)}^k(p) = \bigoplus_s V_{(m)}^{k,s}(p).$$

Following lemma is an immediate consequence from the lemma (2.2).

Lemma 2.3. Denote $V_{(m),p}(n) = \mathbb{C}[a_{p-1}, \ldots, a_{p-n}] \cdot v_p$. Then the map

$$V_{(m),p}(n) \to V_{(m)}(n), \quad a_{i_1} \cdots a_{i_k} v_p \mapsto a_{i_1-p+n} \cdots a_{i_k-p+n} v_n$$

is an isomorphism.

Lemma 2.4.
$$V_{(m)}^{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}(p) = \mathbb{C}[h_1, h_2, \ldots] \cdot a_0^k v_p.$$

Proof. Note that $[h_i, a_j] = \sqrt{2m} a_{i+j}$. We need to prove that for any $p-1 \ge i_1 \ge i_2 \ge \ldots \ge i_k \ge 0$ we have

$$a_{i_1}\cdots a_{i_k}v_p \in \mathbb{C}[h_1, h_2, \ldots] \cdot a_0^k v_p.$$

Let us use the induction on r: the number of such α that $i_{\alpha} \neq 0$.

The case r = 0 is trivial. Suppose our lemma is proved for some p. Then

$$h_{i_{r+1}}(a_{i_1}\cdots a_{i_r}a_0^{k-r}v_p) = \sqrt{2m} (k-r)a_{i_1}\cdots a_{i_r}a_{i_{r+1}}a_0^{k-r-1}v_p + \sqrt{2m} a_0^{k-r} \left(\sum_{j=1}^r a_{i_1}\cdots a_{i_j+i_{r+1}}\cdots a_{i_r}\right)v_p.$$

Thus by induction assumption

$$a_{i_1} \cdots a_{i_r} a_{i_{r+1}} a_0^{k-r-1} v_p \in \mathbb{C}[h_1, h_2, \ldots] \cdot a_0^k v_p.$$

Lemma is proved.

As it was mentioned above if we identify H_{α} with the space Λ of symmetric polynomials in infinite number of variables, then h_i with i > 0 will act on Λ (up to a nonzero constant) as $\frac{\partial}{\partial p_i}$. Thus because of the above lemma and theorem (2.1) it is important to study the space

$$\mathbb{C}\left[\frac{\partial}{\partial p_1}, \frac{\partial}{\partial p_2}, \ldots\right] \cdot J_{\lambda}(m; \mathbf{x})$$

for the rectangular diagram λ .

Consider the polynomial $g \in \mathbb{C}[x_2, x_3, \ldots]$. Denote by $(Lg)(x_1, x_2, \ldots)$ the polynomial obtained from $g(x_2, x_3, \ldots)$ by substituting x_i instead of x_{i+1} , $i = 1, 2, \ldots$

8

Proposition 2.1. Define a collection of maps $T_n : \Lambda \to \Lambda$, n = 1, 2, ...

$$T_n(f(x_1, x_2, \ldots)) = L\left(\frac{\partial^n f}{\partial x_1^n}\Big|_{x_1=0}\right).$$

Then two algebras of operators on Λ are equal:

$$\mathbb{C}\left[\frac{\partial}{\partial p_1}, \frac{\partial}{\partial p_2}, \ldots\right] = \mathbb{C}[T_1, T_2, \ldots].$$

Proof. We prove that for any n > 0

$$\mathbb{C}\left[\frac{\partial}{\partial p_1}, \frac{\partial}{\partial p_2}, \dots, \frac{\partial}{\partial p_n}\right] = \mathbb{C}[T_1, T_2, \dots, T_n].$$

Let $f \in \Lambda$, $f = f(p_1, p_2, ...)$. Then there exist polynomials $b_{j,n}(k_1, ..., k_j)$, $j = 1, ..., n, k_j \in \mathbb{N}$, such that

(8)
$$\frac{\partial^n f}{\partial x_1^n} = \sum_{j=1}^n \sum_{\substack{k_1, \dots, k_j \ge 1 \\ k_1 + \dots + k_j \ge n}} \frac{\partial^j f}{\partial p_{k_1} \cdots \partial p_{k_j}} b_{j,n}(k_1, \dots, k_j) x_1^{k_1 + \dots + k_j - n},$$
$$b_{1,n}(k_1) = k_1(k_1 - 1) \cdots (k_1 - n + 1).$$

We prove (8) by induction on n. For n = 1 we have

$$\frac{\partial f}{\partial x_1} = \sum_{k \ge 1} \frac{\partial f}{\partial p_k} \frac{\partial p_k}{\partial x_1} = \sum_{k \ge 1} \frac{\partial f}{\partial p_k} k x_1^{k-1}.$$

Now suppose (8) is true for n. Note that

$$\frac{\partial}{\partial x_1} \left(\sum_{k \ge n} \frac{\partial f}{\partial p_k} k(k-1) \cdots (k-n+1) x_1^{k-n} \right) =$$

$$= \sum_{k \ge n+1} \frac{\partial f}{\partial p_k} k(k-1) \cdots (k-n) x_1^{k-n-1} +$$

$$+ \sum_{\substack{k_1, k_2 \ge 1\\k_1+k_2 \ge n+1}} \frac{\partial^2 f}{\partial p_{k_1} \partial p_{k_2}} k_2 k_1 (k_1-1) \cdots (k_1-n+1) x_1^{k_1+k_2-n-1}.$$

In addition

$$\frac{\partial}{\partial x_1} \left(\sum_{\substack{k_1 + \dots + k_j \ge n \\ k_1, \dots, k_j \ge 1}} \frac{\partial^j f}{\partial p_{k_1} \cdots \partial_{k_j}} b_{j,n}(k_1, \dots, k_j) x_1^{k_1 + \dots + k_j - n} \right) = \sum_{\substack{k_1 + \dots + k_j \ge n \\ k_1, \dots, k_j \ge 1}} \frac{\partial^j f}{\partial p_{k_1} \cdots \partial p_{k_j}} b_{j,n}(k_1, \dots, k_j) (k_1 + \dots + k_j - n) x_1^{k_1 + \dots + k_j - n - 1} + \sum_{\substack{k_1 + \dots + k_j \ge n \\ k_1, \dots, k_j \ge 1}} \frac{\partial^{j+1} f}{\partial p_{k_1} \cdots \partial p_{k_{j+1}}} k_{j+1} b_{j,n}(k_1, \dots, k_j) x_1^{k_1 + \dots + k_{j+1} - n - 1}.$$

Note that the sum in the last line ranges over k_1, \ldots, k_{j+1} with $k_1 + \ldots + k_{j+1} \ge n+1$. Thus (8) is checked. Now consider $L\left(\frac{\partial^n f}{\partial x_1^n}\Big|_{x_1=0}\right)$. Because of $L\left(p_i\Big|_{x_i=0}\right) = p_i$ we obtain

(9)
$$T_n(f) = L\left(\frac{\partial^n f}{\partial x_1^n}\Big|_{x_1=0}\right) =$$
$$= \frac{\partial f}{\partial p_n} n! + \sum_{j=2}^n \sum_{k_1+\dots+k_j=n} \frac{\partial^j f}{\partial p_{k_1}\cdots\partial p_{k_j}} b_{j,n}(k_1,\dots,k_j)$$

Now we can prove by induction on n that

(10)
$$\mathbb{C}\left[\frac{\partial}{\partial p_1},\ldots,\frac{\partial}{\partial p_n}\right] = \mathbb{C}[T_1,\ldots,T_n].$$

For n = 1 we have $T_1 f = L\left(\frac{\partial f}{\partial x_1}\Big|_{x_1=0}\right) = \frac{\partial f}{\partial p_1}$. In addition if we know (10) for n, (9) gives us the statement for n + 1. Proposition is proved.

Remark 2.1. For any $f \in \Lambda$

$$f(x_1, x_2, x_3, \ldots) = f(x_2, x_3, \ldots) + \sum_{k=1}^{\infty} \frac{1}{k!} x_1^k(T_k f)(x_2, x_3, \ldots).$$

Recall the definition of skew Jack polynomials $J_{\lambda/\mu}(\mathbf{x})$. (From now on we suppress the coupling constant m in the notation for Jack and skew Jack polynomials.) Define $f_{\mu\nu}^{\lambda}$ as follows: $J_{\mu}(\mathbf{x})J_{\nu}(\mathbf{x})\langle J_{\mu}, J_{\mu}\rangle^{-1}\langle J_{\nu}, J_{\nu}\rangle^{-1} = \sum_{\lambda}\langle J_{\lambda}, J_{\lambda}\rangle^{-1}f_{\mu\nu}^{\lambda}J_{\lambda}(\mathbf{x}).$ Here λ, μ, ν are some partitions. Then

$$J_{\lambda/\mu}(\mathbf{x}) = \sum_{\nu} f_{\mu\nu}^{\lambda} J_{\nu}(\mathbf{x}), \quad J_{\lambda/\mu}(x_1) = \sum_{\nu} f_{\mu\nu}^{\lambda} J_{\nu}(x_1)$$

(surely the summation in the second sum ranges over ν with $l(\nu) \leq 1$). For two partitions μ, λ we write $\mu \subset \lambda$ if the Young diagram of μ is a subset of the Young diagram of λ . Note that $J_{\lambda/\mu}(\mathbf{x}) = 0$ unless $\mu \subset \lambda$.

Following proposition is proved in [Mac].

Proposition 2.2.

a) $J_{\lambda}(x_1, x_2, \ldots) = \sum_{\mu} J_{\lambda/\mu}(x_1) J_{\mu}(x_2, x_3, \ldots).$ b) Suppose that $\mu \subset \lambda$. Then $J_{\lambda/\mu}(x_1) = 0$ unless there exists such i that $\lambda_j = \mu_j$ for $j \neq i$. In the latter case $J_{\lambda/\mu}(x_1) = \phi_{\lambda\mu} x_1^{\lambda_i - \mu_i}$, $\phi_{\lambda\mu}$ is some nonzero constant.

Remark 2.2. Note that $\phi_{\lambda,\mu}$ depend on the coupling constant α and do not vanish for α being natural number.

Corollary 2.1. $T_k(J_\lambda(\mathbf{x})) = k! \sum_{\mu} \phi_{\lambda\mu} J_\mu(\mathbf{x})$, where the sum ranges over such partitions μ that there exists i with $\lambda_i - \mu_i = k$, $\lambda_j = \mu_j$ if $j \neq i$.

Proposition 2.3. Let $\lambda = (s^r) = (\overbrace{s, s, \dots, s}^r)$. Then $\mathbb{C}\left[\frac{\partial}{\partial p_1}, \frac{\partial}{\partial p_2}, \dots\right] \cdot J_{\lambda}(\mathbf{x})$ is a linear span of $J_{\mu}(\mathbf{x})$ with $\mu \subset \lambda$.

Proof. Because of the corollary (2.1) and proposition (2.2) we only need to prove that $J_{\mu}(\mathbf{x}) \hookrightarrow \mathbb{C}[T_1, T_2, \ldots] \cdot J_{\lambda}(\mathbf{x})$ for any $\mu \subset \lambda$. Let $\mu = (\mu_1 \ge \ldots \ge \mu_r), \mu_1 \le s$. We use the decreasing induction on such j that $\mu_j < s$ but $\mu_{j-1} = s$.

10

In the case j - 1 = r we have $\mu = \lambda$. Now let $j \leq r$. Let $\tilde{\mu}$ be partition with

$$\tilde{\mu}_j = s, \quad \tilde{\mu}_i = \mu_i \text{ for } i \neq j.$$

Then by induction assumption $J_{\tilde{\mu}}(\mathbf{x}) \in \mathbb{C}[T_1, T_2, \ldots] \cdot J_{\lambda}(\mathbf{x})$. Apply $T_{s-\mu_j}$ to $J_{\tilde{\mu}}(\mathbf{x})$.

$$\frac{1}{(s-\mu_j)!}T_{s-\mu_j}J_{\tilde{\mu}}(\mathbf{x}) = \phi_{\tilde{\mu}\mu}J_{\mu}(\mathbf{x}) + \sum_{\nu}\phi_{\tilde{\mu}\nu}J_{\nu}(\mathbf{x}),$$

where the sum in the right hand side ranges over such ν that there exists i > j with

 $\mu_i - \nu_i = s - \mu_j, \quad \nu_\alpha = \tilde{\mu}_\alpha \text{ for } \alpha \neq i.$

But for any such ν we have $\nu_j = s$ and thus $J_{\nu}(\mathbf{x}) \in \mathbb{C}[T_1, T_2, \ldots] \cdot J_{\lambda}(\mathbf{x})$. This gives us $J_{\mu}(\mathbf{x}) \in \mathbb{C}[T_1, T_2, \ldots] \cdot J_{\lambda}(\mathbf{x})$ (because $\phi_{\tilde{\mu}\mu} \neq 0$). Proposition is proved. \Box

Recall the space $V_{(m)}^{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}(p).$

Theorem 2.2. $V_{(m)}^{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}(p)$ has the basis of the form

$$\widehat{J}_{\mu}(\mathbf{x}), \quad \mu \subset ((p-(k-1)m-1)^k).$$

Proof. By lemma (2.4) we have

$$V_{(m)}^{\frac{-p}{\sqrt{2m}} + k\sqrt{2m}}(p) = \mathbb{C}[h_1, h_2, \ldots] \cdot a_0^k v_p.$$

Recall that if we identify some H_{μ} with the algebra of symmetric polynomials, operators h_i , i > 0 will act (up to a nonzero constant) as $\frac{\partial}{\partial p_i}$. Thus our theorem is a consequence of the theorem (2.1) and proposition (2.3).

3. DIMENSIONS AND CHARACTERS

In this section we find the dimensions and characters of $V_{(m)}(p)$ and of the space of coinvariants.

For partition μ denote $\deg_q \mu = q^{\sum \mu_i}$.

Lemma 3.1. Let
$$\operatorname{ch}_{s,r}(q) = \sum_{\mu \subset (s^r)} \deg_q \mu$$
. Then $\operatorname{ch}_{s,r}(q) = {\binom{s+r}{r}}_q$.
Proof. First note that $\operatorname{ch}_{s,r}(q) = 1 + q + \dots + q^s = {\binom{s+1}{r}}_q$. In addition

Proof. First note that $\operatorname{ch}_{s,1}(q) = 1 + q + \ldots + q^s = \binom{s+1}{1}_q$. In addition

$$ch_{s,r}(q) = \sum_{s \ge \mu_1 \ge \dots \ge \mu_r \ge 0} q^{\sum_{i=1}^r \mu_i} = \sum_{j=0}^r \sum_{\substack{\mu_r = j \\ s \ge \mu_1 \ge \dots \ge \mu_r}} q^{\sum_{i=1}^r \mu_i} = \sum_{j=0}^r q^{rj} ch_{s-j,r-1}.$$

To finish the proof it is enough to note that

$$\binom{s+r}{r}_q = \sum_{j=0}^s q^{jr} \binom{s+r-j-1}{r-1}_q.$$

This finishes the proof of the lemma.

Proposition 3.1.

- a) $\operatorname{ch}(z,q,V_{(m)}(p)) = z^{\deg_z v_p} \sum_{k \ge 0} z^{k\sqrt{2m}} q^{\deg_q v_{p-2km}} {\binom{p-(m-1)(k-1)}{k}}_q.$
- b) Let $F_n^{(m)}$ be natural numbers defined by

$$F_i^{(m)} = i + 1, \ i = 0, 1, \dots, m - 1; \ F_{i+m}^{(m)} = F_{i+m-1}^{(m)} + F_i^{(m)}$$

Then dim $V_{(m),p}(n) = F_n^{(m)}$.

$$\square$$

Proof. Consider the element $\widehat{J}_{\lambda}(\mathbf{x}) \in H_{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}$. Note that

$$L_0\widehat{J}_{\lambda}(\mathbf{x}) = (\deg_q v_{p-2km} + \sum_i \lambda_i)\widehat{J}_{\lambda}(\mathbf{x}).$$

Thus theorem (2.2) and lemma (3.1) give us point a). Note that b) follows from a), because

$$\dim V_{(m)}(p+m) = \sum_{k \ge 0} \binom{p+m-(m-1)(k-1)}{k} = \sum_{k \ge 0} \binom{p+m-1-(m-1)(k-1)}{k} + \sum_{k \ge 0} \binom{p+m-1-(m-1)(k-1)}{k-1} = \dim V_{(m)}(p+m-1) + \sum_{k \ge 0} \binom{p-(m-1)(k-1)}{k} = \dim V_{(m)}(p+m-1) + \dim V_{(m)}(p).$$

Proposition is proved.

Remark 3.1. There is an obvious embedding $V_{(m),p}(p-1) \hookrightarrow V_{(m)}(p)$. Consider the quotient $V_{(m)}(p)/V_{(m),p}(p-1)$. Natural way to prove the formula for the dimension of $V_{(m)}(p)$ is to identify above quotient with the space $V_{(m)}(p-m)$. This was done in [FF1] for m = 1, but the straightforward generalization failed for m > 1.

In the following proposition we find the dimension and character of the space of coinvariants for $V_{(m),m-1}$. We denote $V_{(m),m-1}$ simply by $V_{(m)}$.

Proposition 3.2. Define $C_{(m)}(n) = V_{(m)}/span\{a_iV_{(m)}, i \leq -m-n\}.$ a) dim $C_{(m)}(n) = F_n^{(2m)}.$

b) Note that $C_{(m)}(n)$ inherit bigrading from $V_{(m)}$. Thus its character is well-defined. We have

$$\operatorname{ch}^{\prime}(z,q,C_{(m)}(n)) = z^{\deg_{z} v_{1-m}} q^{\deg_{q} v_{1-m}} \sum_{k \ge 0} z^{k\sqrt{2m}} q^{mk^{2}} \binom{n - (2m - 1)(k - 1)}{k}_{q}.$$

Proof. a) follows from the statement (see corollary (1.1)) that the images of vectors

$$a_{-i_1} \cdots a_{-i_k} v_{1-m}, \ i_{\alpha+1} - i_{\alpha} \ge 2m, \ i_1 \ge m, \ i_k \le n+m-1; \ k=0,1,\dots$$

form a basis of $C_{(m)}(n)$.

To prove b) note that it follows from the corollary (1.1) that for $n \ge 2m$

$$\begin{split} \mathrm{ch}(z,q,C_{(m)}(n)) &= \mathrm{ch}(z,q,C_{(m)}(n-1)) + \\ & z^{\deg_z v_{1-m}} q^{\deg_q v_{1-m}} z^{\sqrt{2m}} q^{m+n-1} \mathrm{ch}(z,q,C_{(m)}(n-2m)). \end{split}$$

It is straightforward to check that this relation holds for the right hand side of (11).

4. Semi-infinite bases

In this section we construct monomial bases for the spaces $L_{(m),i}$.

Lemma 4.1. Recall algebra $A_{(m)} = \mathbb{C}[\xi_0, \xi_1, \ldots]/\langle (\xi(z)^{(i)})^2 \rangle_{0 \le i \le m-1}$. Consider the subspace $A^1_{(m)} \hookrightarrow A_{(m)}, A^1_{(m)} = \xi_0 A_{(m)}$. Then the map

$$\rho: A_{(m)} \to A^1_{(m)}, \quad \xi_{i_1} \cdots \xi_{i_k} \mapsto \xi_0 \xi_{i_1+2m} \cdots \xi_{i_k+2m}$$

is well-defined isomorphism.

Proof. Recall the fermionic realization $A_{(m)} \hookrightarrow F$ from section 1. Note that

$$\xi_0 = \psi_1(0) \cdots \psi_{2m}(0).$$

Thus we obtain that $\xi_0 \xi_i = 0$ for $i = 0, \dots, 2m - 1$ and ρ is an isomorphism. \Box

Lemma 4.2. Consider a map $V_{(m),p} \to A_{(m)}$:

$$a_{j_1}\cdots a_{j_s}v_p\mapsto \xi_{-j_1+p-1}\cdots \xi_{-j_s+p-1}.$$

Then this map is well-defined isomorphism.

Proof. Follows from lemmas (2.1) and (2.2).

Corollary 4.1. $L_{(m),i}$ has a basis labeled by the sequences $(s_j)_{j\in\mathbb{Z}}, s_j \in \mathbb{N} \cup \{0\}$ with the following properties:

- (1) There exists such $n \in \mathbb{Z}$ that $s_{-i+2mk-1} = 1$ for $k \ge n$ and $s_j = 0$ if $(j \ge -i + 2mn 1 \text{ and } j + i + 1 \text{ is not divisible on } 2m)$.
- (2) There exists such $N \in \mathbb{Z}$ that $s_j = 0$ for j < N.
- (3) For any $j \in \mathbb{Z}$ $s_j + \ldots + s_{j+2m-1} \leq 1$.

Proof. Rescale vectors $v_p = \left|\frac{-p}{\sqrt{2m}}\right\rangle$ to satisfy equations $a_{p-1}v_p = v_{p-2m}$. Attach to each sequence $(s_j)_{j\in\mathbb{Z}}$, satisfying properties (1), (2), (3), the following vector (n comes from the property (1)):

(12)
$$\left(\prod_{j<-i+2mn} a_j^{s_j}\right) v_{-i+2nm} \in L_{(m),i}.$$

Then because of (1) for any $n_1 > n$

$$\left(\prod_{j<-i+2n_1m} a_j^{s_j}\right) v_{-i+2n_1m} = \left(\prod_{j<-i+2nm} a_j^{s_j}\right) v_{-i+2nm}$$

Note that $L_{(m),i}$ is the limit $\lim_{n\to\infty} V_{(m),-i+2nm}$:

(13)
$$L_{(m),i} = \ldots \hookrightarrow V_{(m),-i} \hookrightarrow V_{(m),-i+2m} \hookrightarrow V_{(m),-i+4m} \hookrightarrow \ldots$$

and each of $V_{(m),i}$ is isomorphic to $A_{(m)}$ in the sense of lemma (4.2). In addition embeddings (13) are compatible with the isomorphism from the lemma (4.1). Thus corollary (1.1) gives us that vectors (12) form basis of $L_{(m),i}$.

5. Discussion

5.1. Fusion procedure. Recall algebra $A_{(m)}$. Let $A_{(m)}(n) \hookrightarrow A_{(m)}$ be its finitization: subalgebra, generated by ξ_0, \ldots, ξ_{n-1} . Then

$$A_{(m)}(n) \simeq \mathbb{C}[\xi_0, \dots, \xi_{n-1}]/I_{(m)}(n),$$

where $I_{(m)}(n)$ is some ideal in $\mathbb{C}[\xi_0, \ldots, \xi_{n-1}]$. Now let $Z = (z_1, \ldots, z_n) \in \mathbb{C}^n$, $z_i \neq z_j$ for $i \neq j$. Let $I_{(m),Z}(n) \hookrightarrow \mathbb{C}[\xi_0, \ldots, \xi_{n-1}]$ be the ideal, generated by the following elements:

 $(\xi_0 + \xi_1 z_i + \ldots + \xi_{n-1} z_i^{n-1})(\xi_0 + \xi_1 z_j + \ldots + \xi_{n-1} z_j^{n-1}), \ 1 \le i \le j \le n, \ |i-j| < m.$

It was proved in [FF1] that if $Z(\varepsilon) = (z_1(\varepsilon), \ldots, z_n(\varepsilon)) \in \mathbb{C}^n$ is a family of points of \mathbb{C}^n with the properties $z_i(\varepsilon) \neq z_j(\varepsilon)$ $(i \neq j)$ and $\lim_{\varepsilon \to 0} z_i(\varepsilon) = 0$ $(1 \le i \le n)$, then $\lim_{\varepsilon \to 0} I_{(1),Z(\varepsilon)}(n) = I_{(1)}(n)$. Following conjecture generalizes this limit (fusion) procedure to the case of general m.

Conjecture 5.1. Let $Z(\varepsilon) = (z_1(\varepsilon), \ldots, z_n(\varepsilon)) \in \mathbb{C}^n$ be a family with the following properties:

(1)
$$z_i(\varepsilon) \neq z_j(\varepsilon)$$
 for $i \neq j$ and $\lim_{\varepsilon \to 0} z_i(\varepsilon) = 0, \ 1 \le i \le n.$
(2)
$$\lim_{\varepsilon \to 0} \frac{z_2(\varepsilon) - z_1(\varepsilon)}{z_1(\varepsilon)} = 0, \quad \lim_{\varepsilon \to 0} \frac{z_{i+1}(\varepsilon) - z_i(\varepsilon)}{z_i(\varepsilon) - z_{i-1}(\varepsilon)} = 0, \ i = 2, \dots, n-1.$$

Then $\lim_{\varepsilon \to 0} I_{(m),Z(\varepsilon)}(n) = I_{(m)}(n).$

5.2. Finite-dimensional approximation. Consider the following version of the above fusion procedure. Let $R_{(m)}(n)$ be the following ring:

$$R_{(m)}(n) = \mathbb{C}[y_1, \dots, y_n] / \langle y_i y_j \rangle_{|i-j| < m}.$$

For $Z = (z_1, \ldots, z_n) \in \mathbb{C}^n$ with pairwise distinct z_i consider the set of generators of the latter ring:

$$a_i = \sum_{j=1}^n z_j^i y_j, \quad i = 0, \dots, n-1$$

Then for some ideal $\tilde{I}_{(m),Z}(n) \hookrightarrow \mathbb{C}[a_0,\ldots,a_{n-1}]$ we have

$$R_{(m)}(n) \simeq \mathbb{C}[a_0, \dots, a_{n-1}]/\tilde{I}_{(m),Z}(n).$$

Denote by $\tilde{I}_{(m)}(n)$ the limit $\lim_{Z\to 0} \tilde{I}_{(m),Z}(n)$, where Z goes to 0 with the restrictions from the conjecture 5.1.

Lemma 5.1. The homomorphism of algebras

(14)
$$\mathbb{C}[\xi_0, \dots, \xi_{n-1}]/I_{(m)}(n) \to \mathbb{C}[a_0, \dots, a_{n-1}]/\tilde{I}_{(m)}(n),$$

sending ξ_i to a_{n-1-i} , is an isomorphism.

Denote the right hand side of (14) by $W_{(m)}(n)$. Above lemma and conjecture (5.1) allows us to identify $W_{(m)}(n)$ with $V_{(m)}(n)$. Thus we obtain that $L_{(m),i}$ can be constructed as limit of the finite-dimensional spaces:

$$L_{(m),i} = W_{(m)}(2m-i) \hookrightarrow W_{(m)}(4m-i) \hookrightarrow \dots$$

as in the case m = 1 (see [FF2]).

Thus starting from the finite-dimensional spaces, obtained by the fusion procedure, one can construct the infinite-dimensional space as an inductive limit and endow this limit with the structure of the representation of the lattice vertex algebra. 5.3. Reconstruction of Jack polynomials from the fusion product $V_{(m)}(p)$. Let $v \in V_{(m)}^{\frac{-p}{\sqrt{2m}}+k\sqrt{2m}}(p)$ be a basis vector of the form $\widehat{J}_{\mu}(m;\mathbf{x})$ for some μ . Note that Jack polynomial $J_{\mu}(m;\mathbf{x})$ can be reconstructed from v using the action of the annihilating Heisenberg operators.

Namely, let $d = \deg_q v - \deg_q v_{p-2mk}$. Pick some $i_1, \ldots, i_r > 0$ with $i_1 + \cdots + i_r = d$. Then

(15)
$$h_{i_1} \cdots h_{i_k} v = c(i_1, \dots, i_k) v_{p-2mk}.$$

Recall that after the identification of any H_s with the space of symmetric polynomials operators h_i with i > 0 act as $\sqrt{\frac{2}{m}}i\frac{\partial}{\partial p_i}$. Thus (15) gives us

$$\frac{\partial}{\partial p_{i_1}}\cdots\frac{\partial}{\partial p_{i_k}}J_{\mu}(m;\mathbf{x}) = \left(\frac{m}{2}\right)^{k/2}\frac{c(i_1,\ldots,i_k)}{i_1\cdots i_k}$$

So the numbers $c(i_1, \ldots, i_k)$ allow us to reconstruct the corresponding Jack polynomial. (Note that in the case m = 1, where everything is proved for the fusion products, Jack polynomials reduce to Schur polynomials).

5.4. The odd case. Consider the case of vertex operator $\phi_{\sqrt{2m-1}}(z)$. Let a_i be its Fourier components, $a_i a_j = -a_j a_i$. Let L be some irreducible representation of the corresponding vertex algebra, $v \in L$ some extremal vector. Take such number p that $a_{p-1}v \neq 0$ and $a_kv = 0$ for any $k \geq p$. Then one can consider the principle subspace $U = \bigwedge(a_{p-1}, a_{p-2}, \ldots) \cdot v$ and its finitization $U(n) = \bigwedge(a_{p-1}, \ldots, a_{p-n}) \cdot v$. The study of U can be provided by the same means as in the even case. At the same time one needs some other tools to understand the structure of U(n). We have a conjecture concerning its dimension.

Conjecture 5.2. dim $U(n) = F_n^{(m)}$.

References

- [FL] B.Feigin, S.Loktev, On generalized Kostka polynomials and quantum Verlinde rule, Differential topology, infinite-dimensional Lie algebras and applications, Amer. Math. Soc. Transl. Ser. 2, 194, 61-79.
- [FF1] B.Feigin, E.Feigin, Q-characters of the tensor products in sl₂ case, Mosc. Math. J. 2 (2002), no. 3, 567-588.
- [FF2] B.Feigin, E.Feigin, Integrable siz modules as an infinite tensor products. In: S.Lando and O.Sheinman (eds.), Fundamental mathematics today (in honor of the 10th anniversary of the Independent University of Moscow), IUM, MCCME 2003 (Russian). Preprint version available at http://xxx.lanl.gov/abs/math.QA/0205281.
- [FS] B.Feigin, A.Stoyanovski, Quasi-particles models for the representations of Lie algebras and geometry of flag manifold, available at http://xxx.lanl.gov/abs/hep-th/9308079, RIMS 942;
- Functional models of the representations of current algebras, and the semi-infinite Schubert cells, Functsional. Anal. i Prilozhen. **28** (1994), no.1, 68-90, 96 (Russian). English translation in Funct. Anal. Appl. **28** (1994), no.1, 55-72.
- [FJM] B.Feigin, M.Jimbo and T.Miwa, Vertex operator algebra arising from the minimal series M(3,p) and monomial basis, MathPhys Odyssey 2001: integrable models and beyond, In honor of Barry M.McCoy, M.Kashiwara and T.Miwa eds., Birkhauser.
- [CP] V.Chari, A.Pressley, Weyl modules for classical and quantum affine algebras, Represent. Theory 5 (2001), 191-223 (electronic).

Preprint version available at http://xxx.lanl.gov/abs/math.QA/0004174.

[AMOS] H.Awata, Y.Matsuo, S.Odake, J.Shiraishi, Excited states of Calogero-Sutherland Model and Singular vectors of the W_N algebra, Nuclear Phys. B, 449 (1995), no. 4, 665-674. Preprint version available at http://xxx.lanl.gov/abs/hep-th/9503043

B.FEIGIN AND E.FEIGIN

[TY] A.Tsuchiya, Y.Kanie, Fock space representations of the Virasoro algebra – Interwining operators. Publ. RIMS Kyoto Univ. 22, 259-327 (1986).

[MY] K.Mimachi, Y.Yamada, Singular vectors of the Virasoro algebra in terms of Jack symmetric polynomials, Comm. Math. Phys., 174 (1995), no. 2, 447-455.

[Mac] I.G.Macdonald, Symmetric functions and Hall polynomials, 2nd ed., Oxford University Press, New York, 1995.

[D] C.Dong, Vertex algebras associated with even lattices, J.Algebra 161 (1993), 245-265.

BORIS FEIGIN: Russia, Chernogolovka 142432, Landau Institute for Theoretical Physics AND Russia, Moscow, Independent University of Moscow, Bol'shoi Vlas'evskii, 7 E-mail address: feigin@mccme.ru

EVGENY FEIGIN: Russia, Moscow, Independent University of Moscow, Bol'shoi Vlas'evsky, 7 AND Russia, Moscow, Moscow State University, Mech-Math Faculty, Department of Higher Algebra, Leninskie gori, 1

E-mail address: evgfeig@mccme.ru