

Smooth Lie groups over local fields of positive characteristic need not be analytic

Helge Glöckner

Abstract

We describe finite-dimensional smooth Lie groups over local fields of positive characteristic which do not admit an analytic Lie group structure compatible with the given topological group structure, and C^n -Lie groups without a compatible C^{n+1} -Lie group structure, for each $n \geq 1$. We also present examples of non-analytic, smooth automorphisms, as well as C^n -automorphisms which fail to be C^{n+1} .

Subject classification: 22E20 (main); 20G25, 22D05, 22D45, 26E30.

Key words: smooth Lie group, analytic Lie group, totally disconnected group, finite order differentiability, local field, positive characteristic, contractive automorphism, scale function, tidy subgroup, Willis theory, non-archimedean analysis

Introduction

It is a classical fact (the core of which goes back to Schur, [14]) that every finite-dimensional real Lie group of class C^k (where $k \in \mathbb{N} \cup \{\infty\}$) admits a C^k -compatible analytic Lie group structure; see [12, §4.4] and the references given there for further information. Based on Lazard's characterization of analytic p -adic Lie groups within the class of locally compact groups [10], it was shown in [6] that also every finite-dimensional C^k -Lie group over a local field of characteristic 0 admits a C^k -compatible \mathbb{K} -analytic Lie group structure. Here, C^k -maps and Lie groups are understood in the sense of [2], where a setting of differential calculus over arbitrary topological fields was developed. A mapping between finite-dimensional vector spaces over a local field is C^1 if and only if it is strictly differentiable at each point [7, Rem. 5.4], where strict differentiability is a classical concept, [3, 1.2.2].

In the present article, we show by example that the situation is entirely different in the case of positive characteristic. We prove:

Main Theorem. *Let \mathbb{K} be a local field of positive characteristic. Then:*

- (a) *There exists a one-dimensional smooth \mathbb{K} -Lie group which does not admit a \mathbb{K} -analytic Lie group structure compatible with its topological group structure.*
- (b) *For each positive integer n , there exists a one-dimensional \mathbb{K} -Lie group of class C^n which does not admit a \mathbb{K} -Lie group structure of class C^{n+1} compatible with the given topological group structure.*

(See Theorem 5.2 and Theorem 6.1). We also describe a C^n -automorphism of a finite-dimensional \mathbb{K} -analytic Lie group which is not C^{n+1} (Proposition 3.4), and a non-analytic C^∞ -automorphism (Proposition 3.7).

As a tool, we use concepts from George Willis' structure theory of totally disconnected, locally compact groups as developed in [18], [19] and analyzed for Lie groups over local fields in [5] and [9].

While it is quite justified to restrict attention to analytic Lie groups in the real and p -adic cases, our results make it clear that this habit is inappropriate in the case of positive characteristic. Because genuinely larger classes of groups can be captured in this way, it may very well be of interest to study Lie groups over such fields which are merely smooth, or merely C^n . It is not hopeless to prove substantial results in the absence of analyticity. For example, it is possible to discuss contraction groups (opposite subgroups), tidy subgroups and the scale function for automorphisms of mere C^1 -Lie groups, [9].

1 Preliminaries and Notation

We recall some definitions and basic facts from differential calculus (and linear algebra) over local fields, and set up our notational conventions.

1.1 Let E and F be (Hausdorff) topological vector spaces over a non-discrete topological field \mathbb{K} , $U \subseteq E$ be open, and $f: U \rightarrow F$ be a map. Following [2], we call f a C^1 -map if it is continuous and there exists a (necessarily unique) continuous map $f^{[1]}: U^{[1]} \rightarrow F$ on $U^{[1]} := \{(x, y, t) \in U \times E \times \mathbb{K} : x + ty \in U\}$ such that $f^{[1]}(x, y, t) = \frac{1}{t}(f(x + ty) - f(x))$ for all $(x, y, t) \in U^{[1]}$ with $t \neq 0$. Then $f'(x): E \rightarrow F$, $y \mapsto f'(x).y := f^{[1]}(x, y, 0)$ is a continuous linear map, for each $x \in U$ [2, Prop. 2.2]. Inductively, f is called C^{n+1} if it is C^1 and $f^{[1]}$ is C^n ; it is C^∞ (or *smooth*) if it is C^n for all n .

1.2 For maps between open subsets of finite-dimensional real vector spaces, the present definitions of C^n -maps are equivalent to the usual ones (cf. [2, Prop. 7.4 and 7.7]).

1.3 Compositions of composable C^r -maps are C^r [2, Prop. 4.5], the Chain Rule holds [2, Prop. 3.1], and being C^r is a local property, for each $r \in \mathbb{N} \cup \{\infty\}$ [2, La. 4.9]. This makes it possible to define C^r -manifolds modeled on a topological \mathbb{K} -vector space E , in the usual way. A \mathbb{K} -Lie group of class C^r is a group, equipped with a C^r -manifold structure modeled on a topological \mathbb{K} -vector space, which makes the group operations C^r . If G is a C^r -Lie group, we write $L(G) := T_1(G)$ for its (geometric) tangent space at the identity.¹ Given a homomorphism $\alpha : G \rightarrow H$ which is at least C^1 , as usual we abbreviate $L(\alpha) := T_1(\alpha)$.

1.4 For the definition of \mathbb{K} -analytic maps between open subsets of Banach spaces over a complete valued field \mathbb{K} , we refer to [3] (see also [15] for the finite-dimensional case). Any \mathbb{K} -analytic map is C^∞ by [2, Prop. 7.20]. Hence every \mathbb{K} -analytic Lie group modeled on a Banach space can also be considered as a C^∞ -Lie group in the sense of **1.3**.

For functions of a single variable, an equivalent description of the C^n -property is available, which is much easier to work with. This is essential for our constructions of pathological homomorphisms and Lie groups, which necessitate to verify by hand that certain mappings defined on open subsets of the ground field are C^n . It is also clear from the equivalent description that a map between open subsets of an ultrametric field \mathbb{K} is C^n in our sense if and only if it is so in the sense of [13].

Lemma 1.5 (cf. [2, Prop. 6.9]). *Let $U \subseteq \mathbb{K}$ be an open subset of a topological field \mathbb{K} and F be a topological \mathbb{K} -vector space. Given a map $f : U \rightarrow F$, we set $f^{<0>} := f$. Then the following holds:*

- (a) *The map f is C^1 if and only if it is continuous and there exists a (necessarily unique) continuous map $f^{<1>} : U \times U \rightarrow F$ such that*

$$f^{<1>}(x_1, x_2) = \frac{1}{x_1 - x_2}(f(x_1) - f(x_2))$$

for all $x_1, x_2 \in U$ such that $x_1 \neq x_2$.

¹If $r \geq 3$, then $L(G)$ has a natural Lie algebra structure, but we shall not use it.

- (b) Having characterized C^k -maps and defined $f^{<k>} : U^{k+1} \rightarrow F$ for all $k = 0, \dots, n-1$, where $n \in \mathbb{N}$, we have: f is C^n if and only if f is C^{n-1} and there exists a continuous map $f^{<n>} : U^{n+1} \rightarrow F$ such that

$$\begin{aligned} & f^{<n>}(x_1, x_2, \dots, x_{n+1}) \\ &= \frac{1}{x_1 - x_2} \left(f^{<n-1>}(x_1, x_3, \dots, x_{n+1}) - f^{<n-1>}(x_2, x_3, \dots, x_{n+1}) \right) \end{aligned}$$

for all $x_1, \dots, x_{n+1} \in U$ such that $x_1 \neq x_2$.

Here $f^{<n>}$ is uniquely determined and is symmetric in its $n+1$ arguments. Furthermore, $n! D^n f(x) = \frac{d^n f}{dx^n}(x) =: f^{(n)}(x)$ for all $x \in U$, where $D^n f(x) := f^{<n>}(x, \dots, x)$. \square

For further information on differential calculus over topological fields and the corresponding manifolds and Lie groups, we refer to [2] and [6]–[8].

1.6 Recall that a *local field* is a totally disconnected, locally compact, non-discrete topological field. On any local field, there exists a *natural absolute value* $|\cdot| : \mathbb{K} \rightarrow [0, \infty[$, defined via $|0| := 0$ and $|x| := \Delta_{\mathbb{K}}(\lambda_x)$ for $x \neq 0$, where $\lambda_x : \mathbb{K} \rightarrow \mathbb{K}$, $\lambda_x(y) := xy$ is multiplication by x and $\Delta_{\mathbb{K}} : \text{Aut}(\mathbb{K}, +) \rightarrow]0, \infty[$ the modular function. Then $|\mathbb{K}^\times| = \langle q \rangle \leq (]0, \infty[, \cdot)$ for some prime power q , called the *module of \mathbb{K}* . For further information on local fields, see [17].

The following lemma provides information concerning a special type of linear maps which will play a crucial role in our constructions.

Lemma 1.7 *Let V be a finite-dimensional vector space over a local field \mathbb{K} with module q . Let $|\cdot| : \overline{\mathbb{K}} \rightarrow [0, \infty[$ be the unique extension of the natural absolute value on \mathbb{K} to an absolute value on the algebraic closure $\overline{\mathbb{K}}$ of \mathbb{K} . Let $\alpha \in \text{GL}(V)$ be a \mathbb{K} -linear automorphism of V , with eigenvalues $\lambda_1, \dots, \lambda_d$ in $\overline{\mathbb{K}}$, where $d := \dim_{\mathbb{K}}(V)$. Suppose that $|\lambda_j| > 1$ for each $j \in \{1, \dots, d\}$ and $\prod_{j=1}^d |\lambda_j| = q$. Consider V as a $\mathbb{K}[T]$ -module via $(\sum_{j=0}^n a_j T^j) \cdot v := \sum_{j=0}^n a_j A^j v$. Then*

- (a) V is an irreducible $\mathbb{K}[T]$ -module;
- (b) $|\lambda_j| = q^{1/d}$ for each $j \in \{1, \dots, d\}$; and
- (c) There exists an ultrametric norm $\|\cdot\|$ on V such that $\|\alpha(x)\| = q^{1/d} \|x\|$ for each $x \in V$.

Proof. (a) Let $\{0\} \neq U \subseteq V$ be a $\mathbb{K}[T]$ -submodule of V ; we show that $U = V$. To this end, let $e := \dim_{\mathbb{K}}(U) \in \{1, \dots, d\}$ and note that $\lambda_{j_1}, \dots, \lambda_{j_e}$ are the eigenvalues of $\alpha|_U^U$ in $\overline{\mathbb{K}}$, for suitable integers $1 \leq j_1 < j_2 < \dots < j_e \leq d$. Set $J := \{j_k : k = 1, \dots, e\}$ and $r := \prod_{j \notin J} |\lambda_j|$. Then $r \geq 1$, with equality if and only if $J = \{1, \dots, d\}$ (and hence $U = V$). But

$$q = |\det(\alpha)| = \prod_{j=1}^d |\lambda_j| = r \prod_{j \in J} |\lambda_j|,$$

where $1 < \prod_{j \in J} |\lambda_j| = |\det(\alpha|_U^U)| \in q^{\mathbb{Z}}$ and thus $\prod_{j \in J} |\lambda_j| \geq q$. Hence $\prod_{j \in J} |\lambda_j| = q$ and $r = 1$, entailing that $U = V$.

(b) For each j , let $(V_{\overline{\mathbb{K}}})_{\lambda_j} := \{v \in V_{\overline{\mathbb{K}}} : (\alpha_{\overline{\mathbb{K}}} - \lambda_j)^d v = 0\}$ be the generalized eigenspace of $\alpha_{\overline{\mathbb{K}}}$ in $V_{\overline{\mathbb{K}}}$ corresponding to λ_j , where $V_{\overline{\mathbb{K}}}$ is the $\overline{\mathbb{K}}$ -vector space obtained from V by extension of scalars. Let $R := \{|\lambda_j| : j \in \{1, \dots, d\}\}$; given $\rho \in R$ define $J_\rho := \{j \in \{1, \dots, d\} : |\lambda_j| = \rho\}$ and $W_\rho := \bigoplus_{j \in J_\rho} (V_{\overline{\mathbb{K}}})_{\lambda_j}$. By [11, Ch. II, (1.0), p.81], each W_ρ is defined over \mathbb{K} , i.e., $W_\rho = (V_\rho)_{\overline{\mathbb{K}}}$. Note that $V_\rho := W_\rho \cap V$ is α -invariant and hence a $\mathbb{K}[T]$ -submodule of V . Thus $V = \bigoplus_{\rho \in R} V_\rho$ is a direct sum of $\mathbb{K}[T]$ -submodules. By (a), such a decomposition is only possible if $R = \{\rho\}$ is a singleton and thus $V = V_\rho$. Now $q = |\lambda_1| \cdots |\lambda_d| = \rho^d$ entails $\rho = q^{1/d}$.

(c) We choose a $\overline{\mathbb{K}}$ -basis v_1, \dots, v_d of $V_{\overline{\mathbb{K}}}$ with respect to which $\alpha_{\overline{\mathbb{K}}}$ has Jordan canonical form, and let $\|\cdot\|$ be the maximum norm on $V_{\overline{\mathbb{K}}}$ with respect to this basis. Since $q^{1/d} > 1$ and $|1| = 1$, the ultrametric inequality entails that $\|\alpha_{\overline{\mathbb{K}}}(v_j)\| = q^{1/d} \|v_j\|$ for each $j \in \{1, \dots, d\}$ (cf. [5, p.209, Eqn. (1)]). Hence $\|\alpha_{\overline{\mathbb{K}}}(x)\| = q^{1/d} \|x\|$ for each $x \in V_{\overline{\mathbb{K}}}$, using that $\|\cdot\|$ is a maximum norm. Now the restriction of $\|\cdot\|$ to V is norm with the desired property. \square

1.8 Conventions. Throughout the article, q is a prime power, \mathbb{F}_q a finite field of cardinality q and $\mathbb{K} := \mathbb{F}_q((X))$ the field of formal Laurent series over \mathbb{F}_q , unless said otherwise.² We let $\mathbb{O} := \mathbb{F}_q[[X]]$ be the valuation ring of \mathbb{K} (the ring of formal power series over \mathbb{F}_q). We let $|\cdot| : \mathbb{K} \rightarrow [0, \infty[$ be the natural absolute value on \mathbb{K} , given by $|\sum_{k=n}^{\infty} a_k X^k| := q^{-n}$ for $a_n, a_{n+1}, \dots \in \mathbb{F}_q$ with $a_n \neq 0$. We fix an algebraic closure $\overline{\mathbb{K}}$ and use the same symbol, $|\cdot|$, for the unique extension of the absolute value $|\cdot|$ on \mathbb{K} to an absolute value on $\overline{\mathbb{K}}$. We use \mathbb{R} and \mathbb{Z} in their usual meanings and write $\mathbb{N} := \{1, 2, \dots\}$, $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

²Up to isomorphism, every local field of positive characteristic is of this form [17].

In the following, by a ‘‘Lie group’’ we always mean a Lie group modeled on a finite-dimensional vector space. A C^r -*automorphism* of a C^r -Lie group G is an automorphism $\alpha : G \rightarrow G$ such that both α and its inverse are C^r . The same convention applies to \mathbb{K} -analytic automorphisms, \mathbb{K} -analytic diffeomorphisms and C^r -diffeomorphisms. Given a topological group G , its group of bicontinuous automorphisms will be denoted by $\text{Aut}(G)$. If $(E, \|\cdot\|)$ is a normed \mathbb{K} -vector space, we set $B_r^E := \{x \in E : \|x\| < r\}$ for $r > 0$; if E is understood, we abbreviate $B_r := B_r^E$.

2 Criteria making a map C^n , or preventing it

In this section, we describe simple criteria ensuring that a mapping is C^n , resp., that it is not C^n .

Lemma 2.1 *Let $n \in \mathbb{N}$ and $f : U \rightarrow \mathbb{K}$ be a map, defined on an open subset $U \subseteq \mathbb{K}$. Suppose that there exist real numbers $\theta > n$ and $b > 0$ such that*

$$|f(y) - f(x)| \leq b|y - x|^\theta \quad \text{for all } x, y \in U. \quad (1)$$

Then f is C^n and $D^k f(x) = 0$ for all $k \in \{1, \dots, n\}$ and $x \in U$.

Proof. This is a special case of [13, Thm. 29.12]. □

Lemma 2.2 *Let $E \neq \{0\}$ and F be normed \mathbb{K} -vector spaces and $f : U \rightarrow F$ be a mapping on an open subset $U \subseteq E$. Let $n \in \mathbb{N}_0$. Suppose that there exist $x \in U$, $\theta \in]n, n + 1[$ and positive real numbers a, b such that*

$$a\|y - x\|^\theta \leq \|f(y) - f(x)\| \leq b\|y - x\|^\theta \quad (2)$$

for all $y \in U$. Then f is not of class C^{n+1} .

Proof. There exists $0 \neq y \in E$ such that $x + \mathbb{O}y \subseteq U$. Then $g : \mathbb{O} \rightarrow F$, $g(t) := f(x + ty) - f(x)$ satisfies

$$A|t|^\theta \leq \|g(t)\| \leq B|t|^\theta \quad \text{for all } t \in \mathbb{O}, \quad (3)$$

with $A := a\|y\|^\theta$, $B := b\|y\|^\theta$. It suffices to show that g is not C^{n+1} . To reach a contradiction, assume that g is C^{n+1} . By [2, Thm. 5.1], g admits a Taylor expansion of order $n + 1$ around 0:

$$g(t) = \sum_{j=0}^{n+1} a_j t^j + t^{n+1} \rho(t) \quad \text{for all } t \in \mathbb{O},$$

where $\rho: \mathbb{O} \rightarrow F$ is continuous and satisfies $\rho(0) = 0$. But such an expansion is incompatible with the asymptotic described in (3). To see this, note first that $a_k = 0$ for $k = 0, \dots, n$. Indeed, $g(0) = 0$ entails that $a_0 = 0$, and if we already know that $a_0 = \dots = a_{k-1} = 0$, letting $t \rightarrow 0$ in $t^{-k}g(t) = a_k + \sum_{j=k+1}^{n+1} t^{j-k}a_j + t^{n+1-k}\rho(t)$ we find that also $a_k = 0$ (since $t^{-k}g(t) \rightarrow 0$, by (3)). Hence $g(t) = (a_{n+1} + \rho(t))t^{n+1}$ and thus

$$A \leq \frac{\|g(t)\|}{|t|^\theta} = \|a_{n+1} + \rho(t)\| \cdot |t|^{n+1-\theta},$$

where the right hand side tends to 0 as $t \rightarrow 0$. This contradicts $A > 0$. \square

3 Automorphisms with specific properties

In this section, we discuss a family of automorphisms of the additive topological group $\mathbb{O} := \mathbb{F}_q[[X]]$, some of which have specific pathological properties.

3.1 Given a strictly ascending sequence $\lambda := (\ell_j)_{j \in \mathbb{N}_0}$ of positive integers $\ell_0 < \ell_1 < \ell_2 < \dots$, we define a mapping $\beta: \mathbb{O} \rightarrow \mathbb{O}$ via

$$\beta\left(\sum_{j=0}^{\infty} a_j X^j\right) := \sum_{j=0}^{\infty} a_j X^{\ell_j} \quad \text{for } (a_j)_{j \in \mathbb{N}_0} \in \mathbb{F}_q^{\mathbb{N}_0}.$$

We define $\alpha: \mathbb{O} \rightarrow \mathbb{O}$ via $\alpha(z) := z + \beta(z)$.

Lemma 3.2 *The map β from 3.1 is a continuous homomorphism, and α is an automorphism of the topological group $(\mathbb{O}, +)$.*

Proof. Clearly β and α are homomorphisms of groups. Furthermore, β is continuous, since $\mathbb{O} = \mathbb{F}_q^{\mathbb{N}_0}$ is equipped with the product topology and each coordinate of $\beta(z)$ is either constant or only depends on one coordinate of z .

To see that α is a bijection, let $z = \sum_{j=0}^{\infty} a_j X^j$ and $w \in \sum_{j=0}^{\infty} b_j X^j$ be elements of \mathbb{O} , where $a_j, b_j \in \mathbb{F}_q$. Equating coefficients, we see that $\alpha(z) = w$ holds if and only if $a_0 = b_0$ and

$$(\forall i \in \mathbb{N}) \quad a_i = \begin{cases} b_i & \text{if } i \neq \ell_j \text{ for each } j \in \mathbb{N}_0; \\ b_i - a_j & \text{if } i = \ell_j \text{ for some } j \in \mathbb{N}_0. \end{cases} \quad (4)$$

Since $j < i$ in the second case of (4), given w we can solve for a_0, a_1, \dots in turn and deduce that $z \in \mathbb{O}$ with $\alpha(z) = w$ exists and is uniquely determined. Thus α is an isomorphism of groups. It is clear from the preceding formula that, for fixed i , the i th coordinate a_i of $z = \alpha^{-1}(w)$ only depends on finitely many coordinates, b_0, b_1, \dots, b_i , of w . Hence α^{-1} is continuous. \square

Lemma 3.3 *If there exist real numbers $\theta > 0$ and $K > 0$ such that*

$$j\theta - K \leq \ell_j \leq j\theta + K \quad \text{for all } j \in \mathbb{N}_0, \quad (5)$$

then there exist positive real numbers $a < b$ such that

$$a|y - x|^\theta \leq |\beta(y) - \beta(x)| \leq b|y - x|^\theta, \quad \text{for all } x, y \in \mathbb{O}.$$

Proof. We claim that the assertion holds with $a := \min\{q^{-\ell_0}, q^{-K}\}$ and $b := q^K$. Since $\beta(y) - \beta(x) = \beta(y - x)$, we may assume without loss of generality that $x = 0$. We distinguish three cases.

1. If $y = 0$, then the assertion is trivial.
2. If $|y| = 1$, then $|\beta(y)| = q^{-\ell_0} \geq a = a|y|^\theta$ and $b|y|^\theta = b \geq 1 \geq q^{-\ell_0} = |\beta(y)|$.
3. If $|y| = q^{-j}$ with $j \in \mathbb{N}$, then $\ell_j/j = \theta + \varepsilon_j$ for some $\varepsilon_j \in [-K/j, K/j]$, by (5). Thus

$$|\beta(y)| = q^{-\ell_j} = (q^{-j})^{\ell_j/j} = |y|^{\ell_j/j} = |y|^{\theta + \varepsilon_j} = |y|^\theta |y|^{\varepsilon_j},$$

where $|y|^{\varepsilon_j} = q^{-j\varepsilon_j}$ with $j\varepsilon_j \in [-K, K]$ and so $|y|^{\varepsilon_j} \in [q^{-K}, q^K]$. Hence indeed $a|y|^\theta \leq q^{-K}|y|^\theta \leq |\beta(y)| \leq q^K|y|^\theta = b|y|^\theta$. \square

The following proposition provides C^n -automorphisms of the 1-dimensional \mathbb{K} -analytic Lie group $(\mathbb{O}, +)$ which are not C^{n+1} .

Proposition 3.4 *Given $n \in \mathbb{N}$, choose $\theta \in]n, n+1[$ and define $\ell_j := 1 + [j\theta]$ for each $j \in \mathbb{N}_0$, where $[.]$ is the Gauß bracket (integer part). Let $\alpha: \mathbb{O} \rightarrow \mathbb{O}$ be as in 3.1. Then α is a C^n -automorphism of $(\mathbb{O}, +)$ which is not a C^{n+1} -map.*

Proof. Since $j\theta \leq \ell_j \leq j\theta + 1$ for each $j \in \mathbb{N}_0$, Lemma 3.3 provides real numbers $0 < a < b$ such that $a|y - x|^\theta \leq |\beta(y) - \beta(x)| \leq b|y - x|^\theta$ for all $x, y \in \mathbb{O}$. Hence β is C^n (by Lemma 2.1) but not C^{n+1} (by Lemma 2.2).

Furthermore, $\beta'(x) = D^1\beta(x) = 0$ for all $x \in \mathbb{O}$, by Lemma 2.1. Hence $\alpha = \text{id}_{\mathbb{O}} + \beta$ is C^n -map which is not C^{n+1} , and such that $\alpha'(x) = 1$ for each $x \in \mathbb{O}$. As α is an automorphism of groups by Lemma 3.2 and hence a bijection, the Inverse Function Theorem for C^n -maps ([7, Thm. 7.3 & Rem. 5.4] or [13, Thms. 27.5 & 77.2]) implies that α is a C^n -diffeomorphism. \square

To identify suitable smooth mappings as non-analytic, we shall use the following simple fact (see [3, 4.2.3 and 3.2.5]):

Lemma 3.5 *Let E and F be ultrametric Banach spaces over \mathbb{K} , $U \subseteq E$ be an open 0-neighbourhood and $f: U \rightarrow F$ be a \mathbb{K} -analytic function such that*

$$\lim_{x \rightarrow 0} \|f(x)\| / \|x\|^n = 0 \quad \text{for each } n \in \mathbb{N}_0.$$

Then there exists a 0-neighbourhood $W \subseteq U$ such that $f|_W = 0$. \square

We find it convenient to formulate another preparatory lemma.

Lemma 3.6 *In 3.1, choose $\lambda = (\ell_j)_{j \in \mathbb{N}_0}$ such that $\ell_j/j \rightarrow \infty$ as $j \rightarrow \infty$. Then, for each $n \in \mathbb{N}$, there exists a real number $c_n > 0$ such that*

$$|\beta(y) - \beta(x)| \leq c_n |y - x|^n \quad \text{for all } x, y \in \mathbb{O}. \quad (6)$$

Proof. Let $n \in \mathbb{N}$. Since $\ell_j/j \rightarrow \infty$, there exists $k \in \mathbb{N}$ such that $\ell_j/j \geq n$ for all $j \geq k$. We claim that (6) holds with $c_n := q^{kn}$. It suffices to verify (6) for $x = 0$, and we may assume that $y \neq 0$ (cf. proof of Lemma 3.3). If $|y| = q^{-j}$ with $j \geq k$, then $|\beta(y)| = q^{-\ell_j} = (q^{-j})^{\ell_j/j} \leq (q^{-j})^n = |y|^n \leq c_n |y|^n$, as required. If $|y| = q^{-j}$ with $j \in \{0, 1, \dots, k-1\}$, then $|\beta(y)| \leq 1 \leq q^{kn} q^{-jn} = c_n |y|^n$ as well. \square

We now provide examples of smooth, non-analytic automorphisms of the 1-dimensional \mathbb{K} -analytic Lie group $(\mathbb{O}, +)$.

Proposition 3.7 *In 3.1, choose $\lambda = (\ell_j)_{j \in \mathbb{N}_0}$ such that $\ell_j/j \rightarrow \infty$ as $j \rightarrow \infty$. Then the following holds:*

- (a) α is a C^∞ -automorphism of $(\mathbb{O}, +)$.
- (b) α is not \mathbb{K} -analytic.

Proof. (a) Lemmas 3.6 and 2.1 show that β is smooth, with

$$D^j \beta(x) = 0 \quad \text{for all } j \in \mathbb{N} \text{ and } x \in \mathbb{O}. \quad (7)$$

As in the proof of Proposition 3.4, we infer that α is a C^∞ -automorphism.

(b) Since $\beta = \alpha - \text{id}_{\mathbb{O}}$ where $\text{id}_{\mathbb{O}}$ is \mathbb{K} -analytic, it suffices to show that β is not \mathbb{K} -analytic. However, if β was \mathbb{K} -analytic, then β would vanish on some 0-neighbourhood, by (6) and Lemma 3.5. As β is injective, we have reached a contradiction. \square

Note that functions f of one variable with $D^j f = 0$ for small (or all) positive integers are familiar objects in non-archimedean analysis. Examples of such functions in the p -adic case (which are *not* homomorphisms) are described (or suggested) in Example 26.4 as well as Exercises 28.F, 29.F and 29.G in [13]. Our definition of β in **3.1** was stimulated by these examples.

4 Calibrations obtained from contractions

In this section, we provide techniques which shall enable us to check that certain Lie groups constructed with the help of the pathological automorphisms from Section 3 do not admit better-behaved Lie group structures compatible with their given topological group structure. The basic idea is that suitable contractive automorphisms of \mathbb{K} -Lie groups determine a “calibration” which can be used to estimate the norms of elements close to the identity (with respect to a given chart taking 1 to 0). As a tool, we shall employ concepts from the structure theory of totally disconnected, locally compact groups developed in [18], [19] and analyzed for Lie groups over local fields in [5] and [9]. We briefly recall the relevant definitions.

4.1 Let G be a totally disconnected, locally compact topological group and α be a bicontinuous automorphism of G . Then there exists a compact open subgroup $U \subseteq G$ which is *tidy for* α , meaning that the following holds:

T1 $U = U_+ U_-$, where $U_\pm := \bigcap_{k \in \mathbb{N}_0} \alpha^{\pm k}(U)$; and

T2 The subgroup $U_{++} := \bigcup_{k \in \mathbb{N}_0} \alpha^k(U_+)$ is closed in G .

Here $\alpha(U_+) \supseteq U_+$ and $\alpha(U_-) \subseteq U_-$. The index $s_G(\alpha) := [\alpha(U_+) : U_+]$ is finite and is independent of the tidy subgroup; it is called the *scale of* α . If U is tidy for α , then it is also tidy for α^{-1} . (See [18] and [19]).

Let us say that a bicontinuous automorphism $\alpha : G \rightarrow G$ is *contractive* if $\alpha^n(x) \rightarrow 1$ as $n \rightarrow \infty$, for each $x \in G$. A sequence $(U_n)_{n \in \mathbb{N}_0}$ of identity neighbourhoods is called a *fundamental sequence* if $U_0 \supseteq U_1 \supseteq U_2 \supseteq \dots$ and every identity neighbourhood of G contains some U_n .

4.2 If α is contractive and $U \subseteq G$ a compact identity neighbourhood, then $\{\alpha^n(U) : n \in \mathbb{N}_0\}$ is a basis for the filter of identity neighbourhoods of G , as a consequence of [16, Prop. 2.1].

Lemma 4.3 *Let α be a contractive automorphism of a totally disconnected, locally compact topological group G . Then $s_G(\alpha) = 1$, and the following holds:*

- (a) *There exists a compact, open subgroup $U \subseteq G$ such that $\alpha(U) \subseteq U$.*
- (b) *A compact, open subgroup $U \subseteq G$ is tidy for α if and only if $\alpha(U) \subseteq U$. In this case, $U_- = U$,*

$$U_+ := \bigcap_{k \in \mathbb{N}_0} \alpha^k(U) = \{1\}, \quad \text{and} \quad U_{--} := \bigcup_{k \in \mathbb{N}_0} \alpha^{-k}(U_-) = G. \quad (8)$$

- (c) *$(\alpha^n(U))_{n \in \mathbb{N}_0}$ is a fundamental sequence of identity neighbourhoods for G , for each subgroup $U \subseteq G$ which is tidy for α . Each of the groups $\alpha^n(U)$ is tidy for α . Hence G has arbitrarily small subgroups tidy for α .*
- (d) *If G is a C^1 -Lie group over a local field \mathbb{K} and α is a contractive C^1 -automorphism, let $\beta := L(\alpha)$ be the associated \mathbb{K} -linear automorphism of $L(G)$. Then $|\lambda| < 1$ for every eigenvalue λ of β in $\overline{\mathbb{K}}$.*

Proof. (a) As tidy subgroups always exist, (a) follows from (b).

(b) If $U \subseteq G$ is tidy for α , then $\alpha^n(x) \rightarrow 1$ as $n \rightarrow \infty$ for $x \in U$ entails that $x \in U_-$ (cf. [18, La. 9]). Hence $U = U_-$ and hence $\alpha(U) \subseteq U$. If, conversely, $U \subseteq G$ is a compact open subgroup such that $\alpha(U) \subseteq U$, then $\alpha^{-1}(U) \supseteq U$ and thus $U_- = U$. Since G is Hausdorff, **4.2** shows that $U_+ = \bigcap_{n \geq 0} \alpha^n(U) = \{1\}$ and hence $U_{++} = \{1\}$, which is closed. Thus U satisfies **T1** and **T2**. Since $U_+ = \{1\}$, furthermore $s_G(\alpha) = [\{1\} : \{1\}] = 1$. To complete the proof of (8), let $x \in G$. Then $\alpha^n(x) \in U$ for some $n \in \mathbb{N}_0$ and thus $x \in \alpha^{-n}(U) = \alpha^{-n}(U_-) \subseteq U_{--}$. Hence $U_{--} = G$.

(c) As $\alpha^{n+1}(U) \subseteq \alpha^n(U)$ for each n , the claims follow from (b) and **4.2**.

(d) As G has small subgroups tidy for α , results from [9] show that $G_\alpha := \{x \in G : \alpha^n(x) \rightarrow 1 \text{ as } n \rightarrow \infty\}$, $G_{\alpha^{-1}} := \{x \in G : \alpha^{-n}(x) \rightarrow 1 \text{ as } n \rightarrow \infty\}$ and $G_0 := \{x \in G : \alpha^{\mathbb{Z}}(x) \text{ is relatively compact}\}$ are C^1 -Lie subgroups of G , and that the multiplication map

$$G_\alpha \times G_0 \times G_{\alpha^{-1}} \rightarrow G, \quad (x, y, z) \mapsto xyz$$

is a C^1 -diffeomorphism onto an open subset of G . As $G_\alpha = G$, this implies that $G_0 = G_{\alpha^{-1}} = \{1\}$, whence $L(G_0) = L(G_{\alpha^{-1}}) = \{0\}$. However, it is also shown in [9] that $L(G_{\alpha^{-1}})_{\overline{\mathbb{K}}}$ (resp., $L(G_0)_{\overline{\mathbb{K}}}$) is the direct sum of all the generalized eigenspaces of $\beta_{\overline{\mathbb{K}}}$ to eigenvalues of absolute value > 1 (resp., of absolute value 1). Hence such eigenvalues cannot exist. \square

In view of Lemma 4.3 (b) and (c), the following definition makes sense:

Definition 4.4 Let G be a totally disconnected, locally compact group, α be a contractive automorphism of G and $U \subseteq G$ a compact, open subgroup such that $\alpha(U) \subseteq U$. The *calibration of G associated with α and U* is the function $\kappa: G \rightarrow \mathbb{Z} \cup \{\infty\}$ defined via

$$\kappa(x) := \begin{cases} \infty & \text{if } x = 1; \\ n & \text{if } x \in \alpha^n(U) \setminus \alpha^{n-1}(U). \end{cases} \quad (9)$$

4.5 Let G be a C^1 -Lie group over $\mathbb{K} = \mathbb{F}_q((X))$ and $\alpha: G \rightarrow G$ be a contractive C^1 -automorphism of G such that $s_G(\alpha^{-1}) = q$. Let $\phi: P \rightarrow Q \subseteq L(G)$ be a chart for G around 1 such that $\phi(1) = 0$ and $T_1(\phi) = \text{id}_{L(G)}$. Let $\|\cdot\|$ be an ultrametric norm on $L(G)$. For each $r > 0$, let B_r be the ball of radius r in $(L(G), \|\cdot\|)$; let $\rho > 0$ such that $B_\rho \subseteq Q$, and set $W_r := \phi^{-1}(B_r)$ for $0 < r \leq \rho$. Let U be a compact open subgroup of G such that $\alpha(U) \subseteq U$.

Lemma 4.6 *In the situation of 4.5, there exist $N \in \mathbb{N}_0$ and real numbers $a \leq b$ such that $q^b \leq \rho$ and*

$$W_{q^{a-n/d}} \subseteq \alpha^{n+N}(U) \subseteq W_{q^{b-n/d}} \quad \text{for all } n \in \mathbb{N}_0. \quad (10)$$

Proof. Let $\lambda_1, \dots, \lambda_d$ (where $d := \dim_{\mathbb{K}} L(G)$) be the eigenvalues of $L(\alpha^{-1})$ in $\overline{\mathbb{K}}$, each of which has absolute value > 1 as a consequence of Lemma 4.3 (d). Since G has small subgroups tidy for α (Lemma 4.3 (c)), it also has small subgroups tidy for α^{-1} . Hence

$$q = s_G(\alpha^{-1}) = \prod_{\substack{j \in \{1, \dots, d\} \\ \text{s.t. } |\lambda_j| \geq 1}} |\lambda_j| = \prod_{j=1}^d |\lambda_j|, \quad (11)$$

using the main result of [9] to obtain the second equality. Now Lemma 1.7 shows that $|\lambda_j| = q^{1/d}$ for each $j \in \{1, \dots, d\}$, and it provides an ultrametric norm $\|\cdot\|'$ on $L(G)$ such that $\|L(\alpha^{-1}).x\| = q^{1/d}\|x\|'$ for each $x \in L(G)$. Hence

$$\|\beta(x)\|' = q^{-1/d} \|x\|', \quad \text{for each } x \in L(G),$$

where $\beta := L(\alpha)$. Set $B'_r := \{x \in L(G) : \|x\|' < r\}$ for $r > 0$. There exists an open 0-neighbourhood $R \subseteq Q$ such that $\alpha(\phi^{-1}(R)) \subseteq P$. Then

$$\gamma : R \rightarrow Q, \quad \gamma(x) := \phi(\alpha(\phi^{-1}(x)))$$

expresses α in local coordinates. We have $\gamma'(0) = T_1(\phi)L(\alpha)T_1(\phi)^{-1} = L(\alpha) = \beta$ as $T_1(\phi) = \text{id}_{L(G)}$. Thus $\gamma'(0) = \beta$ is invertible. Now the Inverse Function Theorem in the quantitative form [7, Prop. 7.1 (b)'] shows that there exists $\delta > 0$ with $B'_\delta \subseteq R$ such that $\gamma(B'_r) = \beta(B'_r) = B'_{rq^{-1/d}}$ for each $r \in]0, \delta]$ and hence

$$\gamma^n(B'_r) = B'_{rq^{-n/d}} \quad \text{for all } r \in]0, \delta] \text{ and } n \in \mathbb{N}_0. \quad (12)$$

All ultrametric norms on the finite-dimensional \mathbb{K} -vector space $L(G)$ being equivalent [13, Thm. 13.3], there exists $c \geq 1$ such that $c^{-1}\|\cdot\| \leq \|\cdot\|' \leq c\|\cdot\|$ and thus $B_{c^{-1}r} \subseteq B'_r \subseteq B_{cr}$ for each $r > 0$. Set $v := \min\{1, \frac{\rho}{c\delta}\}$. Then $B'_{v\delta} \subseteq B'_\delta \subseteq R$. Choose $N \in \mathbb{N}$ such that $\alpha^N(U) \subseteq \phi^{-1}(B'_{v\delta})$. There exists $u \in]0, v]$ such that $B'_{u\delta} \subseteq \phi(\alpha^N(U))$. Then

$$B'_{u\delta q^{-n/d}} \subseteq \gamma^n(\phi(\alpha^N(U))) \subseteq B'_{v\delta q^{-n/d}} \quad \text{for each } n \in \mathbb{N}_0,$$

by (12). Therefore $B_{c^{-1}u\delta q^{-n/d}} \subseteq \gamma^n(\phi(\alpha^N(U))) \subseteq B_{cv\delta q^{-n/d}}$. Using that $\gamma^n(\phi(\alpha^N(U))) = \phi(\alpha^{n+N}(U))$, we obtain

$$B_{c^{-1}u\delta q^{-n/d}} \subseteq \phi(\alpha^{n+N}(U)) \subseteq B_{cv\delta q^{-n/d}} \quad \text{for each } n \in \mathbb{N}_0. \quad (13)$$

Note that $cv\delta q^{-n/d} \leq cv\delta \leq \frac{c\delta\rho}{c\delta} = \rho$ be definition of v . Applying ϕ^{-1} to (13), we see that (10) holds with $a := \log_q(\frac{u\delta}{c})$ and $b := \log_q(cv\delta)$. \square

In the following lemma, we retain the situation of Lemma 4.6; in particular, $d := \dim_{\mathbb{K}} L(G)$. For $\|\cdot\|$ as in 4.5, define $\mu : L(G) \rightarrow \mathbb{R} \cup \{\infty\}$, $\mu(x) := -\log_q \|x\|$ (where $\log_q(0) := -\infty$). Thus $\|x\| = q^{-\mu(x)}$. The calibration $\kappa : G \rightarrow \mathbb{Z} \cup \{\infty\}$ associated with α and U (as in (9)) can be used to approximate μ on a 0-neighbourhood, up to a bounded error:

Lemma 4.7 (Calibration Lemma) *There exists $R \in [0, \infty[$ such that*

$$\mu(\phi(x))d - R \leq \kappa(x) \leq \mu(\phi(x))d + R, \quad \text{for each } x \in \alpha^N(U). \quad (14)$$

Proof. Let $x \in \alpha^N(U) \setminus \{1\}$. Then $N \leq \kappa(x) < \infty$, entailing that $n := \kappa(x) - N \in \mathbb{N}_0$. Using (10), we obtain

$$\begin{aligned} x \in \alpha^{\kappa(x)}(U) &= \alpha^{n+N}(U) \subseteq W_{q^{b-n/d}} && \text{and} \\ x \notin \alpha^{\kappa(x)+1}(U) &= \alpha^{n+1+N}(U) \supseteq W_{q^{a-(n+1)/d}}. \end{aligned}$$

Hence $q^{a-(n+1)/d} \leq \|\phi(x)\| < q^{b-n/d}$ and thus $(n+1)/d - a \geq \mu(\phi(x)) > n/d - b$, whence $n+1 - ad \geq \mu(\phi(x))d > n - bd$ and therefore

$$\kappa(x) - N + 1 - ad \geq \mu(\phi(x))d > \kappa(x) - N - bd.$$

Hence the assertion holds with $R := \max\{0, 1 - ad - N, N + bd\}$. \square

The following immediate consequence of Lemma 4.7 is what we actually need:

Proposition 4.8 *Let G and G^* be C^1 -Lie groups over \mathbb{K} , of dimensions d and d^* , resp., such that $G = G^*$ as a topological group. Let $\phi : P \rightarrow Q \subseteq L(G)$ and $\phi^* : P^* \rightarrow Q^*$ be charts of G and G^* around 1 taking 1 to 0 and such that $T_1(\phi) = \text{id}_{L(G)}$ and $T_1(\phi^*) = \text{id}_{L(G^*)}$. Let $\|\cdot\|$ and $\|\cdot\|_*$ be ultrametric norms on $L(G)$ and $L(G^*)$, respectively. If there exists a contractive automorphism α of G which is a C^1 -automorphism for both G and G^* and such that $s_G(\alpha^{-1}) = q$, then there exists an identity neighbourhood $W \subseteq P \cap P^*$ and positive real numbers $a < b$ such that*

$$a \|\phi(x)\|^{d/d^*} \leq \|\phi^*(x)\|_* \leq b \|\phi(x)\|^{d/d^*} \quad \text{for each } x \in W. \quad \square$$

It may very well happen that $d \neq d^*$ in the situation of Proposition 4.8, and in fact any combination of positive integers $(d, d^*) \in \mathbb{N}^2$ can occur, as the following example shows.

Example 4.9 Let $G := (\mathbb{K}, +)$, equipped with its usual one-dimensional \mathbb{K} -analytic Lie group structure. Then the (linear) map

$$\tau : \mathbb{K} \rightarrow \mathbb{K}, \quad \tau(z) := X \cdot z$$

apparently is a \mathbb{K} -analytic automorphism of G . The isomorphism of topological groups

$$\phi^* : \mathbb{K} \rightarrow \mathbb{K}^2, \quad \sum_j a_j X^j \mapsto \left(\sum_j a_{2j} X^j, \sum_j a_{2j+1} X^j \right)$$

can be used as a global chart for \mathbb{K} turning it into a 2-dimensional \mathbb{K} -analytic Lie group G^* isomorphic to $(\mathbb{K}^2, +)$. Note that τ also is a \mathbb{K} -analytic automorphism of G^* , because $\phi^* \circ \tau \circ (\phi^*)^{-1} : \mathbb{K}^2 \rightarrow \mathbb{K}^2$ is the \mathbb{K} -linear map

$$\begin{pmatrix} u \\ v \end{pmatrix} \mapsto \begin{pmatrix} 0 & X \\ 1 & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}.$$

By the following lemma, τ is contractive and $s_G(\tau^{-1}) = q$. Hence, we are in the situation of Proposition 4.8.

Variants: Given any positive integer d^* , we obtain an isomorphism of topological groups $\mathbb{K} \cong \mathbb{K}^{d^*}$ if we replace $2\mathbb{Z}$ and $2\mathbb{Z} + 1$ with $d^*\mathbb{Z}$ and its cosets in the preceding construction. This enables us to turn \mathbb{K} into a \mathbb{K} -analytic Lie group G^* of dimension d^* , such that τ is a \mathbb{K} -analytic automorphism of G^* .

We used the following simple fact:

Lemma 4.10 τ is a contractive automorphism of $(\mathbb{K}, +)$ with $s_{\mathbb{K}}(\tau^{-1}) = q$.

Proof. Because $|\tau^n(z)| = |X^n z| = q^{-n}|z|$ for each $z \in \mathbb{K}$ and $n \in \mathbb{N}$, clearly τ is contractive. Since $\tau(\mathbb{O}) = X\mathbb{O} \subseteq \mathbb{O}$, Lemma 4.3 (b) shows that \mathbb{O} is tidy for τ , with $\mathbb{O}_+ = \{0\}$ and $\mathbb{O}_- = \mathbb{O}$. Therefore $s_{\mathbb{K}}(\tau^{-1}) = [\tau^{-1}(\mathbb{O}) : \mathbb{O}] = [X^{-1}\mathbb{O} : \mathbb{O}] = |\mathbb{F}_q| = q$. \square

5 Examples of non-analytic C^∞ -Lie groups

In this section, we present examples of smooth \mathbb{K} -Lie groups which cannot be turned into \mathbb{K} -analytic Lie groups.

5.1 Let $\lambda = (\ell_j)_{j \in \mathbb{N}_0}$, α and β be as in **3.1**. Every element $z = \sum_{j=-n}^{\infty} a_j X^j \in \mathbb{K}$ (where $n \in \mathbb{N}$) can be written as $z = z' + z''$ with $z' := \sum_{j=-n}^{-1} a_j X^j$ and $z'' := \sum_{j=0}^{\infty} a_j X^j$. We define

$$\bar{\beta} : \mathbb{K} \rightarrow \mathbb{K}, \quad \bar{\beta}(z) := \beta(z'') \quad \text{and}$$

$$\bar{\alpha}: \mathbb{K} \rightarrow \mathbb{K}, \quad \bar{\alpha}(z) := z' + \alpha(z'').$$

Thus $\bar{\alpha}|_{\mathbb{O}} = \alpha$, $\bar{\beta}|_{\mathbb{O}} = \beta$, and $\bar{\alpha} = \text{id}_{\mathbb{K}} + \bar{\beta}$. Clearly $\bar{\beta}$ is a continuous homomorphism and $\bar{\alpha}$ a bicontinuous automorphism of $(\mathbb{K}, +)$. We define

$$\tau: \mathbb{K} \rightarrow \mathbb{K}, \quad \tau(z) := X \cdot z$$

and recall from **4.9** and Lemma 4.10 that τ is a \mathbb{K} -analytic, contractive automorphism of $(\mathbb{K}, +)$ such that $s_{\mathbb{K}}(\tau^{-1}) = q$. We equip

$$G := \mathbb{K} \rtimes \langle \bar{\alpha}, \tau \rangle \tag{15}$$

with the uniquely determined topology making it a topological group with \mathbb{K} (in its usual topology) as an open, normal subgroup; here $\langle \bar{\alpha}, \tau \rangle$ denotes the subgroup of $\text{Aut}(\mathbb{K}, +)$ generated by $\bar{\alpha}$ and τ .

Theorem 5.2 *For $\lambda = (\ell_j)_{j \in \mathbb{N}_0}$ as in Proposition 3.7, the topological group G described in (15) has the following properties:*

- (a) *G admits a 1-dimensional smooth \mathbb{K} -Lie group structure compatible with the given topological group structure.*
- (b) *G does not admit a \mathbb{K} -analytic Lie group structure compatible with the given topological group structure.*

Proof. (a) We equip $(\mathbb{K}, +)$ with its usual 1-dimensional C^∞ -Lie group structure. Since the restriction of $\bar{\alpha}$ to the open subset $\mathbb{O} \subseteq \mathbb{K}$ coincides with α , which is a C^∞ -automorphism of \mathbb{O} by Proposition 3.7, we see that $\bar{\alpha}$ is a C^∞ -automorphism of $(\mathbb{K}, +)$. Apparently also τ is a C^∞ -automorphism of $(\mathbb{K}, +)$, and hence so is each $\gamma \in \langle \bar{\alpha}, \tau \rangle$. Now standard arguments provide a (uniquely determined) C^∞ -manifold structure on G making it a smooth Lie group and such that \mathbb{K} , with its given smooth manifold structure, is an open submanifold of G (see [8, Prop. 1.18]).

- (b) The proof has two parts, the first of which can be re-used later.

5.3 Let us assume, to the contrary, that there exists a \mathbb{K} -analytic manifold structure on the group G making it a \mathbb{K} -analytic Lie group G^* and compatible with the given topology. We let $L(G^*)$ be the Lie algebra of G^* and set $d := \dim_{\mathbb{K}} L(G^*) \in \mathbb{N}$. The open subgroup \mathbb{K} of G^* inherits a \mathbb{K} -analytic Lie group structure from G^* . We write H for \mathbb{K} , equipped with this d -dimensional

\mathbb{K} -analytic Lie group structure. Thus $L(H) = L(G^*)$, and both τ and $\bar{\alpha}$ are \mathbb{K} -analytic automorphisms of H . We let $\phi: P \rightarrow Q \subseteq L(H)$ be a chart for H around 0 such that $\phi(0) = 0$ and $T_0(\phi) = \text{id}_{L(H)}$. Furthermore, we choose an ultrametric norm $\|\cdot\|$ on $L(H)$. Using Proposition 4.8, we find $N \in \mathbb{N}_0$ and positive real numbers $a < b$ such that $X^N\mathbb{O} \subseteq P$ and

$$a \|\phi(z)\|^d \leq |z| \leq b \|\phi(z)\|^d \quad \text{for each } z \in X^N\mathbb{O}. \quad (16)$$

Set $W := \phi(X^N\mathbb{O})$. Since $\beta(X^N\mathbb{O}) \subseteq X^N\mathbb{O}$, we can define $\gamma: W \rightarrow W$, $\gamma(w) := \phi(\beta(\phi^{-1}(w)))$.

5.4 Using Lemma 3.5, we want to show that γ is not analytic. Given $n \in \mathbb{N}$, there exists $k \geq N$ such that $\ell_j \geq (n+1)j$ for each $j \geq k$ and hence

$$|\beta(z)| \leq |z|^{n+1} \quad \text{for each } z \in X^k\mathbb{O}. \quad (17)$$

For $w \in W$ such that $\|w\| \leq q^{-k/d}b^{-1/d}$, we have $|\phi^{-1}(w)| \leq b\|w\|^d \leq q^{-k}$ (using (16)) and hence $|\beta(\phi^{-1}(w))| \leq |\phi^{-1}(w)|^{n+1}$, by (17). As a consequence,

$$\|\gamma(w)\| = \|\phi(\beta(\phi^{-1}(w)))\| \leq a^{-1/d}|\phi^{-1}(w)|^{(n+1)/d} \leq a^{-1/d}b^{(n+1)/d}\|w\|^{n+1},$$

entailing that $\|\gamma(w)\|/\|w\|^n \rightarrow 0$ as $w \rightarrow 0$. Since γ is injective, Lemma 3.5 shows that γ cannot be analytic. As a consequence, $\bar{\beta}: H \rightarrow H$ is not analytic either, nor is $\bar{\alpha} = \text{id}_H + \bar{\beta}: H \rightarrow H$. But $\bar{\alpha}: H \rightarrow H$ has to be analytic, being the restriction of an inner automorphism of the analytic Lie group G^* to its open, normal subgroup H . We have reached a contradiction. \square

Remark 5.5 In particular, Theorem 5.2 shows that there exist smooth Lie groups over local fields of positive characteristic without a *smoothly* compatible analytic Lie group structure.

6 Examples of C^n -Lie groups that are not C^{n+1}

A variant of the proof of Theorem 5.2 shows the following:

Theorem 6.1 *Given $n \in \mathbb{N}$, let θ and $\lambda = (\ell_j)_{j \in \mathbb{N}_0}$ be as in Proposition 3.4. Then the topological group G from (15) has the following properties:*

- (a) *G admits a 1-dimensional \mathbb{K} -Lie group structure of class C^n compatible with the given topological group structure.*

(b) G does not admit a \mathbb{K} -Lie group structure of class C^{n+1} compatible with the given topological group structure.

Proof. The proof of Theorem 5.2(a) and the first half of the proof of Theorem 5.2(b) (viz. **5.3**) carry over to the present situation, if we replace the word “smooth” with “ C^n ” and “ \mathbb{K} -analytic” with “ C^{n+1} ” there. Thus Part (a) of the present theorem holds. To complete the proof of (b), we retain the notations from **5.3**. Using Lemma 2.2, we want to show that γ is not C^{n+1} (whence neither is $\bar{\beta}$, nor $\bar{\alpha}$, which is absurd). By Lemma 3.3, there exist positive real numbers $A < B$ such that

$$A|z|^\theta \leq |\beta(z)| \leq B|z|^\theta, \quad \text{for each } z \in \mathbb{O}. \quad (18)$$

For each $w \in W$, we then have

$$\begin{aligned} \|\gamma(w)\| &= \|\phi(\beta(\phi^{-1}(w)))\| \leq a^{-1/d} |\beta(\phi^{-1}(w))|^{1/d} \\ &\leq a^{-1/d} B^{1/d} |\phi^{-1}(w)|^{\theta/d} \leq a^{-1/d} b^{\theta/d} B^{1/d} \|w\|^\theta \quad \text{and} \end{aligned}$$

$$\begin{aligned} \|\gamma(w)\| &= \|\phi(\beta(\phi^{-1}(w)))\| \geq b^{-1/d} |\beta(\phi^{-1}(w))|^{1/d} \\ &\geq b^{-1/d} A^{1/d} |\phi^{-1}(w)|^{\theta/d} \geq b^{-1/d} A^{1/d} a^{\theta/d} \|w\|^\theta, \end{aligned}$$

using (16) and (18). Set $C := b^{-1/d} A^{1/d} a^{\theta/d}$ and $D := a^{-1/d} b^{\theta/d} B^{1/d}$. Then

$$C\|w\|^\theta \leq \|\gamma(w)\| \leq D\|w\|^\theta \quad \text{for each } w \in W,$$

whence Lemma 2.2 shows that γ is not C^{n+1} , contradiction. \square

The following problems remain open:

Problem 1 Does every smooth \mathbb{K} -Lie group have an open subgroup which admits a smoothly compatible analytic Lie group structure?

Problem 2 Given $n \in \mathbb{N}$, does every \mathbb{K} -Lie group of class C^n have an open subgroup which admits a C^n -compatible \mathbb{K} -Lie group structure of class C^{n+1} ?

Acknowledgements. The present research was carried out at the University of Newcastle (N.S.W.) in August 2004, supported by DFG grant 447 AUS-113/22/0-1 and ARC grant LX 0349209. The author profited from discussions with George Willis and Udo Baumgartner. Remark 5.5 answers a question posed to the author by Linus Kramer (Darmstadt) in 2003.

References

- [1] Baumgartner, U. and G. A. Willis, *Contraction groups and scales of automorphisms of totally disconnected locally compact groups*, to appear in Israel J. Math. (cf. arXiv:math.GR/0302095).
- [2] Bertram, W., H. Glöckner and K.-H. Neeb, *Differential calculus over general base fields and rings*, Expo. Math. **22** (2004), 213–282.
- [3] Bourbaki, N., “Variétés différentielles et analytiques. Fascicule de résultats,” Hermann, Paris, 1967.
- [4] —, “Lie Groups and Lie Algebras” (Chapters 1–3), Springer-Verlag, 1989.
- [5] Glöckner, H., *Scale functions on p -adic Lie groups*, Manuscr. Math. **97** (1998), 205–215.
- [6] —, *Every smooth p -adic Lie group admits a compatible analytic structure*, to appear in Forum Math. (cf. arXiv:math.GR/0312113).
- [7] —, *Implicit functions from topological vector spaces to Banach spaces*, preprint, March 2003, arXiv:math.GM/0303320.
- [8] —, *Lie groups over non-discrete topological fields*, preprint, August 2004, arXiv:math.GR/0408008.
- [9] —, *Scale functions on Lie groups over local fields of positive characteristic*, in preparation (will be placed in the arXiv in September 2004).
- [10] Lazard, M., *Groupes analytiques p -adiques*, IHES Publ. Math. **26** (1965), 389–603.
- [11] Margulis, G. A., “Discrete Subgroups of Semisimple Lie Groups,” Springer-Verlag, 1991.
- [12] Montgomery, D. and L. Zippin, “Topological Transformation Groups,” Interscience, New York, 1955.
- [13] Schikhof, W. H., “Ultrametric Calculus,” Cambridge University Press, 1984.
- [14] Schur, F., *Zur Theorie der endlichen Transformationsgruppen*, Math. Ann. **38** (1891), 263–286; cf. JFM 23.0381.01
- [15] Serre, J.-P., “Lie Algebras and Lie Groups,” Springer-Verlag, 1992.
- [16] Wang, J. S. P., *The Mautner phenomenon for p -adic Lie groups*, Math. Z. **185** (1984), 403–412.
- [17] Weil, A., “Basic Number Theory,” Springer-Verlag, 1967.
- [18] Willis, G. A., *The structure of totally disconnected, locally compact groups*, Math. Ann. **300** (1994), 341–363.
- [19] —, *Further properties of the scale function on a totally disconnected group*, J. Algebra **237** (2001), 142–164.

Helge Glöckner, TU Darmstadt, Fachbereich Mathematik AG 5, Schlossgartenstr. 7,
64289 Darmstadt, Germany. E-Mail: gloeckner@mathematik.tu-darmstadt.de