

# COUNTEREXAMPLES TO THE NEGGERS-STANLEY CONJECTURE

PETTER BRÄNDÉN

ABSTRACT. The Neggers-Stanley conjecture (also known as the Poset conjecture) asserts that the polynomial counting the linear extensions of a partially ordered set on  $\{1, 2, \dots, p\}$  by their number of descents has real zeros only. We provide counterexamples to this conjecture.

## 1. THE NEGGERS-STANLEY CONJECTURE

Let  $P$  be a poset (partially ordered set) on  $[p] := \{1, 2, \dots, p\}$ . We will use the symbol  $<$  to denote the usual order on the integers and the symbol  $\prec$  to denote the partial order on  $P$ . The *Jordan-Hölder set*,  $\mathcal{L}(P)$ , of  $P$  is the set of permutations  $\pi$  of  $[p]$  which are linear extensions of  $P$ , i.e., if  $i \prec j$  then  $\pi^{-1}(i) < \pi^{-1}(j)$ . A *descent* in a permutation  $\pi$  is an index  $i \in [p-1]$  such that  $\pi(i) > \pi(i+1)$ . Let  $\text{des}(\pi)$  denote the number of descents in  $\pi$ . The  $W$ -polynomial of  $P$  is defined by

$$W(P, t) = \sum_{\pi \in \mathcal{L}(P)} t^{\text{des}(\pi)}.$$

The  $W$ -polynomials appear naturally in many combinatorial structures, see [2, 7, 8], and are connected to Hilbert series of the Stanley-Reisner rings of simplicial complexes [10, Section III.7] and algebras with straightening laws [9, Thm. 5.2.].

**Example 1.** Let  $P_{2,2}$  be as in Fig. 1. Then

$$\mathcal{L}(P_{2,2}) = \{(1, 3, 2, 4), (1, 3, 4, 2), (3, 1, 2, 4), (3, 1, 4, 2), (3, 4, 1, 2)\}$$

so  $W(P_{2,2}, t) = 4t + t^2$ .


When  $P$  has no relations, that is, when  $P$  is the anti-chain on  $[p]$ , then  $W(P, t)$  is the  $p$ th *Eulerian polynomial*. The Eulerian polynomials are known [3] to have all zeros real. This is an instance where the Neggers-Stanley conjecture, also known as the Poset conjecture, holds:

**Conjecture 1** (Neggers-Stanley). *Let  $P$  be a poset on  $[p]$ . Then all zeros of  $W(P, t)$  are real.*

---

2000 *Mathematics Subject Classification*. Primary 06A07, 26C10.

*Key words and phrases*. Neggers-Stanley conjecture, partially ordered set, linear extension, real roots.

FIGURE 1. The posets  $P_{2,2}$  and  $P_{3,4}$ .

A poset  $P$  is naturally labeled if  $i \prec j$  implies  $i < j$ . The above conjecture was made by J. Neggers [4] in 1978 for naturally labeled posets and by R.P. Stanley in 1986 for arbitrary posets on  $[p]$ . It has been proved for some special cases, see [2, 11].

If a polynomial  $p(t) = a_0 + a_1t + \cdots + a_nt^n$  with non-negative coefficients is real-rooted it follows that the sequence  $\{a_i\}_{i=0}^n$  is *unimodal* i.e., there is a  $d$  such that  $a_0 \leq a_1 \leq \cdots \leq a_{d-1} \leq a_d \geq a_{d+1} \geq \cdots \geq a_n$ . This consequence of the Neggers-Stanley conjecture was recently proved [6] (see also [1]) for an important class of naturally labeled posets, namely *graded posets*. A naturally labeled poset  $P$  is graded if all maximal chains in  $P$  have the same length.

## 2. COUNTEREXAMPLES TO THE NEGGERS-STANLEY CONJECTURE

Let  $\mathbf{m} \sqcup \mathbf{n}$  be the disjoint union of the chains  $1 \prec 2 \prec \cdots \prec m$  and  $m+1 \prec n+2 \prec \cdots \prec m+n$ . Simion [7] proved that  $W(\mathbf{m} \sqcup \mathbf{n}, t)$  has real and simple zeros only. Let  $P_{m,n}$  be the poset obtained by adding the relation  $m+1 \prec m$  to the relations in  $\mathbf{m} \sqcup \mathbf{n}$ , see Fig. 1. The only linear extension of  $\mathbf{m} \sqcup \mathbf{n}$  which is not a linear extension of  $P_{m,n}$  is  $(1, 2, \dots, m+n)$ , which gives

$$W(\mathbf{m} \sqcup \mathbf{n}, t) = 1 + W(P_{m,n}, t).$$

Let  $\pi$  be a permutation. We say that  $\pi(i)$  is a *descent top* and that  $\pi(i+1)$  is a *descent bottom* if  $i$  is a descent in  $\pi$ . A permutation in  $\mathcal{L}(\mathbf{m} \sqcup \mathbf{n})$  is uniquely determined by its descent tops (which are necessarily elements of  $[m+n] \setminus [m]$ ) and its descent bottoms (which are elements of  $[m]$ ). It follows that the number of permutations in  $\mathcal{L}(\mathbf{m} \sqcup \mathbf{n})$  with exactly  $k$  descents is  $\binom{m}{k} \binom{n}{k}$ , so we have

$$W(\mathbf{m} \sqcup \mathbf{n}, t) = \sum_{k=0}^n \binom{m}{k} \binom{n}{k} t^k.$$

**Theorem 1.** *Let  $M > 0$  be an integer. There is an integer  $N = N(M)$  such that  $W(P_{m,n}, t)$  has more than  $M$  non-real zeros whenever  $\min(m, n) > N$ .*

*Proof.* Recall that the *Bessel function* of order 0 is given by

$$J_0(z) = \frac{2}{\pi} \int_0^1 \frac{\cos(zt)}{\sqrt{1-t^2}} dt = \sum_{k=0}^{\infty} \frac{1}{k!k!} \left(\frac{-z^2}{2}\right)^k. \quad (1)$$

It follows from (1) that  $J_0(z)$  has infinitely many zeros all of which are real and simple and that  $|J_0(t)| \leq 1$  for all real  $t$  with equality only if  $t = 0$ . Hence the function

$$F(z) = \sum_{k=0}^{\infty} \frac{1}{k!k!} z^k.$$

has infinitely many zeros all of which are negative and simple and  $|F(t)| < 1$  for  $t < 0$ .

Let  $f_{m,n}(t) = W(P_{m,n}, t/mn)$ . Then

$$f_{m,n}(t) = \sum_{k=1}^{\min(m,n)} \frac{\gamma_{m,k} \gamma_{n,k}}{k!k!} t^k,$$

where  $\gamma_{n,k} = (1 - \frac{1}{n})(1 - \frac{2}{n}) \cdots (1 - \frac{k-1}{n})$ . Let  $\{(m_j, n_j)\}_{j=1}^{\infty} \subset \mathbb{N} \times \mathbb{N}$  be a sequence such that  $\lim_{j \rightarrow \infty} \min(m_j, n_j) = \infty$ . Then

$$\lim_{j \rightarrow \infty} f_{m_j, n_j}(z) = F(z) - 1,$$

where the sequence converges uniformly on all compact subsets of  $\mathbb{C}$ . Let  $(-a, 0)$  be an interval containing more than  $M$  zeros of  $F(z)$ . It follows from Hurwitz's theorem [5, Thm. 1.3.8] that  $f_{m_j, n_j}(z) + 1$  has more than  $M$  zeros in  $(-a, 0)$  for sufficiently large  $j$  and by continuity that  $|f_{m_j, n_j}(z) + 1| < 1$  for  $z \in (-a, 0)$ . Thus by subtracting 1 from  $f_{m_j, n_j}(z) + 1$  we will lose at least  $M$  real zeros.  $\square$

By applying Sturm's Theorem [5, Section 10.5] one can find specific counterexamples. The polynomial  $W(P_{11,11}, t)$  has two non-real zeros which are approximately

$$z = -0.10902 \pm 0.01308i.$$

A counterexample with a polynomial of lower degree is

$$W(P_{36,6}, t) = 216t + 9450t^2 + 142800t^3 + 883575t^4 + 2261952t^5 + 1947792t^6.$$

This polynomial has two non-real zeros.

**Remark 1.** It should be noted that our counterexamples are not naturally labeled, since we have  $n + 1 \prec n$ . The Neggers-Stanley conjecture is therefore still open for naturally labeled posets.

## REFERENCES

- [1] P. Brändén, Sign-graded posets, unimodality of  $W$ -polynomials and the Charney-Davis conjecture. *arXiv preprint: math.CO/0406019*
- [2] F. Brenti, Unimodal, Log-Concave and Pólya Frequency Sequences in Combinatorics. *Memoirs Amer. Math. Soc.* **413**, Amer. Math. Soc., Providence, 1989.
- [3] L.H. Harper, Stirling behavior is asymptotically normal. *Ann. Math. Statist.* **38** (1967), 410–414.
- [4] J. Neggers, Representations of finite partially ordered sets. *J. Comb. Inf. Syst. Sci.* **3** (1978), 113–133.
- [5] Q.I. Rahman and G. Schmeisser, Analytic theory of polynomials. volume 26 of *London Mathematical Society Monographs. New Series*. The Clarendon Press Oxford University Press, Oxford, 2002.
- [6] V. Reiner and V. Welker, On the Charney-Davis and Neggers-Stanley conjectures. *preprint available at [www.math.umn.edu/~reiner](http://www.math.umn.edu/~reiner)*.
- [7] R. Simion, A multiindexed Sturm sequence of polynomials and unimodality of certain combinatorial sequences. *J. Comb. Theory, Ser. A* **36** (1984), 15–22.
- [8] R.P. Stanley, Ordered structures and partitions. *Memoirs Amer. Math. Soc.* **119**, Amer. Math. Soc., Providence, 1972.
- [9] R.P. Stanley, Hilbert functions of graded algebras. *Adv. Math.* **28** (1978), 57–83.
- [10] R.P. Stanley, Combinatorics and commutative algebra. volume 119 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1996
- [11] D. Wagner, Enumeration of functions from posets to chains. *Eur. J. Comb.* **13** (1992), 313–324.

DEPARTMENT OF MATHEMATICS, CHALMERS UNIVERSITY OF TECHNOLOGY  
AND GÖTEBORG UNIVERSITY, S-412 96 GÖTEBORG, SWEDEN

*E-mail address:* `branden@math.chalmers.se`