

Bi-relative algebraic K -theory and topological cyclic homology

Thomas Geisser and Lars Hesselholt

Introduction

It was recently proved by Cortiñas [5] that rationally bi-relative algebraic K -theory and bi-relative cyclic homology agree. In this paper we show that with finite coefficients bi-relative algebraic K -theory and bi-relative topological cyclic homology agree. As an application, we show that for a possibly singular curve over a field k of positive characteristic p , the cyclotomic trace induces an isomorphism of the p -adic algebraic K -groups and the p -adic topological cyclic homology groups in degrees greater than or equal to r where $[k : k^p] = p^r$. As a further application, we show that the difference between the p -adic K -groups of the integral group ring of a finite group and the p -adic K -groups of a maximal \mathbb{Z} -order in the rational group algebra can be expressed entirely in terms of topological cyclic homology.

Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . In general the induced map of relative K -groups $f_*: K_*(A, I) \rightarrow K_*(B, f(I))$ is not an isomorphism and one defines the bi-relative K -groups so as to form a natural long-exact sequence

$$\dots \rightarrow K_q(A, B, I) \rightarrow K_q(A, I) \xrightarrow{f_*} K_q(B, f(I)) \xrightarrow{\partial} K_{q-1}(A, B, I) \rightarrow \dots$$

It was proved recently by Cortiñas [5] that the trace map

$$K_q(A) \otimes \mathbb{Q} \rightarrow \mathrm{HC}_q^-(A \otimes \mathbb{Q})$$

from rational K -theory of A to negative cyclic homology of $A \otimes \mathbb{Q}$ gives rise to an isomorphism of the associated bi-relative theories. The purpose of the present paper is to prove the following analogous result for K -theory with finite coefficients.

THEOREM A. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . Then the map of bi-relative theories induced from the cyclotomic trace*

$$K_q(A, B, I, \mathbb{Z}/p^v) \rightarrow \{\mathrm{TC}_q^n(A, B, I; p, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, for all integers q .

The first named author was partially supported by COE, the Sloan Foundation, and the National Science Foundation. The second named author was partially supported by the National Science Foundation.

We recall from Bass [1, Thm. XII.8.3] that the common pro-abelian group is zero, for $q \leq 0$. We also remark that Thm. A implies the slightly weaker statement that the map of spectra induced by the cyclotomic trace

$$K(A, B, I) \rightarrow \mathrm{TC}(A, B, I; p) = \operatorname{holim}_n \mathrm{TC}^n(A, B, I; p)$$

becomes a weak equivalence after p -completion. However, for the proof, it is essential that we work with the pro-abelian groups and not pass to the limit. We also remark that it is possible to combine Thm. A and the theorem of Cortiñas [5] and express the integral bi-relative groups $K_q(A, B, I)$ in terms of the topological cyclic homology and negative cyclic homology.

We can use Thm. A to prove the following result. Indeed, Thm. A allows us to reduce to the smooth case, which we proved earlier [9, Thm. 4.2.2]. This application of Thm. A follows Geller, Reid, and Weibel [10, 11], where a weaker version of the theorem of Cortiñas was first stated as a conjecture.

THEOREM B. *Let $X \rightarrow \operatorname{Spec} k$ be any curve over a field of positive characteristic p , and let $0 \leq r \leq \infty$ be given by $[k : k^p] = p^r$. Then for all $q \geq r$, the cyclotomic trace induces an isomorphism of pro-abelian groups*

$$K_q(X, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(X; p, \mathbb{Z}/p^v)\}_{n \geq 1}.$$

We remark again that Thm. B implies the slightly weaker statement that the map of spectra induced by the cyclotomic trace

$$K(X) \rightarrow \mathrm{TC}(X; p) = \operatorname{holim}_n \mathrm{TC}^n(X; p)$$

becomes a weak equivalence after p -completion and passage to $(r - 1)$ -connected covers. The topological cyclic homology groups on the right-hand side of the statement of Thm. B can often be effectively calculated. As an example, we have the following result; compare [10]. The proof is similar to the proof of [13, Thm. E] and [14, Thm. A] and will be given elsewhere.

THEOREM C. *Let k be a regular \mathbb{F}_p -algebra, and let $X = \operatorname{Spec} k[x, y]/(xy)$ be the coordinate axes in the affine plane. Then for all positive integers q , there is a canonical isomorphism of pro-abelian groups*

$$\{\mathrm{TC}_q^n(X; p)\}_{n \geq 1} \xleftarrow{\sim} \bigoplus_{m \geq 1} \mathbf{W}_m \Omega_k^{q-2m}$$

where $\mathbf{W}_m \Omega_k^j$ is the group of big de Rham-Witt j -forms on $\operatorname{Spec} k$ [14, 12].

We wish to point out that, at present, the structure of the K -groups of singular schemes is very far from understood. In particular, there is currently no motivic cohomology theory that can explain the structure of the K -groups that we find here. To illustrate this, let k be a perfect field of positive characteristic p . Then Thm. B and C show that the p -adic K -groups of $X = \operatorname{Spec} k[x, y]/(xy)$ are concentrated in even degrees with the group in positive degree $2m$ canonically isomorphic to the ring $\mathbf{W}_m(k)$ of big Witt vectors of length m in k . In particular, the p -adic K -groups of X cannot be generated from $K_1(X)$ by means of products and transfers as is the case for smooth schemes. We remark that the group $K_2(X)$ was evaluated previously by Dennis and Krusemeyer [7].)

Finally, let $A = \mathbb{Z}[G]$ be the integral group ring of a finite group and $B = \mathfrak{M}$ a maximal \mathbb{Z} -order of the rational group algebra. Then Thm. A shows that, for every

prime p , the difference between the p -adic K -groups of A and B can be expressed entirely in terms of topological cyclic homology. Indeed, if we choose B such that $A \subset B$, then the ideal $I = |G| \cdot B$ is a common ideal of both A and B , and the quotient rings A/I and B/I are finite. The precise relationship is most clearly expressed for G a finite p -group where we obtain the following result. We are grateful to Holger Reich for drawing our attention to this application of Thm. A.

THEOREM D. *Let $\mathbb{Z}[G]$ be the integral group ring of a finite p -group, and let $\mathfrak{M} \subset \mathbb{Q}[G]$ be a maximal \mathbb{Z} -order of the rational group algebra such that $\mathbb{Z}[G] \subset \mathfrak{M}$. Then there is a natural sequence of pro-abelian groups*

$$\cdots \rightarrow K_q(\mathbb{Z}[G], \mathbb{Z}/p^v) \rightarrow \left\{ \begin{array}{c} K_q(\mathfrak{M}, \mathbb{Z}/p^v) \\ \oplus \\ \mathrm{TC}_q^n(\mathbb{Z}[G]; p, \mathbb{Z}/p^v) \end{array} \right\}_{n \geq 1} \rightarrow \{\mathrm{TC}_q^n(\mathfrak{M}; p, \mathbb{Z}/p^v)\}_{n \geq 1} \rightarrow \cdots$$

which is exact for non-negative integers q .

We again remark that Thm. D implies the slightly weaker result that the following square of spectra becomes homotopy-cartesian after p -completion and passage to connective covers.

$$\begin{array}{ccc} K(\mathbb{Z}[G]) & \longrightarrow & \mathrm{TC}(\mathbb{Z}[G]; p) \\ \downarrow & & \downarrow \\ K(\mathfrak{M}) & \longrightarrow & \mathrm{TC}(\mathfrak{M}; p). \end{array}$$

The strength of Thm. D lies in the fact that the maximal order \mathfrak{M} is a regular ring whose K -groups fit in the following localization sequence [23, Thm. 1.17].

$$\cdots \rightarrow \bigoplus_{\ell} K_q(\mathfrak{M}_{\ell}/J_{\ell}) \rightarrow K_q(\mathfrak{M}) \rightarrow K_q(\mathbb{Q}[G]) \rightarrow \cdots$$

The sum on the left-hand side runs over all rational primes ℓ and $J_{\ell} \subset \mathfrak{M}_{\ell}$ is the Jacobson radical in the ℓ -adic completion of \mathfrak{M} . The rings $\mathbb{Q}[G]$ and $\mathfrak{M}_{\ell}/J_{\ell}$ are semi-simple.

We briefly discuss the proof of Thm. A which, in outline, is similar to the proof of the theorem of Cortiñas [5]. The proof uses two key observations which are due to Cuntz and Quillen [6]. The first observation is that it suffices to prove Thm. A in the case where the non-unital associative ring I can be embedded as a two-sided ideal $I \subset T$ of a free unital associative ring. The second observation is that, in this case, the pro-algebra $\{I^m \otimes_{\mathbb{F}_p}\}_{m \geq 1}$ is H -unital. By a generalization of the excision theorem of Suslin-Wodzicki [25, 26], we then show that the pro-abelian groups

$$\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1}$$

are zero, for all integers q . The proof is given in paragraph 1 below. Similarly, we show in paragraph 2 that the pro-abelian groups

$$\{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1}$$

are zero, for all integers q . Hence, if I can be embedded as an ideal of a free unital associative ring, the cyclotomic trace induces an isomorphism of pro-abelian groups

in the top row of the following diagram.

$$\begin{array}{ccc}
\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1} & \xrightarrow{\sim} & \{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1} \\
\downarrow & & \downarrow \\
K_q(A, B, I, \mathbb{Z}/p^v) & \longrightarrow & \{\mathrm{TC}_q^n(A, B, I; p, \mathbb{Z}/p^v)\}_{n \geq 1}.
\end{array}$$

Finally, it follows from a theorem of McCarthy [22] and from [8, Thm. 2.2.1] that also the lower horizontal map is an isomorphism of pro-abelian groups. The details of this argument are given in paragraph 3 below. This completes the outline of the proof of Thm. A.

A pro-object of a category \mathcal{C} is a functor from a directed partially ordered set to the category \mathcal{C} , and a *strict* map between two pro-objects with the same indexing set is a natural transformation. A general map from a pro-object $X = \{X_i\}_{i \in I}$ to a pro-object $Y = \{Y_j\}_{j \in J}$ is an element of the set

$$\mathrm{Hom}_{\mathrm{pro}\text{-}\mathcal{C}}(X, Y) = \lim_J \mathrm{colim}_I \mathrm{Hom}_{\mathcal{C}}(X_i, Y_j).$$

In particular, a pro-object $X = \{X_i\}_{i \in I}$ in a category with a null-object is zero if for all $i \in I$, there exists $i' \geq i$ such that the map $X_{i'} \rightarrow X_i$ is zero.

This paper was written while the authors visited the University of Tokyo. We would like to express our sincere gratitude to the university and to Takeshi Saito for their kind hospitality and for the stimulating atmosphere they provided. Finally, it is a particular pleasure to acknowledge the help we have had from G. Cortiñas. He long ago suggested to us to prove the main theorem of this paper by adapting the integral excision theorem of Suslin [25] to the pro-setting. He also pointed out a gap in our arguments in an earlier version of this paper.

1. K-theory

1.1. In this paragraph we prove the following variant of the excision theorem of Suslin-Wodzicki [26, 25]. Our treatment closely follows Suslin [25]. If I is a non-unital associative ring, we write $\mathbb{Z} \ltimes I$ for the unital associative ring given by the product $\mathbb{Z} \times I$ with multiplication $(a, x) \cdot (a', x') = (aa', ax' + xa' + xx')$.

THEOREM 1.1.1. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . Suppose that for all positive integers q , the pro-abelian group*

$$\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}_{n \geq 1}$$

is zero. Then for all integers q , the map of pro-abelian groups

$$\{K_q(A, I^n, \mathbb{Z}/p^v)\}_{n \geq 1} \rightarrow \{K_q(B, f(I)^n, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism.

We will use a *functorial* version of the plus-construction of Quillen. Hence,

$$\lambda: X \rightarrow X^+$$

will be a natural transformation of functors from the category of pointed spaces and base-point preserving continuous maps to itself. For example, one can use the functorial localization [16] with respect to a pointed space W whose reduced

integral homology is trivial and whose fundamental group admits a non-trivial group homomorphism to $GL(\mathbb{Z})$ [2].

Let A be a unital associative ring, and let $I \subset A$ be a two-sided ideal. Let $GL(I)$ and $\overline{GL}(A/I)$ be the kernel and image of the map $GL(A) \rightarrow GL(A/I)$ induced by the canonical projection. We let $F(A, I)$ be the homotopy fiber of the canonical map $BGL(A)^+ \rightarrow B\overline{GL}(A/I)^+$, and let $X(A, I)$ be the homotopy pull-back of $B\overline{GL}(A/I)$ and $BGL(A)$ over $B\overline{GL}(A/I)^+$. We then have the following commutative diagram of pointed spaces and base-point preserving continuous maps where the rows are fiber sequences.

$$(1.1.2) \quad \begin{array}{ccccc} BGL(I) & \longrightarrow & BGL(A) & \longrightarrow & B\overline{GL}(A/I) \\ \downarrow & & \downarrow & & \parallel \\ F(A, I) & \longrightarrow & X(A, I) & \longrightarrow & B\overline{GL}(A/I) \\ \parallel & & \downarrow & & \downarrow \\ F(A, I) & \longrightarrow & BGL(A)^+ & \longrightarrow & B\overline{GL}(A/I)^+. \end{array}$$

We consider the relative Serre spectral sequence

$$(1.1.3) \quad \begin{aligned} E_{s,t}^2 &= H_s(\overline{GL}(A/I), H_t(F(A, I), BGL(I), \mathbb{Z}/p)) \\ &\Rightarrow H_{s+t}(X(A, I), BGL(A), \mathbb{Z}/p) \end{aligned}$$

which converges to zero. Indeed, both of the maps $BGL(A) \rightarrow BGL(A)^+$ and $X(A, I) \rightarrow BGL(A)^+$ induce isomorphisms on homology, and hence so does the map $BGL(A) \rightarrow X(A, I)$. Hence, the relative homology groups, defined as the reduced homology groups of the mapping cone, are zero. The diagram (1.1.2) and the spectral sequence (1.1.3) are both functorial in the pair $I \subset A$. Hence, if we consider the pro-ideal $\{I^n\}_{n \geq 1}$ of powers of the ideal $I \subset A$, we obtain a spectral sequence of pro-abelian groups.

LEMMA 1.1.4. *If the pro-abelian group $\{H_t(F(A, I^n), BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1}$ is zero, for $0 \leq t < p$, then the pro-abelian group*

$$\{H_s(\overline{GL}(A/I^n), H_p(F(A, I), BGL(I)), \mathbb{Z}/p)\}_{n \geq 1}$$

is zero, for $0 \leq s \leq 1$. If, in addition, the canonical map

$$\{H_{p-1}(BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1} \rightarrow \{H_0(\overline{GL}(A/I^n), H_{p-1}(BGL(I^n), \mathbb{Z}/p))\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, then the pro-abelian group

$$\{H_p(F(A, I^n), BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is zero.

PROOF. We shall omit the \mathbb{Z}/p -coefficients from the homology throughout the proof. The first claim follows immediately from the spectral sequence. To prove the second claim, we show that in the exact sequence of pro-abelian groups

$$\{H_p(F(A, I^n))\} \rightarrow \{H_p(F(A, I^n), BGL(I^n))\} \rightarrow \{H_{p-1}(BGL(I^n))\},$$

both maps are equal to zero. We first show that the right-hand map is zero, or equivalently, that in the following exact sequence of pro-abelian groups, the right-hand map is injective.

$$\{H_p(F(A, I^n), BGL(I^n))\} \rightarrow \{H_{p-1}(BGL(I^n))\} \rightarrow \{H_{p-1}(F(A, I^n))\} \rightarrow 0.$$

The right-hand map is surjective by the hypothesis of the lemma, and hence, the induced sequence of $\{\overline{GL}(A/I^n)\}$ -coinvariants is again exact. By the first statement of the lemma, the pro-abelian group

$$\{H_0(\overline{GL}(A/I^n), H_p(F(A, I^n), BGL(I^n)))\}$$

is zero, and therefore the map of pro-abelian groups

$$\{H_0(\overline{GL}(A/I^n), H_{p-1}(BGL(I^n)))\} \rightarrow \{H_0(\overline{GL}(A/I^n), H_{p-1}(F(A, I^n)))\}$$

is injective. We consider the diagram of pro-abelian groups.

$$\begin{array}{ccc} \{H_{p-1}(BGL(I^n))\} & \longrightarrow & \{H_{p-1}(F(A, I^n))\} \\ \downarrow \sim & & \downarrow \\ \{H_0(\overline{GL}(A/I^n), H_{p-1}(BGL(I^n)))\} & \hookrightarrow & \{H_0(\overline{GL}(A/I^n), H_{p-1}(F(A, I^n)))\}. \end{array}$$

The left-hand vertical map is an isomorphism by assumption. Hence, the upper horizontal map is injective as desired.

It remains to show that the left-hand map in the sequence at the beginning of the proof is zero. This is equivalent to showing that in the following exact sequence of pro-abelian groups, the left-hand map is surjective.

$$\{H_p(BGL(I^n))\} \rightarrow \{H_p(F(A, I^n))\} \rightarrow \{H_p(F(A, I^n), BGL(I^n))\} \rightarrow 0.$$

The surjectivity of the right-hand map was proved above. The induced sequence of $\{\overline{GL}(A/I^n)\}$ -coinvariants is again exact, and by the first statement of the lemma, the group of $\{\overline{GL}(A/I^n)\}$ -coinvariants of the right-hand term is zero. Hence, the map of pro-abelian groups

$$\{H_0(\overline{GL}(A/I^n), H_p(BGL(I^n)))\} \rightarrow \{H_0(\overline{GL}(A/I^n), H_p(F(A, I^n)))\}$$

is surjective. We now consider the following diagram of pro-abelian groups.

$$\begin{array}{ccc} \{H_p(BGL(I^n))\} & \longrightarrow & \{H_p(F(A, I^n))\} \\ \downarrow & & \downarrow \sim \\ \{H_0(\overline{GL}(A/I^n), H_p(BGL(I^n)))\} & \twoheadrightarrow & \{H_0(\overline{GL}(A/I^n), H_p(F(A, I^n)))\}. \end{array}$$

The left-hand vertical map and the lower horizontal map are surjective, and we claim that the right-hand vertical map is an isomorphism. Indeed, since $X(A, I^n)$ is a homotopy pull-back, the action of $\overline{GL}(A/I^n) = \pi_1(B\overline{GL}(A/I^n))$ on $H_p(F(A, I^n))$ is induced from the action of $\pi_1(B\overline{GL}(A/I^n)^+)$ on $H_p(F(A, I^n))$. But

$$F(A, I^n) \rightarrow BGL(A)^+ \rightarrow B\overline{GL}(A/I^n)^+$$

is a fiber sequence of infinite loop spaces, and therefore the fundamental group of the base acts trivially on the homology of the fiber [28]. This proves that the top horizontal map in the diagram above is surjective as desired. \square

The group of column vectors $M_{\infty,1}(I^n)$ is naturally a left $GL(I^n)$ -module, so we can form the semi-direct product group $GL(I^n) \ltimes M_{\infty,1}(I^n)$. We shall show in Prop. 1.2.7 below that the map induced by the canonical inclusion

$$\{H_q(GL(I^n), \mathbb{Z}/p)\} \rightarrow \{H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), \mathbb{Z}/p)\}$$

is an isomorphism of pro-abelian groups. But first we show that this implies the following result.

PROPOSITION 1.1.5. *Let A be a unital associative ring, let $I \subset A$ be a two-sided ideal, and suppose that for all positive integers q , the pro-abelian group*

$$\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}_{n \geq 1}$$

is zero. Then for all integers q , the pro-abelian group

$$\{H_q(F(A, I^n), BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is zero.

PROOF. For notational convenience we omit the \mathbb{Z}/p -coefficients for the homology throughout the proof. We shall show that the hypothesis implies that for all integers q , the canonical map of pro-abelian groups

$$\{H_q(BGL(I^n))\}_{n \geq 1} \rightarrow \{H_0(\overline{GL}(A/I^n), H_q(BGL(I^n)))\}_{n \geq 1}$$

is an isomorphism. The proposition then follows from Lemma 1.1.4 by induction on q starting from the trivial case $q = 0$.

It suffices, by the proof of [26, Cor. 1.6], to show that for all $n \geq 1$, there exists $m \geq n$ such that the image of

$$\iota_{m,n*} : H_q(GL(I^m)) \rightarrow H_q(GL(I^n))$$

is $GL(\mathbb{Z})$ -invariant. To this end we consider the following homomorphisms.

$$H_q(GL(I^n)) \xrightarrow{\sigma_*} H_q(GL(I^n) \ltimes M_{\infty,1}(I^n)) \xrightarrow{j_*} H_q(GL(I^n)).$$

The right-hand map is induced from the embedding

$$j(\alpha, v) = \begin{pmatrix} 1 & 0 \\ v & \alpha \end{pmatrix}.$$

By *loc.cit.*, Lemma 1.4, the composite map $\varphi_* = j_* \circ \sigma_*$ is injective. Hence, it suffices to show that the image of the composite map

$$H_q(GL(I^m)) \xrightarrow{\iota_{m,n*}} H_q(GL(I^n)) \xrightarrow{\varphi_*} H_q(GL(I^n))$$

is invariant under the adjoint action by the subgroup

$$\begin{pmatrix} 1 & 0 \\ 0 & GL(\mathbb{Z}) \end{pmatrix} \subset GL(\mathbb{Z}).$$

In effect, we will show that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the adjoint action by the full group $GL(\mathbb{Z})$. The group $GL(\mathbb{Z})$ is generated by the elementary matrices $e_{k,1}(1)$ and $e_{1,k}(1)$, $k \geq 1$, and the diagonal matrix $\mathrm{diag}(-1, 1, 1, \dots)$. It is clear that the image of φ_* is invariant under the action of the diagonal matrix $\mathrm{diag}(-1, 1, 1, \dots)$. It remains to show that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant also under the action of the elementary matrices $e_{k,1}(1)$ and $e_{1,k}(1)$.

The elementary matrices $e_{k,1}(1)$ are in the image of the composite

$$M_{\infty,1}(\mathbb{Z}) \xrightarrow{\gamma} GL(\mathbb{Z}) \ltimes M_{\infty,1}(\mathbb{Z}) \xrightarrow{j} GL(\mathbb{Z}),$$

where the left-hand map takes β to $(1, \beta)$. We show in Prop. 1.2.7 below there exists $m \geq n$ such that the following map of relative homology groups is zero.

$$\iota_{m,n*}: H_q(GL(I^m) \ltimes M_{\infty,1}(I^m), GL(I^m)) \rightarrow H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), GL(I^n)).$$

Let $\beta \in GL(\mathbb{Z})$ be an element in the image of $j \circ \gamma$ and consider the following diagram.

$$\begin{array}{ccc} H_q(GL(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n)) \\ \downarrow \sigma_* & & \downarrow \sigma_* \\ H_q(GL(I^m) \ltimes M_{\infty,1}(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n) \ltimes M_{\infty,1}(I^n)) \\ \sim \downarrow \text{Ad}(\beta)_* & & \sim \downarrow \text{Ad}(\beta)_* \\ H_q(GL(I^m) \ltimes M_{\infty,1}(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n) \ltimes M_{\infty,1}(I^n)) \\ \downarrow \pi_* & & \downarrow \pi_* \\ H_q(GL(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n)). \end{array}$$

The vertical compositions both are the identity homomorphism and, by the choice of $m \geq n$, the image the second horizontal homomorphism $\iota_{m,n*}$ is contained in the image the top right hand vertical homomorphism σ_* . This shows that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the action of β in the image of $j \circ \gamma$. In particular, the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the action of the elementary matrices $e_{k,1}(1)$. One shows in a similar way that there exists (a possibly larger) $m \geq n$ such that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the action of the elementary matrices $e_{1,k}(1)$. This completes the proof. \square

PROOF OF THM. 1.1.1. It suffices by simple induction to treat the case $v = 1$. Let $f: A \rightarrow B$ be as in the statement. By Prop. 1.1.5, the induced map

$$\{H_q(F(A, I^n), \mathbb{Z}/p)\}_{n \geq 1} \rightarrow \{H_q(F(B, f(I)^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, and [24, Cor. 5.8] then shows that also the induced map

$$\{\pi_q(F(A, I^n), \mathbb{Z}/p)\}_{n \geq 1} \rightarrow \{\pi_q(F(B, f(I)^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups. \square

1.2. The main result of this section is Prop. 1.2.7 below which was used in the proof of Prop. 1.1.5 above. We begin with some general theory.

Let A be a unital associative ring, and let $I \subset A$ be a two-sided ideal. Then the powers $\{I^n\}_{n \geq 1}$ form a pro-associative ring without unit. We will say that a pro-abelian group $\{M_n\}_{n \geq 1}$ is a left module over $\{I^n\}_{n \geq 1}$ if for all $n \geq 1$, M_n is a

left I^n -module and if for all $m \geq n \geq 1$, the following diagram commutes.

$$\begin{array}{ccc} I^m \otimes M_m & \xrightarrow{\mu_m} & M_m \\ \downarrow \iota_{m,n} \otimes \lambda_{m,n} & & \downarrow \lambda_{m,n} \\ I^n \otimes M_n & \xrightarrow{\mu_n} & M_n. \end{array}$$

By a homomorphism from a left $\{I^n\}$ -module $\{M_n\}$ to a left $\{I^n\}$ -module $\{M'_n\}$ we mean a strict map of pro-abelian groups $f: \{M_n\} \rightarrow \{M'_n\}$ such that for all $n \geq 1$, the map $f_n: M_n \rightarrow M'_n$ is an I^n -module homomorphism. We will say that a left $\{I^n\}$ -module $\{P_n\}$ is *pseudo-free* if for all $n \geq 1$, there are I^n -module isomorphisms

$$\varphi_n: I^n \otimes L_n \xrightarrow{\sim} P_n$$

with L_n a free abelian group, and if for all $m \geq n \geq 1$, there are maps of abelian groups $\sigma_{m,n}: L_m \rightarrow L_n$ such that the following diagrams commute.

$$\begin{array}{ccc} I^m \otimes L_m & \xrightarrow{\varphi_m} & P_m \\ \downarrow \iota \otimes \sigma_{m,n} & & \downarrow \lambda_{m,n} \\ I^n \otimes L_n & \xrightarrow{\varphi_n} & P_n. \end{array}$$

A homomorphism $f: \{P_n\} \rightarrow \{M_n\}$ from a pseudo-free $\{I^n\}$ -module to an arbitrary $\{I^n\}$ -module is said to be

(i) *special* if there exists a strict map of pro-abelian groups $g: \{L_n\} \rightarrow \{M_n\}$ such that for all $n \geq 1$, the composite

$$I^n \otimes L_n \xrightarrow{\varphi_n} P_n \xrightarrow{f_n} M_n$$

maps $a \otimes x$ to $ag_n(x)$.

(ii) *p^v -special* if the $\{I^n\}$ -module $\{P_n\}$ admits a strict direct sum decomposition

$$\iota' \oplus \iota'': \{P'_n\} \oplus \{P''_n\} \xrightarrow{\sim} \{P_n\}$$

such that $\{P'_n\}$ and $\{P''_n\}$ are pseudo-free $\{I^n\}$ -modules, such that $f \circ \iota'$ is special, and such that $f \circ \iota'' = p^v g$, for some homomorphism $g: \{P''_n\} \rightarrow \{M_n\}$.

LEMMA 1.2.1. *Let $\{P'_n\}$ be a left $\{I^n\}$ -module. Then there exists a special homomorphism $f: \{P_n\} \rightarrow \{P'_n\}$ from a pseudo-free left $\{I^n\}$ -module $\{P_n\}$ such that the image of f_n is equal to $I^n \cdot P'_n \subset P'_n$.*

PROOF. Let $\mathbb{Z}\{P'_n\}$ be the free abelian group generated by the set of elements of P'_n . We define P_n be the extended left I^n -module

$$P_n = I^n \otimes \bigoplus_{k \geq n} \mathbb{Z}\{P'_k\},$$

and let $\lambda_{m,n}: P_m \rightarrow P_n$ be the map

$$I^m \otimes \bigoplus_{k \geq m} \mathbb{Z}\{P'_k\} \xrightarrow{\iota_{m,n} \otimes \text{incl}} I^n \otimes \bigoplus_{k \geq n} \mathbb{Z}\{P'_k\}.$$

Then $\{P_n\}$ is a pseudo-free $\{I^n\}$ -module and the canonical map $f: \{P_n\} \rightarrow \{P'_n\}$ is a special homomorphism such that the image of f_n is equal to $I^n \cdot P'_n \subset P'_n$. \square

Let $\{P_n\}$ be a pseudo-free $\{I^n\}$ -module. An augmented complex of $\{I^n\}$ -modules over $\{P_n\}$ is, by definition, a sequence of $\{I^n\}$ -modules and homomorphisms

$$\dots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\}$$

such that $d_n \circ d_n$ and $\epsilon_n \circ d_n$ are equal to zero, for all $n \geq 1$. We say that the augmented complex $\epsilon: \{C_{n,*}\} \rightarrow \{P_n\}$ is

(i) *special* if the $\{C_{n,q}\}$ are pseudo-free, if ϵ is special, and if the homology of the augmented complex $\epsilon_n: C_{n,*} \rightarrow P_n$ is annihilated by I^n .

(ii) *p^v -special* if the $\{C_{n,q}\}$ are pseudo-free, if ϵ is p^v -special, and if the homology of the augmented complex $\epsilon_n: C_{n,*} \rightarrow P_n$ is annihilated by both I^n and p^v .

COROLLARY 1.2.2. *Let $\{P_n\}$ be an pseudo-free $\{I^n\}$ -module. Then there exists both a special augmented complex and a p^v -special augmented complex over $\{P_n\}$.*

We remark that the identity homomorphism of a pseudo-free module is usually not special, and hence, does not constitute an augmented special complex.

PROOF. It is proved by easy induction based on Lemma 1.2.1 that there exists a special augmented complex

$$\dots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\}.$$

Then the total complex of the double-complex

$$\begin{array}{ccccccc} \dots & \xrightarrow{d} & \{C_{n,2}\} & \xrightarrow{d} & \{C_{n,1}\} & \xrightarrow{d} & \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\} \\ & & \downarrow p^v & & \downarrow p^v & & \downarrow p^v \\ \dots & \xrightarrow{d} & \{C_{n,2}\} & \xrightarrow{d} & \{C_{n,1}\} & \xrightarrow{d} & \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\}. \end{array}$$

is a p^v -special augmented complex. \square

PROPOSITION 1.2.3. *Let $I \subset A$ be a two-sided ideal of a free unital associative ring, and suppose that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \times (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}$ are zero, for all positive integers q . Then every p^v -special augmented complex*

$$\dots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\},$$

is exact as an augmented complex of pro-abelian groups.

PROOF. By induction, the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \times (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}_{n \geq 1}$ are zero, for all $v \geq 1$. We show by induction on $q \geq -1$ that the following complex of pro-abelian groups has zero homology.

$$\dots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{d} \{C_{n,-1}\} = \{P_n\}$$

We let $Z_{n,q}$ and $B_{n,q-1}$ be the kernel and image of $d_n: C_{n,q} \rightarrow C_{n,q-1}$ and note that since both I^n and p^v annihilate $H_q(C_{n,*}) = Z_{n,q}/B_{n,q}$, we have

$$I^n Z_{n,q} + p^v Z_{n,q} \subset B_{n,q}.$$

We first consider $q = -1$. Then $Z_{n,-1} = P_n$, and hence $I^n P_n + p^v P_n \subset B_{n,-1}$. But, by the definition of a special homomorphism, $B_{n,-1} \subset I^n P_n + p^v P_n$, and therefore

$B_{n,-1} = I^n P_n + p^v P_n$. Hence,

$$\begin{aligned} H_{-1}(C_{n,*}) &\xleftarrow{\sim} \operatorname{Tor}_0^{\mathbb{Z} \ltimes (I^n)}(P_n, \mathbb{Z}/p^v) \xleftarrow{\sim} \operatorname{Tor}_0^{\mathbb{Z} \ltimes (I^n)}(I^n \otimes M_n, \mathbb{Z}/p^v) \\ &\xleftarrow{\sim} \operatorname{Tor}_0^{\mathbb{Z} \ltimes (I^n)}(I^n, \mathbb{Z}/p^v) \otimes M_n \xleftarrow{\sim} \operatorname{Tor}_1^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p^v) \otimes M_n. \end{aligned}$$

It follows that the pro-abelian group $\{H_{-1}(C_{n,*})\}$ is zero as desired.

Next, we assume that the pro-abelian group $\{H_i(C_{n,*})\}$ is zero, for $i < q$, and show that the pro-abelian group $\{H_q(C_{n,*})\}$ is zero. The assumption is equivalent to the statement that the following sequence of pro-abelian groups is exact.

$$0 \rightarrow \{Z_{n,q}\} \rightarrow \{C_{n,q}\} \rightarrow \cdots \rightarrow \{C_{n,0}\} \rightarrow \{P_n\} \rightarrow 0.$$

This gives rise to the following spectral sequence of pro-abelian groups.

$$\begin{aligned} E_{s,t}^1 &= \begin{cases} \{\operatorname{Tor}_t^{\mathbb{Z} \ltimes (I^n)}(C_{n,s}, \mathbb{Z}/p^v)\}, & \text{if } 0 \leq s \leq q, \\ \{\operatorname{Tor}_t^{\mathbb{Z} \ltimes (I^n)}(Z_{n,q}, \mathbb{Z}/p^v)\}, & \text{if } s = q+1, \\ \{0\}, & \text{if } s > q+1, \end{cases} \\ &\Rightarrow \{\operatorname{Tor}_{s+t}^{\mathbb{Z} \ltimes (I^n)}(P_n, \mathbb{Z}/p^v)\}. \end{aligned}$$

Since $\{C_{n,s}\}$ is pseudo-free, we have

$$\begin{aligned} \operatorname{Tor}_t^{\mathbb{Z} \ltimes (I^n)}(C_{n,s}, \mathbb{Z}/p^v) &\xleftarrow{\sim} \operatorname{Tor}_t^{\mathbb{Z} \ltimes (I^n)}(I^n \otimes M_{n,s}, \mathbb{Z}/p^v) \\ &\xleftarrow{\sim} \operatorname{Tor}_t^{\mathbb{Z} \ltimes (I^n)}(I^n, \mathbb{Z}/p^v) \otimes M_{n,s} \xleftarrow{\sim} \operatorname{Tor}_{t+1}^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p^v) \otimes M_{n,s}. \end{aligned}$$

It follows that the pro-abelian groups $E_{s,t}^1$ are zero, for $0 \leq s \leq q$. Hence the edge-homomorphism of the spectral sequence

$$\{\operatorname{Tor}_{q+1}^{\mathbb{Z} \ltimes (I^n)}(P_n, \mathbb{Z}/p^v)\} \rightarrow \{\operatorname{Tor}_0^{\mathbb{Z} \ltimes (I^n)}(Z_{n,q}, \mathbb{Z}/p^v)\} = \{Z_{n,q}/(I^n Z_{n,q} + p^v Z_{n,q})\}$$

is an isomorphism of pro-abelian groups, and since the pro-abelian group on the left is zero, so is the one on the right. Finally, since $I^n Z_{n,q} + p^v Z_{n,q} \subset B_{n,q}$, the group $Z_{n,q}/(I^n Z_{n,q} + p^v Z_{n,q})$ surjects onto the group $Z_{n,q}/B_{n,q} = H_q(C_{n,*})$, and hence, the pro-abelian group $\{H_q(C_{n,*})\}$ is zero as stated. \square

PROPOSITION 1.2.4. *Let $I \subset A$ be a two-sided ideal of a unital associative, and suppose that the pro-abelian groups $\{\operatorname{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}$ are zero, for all positive integers q . Let F_i , $i \geq 0$, be a sequence of possibly non-additive functors F_i , $i \geq 0$, from the category of $\{I^n\}$ -modules to the category of pro-abelian groups that satisfies the following conditions (i)–(iii).*

(i) *For every $\{I^n\}$ -module $\{P_n\}$, the pro-abelian group $F_0(\{P_n\})$ is zero.*

(ii) *If $\epsilon: \{P_n[-]\} \rightarrow \{P_n\}$ is an augmented simplicial $\{I^n\}$ -module such that the associated augmented complex of pro-abelian groups is exact, then there is a spectral sequence of pro-abelian groups*

$$E_{s,t}^1 = F_t(\{P_n[s]\}) \Rightarrow F_{s+t}(\{P_n\})$$

such that the edge-homomorphism $F_t(\{P_n[0]\}) \rightarrow F_t(\{P_n\})$ is the map induced by the augmentation $\epsilon: \{P_n[0]\} \rightarrow \{P_n\}$.

(iii) *For every $i \geq 0$, there exists $v \geq 0$ such that the map of pro-abelian groups*

$$f_*: F_i(\{P_n\}) \rightarrow F_i(\{P'_n\})$$

induced by a p^v -special homomorphism $f: \{P_n\} \rightarrow \{P'_n\}$ is zero.

Then for every pseudo-free $\{I^n\}$ -module $\{P_n\}$ and for every $i \geq 0$, the pro-abelian group $F_i(\{P_n\})$ is zero.

PROOF. We prove the statement by induction on $i \geq 0$. The case $i = 0$ is trivial, so we assume the statement for $i - 1$ and prove it for i . We choose v such that for all $0 \leq t \leq i$, the map of pro-abelian groups

$$f_*: F_t(\{P_n\}) \rightarrow F_t(\{P'_n\})$$

induced by a p^v -special homomorphism $f: \{P_n\} \rightarrow \{P'_n\}$ is zero. Let $\{P_n\}$ be a pseudo-free $\{I^n\}$ -module and choose a p^v -special complex $\epsilon: \{C_{n,*}\} \rightarrow \{P_n\}$ over $\{P_n\}$. By Prop. 1.2.3 this augmented complex is exact as an augmented complex of pro-abelian groups. We let $\epsilon: \{P_n[-]\} \rightarrow \{P_n\}$ be the augmented simplicial $\{I^n\}$ -module obtained from the augmented complex $\epsilon: \{C_{n,*}\} \rightarrow \{P_n\}$ by the Dold-Kan correspondance [29]. By assumption, we then have the following spectral sequence of pro-abelian groups.

$$E_{s,t}^1 = F_t(\{P_n[s]\}) \Rightarrow F_{s+t}(\{P_n\}).$$

By the inductive hypothesis, the pro-abelian group $E_{s,t}^1$ is zero, for $0 \leq t < i$, and hence, the edge-homomorphism

$$F_i(\{P_n[0]\}) \twoheadrightarrow F_i(\{P_n\})$$

is an epimorphism of pro-abelian groups. But the edge-homomorphism is equal to the map induced by the special homomorphism $\epsilon: \{P_n[0]\} \rightarrow \{P_n\}$ and therefore is zero by assumption. This completes the proof. \square

If a group G acts from the left on a group H , the semi-direct product group $G \ltimes H$ is defined to be the set $G \times H$ with multiplication

$$(g, h) \cdot (g', h') = (gg', (g'^{-1}h)h').$$

The projection $\pi: G \ltimes H \rightarrow G$ and the map $\sigma: G \rightarrow G \ltimes H$ given by $\sigma(g) = (g, 1)$ are both group-homomorphisms and $\pi \circ \sigma = \text{id}$. Hence, the homology of the semi-direct product group decomposes canonically as a direct sum

$$H_q(G \ltimes H) \xleftarrow{\sim} H_q(G) \oplus H_q(G \ltimes H, G).$$

PROPOSITION 1.2.5. Let $\{G_n\}$ be a pro-group, and let $\epsilon: \{H_n[-]\} \rightarrow \{H_n\}$ be a strict augmented simplicial pro-group with a strict left $\{G_n\}$ -action. Suppose that the associated complex of pro-abelian groups

$$\cdots \rightarrow \{\mathbb{Z}\{H_n[2]\}\} \rightarrow \{\mathbb{Z}\{H_n[1]\}\} \rightarrow \{\mathbb{Z}\{H_n[0]\}\} \xrightarrow{\epsilon} \{\mathbb{Z}\{H_n\}\} \rightarrow 0$$

is exact. Then there is a natural spectral sequence of pro-abelian groups

$$E_{s,t}^1 = \{H_t(G_n \ltimes H_n[s], G_n, \mathbb{Z}/p)\} \Rightarrow \{H_{s+t}(G_n \ltimes H_n, G_n, \mathbb{Z}/p)\},$$

and the edge-homomorphism

$$\{H_t(G_n \ltimes H_n[0], G_n, \mathbb{Z}/p)\} \rightarrow \{H_t(G_n \ltimes H_n, G_n, \mathbb{Z}/p)\}$$

is equal to the map induced from ϵ .

PROOF. Let $\epsilon: \mathbb{Z}\{EG[-]\} \rightarrow \mathbb{Z}$ be the augmented simplicial free abelian group generated by the standard augmented simplicial model $EG[-] \rightarrow *$ of a contractible space with a free G -action. The associated augmented complex

$$\epsilon: \mathbb{Z}\{EG[-]\}_* \rightarrow \mathbb{Z}$$

is a natural resolution of \mathbb{Z} by free (right) $\mathbb{Z}[G]$ -modules. Suppose that M_* is a complex of left G -modules. We define $C_*(G; M_*)$ to be the total complex of the double-complex $\mathbb{Z}\{EG[-]\}_* \otimes_{\mathbb{Z}[G]} M_*$ and the hyper-homology of G with coefficients in M_* to be the homology

$$H_*(G; M_*) = H_*(C_*(G; M_*)).$$

There is a functorial spectral sequence

$$(1.2.6) \quad E_{s,t}^2 = H_s(G; H_t(M_*)) \Rightarrow H_{s+t}(G; M_*)$$

with $H_t(M_*)$ considered as complex concentrated in degree zero.

If G acts from the left on a group H , and if M_* is a complex of left $G \ltimes H$ -modules, then there is a canonical isomorphism of complexes

$$C_*(G \ltimes H; M_*) \xleftarrow{\sim} C_*(G; C_*(H; M_*))$$

where the G -action on $C_*(H; M_*)$ is induced from the actions by G on H and M_* . Moreover, if $\tilde{C}_*(H; M_*)$ denotes the reduced complex defined as the degree-wise cokernel of the canonical map $C_*(\{1\}; M_*) \rightarrow C_*(H; M_*)$, then there is a canonical isomorphism of complexes

$$C_*(G \ltimes H, G; M_*) \xleftarrow{\sim} C_*(G; \tilde{C}_*(H; M_*)).$$

We shall only consider this isomorphism in the case where $M_* = \mathbb{Z}/p$ is the complex that consists of the trivial $G \times H$ -module \mathbb{Z}/p concentrated in degree zero.

Let $\epsilon: H[-] \rightarrow H$ be an augmented simplicial group with a left G -action, and let $\tilde{C}_*(H[-]; \mathbb{Z}/p)_*$ denote the total-complex of the double-complex associated with the simplicial complex $\tilde{C}_*(H[-]; \mathbb{Z}/p)$. Then the filtration in the simplicial direction gives rise to a functorial spectral sequence of hyper-homology groups

$$E_{s,t}^1 = H_t(G; \tilde{C}_*(H[s]; \mathbb{Z}/p)) \Rightarrow H_{s+t}(G; \tilde{C}_*(H[-]; \mathbb{Z}/p)_*).$$

Suppose now that $\{G_n\}$ and $\epsilon: \{H_n[-]\} \rightarrow \{H_n\}$ are as in the statement. Then we obtain a spectral sequence of (strict) pro-abelian groups

$$E_{s,t}^1 = \{H_t(G_n; \tilde{C}_*(H_n[s]; \mathbb{Z}/p))\} \Rightarrow \{H_{s+t}(G_n; \tilde{C}_*(H_n[-]; \mathbb{Z}/p)_*)\}.$$

The augmentation induces a (strict) map of pro-abelian groups

$$\epsilon_*: \{H_q(G_n; \tilde{C}_*(H_n[-]; \mathbb{Z}/p)_*)\} \rightarrow \{H_q(G_n; \tilde{C}_*(H_n; \mathbb{Z}/p))\},$$

and we claim that this map is an isomorphism. Indeed, the hypothesis of the proposition implies that the map induced by the augmentation

$$\epsilon_*: \{H_t(\tilde{C}_*(H_n[-]; \mathbb{Z}/p)_*)\} \rightarrow \{\tilde{H}_t(H_n; \mathbb{Z}/p)\}$$

is an isomorphism of pro-abelian groups for all t . The claim then follows from the spectral sequence (1.2.6) which is located in the first quadrant. This proves that there exists a spectral sequence as stated in the proposition. Finally, it is clear from the construction that the edge-homomorphism is as stated. \square

Let A be a unital associative ring, let $I \subset A$ be a two-sided ideal, and let $\{P_n\}$ be a left $\{I^n\}$ -module. Then the group of column vectors $M_{\infty,1}(P_n)$ is naturally a left $GL(I^n)$ -module, and we consider the canonical inclusion of $GL(I^n)$ in the semi-direct product $GL(I^n) \ltimes M_{\infty,1}(P_n)$. The relative homology groups form a sequence of non-additive functors

$$F_i(\{P_n\}) = \{H_i(GL(I^n) \ltimes M_{\infty,1}(P_n), GL(I^n)), \mathbb{Z}/p\}$$

from the category of left $\{I^n\}$ -modules to the category of pro-abelian groups. This sequence of functors trivially satisfies the condition (i) of Prop. 1.2.4 and it follows from Prop. 1.2.5 that it also satisfies the condition (ii). Finally, it follows from [25, Cor. 6.6] that this sequence of functors also satisfies the condition (iii). Hence we obtain the following result that was used in the proof of Prop. 1.1.5 above.

PROPOSITION 1.2.7. *Let $I \subset A$ be a two-sided ideal of a unital associative ring, and suppose that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}$ are zero, for all positive integers q . Then the map induced by the canonical inclusion*

$$\{H_q(GL(I^n), \mathbb{Z}/p)\} \rightarrow \{H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), \mathbb{Z}/p)\}$$

is an isomorphism of pro-abelian groups, for all $q \geq 0$.

PROOF. Indeed, the module $\{I^n\}$ is pseudo-free, and hence, Prop. 1.2.4 shows that the pro-abelian groups

$$\{H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), GL(I^n), \mathbb{Z}/p)\}$$

are zero, for all $q \geq 0$. \square

2. Topological cyclic homology

2.1. This paragraph is devoted to the proof of the following result.

THEOREM 2.1.1. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . Suppose that I can be embedded as a two-sided ideal $I \subset T$ of a free unital associative ring. Then the induced map*

$$\{\mathrm{TC}_q^n(A, I^m; p, \mathbb{Z}/p^v)\}_{m \geq 1} \rightarrow \{\mathrm{TC}_q^n(B, f(I)^m; p, \mathbb{Z}/p^v)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups, for all integers q and positive integers n .

We first explain that theorem 2.1.1 follows from the analogous result for topological Hochschild homology. The reader is referred to [15] and [12] for further details on topological Hochschild and topological cyclic homology. Associated with the unital associative ring A one has the topological Hochschild spectrum $T(A)$ which is a object of the \mathbb{T} -stable homotopy category. The equivariant homotopy groups

$$\mathrm{TR}_q^n(A; p) = [S^q \wedge (\mathbb{T}/C_{p^{n-1}})_+, T(A)]_{\mathbb{T}}$$

come equipped with an additional *restriction* map

$$\mathrm{TR}_q^n(A; p) \xrightarrow{R} \mathrm{TR}_q^{n-1}(A; p)$$

whose definition uses the additional *cyclotomic* structure of $T(A)$. We consider the family of groups $\{\mathrm{TR}_q^n(A; p)\}_{n \geq 1}$ as a pro-abelian group with the restriction map as the structure map. The group $\mathrm{TR}_q^1(A; p)$ is equal to the q th topological Hochschild group $\mathrm{THH}_q(A)$. In general, there is are natural long-exact sequences

$$(2.1.2) \quad \cdots \rightarrow \mathbb{H}_q(C_{p^{n-1}}, T(A)) \rightarrow \mathrm{TR}_q^n(A; p) \xrightarrow{R} \mathrm{TR}_q^{n-1}(A; p) \rightarrow \cdots$$

and natural spectral sequences

$$(2.1.3) \quad E_{s,t}^2 = H_s(C_{p^{n-1}}, \mathrm{THH}_t(A)) \Rightarrow \mathbb{H}_{s+t}(C_{p^{n-1}}, T(A)).$$

The canonical projection $\mathbb{T}/C_{p^{n-2}} \rightarrow \mathbb{T}/C_{p^{n-1}}$ induces the *Frobenius* map of equivariant homotopy groups

$$F: \mathrm{TR}_q^n(A; p) \rightarrow \mathrm{TR}_q^{n-1}(A; p),$$

and the topological cyclic homology groups $\mathrm{TC}_q^n(A; p)$ are defined so as to form a long-exact sequence

$$(2.1.4) \quad \dots \rightarrow \mathrm{TC}_q^n(A; p) \rightarrow \mathrm{TR}_q^n(A; p) \xrightarrow{R-F} \mathrm{TR}_q^{n-1}(A; p) \xrightarrow{\partial} \mathrm{TC}_{q-1}^n(A; p) \rightarrow \dots$$

As an immediate consequence of the long-exact sequences (2.1.2) and (2.1.4) and the spectral sequence (2.1.3) we note that if a ring-homomorphism $f: A \rightarrow A'$ induces an isomorphism $f_*: \mathrm{THH}_*(A) \xrightarrow{\sim} \mathrm{THH}_*(A')$, then it induces an isomorphism

$$f_*: \mathrm{TC}_q^n(A; p) \xrightarrow{\sim} \mathrm{TC}_q^n(A'; p),$$

for all $n \geq 1$.

Let A be a unital associative ring, and let $I \subset A$ a two-sided ideal. Then there is an exact triangle in the \mathbb{T} -stable category

$$(2.1.5) \quad T(A, I) \rightarrow T(A) \rightarrow T(A/I) \xrightarrow{\partial} T(A, I)[-1].$$

We recall below a particular construction of this sequence by Bökstedt and Madsen [3, Prop. 10.2] which has the additional property that each term in the sequence is *cyclotomic*. This shows that there are induced long-exact sequences

$$\dots \rightarrow \mathrm{TC}_q^n(A, I; p) \rightarrow \mathrm{TC}_q^n(A; p) \rightarrow \mathrm{TC}_q^n(A/I; p) \xrightarrow{\partial} \mathrm{TC}_{q-1}^n(A, I; p) \rightarrow \dots$$

and relative versions of the long-exact sequences (2.1.2) and (2.1.4) and the spectral sequence (2.1.3). In particular, to prove theorem 2.1.1 above, it will suffice to prove the following result.

THEOREM 2.1.6. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . Suppose that I can be embedded as a two-sided ideal $I \subset T$ of a free unital associative ring. Then the induced map*

$$\{\mathrm{THH}_q(A, I^m, \mathbb{Z}/p)\}_{m \geq 1} \rightarrow \{\mathrm{THH}_q(B, f(I)^m, \mathbb{Z}/p)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups, for all integers q .

We remark that theorem 2.1.6 is a statement about the homotopy groups of the underlying non-equivariant spectrum $\mathrm{THH}(A, I)$ of the \mathbb{T} -equivariant spectrum $T(A, I)$. To prove it, we construct a filtration

$$\mathrm{Fil}_0 \mathrm{THH}(A, I) \subset \mathrm{Fil}_1 \mathrm{THH}(A, I) \subset \mathrm{Fil}_2 \mathrm{THH}(A, I) \subset \dots \subset \mathrm{THH}(A, I)$$

and consider the associated spectral sequence

$$E_{s,t}^1 = H_{s+t}(\mathrm{gr}_s \mathrm{THH}(A, I), k) \Rightarrow H_{s+t}(\mathrm{THH}(A, I), k)$$

for the spectrum homology of $\mathrm{THH}(A, I)$ with coefficients in a field k . By a variant of the proof of Wodzicki's excision theorem for Hochschild homology [30, Thm. 3.1] we show that, in the situation of theorem 2.1.6, the induced map of E^1 -terms is an isomorphism of pro-abelian groups.

2.2. In this section, we recall from [3, Appendix] the construction of the exact triangle (2.1.5) in the \mathbb{T} -stable category.

The ring A determines a symmetric ring spectrum E^0 , and the ideal $I \subset A$ determines a symmetric spectrum E^1 which is a unital symmetric E^0 - E^0 -bimodule [17]. To briefly recall this, let $S^1[-] = \Delta^1[-]/\partial\Delta^1[-]$ be the standard simplicial circle, and let $S^i[-] = S^1[-] \wedge \cdots \wedge S^1[-]$ be the i -fold smash product. Then

$$(E^0)_i = |[k] \mapsto A \otimes_{\mathbb{Z}} (\mathbb{Z}\{S_k^i\}/\mathbb{Z}\{s_{0,k}\})|,$$

$$(E^1)_i = |[k] \mapsto I \otimes_{\mathbb{Z}} (\mathbb{Z}\{S_k^i\}/\mathbb{Z}\{s_{0,k}\})|$$

are Eilenberg-MacLane spaces for A and I , respectively, concentrated in degree i . Here $\mathbb{Z}\{S_k^i\}$ is the free abelian group generated by the set of k -simplices S_k^i and $\mathbb{Z}\{s_{0,k}\}$ is the subgroup generated by the base-point. The action of the symmetric group Σ_i by permutation of the i smash factors of $S^i[-]$ induces an action by Σ_i on $(E^e)_i$. The unit and multiplication maps in A give rise to maps

$$\mu_{i,i'}: (E^0)_i \wedge (E^0)_{i'} \rightarrow (E^0)_{i+i'}, \quad \eta_i: S^i \rightarrow (E^0)_i,$$

which make E^0 a symmetric ring spectrum, and the left and right multiplication of A on I give rise to maps

$$\mu_{i,i'}: (E^0)_i \wedge (E^1)_{i'} \rightarrow (E^1)_{i+i'}, \quad \mu_{i,i'}: (E^1)_i \wedge (E^0)_{i'} \rightarrow (E^1)_{i+i'}$$

which make E^1 a unital symmetric E^0 - E^0 -bimodule. Moreover, the canonical inclusion of I in A induces a map $\rho: E^1 \rightarrow E^0$ of E^0 - E^0 -bimodules.

We let \mathcal{I} be the product category $\mathcal{I}_1 \times \mathcal{I}_2$, where \mathcal{I}_1 is the category with objects the finite sets $\underline{i} = \{1, 2, \dots, i\}$, $i \geq 1$, and the empty set $\underline{0}$ and with morphisms all injective maps, and where \mathcal{I}_2 is the category with objects the integers 0 and 1 and with a single non-identity morphism from 1 to 0. The categories \mathcal{I}_1 and \mathcal{I}_2 have strict monoidal structures defined as follows. The monoidal product on \mathcal{I}_1 associates to the pair of objects $(\underline{i}, \underline{i}')$ the object $\underline{i} \sqcup \underline{i}' = \underline{i+i}'$ and to the pair of morphisms $(f: \underline{i} \rightarrow \underline{j}, f': \underline{i}' \rightarrow \underline{j}')$ the morphism $f \sqcup f': \underline{i} \sqcup \underline{i}' \rightarrow \underline{j} \sqcup \underline{j}'$ given by

$$(f \sqcup f')(s) = \begin{cases} f(s) & \text{if } 1 \leq s \leq i \\ f'(s-i) + j & \text{if } i+1 \leq s \leq i+i', \end{cases}$$

and the monoidal product on \mathcal{I}_2 associates to the pair of integers (ϵ, ϵ') the maximum $\epsilon \sqcup \epsilon' = \max\{\epsilon, \epsilon'\}$. The product category $\mathcal{I} = \mathcal{I}_1 \times \mathcal{I}_2$ has a strict monoidal structure given by the component-wise monoidal product.

Let X be a pointed space. There is a functor $G_k(A; X)$ from \mathcal{I}^{k+1} to the category of pointed spaces that on objects is given by

$$G_k(A; X)(\underline{i}_0, \epsilon_0; \dots; \underline{i}_k, \epsilon_k) = F(S^{i_0} \wedge \cdots \wedge S^{i_k}, (E^{\epsilon_0})_{i_0} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge X),$$

where the right-hand side is the space of continuous base-point preserving maps with the compact-open topology. Let $\iota: \underline{i}_r \rightarrow \underline{i}'_r$ be the standard inclusion and write $i'_r = i_r + j_r$. Then $G_k(A; X)(\underline{i}_0, \epsilon_0; \dots; \iota, \epsilon_r; \dots; \underline{i}_k, \epsilon_k)$ takes

$$S^{i_0} \wedge \cdots \wedge S^{i_r} \wedge \cdots \wedge S^{i_k} \xrightarrow{f} (E^{\epsilon_0})_{i_0} \wedge \cdots \wedge (E^{\epsilon_r})_{i_r} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge X$$

to the composite

$$\begin{aligned}
& S^{i_0} \wedge \cdots \wedge S^{i'_r} \wedge \cdots \wedge S^{i_k} \xrightarrow{\sim} S^{i_0} \wedge \cdots \wedge S^{i_r} \wedge \cdots \wedge S^{i_k} \wedge S^{j_r} \\
& \xrightarrow{f \wedge \eta_{j_r}} (E^{\epsilon_0})_{i_0} \wedge \cdots \wedge (E^{\epsilon_r})_{i_r} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge (E^0)_{j_r} \wedge X \\
& \xrightarrow{\sim} (E^{\epsilon_0})_{i_0} \wedge \cdots \wedge (E^{\epsilon_r})_{i_r} \wedge (E^0)_{j_r} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge X \\
& \xrightarrow{\mu_{i_r, j_r}} (E^{\epsilon_0})_{i_0} \wedge \cdots \wedge (E^{\epsilon_r})_{i'_r} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge X,
\end{aligned}$$

where the first and third maps are the canonical isomorphisms, and where we have suppressed identity maps. A general morphism $\alpha: \underline{i}_r \rightarrow \underline{i}'_r$ can be written non-uniquely as a composition $\alpha = \sigma \circ \iota$, where $\iota: \underline{i}_r \rightarrow \underline{i}'_r$ is the standard inclusion and $\sigma: \underline{i}'_r \rightarrow \underline{i}'_r$ an automorphism. The symmetric group $\Sigma_{i'_r}$ acts on $S^{i'_r}$ and $(E^{\epsilon_r})_{i'_r}$ and by conjugation on $G_k(A; X)(\underline{i}_0, \epsilon_0; \dots; \underline{i}'_r, \epsilon_r; \dots; \underline{i}_k, \epsilon_k)$, and this defines the map $G_k(A; X)(\underline{i}_0, \epsilon_0; \dots; \sigma, \epsilon_r; \dots; \underline{i}_k, \epsilon_k)$. Finally, let $\beta: 1 \rightarrow 0$ be the unique non-identity morphism. Then $G_k(A; X)(\underline{i}_0, \epsilon_0; \dots; \underline{i}_r, \beta; \dots; \underline{i}_k, \epsilon_k)$ takes

$$S^{i_0} \wedge \cdots \wedge S^{i_r} \wedge \cdots \wedge S^{i_k} \xrightarrow{f} (E^{\epsilon_1})_{i_0} \wedge \cdots \wedge (E^1)_{i_r} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge X$$

to the composite

$$\begin{aligned}
& S^{i_0} \wedge \cdots \wedge S^{i_r} \wedge \cdots \wedge S^{i_k} \xrightarrow{f} (E^{\epsilon_0})_{i_0} \wedge \cdots \wedge (E^1)_{i_r} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge X \\
& \xrightarrow{\rho} (E^{\epsilon_0})_{i_0} \wedge \cdots \wedge (E^0)_{i_r} \wedge \cdots \wedge (E^{\epsilon_k})_{i_k} \wedge X.
\end{aligned}$$

This defines the functor $G_k(A; X)$ on morphisms.

We now define a cyclic pointed space $\mathrm{THH}(A; X)'[-]$ whose pointed space of k -simplices is given by the following homotopy colimit of pointed spaces. (We refer to [4, §XII] for the definition of homotopy colimits).

$$\mathrm{THH}(A; X)'[k] = \mathrm{hocolim}_{\mathcal{I}^{k+1}} G_k(A; X).$$

The homotopy type of the individual spaces in the limit system approximates the homotopy type of the homotopy colimit in the sense that, if X is m -connected, then the canonical map

$$G_k(A; X)(\underline{i}_0, 0; \dots; \underline{i}_k, 0) \rightarrow \mathrm{THH}(A; X)'[k]$$

is $(i_0 + \cdots + i_k + m)$ -connected [19, Lemma 2.3.7]. The cyclic structure maps

$$\begin{aligned}
d_r &: \mathrm{THH}(A; X)'[k] \rightarrow \mathrm{THH}(A; X)'[k-1], & 0 \leq r \leq k, \\
s_r &: \mathrm{THH}(A; X)'[k] \rightarrow \mathrm{THH}(A; X)'[k+1], & 0 \leq r \leq k, \\
t_k &: \mathrm{THH}(A; X)'[k] \rightarrow \mathrm{THH}(A; X)'[k]
\end{aligned}$$

are defined to be the following composites.

$$\begin{aligned}
& \mathrm{hocolim}_{I^{k+1}} G_k(A; X) \xrightarrow{\delta_r} \mathrm{hocolim}_{I^{k+1}} G_{k-1}(A; X) \circ \mathfrak{d}_r \rightarrow \mathrm{hocolim}_{I^k} G_{k-1}(A; X), \\
& \mathrm{hocolim}_{I^{k+1}} G_k(A; X) \xrightarrow{\sigma_r} \mathrm{hocolim}_{I^{k+1}} G_{k+1}(A; X) \circ \mathfrak{s}_r \rightarrow \mathrm{hocolim}_{I^{k+2}} G_{k+1}(A; X), \\
& \mathrm{hocolim}_{I^{k+1}} G_k(A; X) \xrightarrow{\tau_r} \mathrm{hocolim}_{I^{k+1}} G_{k-1}(A; X) \circ \mathfrak{t}_k \rightarrow \mathrm{hocolim}_{I^{k+1}} G_k(A; X).
\end{aligned}$$

The left-hand maps are induced by natural transformations δ_r , σ_r , and τ_k , respectively, and the right-hand maps are the canonical maps. The functors \mathfrak{d}_r , \mathfrak{s}_r , and

t_k are given by

$$\begin{aligned} d_r(\underline{i}_0, \epsilon_0; \dots; \underline{i}_k, \epsilon_k) &= (\underline{i}_0, \epsilon_0; \dots; \underline{i}_r \sqcup \underline{i}_{r+1}, \epsilon_r \sqcup \epsilon_{r+1}); \dots; \underline{i}_k, \epsilon_k), \quad 0 \leq r < k, \\ &= (\underline{i}_k \sqcup \underline{i}_0, \epsilon_r \sqcup \epsilon_0; \underline{i}_1, \epsilon_1; \dots; \underline{i}_{k-1}, \epsilon_{k-1}), \quad r = k, \\ s_r(\underline{i}_0, \epsilon_0; \dots; \underline{i}_k, \epsilon_k) &= (\underline{i}_0, \epsilon_0; \dots; \underline{i}_r, \epsilon_r; \underline{0}, 0; \underline{i}_{r+1}, \epsilon_{r+1}; \dots; \underline{i}_k, \epsilon_k), \quad 0 \leq r \leq k, \\ t_k(\underline{i}_0, \epsilon_0; \dots; \underline{i}_k, \epsilon_k) &= (\underline{i}_k, \epsilon_k; \underline{i}_0, \epsilon_0; \dots; \underline{i}_{k-1}, \epsilon_{k-1}). \end{aligned}$$

The natural transformation δ_r takes the map

$$S^{i_0} \wedge \dots \wedge S^{i_r} \wedge \dots \wedge S^{i_k} \xrightarrow{f} (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_r})_{i_r} \wedge \dots \wedge (E^{\epsilon_k})_{i_k} \wedge X$$

to the composite

$$\begin{aligned} S^{i_0} \wedge \dots \wedge S^{i_r+i_{r+1}} \wedge \dots \wedge S^{i_k} &\xrightarrow{\sim} S^{i_0} \wedge \dots \wedge S^{i_r} \wedge S^{i_{r+1}} \wedge \dots \wedge S^{i_k} \\ &\xrightarrow{f} (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_r})_{i_r} \wedge (E^{\epsilon_{r+1}})_{i_{r+1}} \wedge \dots \wedge (E^{\epsilon_k})_{i_k} \wedge X \\ &\xrightarrow{\mu_{i_r, i_{r+1}}} (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_r \sqcup \epsilon_{r+1}})_{i_r+i_{r+1}} \wedge \dots \wedge (E^{\epsilon_k})_{i_k} \wedge X, \end{aligned}$$

if $0 \leq r < k$, and to the composite

$$\begin{aligned} S^{i_k+i_0} \wedge S^{i_1} \wedge \dots \wedge S^{i_{k-1}} &\xrightarrow{\sim} S^{i_0} \wedge \dots \wedge S^{i_{k-1}} \wedge S^{i_k} \\ &\xrightarrow{f} (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_{k-1}})_{i_{k-1}} \wedge (E^{\epsilon_k})_{i_k} \wedge X \\ &\xrightarrow{\sim} (E^{\epsilon_k})_{i_k} \wedge (E^{\epsilon_0})_{i_0} \wedge (E^{\epsilon_1})_{i_1} \wedge \dots \wedge (E^{\epsilon_{k-1}})_{i_{k-1}} \wedge X \\ &\xrightarrow{\mu_{i_k, i_0}} (E^{\epsilon_k \sqcup \epsilon_0})_{i_k+i_0} \wedge (E^{\epsilon_1})_{i_1} \wedge \dots \wedge (E^{\epsilon_{k-1}})_{i_{k-1}} \wedge X, \end{aligned}$$

if $r = k$. Here the unnamed isomorphisms are the canonical ones. Similarly, the natural transformations σ_r and τ_k take the map f above to the composites

$$\begin{aligned} S^{i_0} \wedge \dots \wedge S^{i_r} \wedge S^0 \wedge S^{i_{r+1}} \wedge \dots \wedge S^{i_k} &\xrightarrow{\sim} S^{i_0} \wedge \dots \wedge S^{i_k} \wedge S^0 \\ &\xrightarrow{f \wedge \eta_0} (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_r})_{i_r} \wedge (E^{\epsilon_{r+1}})_{i_{r+1}} \wedge \dots \wedge (E^{\epsilon_k})_{i_k} \wedge X \wedge (E^0)_0 \\ &\xrightarrow{\sim} (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_r})_{i_r} \wedge (E^0)_0 \wedge (E^{\epsilon_{r+1}})_{i_{r+1}} \wedge \dots \wedge (E^{\epsilon_k})_{i_k} \wedge X \end{aligned}$$

and

$$\begin{aligned} S^{i_k} \wedge S^{i_0} \wedge \dots \wedge S^{i_{k-1}} &\xrightarrow{\sim} S^{i_0} \wedge \dots \wedge S^{i_{k-1}} \wedge S^{i_k} \\ &\xrightarrow{f} (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_{k-1}})_{i_{k-1}} \wedge (E^{\epsilon_k})_{i_k} \wedge X \\ &\xrightarrow{\sim} (E^{\epsilon_k})_{i_k} \wedge (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_{k-1}})_{i_{k-1}} \wedge X. \end{aligned}$$

This defines the cyclic pointed space $\mathrm{THH}(A; X)'[-]$, and we let

$$\mathrm{THH}(A; X)' = |[k] \mapsto \mathrm{THH}(A; X)'[k]|$$

be the geometric realization.

We compare the pointed space $\mathrm{THH}(A; X)'$ and to the usual topological Hochschild space $\mathrm{THH}(A; X)$ which is defined as follows. Let

$$\iota_0: (\mathcal{I}_1)^{k+1} \rightarrow \mathcal{I}^{k+1}$$

the inclusion functor that takes $(\underline{i}_0, \dots, \underline{i}_k)$ to $(\underline{i}_0, 0; \dots; \underline{i}_k, 0)$. Then

$$\mathrm{THH}(A; X)[k] = \mathrm{hocolim}_{(\mathcal{I}_1)^{k+1}} G_k(A; X) \circ (\iota_0)^{k+1},$$

and since the functors \mathfrak{d}_r , \mathfrak{s}_r , and \mathfrak{t}_r restrict to functors $\mathfrak{d}_r: (\mathcal{I}_1)^{k+1} \rightarrow (\mathcal{I}_1)^k$, $\mathfrak{s}_r: (\mathcal{I}_1)^{k+1} \rightarrow (\mathcal{I}_1)^{k+2}$, and $\mathfrak{t}_k: (\mathcal{I}_1)^{k+1} \rightarrow (\mathcal{I}_1)^{k+1}$, these pointed spaces form a cyclic pointed space $\mathrm{THH}(A; X)[-]$. By definition,

$$\mathrm{THH}(A; X) = |[k] \mapsto \mathrm{THH}(A; X)[k]|$$

is the geometric realization. The inclusion functor $\iota_0: \mathcal{I}_1 \rightarrow \mathcal{I}$ is cofinal in the sense of [4, §XXI], and therefore, the canonical map

$$\mathrm{THH}(A; X)[k] \xrightarrow{\sim} \mathrm{THH}(A; X)'[k]$$

is a weak equivalence of pointed spaces. Finally, since both cyclic pointed spaces are *proper* in the sense of [21], the induced map of geometric realizations

$$\mathrm{THH}(A; X) \rightarrow \mathrm{THH}(A; X)'$$

is a weak equivalence of pointed spaces.

Finally, we define $T(A)'$ to be the following orthogonal \mathbb{T} -spectrum [20]. Let λ be a finite dimensional orthogonal \mathbb{T} -representation and S^λ the one-point compactification. Then the λ th space of $T(A)'$ is the pointed space

$$T(A)'_\lambda = \mathrm{THH}(A; S^\lambda)'$$

with the diagonal \mathbb{T} -action induced from the $\mathbb{T} \times \mathbb{T}$ -action, where the action of one factor is induced from the \mathbb{T} -action on λ and the action of the other factor is the canonical \mathbb{T} -action on the geometric realization of a cyclic space [18, 7.1.9]. Similarly, there is an orthogonal \mathbb{T} -spectrum $T(A)$ with λ th space $\mathrm{THH}(A; S^\lambda)$, and the canonical map of orthogonal \mathbb{T} -spectra

$$T(A) \rightarrow T(A)'$$

is a weak equivalence.

2.3. We consider the full subcategory

$$\mathrm{Fil}_s(\mathcal{I}^{k+1}) \subset \mathcal{I}^{k+1}, \quad 0 \leq s \leq k+1,$$

with objects the tuples $(\underline{i}_0, \epsilon_0; \dots; \underline{i}_k, \epsilon_k)$ such that at least s of $\epsilon_0, \dots, \epsilon_k$ are equal to 0, and define the pointed subspace

$$\mathrm{Fil}_s \mathrm{THH}(A; X)'[k] \subset \mathrm{THH}(A; X)'[k]$$

to the homotopy colimit of the restriction of the functor $G_k(A; X)$ to the subcategory $\mathrm{Fil}_s(\mathcal{I}^{k+1})$. The canonical inclusion

$$\mathrm{Fil}_{s-1} \mathrm{THH}(A; X)'[k] \hookrightarrow \mathrm{Fil}_s \mathrm{THH}(A; X)'[k]$$

has the homotopy extension property, and hence, the canonical map from the mapping cone to the filtration quotient

$$\mathrm{gr}_s \mathrm{THH}(A; X)'[k] = \mathrm{Fil}_s \mathrm{THH}(A; X)'[k] / \mathrm{Fil}_{s-1} \mathrm{THH}(A; X)'[k]$$

is a weak equivalence. To express the homotopy type of the filtration quotient, we need an analog of the hyper-cube lemma [30, Lemma A.1].

We define a *weight* of rank k to be a function

$$\epsilon: \{0, 1, \dots, k\} \rightarrow \{1, 0, -1\},$$

and we write $|\epsilon|_v$ for the number of $0 \leq i \leq k$ with $\epsilon(i) = v$. There is a partial ordering on the set of weights of rank k where $\epsilon \leq \epsilon'$ if $\epsilon(i) \geq \epsilon'(i)$, for all $0 \leq i \leq k$. Let $E^0 \rightarrow E^{-1}$ be the map of unital symmetric ring spectra associated with the

canonical projection $A \rightarrow A/I$, and let $E^1 \rightarrow E^0$ be the map of unital symmetric E^0 - E^0 -bimodules associated with the canonical inclusion $I \rightarrow A$. If ϵ is a weight of rank k and if X is a pointed space, we let $G_k(A, \epsilon; X)$ be the functor from $(\mathcal{I}_1)^{k+1}$ to pointed spaces that on objects is given by

$$G_k(A, \epsilon; X)(\underline{i}_0, \dots, \underline{i}_k) = F(S^{i_0} \wedge \dots \wedge S^{i_k}, (E^{\epsilon_0})_{i_0} \wedge \dots \wedge (E^{\epsilon_k})_{i_k} \wedge X)$$

and that on morphisms is defined analogously to the functor $G_k(A; X)$ considered above. Let $T(A, \epsilon; X)$ be the homotopy colimit

$$T(A, \epsilon; X)[k] = \operatorname{hocolim}_{(\mathcal{I}_1)^{k+1}} G_k(A, \epsilon; X).$$

Then we have the following analog of [30, Lemma A.1].

LEMMA 2.3.1. *For every $s \geq 0$, there is a canonical weak equivalence*

$$\bar{\Phi}_s: \operatorname{gr}_s \operatorname{THH}(A; X)'[k] \xrightarrow{\sim} \bigvee_{\epsilon \in \mathcal{P}'(k, s)} T(A, \epsilon; X)[k],$$

where the wedge is over all weights ϵ of rank k with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$.

PROOF. We first define the map of the statement. The category $\operatorname{Fil}_s(\mathcal{I}^{k+1})$ is the product of the category $(\mathcal{I}_1)^{k+1}$ and the partially ordered set $\mathcal{P}(k, s)$ of weights ϵ of rank k with $|\epsilon|_{-1} = 0$ and $|\epsilon|_0 \leq s$. Moreover, if $\iota_\epsilon: (\mathcal{I}_1)^{k+1} \rightarrow \mathcal{I}^{k+1}$ is the inclusion of categories that to $(\underline{i}_0, \dots, \underline{i}_k)$ associates $(\underline{i}_0, \epsilon_0; \dots; \underline{i}_k, \epsilon_k)$, then

$$G_k(A, \epsilon; X) = G_k(A, I; X) \circ \iota_\epsilon,$$

and hence, there is a canonical homomorphism

$$\operatorname{hocolim}_{\mathcal{P}(k, s)} T(A, \epsilon; X)[k] \xrightarrow{\sim} \operatorname{Fil}_s \operatorname{THH}(A, I; X)'[k].$$

Let $\mathcal{P}'(k, s)$ be the subset of weights of rank k with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$. We define a map of pointed spaces

$$\Phi_s: \operatorname{hocolim}_{\mathcal{P}(k, s)} T(A, \epsilon; X) \rightarrow \prod_{\epsilon' \in \mathcal{P}'(k, s)} T(A, \epsilon'; X)$$

whose image is contained in the corresponding wedge sum and whose restriction to

$$\operatorname{hocolim}_{\mathcal{P}(k, s-1)} T(A, I, \epsilon; X) \subset \operatorname{hocolim}_{\mathcal{P}(k, s)} T(A, I, \epsilon; X)$$

is the constant map. We recall from [4, §XII] that to give the map Φ_s is equivalent to giving, for each $\epsilon' \in \mathcal{P}'(k, s)$, a compatible family of maps

$$\Phi_{s, \epsilon', \epsilon}: T(A, \epsilon; X) \wedge |\epsilon/\mathcal{P}(k, s)|_+ \rightarrow T(A, \epsilon'; X),$$

where the second smash factor on the left-hand side is the geometric realization of the nerve of the under-category $\epsilon/\mathcal{P}(k, s)$. If $\epsilon = \epsilon_0$ is the unique weight of rank k with $\epsilon_0 \leq \epsilon'$ and $|\epsilon_0|_0 = s$, the under-category $\epsilon_0/\mathcal{P}(s, k)$ is the discrete category with the single object $\epsilon_0 \leq \epsilon_0$, and we let

$$\Phi_{s, \epsilon_0, \epsilon'}: T(A, \epsilon_0; X) \rightarrow T(A, \epsilon'; X)$$

be the canonical map, and if $\epsilon \neq \epsilon_0$, we let $\Phi_{s, \epsilon, \epsilon'}$ be the constant map. To see that these maps are compatible, suppose that $\epsilon \leq \epsilon_0$. Then either $\epsilon = \epsilon_0$ or $|\epsilon|_0 < s$. In the former case, there is nothing to check, and in the latter case, the map

$$T(A, \epsilon; X) \rightarrow T(A, \epsilon_0; X) \xrightarrow{\Phi_{s, \epsilon_0, \epsilon'}} T(A, \epsilon'; X)$$

is equal to the constant map, since, on at least one smash factor, the composite map is induced by the composite $I \rightarrow A \rightarrow A/I$, which is constant equal to zero. Hence, we obtain the desired map Φ_s . It is also clear that this map factors to give the map $\bar{\Phi}_s$ of the statement.

The proof that $\bar{\Phi}_s$ is a weak equivalence is by induction on $k \geq 0$ and is similar to the proof of [30, Lemma A.1]. Suppose that $\epsilon' \leq \epsilon \leq \epsilon''$ are such that for some $0 \leq v \leq k$, $\epsilon'(v) = 1$, $\epsilon(v) = 0$, and $\epsilon''(v) = -1$, and that for all $0 \leq i \leq k$ with $i \neq v$, $\epsilon'(i) = \epsilon(i) = \epsilon''(i)$. Then the composite map

$$T(A, \epsilon'; X)[k] \rightarrow T(A, \epsilon; X)[k] \rightarrow T(A, \epsilon''; X)[k]$$

is the constant map and the induced map from pointed space on the left-hand side to the mapping fiber of the right-hand map is a weak equivalence. \square

The functor \mathfrak{d}_r takes $\text{Fil}_s(\mathcal{I}^{k+1})$ to $\text{Fil}_s(\mathcal{I}^k)$, for $0 \leq s < k$, and to $\text{Fil}_{s-1}(\mathcal{I}^k)$, for $k \leq s \leq k+1$, and the functor \mathfrak{s}_r takes $\text{Fil}_s(\mathcal{I}^{k+1})$ to $\text{Fil}_{s+1}(\mathcal{I}^{k+2})$, for all $0 \leq s \leq k+1$. The functor \mathfrak{t}_k restricts to an automorphism $\text{Fil}_s(\mathcal{I}^{k+1})$. It follows that the cyclic structure maps restrict to maps of pointed spaces

$$\begin{aligned} d_r &: \text{Fil}_s \text{THH}(A; X)'[k] \rightarrow \text{Fil}_s \text{THH}(A; X)'[k-1], & 0 \leq s < k, \\ d_r &: \text{Fil}_s \text{THH}(A; X)'[k] \rightarrow \text{Fil}_{s-1} \text{THH}(A; X)'[k-1], & k \leq s \leq k+1, \\ s_r &: \text{Fil}_s \text{THH}(A; X)'[k] \rightarrow \text{Fil}_{s+1} \text{THH}(A; X)'[k+1], & 0 \leq s \leq k+1, \\ t_k &: \text{Fil}_s \text{THH}(A; X)'[k] \rightarrow \text{Fil}_s \text{THH}(A; X)'[k]. \end{aligned}$$

In particular, the pointed spaces

$$\text{THH}(A, I; X)[k] = \text{Fil}_k \text{THH}(A; X)'[k]$$

form a cyclic subspace of $\text{THH}(A; X)'[-]$. We let $\text{THH}(A, I; X)$ be the geometric realization and $T(A, I)$ the orthogonal \mathbb{T} -spectrum with λ th space $\text{THH}(A, I; S^\lambda)$. The following result is [3, Prop. 10.2].

PROPOSITION 2.3.2. *There is an exact triangle of orthogonal \mathbb{T} -spectra*

$$T(A, I) \rightarrow T(A) \rightarrow T(A/I) \xrightarrow{\partial} T(A, I)[-1]$$

which, in addition, is a sequence of cyclotomic spectra.

PROOF. The proof that $T(A, I)$ is cyclotomic is completely analogous to the proof in [13, Prop. 1.5] that $T(A)$ is cyclotomic. To prove that there is an exact triangle as stated, it suffices to show that the canonical map

$$\text{THH}(A; X)'[k] / \text{THH}(A, I; X)[k] \rightarrow \text{THH}(A/I; X)[k]$$

is a weak equivalence. This is the case $s = k$ of Lemma 2.3.1. \square

We consider the induced filtration of the pointed spaces $\text{THH}(A, I; X)[k]$,

$$\text{Fil}_s \text{THH}(A, I; X)[k] = \begin{cases} \text{Fil}_s \text{THH}(A; X)'[k], & \text{if } k \geq s, \\ \text{Fil}_k \text{THH}(A; X)'[k], & \text{if } k \leq s. \end{cases}$$

For a fixed non-negative integer s , the pointed subspaces

$$\text{Fil}_s \text{THH}(A, I; X)[k] \subset \text{THH}(A, I; X)[k]$$

are preserved by the face maps but not by the degeneracy operators. However, we can still form the thick realization

$$\mathrm{Fil}_s \mathrm{THH}(A, I; X)' = \|[k] \mapsto \mathrm{Fil}_s \mathrm{THH}(A, I; X)[k]\|.$$

To briefly recall this, let Δ be the simplicial index category whose objects are the finite ordered sets

$$[k] = \{1, 2, \dots, k\}, \quad k \geq 0,$$

and whose morphisms are all order-preserving maps. Let $\Delta_{\mathrm{mon}} \subset \Delta$ be the subcategory which has the same objects but whose morphisms are all *injective* order-preserving maps. Let $X[-]$ be a functor from Δ^{op} (resp. from $\Delta_{\mathrm{mon}}^{\mathrm{op}}$) to the category of pointed spaces. Then the geometric realization $\|[k] \mapsto X[k]\|$ (resp. the thick realization $\|[k] \mapsto X[k]\|$) is defined to be the coequalizer of the maps

$$\bigvee X[k'] \wedge (\Delta^k)_+ \rightrightarrows \bigvee X[k] \wedge (\Delta^k)_+$$

where the wedge-sums on the right and left run over all objects and morphisms of Δ (resp. of Δ_{mon}), respectively, and where, on the summand on the left indexed by the morphism $\theta: [k] \rightarrow [k']$, the two maps are given by the maps $X[k'] \wedge \theta_*$ and $\theta^* \wedge (\Delta^k)_+$, respectively. We recall that if $X[-]$ is a simplicial pointed spaces, that is, a functor from Δ^{op} to pointed spaces, then the canonical projection

$$\|[k] \mapsto X[k]\| \rightarrow \|[k] \mapsto X[k]\|$$

is a weak equivalence, provided that the degeneracy maps $s_i: X[k-1] \rightarrow X[k]$ have the homotopy extension property [21]. We also note that if $X[-]$ is (the underlying simplicial pointed space of) a cyclic pointed space, then the geometric realization has a canonical \mathbb{T} -action [18, 7.1.9] but the thick realization does not. Finally, geometric realization and thick realization both commute with forming the smash product with a pointed space.

Let $\mathrm{THH}(A, I)$ be the underlying non-equivariant orthogonal spectrum of the orthogonal \mathbb{T} -spectrum $T(A, I)$. If λ is a real inner-product space (or, equivalently, an orthogonal representation of the trivial group), then

$$\mathrm{THH}(A, I)_\lambda = \|[k] \mapsto \mathrm{THH}(A, I; S^\lambda)[k]\|.$$

Similarly, we let $\mathrm{THH}(A, I)'$ be the orthogonal spectrum whose λ th space is

$$\mathrm{THH}(A, I)'_\lambda = \|[k] \mapsto \mathrm{THH}(A, I; S^\lambda)[k]\|.$$

Then the canonical projection is a weak equivalence of orthogonal spectra

$$\mathrm{THH}(A, I)' \xrightarrow{\sim} \mathrm{THH}(A, I),$$

but the left-hand side admits a filtration

$$\mathrm{Fil}_0 \mathrm{THH}(A, I)' \subset \mathrm{Fil}_1 \mathrm{THH}(A, I)' \subset \dots \subset \mathrm{THH}(A, I)'$$

by the sub-orthogonal spectra with λ th space

$$\mathrm{Fil}_s \mathrm{THH}(A, I)'_\lambda = \|[k] \mapsto \mathrm{Fil}_s \mathrm{THH}(A, I; S^\lambda)[k]\|.$$

Let $E_{\mathbb{F}_p}$ be the commutative unital ring spectrum associated with the field \mathbb{F}_p . The homology of a spectrum X with \mathbb{F}_p -coefficients is defined by

$$H_*(X, \mathbb{F}_p) = \pi_*(X \wedge E_{\mathbb{F}_p}).$$

The filtration of $\mathrm{THH}(A, I)'$ gives rise to a spectral sequence

$$E_{s,t}^1(A, I) = H_{s+t}(\mathrm{gr}_s \mathrm{THH}(A, I)', \mathbb{F}_p) \Rightarrow H_{s+t}(\mathrm{THH}(A, I)', \mathbb{F}_p).$$

Moreover, for each $s \geq 0$, we have a spectral sequence

$$E_{u,v}^1(A, I; s) = H_v(\mathrm{gr}_s \mathrm{THH}(A, I)[u], \mathbb{F}_p) \Rightarrow E_{s,u+v-s}^1(A, I),$$

where the d^1 -differential in the latter sequence is induced by the face maps

$$d_r: \mathrm{gr}_s \mathrm{THH}(A, I)[u] \rightarrow \mathrm{gr}_s \mathrm{THH}(A, I)[u-1].$$

We define a map

$$\bigvee_{\epsilon \in \mathcal{P}'(k,s)} T(A, \epsilon; X)[k] \xrightarrow{d_r} \bigvee_{\epsilon' \in \mathcal{P}'(k-1,s)} T(A, \epsilon'; X)[k-1].$$

Let $\epsilon \in \mathcal{P}'(k, s)$ be a weight of rank k with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$, and let $0 \leq r \leq k$. We define a weight $\partial_r(\epsilon)$ of rank $k-1$ with $|\partial_r(\epsilon)|_0 = 0$ and $|\partial_r(\epsilon)|_{-1} \leq s$ as follows. If $0 \leq r < k$, we let

$$\partial_r(\epsilon)(i) = \begin{cases} \epsilon(i) & \text{for } 0 \leq i < r, \\ \max\{\epsilon(r), \epsilon(r+1)\} & \text{for } i = r, \\ \epsilon(i+1) & \text{for } r < i \leq k-1, \end{cases}$$

and if $r = k$, we let

$$\partial_k(\epsilon)(i) = \begin{cases} \max\{\epsilon(k), \epsilon(0)\} & \text{for } i = 0, \\ \epsilon(i) & \text{for } 0 < i \leq k-1. \end{cases}$$

If $\epsilon \in \mathcal{P}'(k, s)$ is such that $|\partial_r(\epsilon)|_{-1} < s$, we define the component $d_{r,\epsilon}$ of the map d_r to be the constant map. If $\epsilon \in \mathcal{P}'(k, s)$ is such that $|\partial_r(\epsilon)|_{-1} = s$, we define the component $d_{r,\epsilon}$ of the map d_r to be the composite

$$T(A, \epsilon; X)[k] \xrightarrow{d'_r} T(A, \partial_r(\epsilon); X)[k-1] \hookrightarrow \bigvee_{\epsilon' \in \mathcal{P}'(k-1,s)} T(A, \epsilon'; X)[k-1],$$

where the right-hand map is the canonical inclusion, and where the left-hand map is the following composite map.

$$\begin{aligned} \mathrm{hocolim}_{(\mathbb{Z}_1)^{k+1}} G_k(A, \epsilon; X) &\xrightarrow{\delta_r} \mathrm{hocolim}_{(\mathbb{Z}_1)^{k+1}} G_{k-1}(A, \partial_r(\epsilon); X) \circ \mathfrak{D}_r \\ &\longrightarrow \mathrm{hocolim}_{(\mathbb{Z}_1)^k} G_{k-1}(A, \partial_r(\epsilon); X). \end{aligned}$$

Here the first map is induced by a natural transformation defined in a manner completely analogous to the natural transformation δ_r we considered earlier, and the second map is the canonical map.

ADDENDUM 2.3.3. *For all $0 \leq r \leq k$, the following diagram commutes.*

$$\begin{array}{ccc} \mathrm{gr}_s \mathrm{THH}(A; X)'[k] & \xrightarrow[\sim]{\Phi_s} & \bigvee T(A, \epsilon; X)[k] \\ \downarrow d_r & & \downarrow d_r \\ \mathrm{gr}_s \mathrm{THH}(A; X)'[k-1] & \xrightarrow[\sim]{\Phi_s} & \bigvee T(A, \epsilon; X)[k-1]. \end{array}$$

Here the wedge sums on the right-hand side run over all weights of rank k and $k-1$, respectively, with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$.

PROOF. This is clear from the proof of Lemma 2.3.1. □

PROOF OF THEOREM 2.1.6. We let $f: A \rightarrow B$ and $I \subset A$ be as in the statement of Thm. 2.1.6. To prove it, we will prove the stronger statement that for all integers q , the canonical map

$$\{\mathrm{THH}_q(I^m, \mathbb{Z}/p^v)\}_{m \geq 1} \rightarrow \{\mathrm{THH}_q(A, I^m, \mathbb{Z}/p^v)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups. It suffices to show that for all integers q , the corresponding map on homology with \mathbb{F}_p -coefficients

$$\{H_q(\mathrm{THH}(I^m)', \mathbb{F}_p)\}_{m \geq 1} \rightarrow \{H_q(\mathrm{THH}(A, I^m)', \mathbb{F}_p)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups [24, Cor. 5.8]. By Addendum 2.3.3, the map

$$H_q(\mathrm{THH}(I^m)', \mathbb{F}_p) \rightarrow H_q(\mathrm{THH}(A, I^m)', \mathbb{F}_p)$$

is equal to the edge-homomorphism of the spectral sequence

$$E_{s,t}^1(A, I^m) = H_{s+t}(\mathrm{gr}_s \mathrm{THH}(A, I^m)', \mathbb{F}_p) \Rightarrow H_{s+t}(\mathrm{THH}(A, I^m)', \mathbb{F}_p).$$

Hence, it suffices to show that the pro-abelian groups $\{E_{s,t}^1(A, I^m)\}_{m \geq 1}$ are zero, if $s > 0$. To this end, we show that for all $s > 0$ and for all u and v , the pro-abelian groups $\{E_{u,v}^2(A, I^m; s)\}_{m \geq 1}$ are zero. Let ϵ be a weight of rank u , and let \mathcal{A}_*^1 and \mathcal{A}_*^{-1} denote the spectrum homology with \mathbb{F}_p -coefficients of I^m and A/I^m . Then the Künneth formula gives a canonical isomorphism of graded \mathbb{F}_p -vector spaces

$$H_*(T(A, \epsilon)[u], \mathbb{F}_p) \xleftarrow{\sim} \mathcal{A}_*^{\epsilon_0} \otimes \cdots \otimes \mathcal{A}_*^{\epsilon_u},$$

and hence, Lemma 2.3.1 gives a canonical isomorphism of graded \mathbb{F}_p -vectors spaces

$$E_{u,*}^1(A, I^m; s) \xleftarrow{\sim} \bigoplus_{\epsilon \in \mathbb{P}'(u,s)} \mathcal{A}_*^{\epsilon_0} \otimes \cdots \otimes \mathcal{A}_*^{\epsilon_u}$$

with the d^1 -differential given by Addendum 2.3.3. We analyze the E^1 -term following [30, Lemma 3.3.]. Let ℓ be a positive integer and $\mathcal{M} = (n_0, \dots, n_\ell)$ a sequence of non-negative integers such that $n_i > 0$, for $0 < i < \ell$, and write $|\mathcal{M}| = n_0 + \cdots + n_\ell$ and $\ell(\mathcal{M}) = \ell$. We let $B_*(\mathcal{A}_*^1)$ be the bar-complex of the graded non-unital associative ring \mathcal{A}_*^1 and consider \mathcal{A}_*^{-1} as a complex with zero differential. (The bar-complex of a non-unital associative ring R is the chain-complex $B_*(R)$ that is zero in degree zero and the n -fold tensor-product $R \otimes \cdots \otimes R$ in degree $n > 0$ and with differential

$$b'(a_1 \otimes \cdots \otimes a_n) = \sum_{0 < i < n} (-1)^i a_1 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_n.$$

Then, given \mathcal{M} , we let $Y_*(\mathcal{M})$ be the complex

$$B_*(\mathcal{A}_*^1) \otimes \cdots \otimes B_*(\mathcal{A}_*^1) \otimes \mathcal{A}_*^{-1} \otimes \cdots \otimes \mathcal{A}_*^{-1},$$

where there are $\ell(\mathcal{M})$ tensor-factors $B_*(\mathcal{A}_*^1)$ and $|\mathcal{M}|$ tensor-factors \mathcal{A}_*^{-1} . We consider the direct sum decomposition

$$E_{u,*}^1(A, I^m, s) = E_{u,*}^1(A, I^m, s)' \oplus E_{u,*}^1(A, I^m, s)'',$$

where $E_{u,*}^1(A, I^m, s)'$ is the graded F -vector generated by the elements $\rho_1 \otimes \cdots \otimes \rho_u$ such that $\rho_1, \dots, \rho_s \in \mathcal{A}_*^1$ and $\rho_{s+1}, \dots, \rho_u \in \mathcal{A}_*^{-1}$, and where $E_{u,*}^1(A, I^m, s)''$ is the graded F -vector space generated by the elements $\rho_1 \otimes \cdots \otimes \rho_u$ such that there exists $1 \leq i < j \leq u$ with $\rho_i \in \mathcal{A}_*^1$ and $\rho_j \in \mathcal{A}_*^{-1}$. The d^1 -differential preserves this decomposition. It is clear that, as a complex,

$$E^1(A, I^m, s)' = \mathcal{A}_*^{-1} \otimes \cdots \otimes \mathcal{A}_*^{-1} \otimes B_*(\mathcal{A}_*^1),$$

where there are s tensor-factors \mathcal{A}_*^{-1} . There is a filtration by subcomplexes

$$\mathrm{Fil}_0 E^1(A, I^m, s)'' \subset \mathrm{Fil}_1 E^1(A, I^m, s)'' \subset \cdots \subset E^1(A, I^m, s)''$$

where $\mathrm{Fil}_p E_{u,*}^1(A, I^m, s)''$ is the graded \mathbb{F}_p -vector space generated by $\rho_1 \otimes \cdots \otimes \rho_u$ such that $\rho_j \in \mathcal{A}_*^{-1}$ for some $u - p \leq j \leq u$. Then there is an isomorphism of complexes

$$\Psi: \bigoplus Y_*(\mathcal{M}) \otimes \mathcal{A}_*^1 \otimes \cdots \otimes \mathcal{A}_*^1 \xrightarrow{\sim} \mathrm{gr}_p E^1(A, I^m, s)''$$

where the sum runs over all \mathcal{M} with $|\mathcal{M}| = u$, with $\ell(\mathcal{M}) \geq 1$, and with $n_\ell \geq 1$, and where there are p tensor-factors \mathcal{A}_*^1 . The isomorphism is given by

$$\begin{aligned} \Psi(x_{(k_0)} \otimes \cdots \otimes x_{(k_{\ell-1})} \otimes b_{(n_0)} \otimes \cdots \otimes b_{(n_\ell)} \otimes x_{(p)}) \\ = b_{(n_0)} \otimes x_{(k_0)} \otimes \cdots \otimes b_{(n_{\ell-1})} \otimes x_{(k_{\ell-1})} \otimes b_{(n_\ell)} \otimes x_{(p)}, \end{aligned}$$

where we write

$$\begin{aligned} b_{(n_i)} &= b_{n_1+\cdots+n_{i-1}+1} \otimes \cdots \otimes b_{n_1+\cdots+n_i}, & b_v &\in \mathcal{A}_*^{-1}, \\ x_{(k_i)} &= x_{k_1+\cdots+k_{i-1}+1} \otimes \cdots \otimes x_{k_1+\cdots+k_i}, & x_v &\in \mathcal{A}_*^1, \\ x_{(p)} &= x_{k_1+\cdots+k_{\ell-1}+1} \otimes \cdots \otimes x_{k_1+\cdots+k_{\ell-1}+p}, & x_v &\in \mathcal{A}_*^1, \end{aligned}$$

Thm. 2.1.6 is now a consequence of Addendum 3.1.4 below. \square

3. Proof of Theorems A and B

3.1. In this section we complete the proof of Thm. A. The following result is similar to the proof of [6, Thm. 5.3].

LEMMA 3.1.1. *If the statement Thm. A holds whenever I can be embedded as a two-sided ideal of a free unital associative ring, then it holds in general.*

PROOF. Suppose that X is a functor from unital associative rings to symmetric spectra that satisfies that for every pair of rings (A, B) the canonical map

$$X(A \times B) \rightarrow X(A) \times X(B)$$

is a weak equivalence. We can then extend X to a functor X' defined on non-unital associative rings as follows. If I is a non-unital associative ring, we define $X'(I)$ to be the homotopy fiber of the map $X(\mathbb{Z} \ltimes I) \rightarrow X(\mathbb{Z})$ induced by the canonical projection. If A is a unital associative ring, we have a canonical isomorphism of rings $\mathbb{Z} \ltimes A \xrightarrow{\sim} \mathbb{Z} \times A$, and hence, we obtain a weak equivalence

$$X(\mathbb{Z} \ltimes A) \xrightarrow{\sim} X(\mathbb{Z} \times A) \xrightarrow{\sim} X(\mathbb{Z}) \times X(A)$$

of symmetric spectra over $X(\mathbb{Z})$. These maps induce a weak equivalence of the homotopy fiber $X'(A)$ of the map $X(\mathbb{Z} \ltimes A) \rightarrow X(\mathbb{Z})$ and the homotopy fiber $X''(A)$ of the canonical projection from $X(\mathbb{Z}) \times X(A)$ to $X(\mathbb{Z})$. This shows that we have natural weak equivalences

$$X'(A) \xrightarrow{\sim} X''(A) \xleftarrow{\sim} X(A),$$

and hence, the functor X' extends, up to canonical weak equivalence, the functor X to non-unital associative rings as claimed. In the following, we will write X instead of X' for the extended functor.

Let $X^n(A)$ be the homotopy fiber of the cyclotomic trace

$$K(A) \rightarrow \mathrm{TC}^n(A; p).$$

Then Thm. A is equivalent to the statement that for every unital associative ring A , for every two-sided ideal $I \subset A$, and for all integers q , the canonical map

$$\{X_q^n(I, \mathbb{Z}/p^v)\}_{n \geq 1} \rightarrow \{X_q^n(A, I, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups. We let FA denote the free associative ring generated by the underlying set of A and consider the following diagram of non-unital associative rings. The rows and columns of the underlying diagram of abelian groups are all short-exact sequences.

$$\begin{array}{ccccc} J(A, I) & \longrightarrow & J(A) & \longrightarrow & J(A/I) \\ \downarrow & & \downarrow & & \downarrow \\ F(A, I) & \longrightarrow & F(A) & \longrightarrow & F(A/I) \\ \downarrow & & \downarrow & & \downarrow \\ I & \longrightarrow & A & \longrightarrow & A/I. \end{array}$$

Since the non-unital rings in the top row can be embedded as ideals of free unital associative rings, the diagram gives rise to the following map of long-exact sequences of pro-abelian groups.

$$\begin{array}{ccc} \vdots & & \vdots \\ \downarrow & & \downarrow \\ \{X_q^n(J(A, I), \mathbb{Z}/p^v)\} & \xrightarrow{f'_*} & \{X_q^n(J(A), J(A, I), \mathbb{Z}/p^v)\} \\ \downarrow & & \downarrow \\ \{X_q^n(F(A, I), \mathbb{Z}/p^v)\} & \xrightarrow{f_*} & \{X_q^n(F(A), F(A, I), \mathbb{Z}/p^v)\} \\ \downarrow & & \downarrow \\ \{X_q^n(I, \mathbb{Z}/p^v)\} & \xrightarrow{f''_*} & \{X_q^n(A, I, \mathbb{Z}/p^v)\} \\ \downarrow & & \downarrow \\ \vdots & & \vdots \end{array}$$

Finally, since the non-unital associative rings $J(A, I)$ and $F(A, I)$ can be embedded in a free unital associative ring, the maps labelled f'_* and f_* are isomorphisms of pro-abelian groups. Hence, so is the map labelled f''_* . This completes the proof. \square

We recall the following result of Cuntz and Quillen [6, Prop. 4.1]. Let k be a field, and let $I \subset T$ be a two-sided ideal of a free unital associative k -algebra. Then there exists a k -linear section

$$(3.1.2) \quad \alpha: I \rightarrow T \otimes_k I$$

of the multiplication $\mu: T \otimes_k I \rightarrow I$ such that the following diagram commutes.

$$\begin{array}{ccc} T \otimes_k I & \xrightarrow{\text{id} \otimes_k \alpha} & T \otimes_k T \otimes_k I \\ \downarrow \mu & & \downarrow \mu \otimes_k \text{id} \\ I & \xrightarrow{\alpha} & T \otimes_k I. \end{array}$$

This is stated in *loc.cit.* for I is a two-sided ideal of free unital associative \mathbb{C} -algebra, but the proof works for every ground field. The diagram shows, in particular, that the map α takes I^2 to $I \otimes_k I$. We use this to prove the following result.

PROPOSITION 3.1.3. *Let $I \subset T$ be a two-sided ideal of a free unital associative ring. Then the pro-abelian groups*

$$\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^m)}(\mathbb{Z}, \mathbb{Z}/p^v)\}_{m \geq 1}$$

are zero, for all positive integers q .

PROOF. We recall that the bar-complex $B_*(I)$ of the non-unital associative ring I is defined to be the chain complex that is zero in degree zero and the n -fold tensor product $I \otimes \cdots \otimes I$ in positive degree n and with differential

$$b'(x_1 \otimes \cdots \otimes x_n) = \sum_{0 < i < n} (-1)^i x_1 \otimes \cdots \otimes x_i x_{i+1} \otimes \cdots \otimes x_n.$$

Since $\mathbb{Z} \ltimes (I^m)$ is flat over \mathbb{Z} , the group $\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^m)}(\mathbb{Z}, \mathbb{Z}/p^v)$ is canonically isomorphic to the q th homology group of the bar-complex $B_*(I^m) \otimes \mathbb{Z}/p^v$. Hence, it suffices to show that for every $m \geq 1$, there exists $m' \geq m$ such that the chain-map induced from the canonical inclusion

$$B_*(I^{m'}) \otimes \mathbb{Z}/p^v \rightarrow B_*(I^m) \otimes \mathbb{Z}/p^v$$

induces the zero map on homology. Since also $I^m \subset T$ is a two-sided ideal of a free unital associative ring, it will suffice to treat the case $m = 1$. By simple induction, it also suffices to consider the case $v = 1$.

The tensor-product $I \otimes \mathbb{F}_p$ is a non-unital associative \mathbb{F}_p -algebra, and there is a canonical isomorphism of complexes of \mathbb{F}_p -vector spaces

$$B_*(I) \otimes \mathbb{F}_p \xrightarrow{\sim} B_*(I \otimes \mathbb{F}_p).$$

Let $(I \otimes \mathbb{F}_p)'$ and $\overline{I \otimes \mathbb{F}_p}$ be the kernel and image of the map $I \otimes \mathbb{F}_p \rightarrow T \otimes \mathbb{F}_p$ such that we have the following extension of non-unital associative \mathbb{F}_p -algebras.

$$(I \otimes \mathbb{F}_p)' \rightarrow I \otimes \mathbb{F}_p \rightarrow \overline{I \otimes \mathbb{F}_p}.$$

The non-unital associative \mathbb{F}_p -algebra $\overline{I \otimes \mathbb{F}_p}$ is a two-sided ideal of the free unital associative \mathbb{F}_p -algebra $T \otimes \mathbb{F}_p$. Let

$$\alpha: \overline{I \otimes \mathbb{F}_p} \rightarrow (T \otimes \mathbb{F}_p) \otimes (\overline{I \otimes \mathbb{F}_p})$$

be the section of the multiplication from (3.1.2). Then the formula

$$s(x_1 \otimes \cdots \otimes x_n) = (-1)^n x_1 \otimes \cdots \otimes x_{n-1} \otimes \alpha(x_n)$$

defines a null-homotopy of the chain map

$$B_*(\overline{I^2 \otimes \mathbb{F}_p}) = B_*((\overline{I \otimes \mathbb{F}_p})^2) \rightarrow B_*(\overline{I \otimes \mathbb{F}_p})$$

induced by the canonical inclusion.

We next let $K_*(I \otimes \mathbb{F}_p)$ be the kernel complex of the following chain-map induced by the canonical projection.

$$B_*(I \otimes \mathbb{F}_p) \rightarrow B_*(\overline{I \otimes \mathbb{F}_p}).$$

The \mathbb{F}_p -vector space $K_n(I \otimes \mathbb{F}_p)$ is generated by the tensors $x_1 \otimes \cdots \otimes x_n$, where at least one of x_1, \dots, x_n in $(I \otimes \mathbb{F}_p)'$. We claim that the chain-map

$$K_*(I^2 \otimes \mathbb{F}_p) \rightarrow K_*(I \otimes \mathbb{F}_p)$$

induced by the canonical inclusion is equal to zero. This follows immediately from the statement that the following map induced by the canonical inclusion is zero.

$$\begin{array}{ccc} (I^2 \otimes \mathbb{F}_p)' & \longrightarrow & (I \otimes \mathbb{F}_p)' \\ \parallel & & \parallel \\ (I^2 \cap pT)/pI^2 & \longrightarrow & (I \cap pT)/pI. \end{array}$$

Hence, it suffices to show that $I^2 \cap pT \subset pI$, or equivalently, that if $x \in I$ and $y \in I$, and if $xy \in pT$, then $xy \in pI$. We claim that $xy \in pT$ implies that $x \in pT$ or $y \in pT$. If $x \in pT$, we write $x = px'$ with $x' \in T$. Then $xy = px'y$ and since I is a left ideal, $x'y \in I$. Similarly, if $y \in pT$, we write $y = py'$ with $y' \in T$. Then $xy = pxy'$, and since I is a right ideal, $xy' \in I$. In either case, $xy \in pI$ as desired.

It remains to prove that $xy \in pT$ implies that $x \in pT$ or $y \in pT$, or equivalently, that if the class $\bar{x}\bar{y} \in T/pT$ is zero then one of the classes $\bar{x} \in T/pT$ and $\bar{y} \in T/pT$ is zero. Suppose conversely that both $\bar{x} \in T/pT$ and $\bar{y} \in T/pT$ are non-zero. The ring T/pT is a free unital associative \mathbb{F}_p -algebra. Hence, we can write \bar{x} and \bar{y} uniquely as sums

$$\bar{x} = \sum_{\omega} a_{\omega} \omega, \quad \bar{y} = \sum_{\omega} b_{\omega} \omega,$$

where ω runs over all finite words with letters in a set of generators of T/pT , and where $a_{\omega}, b_{\omega} \in \mathbb{F}_p$ are zero for all but finitely many ω . The multiplication in T/pT concatenates words and multiplies coefficients. Let $\omega_1, \dots, \omega_m$ and $\omega'_1, \dots, \omega'_n$ be shortest words that occur with non-zero coefficients in \bar{x} and \bar{y} , respectively. Then the words $\omega_1\omega'_1, \dots, \omega_1\omega'_n, \dots, \omega_m\omega'_1, \dots, \omega_m\omega'_n$ are all different and are exactly the shortest words that occur with a non-zero coefficient in $\bar{x}\bar{y}$. In particular, $\bar{x}\bar{y} \in T/pT$ is non-zero as desired.

Finally, we consider now the following diagram with long-exact columns.

$$\begin{array}{ccccc} \vdots & & \vdots & & \vdots \\ \downarrow & & \downarrow & & \downarrow \\ H_q(K_*(I^{4m} \otimes \mathbb{F}_p)) & \xrightarrow{0} & H_q(K_*(I^{2m} \otimes \mathbb{F}_p)) & \xrightarrow{0} & H_q(K_*(I^m \otimes \mathbb{F}_p)) \\ \downarrow & & \downarrow & & \downarrow \\ H_q(B_*(I^{4m} \otimes \mathbb{F}_p)) & \longrightarrow & H_q(B_*(I^{2m} \otimes \mathbb{F}_p)) & \longrightarrow & H_q(B_*(I^m \otimes \mathbb{F}_p)) \\ \downarrow & & \downarrow & & \downarrow \\ H_q(B_*(\overline{I^{4m} \otimes \mathbb{F}_p})) & \xrightarrow{0} & H_q(B_*(\overline{I^{2m} \otimes \mathbb{F}_p})) & \xrightarrow{0} & H_q(B_*(\overline{I^m \otimes \mathbb{F}_p})) \\ \downarrow & & \downarrow & & \downarrow \\ \vdots & & \vdots & & \vdots \end{array}$$

The top and bottom horizontal maps are zero by what was proved above, and a diagram chase then shows that the composite of the two middle horizontal maps is zero. This completes the proof. \square

The following variant of Prop. 3.1.3 above was used in the proof of Thm.2.1.6.

ADDENDUM 3.1.4. *Let T be a free unital associative ring, and let $I \subset T$ be a two-sided ideal. Let $\mathcal{A}_*(m)$ be the homology with \mathbb{F}_p -coefficients of the non-unital ring spectrum associated with the non-unital associative ring I^m . Then for all positive integers m , the canonical map of bar-complexes*

$$B_*(\mathcal{A}_*(4m)) \rightarrow B_*(\mathcal{A}_*(m))$$

induces the zero map on homology.

PROOF. Let \mathcal{A}_* denote the homology with \mathbb{F}_p -coefficients of the unital ring spectrum associated with the ring of integers. Since I^m is free as an abelian group, we have a canonical isomorphism of graded non-unital associative \mathbb{F}_p -algebras

$$\theta_m: \mathcal{A}_* \otimes I^m \xrightarrow{\sim} \mathcal{A}_*(m),$$

with component-wise multiplication on the left-hand side. Since the graded associative \mathbb{F}_p -algebra \mathcal{A}_* is unital, the \mathbb{F}_p -linear map

$$\alpha: \mathcal{A}_* \rightarrow \mathcal{A}_* \otimes \mathcal{A}_*$$

given by $\alpha(a) = a \otimes 1$ is a section of the multiplication $\mu: \mathcal{A}_* \otimes \mathcal{A}_* \rightarrow \mathcal{A}_*$ and the following diagram commutes.

$$\begin{array}{ccc} \mathcal{A}_* \otimes \mathcal{A}_* & \xrightarrow{\text{id} \otimes \alpha} & \mathcal{A}_* \otimes \mathcal{A}_* \otimes \mathcal{A}_* \\ \downarrow \mu & & \downarrow \mu \otimes \text{id} \\ \mathcal{A}_* & \xrightarrow{\alpha} & \mathcal{A}_* \otimes \mathcal{A}_* \end{array}$$

The rest of the proof exactly follows the proof of Prop. 3.1.3. \square

LEMMA 3.1.5. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto an ideal of B . Suppose that the cyclotomic trace induces an isomorphism of pro-abelian groups*

$$\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1} \xrightarrow{\sim} \{\text{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1},$$

for all integers q . Then the map induced by the cyclotomic trace

$$K_q(A, B, I, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\text{TC}_q^n(A, B, I; p, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, for all integers q .

PROOF. We note that for all positive integers m and n , we have a commutative diagram of spectra in which the columns are exact triangles.

$$\begin{array}{ccc}
K(A, B, I^m) & \longrightarrow & \mathrm{TC}^n(A, B, I^m; p) \\
\downarrow & & \downarrow \\
K(A, B, I) & \longrightarrow & \mathrm{TC}^n(A, B, I; p) \\
\downarrow & & \downarrow \\
K(A/I^m, B/I^m, I/I^m) & \longrightarrow & \mathrm{TC}^n(A/I^m, B/I^m, I/I^m; p) \\
\downarrow & & \downarrow \\
K(A, B, I^m)[-1] & \longrightarrow & \mathrm{TC}^n(A, B, I^m; p)[-1].
\end{array}$$

The induced diagram of homotopy groups with \mathbb{Z}/p^v -coefficients gives rise, as m and n vary, to a map of long-exact sequences of pro-abelian groups. The map of pro-abelian groups induced by the top horizontal map in the diagram above is an isomorphism by assumption, and we wish to show that the map of pro-abelian groups induced by the second horizontal map in the diagram above is an isomorphism. By a five-lemma argument this is equivalent to showing that the map of pro-abelian groups induced from the third horizontal map in the diagram above is an isomorphism. But it follows from [22] and from [8, Thm. 2.2.1] that for $m \geq 1$ fixed, the cyclotomic trace induces an isomorphism of pro-abelian groups

$$K_q(A/I^m, B/I^m, I/I^m, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(A/I^m, B/I^m, I/I^m; p, \mathbb{Z}/p^v)\}_{n \geq 1}.$$

But then the map of pro-abelian groups obtained by also allowing $m \geq 1$ to vary is an isomorphism. This completes the proof. \square

PROOF OF THM. A. By Lemma 3.1.1, we can assume that the non-unital associative ring I can be embedded as a two-sided ideal $I \subset T$ of a free unital associative ring. Moreover, by Lemma 3.1.5 it suffices to show that for all integers q , the cyclotomic trace induces an isomorphism of pro-abelian groups

$$\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1} \xrightarrow{\sim} \{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1}.$$

We claim that both pro-abelian groups are zero. Indeed, by Thm. 2.1.1 the pro-abelian groups $\{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m \geq 1}$ are zero, for all integers q and all positive integers n . Hence, the pro-abelian groups $\{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1}$ are zero, for all integers q as claimed. Finally, in order to show that also the pro-abelian groups $\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1}$ are zero, for all integers q , it suffices by Thm. 1.1.1 to show that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^m)}(\mathbb{Z}, \mathbb{Z}/p^v)\}_{m \geq 1}$ are zero, for all positive integers q . This was proved in Prop. 3.1.3 above. \square

3.2. In this section we prove Theorems B and D of the introduction.

PROOF OF THM. B. Let $X \rightarrow \mathrm{Spec} k$ be a curve over a field k , that is, a separated morphism of finite type with X of dimension one. We wish to show that for all $q \geq r$, the cyclotomic trace

$$K_q(X, \mathbb{Z}/p^v) \rightarrow \{\mathrm{TC}_q^n(X; p, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups. We first reduce to the case where X is affine. So assume that the theorem holds for affine curves. A covering $\{U_i\}$ of X by affine open subschemes gives rise to spectral sequences of pro-abelian groups

$$E_{s,t}^2 = \check{H}^{-s}(\{U_i\}, K_t(-, \mathbb{Z}/p^v)) \Rightarrow K_{s+t}(X, \mathbb{Z}/p^v)$$

and

$$E_{s,t}^2 = \{\check{H}^{-s}(\{U_i\}, \mathrm{TC}_t^n(-; p, \mathbb{Z}/p^v))\}_{n \geq 1} \Rightarrow \{\mathrm{TC}_{s+t}^n(X; p, \mathbb{Z}/p^v)\}_{n \geq 1}.$$

Indeed, for K -theory, this follows from [27, Thm. 10.3], and for topological cyclic homology, this follows directly from the definition of the topological cyclic homology of a scheme [9, Def. 3.2.3]. The cyclotomic trace induces a map of spectral sequences which, by assumption, is an isomorphism on $E_{s,t}^2$ for $t \geq r$. But then the cyclotomic trace also induces an isomorphism of the pro-abelian groups in the abutments in degrees $q \geq r$. So the theorem holds for X .

So assume that X is affine. By an argument similar to the proof of Lemma 3.1.5 above, we can further assume that X is reduced. By induction on the number of irreducible components in X , we can further reduce to the case where X is irreducible. Indeed, let Z and Z' be two proper closed subschemes of X and suppose that Z and Z' cover X , or equivalently, that the corresponding ideals I and I' of the coordinate ring A intersect trivially. Then it follows from Thm. A that the cyclotomic trace induces an isomorphism of pro-abelian groups

$$K_q(A, A/I', I, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(A, A/I', I; p, \mathbb{Z}/p^v)\}_{n \geq 1},$$

for all integers q . This proves the induction step.

Finally, let $X \rightarrow \mathrm{Spec} k$ be a reduced and irreducible affine curve. The coordinate ring A is an integral domain, and we let B be the integral closure. We claim that the conductor ideal $I = \{a \in A \mid aB \subset A\}$, which is the largest ideal of A that is also an ideal of B , is non-zero. Indeed, the extension $A \subset B$ is finite, so we can write $B = b_1A + \cdots + b_mA$. Since A and B have the same quotient field, there are elements $a_i \in A$ such that $a_i b_i \in A$, $1 \leq i \leq m$. But then the product $a_1 \cdots a_m$ is in I . The ring B is a smooth k -algebra, and the quotients A/I and B/I are finite k -algebras. Hence, we know from [9, Thm. 4.2.2] and from [22] and [8, Thm. 2.2.1] that the cyclotomic trace induces an isomorphism of pro-abelian groups

$$K_q(R, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(R; p, \mathbb{Z}/p^v)\}_{n \geq 1},$$

for $q \geq r$, if R is one of B , A/I , and B/I . But then Thm. A shows that this map is an isomorphism, for $q \geq r$, also if $R = A$. \square

PROOF OF THM. D. The ideal $I = |G| \cdot \mathfrak{M}$ is a common ideal of $A = \mathbb{Z}[G]$ and $B = \mathfrak{M}$ [1, Cor. XI.1.2], and hence, we can apply Thm. A. It follows that the exactness of the sequence in the statement of Thm. D is equivalent to the exactness of the corresponding sequence with $\mathbb{Z}[G]$ and \mathfrak{M} replaced by $\mathbb{Z}[G]/I$ and \mathfrak{M}/I . The latter are both artinian rings whose quotient by the Jacobson radical is a semi-simple \mathbb{F}_p -algebra. Hence, the exactness of the latter sequence follows from [13, Thm. 5.1(i)] and from [8, Thm. 2.2.1]. \square

References

- [1] H. Bass, *Algebraic K-theory*, W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [2] A. J. Berrick and W. G. Dwyer, *The spaces that define algebraic K-theory*, *Topology* **39** (2000), 225–228.
- [3] M. Bökstedt and I. Madsen, *Topological cyclic homology of the integers*, *K-theory* (Strasbourg, 1992), *Astérisque*, vol. 226, 1994, pp. 57–143.
- [4] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, *Lecture Notes in Math.*, vol. 304, Springer-Verlag, New York, 1972.
- [5] G. Cortiñas, *The obstruction to excision in K-theory and in cyclic homology*, Preprint 2001, math.KT/0111096.
- [6] J. Cuntz and D. Quillen, *Excision in bivariant periodic cyclic cohomology*, *Invent. Math.* **127** (1997), 67–98.
- [7] R. K. Dennis and M. I. Krusemeyer, $K_2(A[X, Y]/(XY))$, *a problem of Swan, and related computations*, *J. Pure Appl. Alg.* **15** (1979), 125–148.
- [8] T. Geisser and L. Hesselholt, *On the K-theory and topological cyclic homology of smooth schemes over a discrete valuation ring*, *Trans. Amer. Math. Soc.*, to appear.
- [9] ———, *Topological cyclic homology of schemes*, *K-theory* (Seattle, 1997), *Proc. Symp. Pure Math.*, vol. 67, 1999, pp. 41–87.
- [10] S. Geller, L. Reid, and C. A. Weibel, *The cyclic homology and K-theory of curves*, *Bull. Amer. Math. Soc.* **15** (1986), no. 2, 205–208.
- [11] ———, *The cyclic homology and K-theory of curves*, *J. reine angew. Math.* **393** (1989), 39–90.
- [12] L. Hesselholt, *K-theory of truncated polynomial algebras*, *Handbook of K-theory*, Springer-Verlag, New York, to appear.
- [13] L. Hesselholt and I. Madsen, *On the K-theory of finite algebras over Witt vectors of perfect fields*, *Topology* **36** (1997), 29–102.
- [14] ———, *On the K-theory of nilpotent endomorphisms*, *Homotopy methods in algebraic topology* (Boulder, CO, 1999), *Contemp. Math.*, vol. 271, Amer. Math. Soc., Providence, RI, 2001, pp. 127–140.
- [15] ———, *On the K-theory of local fields*, *Ann. of Math.* **158** (2003), 1–113.
- [16] P. S. Hirschhorn, *Model categories and their localizations*, *Mathematical Surveys and Monographs*, vol. 99, American Mathematical Society, Providence, RI, 2003.
- [17] M. Hovey, B. Shipley, and J. Smith, *Symmetric spectra*, *J. Amer. Math. Soc.* **13** (2000), 149–208.
- [18] J.-L. Loday, *Cyclic homology*, *Grundlehren der mathematischen Wissenschaften*, vol. 301, Springer-Verlag, New York, 1992.
- [19] I. Madsen, *Algebraic K-theory and traces*, *Current Developments in Mathematics*, 1995, International Press, Cambridge, MA, 1996, pp. 191–321.
- [20] M. A. Mandell and J. P. May, *Equivariant orthogonal spectra and S-modules*, *Mem. Amer. Math. Soc.*, vol. 159, Amer. Math. Soc., Providence, RI, 2002.
- [21] J. P. May, *The geometry of iterated loop spaces*, *Lecture Notes in Math.*, vol. 271, Springer-Verlag, New York, 1972.
- [22] R. McCarthy, *Relative algebraic K-theory and topological cyclic homology*, *Acta Math.* **179** (1997), 197–222.
- [23] R. Oliver, *Whitehead groups of finite groups*, *London Mathematical Society Lecture Note Series*, vol. 132, Cambridge University Press, Cambridge, 1988.
- [24] I. A. Panin, *On a theorem of Hurewicz and K-theory of complete discrete valuation rings*, *Math. USSR Izvestiya* **29** (1987), 119–131.
- [25] A. A. Suslin, *Excision in the integral algebraic K-theory*, *Proc. Steklov Inst. Math.* **208** (1995), 255–279.
- [26] A. A. Suslin and M. Wodzicki, *Excision in algebraic K-theory*, *Ann. of Math.* **136** (1992), 51–122.
- [27] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, *Grothendieck Festschrift, Volume III*, *Progress in Mathematics*, vol. 88, 1990, pp. 247–435.
- [28] J. B. Wagoner, *Delooping classifying spaces in algebraic K-theory*, *Topology* **11** (1972), 349–370.

- [29] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.
- [30] M. Wodzicki, *Excision in cyclic homology and in rational algebraic K-theory*, Ann. of Math. **129** (1989), 591–639.

UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CALIFORNIA

E-mail address: `geisser@math.usc.edu`

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS

E-mail address: `larsh@math.mit.edu`