

Bi-relative algebraic K -theory and topological cyclic homology

Thomas Geisser and Lars Hesselholt

Introduction

It is well-known that algebraic K -theory preserves products of rings. However, in general, algebraic K -theory does not preserve fiber-products of rings, and bi-relative algebraic K -theory measures the deviation. It was proved by Cortiñas [5] that, rationally, bi-relative algebraic K -theory and bi-relative cyclic homology agree. In this paper, we show that, with finite coefficients, bi-relative algebraic K -theory and bi-relative topological cyclic homology agree. As an application, we show that for a possibly singular curve over a perfect field k of positive characteristic p , the cyclotomic trace map induces an isomorphism of the p -adic algebraic K -groups and the p -adic topological cyclic homology groups in non-negative degrees. As a further application, we show that the difference between the p -adic K -groups of the integral group ring of a finite group and the p -adic K -groups of a maximal $\mathbb{Z}$ -order in the rational group algebra can be expressed entirely in terms of topological cyclic homology.

Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . In general the induced map of relative K -groups $f_*: K_*(A, I) \rightarrow K_*(B, f(I))$ is not an isomorphism and one defines the bi-relative K -groups so as to form a natural long-exact sequence

$$\dots \rightarrow K_q(A, B, I) \rightarrow K_q(A, I) \xrightarrow{f_*} K_q(B, f(I)) \xrightarrow{\partial} K_{q-1}(A, B, I) \rightarrow \dots$$

It was proved recently by Cortiñas [5] that the trace map

$$K_q(A) \otimes \mathbb{Q} \rightarrow \mathrm{HC}_q^-(A \otimes \mathbb{Q})$$

from rational K -theory of A to negative cyclic homology of $A \otimes \mathbb{Q}$ gives rise to an isomorphism of the associated bi-relative theories. The purpose of this paper is to prove the following analogous result for K -theory with finite coefficients.

THEOREM A. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal*

The first named author was partially supported by COE (Japan), and the National Science Foundation. The second named author was partially supported by the National Science Foundation.

of B . Then the map of bi-relative theories induced from the cyclotomic trace map

$$K_q(A, B, I, \mathbb{Z}/p^v) \rightarrow \{\mathrm{TC}_q^n(A, B, I; p, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, for all integers q .

We recall from Bass [1, Thm. XII.8.3] that the common pro-abelian group is zero, for $q \leq 0$. We also remark that Thm. A implies the slightly weaker statement that the map of spectra induced by the cyclotomic trace map

$$K(A, B, I) \rightarrow \mathrm{TC}(A, B, I; p) = \mathrm{holim}_n \mathrm{TC}^n(A, B, I; p)$$

becomes a weak equivalence after p -completion. However, for the proof, it is essential that we work with the pro-abelian groups and not pass to the limit. We also remark that it is possible to combine Thm. A and the theorem of Cortiñas [5] and express the integral bi-relative groups $K_q(A, B, I)$ in terms of the topological cyclic homology and negative cyclic homology.

A weaker version of the theorem of Cortiñas was first stated as a conjecture by Geller, Reid, and Weibel [10, 11] and used to evaluate the rational algebraic K -theory of curves over a field of characteristic zero. In a similar manner we obtain the following result from Thm. A.

THEOREM B. *Let $X \rightarrow \mathrm{Spec} k$ be any curve over a field of positive characteristic p , and let $0 \leq r \leq \infty$ be given by $[k : k^p] = p^r$. Then the cyclotomic trace map*

$$K_q(X, \mathbb{Z}/p^v) \rightarrow \{\mathrm{TC}_q^n(X; p, \mathbb{Z}/p^v)\}_{n \geq 1}.$$

is an isomorphism of pro-abelian groups, for all $q \geq r$.

We use Thm. A to reduce to the smooth case which we have proved earlier [9, Thm. 4.2.2]. We remark again that Thm. B implies the slightly weaker statement that the map of spectra induced by the cyclotomic trace map

$$K(X) \rightarrow \mathrm{TC}(X; p) = \mathrm{holim}_n \mathrm{TC}^n(X; p)$$

becomes a weak equivalence after p -completion and passage to $(r - 1)$ -connected covers. The topological cyclic homology groups on the right-hand side of the statement of Thm. B can often be effectively calculated. As an example, we have the following result; compare [10]. The proof is similar to the proof of [14, Thm. E] and [15, Thm. A] and will appear in [13].

THEOREM C. *Let k be a regular $\mathbb{F}_p$ -algebra, and let $X = \mathrm{Spec} k[x, y]/(xy)$ be the coordinate axes in the affine plane. Then for all positive integers q , there is a canonical isomorphism of pro-abelian groups*

$$\{\mathrm{TC}_q^n(X; p)\}_{n \geq 1} \xleftarrow{\sim} \bigoplus_{m \geq 1} \mathbf{W}_m \Omega_k^{q-2m}$$

where $\mathbf{W}_m \Omega_k^j$ is the group of big de Rham-Witt j -forms on $\mathrm{Spec} k$ [15, 12].

We remark that the group $K_2(X)$ was evaluated previously by Dennis and Krusemeyer [7]. We also wish to point out that, at present, the structure of the K -groups of singular schemes is very far from understood. In particular, there is currently no motivic cohomology theory that can explain the structure of the K -groups that we find here. To illustrate this, let k be a perfect field of positive characteristic p . Then Thm. B and C show that the p -adic K -groups of $X = \mathrm{Spec} k[x, y]/(xy)$ are concentrated in even degrees with the group in positive degree

$2m$ canonically isomorphic to the ring $\mathbf{W}_m(k)$ of big Witt vectors of length m in k . In particular, the p -adic K -groups of X cannot be generated from $K_1(X)$ by means of products and transfers as is the case for smooth schemes.

Finally, let $A = \mathbb{Z}[G]$ be the integral group ring of a finite group and $B = \mathfrak{M}$ a maximal $\mathbb{Z}$ -order of the rational group algebra. Then Thm. A shows that, for every prime p , the difference between the p -adic K -groups of A and B can be expressed entirely in terms of topological cyclic homology. Indeed, if we choose B such that $A \subset B$, then the ideal $I = |G| \cdot B$ is a common ideal of both A and B , and the quotient rings A/I and B/I are finite. The precise relationship is most clearly expressed for G a finite p -group where we obtain the following result. We are grateful to Holger Reich for drawing our attention to this application of Thm. A.

THEOREM D. *Let $\mathbb{Z}[G]$ be the integral group ring of a finite p -group, and let $\mathfrak{M} \subset \mathbb{Q}[G]$ be a maximal $\mathbb{Z}$ -order of the rational group algebra such that $\mathbb{Z}[G] \subset \mathfrak{M}$. Then there is a natural sequence of pro-abelian groups*

$$\cdots \rightarrow K_q(\mathbb{Z}[G], \mathbb{Z}/p^v) \rightarrow \left\{ \begin{array}{c} K_q(\mathfrak{M}, \mathbb{Z}/p^v) \\ \oplus \\ \mathrm{TC}_q^n(\mathbb{Z}[G]; p, \mathbb{Z}/p^v) \end{array} \right\}_{n \geq 1} \rightarrow \{\mathrm{TC}_q^n(\mathfrak{M}; p, \mathbb{Z}/p^v)\}_{n \geq 1} \rightarrow \cdots$$

which is exact for non-negative integers q .

We again remark that Thm. D implies the slightly weaker result that the following square of spectra becomes homotopy-cartesian after p -completion and passage to connective covers.

$$\begin{array}{ccc} K(\mathbb{Z}[G]) & \longrightarrow & \mathrm{TC}(\mathbb{Z}[G]; p) \\ \downarrow & & \downarrow \\ K(\mathfrak{M}) & \longrightarrow & \mathrm{TC}(\mathfrak{M}; p). \end{array}$$

The strength of Thm. D lies in the fact that the maximal order $\mathfrak{M}$ is a regular ring whose K -groups fit in the following localization sequence [20, Thm. 1.17].

$$\cdots \rightarrow \bigoplus_{\ell} K_q(\mathfrak{M}_{\ell}/J_{\ell}) \rightarrow K_q(\mathfrak{M}) \rightarrow K_q(\mathbb{Q}[G]) \rightarrow \cdots$$

The sum on the left-hand side runs over all rational primes ℓ and $J_{\ell} \subset \mathfrak{M}_{\ell}$ is the Jacobson radical in the ℓ -adic completion of $\mathfrak{M}$. The rings $\mathbb{Q}[G]$ and $\mathfrak{M}_{\ell}/J_{\ell}$ are semi-simple.

We briefly discuss the proof of Thm. A which, in outline, is similar to the proof of the theorem of Cortiñas [5]. The proof uses two key observations which are due to Cuntz and Quillen [6]. The first observation is that it suffices to prove Thm. A in the case where the non-unital associative ring I can be embedded as a two-sided ideal $I \subset T$ of a free unital associative ring. The second observation is that, in this case, the pro-algebra $\{I^m \otimes \mathbb{F}_p\}_{m \geq 1}$ is H -unital. By a generalization of the excision theorem of Suslin-Wodzicki [24, 25], we then show that the pro-abelian groups

$$\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1}$$

are zero, for all integers q . The proof is given in paragraph 1 below. Similarly, we show in paragraph 2 that the pro-abelian groups

$$\{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1}$$

are zero, for all integers q . Hence, if I can be embedded as an ideal of a free unital associative ring, the cyclotomic trace map induces an isomorphism of pro-abelian groups in the top row of the following diagram.

$$\begin{array}{ccc} \{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1} & \xrightarrow{\sim} & \{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1} \\ \downarrow & & \downarrow \\ K_q(A, B, I, \mathbb{Z}/p^v) & \longrightarrow & \{\mathrm{TC}_q^n(A, B, I; p, \mathbb{Z}/p^v)\}_{n \geq 1}. \end{array}$$

Finally, it follows from a theorem of McCarthy [19] and from [8, Thm. 2.2.1] that also the lower horizontal map is an isomorphism of pro-abelian groups. The details of this argument are given in paragraph 3 below. This completes the outline of the proof of Thm. A.

A pro-object of a category $\mathcal{C}$ is a functor from a directed partially ordered set to the category $\mathcal{C}$, and a *strict* map between two pro-objects with the same indexing set is a natural transformation. A general map from a pro-object $X = \{X_i\}_{i \in I}$ to a pro-object $Y = \{Y_j\}_{j \in J}$ is an element of the set

$$\mathrm{Hom}_{\mathrm{pro}\text{-}\mathcal{C}}(X, Y) = \lim_J \mathrm{colim}_I \mathrm{Hom}_{\mathcal{C}}(X_i, Y_j).$$

In particular, a pro-object $X = \{X_i\}_{i \in I}$ in a category with a null-object is zero if for all $i \in I$, there exists $i' \geq i$ such that the map $X_{i'} \rightarrow X_i$ is zero.

This paper was written while the authors visited the University of Tokyo. We would like to express our sincere gratitude to the university and to Takeshi Saito for their kind hospitality and for the stimulating atmosphere they provided. Finally, it is a particular pleasure to acknowledge the help we have had from Cortiñas. He long ago suggested to us to prove the main theorem of this paper by adapting the integral excision theorem of Suslin [24] to the pro-setting. He also pointed out a gap in our arguments in an earlier version of this paper.

1. K-theory

1.1. In this paragraph we prove the following variant of the excision theorem of Suslin-Wodzicki [25, 24]. Our treatment closely follows Suslin [24]. If I is a non-unital associative ring, we write $\mathbb{Z} \ltimes I$ for the unital associative ring given by the product $\mathbb{Z} \times I$ with multiplication $(a, x) \cdot (a', x') = (aa', ax' + xa' + xx')$.

THEOREM 1.1.1. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . Suppose that for all positive integers q , the pro-abelian group*

$$\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}_{n \geq 1}$$

is zero. Then for all integers q , the map of pro-abelian groups

$$\{K_q(A, I^n, \mathbb{Z}/p^v)\}_{n \geq 1} \rightarrow \{K_q(B, f(I)^n, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism.

We will use a *functorial* version of the plus-construction of Quillen. Hence,

$$\lambda: X \rightarrow X^+$$

will be a natural transformation of functors from the category of pointed spaces and base-point preserving continuous maps to itself. For example, one can use the functorial localization [17] with respect to a pointed space W whose reduced

integral homology is trivial and whose fundamental group admits a non-trivial group homomorphism to $GL(\mathbb{Z})$ [2].

Let A be a unital associative ring, and let $I \subset A$ be a two-sided ideal. Let $GL(I)$ and $\overline{GL}(A/I)$ be the kernel and image of the map $GL(A) \rightarrow GL(A/I)$ induced by the canonical projection. We let $F(A, I)$ be the homotopy fiber of the canonical map $BGL(A)^+ \rightarrow \overline{BGL}(A/I)^+$, and let $X(A, I)$ be the homotopy pullback of $\overline{BGL}(A/I)$ and $BGL(A)$ over $\overline{BGL}(A/I)^+$. We then have the following commutative diagram of pointed spaces and base-point preserving continuous maps where the rows are fiber sequences.

$$(1.1.2) \quad \begin{array}{ccccc} BGL(I) & \longrightarrow & BGL(A) & \longrightarrow & \overline{BGL}(A/I) \\ \downarrow & & \downarrow & & \parallel \\ F(A, I) & \longrightarrow & X(A, I) & \longrightarrow & \overline{BGL}(A/I) \\ \parallel & & \downarrow & & \downarrow \\ F(A, I) & \longrightarrow & BGL(A)^+ & \longrightarrow & \overline{BGL}(A/I)^+. \end{array}$$

We consider the relative Serre spectral sequence

$$(1.1.3) \quad \begin{aligned} E_{s,t}^2 &= H_s(\overline{GL}(A/I), H_t(F(A, I), BGL(I), \mathbb{Z}/p)) \\ &\Rightarrow H_{s+t}(X(A, I), BGL(A), \mathbb{Z}/p) \end{aligned}$$

which converges to zero. Indeed, both of the maps $BGL(A) \rightarrow BGL(A)^+$ and $X(A, I) \rightarrow BGL(A)^+$ induce isomorphisms on homology, and hence so does the map $BGL(A) \rightarrow X(A, I)$. Hence, the relative homology groups, defined as the reduced homology groups of the mapping cone, are zero. The diagram (1.1.2) and the spectral sequence (1.1.3) are both functorial in the pair $I \subset A$. Hence, if we consider the pro-ideal $\{I^n\}_{n \geq 1}$ of powers of the ideal $I \subset A$, we obtain a spectral sequence of pro-abelian groups.

LEMMA 1.1.4. *If the pro-abelian group $\{H_t(F(A, I^n), BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1}$ is zero, for $0 \leq t < p$, then the pro-abelian group*

$$\{H_s(\overline{GL}(A/I^n), H_p(F(A, I), BGL(I)), \mathbb{Z}/p)\}_{n \geq 1}$$

is zero, for $0 \leq s \leq 1$. If, in addition, the canonical map

$$\{H_{p-1}(BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1} \rightarrow \{H_0(\overline{GL}(A/I^n), H_{p-1}(BGL(I^n), \mathbb{Z}/p))\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, then the pro-abelian group

$$\{H_p(F(A, I^n), BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is zero.

PROOF. We shall omit the $\mathbb{Z}/p$ -coefficients from the homology throughout the proof. The first claim follows immediately from the spectral sequence. To prove the second claim, we show that in the exact sequence of pro-abelian groups

$$\{H_p(F(A, I^n))\} \rightarrow \{H_p(F(A, I^n), BGL(I^n))\} \rightarrow \{H_{p-1}(BGL(I^n))\},$$

both maps are equal to zero. We first show that the right-hand map is zero, or equivalently, that in the following exact sequence of pro-abelian groups, the right-hand map is injective.

$$\{H_p(F(A, I^n), BGL(I^n))\} \rightarrow \{H_{p-1}(BGL(I^n))\} \rightarrow \{H_{p-1}(F(A, I^n))\} \rightarrow 0.$$

The right-hand map is surjective by the hypothesis of the lemma, and hence, the induced sequence of $\{\overline{GL}(A/I^n)\}$ -coinvariants is again exact. By the first statement of the lemma, the pro-abelian group

$$\{H_0(\overline{GL}(A/I^n), H_p(F(A, I^n), BGL(I^n)))\}$$

is zero, and therefore the map of pro-abelian groups

$$\{H_0(\overline{GL}(A/I^n), H_{p-1}(BGL(I^n)))\} \rightarrow \{H_0(\overline{GL}(A/I^n), H_{p-1}(F(A, I^n)))\}$$

is injective. We consider the diagram of pro-abelian groups.

$$\begin{array}{ccc} \{H_{p-1}(BGL(I^n))\} & \longrightarrow & \{H_{p-1}(F(A, I^n))\} \\ \downarrow \sim & & \downarrow \\ \{H_0(\overline{GL}(A/I^n), H_{p-1}(BGL(I^n)))\} & \hookrightarrow & \{H_0(\overline{GL}(A/I^n), H_{p-1}(F(A, I^n)))\}. \end{array}$$

The left-hand vertical map is an isomorphism by assumption. Hence, the upper horizontal map is injective as desired.

It remains to show that the left-hand map in the sequence at the beginning of the proof is zero. This is equivalent to showing that in the following exact sequence of pro-abelian groups, the left-hand map is surjective.

$$\{H_p(BGL(I^n))\} \rightarrow \{H_p(F(A, I^n))\} \rightarrow \{H_p(F(A, I^n), BGL(I^n))\} \rightarrow 0.$$

The surjectivity of the right-hand map was proved above. The induced sequence of $\{\overline{GL}(A/I^n)\}$ -coinvariants is again exact, and by the first statement of the lemma, the group of $\{\overline{GL}(A/I^n)\}$ -coinvariants of the right-hand term is zero. Hence, the map of pro-abelian groups

$$\{H_0(\overline{GL}(A/I^n), H_p(BGL(I^n)))\} \rightarrow \{H_0(\overline{GL}(A/I^n), H_p(F(A, I^n)))\}$$

is surjective. We now consider the following diagram of pro-abelian groups.

$$\begin{array}{ccc} \{H_p(BGL(I^n))\} & \longrightarrow & \{H_p(F(A, I^n))\} \\ \downarrow \twoheadrightarrow & & \downarrow \sim \\ \{H_0(\overline{GL}(A/I^n), H_p(BGL(I^n)))\} & \twoheadrightarrow & \{H_0(\overline{GL}(A/I^n), H_p(F(A, I^n)))\}. \end{array}$$

The left-hand vertical map and the lower horizontal map are surjective, and we claim that the right-hand vertical map is an isomorphism. Indeed, since $X(A, I^n)$ is a homotopy pull-back, the action of $\overline{GL}(A/I^n) = \pi_1(B\overline{GL}(A/I^n))$ on $H_p(F(A, I^n))$ is induced from the action of $\pi_1(B\overline{GL}(A/I^n)^+)$ on $H_p(F(A, I^n))$. But

$$F(A, I^n) \rightarrow BGL(A)^+ \rightarrow B\overline{GL}(A/I^n)^+$$

is a fiber sequence of infinite loop spaces, and therefore the fundamental group of the base acts trivially on the homology of the fiber [27]. This proves that the top horizontal map in the diagram above is surjective as desired. $\square$

The group of column vectors $M_{\infty,1}(I^n)$ is naturally a left $GL(I^n)$ -module, so we can form the semi-direct product group $GL(I^n) \ltimes M_{\infty,1}(I^n)$. We shall show in Prop. 1.2.7 below that the map induced by the canonical inclusion

$$\{H_q(GL(I^n), \mathbb{Z}/p)\} \rightarrow \{H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), \mathbb{Z}/p)\}$$

is an isomorphism of pro-abelian groups. But first we show that this implies the following result.

PROPOSITION 1.1.5. *Let A be a unital associative ring, let $I \subset A$ be a two-sided ideal, and suppose that for all positive integers q , the pro-abelian group*

$$\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}_{n \geq 1}$$

is zero. Then for all integers q , the pro-abelian group

$$\{H_q(F(A, I^n), BGL(I^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is zero.

PROOF. For notational convenience we omit the $\mathbb{Z}/p$ -coefficients for the homology throughout the proof. We shall show that the hypothesis implies that for all integers q , the canonical map of pro-abelian groups

$$\{H_q(BGL(I^n))\}_{n \geq 1} \rightarrow \{H_0(\overline{GL}(A/I^n), H_q(BGL(I^n)))\}_{n \geq 1}$$

is an isomorphism. The proposition then follows from Lemma 1.1.4 by induction on q starting from the trivial case $q = 0$.

It suffices, by the proof of [25, Cor. 1.6], to show that for all $n \geq 1$, there exists $m \geq n$ such that the image of

$$\iota_{m,n*} : H_q(GL(I^m)) \rightarrow H_q(GL(I^n))$$

is $GL(\mathbb{Z})$ -invariant. To this end we consider the following homomorphisms.

$$H_q(GL(I^n)) \xrightarrow{\sigma_*} H_q(GL(I^n) \ltimes M_{\infty,1}(I^n)) \xrightarrow{j_*} H_q(GL(I^n)).$$

The right-hand map is induced from the embedding

$$j(\alpha, v) = \begin{pmatrix} 1 & 0 \\ v & \alpha \end{pmatrix}.$$

By *loc.cit.*, Lemma 1.4, the composite map $\varphi_* = j_* \circ \sigma_*$ is injective. Hence, it suffices to show that the image of the composite map

$$H_q(GL(I^m)) \xrightarrow{\iota_{m,n*}} H_q(GL(I^n)) \xrightarrow{\varphi_*} H_q(GL(I^n))$$

is invariant under the adjoint action by the subgroup

$$\begin{pmatrix} 1 & 0 \\ 0 & GL(\mathbb{Z}) \end{pmatrix} \subset GL(\mathbb{Z}).$$

In effect, we will show that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the adjoint action by the full group $GL(\mathbb{Z})$. The group $GL(\mathbb{Z})$ is generated by the elementary matrices $e_{k,1}(1)$ and $e_{1,k}(1)$, $k \geq 1$, and the diagonal matrix $\mathrm{diag}(-1, 1, 1, \dots)$. It is clear that the image of φ_* is invariant under the action of the diagonal matrix $\mathrm{diag}(-1, 1, 1, \dots)$. It remains to show that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant also under the action of the elementary matrices $e_{k,1}(1)$ and $e_{1,k}(1)$.

The elementary matrices $e_{k,1}(1)$ are in the image of the composite

$$M_{\infty,1}(\mathbb{Z}) \xrightarrow{\gamma} GL(\mathbb{Z}) \ltimes M_{\infty,1}(\mathbb{Z}) \xrightarrow{j} GL(\mathbb{Z}),$$

where the left-hand map takes β to $(1, \beta)$. We show in Prop. 1.2.7 below there exists $m \geq n$ such that the following map of relative homology groups is zero.

$$\iota_{m,n*} : H_q(GL(I^m) \ltimes M_{\infty,1}(I^m), GL(I^m)) \rightarrow H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), GL(I^n)).$$

Let $\beta \in GL(\mathbb{Z})$ be an element in the image of $j \circ \gamma$ and consider the following diagram.

$$\begin{array}{ccc}
H_q(GL(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n)) \\
\downarrow \sigma_* & & \downarrow \sigma_* \\
H_q(GL(I^m) \ltimes M_{\infty,1}(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n) \ltimes M_{\infty,1}(I^n)) \\
\sim \downarrow \text{Ad}(\beta)_* & & \sim \downarrow \text{Ad}(\beta)_* \\
H_q(GL(I^m) \ltimes M_{\infty,1}(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n) \ltimes M_{\infty,1}(I^n)) \\
\downarrow \pi_* & & \downarrow \pi_* \\
H_q(GL(I^m)) & \xrightarrow{\iota_{m,n*}} & H_q(GL(I^n)).
\end{array}$$

The vertical compositions both are the identity homomorphism and, by the choice of $m \geq n$, the image the second horizontal homomorphism $\iota_{m,n*}$ is contained in the image the top right hand vertical homomorphism σ_* . This shows that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the action of β in the image of $j \circ \gamma$. In particular, the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the action of the elementary matrices $e_{k,1}(1)$. One shows in a similar way that there exists (a possibly larger) $m \geq n$ such that the image of $\varphi_* \circ \iota_{m,n*}$ is invariant under the action of the elementary matrices $e_{1,k}(1)$. This completes the proof. $\square$

PROOF OF THM. 1.1.1. It suffices by simple induction to treat the case $v = 1$. Let $f: A \rightarrow B$ be as in the statement. By Prop. 1.1.5, the induced map

$$\{H_q(F(A, I^n), \mathbb{Z}/p)\}_{n \geq 1} \rightarrow \{H_q(F(B, f(I)^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, and [21, Cor. 5.8] then shows that also the induced map

$$\{\pi_q(F(A, I^n), \mathbb{Z}/p)\}_{n \geq 1} \rightarrow \{\pi_q(F(B, f(I)^n), \mathbb{Z}/p)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups. $\square$

1.2. The main result of this section is Prop. 1.2.7 below which was used in the proof of Prop. 1.1.5 above. We begin with some general theory.

Let A be a unital associative ring, and let $I \subset A$ be a two-sided ideal. Then the powers $\{I^n\}_{n \geq 1}$ form a pro-associative ring without unit. We will say that a pro-abelian group $\{M_n\}_{n \geq 1}$ is a left module over $\{I^n\}_{n \geq 1}$ if for all $n \geq 1$, M_n is a left I^n -module and if for all $m \geq n \geq 1$, the following diagram commutes.

$$\begin{array}{ccc}
I^m \otimes M_m & \xrightarrow{\mu_m} & M_m \\
\downarrow \iota_{m,n} \otimes \lambda_{m,n} & & \downarrow \lambda_{m,n} \\
I^n \otimes M_n & \xrightarrow{\mu_n} & M_n.
\end{array}$$

By a homomorphism from a left $\{I^n\}$ -module $\{M_n\}$ to a left $\{I^n\}$ -module $\{M'_n\}$ we mean a strict map of pro-abelian groups $f: \{M_n\} \rightarrow \{M'_n\}$ such that for all $n \geq 1$, the map $f_n: M_n \rightarrow M'_n$ is an I^n -module homomorphism. We will say that a left $\{I^n\}$ -module $\{P_n\}$ is *pseudo-free* if for all $n \geq 1$, there are I^n -module isomorphisms

$$\varphi_n: I^n \otimes L_n \xrightarrow{\sim} P_n$$

with L_n a free abelian group, and if for all $m \geq n \geq 1$, there are maps of abelian groups $\sigma_{m,n}: L_m \rightarrow L_n$ such that the following diagrams commute.

$$\begin{array}{ccc} I^m \otimes L_m & \xrightarrow{\varphi_m} & P_m \\ \downarrow \iota \otimes \sigma_{m,n} & & \downarrow \lambda_{m,n} \\ I^n \otimes L_n & \xrightarrow{\varphi_n} & P_n. \end{array}$$

A homomorphism $f: \{P_n\} \rightarrow \{M_n\}$ from a pseudo-free $\{I^n\}$ -module to an arbitrary $\{I^n\}$ -module is said to be

(i) *special* if there exists a strict map of pro-abelian groups $g: \{L_n\} \rightarrow \{M_n\}$ such that for all $n \geq 1$, the composite

$$I^n \otimes L_n \xrightarrow{\varphi_n} P_n \xrightarrow{f_n} M_n$$

maps $a \otimes x$ to $ag_n(x)$.

(ii) p^v -*special* if the $\{I^n\}$ -module $\{P_n\}$ admits a strict direct sum decomposition

$$\iota' \oplus \iota'': \{P'_n\} \oplus \{P''_n\} \xrightarrow{\sim} \{P_n\}$$

such that $\{P'_n\}$ and $\{P''_n\}$ are pseudo-free $\{I^n\}$ -modules, such that $f \circ \iota'$ is special, and such that $f \circ \iota'' = p^v g$, for some homomorphism $g: \{P''_n\} \rightarrow \{M_n\}$.

LEMMA 1.2.1. *Let $\{P'_n\}$ be a left $\{I^n\}$ -module. Then there exists a special homomorphism $f: \{P_n\} \rightarrow \{P'_n\}$ from a pseudo-free left $\{I^n\}$ -module $\{P_n\}$ such that the image of f_n is equal to $I^n \cdot P'_n \subset P'_n$.*

PROOF. Let $\mathbb{Z}\{P'_n\}$ be the free abelian group generated by the set of elements of P'_n . We define P_n be the extended left I^n -module

$$P_n = I^n \otimes \bigoplus_{k \geq n} \mathbb{Z}\{P'_k\},$$

and let $\lambda_{m,n}: P_m \rightarrow P_n$ be the map

$$I^m \otimes \bigoplus_{k \geq m} \mathbb{Z}\{P'_k\} \xrightarrow{\iota_{m,n} \otimes \text{incl}} I^n \otimes \bigoplus_{k \geq n} \mathbb{Z}\{P'_k\}.$$

Then $\{P_n\}$ is a pseudo-free $\{I^n\}$ -module and the canonical map $f: \{P_n\} \rightarrow \{P'_n\}$ is a special homomorphism such that the image of f_n is equal to $I^n \cdot P'_n \subset P'_n$. $\square$

Let $\{P_n\}$ be a pseudo-free $\{I^n\}$ -module. An augmented complex of $\{I^n\}$ -modules over $\{P_n\}$ is, by definition, a sequence of $\{I^n\}$ -modules and homomorphisms

$$\dots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\}$$

such that $d_n \circ d_n$ and $\epsilon_n \circ d_n$ are equal to zero, for all $n \geq 1$. We say that the augmented complex $\epsilon: \{C_{n,*}\} \rightarrow \{P_n\}$ is

(i) *special* if the $\{C_{n,q}\}$ are pseudo-free, if ϵ is special, and if the homology of the augmented complex $\epsilon_n: C_{n,*} \rightarrow P_n$ is annihilated by I^n .

(ii) p^v -*special* if the $\{C_{n,q}\}$ are pseudo-free, if ϵ is p^v -special, and if the homology of the augmented complex $\epsilon_n: C_{n,*} \rightarrow P_n$ is annihilated by both I^n and p^v .

COROLLARY 1.2.2. *Let $\{P_n\}$ be an pseudo-free $\{I_n\}$ -module. Then there exists both a special augmented complex and a p^v -special augmented complex over $\{P_n\}$.*

We remark that the identity homomorphism of a pseudo-free module is usually not special, and hence, does not constitute an augmented special complex.

PROOF. It is proved by easy induction based on Lemma 1.2.1 that there exists a special augmented complex

$$\cdots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\}.$$

Then the total complex of the double-complex

$$\begin{array}{ccccccc} \cdots & \xrightarrow{d} & \{C_{n,2}\} & \xrightarrow{d} & \{C_{n,1}\} & \xrightarrow{d} & \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\} \\ & & \downarrow p^v & & \downarrow p^v & & \downarrow p^v \\ \cdots & \xrightarrow{d} & \{C_{n,2}\} & \xrightarrow{d} & \{C_{n,1}\} & \xrightarrow{d} & \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\}. \end{array}$$

is a p^v -special augmented complex. $\square$

PROPOSITION 1.2.3. *Let $I \subset A$ be a two-sided ideal of a unital associative ring, and suppose that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \times (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}$ are zero, for all positive integers q . Then every p^v -special augmented complex*

$$\cdots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{\epsilon} \{P_n\},$$

is exact as an augmented complex of pro-abelian groups.

PROOF. By induction, the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \times (I^n)}(\mathbb{Z}, \mathbb{Z}/p^v)\}_{n \geq 1}$ are zero, for all $v \geq 1$. We show by induction on $q \geq -1$ that the following complex of pro-abelian groups has zero homology.

$$\cdots \xrightarrow{d} \{C_{n,2}\} \xrightarrow{d} \{C_{n,1}\} \xrightarrow{d} \{C_{n,0}\} \xrightarrow{d} \{C_{n,-1}\} = \{P_n\}$$

We let $Z_{n,q}$ and $B_{n,q-1}$ be the kernel and image of $d_n: C_{n,q} \rightarrow C_{n,q-1}$ and note that since both I^n and p^v annihilate $H_q(C_{n,*}) = Z_{n,q}/B_{n,q}$, we have

$$I^n Z_{n,q} + p^v Z_{n,q} \subset B_{n,q}.$$

We first consider $q = -1$. Then $Z_{n,-1} = P_n$, and hence $I^n P_n + p^v P_n \subset B_{n,-1}$. But, by the definition of a special homomorphism, $B_{n,-1} \subset I^n P_n + p^v P_n$, and therefore $B_{n,-1} = I^n P_n + p^v P_n$. Hence,

$$\begin{aligned} H_{-1}(C_{n,*}) &\xleftarrow{\sim} \mathrm{Tor}_0^{\mathbb{Z} \times (I^n)}(P_n, \mathbb{Z}/p^v) \xleftarrow{\sim} \mathrm{Tor}_0^{\mathbb{Z} \times (I^n)}(I^n \otimes M_n, \mathbb{Z}/p^v) \\ &\xleftarrow{\sim} \mathrm{Tor}_0^{\mathbb{Z} \times (I^n)}(I^n, \mathbb{Z}/p^v) \otimes M_n \xleftarrow{\sim} \mathrm{Tor}_1^{\mathbb{Z} \times (I^n)}(\mathbb{Z}, \mathbb{Z}/p^v) \otimes M_n. \end{aligned}$$

It follows that the pro-abelian group $\{H_{-1}(C_{n,*})\}$ is zero as desired.

Next, we assume that the pro-abelian group $\{H_i(C_{n,*})\}$ is zero, for $i < q$, and show that the pro-abelian group $\{H_q(C_{n,*})\}$ is zero. The assumption is equivalent to the statement that the following sequence of pro-abelian groups is exact.

$$0 \rightarrow \{Z_{n,q}\} \rightarrow \{C_{n,q}\} \rightarrow \cdots \rightarrow \{C_{n,0}\} \rightarrow \{P_n\} \rightarrow 0.$$

This gives rise to the following spectral sequence of pro-abelian groups.

$$\begin{aligned} E_{s,t}^1 &= \begin{cases} \{\mathrm{Tor}_t^{\mathbb{Z} \times (I^n)}(C_{n,s}, \mathbb{Z}/p^v)\}, & \text{if } 0 \leq s \leq q, \\ \{\mathrm{Tor}_t^{\mathbb{Z} \times (I^n)}(Z_{n,q}, \mathbb{Z}/p^v)\}, & \text{if } s = q+1, \\ \{0\}, & \text{if } s > q+1, \end{cases} \\ &\Rightarrow \{\mathrm{Tor}_{s+t}^{\mathbb{Z} \times (I^n)}(P_n, \mathbb{Z}/p^v)\}. \end{aligned}$$

Since $\{C_{n,s}\}$ is pseudo-free, we have

$$\begin{aligned} \mathrm{Tor}_t^{\mathbb{Z}^\times(I^n)}(C_{n,s}, \mathbb{Z}/p^v) &\xleftarrow{\sim} \mathrm{Tor}_t^{\mathbb{Z}^\times(I^n)}(I^n \otimes M_{n,s}, \mathbb{Z}/p^v) \\ &\xleftarrow{\sim} \mathrm{Tor}_t^{\mathbb{Z}^\times(I^n)}(I^n, \mathbb{Z}/p^v) \otimes M_{n,s} \xleftarrow{\sim} \mathrm{Tor}_{t+1}^{\mathbb{Z}^\times(I^n)}(\mathbb{Z}, \mathbb{Z}/p^v) \otimes M_{n,s}. \end{aligned}$$

It follows that the pro-abelian groups $E_{s,t}^1$ are zero, for $0 \leq s \leq q$. Hence the edge-homomorphism of the spectral sequence

$$\{\mathrm{Tor}_{q+1}^{\mathbb{Z}^\times(I^n)}(P_n, \mathbb{Z}/p^v)\} \rightarrow \{\mathrm{Tor}_0^{\mathbb{Z}^\times(I^n)}(Z_{n,q}, \mathbb{Z}/p^v)\} = \{Z_{n,q}/(I^n Z_{n,q} + p^v Z_{n,q})\}$$

is an isomorphism of pro-abelian groups, and since the pro-abelian group on the left is zero, so is the one on the right. Finally, since $I^n Z_{n,q} + p^v Z_{n,q} \subset B_{n,q}$, the group $Z_{n,q}/(I^n Z_{n,q} + p^v Z_{n,q})$ surjects onto the group $Z_{n,q}/B_{n,q} = H_q(C_{n,*})$, and hence, the pro-abelian group $\{H_q(C_{n,*})\}$ is zero as stated. $\square$

PROPOSITION 1.2.4. *Let $I \subset A$ be a two-sided ideal of a unital associative ring, and suppose that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z}^\times(I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}$ are zero, for all positive integers q . Let F_i , $i \geq 0$, be a sequence of possibly non-additive functors F_i , $i \geq 0$, from the category of $\{I^n\}$ -modules to the category of pro-abelian groups that satisfies the following conditions (i)–(iii).*

(i) *For every $\{I^n\}$ -module $\{P_n\}$, the pro-abelian group $F_0(\{P_n\})$ is zero.*

(ii) *If $\epsilon: \{P_n[-]\} \rightarrow \{P_n\}$ is an augmented simplicial $\{I^n\}$ -module such that the associated augmented complex of pro-abelian groups is exact, then there is a spectral sequence of pro-abelian groups*

$$E_{s,t}^1 = F_t(\{P_n[s]\}) \Rightarrow F_{s+t}(\{P_n\})$$

such that the edge-homomorphism $F_t(\{P_n[0]\}) \rightarrow F_t(\{P_n\})$ is the map induced by the augmentation $\epsilon: \{P_n[0]\} \rightarrow \{P_n\}$.

(iii) *For every $i \geq 0$, there exists $v \geq 0$ such that the map of pro-abelian groups*

$$f_*: F_i(\{P_n\}) \rightarrow F_i(\{P'_n\})$$

induced by a p^v -special homomorphism $f: \{P_n\} \rightarrow \{P'_n\}$ is zero.

Then for every pseudo-free $\{I^n\}$ -module $\{P_n\}$ and for every $i \geq 0$, the pro-abelian group $F_i(\{P_n\})$ is zero.

PROOF. We prove the statement by induction on $i \geq 0$. The case $i = 0$ is trivial, so we assume the statement for $i - 1$ and prove it for i . We choose v such that for all $0 \leq t \leq i$, the map of pro-abelian groups

$$f_*: F_t(\{P_n\}) \rightarrow F_t(\{P'_n\})$$

induced by a p^v -special homomorphism $f: \{P_n\} \rightarrow \{P'_n\}$ is zero. Let $\{P_n\}$ be a pseudo-free $\{I^n\}$ -module and choose a p^v -special complex $\epsilon: \{C_{n,*}\} \rightarrow \{P_n\}$ over $\{P_n\}$. By Prop. 1.2.3 this augmented complex is exact as an augmented complex of pro-abelian groups. We let $\epsilon: \{P_n[-]\} \rightarrow \{P_n\}$ be the augmented simplicial $\{I^n\}$ -module obtained from the augmented complex $\epsilon: \{C_{n,*}\} \rightarrow \{P_n\}$ by the Dold-Kan correspondance [28]. By assumption, we then have the following spectral sequence of pro-abelian groups.

$$E_{s,t}^1 = F_t(\{P_n[s]\}) \Rightarrow F_{s+t}(\{P_n\}).$$

By the inductive hypothesis, the pro-abelian group $E_{s,t}^1$ is zero, for $0 \leq t < i$, and hence, the edge-homomorphism

$$F_i(\{P_n[0]\}) \rightarrow F_i(\{P_n\})$$

is an epimorphism of pro-abelian groups. But the edge-homomorphism is equal to the map induced by the special homomorphism $\epsilon: \{P_n[0]\} \rightarrow \{P_n\}$ and therefore is zero by assumption. This completes the proof. $\square$

If a group G acts from the left on a group H , the semi-direct product group $G \ltimes H$ is defined to be the set $G \times H$ with multiplication

$$(g, h) \cdot (g', h') = (gg', (g'^{-1}h)h').$$

The projection $\pi: G \ltimes H \rightarrow G$ and the map $\sigma: G \rightarrow G \ltimes H$ given by $\sigma(g) = (g, 1)$ are both group-homomorphisms and $\pi \circ \sigma = \text{id}$. Hence, the homology of the semi-direct product group decomposes canonically as a direct sum

$$H_q(G \ltimes H) \xleftarrow{\sim} H_q(G) \oplus H_q(G \ltimes H, G).$$

PROPOSITION 1.2.5. *Let $\{G_n\}$ be a pro-group, and let $\epsilon: \{H_n[-]\} \rightarrow \{H_n\}$ be a strict augmented simplicial pro-group with a strict left $\{G_n\}$ -action. Suppose that the associated complex of pro-abelian groups*

$$\cdots \rightarrow \{\mathbb{Z}\{H_n[2]\}\} \rightarrow \{\mathbb{Z}\{H_n[1]\}\} \rightarrow \{\mathbb{Z}\{H_n[0]\}\} \xrightarrow{\epsilon} \{\mathbb{Z}\{H_n\}\} \rightarrow 0$$

is exact. Then there is a natural spectral sequence of pro-abelian groups

$$E_{s,t}^1 = \{H_t(G_n \ltimes H_n[s], G_n, \mathbb{Z}/p)\} \Rightarrow \{H_{s+t}(G_n \ltimes H_n, G_n, \mathbb{Z}/p)\},$$

and the edge-homomorphism

$$\{H_t(G_n \ltimes H_n[0], G_n, \mathbb{Z}/p)\} \rightarrow \{H_t(G_n \ltimes H_n, G_n, \mathbb{Z}/p)\}$$

is equal to the map induced from ϵ .

PROOF. Let $\epsilon: \mathbb{Z}\{EG[-]\} \rightarrow \mathbb{Z}$ be the augmented simplicial free abelian group generated by the standard augmented simplicial model $EG[-] \rightarrow *$ of a contractible space with a free G -action. The associated augmented complex

$$\epsilon: \mathbb{Z}\{EG[-]\}_* \rightarrow \mathbb{Z}$$

is a natural resolution of $\mathbb{Z}$ by free (right) $\mathbb{Z}[G]$ -modules. Suppose that M_* is a complex of left G -modules. We define $C_*(G; M_*)$ to be the total complex of the double-complex $\mathbb{Z}\{EG[-]\}_* \otimes_{\mathbb{Z}[G]} M_*$ and the hyper-homology of G with coefficients in M_* to be the homology

$$H_*(G; M_*) = H_*(C_*(G; M_*)).$$

There is a functorial spectral sequence

$$(1.2.6) \quad E_{s,t}^2 = H_s(G; H_t(M_*)) \Rightarrow H_{s+t}(G; M_*)$$

with $H_t(M_*)$ considered as complex concentrated in degree zero.

If G acts from the left on a group H , and if M_* is a complex of left $G \ltimes H$ -modules, then there is a canonical isomorphism of complexes

$$C_*(G \ltimes H; M_*) \xleftarrow{\sim} C_*(G; C_*(H; M_*))$$

where the G -action on $C_*(H; M_*)$ is induced from the actions by G on H and M_* . Moreover, if $\tilde{C}_*(H; M_*)$ denotes the reduced complex defined as the degree-wise cokernel of the canonical map $C_*(\{1\}; M_*) \rightarrow C_*(H; M_*)$, then there is a canonical isomorphism of complexes

$$C_*(G \ltimes H, G; M_*) \xleftarrow{\sim} C_*(G; \tilde{C}_*(H; M_*)).$$

We shall only consider this isomorphism in the case where $M_* = \mathbb{Z}/p$ is the complex that consists of the trivial $G \times H$ -module $\mathbb{Z}/p$ concentrated in degree zero.

Let $\epsilon: H[-] \rightarrow H$ be an augmented simplicial group with a left G -action, and let $\tilde{C}_*(H[-]; \mathbb{Z}/p)_*$ denote the total-complex of the double-complex associated with the simplicial complex $\tilde{C}_*(H[-]; \mathbb{Z}/p)$. Then the filtration in the simplicial direction gives rise to a functorial spectral sequence of hyper-homology groups

$$E_{s,t}^1 = H_t(G; \tilde{C}_*(H[s]; \mathbb{Z}/p)) \Rightarrow H_{s+t}(G; \tilde{C}_*(H[-]; \mathbb{Z}/p)_*).$$

Suppose now that $\{G_n\}$ and $\epsilon: \{H_n[-]\} \rightarrow \{H_n\}$ are as in the statement. Then we obtain a spectral sequence of (strict) pro-abelian groups

$$E_{s,t}^1 = \{H_t(G_n; \tilde{C}_*(H_n[s]; \mathbb{Z}/p))\} \Rightarrow \{H_{s+t}(G_n; \tilde{C}_*(H_n[-]; \mathbb{Z}/p)_*)\}.$$

The augmentation induces a (strict) map of pro-abelian groups

$$\epsilon_*: \{H_q(G_n; \tilde{C}_*(H_n[-]; \mathbb{Z}/p)_*)\} \rightarrow \{H_q(G_n; \tilde{C}_*(H_n; \mathbb{Z}/p))\},$$

and we claim that this map is an isomorphism. Indeed, the hypothesis of the proposition implies that the map induced by the augmentation

$$\epsilon_*: \{H_t(\tilde{C}_*(H_n[-]; \mathbb{Z}/p)_*)\} \rightarrow \{\tilde{H}_t(H_n; \mathbb{Z}/p)\}$$

is an isomorphism of pro-abelian groups for all t . The claim then follows from the spectral sequence (1.2.6) which is located in the first quadrant. This proves that there exists a spectral sequence as stated in the proposition. Finally, it is clear from the construction that the edge-homomorphism is as stated. $\square$

Let A be a unital associative ring, let $I \subset A$ be a two-sided ideal, and let $\{P_n\}$ be a left $\{I^n\}$ -module. Then the group of column vectors $M_{\infty,1}(P_n)$ is naturally a left $GL(I^n)$ -module, and we consider the canonical inclusion of $GL(I^n)$ in the semi-direct product $GL(I^n) \ltimes M_{\infty,1}(P_n)$. The relative homology groups form a sequence of non-additive functors

$$F_i(\{P_n\}) = \{H_i(GL(I^n) \ltimes M_{\infty,1}(P_n), GL(I^n)), \mathbb{Z}/p)\}$$

from the category of left $\{I^n\}$ -modules to the category of pro-abelian groups. This sequence of functors trivially satisfies the condition (i) of Prop. 1.2.4 and it follows from Prop. 1.2.5 that it also satisfies the condition (ii). Finally, it follows from [24, Cor. 6.6] that this sequence of functors also satisfies the condition (iii). Hence we obtain the following result that was used in the proof of Prop. 1.1.5 above.

PROPOSITION 1.2.7. *Let $I \subset A$ be a two-sided ideal of a unital associative ring, and suppose that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}$ are zero, for all positive integers q . Then the map induced by the canonical inclusion*

$$\{H_q(GL(I^n), \mathbb{Z}/p)\} \rightarrow \{H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), \mathbb{Z}/p)\}$$

is an isomorphism of pro-abelian groups, for all $q \geq 0$.

PROOF. Indeed, the module $\{I^n\}$ is pseudo-free, and hence, Prop. 1.2.4 shows that the pro-abelian groups

$$\{H_q(GL(I^n) \ltimes M_{\infty,1}(I^n), GL(I^n), \mathbb{Z}/p)\}$$

are zero, for all $q \geq 0$. $\square$

2. Topological cyclic homology

2.1. This paragraph is devoted to the proof of the following result.

THEOREM 2.1.1. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . Suppose that for all positive integers q , the pro-abelian group*

$$\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}_{n \geq 1}$$

is zero. Then for all integers q and positive integers n , the induced map

$$\{\mathrm{TC}_q^n(A, I^m; p, \mathbb{Z}/p^v)\}_{m \geq 1} \rightarrow \{\mathrm{TC}_q^n(B, f(I)^m; p, \mathbb{Z}/p^v)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups.

We first explain that Thm. 2.1.1 follows from the analogous result for topological Hochschild homology. The reader is referred to [16] and [12] for further details on topological Hochschild and topological cyclic homology. Associated with the unital associative ring A one has the topological Hochschild spectrum $T(A)$ which is a object of the $\mathbb{T}$ -stable homotopy category. The equivariant homotopy groups

$$\mathrm{TR}_q^n(A; p) = [S^q \wedge (\mathbb{T}/C_{p^{n-1}})_+, T(A)]_{\mathbb{T}}$$

come equipped with an additional *restriction* map

$$\mathrm{TR}_q^n(A; p) \xrightarrow{R} \mathrm{TR}_q^{n-1}(A; p)$$

whose definition uses the *cyclotomic* structure of $T(A)$. We consider the family of groups $\{\mathrm{TR}_q^n(A; p)\}_{n \geq 1}$ as a pro-abelian group with the restriction map as the structure map. The group $\mathrm{TR}_q^1(A; p)$ is equal to the q th topological Hochschild group $\mathrm{TH}_q(A)$. In general, there are natural long-exact sequences

$$(2.1.2) \quad \cdots \rightarrow \mathbb{H}_q(C_{p^{n-1}}, T(A)) \rightarrow \mathrm{TR}_q^n(A; p) \xrightarrow{R} \mathrm{TR}_q^{n-1}(A; p) \rightarrow \cdots$$

and natural spectral sequences

$$(2.1.3) \quad E_{s,t}^2 = H_s(C_{p^{n-1}}, \mathrm{TH}_t(A)) \Rightarrow \mathbb{H}_{s+t}(C_{p^{n-1}}, T(A)).$$

The canonical projection $\mathbb{T}/C_{p^{n-2}} \rightarrow \mathbb{T}/C_{p^{n-1}}$ induces the *Frobenius* map of equivariant homotopy groups

$$F: \mathrm{TR}_q^n(A; p) \rightarrow \mathrm{TR}_q^{n-1}(A; p),$$

and the topological cyclic homology groups $\mathrm{TC}_q^n(A; p)$ are defined so as to form a long-exact sequence

$$(2.1.4) \quad \cdots \rightarrow \mathrm{TC}_q^n(A; p) \rightarrow \mathrm{TR}_q^n(A; p) \xrightarrow{R-F} \mathrm{TR}_q^{n-1}(A; p) \xrightarrow{\partial} \mathrm{TC}_{q-1}^n(A; p) \rightarrow \cdots$$

If $I \subset A$ is a two-sided ideal, one has the following natural exact triangle in the $\mathbb{T}$ -stable category.

$$(2.1.5) \quad T(A, I) \rightarrow T(A) \rightarrow T(A/I) \xrightarrow{\delta} T(A, I)[-1].$$

Moreover, Bökstedt and Madsen [3, Prop. 10.2] have given a construction of this triangle with the additional property that it is also a sequence of cyclotomic spectra. It follows that there are natural long-exact sequences

$$\cdots \rightarrow \mathrm{TC}_q^n(A, I; p) \rightarrow \mathrm{TC}_q^n(A; p) \rightarrow \mathrm{TC}_q^n(A/I; p) \xrightarrow{\partial} \mathrm{TC}_{q-1}^n(A, I; p) \rightarrow \cdots$$

and relative versions of the long-exact sequences (2.1.2) and (2.1.4) and the spectral sequence (2.1.3). Hence, to prove Thm. 2.1.1 above, it will suffice to prove the following result.

THEOREM 2.1.6. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto a two-sided ideal of B . Suppose that for all positive integers q , the pro-abelian group*

$$\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^n)}(\mathbb{Z}, \mathbb{Z}/p)\}_{n \geq 1}$$

is zero. Then for all integers q , the induced map

$$\{\mathrm{TH}_q(A, I^m, \mathbb{Z}/p)\}_{m \geq 1} \rightarrow \{\mathrm{TH}_q(B, f(I)^m, \mathbb{Z}/p)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups.

2.2. We remark that Thm. 2.1.6 can be regarded as a statement about the homotopy groups of the underlying non-equivariant spectrum $\mathrm{TH}(A, I)$ of the $\mathbb{T}$ -spectrum $T(A, I)$. We first recall the definition of $\mathrm{TH}(A, I)$. The category of symmetric spectra is a closed model category and the associated homotopy category is the stable homotopy category. The smash product of symmetric spectra defines a closed symmetric monoidal structure which descends to a symmetric monoidal structure on the stable homotopy category.

We shall consider various symmetric spectra that are defined as the realization of semi-simplicial symmetric spectra. To briefly recall this, let $\mathbf{\Delta}_{\mathrm{mon}}$ be the category whose objects are the finite ordered sets

$$[k] = \{1, 2, \dots, k\}, \quad k \geq 0,$$

and whose morphisms are all *injective* order-preserving maps. The morphisms are generated by the coface maps $d^r: [k-1] \rightarrow [k]$, $0 \leq r \leq k$, given by $d^r(i) = i$, for $0 \leq i < r$, and $d^r(i) = i+1$, for $r \leq i < k$. Then a semi-simplicial symmetric spectrum is a functor $X[-]$ from $\mathbf{\Delta}_{\mathrm{mon}}^{\mathrm{op}}$ to the category of symmetric spectra, and the realization is the symmetric spectrum defined to be the coequalizer

$$(2.2.1) \quad \bigvee X[k'] \wedge (\Delta^k)_+ \rightrightarrows \bigvee X[k] \wedge (\Delta^k)_+ \longrightarrow \|[k] \mapsto X[k]\|,$$

where the wedge-sums in the middle and on the left run over all objects and morphisms, respectively, of $\mathbf{\Delta}_{\mathrm{mon}}$. The two maps are given, on the summand on the left indexed by the morphism $\theta: [k] \rightarrow [k']$, by the maps $X[k'] \wedge \theta_*$ and $\theta^* \wedge (\Delta^k)_+$, respectively. There is a natural spectral sequence

$$(2.2.2) \quad E_{s,t}^1 = \pi_t(X[s]) \Rightarrow \pi_{s+t}(\|[k] \mapsto X[k]\|)$$

where the d^1 -differential is given by the alternating sum of the maps induced by the face maps $d_r: X[s] \rightarrow X[s-1]$, $0 \leq r \leq s$.

Let I be a non-unital associative ring. We will abuse notation and write I also for the associated Eilenberg-MacLane spectrum [18, Ex. 1.2.5]. The ring structure gives rise to a map of symmetric spectra

$$\mu: I \wedge I \rightarrow I$$

that makes I a non-unital associative symmetric ring spectrum. We define the topological Hochschild spectrum of I to be the symmetric spectrum

$$\mathrm{TH}(I) = \|[k] \mapsto \mathrm{TH}(I)[k]\|$$

that is the realization of the semi-simplicial symmetric spectrum defined as follows. The symmetric spectrum in degree k is the $(k+1)$ -fold smash product

$$\mathrm{TH}(I)[k] = I^{\wedge(k+1)}$$

and the face maps $d_r: \mathrm{TH}(I)[k] \rightarrow \mathrm{TH}(I)[k-1]$, $0 \leq r \leq k$, are defined by

$$d_r = \begin{cases} I^{\wedge r} \wedge \mu \wedge I^{\wedge(k-1-r)} & 0 \leq r < k, \\ (\mu \wedge I^{\wedge(k-1)}) \circ \tau & r = k. \end{cases}$$

Here τ is the canonical isomorphism that permutes the first k smash factors and the last smash factor. We recall from [23, Thm. 4.2.8] that if A is a unital associative ring, the symmetric spectrum $\mathrm{TH}(A)$ defined here is canonically weakly equivalent to the topological Hochschild spectrum defined by Bökstedt. It is in order to note that the symmetric spectrum $\mathrm{TH}(A)$ defined here is *not* equipped with the additional cyclotomic structure that is necessary for the definition of topological cyclic homology. However, this extra structure is not relevant for the proof of Thm. 2.1.6.

Let A be a unital associative ring, and let $I \subset A$ be a two-sided ideal. We again abuse notation and write A and I for the associated Eilenberg-MacLane spectra. Then A is a unital associative symmetric ring spectrum, I is a unital symmetric A - A -bimodule spectrum, and the map $\rho: I \rightarrow A$ induced by the inclusion of I in A is a map of A - A -bimodule spectra. We define a symmetric spectrum

$$\mathrm{TH}(A)' = \|[k] \mapsto \mathrm{TH}(A)'[k]\|$$

as the realization of a semi-simplicial symmetric spectrum defined as follows. Let $\mathcal{P}[k]$ be the category with objects all tuples $\epsilon = (\epsilon_0, \dots, \epsilon_k)$ of integers $\epsilon_i \in \{0, 1\}$ and with a single morphism from ϵ to ϵ' if $\epsilon_i \geq \epsilon'_i$ for all $0 \leq i \leq k$. Let $E[k]$ be the functor from $\mathcal{P}[k]$ to the category of symmetric spectra that is given on objects and morphisms, respectively, by

$$\begin{aligned} E[k](\epsilon_0, \dots, \epsilon_r, \dots, \epsilon_k) &= E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_r} \wedge \dots \wedge E^{\epsilon_k} \\ E[k](\epsilon_0, \dots, \epsilon_r, \dots, \epsilon_k) &= E^{\epsilon_0} \wedge \dots \wedge \rho \wedge \dots \wedge E^{\epsilon_k}, \end{aligned}$$

where $E^0 = A$ and $E^1 = I$, and where $(\epsilon_0, \dots, \epsilon_r, \dots, \epsilon_k)$ denotes the unique morphism from $(\epsilon_0, \dots, 1, \dots, \epsilon_k)$ to $(\epsilon_0, \dots, 0, \dots, \epsilon_k)$. Then

$$\mathrm{TH}(A)'[k] = \mathrm{hocolim}_{\mathcal{P}[k]} E[k];$$

see [4, Chap. XII] for the definition of homotopy colimits. To define the face maps, we first note that the ring and bimodule spectrum maps define maps of symmetric spectra $\mu_{\epsilon, \epsilon'}: E^\epsilon \wedge E^{\epsilon'} \rightarrow E^{\epsilon \sqcup \epsilon'}$ where $\epsilon \sqcup \epsilon' = \max\{\epsilon, \epsilon'\}$. Then the face maps

$$d_r: \mathrm{TH}(A)'[k] \rightarrow \mathrm{TH}(A)'[k-1], 0 \leq r \leq k,$$

are defined to be the composites

$$\mathrm{hocolim}_{\mathcal{P}[k]} E[k] \xrightarrow{\delta_r} \mathrm{hocolim}_{\mathcal{P}[k]} E[k-1] \circ \mathfrak{d}_r \rightarrow \mathrm{hocolim}_{\mathcal{P}[k-1]} E[k-1],$$

where $\mathfrak{d}_r: \mathcal{P}[k] \rightarrow \mathcal{P}[k-1]$ is the functor

$$\mathfrak{d}_r(\epsilon_0, \dots, \epsilon_k) = \begin{cases} (\epsilon_0, \dots, \epsilon_r \sqcup \epsilon_{r+1}, \epsilon_k) & 0 \leq r < k, \\ (\epsilon_k \sqcup \epsilon_0, \epsilon_1, \dots, \epsilon_{k-1}) & r = k, \end{cases}$$

and where $\delta_r: E[k] \rightarrow E[k-1] \circ \mathfrak{d}_r$ is the natural transformation

$$\delta_r = \begin{cases} E^{\epsilon_0} \wedge \cdots \wedge E^{\epsilon_{r-1}} \wedge \mu_{\epsilon_r, \epsilon_{r+1}} \wedge E^{\epsilon_{r+2}} \wedge \cdots \wedge E^{\epsilon_k} & 0 \leq r < k, \\ (\mu_{\epsilon_k, \epsilon_0} \wedge E^{\epsilon_1} \wedge \cdots \wedge E^{\epsilon_{k-1}}) \circ \tau & r = k. \end{cases}$$

The object $(0, \dots, 0)$ in $\mathcal{P}[k]$ is a final object, and hence, the canonical map

$$\mathrm{TH}(A)[k] = E[k](0, \dots, 0) \rightarrow \mathrm{hocolim}_{\mathcal{P}[k]} E[k] = \mathrm{TH}(A)'[k]$$

is a weak equivalence [4, XII.3.1]. The canonical maps form a map of semi-simplicial symmetric spectra and the spectral sequence (2.2.2) shows that the induced map of realizations $\mathrm{TH}(A) \rightarrow \mathrm{TH}(A)'$ is a weak equivalence of symmetric spectra.

Let $\mathcal{P}'[k] \subset \mathcal{P}[k]$ be the full subcategory whose objects are the tuples ϵ such that not all of $\epsilon_0, \dots, \epsilon_k$ are equal to zero. The symmetric spectrum

$$\mathrm{TH}(A, I)[k] = \mathrm{hocolim}_{\mathcal{P}'[k]} E[k]$$

is a sub-symmetric spectrum of $\mathrm{TH}(A)'[k]$, and the mapping cone of the canonical inclusion is canonically weakly equivalent to $\mathrm{TH}(A/I)[k]$. This follows from the proof of Lemma 2.3.1 below. Moreover, the face maps $d_r: \mathrm{TH}(A)'[k] \rightarrow \mathrm{TH}(A)'[k-1]$, $0 \leq r \leq k$, restrict to face maps $d_r: \mathrm{TH}(A, I)[k] \rightarrow \mathrm{TH}(A, I)[k-1]$, $0 \leq r \leq k$, and we define the symmetric spectrum

$$\mathrm{TH}(A, I) = \|[k] \mapsto \mathrm{TH}(A, I)[k]\|$$

to be the realization. Then the canonical weak equivalences of $\mathrm{TH}(A)$ and $\mathrm{TH}(A)'$ and of $\mathrm{TH}(A/I)$ and the mapping cone of the canonical inclusion of $\mathrm{TH}(A, I)$ in $\mathrm{TH}(A)'$ give rise to the following exact triangle in the stable homotopy category.

$$\mathrm{TH}(A, I) \rightarrow \mathrm{TH}(A) \rightarrow \mathrm{TH}(A/I) \rightarrow \mathrm{TH}(A, I)[-1].$$

This exact triangle is canonically isomorphic to the exact triangle associated with the exact triangle (2.1.5) in the T-stable category.

2.3. The category $\mathcal{P}'[k]$ admits a finite filtration

$$\emptyset \subset \mathrm{Fil}_0 \mathcal{P}'[k] \subset \mathrm{Fil}_1 \mathcal{P}'[k] \subset \cdots \subset \mathrm{Fil}_{k-1} \mathcal{P}'[k] \subset \mathrm{Fil}_k \mathcal{P}'[k] = \mathcal{P}'[k]$$

where $\mathrm{Fil}_s \mathcal{P}'[k]$ is the full sub-category whose objects are the tuples $(\epsilon_0, \dots, \epsilon_k)$ such that at most s of $\epsilon_0, \dots, \epsilon_k$ are equal to 0. We obtain an induced filtration

$$* \subset \mathrm{Fil}_0 \mathrm{TH}(A, I)[k] \subset \mathrm{Fil}_1 \mathrm{TH}(A, I)[k] \subset \cdots \subset \mathrm{Fil}_k \mathrm{TH}(A, I)[k] = \mathrm{TH}(A, I)[k]$$

where $\mathrm{Fil}_s \mathrm{TH}(A, I)[k]$ is the sub-symmetric spectrum of $\mathrm{TH}(A, I)[k]$ defined to be the homotopy colimit of the restriction of the functor $E[k]$ to $\mathrm{Fil}_s \mathcal{P}'[k]$. The functor $\mathfrak{d}_r: \mathcal{P}'[k] \rightarrow \mathcal{P}'[k-1]$ takes $\mathrm{Fil}_s \mathcal{P}'[k]$ to $\mathrm{Fil}_s \mathcal{P}'[k-1]$, for $0 \leq s < k$, and therefore, the face map $d_r: \mathrm{TH}(A, I)[k] \rightarrow \mathrm{TH}(A, I)[k-1]$ takes $\mathrm{Fil}_s \mathrm{TH}(A, I)[k]$ to $\mathrm{Fil}_s \mathrm{TH}(A, I)[k-1]$, for $0 \leq s < k$. Hence, we can form the realization

$$\mathrm{Fil}_s \mathrm{TH}(A, I) = \|[k] \mapsto \mathrm{Fil}_s \mathrm{TH}(A, I)[k]\|$$

and obtain the following filtration of $\mathrm{TH}(A, I)$ by sub-symmetric spectra.

$$* \subset \mathrm{Fil}_0 \mathrm{TH}(A, I) \subset \mathrm{Fil}_1 \mathrm{TH}(A, I) \subset \cdots \subset \mathrm{Fil}_s \mathrm{TH}(A, I) \subset \cdots \subset \mathrm{TH}(A, I).$$

We define the filtration quotient $\mathrm{gr}_s \mathrm{TH}(A, I)[k]$ to be the mapping cone of the canonical inclusion of $\mathrm{Fil}_{s-1} \mathrm{TH}(A, I)[k]$ in $\mathrm{Fil}_s \mathrm{TH}(A, I)[k]$ and let

$$\mathrm{gr}_s \mathrm{TH}(A, I) = \|[k] \mapsto \mathrm{gr}_s \mathrm{TH}(A, I)[k]\|.$$

Since forming mapping cone and realization commute, up to canonical isomorphism, we can also define $\mathrm{gr}_s \mathrm{TH}(A, I)$, up to canonical isomorphism, to be the mapping cone of the inclusion of $\mathrm{Fil}_{s-1} \mathrm{TH}(A, I)$ in $\mathrm{Fil}_s \mathrm{TH}(A, I)$. We evaluate the homotopy type of $\mathrm{gr}_s \mathrm{TH}(A, I)$ and begin by evaluating that of $\mathrm{gr}_s \mathrm{TH}(A, I)[k]$.

Let $E^1 \rightarrow E^0$ be the map of unital symmetric A - A -bimodule spectra associated with the canonical inclusion of I in A , and let $E^0 \rightarrow E^{-1}$ be the map of unital symmetric ring spectra associated with the canonical projection of A onto A/I . We say that a tuple $\epsilon = (\epsilon_0, \dots, \epsilon_k)$ of integers $\epsilon_i \in \{1, 0, -1\}$ is a weight of rank k and write $|\epsilon|_v$ for the number of $0 \leq i \leq k$ with $\epsilon_i = v$. Then we have the following analog of the hyper-cube lemma [29, Lemma A.1].

LEMMA 2.3.1. *For every $0 \leq s \leq k$, there is a canonical weak equivalence*

$$\bar{\Phi}_s: \mathrm{gr}_s \mathrm{TH}(A, I)[k] \xrightarrow{\sim} \bigvee E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_k},$$

where the wedge sum is over all weights ϵ of rank k with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$.

PROOF. There is a partial ordering on the set of weights of rank k where $\epsilon \leq \epsilon'$ if $\epsilon_i \geq \epsilon'_i$, for all $0 \leq i \leq k$. We first define a map of symmetric spectra

$$\Phi_s: \mathrm{Fil}_s \mathrm{TH}(A, I)[k] = \mathrm{hocolim}_{\mathrm{Fil}_s \mathcal{P}'[k]} E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_k} \rightarrow \prod E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_k}$$

where the product on the right-hand side is indexed by the set of weights of rank k with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$. Recall from [4, XII.2.2] that to give the map Φ_s is equivalent to giving, for each weight ϵ of rank k with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$, a compatible family of maps

$$\Phi_{s, \epsilon, \epsilon'}: (E^{\epsilon'_0} \wedge \dots \wedge E^{\epsilon'_k}) \wedge |\epsilon'| / \mathrm{Fil}_s \mathcal{P}'[k]|_+ \rightarrow E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_k},$$

where the second smash factor on the left-hand side is the geometric realization of the nerve of the under-category $\epsilon' / \mathrm{Fil}_s \mathcal{P}'[k]$. If $\epsilon' = \omega$ is the unique weight of rank k with $\omega \leq \epsilon$ and $|\omega|_0 = s$, then the under-category $\omega / \mathrm{Fil}_s \mathcal{P}'[k]$ is the discrete category with the single object $\mathrm{id}_\omega: \omega \rightarrow \omega$, and we let

$$\Phi_{s, \omega, \omega}: E^{\omega_0} \wedge \dots \wedge E^{\omega_k} \rightarrow E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_k}$$

be the canonical map. If $\epsilon' \neq \omega$, we let $\Phi_{s, \epsilon, \epsilon'}$ be the constant map. To see that these maps are compatible, suppose that $\epsilon' \leq \omega$. Then either $\epsilon' = \omega$ or $|\epsilon'|_0 < s$. In the former case, there is nothing to check, and in the latter case, the map

$$E^{\epsilon'_0} \wedge \dots \wedge E^{\epsilon'_k} \rightarrow E^{\omega_0} \wedge \dots \wedge E^{\omega_k} \rightarrow E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_k}$$

is equal to the constant map, since, on at least one smash factor, the composite map is induced by the composite $I \rightarrow A \rightarrow A/I$, which is constant equal to zero. Hence, we obtain the desired map Φ_s . The image of Φ_s is contained in the corresponding wedge sum, and the restriction of Φ_s to $\mathrm{Fil}_{s-1} \mathrm{TH}(A, I)[k]$ is equal to the constant map. Hence, we obtain the map $\bar{\Phi}_s$ of the statement.

The proof that $\bar{\Phi}_s$ is a weak equivalence is by induction on $k \geq 0$ and is similar to the proof of [29, Lemma A.1]. Suppose that $\epsilon' \leq \epsilon \leq \epsilon''$ are such that for some $0 \leq v \leq k$, $\epsilon'(v) = 1$, $\epsilon(v) = 0$, and $\epsilon''(v) = -1$, and that for all $0 \leq i \leq k$ with $i \neq v$, $\epsilon'(i) = \epsilon(i) = \epsilon''(i)$. Then the composite map

$$E^{\epsilon'_0} \wedge \dots \wedge E^{\epsilon'_k} \rightarrow E^{\epsilon_0} \wedge \dots \wedge E^{\epsilon_k} \rightarrow E^{\epsilon''_0} \wedge \dots \wedge E^{\epsilon''_k}$$

is the constant map and the induced map from the mapping cone of left-hand map to the symmetric spectrum on the right-hand side is a weak equivalence. $\square$

We define maps of symmetric spectra

$$D_{r,s}: \bigvee E^{\epsilon_0} \wedge \cdots \wedge E^{\epsilon_k} \rightarrow \bigvee E^{\epsilon_0} \wedge \cdots \wedge E^{\epsilon_{k-1}}, \quad 0 \leq r \leq k,$$

where the wedge sum on the left and right-hand side is over all weights ϵ of rank k and $k-1$, respectively, with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$. Let ϵ be a weight of rank k with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$, and let $0 \leq r \leq k$. We define a weight $\partial_r(\epsilon)$ of rank $k-1$ with $|\partial_r(\epsilon)|_0 = 0$ and with $|\partial_r(\epsilon)|_{-1} \leq s$ as follows. If $0 \leq r < k$, we let

$$\partial_r(\epsilon)_i = \begin{cases} \epsilon_i & \text{for } 0 \leq i < r, \\ \max\{\epsilon_r, \epsilon_{r+1}\} & \text{for } i = r, \\ \epsilon_{i+1} & \text{for } r < i \leq k-1, \end{cases}$$

and if $r = k$, we let

$$\partial_k(\epsilon)_i = \begin{cases} \max\{\epsilon_k, \epsilon_0\} & \text{for } i = 0, \\ \epsilon_i & \text{for } 0 < i \leq k-1. \end{cases}$$

If now ϵ is such that $|\partial_r(\epsilon)|_{-1} < s$, we define the component $D_{r,s,\epsilon}$ of the map $D_{r,s}$ to be the constant map. If ϵ is such that $|\partial_r(\epsilon)|_{-1} = s$, we define the component $D_{r,s,\epsilon}$ of the map $D_{r,s}$ to be the composite

$$E^{\epsilon_0} \wedge \cdots \wedge E^{\epsilon_k} \xrightarrow{\delta_r} E^{\partial_r(\epsilon)_0} \wedge \cdots \wedge E^{\partial_r(\epsilon)_{k-1}} \hookrightarrow \bigvee_{\epsilon' \in \mathcal{P}(k-1,s)} E^{\epsilon'_0} \wedge \cdots \wedge E^{\epsilon'_{k-1}},$$

where the right-hand map is the canonical inclusion. The proof of Lemma 2.3.1 gives the following result.

ADDENDUM 2.3.2. *For all $0 \leq s \leq k$ and $0 \leq r \leq k$, diagram*

$$\begin{array}{ccc} \mathrm{gr}_s \mathrm{TH}(A, I)[k] & \xrightarrow[\sim]{\bar{\Phi}_s} & \bigvee E^{\epsilon_0} \wedge \cdots \wedge E^{\epsilon_k} \\ \downarrow d_r & & \downarrow D_{r,s} \\ \mathrm{gr}_s \mathrm{TH}(A, I)[k-1] & \xrightarrow[\sim]{\bar{\Phi}_s} & \bigvee E^{\epsilon_0} \wedge \cdots \wedge E^{\epsilon_{k-1}} \end{array}$$

commutes. Here the wedge sums on the right-hand side run over all weights of rank k and $k-1$, respectively, with $|\epsilon|_0 = 0$ and $|\epsilon|_{-1} = s$. $\square$

We next evaluate the homotopy type of the realization $\mathrm{gr}_s \mathrm{TH}(A, I)$. We recall that the right-hand side of the statement of Lemma 2.3.1 forms a semi-simplicial symmetric spectrum $\mathrm{gr}_s \mathrm{TH}(A, I)^\dagger[-]$ with face maps

$$D_{r,s}: \mathrm{gr}_s \mathrm{TH}(A, I)^\dagger[k] \rightarrow \mathrm{gr}_s \mathrm{TH}(A, I)^\dagger[k-1], \quad 0 \leq r \leq k,$$

and that by Addendum 2.3.2, the weak equivalences of Lemma 2.3.1 form a map of semi-simplicial symmetric spectra. The induced map of realizations

$$\bar{\Phi}_s: \mathrm{gr}_s \mathrm{TH}(A, I) \rightarrow \mathrm{gr}_s \mathrm{TH}(A, I)^\dagger$$

is a weak equivalence of symmetric spectra as one sees by using the natural spectral sequence (2.2.2). It is clear that for $s = 0$,

$$\mathrm{gr}_0 \mathrm{TH}(A, I)^\dagger = \mathrm{TH}(I).$$

For $s > 0$, we further write

$$\mathrm{gr}_s \mathrm{TH}(A, I)^\dagger[k] = \mathrm{gr}_s \mathrm{TH}(A, I)_0^\dagger[k] \vee \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[k]$$

where the first term on the right-hand side consists of the single wedge summand with weight $\epsilon = (-1, \dots, -1, 1, \dots, 1)$ and where the second term consists of the remaining wedge summands. The face maps preserve this decomposition, and hence, after realization, we have a natural isomorphism of symmetric spectra

$$\mathrm{gr}_s \mathrm{TH}(A, I)^\dagger \xleftarrow{\sim} \mathrm{gr}_s \mathrm{TH}(A, I)_0^\dagger \vee \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger.$$

Following [29, Lemma 3.3] we introduce a further filtration of the semi-simplicial symmetric spectrum $\mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[-]$. Let

$$\mathrm{Fil}'_u \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[k] \subset \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[k]$$

consist of the wedge summands that are indexed by the weights ϵ for which there exists $k - u \leq j \leq k$ with $\epsilon_j = -1$, and let $\mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[k]$ be the filtration quotient. After realization we get an induced filtration of the symmetric spectrum $\mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger$ with filtration quotients

$$\mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger = \|[k] \mapsto \mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[k]\|.$$

We recall that the bar-construction $B(I)$ of the non-unital associative symmetric ring spectrum I is defined to be the realization

$$B(I) = \|[k] \mapsto B(I)[k]\|$$

of the semi-simplicial symmetric spectrum whose symmetric spectrum in degree k is $B(I)[k] = I^{\wedge k}$, if $k > 0$, and the trivial symmetric spectrum $B(I)[0] = *$, if $k = 0$, and whose the face maps are defined by

$$d_r = \begin{cases} I^{\wedge(r-1)} \wedge \mu \wedge I^{\wedge(k-r-1)} & 0 < r < k \\ * & r = 0 \text{ or } r = k. \end{cases}$$

The structure of $\mathrm{gr}_s \mathrm{TH}(A, I)$ is given by the following result.

LEMMA 2.3.3. *There are isomorphisms in the stable homotopy category*

$$\begin{aligned} (A/I)^{\wedge s} \wedge B(I) \wedge S^{s-1} &\xrightarrow{\sim} \mathrm{gr}_s \mathrm{TH}(A, I)_0^\dagger \\ \bigvee B(I)^{\wedge \ell} \wedge (A/I)^{\wedge s} \wedge I^{\wedge u} \wedge S^{s+u-1} &\xrightarrow{\sim} \mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger \end{aligned}$$

where, in the second line, the wedge sum runs over all sequences $(n_0, \dots, n_\ell)$ of non-negative integers of length $\ell \geq 1$ such that $n_i > 0$, for $i > 0$, and such that $n_0 + \dots + n_\ell = s$. The isomorphisms are natural with respect to the pair (A, I) .

PROOF. To establish the first isomorphism, we note that

$$\mathrm{gr}_s \mathrm{TH}(A, I)_0^\dagger[k] = \begin{cases} (A/I)^{\wedge s} \wedge B(I)[k - (s-1)] & k \geq s \\ * & 0 \leq k < s \end{cases}$$

with face map $d_r = (A/I)^{\wedge s} \wedge d_i$, if $r = s - 1 + i$ with $0 < i < k - (s - 1)$, and $d_r = *$, otherwise. The desired isomorphism is then given by an isomorphism of symmetric spectra

$$\alpha: (A/I)^{\wedge s} \wedge B(I) \wedge \Delta^{s-1} / \partial \Delta^{s-1} \xrightarrow{\sim} \mathrm{gr}_s \mathrm{TH}(A, I)_0^\dagger$$

defined as follows. The map α is induced by the maps of symmetric spectra

$$\alpha_k: (A/I)^{\wedge s} \wedge B(I)[k - (s-1)] \wedge \Delta_+^{k-(s-1)} \wedge \Delta_+^{s-1} \rightarrow \mathrm{gr}_s \mathrm{TH}(A, I)_0^\dagger[k] \wedge \Delta_+^k$$

which, in turn, are induced by the maps of pointed spaces

$$\alpha'_k: \Delta_+^{k-(s-1)} \wedge \Delta_+^{s-1} \rightarrow \Delta_+^k$$

that are given by the formula

$$\alpha'_k((u_0, \dots, u_{k-(s-1)}), (t_0, \dots, t_s)) = (t_1, \dots, t_s, t_0 u_0, \dots, t_0 u_{k-(s-1)}).$$

One checks that the maps α_k , $k \geq 0$, factor through the equivalence relation that defines the realization and induce the isomorphism α as stated.

The proof of the second isomorphism is similar but notationally more involved. The symmetric spectrum $\mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[k]$, for $0 \leq k < s$, is trivial, and for $k \geq s$, is equal to

$$\bigvee (A/I)^{\wedge n_0} \wedge B(I)[k_1] \wedge (A/I)^{\wedge n_1} \wedge \dots \wedge B(I)[k_\ell] \wedge (A/I)^{\wedge n_\ell} \wedge I^{\wedge u}$$

where the wedge sum runs over all sequences of non-negative integers $(n_0, \dots, n_\ell)$ and $(k_1, \dots, k_\ell)$ such that $\ell \geq 1$ and $n_i > 0$, for $i > 0$, and such that $n_0 + \dots + n_\ell = s$ and $k_1 + \dots + k_\ell = k - (u + s - 1)$. The face map $D_{r,s}$ takes the summand indexed by $(n_0, \dots, n_\ell)$ and $(k_1, \dots, k_t, \dots, k_\ell)$ to the summand indexed by $(n_0, \dots, n_\ell)$ and $(k_1, \dots, k_t - 1, \dots, k_\ell)$ by the map induced from

$$d_i: B(I)[k_t] \rightarrow B(I)[k_t - 1],$$

if there exists $1 \leq t \leq \ell$ and $0 < i < k_t$ such that

$$r = n_0 + k_1 + n_1 + \dots + k_{t-1} + n_{t-1} - 1 + i,$$

and it annihilates the summand indexed by $(n_0, \dots, n_\ell)$ and $(k_1, \dots, k_\ell)$, otherwise. In particular, the face maps preserve the index $(n_0, \dots, n_\ell)$. Hence, as a semi-simplicial symmetric spectrum, $\mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I)_1^\dagger[-]$ decomposes as a wedge-sum indexed by the sequences $(n_0, \dots, n_\ell)$. Let $E(n_0, \dots, n_\ell)[-]$ denote the wedge-summand corresponding to the index $(n_0, \dots, n_\ell)$. To finish the proof of the lemma, we observe that there is a natural isomorphism of symmetric spectra

$$\beta: B(I)^{\wedge \ell} \wedge (A/I)^{\wedge s} \wedge I^{\wedge u} \wedge \Delta^{s+u-1} / \partial \Delta^{s+u-1} \xrightarrow{\sim} \|[k] \mapsto E(n_0, \dots, n_\ell)[k]\|.$$

The map β is defined in a manner similar to the map α above and is induced from the maps of pointed spaces

$$\beta'_{(k_1, \dots, k_\ell)}: \Delta_+^{k_1} \wedge \dots \wedge \Delta_+^{k_\ell} \wedge \Delta_+^{s+u-1} \rightarrow \Delta_+^k$$

that to the $(\ell + 1)$ -tuple of points with barycentric coordinates

$$((u_{1,0}, \dots, u_{1,k_1}), \dots, (u_{\ell,0}, \dots, u_{\ell,k_\ell}), (t_0, \dots, t_{s+u-1}))$$

associates the point with barycentric coordinates

$$(t_1, \dots, t_{n_0}, t'_0 u_{1,0}, \dots, t'_0 u_{1,k_1}, t_{n_0+1}, \dots, t_{n_0+n_1}, \dots, \dots, t'_0 u_{\ell,0}, \dots, t'_0 u_{\ell,k_\ell}, t_{n_0+\dots+n_{\ell-1}+1}, \dots, t_{s+u-1})$$

where $t'_0 = t_0/\ell$. This completes the proof of the lemma. $\square$

2.4. Let I be a non-unital associative ring, and let also I denote the associated Eilenberg-MacLane spectrum. We proceed to describe the spectrum homology

$$H_*(B(I), \mathbb{Z}/p) = \pi_*(B(I) \wedge \mathbb{Z}/p).$$

We let S be the sphere spectrum and define a unital symmetric ring spectrum $S \ltimes I$ to be the wedge-sum $S \vee I$ with multiplication map

$$(S \vee I) \wedge (S \vee I) \xrightarrow{\sim} S \vee I \vee I \vee I \wedge I \rightarrow S \vee I.$$

The left-hand map is the canonical isomorphism and the right-hand map is given by the canonical inclusion $\iota_1: S \rightarrow S \vee I$ on the first summand, by the canonical inclusion $\iota_2: I \rightarrow S \vee I$ on the second and third summands, and by the composition of the multiplication $\mu: I \wedge I \rightarrow I$ and the canonical inclusion $\iota_2: I \rightarrow S \vee I$ on the last summand. The map $\epsilon: S \ltimes I \rightarrow S$ that annihilates the summand I is a map of unital symmetric ring spectra and a retraction of the unit map $\eta = \iota_1: S \rightarrow S \ltimes I$. We define the two-sided bar-construction to be the geometric realization

$$B(S, S \ltimes I, S) = |[k] \mapsto B(S, S \ltimes I, S)[k]|$$

of the simplicial symmetric spectrum with the symmetric spectrum in degree k defined to be $B(S, S \ltimes I, S)[k] = (S \ltimes I)^{\wedge k}$ and with face and degeneracy maps

$$d_r = \begin{cases} \epsilon \wedge (S \ltimes I)^{\wedge(k-1)} & r = 0 \\ (S \ltimes I)^{\wedge(r-1)} \wedge \mu \wedge (S \ltimes I)^{\wedge(k-r-1)} & 0 < r < k \\ (S \ltimes I)^{\wedge(k-1)} \wedge \epsilon & r = k \end{cases}$$

$$s_r = (S \ltimes I)^{\wedge r} \wedge \eta \wedge (S \ltimes I)^{\wedge(k-r)} \quad 0 \leq r \leq k.$$

The geometric realization of a simplicial symmetric spectrum is formed as in (2.2.1) but with the category $\mathbf{\Delta}_{\text{mon}}$ replaced by the category $\mathbf{\Delta}$ where morphisms are *all* order-preserving maps. The symmetric spectrum $B(S, S \ltimes I, S)$ contains $B(S, S, S) = S$ and $B(I)$ as sub-symmetric spectra, and the canonical map

$$S \vee B(I) \rightarrow B(S, S \ltimes I, S)$$

is an isomorphism. The homology of $B(I)$ is given by the following result.

LEMMA 2.4.1. *Let I be a non-unital associative ring. Then there is a natural first quadrant spectral sequence*

$$E_{s,t}^2 = \text{Tor}_s^{\mathbb{Z} \ltimes I}(\mathbb{Z}, \mathbb{Z}/p) \otimes H_t(\mathbb{Z}, \mathbb{Z}/p) \Rightarrow H_{s+t}(B(S, S \ltimes I, S), \mathbb{Z}/p).$$

PROOF. Let $\epsilon: R \xrightarrow{\sim} I$ be a weak equivalence of non-unital associative simplicial rings such that for all $s \geq 0$, the non-unital associative ring $R[s]$ is a free abelian group. The induced weak equivalence of the associative symmetric ring spectra, which we also write $\epsilon: R \xrightarrow{\sim} I$, gives rise to an equivalence of spectra

$$B(S, S \ltimes R, S) \xrightarrow{\sim} B(S, S \ltimes I, S).$$

The left-hand spectrum is the geometric realization of the simplicial symmetric spectrum that in degree k is given by the k -fold smash product of $S \ltimes R[k]$ and the map is induced by the canonical map $R[k] \rightarrow I$. The skeleton filtration gives rise to a first quadrant spectral sequence

$$E_{s,t}^2 = H_s([k] \mapsto H_t(B(S, S \ltimes R[k], S)[k], \mathbb{Z}/p)) \Rightarrow H_{s+t}(B(S, S \ltimes R, S), \mathbb{Z}/p).$$

Since $R[k]$ is a free abelian group, the canonical maps

$$\begin{aligned} H_t(B(S, S \ltimes R[k], S)[k], \mathbb{Z}/p) &\leftarrow B(\mathbb{Z}, \mathbb{Z} \ltimes R[k], \mathbb{Z})[k] \otimes H_t(\mathbb{Z}, \mathbb{Z}/p) \\ &\rightarrow B(\mathbb{Z}, \mathbb{Z} \ltimes R[k], \mathbb{Z}/p)[k] \otimes H_t(\mathbb{Z}, \mathbb{Z}/p) \end{aligned}$$

are isomorphisms. Therefore, we can write

$$E_{s,t}^2 = H_s(B(\mathbb{Z}, \mathbb{Z} \ltimes R[-], \mathbb{Z}/p)[-]) \otimes H_t(\mathbb{Z}, \mathbb{Z}/p).$$

It remains to prove that

$$H_s(B(\mathbb{Z}, \mathbb{Z} \ltimes R[-], \mathbb{Z}/p)[-]) = \mathrm{Tor}_s^{\mathbb{Z} \ltimes I}(\mathbb{Z}, \mathbb{Z}/p).$$

We recall that the canonical map of simplicial abelian groups

$$B(\mathbb{Z}, \mathbb{Z} \ltimes R[-], \mathbb{Z} \ltimes R[-])[-] \rightarrow \mathbb{Z}$$

is a weak equivalence. Moreover, since $R[k]$ is a free abelian group, for all $k \geq 0$, this map is a projective resolution in the sense of [22, p. 6.7] of the simplicial $\mathbb{Z} \ltimes R[-]$ -module $\mathbb{Z}$. It follows that the simplicial abelian group $B(\mathbb{Z}, \mathbb{Z} \ltimes R[-], \mathbb{Z}/p)$ represents the derived tensor-product $\mathbb{Z} \otimes_{\mathbb{Z} \ltimes R[-]}^{\mathbb{L}} \mathbb{Z}/p$. Finally, the Künneth spectral sequence of [22, §6, Thm. 6(b)] collapses to yield an isomorphism

$$\mathrm{Tor}_s^{\mathbb{Z} \ltimes I}(\mathbb{Z}, \mathbb{Z}/p) \xrightarrow{\sim} H_s(\mathbb{Z} \otimes_{\mathbb{Z} \ltimes R[-]}^{\mathbb{L}} \mathbb{Z}/p).$$

This completes the proof. $\square$

PROOF OF THM. 2.1.6. We prove that the canonical map

$$\{\mathrm{TH}_q(I^m, \mathbb{Z}/p)\}_{m \geq 1} \rightarrow \{\mathrm{TH}_q(A, I^m, \mathbb{Z}/p)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups, for all q . It suffices by [21, Cor. 5.8] to show that the canonical map

$$\{H_q(\mathrm{TH}(I^m), \mathbb{Z}/p)\}_{m \geq 1} \rightarrow \{H_q(\mathrm{TH}(A, I^m), \mathbb{Z}/p)\}_{m \geq 1}$$

is an isomorphism of pro-abelian groups, for all q . This map is the edge-homomorphism of the following first-quadrant spectral sequence of pro-abelian groups.

$$E_{s,t}^1 = \{H_t(\mathrm{gr}_s \mathrm{TH}(A, I^m), \mathbb{Z}/p)\} \Rightarrow \{H_{s+t}(\mathrm{TH}(A, I^m), \mathbb{Z}/p)\}.$$

Hence, it is enough to show that the pro-abelian groups $E_{s,t}^1$ are zero, for $s > 0$. We have a direct sum decomposition

$$E_{s,t}^1 = H_t(\mathrm{gr}_s \mathrm{TH}(A, I^m)_0^\dagger, \mathbb{Z}/p) \oplus H_t(\mathrm{gr}_s \mathrm{TH}(A, I^m)_1^\dagger, \mathbb{Z}/p)$$

and an additional first-quadrant spectral sequence

$${}^v E_{u,v}^1 = \{H_v(\mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I^m)_1^\dagger, \mathbb{Z}/p)\} \Rightarrow \{H_{u+v}(\mathrm{gr}_s \mathrm{TH}(A, I^m)_1^\dagger, \mathbb{Z}/p)\}.$$

It follows that to prove the theorem, it will suffice to show that the pro-abelian groups $\{H_q(\mathrm{gr}_s \mathrm{TH}(A, I^m), \mathbb{Z}/p)_0^\dagger\}$ and $\{H_q(\mathrm{gr}'_u \mathrm{gr}_s \mathrm{TH}(A, I^m), \mathbb{Z}/p)_1^\dagger\}$ are zero, for all positive integers s and all non-negative integers u and q . But this is an immediate consequence of Lemmas 2.3.3 and 2.4.1, since we assumed that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^m)}(\mathbb{Z}, \mathbb{Z}/p)\}$ are zero, for all positive integers q . $\square$

3. Proof of Theorems A and B

3.1. In this section we complete the proof of Thm. A. The following result is similar to the proof of [6, Thm. 5.3].

LEMMA 3.1.1. *If the statement Thm. A holds whenever I can be embedded as a two-sided ideal of a free unital associative ring, then it holds in general.*

PROOF. Suppose that X is a functor from unital associative rings to symmetric spectra that satisfies that for every pair of rings (A, B) the canonical map

$$X(A \times B) \rightarrow X(A) \times X(B)$$

is a weak equivalence. We can then extend X to a functor X' defined on non-unital associative rings as follows. If I is a non-unital associative ring, we define $X'(I)$ to be the homotopy fiber of the map $X(\mathbb{Z} \ltimes I) \rightarrow X(\mathbb{Z})$ induced by the canonical projection. If A is a unital associative ring, the map $\mathbb{Z} \ltimes A \rightarrow \mathbb{Z} \times A$ that takes (x, a) to $(x, x \cdot e + a)$ is a ring-isomorphism, and hence, we obtain a weak equivalence

$$X(\mathbb{Z} \ltimes A) \xrightarrow{\sim} X(\mathbb{Z} \times A) \xrightarrow{\sim} X(\mathbb{Z}) \times X(A)$$

of symmetric spectra over $X(\mathbb{Z})$. These maps induce a weak equivalence of the homotopy fiber $X'(A)$ of the map $X(\mathbb{Z} \ltimes A) \rightarrow X(\mathbb{Z})$ and the homotopy fiber $X''(A)$ of the canonical projection from $X(\mathbb{Z}) \times X(A)$ to $X(\mathbb{Z})$. This shows that we have natural weak equivalences

$$X'(A) \xrightarrow{\sim} X''(A) \xleftarrow{\sim} X(A),$$

and hence, the functor X' extends, up to canonical weak equivalence, the functor X to non-unital associative rings as claimed. In the following, we will write X instead of X' for the extended functor.

Let $X^n(A)$ be the homotopy fiber of the cyclotomic trace

$$K(A) \rightarrow \mathrm{TC}^n(A; p).$$

Then Thm. A is equivalent to the statement that for every unital associative ring A , for every two-sided ideal $I \subset A$, and for all integers q , the canonical map

$$\{X_q^n(I, \mathbb{Z}/p^v)\}_{n \geq 1} \rightarrow \{X_q^n(A, I, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups. We let $F(A)$ denote the free unital associative ring generated by the underlying set of A . There is a canonical surjective ring-homomorphism $F(A) \rightarrow A$. We consider the following diagram of non-unital associative rings, where the rows and columns of the underlying diagram of abelian groups are all short-exact sequences.

$$\begin{array}{ccccc} J(A, I) & \longrightarrow & J(A) & \longrightarrow & J(A/I) \\ \downarrow & & \downarrow & & \downarrow \\ F(A, I) & \longrightarrow & F(A) & \longrightarrow & F(A/I) \\ \downarrow & & \downarrow & & \downarrow \\ I & \longrightarrow & A & \longrightarrow & A/I. \end{array}$$

Since the non-unital rings in the top row can be embedded as ideals of free unital associative rings, the diagram gives rise to the following map of long-exact sequences

of pro-abelian groups.

$$\begin{array}{ccc}
\vdots & & \vdots \\
\downarrow & & \downarrow \\
\{X_q^n(J(A, I), \mathbb{Z}/p^v)\} & \xrightarrow{f'_*} & \{X_q^n(J(A), J(A, I), \mathbb{Z}/p^v)\} \\
\downarrow & & \downarrow \\
\{X_q^n(F(A, I), \mathbb{Z}/p^v)\} & \xrightarrow{f_*} & \{X_q^n(F(A), F(A, I), \mathbb{Z}/p^v)\} \\
\downarrow & & \downarrow \\
\{X_q^n(I, \mathbb{Z}/p^v)\} & \xrightarrow{f''_*} & \{X_q^n(A, I, \mathbb{Z}/p^v)\} \\
\downarrow & & \downarrow \\
\vdots & & \vdots
\end{array}$$

Finally, since the non-unital associative rings $J(A, I)$ and $F(A, I)$ can be embedded in a free unital associative ring, the maps labelled f'_* and f_* are isomorphisms of pro-abelian groups. Hence, so is the map labelled f''_* . This completes the proof. $\square$

We recall the following result of Cuntz and Quillen [6, Prop. 4.1]. Let k be a field, and let $I \subset T$ be a two-sided ideal of a free unital associative k -algebra. Then there exists a k -linear section

$$(3.1.2) \quad \alpha: I \rightarrow T \otimes_k I$$

of the multiplication $\mu: T \otimes_k I \rightarrow I$ such that the following diagram commutes.

$$\begin{array}{ccc}
T \otimes_k I & \xrightarrow{\text{id} \otimes_k \alpha} & T \otimes_k T \otimes_k I \\
\downarrow \mu & & \downarrow \mu \otimes_k \text{id} \\
I & \xrightarrow{\alpha} & T \otimes_k I.
\end{array}$$

This is stated in *loc.cit.* for I is a two-sided ideal of free unital associative $\mathbb{C}$ -algebra, but the proof works for every ground field. The diagram shows, in particular, that the map α takes I^2 to $I \otimes_k I$. We use this to prove the following result.

PROPOSITION 3.1.3. *Let $I \subset T$ be a two-sided ideal of a free unital associative ring. Then the pro-abelian groups*

$$\{\text{Tor}_q^{\mathbb{Z} \ltimes (I^m)}(\mathbb{Z}, \mathbb{Z}/p^v)\}_{m \geq 1}$$

are zero, for all positive integers q .

PROOF. We recall that the bar-complex $B_*(I)$ of the non-unital associative ring I is defined to be the chain complex that is zero in degree zero and the n -fold tensor product $I \otimes \cdots \otimes I$ in positive degree n and with differential

$$b'(x_1 \otimes \cdots \otimes x_n) = \sum_{0 < i < n} (-1)^i x_1 \otimes \cdots \otimes x_i x_{i+1} \otimes \cdots \otimes x_n.$$

Since $\mathbb{Z} \ltimes (I^m)$ is flat over $\mathbb{Z}$, the group $\text{Tor}_q^{\mathbb{Z} \ltimes (I^m)}(\mathbb{Z}, \mathbb{Z}/p^v)$ is canonically isomorphic to the q th homology group of the bar-complex $B_*(I^m) \otimes \mathbb{Z}/p^v$. Hence, it suffices to

show that for every $m \geq 1$, there exists $m' \geq m$ such that the chain-map induced from the canonical inclusion

$$B_*(I^{m'}) \otimes \mathbb{Z}/p^v \rightarrow B_*(I^m) \otimes \mathbb{Z}/p^v$$

induces the zero map on homology. Since also $I^m \subset T$ is a two-sided ideal of a free unital associative ring, it will suffice to treat the case $m = 1$. By simple induction, it also suffices to consider the case $v = 1$.

The tensor-product $I \otimes \mathbb{F}_p$ is a non-unital associative $\mathbb{F}_p$ -algebra, and there is a canonical isomorphism of complexes of $\mathbb{F}_p$ -vector spaces

$$B_*(I) \otimes \mathbb{F}_p \xrightarrow{\sim} B_*(I \otimes \mathbb{F}_p).$$

Let $(I \otimes \mathbb{F}_p)'$ and $\overline{I \otimes \mathbb{F}_p}$ be the kernel and image of the map $I \otimes \mathbb{F}_p \rightarrow T \otimes \mathbb{F}_p$ such that we have the following extension of non-unital associative $\mathbb{F}_p$ -algebras.

$$(I \otimes \mathbb{F}_p)' \rightarrow I \otimes \mathbb{F}_p \rightarrow \overline{I \otimes \mathbb{F}_p}.$$

The non-unital associative $\mathbb{F}_p$ -algebra $\overline{I \otimes \mathbb{F}_p}$ is a two-sided ideal of the free unital associative $\mathbb{F}_p$ -algebra $T \otimes \mathbb{F}_p$. Let

$$\alpha: \overline{I \otimes \mathbb{F}_p} \rightarrow (T \otimes \mathbb{F}_p) \otimes (\overline{I \otimes \mathbb{F}_p})$$

be the section of the multiplication from (3.1.2). Then the formula

$$s(x_1 \otimes \cdots \otimes x_n) = (-1)^n x_1 \otimes \cdots \otimes x_{n-1} \otimes \alpha(x_n)$$

defines a null-homotopy of the chain map

$$B_*(\overline{I^2 \otimes \mathbb{F}_p}) = B_*(\overline{(I \otimes \mathbb{F}_p)^2}) \rightarrow B_*(\overline{I \otimes \mathbb{F}_p})$$

induced by the canonical inclusion.

We next let $K_*(I \otimes \mathbb{F}_p)$ be the kernel complex of the following chain-map induced by the canonical projection.

$$B_*(I \otimes \mathbb{F}_p) \rightarrow B_*(\overline{I \otimes \mathbb{F}_p}).$$

The $\mathbb{F}_p$ -vector space $K_n(I \otimes \mathbb{F}_p)$ is generated by the tensors $x_1 \otimes \cdots \otimes x_n$, where at least one of $x_1, \dots, x_n$ is in $(I \otimes \mathbb{F}_p)'$. We claim that the chain-map

$$K_*(I^2 \otimes \mathbb{F}_p) \rightarrow K_*(I \otimes \mathbb{F}_p)$$

induced by the canonical inclusion is equal to zero. This follows immediately from the statement that the following map induced by the canonical inclusion is zero.

$$\begin{array}{ccc} (I^2 \otimes \mathbb{F}_p)' & \longrightarrow & (I \otimes \mathbb{F}_p)' \\ \parallel & & \parallel \\ (I^2 \cap pT)/pI^2 & \longrightarrow & (I \cap pT)/pI. \end{array}$$

Hence, it suffices to show that $I^2 \cap pT \subset pI$, or equivalently, that if $x \in I$ and $y \in I$, and if $xy \in pT$, then $xy \in pI$. We claim that $xy \in pT$ implies that $x \in pT$ or $y \in pT$. If $x \in pT$, we write $x = px'$ with $x' \in T$. Then $xy = px'y$ and since I is a left ideal, $x'y \in I$. Similarly, if $y \in pT$, we write $y = py'$ with $y' \in T$. Then $xy = pxy'$, and since I is a right ideal, $xy' \in I$. In either case, $xy \in pI$ as desired.

It remains to prove that $xy \in pT$ implies that $x \in pT$ or $y \in pT$, or equivalently, that if the class $\bar{x}\bar{y} \in T/pT$ is zero then one of the classes $\bar{x} \in T/pT$ and $\bar{y} \in T/pT$ is zero. Suppose conversely that both $\bar{x} \in T/pT$ and $\bar{y} \in T/pT$ are non-zero. The

ring T/pT is a free unital associative $\mathbb{F}_p$ -algebra. Hence, we can write $\bar{x}$ and $\bar{y}$ uniquely as sums

$$\bar{x} = \sum_{\omega} a_{\omega} \omega, \quad \bar{y} = \sum_{\omega} b_{\omega} \omega,$$

where ω runs over all finite words with letters in a set of generators of T/pT , and where $a_{\omega}, b_{\omega} \in \mathbb{F}_p$ are zero for all but finitely many ω . The multiplication in T/pT concatenates words and multiplies coefficients. Let $\omega_1, \dots, \omega_m$ and $\omega'_1, \dots, \omega'_n$ be shortest words that occur with non-zero coefficients in $\bar{x}$ and $\bar{y}$, respectively. Then the words $\omega_1 \omega'_1, \dots, \omega_1 \omega'_n, \dots, \omega_m \omega'_1, \dots, \omega_m \omega'_n$ are all different and are exactly the shortest words that occur with a non-zero coefficient in $\bar{x}\bar{y}$. In particular, $\bar{x}\bar{y} \in T/pT$ is non-zero as desired.

Finally, we consider now the following diagram with long-exact columns.

$$\begin{array}{ccccc} \vdots & & \vdots & & \vdots \\ \downarrow & & \downarrow & & \downarrow \\ H_q(K_*(I^{4m} \otimes \mathbb{F}_p)) & \xrightarrow{0} & H_q(K_*(I^{2m} \otimes \mathbb{F}_p)) & \xrightarrow{0} & H_q(K_*(I^m \otimes \mathbb{F}_p)) \\ \downarrow & & \downarrow & & \downarrow \\ H_q(B_*(I^{4m} \otimes \mathbb{F}_p)) & \longrightarrow & H_q(B_*(I^{2m} \otimes \mathbb{F}_p)) & \longrightarrow & H_q(B_*(I^m \otimes \mathbb{F}_p)) \\ \downarrow & & \downarrow & & \downarrow \\ H_q(B_*(\overline{I^{4m} \otimes \mathbb{F}_p})) & \xrightarrow{0} & H_q(B_*(\overline{I^{2m} \otimes \mathbb{F}_p})) & \xrightarrow{0} & H_q(B_*(\overline{I^m \otimes \mathbb{F}_p})) \\ \downarrow & & \downarrow & & \downarrow \\ \vdots & & \vdots & & \vdots \end{array}$$

The top and bottom horizontal maps are zero by what was proved above, and a diagram chase then shows that the composite of the two middle horizontal maps is zero. This completes the proof. $\square$

LEMMA 3.1.4. *Let $f: A \rightarrow B$ be a map of unital associative rings, and let $I \subset A$ be a two-sided ideal such that $f: I \rightarrow f(I)$ is an isomorphism onto an ideal of B . Suppose that the cyclotomic trace induces an isomorphism of pro-abelian groups*

$$\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1} \xrightarrow{\sim} \{\mathrm{TC}_q^m(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1},$$

for all integers q . Then the map induced by the cyclotomic trace

$$K_q(A, B, I, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(A, B, I; p, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups, for all integers q .

PROOF. We note that for all positive integers m and n , we have a commutative diagram of spectra in which the columns are exact triangles.

$$\begin{array}{ccc}
K(A, B, I^m) & \longrightarrow & \mathrm{TC}^n(A, B, I^m; p) \\
\downarrow & & \downarrow \\
K(A, B, I) & \longrightarrow & \mathrm{TC}^n(A, B, I; p) \\
\downarrow & & \downarrow \\
K(A/I^m, B/I^m, I/I^m) & \longrightarrow & \mathrm{TC}^n(A/I^m, B/I^m, I/I^m; p) \\
\downarrow & & \downarrow \\
K(A, B, I^m)[-1] & \longrightarrow & \mathrm{TC}^n(A, B, I^m; p)[-1].
\end{array}$$

The induced diagram of homotopy groups with $\mathbb{Z}/p^v$ -coefficients gives rise, as m and n vary, to a map of long-exact sequences of pro-abelian groups. The map of pro-abelian groups induced by the top horizontal map in the diagram above is an isomorphism by assumption, and we wish to show that the map of pro-abelian groups induced by the second horizontal map in the diagram above is an isomorphism. By a five-lemma argument this is equivalent to showing that the map of pro-abelian groups induced from the third horizontal map in the diagram above is an isomorphism. But it follows from [19] and from [8, Thm. 2.2.1] that for $m \geq 1$ fixed, the cyclotomic trace induces an isomorphism of pro-abelian groups

$$K_q(A/I^m, B/I^m, I/I^m, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(A/I^m, B/I^m, I/I^m; p, \mathbb{Z}/p^v)\}_{n \geq 1}.$$

But then the map of pro-abelian groups obtained by also allowing $m \geq 1$ to vary is an isomorphism. This completes the proof. $\square$

PROOF OF THM. A. By Lemma 3.1.1, we can assume that the non-unital associative ring I can be embedded as a two-sided ideal $I \subset T$ of a free unital associative ring. Moreover, by Lemma 3.1.4 it suffices to show that for all integers q , the cyclotomic trace induces an isomorphism of pro-abelian groups

$$\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1} \xrightarrow{\sim} \{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1}.$$

We claim that both pro-abelian groups are zero. Indeed, by Thm. 2.1.1 the pro-abelian groups $\{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m \geq 1}$ are zero, for all integers q and all positive integers n . Hence, the pro-abelian groups $\{\mathrm{TC}_q^n(A, B, I^m; p, \mathbb{Z}/p^v)\}_{m, n \geq 1}$ are zero, for all integers q as claimed. Finally, in order to show that also the pro-abelian groups $\{K_q(A, B, I^m, \mathbb{Z}/p^v)\}_{m \geq 1}$ are zero, for all integers q , it suffices by Thm. 1.1.1 to show that the pro-abelian groups $\{\mathrm{Tor}_q^{\mathbb{Z} \ltimes (I^m)}(\mathbb{Z}, \mathbb{Z}/p)\}_{m \geq 1}$ are zero, for all positive integers q . This was proved in Prop. 3.1.3 above. $\square$

3.2. In this section we prove Theorems B and D of the introduction.

PROOF OF THM. B. Let $X \rightarrow \mathrm{Spec} k$ be a curve over a field k , that is, a separated morphism of finite type with X of dimension one. We wish to show that for all $q \geq r$, the cyclotomic trace

$$K_q(X, \mathbb{Z}/p^v) \rightarrow \{\mathrm{TC}_q^n(X; p, \mathbb{Z}/p^v)\}_{n \geq 1}$$

is an isomorphism of pro-abelian groups. We first reduce to the case where X is affine. So assume that the theorem holds for affine curves. A covering $\{U_i\}$ of X by affine open subschemes gives rise to spectral sequences of pro-abelian groups

$$E_{s,t}^2 = \check{H}^{-s}(\{U_i\}, K_t(-, \mathbb{Z}/p^v)) \Rightarrow K_{s+t}(X, \mathbb{Z}/p^v)$$

and

$$E_{s,t}^2 = \{\check{H}^{-s}(\{U_i\}, \mathrm{TC}_t^n(-; p, \mathbb{Z}/p^v))\}_{n \geq 1} \Rightarrow \{\mathrm{TC}_{s+t}^n(X; p, \mathbb{Z}/p^v)\}_{n \geq 1}.$$

Indeed, for K -theory, this follows from [26, Thm. 10.3], and for topological cyclic homology, this follows directly from the definition of the topological cyclic homology of a scheme [9, Def. 3.2.3]. The cyclotomic trace induces a map of spectral sequences which, by assumption, is an isomorphism on $E_{s,t}^2$ for $t \geq r$. But then the cyclotomic trace also induces an isomorphism of the pro-abelian groups in the abutments in degrees $q \geq r$. So the theorem holds for X .

So assume that X is affine. By an argument similar to the proof of Lemma 3.1.4 above, we can further assume that X is reduced. By induction on the number of irreducible components in X , we can further reduce to the case where X is irreducible. Indeed, let Z and Z' be two proper closed subschemes of X and suppose that Z and Z' cover X , or equivalently, that the corresponding ideals I and I' of the coordinate ring A intersect trivially. Then it follows from Thm. A that the cyclotomic trace induces an isomorphism of pro-abelian groups

$$K_q(A, A/I', I, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(A, A/I', I; p, \mathbb{Z}/p^v)\}_{n \geq 1},$$

for all integers q . This proves the induction step.

Finally, let $X \rightarrow \mathrm{Spec} k$ be a reduced and irreducible affine curve. The coordinate ring A is an integral domain, and we let B be the integral closure. We claim that the conductor ideal $I = \{a \in A \mid aB \subset A\}$, which is the largest ideal of A that is also an ideal of B , is non-zero. Indeed, the extension $A \subset B$ is finite, so we can write $B = b_1A + \cdots + b_mA$. Since A and B have the same quotient field, there are elements $a_i \in A$ such that $a_i b_i \in A$, $1 \leq i \leq m$. But then the product $a_1 \cdots a_m$ is in I . The ring B is a regular k -algebra, and the quotients A/I and B/I are finite k -algebras. Hence, we know from [9, Thm. 4.2.2] and from [19] and [8, Thm. 2.2.1] that the cyclotomic trace induces an isomorphism of pro-abelian groups

$$K_q(R, \mathbb{Z}/p^v) \xrightarrow{\sim} \{\mathrm{TC}_q^n(R; p, \mathbb{Z}/p^v)\}_{n \geq 1},$$

for $q \geq r$, if R is one of B , A/I , and B/I . But then Thm. A shows that this map is an isomorphism, for $q \geq r$, also if $R = A$. $\square$

PROOF OF THM. D. The ideal $I = |G| \cdot \mathfrak{M}$ is a common ideal of $A = \mathbb{Z}[G]$ and $B = \mathfrak{M}$ [1, Cor. XI.1.2], and hence, we can apply Thm. A. It follows that the exactness of the sequence in the statement of Thm. D is equivalent to the exactness of the corresponding sequence with $\mathbb{Z}[G]$ and $\mathfrak{M}$ replaced by $\mathbb{Z}[G]/I$ and $\mathfrak{M}/I$. The latter are both artinian rings whose quotient by the Jacobson radical is a semi-simple $\mathbb{F}_p$ -algebra. Hence, the exactness of the latter sequence follows from [14, Thm. 5.1(i)] and from [8, Thm. 2.2.1]. $\square$

References

- [1] H. Bass, *Algebraic K-theory*, W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [2] A. J. Berrick and W. G. Dwyer, *The spaces that define algebraic K-theory*, *Topology* **39** (2000), 225–228.

- [3] M. Bökstedt and I. Madsen, *Topological cyclic homology of the integers*, *K-theory* (Strasbourg, 1992), Astérisque, vol. 226, 1994, pp. 57–143.
- [4] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Math., vol. 304, Springer-Verlag, New York, 1972.
- [5] G. Cortiñas, *The obstruction to excision in K-theory and in cyclic homology*, Preprint 2001, math.KT/0111096.
- [6] J. Cuntz and D. Quillen, *Excision in bivariant periodic cyclic cohomology*, *Invent. Math.* **127** (1997), 67–98.
- [7] R. K. Dennis and M. I. Krusemeyer, $K_2(A[X, Y]/(XY))$, a problem of Swan, and related computations, *J. Pure Appl. Alg.* **15** (1979), 125–148.
- [8] T. Geisser and L. Hesselholt, *On the K-theory and topological cyclic homology of smooth schemes over a discrete valuation ring*, *Trans. Amer. Math. Soc.*, to appear.
- [9] ———, *Topological cyclic homology of schemes*, *K-theory* (Seattle, 1997), *Proc. Symp. Pure Math.*, vol. 67, 1999, pp. 41–87.
- [10] S. Geller, L. Reid, and C. A. Weibel, *The cyclic homology and K-theory of curves*, *Bull. Amer. Math. Soc.* **15** (1986), no. 2, 205–208.
- [11] ———, *The cyclic homology and K-theory of curves*, *J. reine angew. Math.* **393** (1989), 39–90.
- [12] L. Hesselholt, *K-theory of truncated polynomial algebras*, *Handbook of K-theory*, Springer-Verlag, New York, to appear.
- [13] ———, *On the K-theory of the coordinate axes in the affine plane over a field of positive characteristic*, in preparation.
- [14] L. Hesselholt and I. Madsen, *On the K-theory of finite algebras over Witt vectors of perfect fields*, *Topology* **36** (1997), 29–102.
- [15] ———, *On the K-theory of nilpotent endomorphisms*, *Homotopy methods in algebraic topology* (Boulder, CO, 1999), *Contemp. Math.*, vol. 271, Amer. Math. Soc., Providence, RI, 2001, pp. 127–140.
- [16] ———, *On the K-theory of local fields*, *Ann. of Math.* **158** (2003), 1–113.
- [17] P. S. Hirschhorn, *Model categories and their localizations*, *Mathematical Surveys and Monographs*, vol. 99, American Mathematical Society, Providence, RI, 2003.
- [18] M. Hovey, B. Shipley, and J. Smith, *Symmetric spectra*, *J. Amer. Math. Soc.* **13** (2000), 149–208.
- [19] R. McCarthy, *Relative algebraic K-theory and topological cyclic homology*, *Acta Math.* **179** (1997), 197–222.
- [20] R. Oliver, *Whitehead groups of finite groups*, *London Mathematical Society Lecture Note Series*, vol. 132, Cambridge University Press, Cambridge, 1988.
- [21] I. A. Panin, *On a theorem of Hurewicz and K-theory of complete discrete valuation rings*, *Math. USSR Izvestiya* **29** (1987), 119–131.
- [22] D. Quillen, *Homotopical algebra*, *Lecture Notes in Math.*, vol. 43, Springer-Verlag, New York, 1967.
- [23] B. Shipley, *Symmetric ring spectra and topological Hochschild homology*, *K-theory* **19** (2000), 155–183.
- [24] A. A. Suslin, *Excision in the integral algebraic K-theory*, *Proc. Steklov Inst. Math.* **208** (1995), 255–279.
- [25] A. A. Suslin and M. Wodzicki, *Excision in algebraic K-theory*, *Ann. of Math.* **136** (1992), 51–122.
- [26] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, *Grothendieck Festschrift, Volume III*, *Progress in Mathematics*, vol. 88, 1990, pp. 247–435.
- [27] J. B. Wagoner, *Delooping classifying spaces in algebraic K-theory*, *Topology* **11** (1972), 349–370.
- [28] C. A. Weibel, *An introduction to homological algebra*, *Cambridge Studies in Advanced Mathematics*, vol. 38, Cambridge University Press, Cambridge, 1994.
- [29] M. Wodzicki, *Excision in cyclic homology and in rational algebraic K-theory*, *Ann. of Math.* **129** (1989), 591–639.

UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CALIFORNIA
E-mail address: `geisser@math.usc.edu`

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MASSACHUSETTS
E-mail address: `larsh@math.mit.edu`