AN INVARIANT FOR OPEN VIRTUAL STRINGS

Daniel S. Silver and Susan G. Williams

ABSTRACT: Extended Alexander groups are used to define an invariant for open virtual strings. Examples of non-commuting open strings and a ribbon-concordance obstruction are given. An example is given of a slice virtual open string that is not ribbon. Definitions are extended to open *n*-strings.

1. Introduction. A classical open string is the image α of a generic immersion $\mathbb{R} \to \mathbb{R}^2$ such that $x \mapsto (x, 0)$ for |x| sufficiently large. When some double points are encircled, α is said to be a virtual open string (or simply, an open string). Two open strings are regarded as the same if they are related by a finite sequence of flat generalized Reidemeister moves (see Figure 1). The set \mathcal{M} of open strings is a semigroup under concatenation.



Figure 1: Flat generalized Reidemeister moves

Gauss diagrams provide an alternative means for defining open strings. Following [T04], we describe an open string by an oriented line with 2m distinguished points partitioned into m ordered pairs. A pair (s, t) is the preimage of an uncircled double point, and the segment of R near s crosses the segment near t transversely from left to right. We visualize each ordered pair as as an arrow above the line starting at the first coordinate of the pair and ending at the second. An equivalence relation on Gauss diagrams, called *homotopy*, is described in [T04], where it is shown that two open strings are the same if and only if their underlying Gauss diagrams are homotopic.

Remarks 1.1. (Non-open) strings are defined by Gauss diagrams in an analogous manner, replacing the line with an oriented circle [T04].

Strings and open strings represent a natural variation of the theory of virtual knots, introduced by L. Kauffman [K99]. Strings correspond to flat virtual knots while open strings can be regarded

Both authors partially supported by NSF grant DMS-0304971.

²⁰⁰⁰ Mathematics Subject Classification. Primary 57M25; Secondary 20F05, 20F34.

as flat virtual 1-tangles [K99]. Our terminology was suggested by V. Turaev, who has borrowed "open string" from physics and "virtual" from Kauffman's theory.



Figure 2: A virtual open string α and its Gauss diagram

In [SW04] we adapted Alexander group techniques of [SW01'] and [SW03] in order to define invariants for *long virtual knots*, open strings for which each uncircled double point is resolved as a classical crossing and the usual generalized Reidemeister moves for virtual knots are allowed. Like open strings, long virtual knots form a semigroup, and the invariant of [SW04] provides the means to show that many pairs of long virtual knots do not commute. Earlier V. Manturov [M04] had used quandle techniques to prove noncommutativity for a particular pair, and [SW04] was motivated by his work.

We modify the invariant of [SW04] to obtain a homomorphism Φ from \mathcal{M} to the group of endomorphisms of a countable rank free group. Using it, we show that \mathcal{M} is non-commutative, thereby answering a question posed in [T04]. In §3 we show that $\Phi(\alpha)$ has a special form whenever α is a ribbon open knot, and we give an example of a slice open knot that isn't ribbon. In the last section, we extend the invariant Φ to open multi-links.

We are grateful to Vladimir Turaev for questions, suggestions and encouragement. We also thank Seiichi Kamada and Naoko Kamada for valuable conversations.

2. Extended Alexander groups and open strings. The Alexander group \mathcal{A}_k of a virtual knot was introduced in [SW01']. It is a special case of R. Crowell's derived group [C84]. When k is classical, \mathcal{A}_k abelianizes to the Alexander module of the knot, a fact that provides justification for the name we have given it. The extended Alexander group $\tilde{\mathcal{A}}_k$, also defined in [SW01'], contains new information, and in many cases it can show that a virtual knot is not classical.

Extended Alexander groups are defined for long virtual knots in [SW04]. In order to obtain an invariant of an open string α , we apply the following observation of Turaev. Give α the induced orientation from \mathbb{R} , directing the open string from left to right. Let D be the associated descending diagram for a long virtual knot k, that is, while traveling along α in the preferred direction, each time an uncircled double point is encounted for the first time, resolve it as a classical crossing in such a way that we travel along the overcrossing arc. Encircled double points are treated as virtual crossings. One easily verifies that if two open strings are related by a flat generalized Reidemeister move, then the associated long virtual knots are related by a corresponding move. Consequently, invariants of k_{α} provide invariants for α .

Each overcrossing arc of the descending diagram D is regarded as a union of two arcs joined at the point of overcrossing. Virtual crossings are disregarded when determining arcs. As in [SW04], associate to k a group $\tilde{\mathcal{A}}_k$. It is described by families of generators $a_{\mathbf{n}}, b_{\mathbf{n}}, c_{\mathbf{n}}, \ldots$, each indexed by vectors $\mathbf{n} \in \mathbb{Z}^2$, corresponding to the arcs of D. To a classical positive crossing labeled as in Figure 3, associate a similarly indexed family of relations of the form

$$a_{\mathbf{n}}b_{\mathbf{n}+u} = c_{\mathbf{n}}a_{\mathbf{n}+u}, \ c_{\mathbf{n}+v} = b_{\mathbf{n}} \tag{2.1}$$

For a negative crossing, the second relator family is replaced by $a_{\mathbf{n}+v} = d_{\mathbf{n}}$. Here u, v are standard generators of \mathbb{Z}^2 .

It is convenient to abbreviate generator families by a, b, c, \ldots , and $a_{\mathbf{n}}b_{\mathbf{n}+u} = c_{\mathbf{n}}a_{\mathbf{n}+u}, a_{\mathbf{n}+v} = d_{\mathbf{n}}$ by $ab^u = ca^u$ and $a^v = d$, respectively. In this notation, $(j, k) \in \mathbb{Z}^2$ is written multiplicatively as $u^j v^k$. We regard $\tilde{\mathcal{A}}_{\ell}$ as a \mathbb{Z}^2 -group, a group together with an action $\mathbb{Z}^2 \to \operatorname{Aut}(\tilde{\mathcal{A}}_k)$.



Figure 3: Relations for extended Alexander group

Let $a_{-\infty}$ and $a_{+\infty}$ be generators corresponding to the arcs that run to $-\infty$ and $+\infty$, respectively. We refer to these as left and right *end generators*. Associate a triple $(\tilde{\mathcal{A}}_{\ell}, a_{-\infty}, a_{+\infty})$. Two such triples are regarded as the same if there is an isomorphism of underlying groups that commutes with the \mathbb{Z}^2 -actions and matches corresponding end generators. As in [SW04], the triple is an invariant of k.

Since the diagram D is descending, it follows easily that $a_{-\infty}$ generates $\tilde{\mathcal{A}}_k$. In particular, $a_{+\infty}$ is uniquely expressible as a reduced word in the free group F on generators $a_{-\infty}$. The correspondence determines an endomorphism $\Phi(\alpha)$ of F that respects the \mathbb{Z}^2 -action. Moreover, as in [SW04], if α_1 and α_2 are two open strings, then $\Phi(\alpha_1 \cdot \alpha_2)$ is the composition of $\Phi(\alpha_1)$ and $\Phi(\alpha_2)$.

We summarize the above results as follows.

Theorem 2.1. The assignment $\alpha \mapsto \Phi(\alpha)$ defines a homomorphism $\Phi : \mathcal{M} \to \text{End}(F)$.

Remark 2.2. Φ is similar in certain respects to the Burau representation for braids. The latter can be regarded as an automorphism of the \mathbb{R} -vector space spanned by the braid inputs. The image of each basis element is seen at the braid outputs. This point of view can be found, for example, in [SW01].

An open string that has no circled double points is called *classical*. The descending diagram D will be trivial in this case. The following criterion is immediate.

Corollary 2.3. If α is a classical open string, then $\Phi(\alpha)$ is the identity map.

Example 2.4 Consider the open strings α_1 and α_2 in Figure 4 together with the associated long virtual knots k_1 and k_2 . The knot diagrams have been labeled with generators of the extended Alexander groups. For notational convenience, we have abbreviated $a_{-\infty}$ by a.

The endomorphism $\Phi(\alpha_1)$ maps a to $\bar{a}^{u^{-1}v^{-2}}\bar{a}^{u^{-2}v^{-1}}a^{u^{-2}v^{-2}}a^{u^{-1}}a^{v^{-1}}$ while $\Phi(\alpha_2)$ maps a to $a^v a^u a^{u^2v^2}\bar{a}^{u^2v}\bar{a}^{uv^2}$, where denotes inversion. By Corollary 2.3, neither α_1 nor α_2 is classical.

We employ the functional notation (x)f instead of f(x) so that the composition "f followed by g" is denoted by fg. It is straightforward to compute $\Phi(\alpha_i \alpha_j) = \Phi(\alpha_i)\Phi(\alpha_j)$, i = 1, 2. Simply substitute $(a)\Phi(\alpha_i)$ for each occurrence of a in $(a)\Phi(\alpha_j)$. One checks that $\Phi(\alpha_1 \cdot \alpha_2)$ is not equal to $\Phi(\alpha_2 \cdot \alpha_1)$. Hence the open strings α_1 and α_2 do not commute.



Figure 4: Non-commuting α_1 (top), α_2 (bottom) and associated long virtual knots

Remark 2.5. In the above example, $a_{+\infty}$ has the form $Wa^{(uv)^{-\omega}}\overline{W}^{uv}$, where W is a word in generators $a_{-\infty}$ and ω is equal the sum of signs of classical crossings. The fact holds generally for any open string, but we will not use it here. The proof is left to the reader.

Example 2.6. The representation Φ is not faithful. Consider the open string β in Figure 5. We leave it to the reader to check that $\Phi(\beta)$ is trivial. On the other hand, if we close the associated long virtual knot k, we obtain a virtual knot \hat{k} that has nontrivial Jones polynomial. The latter claim can be seen easily, since \hat{k} is a connected sum of two simpler virtual knots, each of which has nontrivial Jones polynomial. Hence α is a nontrivial open string.



Figure 5: Open string β

3. Ribbon open strings. As discussed in §1, open strings can be defined either by generic immersions $\mathbb{R} \to \mathbb{R}^2$ or by Gauss diagrams. They can also be defined in terms of generic paths in surfaces, a third perspective that motivates the notion of cobordism. Slice and ribbon open strings can be defined, and as in classical knot theory, ribbon implies slice [T04].

Definition 3.1. An open string is *ribbon* if it has a Gauss diagram for which reflection about the origin takes each arrow to an arrow with opposite direction.

The polynomial invariant in §3.2 of [T04] provides is an elementary obstruction for an open string to be slice (and hence an obstruction for it to be ribbon). It vanishes for the non-ribbon open string α in the following example.

Example 3.2. Consider the ribbon open strings α_3 and α_4 and their Gauss diagrams in Figure 6. Let α be the product $\alpha_3 \cdot \alpha_4$. A Gauss diagram for α is obtained by concatenating the Gauss

diagrams for α_3 and α_4 . Reflecting the Gauss diagram for α and changing the direction of each arrow produces a Gauss diagram for $\alpha_4 \cdot \alpha_3$. Hence if α is ribbon, then $\alpha_3 \cdot \alpha_4$ is equal to $\alpha_4 \cdot \alpha_3$.

Computing $\Phi(\alpha_3)$ and $\Phi(\alpha_4)$, as in Example 2.4 and using similar notation, we find that $(a)\Phi(\alpha_3) = a^v a^u a^{u^2v^2} \bar{a}^{u^2v} \bar{a}^{uv^2}$ while $(a)\Phi(\alpha_4) = a a^{uv^2} a^{u^2v} a^{u^3v^3} \bar{a}^{u^3v^2} \bar{a}^{u^2v^3} \bar{a}^{uv}$. The reader can check that $\Phi(\alpha_3)$ and $\Phi(\alpha_4)$ are non-commuting endomorphisms. Hence α_3 and α_4 do not commute, and so α is not ribbon.

Since ribbon implies slice for open strings, and since a product of slice open strings is slice, α is an example of a slice open string that is not ribbon.



Figure 6: Open strings α_3 (top) and α_4 (bottom)

An obstruction for a general open string α to be ribbon can be described. First compute $\Phi(\alpha)$. Then reflect the Gauss diagram for α in the plane about the *y*-axis and change the direction of each arrow, thereby obtaining a second open string. If its image under Φ does not agree with $\Phi(\alpha)$, then α cannot be ribbon.

Testing to see if a given open string is ribbon requires two, sometimes tedious calculations. However, by considering the abelianized endomorphism Φ , we obtain an obstruction that requires only one, relatively easy calculation, and yet remains very effective. In order to describe it, we let $\phi(\alpha)(u, v) \in \mathbb{Z}[u^{\pm 1}, v^{\pm 1}]$ be the Laurent polynomial, well defined up to multiplication by units $u^{j}v^{k}$, given as the sum of the exponents of $(a)\Phi(\alpha)$. For example, the ribbon open string α_{3} in Example 3.2 has polynomial $v + u + u^{2}v^{2} - u^{2}v - uv^{2}$.

Proposition 3.3. If α is an open ribbon string, then $\phi(\alpha)(u, v) = \phi(\alpha)(v, u)$.

Before proving Proposition 3.3, we note that changing the directions of all arrows of a Gauss diagram for an open string α is equivalent to reflecting α about a line in the plane. (This is easily seen by choosing the line to be the *x*-axis.) We denote the reflected open string by α *. The descending diagrams D, D^* for the associated long virtual knots are also reflections of each other, and the effect of reflection on the invariant Φ is easily described. If W is a word in the generators $a_{\mathbf{n}} = a^{u^i v^k}$, let W^* be the word obtained by writing W in reverse order and then replacing u, v by u^{-1}, v^{-1} , respectively.

Lemma 3.4. $(a)\Phi(\alpha *) = [(a)\Phi(\alpha)]^*$.

Proof. The arc labeling of D^* is obtained from that of D by the involution $x \mapsto x^*$. To see this, it suffices to check that if the arc labelings in Figure 7a satisfy the crossing relation (2.1), then so do those in the reflected diagram of Figure 7b. The calculation is routine.



Figure 7: Labeled crossing diagram and its reflection

Proof of Proposition 3.3. Let k be the long virtual knot with descending diagram associated to α , and let $\kappa(k)$ be the virtual knot obtained by closing k. The difference $\phi(\alpha)(u, v) - 1$ is equal to the polynomial invariant $\Delta_0(u, v)$ for $\kappa(k)$ defined in [SW01']. Reflecting the Gauss diagram of α and closing the associated descending diagram produces a knot that is obtained from $\kappa(\alpha)$ by reversing its orientation and changing the sense of each classical crossing. In [SW03] we used a result of J. Sawollek [S02] to show that this has the effect of replacing $\Delta_0(u, v)$ with $\Delta_0(v^{-1}, u^{-1})$. Combining this result with Lemma 3.4, we see that if α is ribbon, then $\phi(\alpha)(u, v) = \phi(\alpha)(v, u)$.

Example 3.5. Consider the open string α described by the Gauss diagram in Figure 2. Let β be the product $\alpha \cdot \alpha^*$. As in Example 3.2, the slice obstruction in [T04] vanishes. We apply Proposition 3.3 to show that β is not ribbon.

We have $\phi(\beta)(u, v) = \phi(\alpha)(u, v) \cdot \phi(\alpha^*)(u, v) = \phi(\alpha)(u, v) \cdot \phi(\alpha)(u^{-1}, v^{-1})$. One easily computes that $\phi(\alpha)(u, v) = -uv^3 - u^3v^2 + u^3v^3 + u + v^2$, and from this that $\phi(\beta)(u, v)$ is not equal to $\phi(\beta)(v, u)$. Hence γ is not ribbon.

Remark 3.6. The abelian invariant ϕ cannot detect non-commutativity of open strings, and so it is not effective for the construction in Example 3.2.

4. Open virtual multi-strings. Open virtual strings generalize in several ways. For example, we can consider several open components.

An open virtual n-string (or simply open n-string) is the image α of a generic immersion $\mathbb{R} \times \{1, \ldots, n\} \to \mathbb{R}^2$ such that $(x, j) \mapsto (x, \pi_j)$ for |x| sufficiently large. Here (π_1, \ldots, π_n) is a permutation of $(1, \ldots, n)$. Some double points may be encircled, and as in the special case of open strings, when n = 1, two open n-strings are regarded as the same if they are related by a finite sequence of flat generalized Reidemeister moves (Figure 1).

In classical link theory, one often puts labels on components and keeps track of them. Similarly, we assign integers $1, 2, \ldots, n$ (called *colors*) to the components of an open *n*-string α , and we refer to α as a *colored open n-string*. The component of α labeled *i* is the *ith component*. We multiply two colored open *n*-strings only if the colors of each pair of joined components agree.

To each colored open *n*-string α we assign a \mathbb{Z}^{n+1} -group in a manner similar to that of §2. Begin by constructing a descending diagram D for a long virtual link ℓ : While traveling along the first component of α in the preferred direction, each time an uncircled double point of α is encounted for the first time, resolve it as a classical crossing in such a way that we travel along the overcrossing arc. Repeat with the remaining components in turn, similarly resolving the uncircled double points that remain. Double points are treated as virtual crossings. As before, one verifies that if two colored open *n*-strings are related by a flat generalized Reidemeister move, then the associated long virtual links are related by a corresponding move. Again, invariants of ℓ provide invariants of α .

Now define an extended Alexander group $\tilde{\mathcal{A}}_{\ell}$ by associating \mathbb{Z}^{n+1} -group generators to arcs of D. At each classical crossing, associate relators as in Figure 7.



Figure 8: Extended Alexander group relations for an open *n*-string

Let $a_{-\infty}$ and $a_{+\infty}$ be generators corresponding to the arcs on the first component of D that are leftmost and rightmost, respectively. Let $b_{-\infty}$ and $b_{+\infty}$ be similar generators for the second component, and so forth. We will refer to these as left and right *end generators*. Associate a tuple $(\tilde{\mathcal{A}}_{\ell}, a_{-\infty}, b_{-\infty}, \ldots, a_{+\infty}, b_{+\infty}, \ldots)$. Regard two such tuples as the same if there is an isomorphism of underlying groups that commutes with the \mathbb{Z}^{n+1} -actions and matches corresponding end generators. As in [SW04], the tuple is an invariant of ℓ . (This is proven only in the case that ℓ has a single component. However, the general argument is essentially the same.)

By design, the diagram D is descending. It follows as before that $a_{-\infty}, b_{-\infty}, \ldots$ generate $\hat{\mathcal{A}}_{\ell}$. In particular, $a_{+\infty}, b_{+\infty}, \ldots$ are uniquely expressible as words in the free group F on $a_{-\infty}, b_{-\infty}, \ldots$ Again the correspondence determines an endomorphism $\Phi(\alpha)$ of F. Moreover, if α_1 and α_2 are two colored open *n*-strings such that the product $\alpha_1 \cdot \alpha_2$ is defined, then $\Phi(\alpha_1 \cdot \alpha_2)$ is the composition $\Phi(\alpha_1) \circ \Phi(\alpha_2)$.

References.

[C84] R. Crowell, The derived group of a permutation representation, Advances in Math. 53 (1984), 88–124.

[K99] L.H. Kauffman, Virtual knot theory, European J. Comb. 20 (1999), 663–690.

[M04] V. Manturov, Long virtual knots and their invariants II: the commutation problem, preprint.

[S02] J. Sawollek, A formula for the Z-polynomial, preprint, 2002.

[SW01] D.S. Silver and S.G. Williams, A generalized Burau representation for open string links, *Pacific J. Math.* **197** (2001), 241–255.

[SW01'] D.S. Silver and S.G. Williams, Alexander groups and virtual links, J. Knot Theory and its Ramifications 10 (2001), 151–160.

[SW03] D.S. Silver and S.G. Williams, Polynomial invariants of virtual links, J. Knot Theory and its Ramifications 12 (2003), 987–1000.

[SW04] D.S. Silver and S.G. Williams, Alexander groups of long virtual knots, preprint, 2004, arXiv:math.GT/0405460

 $[{\rm T04}]$ V. Turaev, Virtual open strings and their cobordisms, preprint, 2004, arXiv: math.GT/ 0311185 v5