

# HOMOTOPY OPERATIONS AND RATIONAL HOMOTOPY TYPE

DAVID BLANC

## 1. INTRODUCTION

In [HS] and [F1] Halperin, Stasheff, and Félix showed how an inductively-defined sequence of elements in the cohomology of a graded commutative algebra over the rationals can be used to distinguish among the homotopy types of all possible realizations, thus providing a collection of algebraic invariants for distinguishing among rational homotopy types of spaces. There is also a dual version, in the setting of graded Lie algebras (see [O]).

However, these authors provided no homotopy-theoretic interpretation of these invariants, which are defined in terms of differential graded algebras (resp. Lie algebras) and their possible perturbations.

The goal of this paper is to provide such an interpretation, in terms of higher rational homotopy operations, and thus to make sense of the following

**Theorem A.** *For any simply-connected space  $\mathbf{X}$ , there is a sequence of higher homotopy operations taking value in  $\pi_*\mathbf{X}$ , which, together with the rational homotopy Lie algebra  $\pi_{*-1}\mathbf{X} \otimes_{\mathbb{Z}} \mathbb{Q}$  itself, determine the rational homotopy type of  $\mathbf{X}$ .*

(See Theorem 7.14 below). At the same time, we provide a more concrete (rational) version of the general theory of higher homotopy operations provided in [BM].

It should be noted that an integral version of the Lie algebra case is contained in [Bl5] (see also [BG, BDG]), and the mod  $p$  homology analogue of the Halperin-Stasheff-Félix theory appears in [Bl7]. Moreover, in [Bl2] we showed that the (integral) homotopy type of a space  $\mathbf{X}$  is in fact determined by its homotopy groups  $\pi_*\mathbf{X}$ , together with the action of all primary homotopy operations on it, and of certain higher

---

*Date:* October 20, 2001; revised August 31, 2002.

*1991 Mathematics Subject Classification.* Primary 55P62; Secondary 55Q35, 55P15, 18G50.

*Key words and phrases.* rational homotopy type, higher homotopy operations, homology of DGLs, non-associative algebras, obstruction theory.

homotopy operations (see [Bl3, Bl6] for subsequent modifications and improvements).

However, if we are interested only in the *rational* homotopy type of a simply-connected space  $\mathbf{X}$ , Whitehead products are the only non-trivial primary homotopy operations on the rational homotopy groups  $\pi_*\mathbf{X}_{\mathbb{Q}} = \pi_*\mathbf{X} \otimes \mathbb{Q}$ , which, after re-indexing, constitute a graded Lie algebra over  $\mathbb{Q}$ . The relevant higher order operations are also simpler than in the integral case. Thus we hope that the rational version of this theory will be both easier to understand, and more accessible to computation.

Moreover, the higher operations we define are certain subsets of  $\pi_*\mathbf{X}$ , indexed by elements in homology groups of a certain inductively defined collection of differential graded Lie algebras (DGLs) defined below, so we provide an explicit correspondence between our higher operations and the corresponding elements in cohomology groups with coefficients in  $\pi_*\mathbf{X}$  (provided by the Halperin-Stasheff-Félix theory) – a correspondence which was lacking in the integral case.

Finally, while a notion of higher homotopy operations for a differential graded Lie algebra  $L$  has been defined in the special case of higher Whitehead products (also known as “Lie-Massey products” – see [A1, A2, AA, R1, R2, T]), in general it is not clear how to represent such rational operations as *integral* higher order operations in  $\pi_*\mathbf{X}$ , if  $L$  represents the rational homotopy type of a topological space  $\mathbf{X}$ . In order to address this problem, we must consider a somewhat “flabbier” model of rational homotopy than that provided by differential graded Lie algebras, namely a certain class of differential graded non-associative algebras (see section 7 below).

Thus we also provide a (somewhat incomplete) answer to the following question: what additional structure on the ordinary homotopy groups  $\pi_*\mathbf{X}$  of a simply-connected space  $\mathbf{X}$ , beyond the Whitehead products, is needed to determine its homotopy type up to rational equivalence?

**1.1. Notation and conventions.** The ground field for all vector spaces, algebras, and tensor products will be  $\mathbb{Q}$  (the rationals), unless otherwise stated.

$\mathcal{T}_*$  denotes the category of pointed *CW* complexes with base-point preserving maps, and by a *space* we shall always mean an object in  $\mathcal{T}_*$ , which will be denoted by a boldface letter:  $\mathbf{X}, \mathbf{S}^n, \dots$ . The subcategory of 1-connected spaces is denoted by  $\mathcal{T}_1$ , and the rationalization of a space  $\mathbf{X} \in \mathcal{T}_1$  is  $\mathbf{X}_{\mathbb{Q}}$ . The category of rational 1-connected topological spaces is denoted by  $\mathcal{T}_{\mathbb{Q}}$ .

Let  $\Delta$  denote the category of ordered sequences  $\mathbf{n} = \langle 0, 1, \dots, n \rangle$  ( $n \in \mathbb{N}$ ), with order-preserving maps. For any category  $\mathcal{C}$ , we let  $s\mathcal{C}$  denote the category of simplicial objects over  $\mathcal{C}$  – i.e., functors  $\Delta^{op} \rightarrow \mathcal{C}$  (cf. [Ma, §2]); objects therein will be written  $A_\bullet, \dots$ . If we omit the degeneracies, we have a  $\Delta$ -simplicial object, which we denote by  $A_\bullet^\Delta, \dots$ .

The category of non-negatively graded objects over a category  $\mathcal{C}$  will be denoted by  $\text{gr } \mathcal{C}$ , with objects written  $T_*, \dots$ ; we will write  $|x| = p$  if  $x \in T_p$ . An upward shift by one in the indexing will be denoted by  $\Sigma : \text{gr } \mathcal{C} \rightarrow \text{gr } \mathcal{C}$ , so that  $(\Sigma X_*)_{k+1} = X_k$ , and  $(\Sigma X_*)_0 = 0$ . The category of graded vector spaces is denoted by  $\mathcal{V}$ .

The category of chain complexes (over  $\mathbb{Q}$ ) will be denoted by  $d\mathcal{V}$ , and that of double chain complexes by  $dd\mathcal{V}$ . The differential of any differential graded object is written  $\partial$  (to distinguish it from the face maps  $d_i$  of a simplicial object).

If  $\mathcal{C}$  is a closed model category (cf. [Q1, I] or [Q3, II, §1]), we denote by  $\text{ho } \mathcal{C}$  the corresponding homotopy category. If  $X \in \mathcal{C}$  is cofibrant and  $Y \in \mathcal{C}$  is fibrant, we denote by  $[X, Y]_{\mathcal{C}}$  the set of homotopy classes of maps between them.

Let  $\text{Set}$  denote the category of sets,  $\text{Vec}$  the category of vector spaces (over  $\mathbb{Q}$ ),  $\text{Lie}$  the category of Lie algebras, and  $\text{Alg}$  the category of non-associative algebras. We write  $\mathcal{S}$  rather than  $s\text{Set}$  for the category of simplicial sets, and  $\mathcal{S}_*$  for the category of *pointed* simplicial sets.

**1.2. Organization:** In section 2 we review some background material on the Quillen DGL model for rational homotopy theory, and describe a bigraded variant of it; and in section 3 we give some more background on simplicial resolutions. These are applied to the rational context in section 4, where we also define higher order homotopy operations for DGLs. These appear as the obstructions to realizing certain algebraic equivalences, and serve to determine the rational homotopy type of a simply-connected space. We give a first approximation to Theorem A in §4.15.

In section 5 we explain how to translate the usual bigraded and filtered DGL models into simplicial DGLs, which allows us to construct appropriate minimal simplicial resolutions. In section 6 we define the homology and cohomology of a DGL (after Quillen), and show that the obstructions we define above actually take value in the appropriate cohomology groups. Finally, in section 7 we describe a non-associative differential graded algebra model for rational homotopy theory, which facilitates the translation of the higher homotopy operations described

above into integral homotopy operations. We summarize our main results in Theorem 7.14.

1.3. *Acknowledgements.* I would like to thank Ron Livné for arousing my interest in rational higher homotopy operations, Martin Arkowitz and Jean-Michel Lemaire for providing me with copies of hard-to-get theses, and Steve Halperin for some useful comments. I would also like to thank the referee for his remarks and corrections.

## 2. LIE MODELS

In this section we briefly recall some well-known definitions and facts of rational homotopy theory, and describe variants thereof.

2.1. **Differential graded Lie algebras.** Let  $\mathcal{L}$  denote the category of graded Lie algebras, or *GL*'s. An object  $L_* \in \mathcal{L}$  is thus a graded vector space:  $L_* = \bigoplus_{n=0}^{\infty} L_n$  over  $\mathbb{Q}$ , equipped with a bilinear graded product  $[ \ , \ ] : L_p \otimes L_q \rightarrow L_{p+q}$  for each  $p, q, \geq 0$ , such that  $[x, y] = (-1)^{|x||y|+1}[y, x]$  and  $(-1)^{|x||z|}[[x, y], z] + (-1)^{|y||x|}[[y, z], x] + (-1)^{|z||y|}[[z, x], y] = 0$ . We denote by  $\mathcal{L}_0$  the full subcategory of all *connected* graded Lie algebras – that is, those for which  $L_0 = 0$ .

The free graded Lie algebra generated by a graded set  $X_*$  is denoted by  $\mathbb{L}\langle X_* \rangle$ . The functor  $\mathbb{L} : \text{gr Set} \rightarrow \mathcal{L}$  is left adjoint to the forgetful “underlying graded set” functor  $U : \mathcal{L} \rightarrow \text{gr Set}$ , and it factors through  $\mathcal{V}$ : that is,  $\mathbb{L}\langle X_* \rangle = L(\mathbb{V}\langle X_* \rangle)$ , where  $\mathbb{V}\langle X_* \rangle \in \mathcal{V}$  is the graded vector space with basis  $X_*$  and  $L(\mathbb{V}_*)$  is the free Lie algebra on the graded vector space  $V_*$  (defined as the appropriate quotient of the graded tensor algebra).

The category of *differential graded Lie algebras*, or *DGLs*, will be denoted by  $d\mathcal{L}$ , with  $d\mathcal{L}_0$  the subcategory of *connected* Lie algebras (i.e., those with  $L_0 = 0$ ). An object  $L = (L_*, \partial_L) \in d\mathcal{L}$  is a graded Lie algebra  $L_* \in \mathcal{L}$ , together with a differential  $\partial_L = \partial_L^n : L_n \rightarrow L_{n-1}$ , for each  $n > 0$ , such that  $\partial_L^{n-1} \circ \partial_L^n = \{0\}$  and  $\partial_L[x, y] = [\partial_L x, y] + (-1)^{|x|}[x, \partial_L y]$ .

The homology of the underlying chain complex of a DGL  $L = (L_*, \partial)$  will be denoted  $H'_*L$ , to distinguish it from the DGL homology defined in §6.5 below. Because the differential  $\partial$  is a derivation,  $H'_*L$  inherits from  $L$  the structure of a graded Lie algebra.

A morphism of DGLs which induces an isomorphism in homology will be called a *quasi-isomorphism*, or *weak equivalence*, denoted by  $f : L \xrightarrow{\sim} L'$ .

In [Q3, II,§4-5], Quillen defined closed model category structures for the categories  $d\mathcal{L}_0$  and  $s\mathcal{L}ie$ , as well as for topological spaces (and thus for  $\mathcal{T}_{\mathbb{Q}}$ ), and proved:

**2.2. Proposition.** *There are pairs of adjoint functors  $\mathcal{T}_{\mathbb{Q}} \rightleftarrows s\mathcal{L}ie$  and  $s\mathcal{L}ie \rightleftarrows d\mathcal{L}_0$ , which induce equivalences between the corresponding homotopy categories:  $\text{ho}\mathcal{T}_{\mathbb{Q}} \approx \text{ho}(s\mathcal{L}ie) \approx \text{ho}(d\mathcal{L}_0)$ .*

**2.3. Notation.** To every simply-connected space  $\mathbf{X} \in \mathcal{T}_1$  one can thus associate a DGL  $(L_*, \partial_L) \in d\mathcal{L}_0$ , unique up to quasi-isomorphism, which determines its rational homotopy type. We denote any such DGL by  $L_X$ . In particular,  $H'_*(L_X) \cong \pi_{*-1}\mathbf{X} \otimes_{\mathbb{Z}} \mathbb{Q}$ , the rational homotopy algebra of  $\mathbf{X}$ , which we denote by  $\Pi_*^X \in \mathcal{L}$ .

**2.4. Definition.** The graded Lie algebra  $H'_*(L_X)$  does not suffice to determine the rational homotopy type of  $\mathbf{X} \in \mathcal{T}_1$ : in fact, there may be infinitely many DGLs  $\{L^{(n)}\}_{n=1}^{\infty}$  with  $H'_*(L^{(n)}) \cong H'_*(L_X)$ , no two of which are quasi-isomorphic as DGLs; see e.g. [LS]. We shall denote by  $d\mathcal{L}_0(\mathbf{X})$  the full subcategory of  $d\mathcal{L}_0$  whose objects  $A$  satisfy  $H'_*A \cong H'_*(L_X)$ , with the isomorphism in  $\mathcal{L}$  (see [SS], [LS], or [F1] for treatments of the cohomology analogue of  $d\mathcal{L}_0(\mathbf{X})$  in terms of algebraic varieties).

The objects of  $\text{ho } d\mathcal{L}_0(\mathbf{X})$  are thus all rational homotopy types which are indistinguishable from  $\mathbf{X}_{\mathbb{Q}}$  on the primary homotopy operation level. Among these there is a distinguished simplest one: recall that a space  $\mathbf{X}_{\mathbb{Q}} \in \mathcal{T}_{\mathbb{Q}}$  (or its corresponding DGL model  $L_X \in d\mathcal{L}$ ) is called *coformal* (cf. [MN]) if  $L_X$  is weakly equivalent to the trivial DGL  $(L_*, 0)$  (where of course  $L_* = H'_*(L_X)$ ).

**2.5. Minimal models.** Baues and Lemaire (in [BL, Cor. 2.4]; see also [N, Props. 5.6, 8.1 & 8.8]) showed that each connected DGL  $(L_*, \partial)$  has a *minimal model*  $(\hat{L}_*, \hat{\partial})$ , such that  $\hat{L}_*$  is a free graded Lie algebra,  $\hat{\partial} : \hat{L} \rightarrow \hat{L}$  factors through  $[\hat{L}, \hat{L}]$ , and there is a quasi-isomorphism of DGLs  $\varphi : (\hat{L}_*, \hat{\partial}) \rightarrow (L_*, \partial)$  (unique up to chain homotopy). In particular, we can choose such a minimal model  $\hat{L}_X$  for any space  $\mathbf{X} \in \mathcal{T}_1$  (cf. §2.3).

As Neisendorfer observes in [N, §5], in general minimal models do not exist for non-connected DGLs (but see [Me] or [GHT] for ways around this).

**2.6. Bigraded Lie algebras.** A differential bigraded Lie algebra, or *DBGL*, is a bigraded vector space  $L_{*,*} = \bigoplus_{p=0}^{\infty} \bigoplus_{s=0}^{\infty} L_{p,s}$ , equipped

with a differential  $\partial_L = \partial_L^{p,s} : L_{p,s} \rightarrow L_{p-1,s}$  and a bilinear graded product  $[\cdot, \cdot] : L_{p,s} \otimes L_{q,t} \rightarrow L_{p+q,s+t}$  for each  $p, q, s, t \geq 0$  satisfying:

$$\begin{aligned}
(2.7) \quad & [x, y] = (-1)^{(p+s)(q+t)+1} [y, x] \\
& (-1)^{(p+s)(r+u)} [[x, y], z] + (-1)^{(p+s)(q+t)} [[y, z], x] \\
& + (-1)^{(q+t)(r+u)} [[z, x], y] = 0 \\
& \partial_L \circ \partial_L = 0 \\
& \partial_L [x, y] = [\partial_L x, y] + (-1)^{p+s} [x, \partial_L y]
\end{aligned}$$

for  $x \in L_{p,s}$ ,  $y \in L_{q,t}$ , and  $z \in L_{r,u}$ . The category of such DBGLs will be denoted by  $\text{db } \mathcal{L}$ , with  $\text{db } \mathcal{L}_0$  the subcategory with  $L_{p,0} = 0$  for all  $p$ .

**2.8. Definition.** For each DBGL  $(L_{*,*}, \partial_L)$  there is an *associated* DGL  $(L_*, \partial_L)$ , defined  $L_n = \bigoplus_{p+q=n} L_{p,q}$  (same  $\partial_L$ ); some authors re-index  $L_{*,*}$  so that  $\hat{L}_{p,s} = L_{p,p+s}$ , and then  $L_*$  is obtained from  $\hat{L}_{*,*}$  by disregarding the first (homological) grading.

As for ordinary graded Lie algebras, one can define closed model category structures on  $s\mathcal{L}_0$  and  $\text{db } \mathcal{L}_0$  (see [BS, §2], and [Bl4, §4]), and we have the following analogue of [Q3, I, Props. 2.3 & 4.6, Thm. 4.4]:

**2.9. Proposition.** *There are adjoint functors  $s\mathcal{L}_0 \xrightleftharpoons[N^*]{N} \text{db } \mathcal{L}_0$ , which induce equivalences of the corresponding homotopy categories  $\text{ho}(s\mathcal{L}_0) \approx \text{ho}(\text{db } \mathcal{L}_0)$ .  $N^*$  takes free DBGLs to free simplicial graded Lie algebras.*

*Proof.* (We give the proof mainly to fix notation which will be needed later.) Given a simplicial graded Lie algebra  $L_{\bullet,*} \in s\mathcal{L}_0$ , we define the simplicial Lie bracket  $[[\cdot, \cdot]] : L_{p,s} \otimes L_{q,t} \rightarrow L_{p+q,s+t}$  on  $L_{\bullet,*}$  by combining the Lie brackets with the simplicial structure on  $L_{\bullet,*}$  via the Eilenberg-Zilber map:

$$(2.10) \quad [[x, y]] = \sum_{(\sigma, \tau) \in S_{p,q}} (-1)^{\varepsilon(\sigma)+pt} [s_{\tau_q} \dots s_{\tau_1} x, s_{\sigma_p} \dots s_{\sigma_1} y]$$

where  $S_{p,q}$  denotes the set of all  $(p, q)$ -shuffles – that is, partitions of  $\{0, 1, \dots, p+q-1\}$  into disjoint sets  $\sigma = \{\sigma_1, \sigma_2, \dots, \sigma_p\}$  and  $\tau = \{\tau_1, \tau_2, \dots, \tau_q\}$  with  $\sigma_1 < \sigma_2 < \dots < \sigma_p$ ,  $\tau_1 < \tau_2 < \dots < \tau_q$  – and  $\varepsilon(\sigma) = p + \sum_{i=1}^p (\sigma_i - i)$ , so  $(-1)^{\varepsilon(\sigma)}$  is the sign of the permutation corresponding to  $(\sigma, \tau)$ . (See [Mc1, VIII, §8]).

Now let  $(C_{*,*}, \partial)$  be the Moore chain complex (cf. [Ma, §22]) of  $L_{\bullet,*}$ , defined by:

$$(2.11) \quad C_{p,s} = \bigcap_{i=1}^p [\text{Ker}(d_i^p)]_s \quad \text{with } \partial_p = (-1)^s d_0^p|_{C_{p,s}}.$$

**2.12. Lemma.** *If  $A_{\bullet} \in s\mathcal{L}$  is a simplicial graded Lie algebra,  $x \in A_p$  with  $d_i x = 0$  for  $1 \leq i \leq p-1$ , and  $y \in A_q$  with  $d_j y = 0$  for  $1 \leq j \leq q-1$ , then  $d_k(\llbracket x, y \rrbracket) = 0$  for  $1 \leq k \leq p+q-1$ .*

*Proof.* By definition (2.10) we have

$$(2.13) \quad \llbracket x, y \rrbracket = \sum_{(\sigma, \tau) \in S_{p,q}} (-1)^{\varepsilon(\sigma) + p|y|} [s_{\tau_q} \dots s_{\tau_1} x, s_{\sigma_p} \dots s_{\sigma_1} y] \in A_{p+q}.$$

Now for each summand  $w_{\sigma, \tau} := [s_{\tau} x, s_{\sigma} y]$  in (2.13), with  $(\sigma, \tau)$  a  $(p, q)$ -shuffle, there are two cases to consider:

The first is that there exist  $\ell, m$  such that  $\tau_{\ell} = k$ ,  $\sigma_m = k-1$  – in which case there is an associated  $(p, q)$ -shuffle  $(\sigma', \tau')$ , differing from  $(\sigma, \tau)$  only in that  $\tau_{\ell}$  and  $\sigma_m$  are switched, so that  $d_k(w_{\sigma, \tau}) = d_k(w_{\sigma', \tau'})$  but  $(-1)^{\varepsilon(\sigma)} = -(-1)^{\varepsilon(\sigma')}$ , and these pairs thus cancel in the sum (2.13).

In the second case,  $k, k-1 \in \{\sigma_1, \dots, \sigma_p\}$ , say, and then there is some  $0 \leq \ell \leq q$  with  $\tau_{\ell} < k-1$  and  $\tau_{\ell+1} > k$ . Since necessarily  $k+1-p \leq \ell \leq k-1$ , we find that  $d_k s_{\tau} x = s_{\tau_{\ell-1}} \dots s_{\tau_{\ell+1}-1} s_{\tau_{\ell}} \dots s_{\tau_1} d_{k-\ell} x = 0$ .  $\square$

**2.14. Corollary.**  $\llbracket \ , \ ]$  restricts to a bracket  $C_{p,s} \otimes C_{q,t} \rightarrow C_{p+q,s+t}$

Moreover, if we forget the Lie structure, the Moore chain complex functor  $N$  induces an equivalence between the categories of simplicial graded vector spaces and bigraded chain complexes (cf. [Do, Thm 1.9]), with the inverse functor  $\Gamma$  defined for such a chain complex  $(A_{*,*}, \partial)$  by

$$(\Gamma A_{*,*})_{n,s} := \bigoplus_{0 \leq \lambda \leq n} \bigoplus_{I \in \mathcal{J}_{n,\lambda}} A_{n-\lambda, s}$$

(where for each  $n \geq 0$  and  $0 \leq \lambda \leq n$ , we let  $\mathcal{J}_{\lambda, n}$  denote the set of all sequences of  $\lambda$  non-negative integers  $i_1 < \dots < i_{\lambda} (< n)$ ), with the obvious face maps (induced by  $\partial$ ) and degeneracies (see [Ma, p. 95]).

The left adjoint  $N^* : \text{db } \mathcal{L}_0 \rightarrow s\mathcal{L}_0$  to  $N$  is defined

$$N^*((L_{*,*}, \partial)) = L(\Gamma(L_{*,*})) / I(L_{*,*}),$$

where  $L$  is the free graded Lie algebra functor, and  $I(L_{*,*})$  is the ideal generated by  $\llbracket \Gamma(x), \Gamma(y) \rrbracket - \Gamma(\llbracket x, y \rrbracket)$ . The identities (2.7) follow from

the corresponding ones in the singly-graded case and the simplicial identities.  $\square$

### 3. SIMPLICIAL RESOLUTIONS

The proper algebraic setting for defining our higher homotopy operations is a suitable notion of a simplicial resolution of  $\pi_*\mathbf{X}_{\mathbb{Q}}$ :

**3.1. Definition.** Recall that a category of *universal graded algebras* (or variety of graded algebras, in the terminology of [Mc2, V, §6]) is a category  $\mathcal{C}$  in which the objects are graded sets  $X_*$ , together with an action of a fixed set of  $n$ -ary graded operators  $W = \{\omega : X_{k_1} \times X_{k_2} \times \cdots \times X_{k_n} \rightarrow X_m\}$ , satisfying a set of identities  $E$ , and the morphisms are functions on the sets which commute with the operators. Such categories always come equipped with a “free graded algebra” functor  $F : \text{gr Set} \rightarrow \mathcal{C}$ , left adjoint to the “underlying graded set” functor  $U : \mathcal{C} \rightarrow \text{gr Set}$ . In all the examples we shall be concerned with, the objects  $X_*$  will be “underlying-abelian” (see [BS, §2.1.1]), and in fact will have the underlying structure of a graded vector space over  $\mathbb{Q}$ .

Examples include  $\mathcal{L}$ , and the categories of associative (resp. non-associative) graded algebras. Note that any ordinary ungraded category of universal algebras may be thought of as a CUGA with all objects concentrated in degree 0.

**3.2. Definition.** A *free simplicial resolution* of an object  $B$  in a CUGA  $\mathcal{C}$  is a weak equivalence from a cofibrant object  $A_{\bullet} \in s\mathcal{C}$  to the constant simplicial object associated to  $B$  (with respect to the closed model category structure on the category  $s\mathcal{C}$  defined in [Q1, II, §4]). Such resolutions always exist, by [Q1, II, §4]; see section 5 below for a specific construction.

**3.3. Bisimplicial objects.** We shall be interested in a particular type of simplicial resolution, which may be defined for an arbitrary CUGA  $\mathcal{C}$  ((cf. [DKS] and [BS]), though we shall only need it for the case where  $\mathcal{C}$  is a category of ungraded universal algebras, such as *Lie* or *Alg*):

Consider the category  $ss\mathcal{C}$  of bisimplicial objects over  $\mathcal{C}$ . We think of an object  $A_{\bullet\bullet} \in ss\mathcal{C}$  as having *internal* and *external* simplicial structures, with corresponding homotopy group objects  $\pi_t^i A_{\bullet\bullet}$  and  $\pi_s^e A_{\bullet\bullet}$  (each taking value in  $s\mathcal{C}$  – see [BS, App.]). Let  $sF : \text{gr } \mathcal{S} \rightarrow s\mathcal{C}$  denote the free graded algebra functor, extended dimensionwise, and let  $S^n(k)_{\bullet}$  be the graded simplicial set having the simplicial  $n$ -sphere  $S_{\bullet}^n := \Delta[n]/\Delta[n]^{n-1}$  in degree  $k$ . We think of the simplicial graded algebras  $F(S^n(k)_{\bullet})$  as the  $\mathcal{C}$ -spheres, or *models*, for  $s\mathcal{C}$  (cf. [BS, §3.1]). (In the ungraded case one can of course omit the extra degree  $k$ , and

write simply  $F(\mathbf{S}^n)$ ). Similarly, if  $D^n(k)_\bullet$  is the graded simplicial set having  $\Delta[n]$  in degree  $k$ , we can think of  $F(D^n(k)_\bullet)$  as the  $\mathcal{C}$ -disks for  $s\mathcal{C}$ . The full subcategory of  $s\mathcal{C}$  whose objects are weakly equivalent to coproducts of such models will be denoted by  $\mathcal{M}_{\mathcal{C}}$ , or simply  $\mathcal{M}$ .

One can use these models to define the so-called “ $E^2$ -model category structure” for  $ss\mathcal{C}$ , as in [DKS, §5], in which a map  $f : X_{\bullet\bullet} \rightarrow Y_{\bullet\bullet}$  is a weak equivalence if

$$(3.4) \quad f_* : \pi_s \pi_t^i X_{\bullet\bullet} \rightarrow \pi_s \pi_t^i Y_{\bullet\bullet} \text{ is an isomorphism for each } s, t \geq 0$$

We shall not need an explicit description of the fibrations and cofibrations in  $ss\mathcal{C}$ , but only a particular type of cofibrant object, as follows:

**3.5. Definition.** A bisimplicial object  $A_{\bullet\bullet} \in ss\mathcal{C}$  is called  $\mathcal{M}$ -free if for each  $m \geq 0$  there are graded simplicial sets  $X[m]_\bullet \simeq \bigvee_i \mathbf{S}^{n_i}(k_i)_\bullet$  such that  $A_{\bullet,m} \cong F(X[m]_\bullet)$  (so that  $A_{\bullet,m} \in \mathcal{M}$ ), and the external degeneracies of  $A_{\bullet\bullet}$  are induced under  $F$  by maps  $X[m]_\bullet \rightarrow X[m+1]_\bullet$  which are, up to homotopy, the inclusion of sub-coproduct summands. Any  $X_\bullet \in s\mathcal{C}$  may be resolved by an  $\mathcal{M}$ -free bisimplicial algebra  $A_{\bullet\bullet}$  (see [BS, §4.1]); this is called an  $\mathcal{M}$ -free resolution of  $X_\bullet$ .

**3.6. Definition.** The *diagonal* of a bisimplicial object  $A_{\bullet\bullet} \in ss\mathcal{C}$  is a simplicial object  $\text{diag}(A_{\bullet\bullet}) \in s\mathcal{C}$  with  $\text{diag}(A_{\bullet\bullet})_n := A_{n,n}$ , face maps  $d_k = d_k^i \circ d_k^e$ , and degeneracies  $s_j = s_j^i \circ s_j^e$ .

**3.7. Remark.** There is a first quadrant spectral sequence with

$$E_{s,t}^2 = \pi_s^e(\pi_t^i A_{\bullet\bullet}) \Rightarrow \pi_{s+t} \text{diag}(A_{\bullet\bullet})$$

(see [Q2], and compare [BF, Thm B.5]).

Thus in particular if  $A_{\bullet\bullet} \rightarrow X_\bullet$  is a resolution (in the  $E^2$ -model category sense), we see that  $\varepsilon : A_{0,\bullet} \rightarrow X_\bullet$  induces a weak equivalence  $\text{diag}(A_{\bullet\bullet}) \simeq X_\bullet$ .

Moreover, the same is true if we disregard the degeneracies and consider only the  $\Delta$ -bisimplicial resolution  $A_{\bullet\bullet}^\Delta \rightarrow X_\bullet$ .

#### 4. RESOLUTIONS FOR RATIONAL SPACES

Given a simply-connected space  $\mathbf{X} \in \mathcal{T}_1$ , the first approximation to an algebraic description of its rational homotopy type is given by its rational homotopy Lie algebra  $\Pi_*^X := \pi_{*-1} \mathbf{X}_{\mathbb{Q}} \in \mathcal{L}$ .

If  $\mathbf{X}_{\mathbb{Q}}$  were coformal (§2.4), then in particular all higher homotopy operations vanish in  $\pi_* \mathbf{X}_{\mathbb{Q}}$ , and no information beyond  $\Pi_*^X$  itself is needed to determine the rational homotopy type of  $\mathbf{X}$ . The higher

homotopy operations we shall describe may thus be thought of as “obstructions to coformality”, much in the spirit (though not the specific approach) of [HS].

**4.1. Topological resolutions.** To proceed further, we need some kind of a “topological” simplicial object  $C_\bullet$  which realizes a suitable “algebraic” simplicial resolution  $V_{\bullet,*} \rightarrow \Pi_*^X$  in  $s\mathcal{L}$ , in the sense that  $V_{\bullet,*} = \pi_{*-1}C_\bullet$ . The higher homotopy operations we want then arise as the obstructions to realizing the “algebraic” augmentation map  $\pi_{*-1}C_\bullet \rightarrow \Pi_*^X$  topologically.

This can be done using actual topological spaces, as in the integral case (see [Bl2, §7], as simplified in [Bl3, §4.9]), but for rational spaces it is more convenient to use an algebraic model, in a category such as  $d\mathcal{L}$ . To allow us freedom in choosing this model, we give a general definition:

**4.2. Assumptions.** Let  $\text{gr}\mathcal{C}$  be a CUGA (which we may assume to have the underlying structure of a graded vector space), and  $\mathcal{C}$  the category of (ungraded) universal algebras corresponding to objects of  $\text{gr}\mathcal{C}$  concentrated in degree 0. The cases we shall be interested in are  $\mathcal{C} = \mathcal{L}ie$  (with  $\text{gr}\mathcal{C} = \mathcal{L}$ ) and  $\mathcal{C} = \mathcal{A}lg$  (with  $\text{gr}\mathcal{C} = \mathcal{A}$ ).

As shown in [BS, App.], for each simplicial algebra  $A_\bullet \in s\mathcal{C}$ , the graded homotopy object  $\pi_*A_\bullet$  actually takes value in  $\text{gr}\mathcal{C}$ .

For a given  $A_\bullet \in s\mathcal{C}$ , let  $C_{\bullet\bullet} \rightarrow A_\bullet$  be an  $\mathcal{M}_{\mathcal{C}}$ -free resolution (Definition 3.5). In particular, this implies that upon applying the functor  $\pi_*$  we obtain a free simplicial resolution  $\pi_*^i C_{\bullet\bullet}$  (in the “external” direction!) of the graded algebra  $\pi_*A_\bullet$ . In fact, we only need a  $\Delta$ -bisimplicial resolution (§3.7), but we shall nevertheless usually abuse notation by writing  $C_{\bullet\bullet}$  for  $C_{\bullet\bullet}^\Delta$ .

Next, assume we are given another object  $B_\bullet \in s\mathcal{C}$ , together with an isomorphism  $\varphi : \pi_*A_\bullet \cong \pi_*B_\bullet$  (in  $\text{gr}\mathcal{C}$ ). Define a sequence of morphisms  $\psi_n : \pi_*C_{n,\bullet} \rightarrow \pi_*B_\bullet$  by  $\psi_0 := \varphi \circ \varepsilon_\#$  and  $\psi_{n+1} := \psi_n \circ d_0$  (which implies that  $\psi_{n+1} = \psi_n \circ d_i$  for all  $0 \leq i \leq n$ , by the simplicial identities).

We choose once and for all a fixed map  $f_0 : C_{0,\bullet} \rightarrow B_\bullet$  realizing  $\psi_0$  (this is possible because  $C_{\bullet\bullet} \rightarrow A_\bullet$  is  $\mathcal{M}$ -free) and define  $f_n : C_{n,\bullet} \rightarrow B_\bullet$  inductively by setting  $f_{n+1} := f_n \circ d_n$ , so that  $\pi_*(f_n) = \psi_n$  for all  $n \geq 0$ . It is usually most convenient to set  $f|_{\mathcal{D}k} = 0$  for all  $\mathcal{C}$ -disks  $\mathcal{D}k \hookrightarrow C_{0,\bullet}$  (cf. §3.3).

Note that, because  $C_{\bullet\bullet}$  is  $\mathcal{M}$ -free, the maps  $\{\psi_n\}_{n=0}^\infty$  define an augmented  $\Delta$ -simplicial object  ${}^h C_{\bullet\bullet}^\Delta \rightarrow B_\bullet$  in the homotopy category  $\text{ho}(s\mathcal{C})$  – or equivalently, an augmented  $\Delta$ -simplicial object up-to-homotopy.

**4.3. Definition.** Let  $D[n] \in \mathcal{S}_*$  denote the standard simplicial  $n$ -simplex, together with an indexing of its non-degenerate  $k$ -dimensional faces  $D[k]^{(\gamma)}$  by the composite face maps

$$\gamma = d_{i_{n-k+1}} \circ \dots \circ d_{i_n} : \mathbf{n} \rightarrow \mathbf{k}$$

in  $\Delta^{op}$  (cf. [Bl3, §4]). Its  $(n-1)$ -skeleton, which is a simplicial  $(n-1)$ -sphere, is denoted by  $\partial D[n]$ . We shall take  $*$  :=  $D[0]^{(d_0 d_1 d_2 \dots d_{n-1})}$  as the base point of  $D[n] \in \mathcal{S}_*$ , and we choose once and for all a fixed isomorphism  $\varphi^{(\gamma)} : D[k]^{(\gamma)} \rightarrow D[k]$  for each face  $D[k]^{(\gamma)}$  of  $D[n]$  (see, e.g., [Bl3, (4.5)]).

**4.4. Definition.** Given  $Y_\bullet \in s\mathcal{C}$  and a simplicial set  $K_\bullet \in \mathcal{S}$ , we define their *half-smash* (in  $s\mathcal{C}$ ) by:

$$Y_\bullet \rtimes K_\bullet := Y_\bullet \otimes K_\bullet / (\{0\} \otimes K_\bullet)$$

(where  $(Y_\bullet \otimes K_\bullet)_n := \coprod_{x \in K_n} (Y_n)_{(x)}$  – cf. [Q1, II, §1, Prop. 2]).

Similarly, the *smash product* (in  $s\mathcal{C}$ ) of  $Y_\bullet$  with a *pointed* simplicial set  $K_\bullet \in \mathcal{S}_*$  is defined  $Y_\bullet \wedge K_\bullet := Y_\bullet \rtimes K_\bullet / (Y_\bullet \rtimes \{*\})$ , and if  $K_\bullet = \mathbf{S}^r$  (the simplicial sphere), we write  $\Sigma^r Y_\bullet$  for  $Y_\bullet \wedge \mathbf{S}^r$ .

**4.5. Remark.** If  $Y_\bullet = F(\mathbf{S}^n)$  is a  $\mathcal{C}$ -sphere (see §3.3), then  $\Sigma^r Y_\bullet \cong F(\mathbf{S}^{n+r})$  is also a  $\mathcal{C}$ -sphere. In fact, many of the usual properties of spheres in  $\text{ho } \mathcal{T}$  also hold for  $\mathcal{C}$ -spheres – e.g.,  $\pi_r X_\bullet \cong [F(\mathbf{S}^n), X_\bullet]_{s\mathcal{C}}$  for any  $X_\bullet \in s\mathcal{C}$  (cf. [Q1, I, §4]), and  $Y_\bullet \simeq \coprod_i F(\mathbf{S}^{n_i}) \Rightarrow \Sigma^r Y_\bullet \simeq \coprod_i F(\mathbf{S}^{n_i+r})$  (cf. [Q1, I, §3]).

**4.6. Definition.** Under the assumptions of §4.2, for each  $n \in \mathbb{N}$ , we define a  $\partial D[n]$ -compatible sequence to be a sequence of maps  $\{h_k : C_{k,\bullet} \rtimes D[k] \rightarrow B_\bullet\}_{k=0}^{n-1}$ , such that  $h_0 = f_0$  (under the natural identification  $C_{0,\bullet} \rtimes D[0] = C_{0,\bullet}$ ), and for any iterated face maps  $\delta = d_{i_{j+1}} \circ \dots \circ d_{i_n}$  and  $\gamma = d_{i_j} \circ \delta$  ( $0 \leq j < n$ ) we have

$$(4.7) \quad h_j \circ (d_{i_j} \rtimes id) = h_{j+1} \circ (id \rtimes \iota_\delta^\gamma) \quad \text{on } C_{j+1,\bullet} \rtimes D[j],$$

where  $\iota_\delta^\gamma : D[j] \rightarrow D[j+1]$  is the composite  $\iota_\delta^\gamma := \varphi^\delta \circ \iota \circ (\varphi^\gamma)^{-1}$ . Here  $\varphi^\gamma$  and  $\varphi^\delta$  are the isomorphisms of Definition 4.3, and  $\iota : D[j]^{(\gamma)} \rightarrow D[j+1]^{(\delta)}$  is the inclusion (compare [Bl3, Def. 4.10]).

A sequence of maps  $\{h_k : C_{k,\bullet} \rtimes D[k] \rightarrow B_\bullet\}_{k=0}^\infty$  satisfying condition (4.7) for all  $\gamma, \delta$ , and  $n$  is called a  $\partial D[\infty]$ -compatible sequence.

**4.8. Definition.** Given such a  $\partial D[n]$ -compatible sequence  $\{h_k : C_{k,\bullet} \rtimes D[k] \rightarrow B_\bullet\}_{k=0}^{n-1}$  the *induced map*  $\bar{h} : C_{n,\bullet} \rtimes \partial D[n] \rightarrow B_\bullet$  is defined on the “faces”  $C_{n,\bullet} \rtimes D[n-1]^{(d_i)}$  of  $C_{n,\bullet} \rtimes D[n]$  by:  $\bar{h}|_{C_{n,\bullet} \rtimes D[n-1]^{(d_i)}} = h_{n-1} \circ (d_i \rtimes id)$ . The compatibility condition (4.7) above guarantees that  $\bar{h}$  is well-defined.

**4.9. Definition.** For each  $n \geq 2$ , the  $n$ -th order homotopy operation (associated to the choice of  $C_{\bullet\bullet} \rightarrow A_{\bullet}$  in §4.2) is a subset  $\langle\langle n \rangle\rangle$  of the track group  $[\Sigma^{n-1}C_{n,\bullet}, B_{\bullet}]_{s\mathcal{C}}$  defined as follows:

Let  $T_n \subseteq [C_{n,\bullet} \rtimes \partial D[n], B_{\bullet}]_{s\mathcal{C}}$  be the set of homotopy classes of maps  $\bar{h} : C_{n,\bullet} \rtimes \partial D[n] \rightarrow B_{\bullet}$  induced as above by some  $\partial D[n]$ -compatible collection  $\{h_k\}_{k=0}^{n-1}$ . Since each  $C_{n,\bullet}$  is a suspension, up to homotopy, by Remark 4.5, we have a splitting

$$(4.10) \quad C_{n,\bullet} \rtimes \partial D[n] \simeq (\mathbf{S}^{n-1} \wedge C_{n,\bullet}) \amalg C_{n,\bullet}$$

(as for topological spaces). We define  $\langle\langle n \rangle\rangle \subseteq [\Sigma^{n-1}C_{n,\bullet}, B_{\bullet}]_{s\mathcal{C}}$  to be the image under the resulting projection of the subset  $T_n \subseteq [C_{n,\bullet} \rtimes \partial D[n], B_{\bullet}]_{s\mathcal{C}}$ .

Note that the projection of a class  $[\bar{h}] \in T_n$  on the other summand  $[C_{n,\bullet}, B_{\bullet}]_{s\mathcal{C}}$  coming from the splitting (4.10) is just the homotopy class of the map  $f_n$  of §4.2. On the other hand, since  $C_{\bullet\bullet}$  was assumed to be  $\mathcal{M}$ -free, each  $C_{n,\bullet} \simeq \coprod_{k=1}^{\infty} \coprod_{x \in K_{n,k}} F(\mathbf{S}_{(x)}^k)$  is weakly equivalent to a wedge of spheres over some indexing set  $K_{*,*}$ , so  $\Sigma^{n-1}C_{n,\bullet} \simeq \coprod_{k=1}^{\infty} \coprod_{x \in K_{n,k}} F(\mathbf{S}_{(x)}^{k+n-1})$ . Thus

$$(4.11) \quad [\Sigma^{n-1}C_{n,\bullet}, B_{\bullet}]_{s\mathcal{C}} \cong \prod_{k=1}^{\infty} \prod_{x \in K_{n,k}} [F(\mathbf{S}_{(x)}^{k+n-1}), B_{\bullet}]_{s\mathcal{C}},$$

and we shall denote the components of  $\langle\langle n \rangle\rangle$  under this product decomposition by  $\langle\langle n, x \rangle\rangle \subseteq [F(\mathbf{S}_{(x)}^{k+n-1}), B_{\bullet}]_{s\mathcal{C}} = \pi_{k+n-1}B_{\bullet}$ .

**4.12. Definition.** An operation  $\langle\langle k \rangle\rangle$  *vanishes* if it contains the null class. We say that all the lower order operations  $\langle\langle k \rangle\rangle$  ( $2 \leq k < n$ ) *vanish coherently* (cf. [Bl2, Def. 5.7]) if the  $\partial D[m]$ -compatible collections  $\{h_k^{\gamma}\}_{k=0}^{m-1}$  for the various faces  $\gamma$  of  $\partial D[n]$  can be chosen to agree on their intersections, so that they in fact fit together to form a  $\partial D[n+1]$ -compatible collection  $\{h_k\}_{k=0}^n$ .

**4.13. Proposition.** *A necessary condition for the subset  $\langle\langle n \rangle\rangle$  to be non-empty that the lower order operations  $\langle\langle k \rangle\rangle$  vanish for  $2 \leq k < n$ ; a sufficient condition is that they vanish coherently.*

*Proof.* See [BM, Theorem 3.29]. □

**4.14. Remark.** The coherent vanishing of all the operations  $\{\langle\langle n \rangle\rangle\}_{n=2}^{\infty}$  is equivalent, by [BV, Cor. 4.21 & Thm. 4.49] and [Bl2, §4.11], to the *rectifiability* of the augmented  $\Delta$ -simplicial object up-to-homotopy  ${}^h C_{\bullet\bullet}^{\Delta} \rightarrow B_{\bullet}$ : that is, its replacement by augmented  $\Delta$ -simplicial object  $\hat{C}_{\bullet\bullet}^{\Delta} \rightarrow B_{\bullet}$  over  $s\mathcal{C}$  (with the simplicial identities now holding precisely,

in  $s\mathcal{C}$ , rather than just in  $\text{ho}(s\mathcal{C})$ ), such that  $C_{n,\bullet} \simeq \hat{C}_{n,\bullet}$  for each  $n$ .

This in turn implies (by §3.7) that  $\text{diag}(\hat{C}_{\bullet\bullet}^\Delta) \simeq B_\bullet$ ; but since  $\text{diag}(\hat{C}_{\bullet\bullet}^\Delta) \simeq \text{diag}(C_{\bullet\bullet}^\Delta)$ , and  $\text{diag}(C_{\bullet\bullet}^\Delta) \simeq A_\bullet$  by assumption, we conclude that  $A_\bullet \simeq B_\bullet$  if and only if the higher homotopy operations  $\{\langle\langle n \rangle\rangle\}_{n=2}^\infty$  vanish coherently.

**4.15. Summary.** This yields a first approximation to Theorem A, which may be described as follows:

We work in  $\mathcal{C} = \mathcal{L}ie$  (and  $\text{gr}\mathcal{C} = \mathcal{L}$ ). Given a space  $\mathbf{X} \in \mathcal{T}_1$  we consider the simplicial Lie algebra  $B_\bullet$  corresponding to a DGL model  $L_X \in d\mathcal{L}$  for  $\mathbf{X}_\mathbb{Q}$  (under the functors of Proposition 2.2), and let  $\Pi_*^X := \pi_{*-1}\mathbf{X}_\mathbb{Q} \in \mathcal{L}$  be its rational homotopy Lie algebra, with  $A_\bullet \in s\mathcal{L}ie$  the simplicial Lie algebra corresponding to the trivial DGL  $L^{(0)} := (\Pi_*^X, 0)$ . Choose some  $\mathcal{M}_{\mathcal{L}ie}$ -free resolution  $C_{\bullet\bullet} \in ss\mathcal{L}ie$  of  $A_\bullet$ .

$\mathbf{X}$  is coformal if and only if  $A_\bullet \simeq B_\bullet$ , and this happens if and only if all the higher homotopy operations  $\{\langle\langle n \rangle\rangle\}_{n=2}^\infty$  associated to  $C_{\bullet\bullet}$  vanish coherently, by Remark 4.14. If not, let  $n_0$  denote the least  $n \geq 2$  such that  $0 \notin \langle\langle n \rangle\rangle$ .

Note that we can apply the above procedure to any DGL in  $d\mathcal{L}(\mathbf{X})$  (Def. 2.4), not only to  $L_X$ ; and the existence and vanishing or non-vanishing of the higher homotopy operation  $\langle\langle n_0 \rangle\rangle \subset \pi_*\mathbf{X}_\mathbb{Q}$  is a homotopy invariant. Denote by  $\mathcal{H}^{(1)}$  the set of all homotopy types in  $\text{ho } d\mathcal{L}_0(\mathbf{X})$  for which  $\langle\langle n_0 \rangle\rangle$  is defined and has the same value as for  $B_\bullet$  itself (i.e., those DGLs which are indistinguishable from  $L_X$  as far as the primary homotopy operations, and all the higher homotopy operations  $\{\langle\langle n \rangle\rangle\}_{n=2}^{n_0}$  associated to  $C_{\bullet\bullet}$ , can see). For each  $\alpha \in \mathcal{H}^{(1)}$ , choose a representative DGL  $L^{(1,\alpha)}$ .

Next, choose a new  $\mathcal{M}$ -free resolution for the simplicial Lie algebra corresponding to  $L^{(1,\alpha)}$ , and repeat the above procedure, yielding a set of higher homotopy operations  $\langle\langle n_{1,\alpha} \rangle\rangle \subset \pi_*\mathbf{X}_\mathbb{Q}$  which serve as obstructions to the existence of a homotopy equivalence  $L^{(1,\alpha)} \xrightarrow{\simeq} L_X$ . For each such higher operation  $\langle\langle n_{1,\alpha} \rangle\rangle$ , we denote by  $\mathcal{H}^{(2,\alpha)}$  the set of all homotopy types in  $\mathcal{H}^{(1)} \subseteq \text{ho } d\mathcal{L}_0(\mathbf{X})$  for which  $\langle\langle n_{1,\alpha} \rangle\rangle$  has the same value as for  $L_X$ . Now choose representatives  $L^{(2,\alpha,\alpha')}$  for each  $\alpha' \in \mathcal{H}^{(2,\alpha)}$ , and proceed as above.

In this way we obtain a tree  $T_X$  of rational homotopy types in  $\text{ho } d\mathcal{L}_0(\mathbf{X})$ , which also indexes a collection of higher homotopy operations of the form  $\langle\langle n_{k,\alpha_1,\alpha_2,\dots,\alpha_k} \rangle\rangle \subseteq \pi_*\mathbf{X}_\mathbb{Q}$ , and  $\lim_{k \rightarrow \infty} n_k = \infty$  along any branch of the tree  $T_X$ , so that in fact this collection of operations determines the rational homotopy type of  $\mathbf{X}$ .

In [B15], we show how this tree of homotopy types in  $\text{ho } d\mathcal{L}_0(\mathbf{X})$ , and thus the corresponding collection of higher homotopy operations, may be described more effectively in terms of a ‘‘Postnikov tower’’ for an  $\mathcal{M}_{\mathcal{L}ie}$ -free resolution for  $\mathbf{X}$ .

## 5. MINIMAL RESOLUTIONS

We now explain how the bisimplicial theory described in section 4 translates into a differential graded theory, when  $\mathcal{C} = \mathcal{L}$ . In particular, this allows an application of the Halperin-Stasheff-Félicx perturbation theory to our context.

First, it is sometimes convenient to have *minimal*  $\mathcal{M}$ -free resolutions for a DGL, defined for any CUGA  $\mathcal{C}$  as follows:

**5.1. Definition.** Any  $B \in \mathcal{C}$  has a special kind of free simplicial resolution (see §3.2)  $A_\bullet \rightarrow B$ , called a *CW-resolution*, defined as follows (cf. [B11, §5.3]):

Assume that for each  $n \geq 0$ ,  $A_n = F(T_*^n)$  is the free graded algebra on the graded set  $T_*^n$ , and that the degeneracies of  $A_\bullet$  take  $T_*^n$  to  $T_*^{n+1}$ . Let  $\bar{A}_n$  denote the sub-algebra of  $A_n$  generated by the non-degenerate elements in  $T_*^n$ . Then we require that  $d_i|_{\bar{A}_n} = 0$  for  $1 \leq i \leq n$ . The sequence  $\bar{A}_0 = A_0, \bar{A}_1, \dots, \bar{A}_n, \dots$  is called a *CW-basis* for  $A_\bullet$ , and  $\bar{d}_0 = d_0|_{\bar{A}_n}$  is the *attaching map* for  $\bar{A}_n$ .

Such a  $A_\bullet \rightarrow B$  will be called *minimal* if each  $\bar{A}_{n+1}$  is minimal among those free algebras in  $\mathcal{C}$  which map onto the Moore  $n$ -cycles  $Z_n A_\bullet = \text{Ker}(\partial_n)$  (see (2.11)).

**5.2. Definition.** When  $\mathcal{C} = \mathcal{L}$ , the category of graded Lie algebras, it will be more convenient at times to use of the adjoint functors of Proposition 2.9 to replace  $A_{\bullet\bullet} \rightarrow X_\bullet$  by a simplicial DGL  $L_{\bullet,*} \rightarrow X_*$ . In this case the simplicial models are replaced by the corresponding DGLs, namely

- (1) A *d $\mathcal{L}$ - $n$ -sphere*,  $\mathcal{S}_{(x)}^n$ , is a DGL of the form  $(\mathbb{L}\langle X_* \rangle, 0)$  where  $X_*$  is the graded set with  $X_n = \{x\}$  and  $X_i = \emptyset$  for  $i \neq n$ .
- (2) A *d $\mathcal{L}$ - $(n+1)$ -disk*, denoted  $\mathcal{D}n+1_{(x)}$ , is the DGL  $(\mathbb{L}\langle X_* \rangle, \partial_L)$  where  $X_{n+1} = \{x\}$ ,  $X_n = \{\partial_L x\}$ , and  $X_i = \emptyset$  for  $i \neq n, n+1$ . Its *boundary* is the *d $\mathcal{L}$ - $n$ -sphere*  $\partial \mathcal{D}n+1_{(x)} := \mathcal{S}_{(\partial_L x)}^n$ .
- (3) A *two-stage DGL* is a DGL  $(\mathbb{L}\langle X_* \rangle, \partial_L) \in d\mathcal{L}$ , where for some  $n \geq 0$  we have  $X_i = \emptyset$  for  $i \neq n, n+1$ . Any coproduct (in  $d\mathcal{L}$ ) of two-stage DGLs will be called a *free* DGL.

Evidently  $d\mathcal{L}$ -spheres and disks are free DGLs, and any free DGL may be described as the coproduct of  $d\mathcal{L}$ -spheres and disks – more precisely, as a coproduct of  $d\mathcal{L}$ -spheres, disks, and collections of disks with their boundaries identified to a single sphere.

**5.3. Definition.** Following Stover, we define a comonad  $F : d\mathcal{L} \rightarrow d\mathcal{L}$  by setting

$$(5.4) \quad F(B) := \left( \prod_{k=1}^{\infty} \prod_{x \in B_k} \mathcal{D}k_{(x)} \right) / \sim,$$

for any  $B = (B_*, \partial_B) \in d\mathcal{L}$ , where we set  $\mathcal{D}k_{(x)} := \mathcal{S}_{(x)}^k$  if  $\partial_B x = 0$ , and let  $\partial \mathcal{D}k + 1_{(x)} \sim \mathcal{S}_{(\partial_B x)}^k$  if  $\partial_B x \neq 0$ .

Clearly  $F(B)$  is a free DGL, and by iterating  $F$  we obtain a free simplicial DGL  $W_{\bullet,*}$  with  $W_n = F^{n+1}(B)$  (see [Gd, App., §3]), which we call the *canonical free simplicial DGL resolution* of  $B = (B_*, \partial_B)$ , which we denote by  $W_{\bullet,*}(B)$ . Observe that  $W_{\bullet,*}$  (or equivalently, the corresponding bisimplicial Lie algebra  $W_{\bullet\bullet}$ ) is an  $\mathcal{M}$ -free resolution of  $B$ .

**5.5. Remark.** Note that if  $\partial_B \equiv 0$ , by definition (5.4)  $F(B_*, 0)$  has only spheres, and no disks, and thus the canonical resolution  $W_{\bullet,*}(B)$  has  $\partial_{W_n} = 0$  for all  $n \geq 0$ . Thus  $W_{\bullet,*}$  may be identified with the usual canonical resolution of the graded Lie algebra  $B_*$  (coming from the “free graded Lie algebra on underlying graded set” comonad), which we shall denote by  $V_{\bullet,*}(B_*)$ .

However – unlike the canonical resolution – the construction above can be mimicked topologically (cf. [St, §2.3]). Since we want to present our results in a manner which could be generalized (as far as possible) to the integral case, we have chosen the somewhat convoluted description of (5.4).

Note further that by (3.4), if we apply the functor  $H'_*$  to  $W_{\bullet,*} \rightarrow B_*$  – or equivalently, the functor  $\pi_*^i$  to  $W_{\bullet\bullet} \rightarrow B_{\bullet}$  – we obtain a free simplicial resolution of the graded Lie algebra  $L_* := H'_*(B_*, \partial_B)$ .

**5.6. Notation.** If we write  $\langle x \rangle \in F(B)$  for the generator corresponding to an element  $x \in B_*$ , then recursively a typical DGL generator for  $W_n = W_{n,*}$  (in the canonical resolution  $W_{\bullet,*}(B)$ ) is  $\langle \alpha \rangle$ , for  $\alpha \in W_{n-1}$ , so an element of  $W_n$  is a sum of iterated Lie products of elements of  $B_*$ , arranged within  $n + 1$  nested pairs of brackets  $\langle \langle \dots \rangle \rangle$ . With this notation, the  $i$ -th face map of  $W_{\bullet,*}$  is “omit  $i$ -th pair of brackets”, and the  $j$ -th degeneracy map is “repeat  $j$ -th pair of brackets”. The operation of bracketing is defined to be linear  $\langle - \rangle$  is

linear – i.e., we set  $\langle \alpha x + \beta y \rangle = \alpha \langle x \rangle + \beta \langle y \rangle$  for  $\alpha, \beta \in \mathbb{Q}$  and  $x, y \in B$ .

In order to construct minimal  $\mathcal{M}$ -simplicial resolutions, first consider the coformal case:

**5.7. The bigraded model.** Any coformal DGL (§2.4), and in particular  $L = (L_*, 0)$ , has a *bigraded model*  $A_{*,*} \rightarrow L_*$  – that is, a bigraded DGL  $(A_{*,*}, \partial_A)$  (see §2.6) which is minimal in the sense of §2.5 (so in particular free as a graded Lie algebra), along with a quasi-isomorphism  $A_{*,*} \rightarrow L_*$ . The bigraded model is unique up to isomorphism. See [O, Part I] for an explicit construction.

This is just the Lie algebra version of the bigraded model of [HS, §3] (see also [F2]), which is in turn essentially the Tate-Jozefiak resolution (see [J]) of a graded commutative algebra.

$A = (A_*, \partial_A)$  will denote the DGL associated to  $A_{*,*}$  (Definition 2.8); by construction  $A$  is the minimal model (§2.5) for  $L$  (which is not minimal itself, unless  $L_*$  happens to be a *free* graded Lie algebra).

**5.8. Proposition.** *Let  $L = (L_*, 0) \in d\mathcal{L}$  be a coformal DGL, and  $\phi : A_{*,*} \rightarrow L_*$  its bigraded model; then there is an  $\mathcal{M}_{d\mathcal{L}}$ -free simplicial resolution  $C_{\bullet,*} \rightarrow L$ , with a bijection  $\theta : X_{**} \xrightarrow{\sim} C_{\bullet,*}$  between a bigraded set  $X_{**}$  of generators for  $A_{*,*}$  and the set of non-degenerate (§5.1)  $d\mathcal{L}$ -spheres in  $C_{\bullet,*}$ . Moreover,  $H'_*(C_{\bullet,*})$  is a minimal CW-resolution of  $L_* = H'_*(A_{*,*})$ , with CW basis generated by  $\text{Im}(\theta)$ .*

*Proof.* By Proposition 2.9 there is a simplicial graded Lie algebra resolution  $C_{\bullet,*} \rightarrow L_*$  corresponding to  $A_{*,*}$ , and thus a weak equivalence of simplicial graded Lie algebras  $\psi : C_{\bullet,*} \rightarrow V_{\bullet,*} = V_{\bullet,*}(L_*)$  (see §5.5), which is one-to-one because  $A_{*,*}$ , and thus  $C_{\bullet,*}$ , are minimal (cf. [BL, §2]).

Now let  $W_{\bullet,*}$  be the canonical free simplicial DGL resolution of  $A_{*,*}$ ; the fact that  $\phi : A_{*,*} \rightarrow L_*$  is a quasi-isomorphism implies that there is a weak equivalence  $\varphi : V_{\bullet,*} \rightarrow W_{\bullet,*}$  (as well as one in the other direction). The composite  $\varphi \circ \psi : C_{\bullet,*} \rightarrow W_{\bullet,*}$  is again a one-to-one weak equivalence (by minimality); we may therefore think of  $C_{\bullet,*}$  as a sub-simplicial object of  $W_{\bullet,*}$ .

Moreover, there is an embedding of bigraded vector spaces  $\eta : A_{*,*} \rightarrow C_{\bullet,*}$  (see proof of Proposition 2.9), and thus another such embedding  $\theta : A_{*,*} \rightarrow W_{\bullet,*}$ , which may be defined explicitly as follows (using the notation of §5.6):

For  $x \in X_{0,*}$ , set  $\theta(x) = \langle x \rangle \in C_{0,*} = F(A_*)$ . Since  $\phi$  maps  $X_{0,*}$  onto a (minimal) set of Lie algebra generators for  $L_* = H'_*(A_{*,*})$ , each

$\theta(x)$  is a  $\partial_W$ -cycle, so  $C_{0,*}^{(0)} := \coprod_{k=1}^{\infty} \coprod_{x \in X_{0,k}} \mathcal{S}_{(\theta(x))}^k$  is a sub free DGL of  $W_{0,*}$ .

By minimality of  $A_{*,*}$ , any  $x \in X_{n,*}$  ( $n \geq 1$ ) is uniquely determined by  $\partial_A(x) \in A_{n-1,*}$ . Thus if we require  $\theta$  to be multiplicative (with respect to the ordinary bracket in  $A_{*,*}$ , and with respect to the simplicial Lie bracket  $\llbracket \cdot, \cdot \rrbracket$  of (2.10) in  $W_{\bullet,*}$ ), we may define  $\theta : A_{*,*} \rightarrow W_{\bullet,*}$  inductively by

$$(5.9) \quad \theta(x) = \langle \theta(\partial_A(x)) \rangle,$$

and we shall write  $x^{(0)}$  for  $\theta(x)$  if  $x \in X_{**}$ .

By definition (see Proposition 2.9),  $d_0 \circ \theta = \theta \circ \partial_A$ , so for  $x \in X_{n,*}$  ( $n \geq 2$ ) we have  $d_1(x^{(0)}) = d_1(\theta(\partial_A(x))) = \langle d_0 \theta(\partial_A(x)) \rangle = \langle \theta(\partial_A^2(x)) \rangle = 0$ , while  $\varepsilon(d_1(x^{(0)}))$  is a  $\partial_A$ -boundary for  $x \in X_{1,*}$  (where  $\varepsilon : W_{0,*} \rightarrow A_*$  is the augmentation). Thus Lemma 2.12 below implies that for  $x \in X_{n,*}$  ( $n \geq 1$ ) we have  $d_i(x^{(0)}) = 0$  for all  $1 \leq i \leq n-1$ , while  $d_n(x^{(0)})$  is a  $\partial_W$ -boundary.

Therefore, if we set  $C_{n,*}^{(0)} := \coprod_{k=1}^{\infty} \coprod_{x \in X_{n,k}} \mathcal{S}_{(x^{(0)})}^k$  for all  $n \geq 0$ , we see  $\{H'_*(C_{n,*}^{(0)})\}_{n=1}^{\infty}$  is an  $\mathcal{L}$ -CW basis for  $H'_*(C_{\bullet,*})$ .

In order to give an explicit description of  $C_{\bullet,*}^{(0)}$  in terms of  $C_{\bullet,*}^{(0)}$ , we need to know the Lie disks in which  $d_n(x^{(0)})$  (and their faces) lie. By a double induction on  $n \geq 1$  and  $1 \leq r \leq n$ , we shall now define, for all  $x \in X_{n,k}$ , elements  $x^{(r)} \in W_{n-r,k+r}$  such that  $\partial_W(x^{(r)}) = d_{n-r}(x^{(r-1)})$ :

Note that for each  $x \in A_{n,*}$  we have

$$\partial_A(x) = \sum_t a_t \omega_t[y_{i_1}, \dots, y_{i_{m_t}}],$$

where  $\omega_t[\dots]$  is some  $m_t$ -fold iterated Lie bracket,  $y_{i_j} \in X_{n_j,*}$  with  $\sum_{j=1}^{m_t} n_j = n$ , and  $a_t \in \mathbb{Q}$ . Then

$$(5.10) \quad \theta(x) = \langle \theta(\partial_A(x)) \rangle = \left\langle \sum_t a_t \omega_t \llbracket y_{i_1}^{(0)}, \dots, y_{i_{m_t}}^{(0)} \rrbracket \right\rangle,$$

where  $\omega_t[\dots]$  is the same  $m_t$ -fold iterated Lie bracket as above, but now with respect to the simplicial Lie bracket  $\llbracket \cdot, \cdot \rrbracket$ , rather than  $[\cdot, \cdot]$ .

If we set  $x^{(s)} = 0$  for  $i > n$ , we may define  $x^{(s)}$  for  $0 < s \leq n$  inductively by:

$$(5.11) \quad x^{(s)} = \left\langle \sum_t a_t \sum_{\substack{r_1 + \dots + r_{m_t} = s \\ 0 \leq r_j}} \omega_t \llbracket y_{i_1}^{(r_1)}, \dots, y_{i_{m_t}}^{(r_{m_t})} \rrbracket \right\rangle \in C_{n-s, k-n+s}^{(s)}.$$

Thus if we assume by induction that we have chosen  $y_{i_j}^{(r_j)}$  with  $\partial_W(y_{i_j}^{(r_j)}) = d_{n_{j_0}}(y_{i_j}^{(r_j-1)})$ , it follows from Lemma 2.12 below that indeed  $\partial_W(x^{(s+1)}) = d_n(x^{(s)})$  and  $d_i(x^{(s)}) = 0$  for  $0 < i < n$ .

For example,  $y^{(0)} = \langle y \rangle$  and  $\varepsilon(\langle y \rangle) = y \in A_k$  for any  $y \in X_{0,k}$ . Therefore, for  $x \in X_{1,*}$  we have  $\varepsilon d_1(x^{(0)}) = \varepsilon d_0(x^{(0)}) = \partial_A(x)$ , so we may set  $x^{(1)} = \langle x \rangle \in W_{1,k+1}$ , with  $\partial_W(x^{(1)}) = d_1(x^{(0)})$ .

Now if we define by induction

$$C_{n,*}^{(r)} := C_{n,*}^{(r-1)} \amalg \prod_{k=1}^{\infty} \prod_{x \in X_{n+r,k}} \mathcal{D}k + r_{(x^{(r)})},$$

then it is not hard to see that  $C_{\bullet,*}$  is the sub-simplicial graded Lie algebra of  $W_{\bullet,*}$  generated (under the degeneracies of  $W_{\bullet,*}$ ) by  $(C_{n,*}^{(r)})_{r=0}^n$  for all  $n \in \mathbb{N}$ . (In particular, this is closed under face maps and includes  $\text{Im}(\theta)$ , and  $\theta : A_{*,*} \rightarrow C_{\bullet,*}$  is a weak equivalence. The only non-degenerate Lie spheres in  $C_{\bullet,*}$  are those of  $C_{\bullet,*}^{(0)}$ , as required.  $\square$ )

**5.12. The filtered model.** If  $B = (B_*, \partial_B) \in d\mathcal{L}$  is an arbitrary DGL, it no longer has a bigraded model, as in §5.7 above. However, it does have a *filtered model*, constructed as follows:

Let  $L_* := H'_*(B)$  be the homotopy Lie algebra of  $B_*$ , and  $(A_{*,*}, \partial_A)$  the bigraded model for  $(L_*, 0)$ . The filtered model for  $B$  is the free graded Lie algebra  $A_{*,*}$ , equipped with an increasing filtration  $0 = \mathcal{F}^{-1}(A) \subset \mathcal{F}^0(A) \subset \dots \subset \mathcal{F}^r(A) \subset \mathcal{F}^{r+1}(A) \subset \dots$  (defined by  $\mathcal{F}^r(A) := \bigoplus_{i=0}^r A_{i,*}$ ), and a new differential  $D_A = \partial_A + \delta_A$  on  $A_{*,*}$  such that  $\delta_A : A_{n,*} \rightarrow \mathcal{F}^{n-2}(A)$  (Of course,  $D_A$  is still required to be a derivation).

We may decompose  $D_A : A_{n,*} \rightarrow A_{*,*}$  as  $D_A = \partial_0 + \partial_1 + \dots + \partial_{n-1}$ , where  $\partial_r : A_{n,*} \rightarrow A_{n-r-1,*}$  (and  $\partial_0 = \partial_A$ , the original differential of the bigraded model).

See [O, II] or [Har]; this is again the Lie algebra version of a construction of Halperin-Stasheff and Félix in [HS, §4],[F1].

Note that the filtered model is no longer unique, since its construction depends on choices; in particular, it is not necessarily minimal. One again has the associated DGL  $(A_*, D_A)$ , which is quasi-isomorphic to the original DGL  $B$ , and  $A_{*,*}$  is obtained by filtering  $A_*$ .

**5.13. Proposition.** *Let  $B = (B_*, \partial_B)$  be a DGL, and  $(A_{*,*}, D_A)$  a filtered model for  $B$ ; then there is an  $\mathcal{M}_{d\mathcal{L}}$ -free simplicial resolution  $E_{\bullet,*} \rightarrow B$ , with a bijection  $\theta : X_{**} \xrightarrow{\sim} E_{\bullet,*}$  between a bigraded set  $X_{**}$  of generators for  $A_{*,*}$  and the set of non-degenerate  $d\mathcal{L}$ -spheres in  $E_{\bullet,*}$ .*

*Proof.* We start with the minimal  $\mathcal{M}$ -free resolution  $C_{\bullet,*} \rightarrow L_*$  for  $L_* = H'_*(B_*)$ , constructed as in the proof of Proposition 5.8, and deform it into an  $\mathcal{M}$ -free resolution for  $B$ , using the filtered model  $(A_{*,*}, D_A)$  as a guideline. This time we shall embed the resulting  $\mathcal{M}$ -free resolution in the canonical free DGL resolution  $W_{\bullet,*}$  of  $(A_*, D_A)$ , the DGL associated to the filtered model:

For each  $x \in X_{n,k}$  (where  $X_{**}$  is a bigraded set of generators for the bigraded Lie algebra  $A_{*,*}$ , as above), set  $x^{(n)} = \langle x \rangle \in W_{0,k}$ , and let  $D_A(x) = \partial_0(x) + \partial_1(x) + \cdots + \partial_{n-1}(x)$  as above, with

$$\partial_r(x) = \sum_t a_t^{(r)} \omega_t^{(r)}[y_{i_1}, \dots, y_{i_{m_t}}] \in A_{n-r-1,*},$$

where  $\omega_t^{(r)}[\dots]$  is some  $m_t$ -fold iterated Lie bracket, as above, and each  $y_{i_j} \in X_{n_j,*}$  with  $\sum_{j=1}^{m_t} n_j = n - r - 1$ .

If we set  $x^{(s)} = 0$  for  $i > n$ , we may define  $x^{(s)}$  for  $0 < s \leq n$  inductively by:

$$(5.14) \quad x^{(s)} = \left\langle \sum_{r=0}^s \sum_t a_t^{(r)} \sum_{\substack{r_1+\dots+r_{m_t}=s-r \\ 0 \leq r_j}} \omega_t^{(r)} \llbracket y_{i_1}^{(r_1)}, \dots, y_{i_{m_t}}^{(r_{m_t})} \rrbracket \right\rangle \in C_{n-s, k-n+s}^{(s)}$$

Using Lemma 2.12 and the fact that for any  $A_\bullet \in s\mathcal{L}$ ,  $x \in A_p$  and  $y \in A_q$  we have  $d_{p+q}(\llbracket x, y \rrbracket) = \llbracket d_p(x), y \rrbracket + (-1)^q \llbracket x, d_q(y) \rrbracket$ , one may then verify inductively that  $d_{n-s}(x^{(s)}) = \partial_W(x^{(s+1)})$  and  $d_i(x^{(s)}) = 0$  for  $0 < i < n - s$ , for all  $0 \leq s < n$ . The rest of the construction is as in the proof of Proposition 5.8.  $\square$

We have the following analogue of Definition 4.4:

**5.15. Definition.** Given a DGL  $L = (\mathbb{L}\langle X_* \rangle, \partial_L) \in d\mathcal{L}$  and a simplicial set  $A \in \mathcal{S}$ , we define their *half-smash*  $L \rtimes A = (\mathbb{L}\langle Y_* \rangle, \partial')$  by setting  $Y_n := \coprod_{k=0}^n X_k \times \hat{A}_{n-k}$ , where  $\hat{A}_i$  denotes the set of non-degenerate  $i$ -simplices of  $A$ . For  $a \in \hat{A}_k$  and  $x \in X_m$ , we set

$$\partial'(x, a) = \sum_{i=0}^k (-1)^{i+m} (x, d_i a) + (\partial_L x, a)$$

(and extend  $\partial'$  by requiring that it be a derivation).

**5.16. Remark.** In order to apply the obstruction theory described in §4.15, note that all the definitions of section 4 pass over to the DGL setting in a straightforward manner. However, if we now start with the trivial DGL  $A = L_*^{(0)} := (\Pi_*^X, 0)$ , we may take  $C_{\bullet,*} \rightarrow L_*^{(0)}$  to be

the minimal  $\mathcal{M}$ -free resolution of Proposition 5.8, corresponding to the bigraded model for  $(\Pi_*^X, 0)$ , and let  $B = (B_*, \partial_B)$  (corresponding to  $B_\bullet$  in §4.15) be the filtered model for  $L_X$ . We assume that  $A \not\cong B$ .

As explained in §4.15, there is a least  $n_0 \geq 2$  such that  $0 \notin \langle\langle n_0 \rangle\rangle \subseteq H'_*(B)$ , and we write  $\langle\langle n_0 \rangle\rangle = (\langle\langle n_0, x_i \rangle\rangle)_{i \in I}$ , in the notation of 4.9, where  $x_i \in X_{n_0, i}$  and  $\mathcal{S}_{(x_i)}^{k_i}$  are corresponding DGL spheres in  $C_{n_0, *}$  (we include in the index set  $I$  only those coordinates of (4.11) which do not vanish).

Again let  $\mathcal{H}^{(1)}$  denote the set of all homotopy types in  $\text{ho } d\mathcal{L}(\mathbf{X})$  for which  $\langle\langle n_0 \rangle\rangle$  has the same value as for  $L_X$ , and choose a representative  $L^{(1, \alpha)} \in d\mathcal{L}(\mathbf{X})$  for each  $\alpha \in \mathcal{H}^{(1)}$ . By [HS, §3], we may assume  $L^{(1, \alpha)}$  is obtained from  $B$  by perturbation of  $\partial_B$ . Proceeding as in §4.15 we obtain a tree of DGLs  $L^{(k, \alpha_1, \alpha_2, \dots, \alpha_k)} \in d\mathcal{L}(\mathbf{X})$ , and by [B11, Theorem 3.1], we know that  $L^{(k, \alpha_1, \alpha_2, \dots, \alpha_k)}$  may be chosen to agree with  $L_X$  through degree  $n+1$  at least, so  $\text{colim}_n L^{(n)} \simeq L_X$  along any branch of the tree.

Note also that because  $H'_*(C_{\bullet, *})$  is a (minimal) CW resolution of  $H'_*(B)$ , in each case, the maps  $\psi_n : H'_*(C_{n, *}) \rightarrow H'_*(B)$  are null for all  $n \geq 1$  (see §4.2). Thus any  $\partial D[n]$ -compatible collection  $\{h_k\}_{k=0}^{n-1}$  in §4.9 induce a map  $C_{n, *} \wedge \partial D[n] \rightarrow B$  directly, without need of the splitting (4.10).

5.17. *Remark.* The second order operation described in the previous example is actually a secondary Whitehead product. Unlocalized higher order Whitehead products were defined by G. Porter in [P, 1.3], and the relation between this definition and the rational version has been studied by several authors – see [AA], [A1, A2], [R2, R1] and [T, V.1]. However, there are other higher order rational homotopy operations, too:

For example, in the DGL  $L_* = (\mathbb{L}\langle a_1, b_1, c_1, d_1, x_4, y_4, z_4, w_4 \rangle, \partial)$ , with  $\partial(x) = [[b, a], c]$ ,  $\partial(y) = [[b, a], d]$ ,  $\partial(z) = [[d, c], a]$  and  $\partial(w) = [[d, c], b]$ , the cycle  $[x, d] + [y, c] + [z, b] + [w, a]$  represents such an operation. There appears to be no general procedure for representing these as *integral* higher order operations in  $\pi_*\mathbf{X}$ ; we shall offer a (partial) answer to this difficulty in section 7.

## 6. HOMOLOGY OF DGLS

Obstructions in algebraic topology traditionally take values in suitable cohomology groups. In order to show that this holds in our setting, too, we recall Quillen's definition of homology and cohomology in model categories:

**6.1. Definition.** An object  $X$  in a category  $\mathcal{C}$  is said to be *abelian* if it is an abelian group object – that is, if  $\mathrm{Hom}_{\mathcal{C}}(Y, X)$  has a natural abelian group structure for any  $Y \in \mathcal{C}$ . When  $\mathcal{C}$  is  $\mathcal{L}ie$ ,  $\mathcal{A}lg$ ,  $s\mathcal{L}ie$ ,  $s\mathcal{A}lg$ ,  $\mathcal{L}$ , or  $d\mathcal{L}$ , for example, this is equivalent to requiring that all products vanish in  $X$  (cf. [BS, §5.1.3]).

The full subcategory of abelian objects in  $\mathcal{C}$  is denoted by  $\mathcal{C}_{ab} \subset \mathcal{C}$ . In the cases of interest to us, this will itself be an abelian category. It is equivalent to the category  $\mathcal{V}ec$  of vector spaces if  $\mathcal{C} = \mathcal{L}ie$  or  $\mathcal{A}lg$ , to  $\mathcal{V}$  if  $\mathcal{C} = \mathcal{L}$ , to the category  $s\mathcal{V}ec$  of simplicial vector spaces if  $\mathcal{C} = s\mathcal{L}ie$  or  $s\mathcal{A}lg$ , and to the category  $d\mathcal{V}$  if  $\mathcal{C} = d\mathcal{L}$  (see §1.1).

In these cases, we have an *abelianization* functor  $Ab : \mathcal{C} \rightarrow \mathcal{C}_{ab}$ , along with a natural transformation  $\theta : Id \rightarrow Ab$  having the appropriate universal property. In all the examples above,  $Ab(X) = X/I(X)$ , where  $I(X)$  is the ideal in  $X \in \mathcal{C}$  generated by all non-trivial products.

**6.2. Definition.** Let  $\mathcal{C}$  be a category as above, which also has a closed model category structure: in [Q1, II, §5] (or [Q4, §2]), Quillen defines the *homology* of an object  $X \in \mathcal{C}$  to be the total left derived functor  $\mathbf{L}(Ab)$  of  $Ab$ , applied to  $X$  (cf. [Q1, I, §4]).

In more familiar terms, this means that we construct a resolution  $A \rightarrow X$  (i.e., replace  $X$  by a weakly equivalent cofibrant object  $A \in \mathcal{C}$ ), and then define the  $i$ -th homology group of  $X$  by  $H_i X := \pi_i(Ab(A))$ , for an appropriate concept of homotopy groups  $\pi_*$  in  $\mathcal{C}_{ab}$  (see [Q1, II, §4]). One must verify, of course, that this definition is independent of the choice of the resolution  $A \rightarrow X$ .

Similarly, the *cohomology* of  $X$  with coefficients in  $M \in \mathcal{C}_{ab}$  is defined:

$$H^i(X; M) := [\mathbf{L}(Ab)X, \Omega^{i+N} \Sigma^N M]_{\mathcal{C}_{ab}} \quad \text{for } N \quad \text{large enough}$$

(where the loop and suspension functors  $\Omega$  and  $\Sigma$  are defined in [Q1, I, §2]).

Again, in the cases that interest us,  $\Omega$  is essentially the shift operator  $\Sigma^{-1}$  of §1.1, and so the  $i$ -th cohomology group of  $X$  with coefficients in  $M$  is then  $H^i(X; M) := [\Sigma^i Ab(A), M]_{\mathcal{C}_{ab}}$ .

**6.3. Definition.** If  $\mathcal{C}$  itself does not have a closed model category structure, one often defines the homology of  $\mathbf{X} \in \mathcal{C}$  by embedding  $\mathcal{C}$  in some category which does have such a structure, which in many cases may be taken to be  $s\mathcal{C}$ , the category of simplicial objects over  $\mathcal{C}$  (see [Q1, II, §4]). Thus, if  $\iota : \mathcal{C} \hookrightarrow s\mathcal{C}$  is the embedding of categories defined by taking  $\iota(C)$  to be the constant simplicial object equal to  $C$  in all dimensions, then  $H_i(C) := \pi_i(\mathbf{L}(Ab \circ \iota)C)$ .

This is the approach usually taken for  $\mathcal{C} = \mathcal{L}ie, \mathcal{A}lg,$  or  $\mathcal{L}$ : to define the homology of a graded Lie algebra  $L_* \in \mathcal{L}$ , say, one chooses a free simplicial resolution  $A_{\bullet,*} \rightarrow L_*$  (such as the canonical resolution – cf. §5.3), and then calculate the homotopy groups of the simplicial graded vector space  $Ab(A_{\bullet,*}) \in s\mathcal{V}$  (or the homology groups of the bigraded chain complex in  $db\mathcal{V}$  corresponding to  $Ab(A_{\bullet,*})$  – see proof of Proposition 2.9).

6.4. *Remark.* Note that if we apply Definition 6.2 as is to a DGL  $L = (L_*, \partial_L) \in d\mathcal{L}$ , we may take the resolution  $A$  to be the minimal model  $\hat{L} = (\hat{L}_*, \hat{\partial})$  for  $L$  (cf. §2.5), and since its abelianization is just the graded vector space  $Q(\hat{L})$  of indecomposables, and  $Q(\hat{\partial}) = 0$  by definition,  $H_i(L)$  would be isomorphic to the vector space spanned by a set of generators for  $\hat{L}$  in dimension  $i$ .

If we want cohomology with coefficients in an object  $M_* \in d\mathcal{L}_{ab} \approx d\mathcal{V}$  with *trivial* differential – i.e.,  $M_*$  is just a graded vector space – we find

$$H^i(X; M_*) \cong \prod_{j=1}^{\infty} \text{Hom}_{\mathcal{V}ec}(H_j(X), M_{i+j}),$$

by the universal coefficients theorem.

However, since  $L$  is itself graded, we would like  $H_*L$  to be bigraded (with a “homological” degree, as well as a “topological” one). This requires a combination of Definitions 6.2 and 6.3, as follows:

6.5. **Definition.** The *homology*  $H_{*,*}(L_{\bullet})$  of a simplicial Lie algebra  $L_{\bullet} \in s\mathcal{L}ie$  is defined to be the left derived functors of the abelianization, with respect to the  $E^2$ -closed model category structures (§3.3) on  $ss\mathcal{L}ie$  and  $ss\mathcal{L}ie_{ab} \approx dd\mathcal{V}$  respectively. More precisely,

$$(6.6) \quad H_{s,t}(L_{\bullet}) := \pi_s(\mathbf{L}(Ab \circ \iota)L_{\bullet})_t = \pi_s \pi_t^i(AbA_{\bullet\bullet}),$$

where  $A_{\bullet\bullet} \rightarrow L_{\bullet}$  is some  $\mathcal{M}$ -free bisimplicial resolution of  $L_{\bullet}$ .

Similarly, for any DGL  $L \in d\mathcal{L}$  we may define  $H_{s,t}(L) := \pi_s H_t'(Ab(A_{\bullet,*}))$ , for a  $\mathcal{M}_{d\mathcal{L}}$ -free simplicial resolution  $A_{\bullet,*} \rightarrow L$ ; and these two definitions agree under the equivalence of homotopy categories  $\text{ho}(s\mathcal{L}ie) \approx \text{ho}(d\mathcal{L})$  of Proposition 2.2.

The bigraded cohomology of a DGL  $L$  with coefficients in the abelian DGL (i.e., chain complex)  $M$  is defined analogously as

$$H_t^s(L; M) := \pi^s(\text{Hom}_{d\mathcal{L}_{ab}}(Ab(A_{\bullet,*}), M)_t).$$

We note that the homology and cohomology of differential graded (commutative) algebras have been defined by Goodwillie (in [Go]) and

Burghelea & Vigué-Poirrier (in [BV]), in a manner analogous to the traditional definitions of Hochschild homology. See [Lo, §5.3].

**6.7. Proposition.** *For any DGL  $L \in d\mathcal{L}$ , there is a monomorphism of graded vector spaces  $H_{n,t}(L) \hookrightarrow H_{n,t}(L')$ , where  $L' \simeq (H'_*(L), 0)$  is the coformal model for  $L$ ; the same holds for cohomology with trivial coefficients.*

*Proof.* If  $A_{*,*}$  is the bigraded model for  $L'$ , and  $C_{\bullet,*} \rightarrow L'$  the simplicial resolution of Proposition 5.8, then the non-degenerate spheres  $\mathcal{S}_{(x^{(0)})}^k \subset C_{n,t}$ , which correspond to a vector space basis for  $H_{n,t}(L')$ , are in bijective correspondence with the generators  $x \in X_{n,t}$  for  $A_{*,*}$ .

Now let  $B_{*,*}$  be a filtered model for  $L$  obtained by perturbing  $(A_{*,*}, \partial_A)$ , and  $E_{\bullet,*} \rightarrow L$  the associated simplicial resolution of Proposition 5.13: since  $B_{*,*}$  need no longer be minimal (§5.12), a vector space basis for  $H_{n,t}(L)$  now corresponds to a *subset* of the collection of non-degenerate spheres  $\mathcal{S}_{(x^{(0)})}^k \subset E_{n,t}$ , (which are still in bijective correspondence with the generators  $x \in X_{n,t}$  for  $A_{*,*}$  or  $B_{*,*}$ ).  $\square$

Note that this description of the homology implies that  $H_{*,*}(L)$  is indeed just a bigraded version of the DGL homology defined in §6.3.

**6.8. Proposition.** *The collection of higher operations  $\langle\langle n_{k,\alpha_1,\alpha_2,\dots,\alpha_k}, x \rangle\rangle$  which determine the rational homotopy type of  $\mathbf{X} \in \mathcal{T}_1$  (described in §4.15 above) are indexed by elements  $x \in H_{n_k,\alpha_1,\alpha_2,\dots,\alpha_k,t}(L^{(k,\alpha_1,\alpha_2,\dots,\alpha_k)})$  in the homology of the DGLs of §5.16, and take value in the cohomology of these DGLs, with*

$$\langle\langle n_{k,\alpha_1,\alpha_2,\dots,\alpha_k}, x \rangle\rangle \subseteq H_{t+n_k-1}^{n_k,\alpha_1,\alpha_2,\dots,\alpha_k}(L^{(k,\alpha_1,\alpha_2,\dots,\alpha_k)}; \pi_* \mathbf{X}_{\mathbb{Q}}).$$

*Proof.* We may construct a simplicial resolution  $E_{\bullet,*}$  for each successive DGL  $L^{(k)} = L^{(k,\alpha_1,\alpha_2,\dots,\alpha_k)}$ , corresponding to the filtered models obtained as perturbations  $(A_{*,*}, D_A)$  of the bigraded model  $(A_{*,*}, \partial_A)$  for  $L^{(0)}$ , as above. The non-degenerate spheres  $\mathcal{S}_{(x)}^m = \mathcal{S}_{(x^{(0)})}^m \subset E_{m,t}$  which index the higher homotopy operations  $\langle\langle m, x \rangle\rangle$  are thus in bijective correspondence with the generators  $x \in X_{m,t}$  for  $A_{*,*}$ . However, if  $x$  is not minimal – in the sense that  $D_A(x) \notin [A, A]$ , or  $x + \alpha = D_a(y)$  for some  $\alpha \in A_{*,*}$  and  $y \in X_{m+1,t}$  – then we can construct a new simplicial resolution  $E'_{\bullet,*}$  of  $L^{(k)}$  in which  $\mathcal{S}_{(x)}^m$  has been eliminated (though of course new spheres may appear in higher simplicial dimensions). By the universal property of resolutions (i.e., of cofibrant objects in the  $E^2$  model category for  $sd\mathcal{L}$  – see §3.3) there is a map of resolutions  $E_{\bullet,*} \rightarrow E'_{\bullet,*}$ , and there can be no non-vanishing higher operation  $\langle\langle n_{k,\alpha_1,\dots,\alpha_k}, x \rangle\rangle$  which serves as an obstruction to rectifying the augmentation up-to-homotopy  $\varphi : E_{\bullet,*} \rightarrow L^X$ ,

since  $\varphi_n|_{S_{(x)}^m}$  can be factored through  $0 \in E'_{\bullet,*} \rightarrow L^X$ . Thus the only homotopy operations which can appear are those corresponding to *non-trivial* homology classes in  $H_*(L^{(k,\alpha_1,\alpha_2,\dots,\alpha_k)})$ . These yield the requisite cohomology classes by Universal Coefficients.  $\square$

Proposition 6.7 thus implies that we may if we like think of all the higher homotopy operations described in §4.15 (associated to the various deformations of  $L^{(0)}$ ) as lying in one fixed bigraded group  $H_*(L^{(0)}; \pi_* \mathbf{X}_{\mathbb{Q}})$ , which is of course just the usual cohomology of a graded Lie algebra, and is easier to compute than the cohomology of a non-trivial DGL.

## 7. NON-ASSOCIATIVE ALGEBRA MODELS

The DGL higher homotopy operations are unsatisfactory from a topological point of view because there is no obvious way to translate them, in general, into *unlocalized* topological homotopy operations. We now describe an algebraic model for rational homotopy theory which may serve to answer this objection.

**7.1. Non-associative graded algebras.** Let  $\mathcal{A}$  denote the category of *non-associative graded algebras*: an object  $A_* \in \mathcal{A}$ , is just a graded vector space equipped with a bilinear graded product  $A_p \otimes A_q \rightarrow A_{p+q}$ . Let  $\mathbb{A}\langle X \rangle$  denote the free non-associative graded algebra generated by a graded set  $X_*$ . As in §2.1, the functor  $\mathbb{A} : \text{grSet} \rightarrow \mathcal{A}$  factors through  $A : \mathcal{V} \rightarrow \mathcal{A}$ .

$d\mathcal{A}$  will denote the category of differential graded non-associative algebras  $(A_*, \partial_A)$ , called *DGNAs*; the differential  $\partial_A$  must satisfy  $\partial_A \circ \partial_A = 0$  and  $\partial_A(x \cdot y) = \partial_A x \cdot y + (-1)^{|x|} x \cdot \partial_A y$ , as for DGLs.

For simplicity we assume each  $A_* \in \mathcal{A}$ ,  $d\mathcal{A}$  is connected – that is,  $A_0 = \{0\}$ . Again, we have a category  $\text{db}\mathcal{A}$  of differential bigraded non-associative algebras (*DBGNAs*), as in §2.6.

As for any CUGA (§3.1), one can define a closed model category structure on  $s\mathcal{A}$  (see [Q1, II, §4]) and thus by [Bl4, §4] on  $\text{db}\mathcal{A}$ , and one has the expected analogues of Propositions 2.2 and 2.9:

**7.2. Proposition.** *There are adjoint functors  $s\text{Alg}_{N^*}^N d\mathcal{A}$ , which induce equivalences of the corresponding homotopy categories  $\text{ho}(s\text{Alg}) \approx \text{ho}(d\mathcal{A})$ .*

**7.3. Proposition.** *There are adjoint functors  $s\mathcal{A}_{N^*}^N \text{db}\mathcal{A}$ , which induce equivalences of the corresponding homotopy categories  $\text{ho}(s\mathcal{A}) \approx \text{ho}(\text{db}\mathcal{A})$ ; and  $N^*$  takes free DBGNAs to free simplicial non-associative algebras.*

**7.4. Notation.** For any  $(X_*, \cdot) \in \mathcal{A}$ , let  $[x, y]$  denote  $\frac{1}{2}(x \cdot y + (-1)^{|x||y|+1}y \cdot x)$ . We then have  $[y, x] = (-1)^{|x||y|+1}[x, y]$ , so  $(X_*, [, ])$  is now a non-associative graded algebra with a graded-commutative (or: graded skew-symmetric) multiplication. Moreover, any graded derivation  $\partial$  on  $(X_*, \cdot)$  is also a derivation with respect to  $[, ]$ , and any morphism of algebras from  $(X_*, \cdot)$  to a graded-commutative algebra will also respect  $[, ]$ . Therefore we can (and will) assume that our non-associative algebras are all graded-commutative, and denote the product by  $[, ]$ .

Moreover, if  $A_\bullet \in s\mathcal{A}$  is a simplicial graded algebra, we shall also use the notation

$$[x, y] = \sum_{(\sigma, \tau) \in S_{p,q}} (-1)^{\varepsilon(\sigma) + p|y|} [s_{\tau_q} \dots s_{\tau_1} x, s_{\sigma_p} \dots s_{\sigma_1} y]$$

for the corresponding simplicial bracket (compare (2.10)).

**7.5. Definition.** Any simplicial Lie algebra  $L_\bullet \in s\mathcal{L}ie$  is in particular an object in  $sAlg$ ; let  $\iota : d\mathcal{L} \hookrightarrow d\mathcal{A}$  be the inclusion functor. Note that even if each  $L_n$  is free as a Lie algebra, it is not free as a non-associative algebra: a free simplicial resolution  $J_\bullet \rightarrow \iota(L_\bullet)$  in the category  $sAlg$  (see §3.2) will be called a *sAlg-model* for  $L_\bullet$ . Such models can be constructed functorially, for example by a variant of §5.5.

There is also the analogous concept of a *dA-model*  $J_* \in d\mathcal{A}$  of a DGL  $L$ ; we can of course translate back and forth between these two types of models using Proposition 7.2.

Since the DGL  $L = (L_*, \partial_L)$  has an internal grading, and its *dA-model*  $J = (J_*, \partial_J)$  is constructed as a resolution of  $L$ , it is natural to define a second “homological” degree on  $J_*$ , so as to have a *filtered dA-model* (cf. §5.12). If the DGL is trivial (i.e.,  $\partial_L = 0$ ), a *dA-model* for  $(L_*, 0)$  will be a differential bigraded non-associative algebra (DBGNA)  $J = (J_{*,*}, \partial_J)$  (cf. §5.7).

**7.6. Remark.** Define a *Jacobi algebra* to be a DGNA  $J = (J_*, \partial_J) \in d\mathcal{A}$  such that  $H'_* J \in \mathcal{L}$ . In particular, any *dA-model*  $J$  of a DGL  $L$  is a Jacobi algebra, since  $H'_* J \cong H'_* L$ . We denote by  $\mathcal{J} \subset d\mathcal{A}$  the full subcategory of Jacobi algebras. These algebras are clearly related to the strongly homotopy Lie algebras of [SS] (see also [LM]), though in general a Jacobi algebra is just a “Lie algebra up to homotopy”.

Note that even though  $d\mathcal{A}$  itself is a CUGA,  $\mathcal{J}$  apparently is not, and it does not inherit many desirable properties from  $d\mathcal{A}$ : for example,  $\mathcal{J}$  is not closed under the coproduct in  $d\mathcal{A}$ . However, one still has *free* Jacobi algebras, in the following sense:

**7.7. Lemma.** *There is a functor  $J : d\mathcal{V} \rightarrow d\mathcal{A}$ , and a natural transformation  $\theta : J \rightarrow L$  such that:*

- (a)  $JV_*$  is free as an algebra, for any chain complex  $V_*$ ;
- (b)  $\theta_{V_*}$  is a surjective quasi-isomorphism;
- (c) any chain map  $\varphi : V_* \rightarrow K_*$  (where  $V_* \in d\mathcal{V}$  and  $K_* \in \mathcal{J}$ ) extends to a  $d\mathcal{A}$  map  $\hat{\varphi} : JV_* \rightarrow K_*$ .

*Proof.* As noted above, we may define  $J_{*,*} = J(V_*)$  by induction on the homological filtration:

Start with  $J_{0,*} = A(V_*)$  (the free non-associative algebra on the differential graded vector space  $(V_*, \partial_V)$ , with  $\partial_J$  extending  $\partial_V$  as a derivation), and let  $\theta_0 : J_{0,*} \rightarrow L(V_*)$  be the obvious surjection, with  $K_{0,*} = \text{Ker}(\theta_0)$  a two-sided  $d\mathcal{A}$ -ideal of  $J_{0,*}$ .

Choose once and for all some collection of generators  $M_0 = \{\mu_i\}_{i \in I}$  for  $K_{0,*}$  as a  $J_{0,*}$ -bimodule: for each choice of a  $d\mathcal{V}$ -basis  $\{x_\gamma\}_{\gamma \in \Gamma}$  for  $V_*$  – that is, of a graded vector space basis of the form  $\{x_\alpha, \partial_V x_\alpha, x_\beta\}$ , with  $\partial_V x_\beta = 0$  – we may write each  $\mu_i$  as some expression  $\mu_i(x_{\gamma_{i_1}}, \dots, x_{\gamma_{i_n}})$ . If we then choose some other  $d\mathcal{V}$ -basis  $\{x'_{\gamma'}\}_{\gamma' \in \Gamma'}$  for  $V_*$ , again we will have  $\mu'_i := \mu_i(x'_{\gamma'_{i_1}}, \dots, x'_{\gamma'_{i_n}}) \in K_{0,*}$ ; define  $J_{1,*}$  to be the free non-associative algebra on the DG vector subspace of  $K_{0,*}$  spanned by all such “canonical operations”  $\mu'_i$  ( $i \in I$ ), for all possible choices of  $d\mathcal{V}$ -bases  $\{x'_{\gamma'}\}_{\gamma' \in \Gamma'}$  for  $V_*$ . Again one has the obvious augmentation  $J_{1,*} \rightarrow J_{0,*}$  to serve as  $\partial_J$  (with  $\partial_J \circ \partial_J = 0$  by construction), and one takes the kernel  $K_{1,*}$  of this augmentation for the next step.

Proceeding in this way we may define the functor  $J$  by induction on the homological filtration; if the collections of operations  $M_n$  are chosen canonically, the functor itself will be canonical. Properties (a)–(c) are readily verified.  $\square$

**7.8. Example.** Let  $L = (\mathbb{L}\langle X_* \rangle, 0)$  be the trivial free DGL on a graded set  $X_*$ ; in this case the canonical DBGNA model  $(J_{*,*}, \partial_J) = J(X_*, 0)$  for  $L$  may be described in part as follows:

Let  $J_{0,*} = \mathbb{A}\langle X_* \rangle$  (the free non-associative algebra on  $X_*$ ). Since the Jacobi identity holds in  $\mathbb{L}\langle X_* \rangle$ , but not in  $\mathbb{A}\langle X_* \rangle$ , we have  $\mu(x, y, z) = [x, [y, z]] - [[x, y], z] + (-1)^{qr} [[x, z], y] \in K_{0,*}$  for all  $x, y, z \in X_*$ . Thus  $J_{1,*}$  will be generated as a  $J_{0,*}$ -bimodule by the image of  $(J_{0,*})^{\otimes 3}$  under the  $\Sigma_3$ -equivariant multilinear map  $\lambda_3 : J_{i,p} \otimes J_{j,q} \otimes J_{k,r} \rightarrow J_{i+j+k+1, p+q+r}$ . Here the symmetric group  $\Sigma_n$  acts on  $J_{*,p_1} \otimes \dots \otimes J_{*,p_n}$  by permutations, and on  $J_{*,p_1+\dots+p_n}$  via the Koszul sign homomorphism  $\varepsilon_I : \Sigma_n \rightarrow \{1, -1\}$  (defined by letting  $\varepsilon_I((k, k+1)) = (-1)^{i_k i_{k+1} + 1}$  for any adjacent transposition  $(k, k+1) \in$

$\Sigma_n$ ). We set

$$\partial_J(\lambda_3(x \otimes y \otimes z)) = [x, [y, z]] - [[x, y], z] + (-1)^{qr} [[x, z], y].$$

However, there are relations among these elements  $\lambda_3(x \otimes y \otimes z)$ , so we define a  $\Sigma_4$ -equivariant multilinear map  $\lambda_4 : J_{i,p} \otimes J_{j,q} \otimes J_{k,r} \otimes J_{\ell,s} \rightarrow J_{i+j+k+\ell+2, p+q+r+s}$ ,

$$\begin{aligned} \partial_J(\lambda_4(x \otimes y \otimes z \otimes w)) &= [x, \lambda_3(y \otimes z \otimes w)] + [\lambda_3(x \otimes y \otimes z), w] \\ &\quad - (-1)^{|z||w|} [\lambda_3(x \otimes y \otimes w), z] \\ &\quad + (-1)^{|y|(|z|+|w|)} [\lambda_3(x \otimes z \otimes w), y] \\ &\quad - (\lambda_3([x, y] \otimes z \otimes w) + \lambda_3(x \otimes y \otimes [z, w])) \\ &\quad + (-1)^{|y||z|} (\lambda_3([x, z] \otimes y \otimes z) \\ &\quad + \lambda_3(x \otimes z \otimes [y, w])) \\ &\quad - (-1)^{|w|(|y|+|z|)} (\lambda_3([x, w] \otimes y \otimes z) \\ &\quad + \lambda_3(x \otimes w \otimes [y, z])). \end{aligned}$$

In fact, one can define a sequence of “higher Jacobi relations”  $\lambda_n(x_1 \otimes \cdots \otimes x_n)$ , for all  $n \geq 3$ , which yield an explicit construction of  $J(X_*)$  for a the free (graded) Lie algebra  $LX_*$ . See [AB], and compare [LM, 2.1].

**7.9.  $\mathcal{A}$ -homotopy operations.** One can now apply the theory of section 4 verbatim to any space  $\mathbf{X} \in \mathcal{T}_1$  with  $\mathcal{C} = \mathcal{A}lg$  rather than  $\mathcal{L}ie$ , to obtain a sequence of higher homotopy operations as in §4.15 which determining the rational homotopy type of  $\mathbf{X}$  – the only difference being that the simplicial resolutions  $C_{\bullet\bullet}$  of the successive simplicial Lie algebras  $L_{\bullet}^{(k)}$  are now  $\mathcal{M}_{\mathcal{A}lg}$  free resolutions of  $L_{\bullet}^{(k)}$  in  $ss\mathcal{A}lg$ .

This is the reason that the theory of section 3 was stated for an arbitrary CUGA, rather than specifically for  $\mathcal{C} = \mathcal{L}ie$ . The reason that our general theory was stated for *simplicial* rather than *differential graded* universal algebras is that there seems to be no reasonable version of Proposition 7.2 for an arbitrary CUGA.

**7.10. Minimal resolutions.** To make the construction more accessible, it is again useful to have *minimal*  $\mathcal{M}_{\mathcal{A}lg}$  resolutions, as in section 5. For this purpose, we consider a variant of the above approach:

Even though  $\mathcal{J}$  does not inherit a closed model category structure from  $d\mathcal{A}$ , one may define *models* for  $\mathcal{J}$ , in the sense of §3.3, by letting a *Jacobi sphere* be any  $d\mathcal{A}$ -model of a  $\mathcal{L}$ -sphere (§5.2), and more generally let  $\mathcal{M}_{\mathcal{J}}$  denote the full subcategory of  $\mathcal{J}$  consisting of

DGNAs weakly equivalent to objects in  $\mathcal{M}_{d\mathcal{L}}$  – i.e., Jacobi models of DGLs which are (up to homotopy) coproducts of  $d\mathcal{L}$ -spheres.

An  $\mathcal{M}_{\mathcal{J}}$  resolution of a DGL  $L$ , which we shall call simply a *Jacobi resolution*, is then defined to be a free simplicial resolution of DGNAs  $A_{\bullet,*} \rightarrow \iota(L)$  (Def. 3.2), with each  $A_{n,*} \in \mathcal{M}_{\mathcal{J}}$ . Note that such an  $A_{\bullet,*} \rightarrow \iota(L)$  is at the same time also an  $\mathcal{M}_{\mathcal{J}}$ -Jacobi resolution of the  $d\mathcal{A}$ -model  $J_*$  of  $L$ , and it is usually more convenient to think of it as such.

There is a comonad  $F : d\mathcal{A} \rightarrow d\mathcal{A}$  as in (5.4), which yields the *canonical Jacobi resolution*  $U_{\bullet,*}$  for any  $C \in \mathcal{J}$ , as in §5.3. Again we may use the notation of §5.6.

One also has an analogue of Propositions 5.8 and 5.13, as follows:

**7.11. Proposition.** *Let  $B = (B_*, \partial_B) \in d\mathcal{L}$  be any DGL, and  $(A_{*,*}, D_A)$  a filtered model for  $B$ : then there is a Jacobi resolution  $J_{\bullet,*} \rightarrow \iota(B)$ , with a bijection  $\theta : X_{**} \xrightarrow{\sim} J_{\bullet,*}$  between a bigraded set  $X_{**}$  of generators for  $A_{*,*}$  and the set of non-degenerate  $d\mathcal{A}$ -spheres in  $J_{\bullet,*}$ .*

*Proof.* Let  $G = G_B$  be a  $d\mathcal{A}$ -model for the DGL  $B$ , and  $U_{\bullet,*} \rightarrow G$  the canonical Jacobi resolution. As in the proof of Proposition 5.8, we may define a map  $\theta : A_{*,*} \rightarrow U_{\bullet,*}$  inductively by the equation  $\theta(x) = \langle \theta(\partial_A(x)) \rangle$  (compare (5.9)), and we shall again write  $x^{(0)}$  for  $\theta(x)$  if  $x \in X_{**}$  (a set of generators for  $A_{*,*}$ ), and let  $V_{**}$  be the bigraded vector space spanned by  $\theta(X_{**})$ . For simplicity of notation we consider first the case where  $B$  has trivial differential and  $A_{*,*}$  is bigraded (with  $D_A = \partial_A$ ).

For each  $n \in \mathbb{N}$ , define the sub-DGNA  $J_{n,*}^{(0)}$  of  $U_{n,*}$  to be  $J(V_{n,*})$ , in the notation of Lemma 7.7 – that is,  $J_{n,*}^{(0)}$  is the coproduct, in  $\mathcal{J}$ , of a set of Jacobi spheres  $\mathbb{S}_{(x^{(0)})}^k$ , one for each generator  $x \in X_{n,k}$  of  $A_{*,*}$ . By Lemma 7.7(c), it is enough to define the face and degeneracy maps of  $J_{\bullet,*}$  on each  $x$  – where we may use the description of §5.6.

Once again, we want  $d_i(x^{(0)})$  to be a  $\partial_U$ -boundary for each  $1 \leq i \leq n$ ; but the analogues of the elements  $x^{(s)}$  of Propositions 5.8 and 5.13 are more complicated, so we need some definitions:

For each  $0 \leq s \leq n$ , let  $\mathcal{K}_{n,s}$  denote the set of all sequences  $I = (i_1, \dots, i_s)$  of integers  $1 \leq i_1 < \dots < i_s \leq n$ , corresponding to the  $s$ -fold face map  $d_I = d_{i_1} \circ \dots \circ d_{i_s} : \mathbf{n} \rightarrow \mathbf{n} - \mathbf{s}$  in  $\Delta^{op}$  (compare Definition 4.3 and the proof of Proposition 5.8). Given  $I = (i_1, \dots, i_s) \in \mathcal{K}_{n,s}$ , for each  $1 \leq j \leq s$  let  $I(\hat{j}) := (i_1, \dots, \widehat{i_j}, \dots, i_s) \in \mathcal{K}_{n,s-1}$  be obtained from  $I$  by omitting the  $j$ -th entry. By repeatedly using the identity

$d_k d_m = d_{m-1} d_k$  ( $k < m$ ), we can find a unique  $\kappa(j) \in \{1, 2, \dots, n\}$  such that  $d_{\kappa(j)} \circ d_{I(j)} = d_I$ .

For each  $x \in X_{n,k}$ ,  $0 \leq s \leq n$ , and  $I \in \mathcal{K}_{n,s}$ , we want to choose an element  $x^{(s;I)} \in J_{n-s,k-n+s}^{(s)} \subset U_{n-s,k-n+s}$  by induction on  $n-s$ , starting with  $x^{(0;\emptyset)} := x^{(0)} = \theta(x)$ , so that:

$$(7.12) \quad \partial_U(x^{(s;I)}) = \sum_{j=1}^s (-1)^j d_{\kappa(j)}(x^{(s-1;I(j))})$$

for  $s \geq 1$ . (The index  $s$  is not really needed, since  $s = |I|$ , but it is useful for keeping the analogy with the notation of (5.11) in mind.)

Note that since  $d_0 \circ \theta = \theta \circ \partial_A$  no longer holds here (because  $d_0$  is a morphism in  $d\mathcal{A}$ , not in  $d\mathcal{L}$ ), it is not generally true that  $d_1(x^{(0)}) = 0$  for all  $x \in X_{**}$ . However, since applying  $H'_*$  still yields a *CW* resolution  $H'_* J_{\bullet,*}^{(0)} \rightarrow H'_* C = H'_* B$ , (where  $C$  is the  $d\mathcal{A}$ -model for the DGL  $B$ ), we know that  $d_i(x^{(0)})$  must be a  $\partial_U$ -boundary for each  $1 \leq i \leq n$ . Thus we can choose an element  $x^{(1;1)} \in U_{n-1,k-n+1}$  with  $\partial_U(x^{(1;1)}) = d_1(x^{(0)})$  (a special case of (7.12)) – and in fact  $x^{(1;1)}$  may be expressed in terms of the “canonical operations”  $\mu_i$  of Lemma 7.7.

Now let  $\partial_A(x) = \sum_t a_t \omega_t[y_{i_1}, \dots, y_{i_{m_t}}]$  for  $y_{i_j} \in A_{n_j,*}$ , so

$$(7.13) \quad \begin{aligned} x^{(0)} &= \left\langle \sum_t a_t \omega_t \llbracket y_{i_1}^{(0)}, \dots, y_{i_{m_t}}^{(0)} \rrbracket \right\rangle \\ &= \left\langle \sum_t a_t \sum_{(J_1, \dots, J_{m_t})} (-1)^{\varepsilon_t} \omega_t[s_{J_1}(y_{i_1}^{(0)}), \dots, s_{J_{m_t}}(y_{i_{m_t}}^{(0)})] \right\rangle \end{aligned}$$

as in (5.10), where the  $(n-n_j)$ -multi-index  $J_j \subseteq \{0, 1, \dots, n-1\}$  is obtained by repeated shuffles, which also determine the sign  $(-1)^{\varepsilon_t}$  (see (2.10) ff.). Therefore,

$$d_k(x^{(0)}) = \left\langle \sum_t a_t \sum_{(I_1, \dots, I_{m_t})} (-1)^{\varepsilon_t} \omega_t[d_{k-1} s_{I_1}(y_{i_1}^{(0)}), \dots, d_{k-1} s_{I_{m_t}}(y_{i_{m_t}}^{(0)})] \right\rangle.$$

The proof of Lemma 2.12 (which is valid in  $d\mathcal{A}$ , too) implies by induction on  $t \geq 2$  that for each summand

$$v_t = \omega_t[s_{J_1}(y_{i_1}^{(0)}), \dots, s_{J_{m_t}}(y_{i_{m_t}}^{(0)})],$$

there is exactly one  $1 \leq j \leq t$  and  $0 \leq \ell \leq k-1$  such that

$$\begin{aligned} d_{k-1}(v_t) &= \omega_t[s_{J'_1}(y_{i_1}^{(0)}), \dots, s_{J'_{j-1}}(y_{i_{j-1}}^{(0)}), \\ &\quad s_{J''_j} d_\ell(y_{i_j}^{(0)}), s_{J'_{j+1}}(y_{i_{j+1}}^{(0)}), \dots, s_{J'_{m_t}}(y_{i_{m_t}}^{(0)})], \end{aligned}$$

for suitable multi-indices  $J'_1, \dots, J'_{j-1}, J'_{j+1}, \dots, J'_{m_t}$  and  $J''_j$ .

Since  $\ell < k$  and  $n_j < n$ , we may assume by induction that we have defined  $y_{i_j}^{(1;\ell)} \in U_{n_j-1,*}$  such that  $\partial_U(y_{i_j}^{(1;\ell)}) = d_\ell(y_{i_j}^{(0)})$ , and then let  $x^{(1;k)}$  be

$$\left\langle \sum_t \sum_{(J_1, \dots, J_{m_t})} (-1)^{\varepsilon t} \omega_t [s_{J'_1}(y_{i_1}^{(0)}), \dots, s_{J'_{j-1}}(y_{i_{j-1}}^{(0)}), \right. \\ \left. s_{J'_j}(y_{i_j}^{(1;\ell)}), s_{J'_{j+1}}(y_{i_{j+1}}^{(0)}), \dots, s_{J'_{m_t}}(y_{i_{m_t}}^{(0)}) \right\rangle.$$

The rest of the construction of the elements  $x^{(s;I)}$ , as well as the generalization to the filtered case, is similar to that in the proofs of Propositions 5.8 and 5.13.  $\square$

We can now summarize the main result of this paper in the following

**7.14. Theorem.** *Let  $\mathbf{X}$  be a simply connected space, and  $\Pi_*^X := \pi_{*-1} \mathbf{X}_{\mathbb{Q}} \in \mathcal{L}_0$  its rational homotopy Lie algebra. There is a tree  $T_X$  of DGLs  $L^{(k, \alpha_1, \dots, \alpha_k)}$ , starting with  $L^{(0)} \simeq (\Pi_*^X, 0)$ , and for each branch  $\alpha_1, \dots, \alpha_k, \dots$  of  $T_X$ , an increasing sequence of positive integers (or  $\infty$ )  $(n_k = n_{k, \alpha_1, \dots, \alpha_k})_{k=1}^\infty$  such that*

- (a)  $H'_*(L^{(k, \alpha_1, \dots, \alpha_k)}) \cong \Pi_*^X$ ;
- (b) The higher homotopy operations  $\langle\langle m \rangle\rangle \subset H_*^m(L^{(k, \alpha_1, \dots, \alpha_k)}; \Pi_*^X)$  associated to a minimal Jacobi resolution of  $L^{(k, \alpha_1, \dots, \alpha_k)}$  as in §4.9, vanish for  $m < n_k$ .
- (c) The operation  $\langle\langle n_k \rangle\rangle = \langle\langle n_{k, \alpha_1, \dots, \alpha_k} \rangle\rangle \subset H_*^{n_k}(L^{(k)}; \Pi_*^X)$  does not vanish (unless  $n_k = \infty$ ).
- (d) For any  $\alpha_{k+1}$  along the branch of  $(\alpha_1, \dots, \alpha_k)$ , the DGL  $L^{(k+1, \alpha_1, \dots, \alpha_k, \alpha_{k+1})}$  may be chosen to agree with  $L^{(k, \alpha_1, \dots, \alpha_k)}$  in degrees  $\leq n_k + 1$ , so the sequential colimit  $L^{(\infty)} = \text{colim}_k L^{(k)}$ , along any branch, is well defined, and is a DGL model for  $\mathbf{X}$ .

The main difference between the construction described here (in  $s\mathcal{J}$ ) and that of Proposition 5.13 is that the “higher order information” in  $J_{\bullet,*}$  is no longer concentrated in the last face map  $d_n : J_{n,*} \rightarrow J_{n-1,*}$ . As a result, the higher homotopy operations associated to the (minimal) Jacobi resolution (as in section 4) are true simplicial operations, which can be translated more directly into topological ones.

## REFERENCES

- [AB] R. Adin & D. Blanc, “Resolutions of Associative and Lie algebras”, *Can. Math. Bull.* **43** (2000) No. 1, pp. 3-16.
- [A1] C. Allday, “Rational Whitehead products and a spectral sequence of Quillen”, *Pac. J. Math.* **46** (1973) No. 2, pp. 313-323.
- [A2] C. Allday, “Rational Whitehead products and a spectral sequence of Quillen, II”, *Houston J. Math.* **3** (1977), No. 3, pp. 301-308.

- [AA] P.G. Andrews & M. Arkowitz, “Sullivan’s minimal models and higher order Whitehead products”, *Can. J. Math.* **30** (1978) No. 5, pp. 961-982.
- [BL] H.J. Baues & J.-M. Lemaire. “Minimal models in homotopy theory”, *Math. Ann.* **225** (1977), pp. 219-242.
- [B11] D. Blanc, “Derived functors of graded algebras”, *J. Pure Appl. Alg.* **64** (1990) No. 3, pp. 239-262.
- [B12] D. Blanc, “Higher homotopy operations and the realizability of homotopy groups”, *Proc. Lond. Math. Soc. (3)* **70** (1995), pp. 214-240.
- [B13] D. Blanc, “Homotopy operations and the obstructions to being an  $H$ -space”, *Manus. Math.* **88** (1995) No. 4, pp. 497-515.
- [B14] D. Blanc, “New model categories from old”, *J. Pure Appl. Math.* **109** (1996) No. 1, pp. 37-60.
- [B15] D. Blanc, “Algebraic invariants for homotopy types”, *Math. Proc. Camb. Phil. Soc.* **27** (1999), No. 3, pp. 497-523.
- [B16] D. Blanc, “CW simplicial resolutions of spaces, with an application to loop spaces”, *Topology & Appl.* **100** (2000), No. 2-3, pp. 151-175.
- [B17] D. Blanc, “Realizing coalgebras over the Steenrod algebra”, *Topology* **40** (2001), No. 5, pp. 993-1016.
- [BDG] D. Blanc, W.G. Dwyer, & P.G. Goerss, “The realization space of a  $\Pi$ -algebra: a moduli problem in algebraic topology”, preprint 2001.
- [BG] D. Blanc & P.G. Goerss, “Realizing simplicial  $\Pi$ -algebras”, preprint 2002.
- [BM] D. Blanc & M. Markl, “Higher homotopy operations”, to appear in *Math. Z.*
- [BS] D. Blanc & C.S. Stover, “A generalized Grothendieck spectral sequence”, in N. Ray & G. Walker, eds., *Adams Memorial Symposium on Algebraic Topology, Vol. 1*, Lond. Math. Soc. Lec. Notes Ser. **175**, Cambridge U. Press, Cambridge, 1992, pp. 145-161.
- [BV] J.M. Boardman & R.M. Vogt, *Homotopy Invariant Algebraic Structures on Topological Spaces*, SpringerVerlag *Lec. Notes Math.* **347**, Berlin-New York, 1973.
- [BF] A.K. Bousfield & E.M. Friedlander, “Homotopy theory of  $\Gamma$ -spaces, spectra, and bisimplicial sets”, in M.G. Barratt & M.E. Mahowald, eds., *Geometric Applications of Homotopy Theory, II*, Springer-Verlag *Lec. Notes Math.* **658**, Berlin-New York, 1978, pp. 80-130.
- [BV] D. Burghilea & M. Vigué-Poirrier, “Cyclic homology of commutative algebras, I”, in Y. Félix, ed., *Algebraic Topology – Rational Homotopy 1986*, Springer-Verlag *Lec. Notes Math.* **1318**, Berlin-New York, 1987, pp. 51-72.
- [Do] A. Dold, “Homology of symmetric products and other functors of complexes”, *Ann. Math. (2)* **68** (1958), pp. 54-80.
- [DKS] W.G. Dwyer, D.M. Kan, & C.R. Stover, “An  $E^2$  model category structure for pointed simplicial spaces”, *J. Pure & Appl. Alg.* **90** (1993) No. 2, pp. 137-152.
- [F1] Y. Félix, *Dènombrement des types de  $k$ -homotopie: théorie de la déformation*, Mem. Soc. Math. France **3**, Paris, 1980.
- [F2] Y. Félix, “Modèles bifiltrés: une plaque tournant en homotopie rationnelle”, *Can. J. Math.* **33** (1981) No. 6, pp. 1448-1458.
- [GHT] A. Gómez-Tato, S. Halperin & D. Tanré, “Rational homotopy theory for non simply connected spaces”, *preprint* 1996.

- [Gd] R. Godement, *Topologie algébrique et théorie des faisceaux*, Act. Sci. & Ind. No. 1252, Publ. Inst. Math. Univ. Strasbourg **XIII**, Hermann, Paris 1964.
- [Go] T.G. Goodwillie, “Cyclic homology, derivations, and the free loop space”, *Topology* **24** (1985) No. 2, pp. 187-216.
- [HS] S. Halperin & J.D. Stasheff, “Obstructions to homotopy equivalences”, *Adv. in Math.* **32** (1979) No. 3, pp. 233-279.
- [Har] Y. Haralambous, “Perturbation d’algèbres de Lie différentielles et coformalité modérée”, *Bull. Soc. Math. Belg. Sér. B* **43** (1991) No. 1, pp. 59-67.
- [J] T. Jozefiak, “Tate resolutions for commutative graded algebras over a local ring”, *Fund. Math.* **74** (1972) No. 3, pp. 209-231.
- [LM] , T.J. Lada & M. Markl, “Strongly homotopy Lie algebras”, *Comm. in Alg.* **23** (1995) No. 6, pp. 2147-2161.
- [LS] J.-M. Lemaire & F. Sigrist, “Dénombrement de types d’homotopie rationnelle”, *C. R. Acad. Sci., Paris, I* **287** (1978) No. 3, pp. 109-112.
- [Lo] J.-L. Loday, *Cyclic Homology*, Springer-Verlag *Grund. math. Wissens.* **301**, Berlin-New York, 1992.
- [Mc1] S. Mac Lane, *Homology*, Springer-Verlag *Grund. math. Wissens.* **114**, Berlin-New York, 1963.
- [Mc2] S. Mac Lane, *Categories for the Working Mathematician*, Springer-Verlag *Grad. Texts Math.* **5**, Berlin-New York, 1971.
- [Ma] J.P. May, *Simplicial Objects in Algebraic Topology*, U. Chicago Press, Chicago-London, 1967.
- [Me] W. Meier, “Minimal models for nonnilpotent spaces”, *Comm. Math. Helv.* **55** (1980), No. 4, pp. 622-633.
- [MN] T.J. Miller & J.A. Neisendorfer. “Formal and coformal spaces”, *Ill. J. Math.* **22** (1978) No. 4, pp. 565-580.
- [N] J.A. Neisendorfer, “Lie algebras, coalgebras and rational homotopy theory for nilpotent spaces”, *Pac. J. Math.* **74** (1978) No. 2, pp. 429-460.
- [O] A. Oukili, *Sur l’homologie d’une algèbre différentielle de Lie*, Ph.D. thesis, Univ. de Nice, 1978.
- [P] G.J. Porter. “Higher order Whitehead products”, *Topology* **3** (1965) pp. 123-165.
- [Q1] D.G. Quillen, *Homotopical Algebra*, Springer-Verlag *Lec. Notes Math.* **20**, Berlin-New York, 1963.
- [Q2] D.G. Quillen, “Spectral sequences of a double semi-simplicial group”, *Topology* **5** (1966), pp. 155-156.
- [Q3] D.G. Quillen, “Rational homotopy theory”, *Ann. Math.* **90** (1969) No. 2, pp. 205-295.
- [Q4] D.G. Quillen, “On the (co-)homology of commutative rings”, *Applications of Categorical Algebra*, Proc. Symp. Pure Math. **17**, AMS, Providence, RI, 1970, pp. 65-87.
- [R1] V.S. Retakh, “Massey operations in Lie superalgebras and differentials of the Quillen spectral sequence”, *Funk. Ann. i Pril.* **12** (1978) No. 4, pp. 91-92 [transl. in *Funct. Anal. & Applic.* **12** (1978) No. 4, pp. 319-321].
- [R2] V.S. Retakh, “Lie-Massey brackets and  $n$ -homotopically multiplicative maps of differential graded Lie algebras”, *J. Pure & Appl. Alg.* **89** (1993) No. 1-2, pp. 217-229.

- [SS] M. Schlessinger & J.D. Stasheff. “Deformation theory and rational homotopy type”, to appear in *Pub. Math. Inst. Hautes Et. Sci.*
- [St] C.R. Stover, “A Van Kampen spectral sequence for higher homotopy groups”, *Topology* **29** (1990), pp. 9-26.
- [T] D. Tanré, *Homotopie Rationnelle: Modèles de Chen, Quillen, Sullivan*, Springer-Verlag *Lec. Notes Math.* **1025**, Berlin-New York, 1983.

DEPT. OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA 31905, ISRAEL  
*E-mail address:* blanc@math.haifa.ac.il