

# ON THE LINEAR STRAND OF AN EDGE IDEAL

MIKE ROTH AND ADAM VAN TUYL

ABSTRACT. Let  $I(G)$  be the edge ideal associated to a simple graph  $G$ . We study the graded Betti numbers that appear in the linear strand of the minimal free resolution of  $I(G)$ .

## INTRODUCTION

Let  $R = k[x_1, \dots, x_n]$  with  $k$  a field. If  $G$  is a simple graph (no loops or multiple edges, but not necessarily connected) with edge set  $E_G$  and vertex set  $V_G = \{x_1, \dots, x_n\}$ , then associated to  $G$  is the monomial ideal  $I(G) := \langle \{x_i x_j \mid \{x_i, x_j\} \in E_G\} \rangle$  in  $R$  called the *edge ideal of  $G$* . We would like to establish a dictionary between algebraic invariants of  $I(G)$  and numerical data associated to the graph  $G$ . Such a dictionary might then allow us to prove graph theoretical results algebraically, and vice versa. Among the many papers that have studied the properties of edge ideals, we mention [7, 8, 9, 12, 13, 16, 17, 18, 19, 20].

In this paper we investigate the relation between the graded Betti numbers arising from the resolution of  $I(G)$  and numerical information associated to  $G$ . Recall that if  $I$  is a homogeneous ideal, then we can associate to  $I$  a minimal graded free resolution of the form

$$0 \rightarrow \bigoplus_j R(-j)^{\beta_{l,j}(I)} \rightarrow \bigoplus_j R(-j)^{\beta_{l-1,j}(I)} \rightarrow \dots \rightarrow \bigoplus_j R(-j)^{\beta_{0,j}(I)} \rightarrow I \rightarrow 0$$

where  $l \leq n$  and  $R(-j)$  is the  $R$ -module obtained by shifting the degrees of  $R$  by  $j$ . The number  $\beta_{i,j}(I)$ , the  $ij$ th graded Betti number of  $I$ , equals the number of generators of degree  $j$  in the  $i$ th syzygy module. Since  $I(G)$  is a square-free monomial ideal, our principal tool to study the numbers  $\beta_{i,j}(I(G))$  will be Hochster's formula (see Propositions 1.1 and 1.2).

Some relations between the numbers  $\beta_{i,j}(I(G))$  and the properties of  $G$  have already been established. For example, from the construction of  $I(G)$  we see that  $\beta_{0,2}(I(G)) = \#E_G$ , and  $\beta_{0,j}(I(G)) = 0$  for  $j \neq 2$ . Formulas for the Betti numbers of the first syzygy module are given in [3]. In [13] it is shown that  $\beta_{i,2(i+1)}(I(G))$  with  $i \geq 1$  is the number of induced subgraphs of  $G$  consisting of exactly  $i + 1$  disjoint edges. This result, and many other such relations between  $\beta_{i,j}(I(G))$  and  $G$ , can be found in the thesis of Jacques [12].

In this paper we shall focus on the graded Betti numbers of  $I(G)$  that appear in the linear strand. When  $I$  is generated by elements of degree at least  $d$ , then it can be shown that  $\beta_{i,j}(I) = 0$  for all  $j < i + d$ . Thus, the first graded Betti number of

2000 *Mathematics Subject Classification.* 13D40, 13D02, 05C90.

*Key words and phrases.* graphs, resolutions, Betti numbers.

Version: November 8, 2004.

interest is  $\beta_{i,i+d}(I)$  for each  $1 \leq i \leq l$ . The numbers  $\beta_{i,i+d}(I)$  for  $i \geq 0$  describe the *linear strand* of the minimal resolution of  $I$ , i.e., they count the number of linear syzygies appearing in the resolution. It is then of interest to understand what the linear strand can tell us about  $I$  (cf. [4, 5, 15]). Because  $I(G)$  is generated by monomials of degree 2, we will be interested in the numbers  $\beta_{i,i+2}(I(G))$  for  $i \geq 0$ .

The contributions of this paper are as follows. If  $G$  has no induced 4-cycles, then we provide an exact formula for  $\beta_{i,i+2}(I(G))$  in terms of data associated to  $G$ . In particular, our result applies to all forests. As well, we give formulas for  $\beta_{i,i+2}(I(G))$  for  $i = 0, \dots, 3$  for any simple graph  $G$ . The formulas for  $i = 0, 1$  were known, but our proofs are new. When  $G$  has an induced 4-cycle, we give upper and lower bounds for  $\beta_{i,i+2}(I(G))$  based upon the numerical information associated to the graph  $G$ . Using growth bounds on the Betti numbers of lexicographical ideals, we also obtain a purely graph theoretical result by providing a lower bound on the number of triangles in  $G$ .

## 1. THE GRADED BETTI NUMBERS OF AN EDGE IDEAL

The main result of this section is to express  $\beta_{i,j}(I(G))$  in terms of the dimensions of the reduced simplicial homology groups of a simplicial complex constructed from the graph  $G$ .

**1.1. Graphs and simplicial complexes.** Let  $G$  denote a graph with vertex set  $V_G = \{x_1, \dots, x_n\}$  and edge set  $E_G$ . We shall say  $G$  is *simple* if  $G$  has no loops or multiple edges. A simple graph need not be connected. The *degree* of a vertex  $x_i \in V_G$ , denoted  $\deg x_i$ , is the number of edges incident to  $x_i$ .

If  $S \subseteq V_G$ , then the *induced subgraph* of  $G$  on the vertex set  $S$ , denoted  $G_S$ , is the subgraph of  $G$  such that every edge of  $G$  contained in  $S$  is an edge of  $G_S$ . The *complement* of a graph  $G$ , denoted by  $G^c$ , is the graph whose vertex set is the same as  $G$ , but whose edge set is defined by the rule:  $\{x_i, x_j\} \in E_{G^c}$  if and only if  $\{x_i, x_j\} \notin E_G$ . We shall let  $\#\text{comp}(G)$  denote the number of connected components of  $G$ . Furthermore, we let  $\iota(G)$  denote the number of isolated vertices of  $G$ . It is then clear that  $\#\text{comp}(G) \geq \iota(G)$ .

The *cycle* on  $n$  vertices, denoted  $C_n$ , is the graph with  $V_G = \{x_1, \dots, x_n\}$  and  $E_G = \{\{x_1, x_2\}, \{x_2, x_3\}, \dots, \{x_n, x_1\}\}$ . The *wheel*  $W_n$  is the graph obtained by adding a vertex  $z$  to  $C_n$  and then adjoining an edge between  $z$  and every vertex  $x_i \in V_{C_n}$ . Note that  $W_n$  has  $n + 1$  vertices. The *complete graph* on  $n$  vertices, denoted  $K_n$ , is the graph with the property that for  $x_i \neq x_j \in V_{K_n}$ , the edge  $\{x_i, x_j\} \in E_{K_n}$ . The *complete bipartite graph*, denoted  $K_{n,m}$ , is the graph with vertex set  $V_G = \{x_1, \dots, x_n, y_1, \dots, y_m\}$  and edge set  $E_G = \{\{x_i, y_j\} \mid 1 \leq i \leq n, 1 \leq j \leq m\}$ . We shall write  $c_n(G)$ ,  $w_n(G)$ ,  $k_n(G)$ , and  $k_{a,b}(G)$  for the number of induced subgraphs of  $G$  isomorphic to  $C_n$ ,  $W_n$ ,  $K_n$ , and  $K_{a,b}$ , respectively.

A *simplicial complex*  $\Delta$  on a set of vertices  $V = \{x_1, \dots, x_n\}$  is a collection of subsets of  $V$  such that  $\{x_i\} \in \Delta$  for  $i = 1, \dots, n$ , and for each  $F \in \Delta$ , if  $G \subseteq F$ , then  $G \in \Delta$ . Note that  $\emptyset \in \Delta$ . Associated to  $\Delta$  is a square-free monomial ideal  $I_\Delta$  in  $R = k[x_1, \dots, x_n]$ . More precisely, the ideal

$$I_\Delta = \langle \{x_{i_1}x_{i_2} \cdots x_{i_t} \mid \{x_{i_1}, x_{i_2}, \dots, x_{i_t}\} \notin \Delta \} \rangle$$

is the *Stanley-Reisner ideal* of  $\Delta$ , and  $I_\Delta$  is generated by the square-free monomials that correspond to non-faces of  $\Delta$ .

If  $G$  is any graph, the *clique complex* is the simplicial complex  $\Delta(G)$  where  $F = \{x_{i_1}, \dots, x_{i_j}\} \in \Delta(G)$  if and only if  $G_F$  is a complete graph.

If  $G$  is a simple graph with edge ideal  $I(G)$ , then the ideal  $I(G)$  is generated by square-free monomial ideals. So  $I(G)$  is also the defining ideal of a simplicial complex  $\Delta$ , that is,  $I(G) = I_\Delta$ . Specifically,  $I(G)$  is the Stanley-Reisner ideal associated to the clique complex  $\Delta = \Delta(G^c)$ .

**1.2. Graded Betti numbers of edge ideals.** Let  $I$  be a homogeneous ideal of  $R$ . The  $ij$ th graded Betti number of  $I$ , denoted  $\beta_{i,j}(I)$ , is given by

$$\beta_{i,j}(I) = \dim_k \operatorname{Tor}_i^R(I, k)_j.$$

If  $I$  is also a monomial ideal, then  $I$  also has an  $\mathbb{N}^n$ -graded minimal free resolution. As a consequence  $\operatorname{Tor}_i^R(I, k)$  is also  $\mathbb{N}^n$ -graded, and we have

$$\beta_{i,\alpha}(I) = \dim_k \operatorname{Tor}_i^R(I, k)_\alpha \text{ with } \alpha \in \mathbb{N}^n.$$

When  $I = I_\Delta$  is the Stanley-Reisner ideal of a simplicial complex, Hochster [10] described the  $\mathbb{N}^n$ -graded Betti numbers of  $I_\Delta$  in terms of the dimensions of the reduced homology of the simplicial complex  $\Delta$ . To state the result, we require the following notation: if  $m$  is a monomial of  $R = k[x_1, \dots, x_n]$  with  $\deg m = \alpha \in \mathbb{N}^n$ , then we define

$$\operatorname{Tor}_i^R(I_\Delta, k)_m := \operatorname{Tor}_i^R(I_\Delta, k)_\alpha$$

**Proposition 1.1** (Hochster's Formula). *Let  $\Delta$  be a simplicial complex on vertex set  $\{x_1, \dots, x_n\}$  and let  $m$  be a monomial of  $R$ . If  $m$  is square-free, then*

$$\dim_k \operatorname{Tor}_i^R(I_\Delta, k)_m = \dim_k \tilde{H}_{\deg(m)-i-2}(|m|, k)$$

where  $\tilde{H}_j(|m|, k)$  denotes the  $j$ th reduced homology of the full subcomplex  $|m|$  of  $\Delta$  whose vertices correspond to the variables dividing  $m$ . If  $m$  is not square-free, then  $\operatorname{Tor}_i^R(I_\Delta, k)_m$  vanishes.

By specializing this result to edge ideals, we obtain:

**Proposition 1.2.** *Let  $G$  be a simple graph with edge ideal  $I(G)$ . Then*

$$\beta_{i,j}(I(G)) = \sum_{S \subseteq V_G, |S|=j} \dim_k \tilde{H}_{j-i-2}(\Delta(G_S^c), k) \text{ for all } i, j \geq 0.$$

*Proof.* We sketch out the main idea of the proof. Let  $\Delta = \Delta(G^c)$  be the simplicial complex defined by  $I(G)$ . It follows from Proposition 1.1 that

$$\beta_{i,j}(I(G)) = \beta_{i,j}(I_\Delta) = \sum_{m \in M_j, m \text{ is square-free}} \dim_k \tilde{H}_{j-i-2}(|m|, k)$$

where  $M_j$  consists of all the monomials of degree  $j$  in  $R$ . Since  $\deg m = j$  and  $m$  is square-free, the variables that divide  $m$  give a subset  $S \subseteq V_G$  of size  $j$ . Let  $G_S$  denote the induced subgraph of  $G$  on this vertex set  $S$ , and let  $G_S^c$  denote its complement. To finish the proof, it is enough to note that the full subcomplex  $|m|$  of  $\Delta(G^c)$  and  $\Delta(G_S^c)$  are the same simplicial complex.  $\square$

*Remark 1.3.* A similar formula for  $\beta_{i,j}(I(G))$  can be found in [13]. While Proposition 1.2 gives a formula for all the numbers  $\beta_{i,j}(I(G))$ , the formula appears difficult to apply since one has to compute the dimensions of all the homology groups  $\tilde{H}_{j-i-2}(\Delta(G_S^c), k)$  as  $S$  varies over all subsets of  $V_G$  of size  $j$ . Moreover, it is not clear how this formula relates to data associated to  $G$ .

## 2. THE BETTI NUMBERS IN THE LINEAR STRAND OF AN EDGE IDEAL

In this section we shall give some exact formulas for the graded Betti numbers in the linear strand of  $I(G)$  when  $G$  is a simple graph.

**Proposition 2.1.** *Let  $G$  be a simple graph with edge ideal  $I(G)$ . Then*

$$\beta_{i,i+2}(I(G)) = \sum_{S \subseteq V_G, |S|=i+2} (\#\text{comp}(G_S^c) - 1) \text{ for all } i \geq 0.$$

*Proof.* By Proposition 1.2

$$\beta_{i,i+2}(I(G)) = \sum_{S \subseteq V_G, |S|=i+2} \dim_k \tilde{H}_0(\Delta(G_S^c), k).$$

Now  $\dim_k \tilde{H}_0(\Delta(G_S^c), k) = \#\text{comp}(\Delta(G_S^c)) - 1$ . It now suffices to note that the simplicial complex  $\Delta(G_S^c)$  and  $G_S^c$  have the same number of components.  $\square$

*Remark 2.2.* From this reformulation it is clear the numbers  $\beta_{i,i+2}(I(G))$  do not depend upon the characteristic of the field  $k$ . See Katzman [13] for further results on the dependence of  $\beta_{i,j}(I(G))$  upon the characteristic of  $k$ .

*Remark 2.3.* It will sometimes be convenient to write  $\beta_{i,i+2}(I(G))$  as

$$(1) \quad \beta_{i,i+2}(I(G)) = \sum_{\substack{S \subseteq V_G, \\ |S|=i+2, \\ G_S^c \text{ contains an} \\ \text{isolated vertex}}} (\#\text{comp}(G_S^c) - 1) + \sum_{\substack{S \subseteq V_G, \\ |S|=i+2, \\ G_S^c \text{ contains no} \\ \text{isolated vertices}}} (\#\text{comp}(G_S^c) - 1).$$

Although the formula in Proposition 2.1 is defined strictly in terms of the graph  $G$ , the formula has the disadvantage that one must sum over all  $S \subseteq V_G$  of size  $i+2$ . A further reduction can be made if we impose an extra hypothesis on  $G$ . We say that  $G$  has no induced 4-cycle if for every  $S \subseteq V_G$  with  $|S|=4$ , then  $G_S \not\cong C_4$ .

**Proposition 2.4.** *Let  $G$  be a simple graph with edge ideal  $I(G)$ . If  $G$  has no induced 4-cycles, then*

$$\beta_{i,i+2}(I(G)) = \sum_{v \in V_G} \binom{\deg v}{i+1} - k_{i+2}(G) \text{ for all } i \geq 0.$$

*Furthermore, the above formula holds for all simple graphs  $G$  if  $i = 0$  or  $1$ .*

*Proof.* The hypothesis that  $G$  contains no induced 4-cycles is the same as saying that for any subset  $S$  of the vertices, at most one connected component of  $G_S^c$  is larger than a vertex. That is, if  $G_S^c$  has two connected components, each containing an edge, then letting  $S'$  be the set of the four vertices in these two edges we see that  $G_{S'}$  is an induced 4-cycle of  $G_S$ . Therefore, to count  $\#\text{comp}(G_S^c) - 1$  we can simply count the number of isolated vertices in  $G_S^c$ , the “ $-1$ ” term being taken care

of by the component which isn't a vertex; we thus over count by one whenever  $G_S^c$  consists completely of isolated vertices.

For any vertex  $v$ , the number of subsets  $S$  of size  $i + 2$  containing  $v$  such that  $v$  is an isolated vertex in  $G_S^c$  is  $\binom{\deg v}{i+1}$ . To take care of the over count, note that  $G_S^c$  consists of isolated vertices exactly when  $G_S$  is a complete graph on  $i + 2$  vertices. Subtracting the number of times this happens gives the formula above.

Now let  $G$  be any simple graph. To compute  $\beta_{i,i+2}(I(G))$  when  $i = 0$  or  $1$ , we need to count the number of connected components of  $G_S^c$  when  $|S| = 2$  or  $3$ . But for any simple graph on two or three vertices, at most one connected component can be larger than a vertex. The proof is now the same as the one given above.  $\square$

*Remark 2.5.* It follows from the construction of  $I(G)$  that  $\beta_{0,2}(I(G)) = \#E_G$ . Coupling this result with the above formula gives the identity

$$\#E_G = \sum_{v \in V_G} \deg v - k_2(G).$$

But  $k_2(G) = \#E_G$  since each edge is isomorphic to  $K_2$ . We thus recover the well known Degree-Sum Formula which states that  $2\#E_G = \sum_{v \in V_G} \deg v$ . When  $i = 1$ , we have the formula  $\beta_{1,3} = \sum_{v \in V_G} \binom{\deg v}{2} - k_3(G)$ . Since  $k_3(G)$  is simply the number of triangles in  $G$ , we recover the formula of [3].

Since forests do not have cycles, we get an immediate corollary.

**Corollary 2.6.** *Let  $G$  be a forest with edge ideal  $I(G)$ . Then  $\beta_{0,2}(I(G)) = \#E_G$ , and*

$$\beta_{i,i+2}(I(G)) = \sum_{v \in V_G} \binom{\deg v}{i+1} \text{ for all } i \geq 1.$$


*Remark 2.7.* Zheng [20] studied the linear strand of a special class of simplicial complexes that generalized the notion of a tree (due to Faridi [6]). Our formula above agrees with [20, Proposition 3.3] in the case that the simplicial complex is a connected tree in the standard sense.

We now provide formulas for  $\beta_{2,4}(I(G))$  and  $\beta_{3,4}(I(G))$  for any simple graph  $G$ .

**Proposition 2.8.** *Let  $G$  be any simple graph with edge ideal  $I(G)$ . Then*

$$\begin{aligned} \beta_{2,4}(I(G)) &= \sum_{v \in V_G} \binom{\deg v}{3} - k_4(G) + k_{2,2}(G) \text{ and} \\ \beta_{3,5}(I(G)) &= \sum_{v \in V_G} \binom{\deg v}{4} - k_5 + k_{2,3}(G) + w_4(G) + d(G) \end{aligned}$$

where  $d(G)$  is the number of induced subgraphs of  $G$  isomorphic to the graph  $D$  given below:



*Proof.* Suppose  $i = 2$ . We will compute both sums in (1) to find the value of  $\beta_{2,4}(I(G))$ . Suppose  $S \subseteq V_G$  is such that  $G_S^c$  has an isolated vertex. The remaining

three vertices of  $G_S^c$  must form a simple graph. By considering all possible simple graphs on three vertices, we will have

$$\# \text{comp}(G_S^c) - 1 = \iota(G_S^c)$$

except when  $G_S^c$  is the graph of 4 isolated vertices, that is,  $G_S \cong K_4$ , in which case we have

$$\# \text{comp}(G_S^c) - 1 = \iota(G_S^c) - 1.$$

The first sum in the expression for  $\beta_{2,4}(I(G))$  therefore equals

$$\sum_{\substack{S \subseteq V_G, |S|=4, \\ G_S^c \text{ contains an isolated vertex}}} \iota(G_S^c) - \sum_{G_S \cong K_4} 1 = \sum_{v \in V_G} \binom{\deg v}{3} - k_4(G).$$

On the other hand, suppose  $S \subseteq V_G$  is such that  $G_S^c$  has no isolated vertex. We can further assume that  $G_S^c$  is disconnected, because if  $G_S^c$  is connected, no contribution is made to  $\beta_{2,4}(I(G))$  for this subset. But then the only possibility for  $G_S^c$  is two disjoint edges, or equivalently,  $G_S \cong K_{2,2}$ . So the second sum in (1) must equal  $k_{2,2}(G)$ , thus giving the desired formula for  $\beta_{2,4}(I(G))$ .

Suppose now that  $i = 3$ . If  $S \subseteq V_G$  is such that  $|S| = 5$  and  $G_S^c$  has an isolated vertex, then the remaining four vertices form a simple graph. By considering all possible simple graphs on four vertices, we have

$$\# \text{comp}(G_S^c) - 1 = \begin{cases} \iota(G_S^c) + 1 & \text{if } G_S \cong W_4 \\ \iota(G_S^c) - 1 & \text{if } G_S \cong K_5 \\ \iota(G_S^c) & \text{otherwise} \end{cases}.$$

Thus

$$\sum_{\substack{S \subseteq V_G, |S|=5, \\ G_S^c \text{ contains an isolated vertex}}} (\# \text{comp}(G_S^c) - 1) = \sum_{v \in V_G} \binom{\deg v}{4} - k_5(G) + w_4(G).$$

Now suppose  $S$  is such that  $|S| = 5$  but  $G_S^c$  has no isolated vertices. Since such an  $S$  will only contribute to  $\beta_{3,5}(I(G))$  if  $G_S^c$  is disconnected, we can further assume that  $S$  is such that  $G_S^c$  is disconnected. By consider all simple graphs on 5 vertices, we find there are only two possibilities for  $S$ : either  $G_S \cong K_{2,3}$ , or  $G_S$  is isomorphic to the graph  $D$ . In both cases,  $\# \text{comp}(G_S^c) - 1 = 1$ . Hence

$$\sum_{\substack{S \subseteq V_G, |S|=5, \\ G_S^c \text{ contains no isolated vertices}}} (\# \text{comp}(G_S^c) - 1) = k_{2,3}(G) + d(G).$$

If we substitute these values into (1), we get the desired formula for  $\beta_{3,5}(I(G))$ .  $\square$

For any homogeneous ideal  $I \subseteq R$  generated in degree  $d$ , we say  $I$  has a *linear resolution* if  $\beta_{i,j}(I) = 0$  for any  $j \neq i + d$ . In other words, the only non-zero graded Betti numbers of  $I$  are those in the linear strand. When  $I = I(G)$  for some simple graph  $G$ , Fröberg [7] gave a characterization of those ideals having a linear resolution in terms of chordal graphs. We say  $G$  is a *chordal graph* if every cycle of length  $n > 3$  has a chord. Here, if  $\{x_1, x_2\}, \dots, \{x_n, x_1\}$  are the  $n$  edges of a cycle of length  $n$ , we say the cycle has a chord in  $G$  if there exists vertices  $x_i, x_j$  such that  $\{x_i, x_j\}$  is also an edge of  $G$ , but  $j \not\equiv i + 1 \pmod{n}$ .

**Proposition 2.9.** *Let  $G$  be a graph with edge ideal  $I(G)$ . The edge ideal  $I(G)$  has a linear resolution if and only if  $G^c$  is a chordal graph.*

Using this criterion and the results of this section we can completely describe the graded minimal free resolutions of  $I(G)$  when  $G = K_n$  or  $K_{a,b}$ . These results can also be found in [12].

**Example 2.10.** (The resolution of  $I(K_n)$ ) The complement of  $K_n$  is simply  $n$  isolated vertices, and hence,  $I(K_n)$  must have a linear resolution by Proposition 2.9. Moreover, for all  $n \geq 3$ , no induced subgraph of  $K_n$  will be a 4-cycle. Proposition 2.4 then gives  $\beta_{i,i+2}(I(K_n)) = \sum_{v \in V_{K_n}} \binom{\deg v}{i+1} - k_{i+2}(K_n)$  for all  $i \geq 0$ . Each vertex of  $K_n$  has degree  $n-1$ , and  $K_n$  has  $\binom{n}{i+2}$  subgraphs isomorphic to  $K_{i+2}$ . The above expression therefore reduces to

$$\beta_{i,i+2}(I(K_n)) = (i+1) \binom{n}{i+2} \text{ for all } i \geq 0.$$

**Example 2.11.** (The resolution of  $I(K_{a,b})$ ) Let  $G = K_{a,b}$  be a complete bipartite graph. We write the vertex set of  $G$  as  $V_G = \{x_1, \dots, x_a, y_1, \dots, y_b\}$  so that  $E_G = \{\{x_i, y_j\} \mid 1 \leq i \leq a, 1 \leq j \leq b\}$ . For all  $a, b \geq 1$ , the complement of  $G$  is the disjoint union of  $K_a$  and  $K_b$ . Since this graph has no induced cycle of length  $\geq 4$ , the resolution of  $I(G)$  is linear by Proposition 2.9.

Because  $G^c = K_a \cup K_b$ , for any  $S \subseteq V_G$  with  $|S| = i+2$ , we have

$$\#\text{comp}(G_S^c) = \begin{cases} 2 & \text{if } S \cap \{x_1, \dots, x_a\} \neq \emptyset \text{ and } S \cap \{y_1, \dots, y_b\} \neq \emptyset \\ 1 & \text{otherwise} \end{cases}.$$

By Proposition 2.1, to determine  $\beta_{i,i+2}(I(G))$  it therefore suffices to count the number of subsets  $S \subseteq V_G$  with  $|S| = i+2$  and  $\#\text{comp}(G_S^c) = 2$ .

There are  $\binom{a+b}{i+2}$  subsets of  $V_G$  that contain  $i+2$  distinct vertices. Furthermore,  $\binom{a}{i+2}$  of these subsets must contain only vertices among  $\{x_1, \dots, x_a\}$ ; similarly,  $\binom{b}{i+2}$  of these subsets contain only vertices among  $\{y_1, \dots, y_b\}$ . It thus follows that

$$\beta_{i,i+2}(I(K_{a,b})) = \binom{a+b}{i+2} - \binom{a}{i+2} - \binom{b}{i+2} \text{ for all } i \geq 0$$

since the expression on the right hand side counts the number of subsets  $S \subseteq V_G$  with  $|S| = i+2$  and  $S$  contains at least one  $x_i$  and one  $y_j$  vertex.

### 3. BOUNDS ON BETTI NUMBERS IN THE LINEAR STRAND

In this section, we give bounds on  $\beta_{i,i+2}(I(G))$  when  $G$  has an induced 4-cycle.

**Proposition 3.1.** *Let  $G$  be a simple graph with edge ideal  $I(G)$ . Suppose that  $G$  has an induced 4-cycle. Then for all  $i \geq 2$*

$$\beta_{i,i+2}(I(G)) \geq \sum_{v \in V_G} \binom{\deg v}{i+1} - k_{i+2}(G) + k_{2,i}(G) + k_{3,i-1}(G) + \dots + k_{\lfloor \frac{i+2}{2} \rfloor, \lceil \frac{i+2}{2} \rceil}(G).$$

*Proof.* We will use the formula for  $\beta_{i,i+2}(I(G))$  as given in equation (1). Suppose that  $S \subseteq V_G$  is such that  $G_S^c$  contains an isolated vertex. If  $G_S^c$  consists entirely of isolated vertices, i.e.,  $G_S \cong K_{i+2}$ , then

$$\#\text{comp}(G_S^c) - 1 = \iota(G_S^c) - 1.$$

On the hand, if  $G_S^c$  has at least one component consisting of an edge, then

$$\# \text{comp}(G_S^c) - 1 \geq \iota(G_S^c)$$

since the “-1” takes care of the component containing the edge. It thus follows that

$$\sum_{\substack{S \subseteq V_G, \\ |S| = i+2, \\ G_S^c \text{ contains an} \\ \text{isolated vertex}}} (\# \text{comp}(G_S^c) - 1) \geq \sum_{\substack{S \subseteq V_G, \\ |S| = i+2, \\ G_S^c \text{ contains an} \\ \text{isolated vertex}}} \iota(G_S^c) - \sum_{\substack{S \subseteq V_G \\ G_S \cong K_{i+2}}} 1.$$

For any vertex  $v$ , the number of subsets  $S$  of size  $i+2$  containing  $v$  such that  $v$  is an isolated vertex in  $G_S^c$  is  $\binom{\deg v}{i+1}$ . We thus obtain the bound

$$(2) \quad \sum_{\substack{S \subseteq V_G, |S| = i+2, \\ G_S^c \text{ contains an isolated vertex}}} (\# \text{comp}(G_S^c) - 1) \geq \sum_{v \in V_G} \binom{\deg v}{i+1} - k_{i+2}(G).$$

Suppose that  $S \subseteq V_G$  is such that  $G_S^c$  contains no isolated vertices. If  $G_S^c$  is connected, then this subset  $S$  does not contribute to the value of  $\beta_{i,i+2}(I(G))$  because  $\# \text{comp}(G_S^c) - 1 = 0$ . So, we can assume that  $S$  is such that  $G_S^c$  has at least two connected components.

Now if  $S$  is such that  $|S| = i+2$  and  $G_S \cong K_{a,i+2-a}$  for some  $a = 2, \dots, \lfloor \frac{i+2}{2} \rfloor$ , then  $G_S^c$  has exactly two components, and no component is an isolated vertex. So, each such  $S$  contributes exactly 1 to the value of  $\beta_{i,i+2}(I(G))$ , and hence

$$(3) \quad \sum_{\substack{S \subseteq V_G, |S| = i+2, \\ G_S^c \text{ contains no isolated vertices}}} (\# \text{comp}(G_S^c) - 1) \geq k_{2,i}(G) + \dots + k_{\lfloor \frac{i+2}{2} \rfloor, \lceil \frac{i+2}{2} \rceil}(G).$$

By now combining the bounds (2) and (3), we get the desired conclusion.  $\square$

Recall that the elements of  $\mathbb{N}^n$  can be given a total ordering using the lexicographical order defined by  $(a_1, \dots, a_n) > (b_1, \dots, b_n)$  if  $a_1 = b_1, \dots, a_{i-1} = b_{i-1}$  but  $a_i > b_i$ . This induces an ordering on the monomials of  $R$ :  $x_1^{a_1} \dots x_n^{a_n} >_{\text{lex}} x_1^{b_1} \dots x_n^{b_n}$  if  $(a_1, \dots, a_n) > (b_1, \dots, b_n)$ . A monomial ideal  $I$  is a lexicographical ideal if for each  $d \in \mathbb{N}$ , a basis for  $I_d$  is the  $\dim_k I_d$  largest monomials of degree  $d$  with respect to the lexicographical ordering.

**Proposition 3.2.** *Let  $G$  be a simple graph and let  $\{m_1, \dots, m_{\#E_G}\}$  be the  $\#E_G$  largest monomials of degree 2 in  $R$  with respect to the lexicographical ordering. Then*

$$\beta_{i,i+2}(I(G)) \leq \sum_{t=1}^{\#E_G} \binom{u_t - 1}{i}$$

where  $u_t$  is the largest index of a variable dividing  $m_t$ .

*Proof.* Let  $L$  be the lexicographical monomial ideal with the property that  $H_{R/L}$ , the Hilbert function of  $R/L$ , is the same as the Hilbert function of  $R/I(G)$ . By the Bigatti-Hulett-Pardue Theorem (see [1, 11, 14]), we have

$$\beta_{i,j}(I(G)) \leq \beta_{i,j}(L) \text{ for all } i, j \geq 0.$$



In particular,  $\beta_{i,i+2}(I(G)) \leq \beta_{i,i+2}(L)$  for all  $i \geq 0$ . Since  $L$  is generated in degrees  $\geq 2$ , it follows that  $\beta_{i,i+2}(L) = \beta_{i,i+2}(L_2)$  for all  $i \geq 0$  where  $L_2$  is the ideal generated by all the degree 2 monomials in  $L$ . Moreover, since  $H_{R/I(G)}(2) = H_{R/L}(2)$ ,  $L$  has  $\#E_G$  monomials of degree 2. Thus  $L_2 = (m_1, \dots, m_{\#E_G})$  where the  $m_i$ s are the  $\#E_G$  largest monomials of degree 2 with respect to the lexicographical ordering. Note that  $L_2$  is also a lexicographical ideal.

To get the desired conclusion, we use the Eliahou-Kervaire resolution (see [2]) of Borel-fixed ideals (a class of ideals which includes the lexicographical ideals) to compute  $\beta_{i,i+2}(L_2)$ :

$$\beta_{i,i+2}(L_2) = \sum_{t=1}^{\#E_G} \binom{u_t - 1}{i} \text{ for all } i \geq 0$$

where  $u_t$  is the largest index of a variable dividing  $m_t$ .  $\square$

*Remark 3.3.* We can give an explicit description of the  $\#E_G$  largest monomials of degree 2. Let  $j$  be the integer such that

$$n + (n-1) + \dots + n - (j-1) < \#E_G \leq n + (n-1) + \dots + n - j,$$

and let  $l = \#E_G - \sum_{i=1}^{j-1} (n-1)$ . (If  $j = 0$ , then we take  $\sum_{i=1}^{j-1} (n-1) = 0$ .) Then the  $\#E_G$  largest monomials of degree 2 are

$$\{x_1^2, x_1x_2, \dots, x_1x_n, x_2^2, x_2x_3, \dots, x_2x_n, x_3^2, \dots, x_{j+1}^2, \dots, x_{j+1}x_{j+l}\}.$$

Notice that since we are only interested in the indices, which only depend upon  $\#E_G$  and  $\#V_G$ , Proposition 3.2 can be written without reference to monomials.

As a corollary, we derive a lower bound for the number of triangles in a graph.

**Corollary 3.4.** *Let  $G$  be a simple graph, and let  $j$  and  $l$  be defined from  $\#E_G$  as above. Then*

$$\max \left\{ 0, \sum_{v \in V_G} \binom{\deg v}{2} - j \binom{\#V_G}{2} + \binom{j}{j-3} - (j + \dots + (j+l-1)) \right\} \leq k_3(G).$$

*Proof.* Let  $\{x_1^2, x_1x_2, \dots, x_{j+1}^2, \dots, x_{j+1}x_{j+l}\}$  be the  $\#E_G$  largest monomials of degree 2 with respect to lexicographical ordering. By combining Propositions 2.4 and 3.2 we get

$$\beta_{1,3}(I(G)) = \sum_{v \in V_G} \binom{\deg v}{2} - k_3(G) \leq \sum_{t=1}^{\#E_G} u_t - 1$$

where  $u_t$  is the largest index of a variable dividing  $m_i$ . From our description of the monomials, it is now a calculation to check that

$$\sum_{t=1}^{\#E_G} u_t - 1 = j \binom{\#V_G}{2} - \binom{j}{j-3} + (j + \dots + (j+l-1)).$$

(We have used the fact the  $n = \#V_G$ .) After substituting this result into our inequality for  $\beta_{1,3}(I(G))$  and rearranging, we get desired bound.  $\square$

*Acknowledgments.* We would like to thank Huy Tài Hà and David Gregory for their comments and some fruitful discussions. Both authors would like to acknowledge the financial support of NSERC while working on this project. The second author

also would like to thank Queen's University and the Università di Genova for their hospitality while working on this project.

## REFERENCES

- [1] A.M. Bigatti, Upper bounds for the Betti numbers of a given Hilbert function. *Comm. Algebra* **21** (1993) 2317–2334.
- [2] S. Eliahou, M. Kervaire, Minimal resolutions of some monomial ideals. *J. Algebra* **129** (1990) 1–25.
- [3] S. Eliahou, R.H. Villarreal, The second Betti number of an edge ideal. XXXI National Congress of the Mexican Mathematical Society (Hermosillo, 1998), 115–119, *Aportaciones Mat. Comun.*, **25**, Soc. Mat. Mexicana, México, 1999.
- [4] D. Eisenbud, M. Green, K. Hulek, S. Popescu, Restricting linear syzygies: algebra and geometry. (2004) Preprint. [math.AG/0404516](#)
- [5] D. Eisenbud, S. Goto, Linear free resolutions and minimal multiplicity. *J. Algebra* **88** (1984) 89–133.
- [6] S. Faridi, The facet ideal of a simplicial complex. *manuscripta math.* **109** (2002) 159–174.
- [7] R. Fröberg, On Stanley-Reisner rings. In: *Topics in algebra*, Banarch Center Publications, **26** (2) (1990) 57–70.
- [8] J. Herzog, T. Hibi, X. Zheng, Cohen-Macaulay chordal graphs. (2004) Preprint. [math.AC/0407375](#)
- [9] J. Herzog, T. Hibi, X. Zheng, Dirac's theorem on chordal graphs and Alexander duality. (2003) Preprint. [math.AC/0307224](#)
- [10] M. Hochster, Cohen-Macaulay rings, combinatorics, and simplicial complexes. *Ring theory, II* (Proc. Second Conf., Univ. Oklahoma, Norman, Okla., 1975), pp. 171–223. *Lecture Notes in Pure and Appl. Math.*, **26**, Dekker, New York, 1977.
- [11] H. Hulett, Maximum Betti numbers of homogeneous ideals with a given Hilbert function. *Comm. Algebra* **21** (1993) 2335–2350.
- [12] S. Jacques, Betti numbers of graph ideals. Ph.D. Thesis, University of Sheffield, 2004. [math.AC/0410107](#)
- [13] M. Katzman, Characteristic-independence of Betti numbers of graph ideals. (2004) Preprint. [math.AC/0408016](#)
- [14] K. Pardue, Deformation classes of graded modules and maximal Betti numbers. *Illinois J. Math.* **40** (1996) 564–585.
- [15] V. Reiner, V. Welker, Linear syzygies of Stanley-Reisner ideals. *Math. Scand.* **89** (2001) 117–132.
- [16] A. Simis, W.V. Vasconcelos, R.H. Villarreal, On the ideal theory of graphs. *J. Algebra* **167** (1994) 389–416.
- [17] R. H. Villarreal, Rees algebras of edge ideals. *Comm. Algebra* **23** (1995) 3513–3524.
- [18] R. H. Villarreal, Cohen-Macaulay graphs. *manuscripta math.* **66** (1990), no. 3, 277–293.
- [19] R. H. Villarreal, *Monomial algebras*. Monographs and Textbooks in Pure and Applied Mathematics, **238**. Marcel Dekker, Inc., New York, 2001.
- [20] X. Zheng, Resolutions of Facet Ideals. *Comm. Algebra* **32** (2004) 2301–2324.

DEPARTMENT OF MATHEMATICS AND STATISTICS, QUEEN'S UNIVERSITY, KINGSTON, ONTARIO, K7L 3N6

*E-mail address:* [mikeroth@mast.queensu.ca](mailto:mikeroth@mast.queensu.ca)

DEPARTMENT OF MATHEMATICAL SCIENCES, LAKEHEAD UNIVERSITY, THUNDER BAY, ON P7B 5E1, CANADA

*E-mail address:* [avantuy1@sleet.lakeheadu.ca](mailto:avantuy1@sleet.lakeheadu.ca)