EXPLICIT DESCENT VIA 4-ISOGENY ON AN ELLIPTIC CURVE

EDRAY HERBER GOINS

ABSTRACT. We work out the complete descent via 4-isogeny for a family of rational elliptic curves with a rational point of order 4; such a family is of the form $y^2 + xy + ay = x^3 + ax^2$ where $\sqrt{-a} \in \mathbb{Q}^{\times}$. In the process we exhibit the 4-isogeny and the isogenous curve, explicitly present the principal homogeneous spaces, and discuss examples by computing the rank.

1. Introduction

For each $t \in \mathbb{Q}^{\times}$ we have the rational elliptic curve

(1.1)
$$E_t: v^2 = u^3 + (t^2 + 2)u^2 + u \text{ where } [4](-1,t) = \mathcal{O}.$$

This curve is studied in detail in [2], where, among other things, it is shown that the torsion subgroup \mathbb{T} of $E_t(\mathbb{Q})$ is completely classified as being either

$$(1.2) \quad \mathbb{T} \simeq \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{or} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} & \text{if } t = (s^2 - 1)/s \text{ for } s \in \mathbb{Q}^\times, \text{ or} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z} & \text{if } t = (s^2 - 1)/s \text{ and } s = (r^2 - 1)/(2\,r) \text{ for } r \in \mathbb{Q}^\times. \end{cases}$$

On the other hand, the rank of $E_t(\mathbb{Q})$ may be computed by a descent via two-isogeny coming from the rational point [2](-1,t)=(0,0) of order 2, as implemented in computer packages such as MAGMA [1]. Unfortunately, there may be a global obstruction in computing the rank due to nontrivial elements in the Shafarevich-Tate group, so one can only find a lower bound on the rank in general. In this exposition, I show how to compute the rank of $E_t(\mathbb{Q})$ by exploiting instead the rational point (-1,t) of order 4 i.e. I work out a descent via four-isogeny.

The main result of this paper is the following:

Theorem 1.1. Fix $t \in \mathbb{Q}^{\times}$, and denote the elliptic curves

(1.3)
$$E_t: \qquad v^2 = u^3 + (t^2 + 2) u^2 + u,$$

$$E_t': \qquad V^2 = U^3 - 2(t^2 - 4) U^2 + (t^2 + 4)^2 U,$$

$$E_t'': \qquad \mathfrak{v}^2 = \mathfrak{u}^3 + (t^2 - 4) \mathfrak{u}^2 - 4t^2 \mathfrak{u}.$$

(1) There are isogenies $\phi: E_t \to E'_t$ of degree 4 with kernel generated by (-1,t), and $\varphi: E_t \to E''_t$ of degree 2 generated by [2](-1,t) = (0,0).

²⁰⁰⁰ Mathematics Subject Classification. 14G05, 11G05, 11D25. Key words and phrases. elliptic curve, rank, torsion, descent, isogeny.

(2) Upon identifying the Selmer groups as subgroups of $\mathbb{Q}^{\times}/(\mathbb{Q}^{\times})^4$ and $\mathbb{Q}^{\times}/(\mathbb{Q}^{\times})^2$, respectively; and Shafarevich-Tate groups as a collection of homogeneous spaces, there is an exact diagram

$$0 \longrightarrow \frac{E_{t}(\mathbb{Q})}{\hat{\phi}(E'_{t}(\mathbb{Q}))} \stackrel{\delta'}{\longrightarrow} S^{(\hat{\phi})}(E'_{t}/\mathbb{Q}) \longrightarrow \coprod (E'_{t}/\mathbb{Q})[\hat{\phi}] \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad$$

where we have the connecting homomorphisms

(1.5)
$$\delta': \frac{(-1,t) \mapsto -4t^2}{(u,v) \mapsto u^2 + 2(t^2 + 1)u + 1 - 2tv} \qquad \delta'': \frac{(0,0) \mapsto t^2 + 4}{(u,v) \mapsto u}$$

while the maps into the Shafarevich-Tate groups send $d \mapsto \{C'_{d,t}/\mathbb{Q}\}$ and $d \mapsto \{C''_{d,t}/\mathbb{Q}\}$ in terms of

(1.6)
$$C'_{d,t}: \qquad d\left(w - \frac{z^2}{4t^2}\right)z^2 = \left(w^2 - d\right)^2,$$
$$C''_{d,t}: \qquad dW^2 = d^2 + \left(t^2 + 2\right)dZ^2 + Z^4.$$

(3) The composition

(1.7)
$$\psi': C'_{d,t} \xrightarrow{\eta} C''_{d,t} \xrightarrow{\psi''} E_t$$

in terms of the maps η and ψ'' sending

$$(1.8) \quad (z,w) \mapsto \left(-\frac{dz^2}{2t(w^2-d)}, \frac{dz^2(w^2+d)}{2(w^2-d)^2}\right), \qquad (Z,W) \mapsto \left(\frac{d}{Z^2}, -\frac{dW}{Z^3}\right);$$
satisfy the congruence $\delta'(\psi'(P)) \equiv d \mod (\mathbb{Q}^{\times})^4.$

I mention that the article [4] outlines a similar algorithm using a 4-descent by considering a diagram such as

$$0 \longrightarrow \frac{E(\mathbb{Q})}{4E(\mathbb{Q})} \longrightarrow S^{(4)}(E/\mathbb{Q}) \longrightarrow \mathrm{III}(E/\mathbb{Q})[4] \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$0 \longrightarrow \frac{E(\mathbb{Q})}{2E(\mathbb{Q})} \longrightarrow S^{(2)}(E/\mathbb{Q}) \longrightarrow \mathrm{III}(E/\mathbb{Q})[2] \longrightarrow 0$$

(I have learned that Tom Womack is working on implementation of a similar algorithm for a future version of MAGMA.) The paper at hand may be considered a refinement of this algorithm, even though I present the algorithm for a limited family of elliptic curves possessing a 4-isogeny.

I am motivated to implement such an algorithm by the search for elliptic curves E_t with $\mathbb{T} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$ having "large" rank. In [2], there is an elliptic curve with rank 3 corresponding to r = 15/76. Moreover, the author, along with Garikai Campbell, conjectured that if

$$(1.10) r \in \left\{ \frac{15}{56}, \frac{24}{65}, \frac{11}{69}, \frac{7}{88}, \frac{12}{97} \right\}$$

then $E_t(\mathbb{Q})$ will have rank at least 4. Randall Rathbun [5] has informed me that if we assume the validity of the conjectures of Birch and Swinnerton-Dyer then our conjecture is false; in fact the curves have rank not greater than 2. This is apparently because the Shafarevich-Tate groups in the sequence (1.11)

$$0 \longrightarrow \mathrm{III}(E_t/\mathbb{Q})[\phi] \longrightarrow \mathrm{III}(E_t/\mathbb{Q})[4] \stackrel{\phi}{\longrightarrow} \mathrm{III}(E_t'\mathbb{Q})[\hat{\phi}] \longrightarrow 0$$

are quite large. Explicitly, for these five values of r we have $\#\coprod(E'_t\mathbb{Q})[\hat{\phi}] \geq 2^4$.

The authors would like to thank Garikai Campbell, John Cremona, Lloyd Kilford, William Stein, and Randall Rathbun for helpful conversations.

Contents

1. Introduction	1
2. Descent via Four-Isogeny	3
2.1. The 4-Isogeny	3
2.2. The Isogenous Curve	4
2.3. The Connecting Homomorphism	6
2.4. The Weil-Châtelet Group	8
2.5. Selmer and Shafarevich-Tate Groups	9
2.6. Complete Descent via 4-Isogeny	10
3. Descent via Two-Isogeny	10
3.1. The 2-Isogenies	11
3.2. Complete Descent via 2-Isogeny	12
3.3. Modified Descent via 4-Isogeny	13
4. Examples	14
4.1. Example: $\mathbb{T} \simeq \mathbb{Z}/4\mathbb{Z}$.	14
4.2. Example: $\mathbb{T} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$.	15
4.3. Example: $\mathbb{T} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$.	16
References	19

2. Descent via Four-Isogeny

For this section, we fix $t \in \mathbb{Q}^{\times}$ and drop the notation involving t.

2.1. **The 4-Isogeny.** We begin by explicitly describing the 4-isogeny and the isogenous elliptic curve.

Proposition 2.1. The elliptic curve E_t is rationally equivalent to

(2.1)
$$E: y^2 + xy + ay = x^3 + ax^2$$
 where $a = -\frac{1}{4t^2}$, [4] $(0,0) = \mathcal{O}$;

and is 4-isogenous to the elliptic curve

(2.2)
$$E': Y^2 + XY + AY = X^3 + AX^2 \quad where \quad A = \frac{t^2 + 4}{64}.$$

Moreover, the isogeny $\phi: E \to E'$ sends $(0,0) \mapsto \mathcal{O}$.

One can easily show that any rational elliptic curve with a rational point of order 4 is birationally equivalent to a curve in the form above for some $a \in \mathbb{Q}^{\times}$. Indeed, see [7, Exercise 8.13(a)] as well as [3, pg. 217]. The condition $\sqrt{-a} \in \mathbb{Q}^{\times}$ is special to simplify the formulas to follow.

Proof. Starting with the curve $v^2 = u^3 + (t^2 + 2)u^2 + u$, make the substitution

(2.3)
$$x = \frac{u+1}{4t^2}$$
 and $y = \frac{v-tu}{8t^3} \implies (-1, t) \mapsto (0, 0).$

Then E has the Weierstrass equation above. Following the ideas in [8] we produce an isogeny $\phi: E \to E'$ such that the cyclic subgroup

(2.4)
$$E[\phi] = \{(0,0), (-a,0), (0,-a), \mathcal{O}\}\$$

is the kernel. We choose such a map that sends $(x,y) \mapsto (X,Y)$ in terms of

(2.5)
$$X = -A + \left(\frac{t(x+2a)(2y+x+a)}{8x(x+a)}\right)^{2},$$

$$Y = X^{2} - (x+a)^{2} \left(\frac{(2y+x+a)-2tx(x+2a)}{8x(x+a)}\right)^{4}.$$

The new curve E' has a rational point of order 4, but we choose the dual isogeny $\hat{\phi}: E' \to E$ such that

(2.6)
$$E'[\hat{\phi}] = \left\{ \left(-2A, A \frac{2+it}{4} \right), (-A, 0), \left(-2A, A \frac{2-it}{4} \right), \mathcal{O} \right\}$$

where $i = \sqrt{-1}$. This map sends $(X, Y) \mapsto (x, y)$ in terms of

(2.7)
$$x = -a + \left(\frac{X(2Y + X + A)}{2t(X + A)(X + 2A)}\right)^{2},$$
$$y = x^{2} - \left(\frac{2(X + 2A)(2Y + X + A) - tX(X + A)}{4t(X + A)(X + 2A)}\right)^{4}.$$

One checks that $\hat{\phi} \circ \phi = [4]$ is the "multiplication-by-4" map on E, while $\phi \circ \hat{\phi} = [4]$ is the "multiplication-by-4" map on E'.

2.2. **The Isogenous Curve.** The following proposition shows how the size of $E(\mathbb{Q})/4 E(\mathbb{Q})$ is related to the size of $E(\mathbb{Q})/\hat{\phi}(E'(\mathbb{Q}))$.

Proposition 2.2. Let $\phi: E \to E'$ be the isogeny in Proposition 2.1 and $\hat{\phi}: E' \to E$ denote its dual.

(1) The following sequence is exact if and only if $\sqrt{t^2+4} \in \mathbb{Q}^{\times}$:

$$(2.8) 0 \longrightarrow \frac{E'(\mathbb{Q})}{\phi(E(\mathbb{Q}))} \stackrel{\hat{\phi}}{\longrightarrow} \frac{E(\mathbb{Q})}{4E(\mathbb{Q})} \longrightarrow \frac{E(\mathbb{Q})}{\hat{\phi}(E'(\mathbb{Q}))} \longrightarrow 0.$$

- (2) Both $E(\mathbb{Q})$ and $E'(\mathbb{Q})$ have the same rank.
- (3) The torsion subgroup \mathbb{T}' of $E'(\mathbb{Q})$ is $\mathbb{Z}/8\mathbb{Z}$ when $\sqrt{t+\sqrt{t^2+4}} \in \mathbb{Q}^{\times}$, and $\mathbb{Z}/4\mathbb{Z}$ otherwise.

Proof. The dual isogeny induces the exact sequence

$$(2.9) 0 \to \frac{E'(\mathbb{Q})[\hat{\phi}]}{\phi\left(E(\mathbb{Q})[4]\right)} \to \frac{E'(\mathbb{Q})}{\phi\left(E(\mathbb{Q})\right)} \to \frac{E(\mathbb{Q})}{4\,E(\mathbb{Q})} \to \frac{E(\mathbb{Q})}{\hat{\phi}\left(E'(\mathbb{Q})\right)} \to 0.$$

We compute the first term in this sequence. From above, $E'(\mathbb{Q})[\hat{\phi}] = \{(-A,0), \mathcal{O}\}$ has order 2. E[4] is generated by P = (0,0) and

(2.10)
$$Q = \left(-\frac{2a}{1+\alpha}, \frac{8a^3}{(1+\alpha)^2(1+i\alpha)}\right) \text{ where } \alpha = \sqrt[4]{\frac{t^2+4}{t^2}}.$$

One verifies that $\phi(P) = \mathcal{O}$ while $\phi([2]Q) = (-A, 0)$. Clearly $Q \notin E(\mathbb{Q})[4]$, while

$$(2.11) \hspace{1cm} [2]\,Q = \left(-\frac{1+\alpha^2}{8},\ \frac{(1+\alpha^2)^2}{32}\right) \in E(\mathbb{Q})[4] \iff \sqrt{t^2+4} \in \mathbb{Q}^\times.$$

Hence $E'(\mathbb{Q})[\hat{\phi}] = \phi\left(E(\mathbb{Q})[4]\right)$ if and only if t^2+4 is a square.

Second, the statement that isogenous elliptic curves have the same rank is well-known: As $E'(\mathbb{Q})/\phi(E(\mathbb{Q}))$ is finite the rank of $E'(\mathbb{Q})$ cannot be greater than the rank of $E(\mathbb{Q})$. Similarly, $E(\mathbb{Q})/\hat{\phi}(E'(\mathbb{Q}))$ is finite so we must have equality with the rank.

As for the torsion, it is easy to see that $(0,0) \in E'(\mathbb{Q})$ is a rational point of order 4. By Mazur's Theorem, either $E'[2] \subseteq E'(\mathbb{Q})$ or else the torsion subgroup is of the form $\mathbb{Z}/4n\mathbb{Z}$ for n=1,2,3. We begin by showing that not all of the 2-torsion is rational. The points (X,Y) of order 2 satisfy 2Y + X + A = 0, so that

$$(2.12) (X+A)(4X^2+X+A) = -4(Y^2+XY+AY-X^3-AX^2) = 0.$$

(-A,0) is one rational point of order 2 corresponding to the linear factor, but the quadratic factor has discriminant -1 so that polynomial has no rational roots.

Next we show that there are no rational points of order 3. Assume that (X, Y) is such a point. One computes that the 3-division polynomial of E' is

$$\psi_3(X,t) = 3X^4 + (1+4A)X^3 + 3AX^2 + 3A^2X + A^3.$$

We make the substitution

(2.14)
$$\sigma = -\frac{3X + 4A}{X}$$
 and $\tau = \frac{3X + 4A}{X + 2A}t$.

Note that $X(X+2A) \neq 0$ because otherwise (X,Y) would be a point of order 4. Also, $3X+4A \neq 0$: if not, then $(X,Y)=(-4A/3, -A\beta/9)$ in terms of a root of the quadratic $\beta^2+3\beta+48A$. However, this quadratic has discriminant $-3(t^2+1)$ which is not a square, so (X,Y) is not a rational point. Hence σ and τ are well-defined and nonzero. Solving for X and t transforms the 3-division polynomial as

$$(2.15) \qquad \psi_3\left(-\frac{4\,A}{\sigma+3},\,\,\frac{\tau\left(\sigma+1\right)}{2\,\sigma}\right) = \frac{A^3\left(\sigma+1\right)^2}{\sigma\left(\sigma+3\right)^4}\cdot\left[\tau^2-\sigma\left(\sigma+1\right)\left(\sigma-3\right)\right]$$

so that (σ, τ) is a rational point on the elliptic curve $\tau^2 = \sigma(\sigma+1)(\sigma-3)$. However, one checks that the only such (affine) rational points satisfy $\tau = 0$, a contradiction.

Finally, we are in the case where we distinguish between $\mathbb{Z}/4\mathbb{Z}$ and $\mathbb{Z}/8\mathbb{Z}$. Say that R = (X, Y) is a rational point of order 8. Then [2] R is a point of order 4, so without loss of generality say [2] R = P. By considering the X-coordinates we have

$$(2.16) X([2]R) = \frac{X^4 - A(X+A)}{(X+A)(4X^2 + X + A)} = 0 \implies X^2 - \sqrt{A}(X+A) = 0.$$

Hence $\sqrt{A} \in \mathbb{Q}^{\times}$. In fact, make the substitution

(2.17)
$$\gamma = \sqrt{t + \sqrt{t^2 + 4}} \implies t = \frac{\gamma^4 - 4}{2\gamma^2} \quad \text{and} \quad A = \left(\frac{\gamma^4 + 4}{16\gamma^2}\right)^2.$$

The discriminant of the quadratic $X^2 - \sqrt{A}(X + A)$, namely γ^2 , must be a square since its roots are rational; hence $\gamma \in \mathbb{Q}^{\times}$. Then the rational point of order 8 is

$$(2.18) R = \left(\frac{(\gamma^4 + 4)(\gamma^2 + 2\gamma + 2)}{64\gamma^3}, \frac{(\gamma^4 + 4)^2(\gamma^2 + 2\gamma + 2)}{1024\gamma^5}\right).$$

The converse is clear i.e. if $[8]R = \mathcal{O}$ then $\sqrt{t + \sqrt{t^2 + 4}} \in \mathbb{Q}^{\times}$.

2.3. The Connecting Homomorphism. There is a natural bijection between the crossed homomorphisms $H^1\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E\right)$ and $WC\left(E/\mathbb{Q}\right)$, the collection of equivalence classes of homogeneous spaces for E, usually called the Weil-Châtelet group. We will find it more convenient to work with the isogenous curve E': The exact sequence

$$(2.19) 0 \longrightarrow E'[\hat{\phi}] \longrightarrow E' \stackrel{\hat{\phi}}{\longrightarrow} E \longrightarrow 0$$

implies, through Galois cohomology, the exact sequence

$$(2.20) \quad 0 \longrightarrow \frac{E(\mathbb{Q})}{\hat{\phi}\left(E'(\mathbb{Q})\right)} \stackrel{\delta_E}{\longrightarrow} H^1\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E'[\hat{\phi}]\right) \longrightarrow WC\left(E'/\mathbb{Q}\right)$$

via the connecting homomorphism δ_E . We make such a homomorphism explicit by considering a generalization of the Weil pairing.

Proposition 2.3. (1) There is a pairing $e_{\phi}: E'[\hat{\phi}] \times E[\phi] \to \mu_4$ that is bilinear, alternating, non-degenerate, and Galois invariant; which satisfies

(2.21)
$$e_{\phi}(T',T) = \sqrt{-1}$$
 when $T' = \left(-2A, A \frac{2+\sqrt{-1}t}{4}\right), T = (0,0).$

(2) The pairing $b: E(\mathbb{Q})/\hat{\phi}(E'(\mathbb{Q})) \times E[\phi] \to \mathbb{Q}^{\times}/(\mathbb{Q}^{\times})^4$ defined by

(2.22)
$$(x,y), [m](0,0) \mapsto \begin{cases} a^{-m} & (x,y) = (0,0), \\ (x^2 - y)^m & otherwise; \end{cases}$$

is bilinear and non-degenerate on the left.

(3) Upon identifying $E'[\hat{\phi}] \simeq \mu_4$ as $Gal(\overline{\mathbb{Q}}/\mathbb{Q})$ -modules we have

(2.23)
$$H^{1}\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E'[\hat{\phi}]\right) \simeq \mathbb{Q}^{\times}/(\mathbb{Q}^{\times})^{4}.$$

(4) The composition of these maps

$$(2.24) \qquad \frac{E(\mathbb{Q})}{\hat{\phi}\left(E'(\mathbb{Q})\right)} \times E[\phi] \stackrel{b}{\longrightarrow} \frac{\mathbb{Q}^{\times}}{(\mathbb{Q}^{\times})^{4}} \stackrel{\delta_{\mathbb{Q}}}{\longrightarrow} H^{1}\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E'[\hat{\phi}]\right)$$

relates the connecting homomorphisms by

(2.25)
$$e_{\phi}(\delta_E(P)(\sigma), T) = \delta_{\mathbb{Q}}(b(P,T))(\sigma) \text{ for all } \sigma \in Gal(\overline{\mathbb{Q}}/\mathbb{Q}).$$

In particular, we may identify $\delta_E : P \mapsto \delta_{\mathbb{Q}}(b(P,T))$ when $T = (0,0).$

Proof. We follow the construction of the Weil pairing in [7, §8, pgs. 95 – 99], but we make the formulas explicit. Consider the function $f \in \mathbb{Q}(E)$ defined as $f(x,y) = x^2 - y$. One checks that

$$(2.26) x^2 - y = 0 \implies x^4 = y^2 + xy + ay - x^3 - ax^2 = 0 \implies (x, y) = (0, 0);$$

so that the divisor of this function is $\operatorname{div}(f) = 4((0,0)) - 4(\mathcal{O})$. Similarly, we consider the function $g \in \mathbb{Q}(E')$ defined by

(2.27)
$$g(X,Y) = \frac{2(X+2A)(2Y+X+A) - tX(X+A)}{4t(X+A)(X+2A)};$$

the formulas in (2.7) show that $f \circ \hat{\phi} = g^4$. For $P' \in E'(\overline{\mathbb{Q}})$ and $T' \in E'[\hat{\phi}]$ we have

$$(2.28) g\left(P'\oplus T'\right)^4 = f\left(\hat{\phi}(P')\oplus\hat{\phi}(T')\right) = f\left(\hat{\phi}(P')\oplus\mathcal{O}\right) = g\left(P'\right)^4.$$

We define the pairing $e_{\phi}: E'[\hat{\phi}] \times E[\phi] \to \mu_4$ by

(2.29)
$$e_{\phi}\left(T',T\right) = \left(\frac{g\left(P' \oplus T'\right)}{g\left(P'\right)}\right)^{m} \quad \text{where} \quad T = [m]\left(0,0\right).$$

The bilinear, alternating, non-degenerate, and Galois invariant properties follow from the arguments given in $[7, \S 8, pgs. 95 - 99]$. In particular, one verifies that (2.30)

$$T' = \left(-2A, A \frac{2+\sqrt{-1}t}{4}\right), \quad T = (0,0) \implies e_{\phi}(T', T) = \frac{g(P' \oplus T')}{g(P')} = \sqrt{-1}.$$

Consider the map $b: E(\mathbb{Q}) \times E[\phi] \to \mathbb{Q}^{\times}/(\mathbb{Q}^{\times})^4$. First we show it is well-defined. As shown above, $x^2 - y = 0$ if and only if (x, y) = (0, 0). On the other hand, we have by assumed bilinearity

(2.31)
$$b(0,0), [m](0,0) = b([-1](0,0), [m](0,0))^{-1} = a^{-m}.$$

Now we show that this map can be extended to the quotient. When $(x,y) = \hat{\phi}(X,Y)$ then $x^2 - y$ is a fourth-power by (2.7) so that the map extends to a well-defined map $b: E(\mathbb{Q})/\hat{\phi}\left(E'(\mathbb{Q})\right) \times E[\phi] \to \mathbb{Q}^\times/(\mathbb{Q}^\times)^4$. The fact that b is a non-degenerate pairing follows from the same arguments as with the Weil pairing above.

Fix $d \in \mathbb{Q}^{\times}$, and consider the map $\xi : \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to E'[\hat{\phi}]$ defined by

$$(2.32) \quad \xi: \quad \sigma \mapsto \xi_{\sigma} = [m] \left(-2A, \ A \frac{2+it}{4} \right) \quad \text{whenever} \quad \frac{\sigma(\sqrt[4]{d})}{\sqrt[4]{d}} = (-1)^{m/2}.$$

The map ξ is a 1-cocycle, so has a cohomology class $\{\xi\} \in H^1\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E'[\hat{\phi}]\right)$. Identify translation by this point with the Galois action i.e. $(X,Y)^{\sigma}=(X,Y)\oplus \xi_{\sigma}$. From the map $d\mapsto d^4$ we find the well-known Kummer sequence

$$(2.33) 1 \longrightarrow E'[\hat{\phi}] \simeq \mu_4 \longrightarrow \overline{\mathbb{Q}}^{\times} \longrightarrow \overline{\mathbb{Q}}^{\times} \longrightarrow 1$$

of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -modules, so from Galois cohomology we find the exact sequence (2.34)

$$1 \longrightarrow \frac{\mathbb{Q}^{\times}}{(\mathbb{Q}^{\times})^{4}} \stackrel{\delta_{\mathbb{Q}}}{\longrightarrow} H^{1}\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E'[\hat{\phi}]\right) \longrightarrow H^{1}\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), \overline{\mathbb{Q}}^{\times}\right).$$

where $\delta_{\mathbb{Q}}: d \mapsto \{\xi\}$. Hilbert's Theorem 90 states the last term is trivial; hence the third part of the proposition holds.

To show the fourth part of the proposition, fix $P \in E(\mathbb{Q})$, choose $P' \in E'(\overline{\mathbb{Q}})$ satisfying $P = \hat{\phi}(P')$, and denote $\xi = \delta_E(P)$. For any $\sigma \in \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ we have

$$(2.35) e_{\phi}(\delta_{E}(P)(\sigma), T) = e_{\phi}(\xi_{\sigma}, [m](0,0)) = \left(\frac{g(P' \oplus \xi_{\sigma})}{g(P')}\right)^{m} = \left(\frac{g(P')^{\sigma}}{g(P')}\right)^{m} = \left(\delta_{\mathbb{Q}}(d)(\sigma)\right)^{m} = \delta_{\mathbb{Q}}(b(P,T))(\sigma).$$

where we have identified $E'[\hat{\phi}] \simeq \mu_4$ and set $d = f(P) \equiv b(P, (0, 0))$.

2.4. The Weil-Châtelet Group. The full collection of homogeneous spaces in the Weil-Châtelet group is too large for interest in this exposition, so instead we consider the image $WC(E'/\mathbb{Q})[\hat{\phi}]$ of the crossed homomorphisms studied above. Recall that $WC(E'/\mathbb{Q})$ consists of equivalence classes $\{C'/\mathbb{Q}\}$ of certain rational curves, where a class is trivial if and only if $C'(\mathbb{Q})$ is nonempty.

Proposition 2.4. (1) The composite map

$$(2.36) \qquad \frac{\mathbb{Q}^{\times}}{(\mathbb{Q}^{\times})^4} \xrightarrow{\delta_{\mathbb{Q}}} H^1\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E'[\hat{\phi}]\right) \longrightarrow WC\left(E'/\mathbb{Q}\right)$$

sends $d \mapsto \{C'_d/\mathbb{Q}\}$, where

(2.37)
$$C'_d: \quad d(w+az^2)z^2 = (w^2-d)^2.$$

(2) There is an isomorphism $\theta: C'_d \to E'$ defined over $\overline{\mathbb{Q}}$ such that

$$(2.38) \psi': C'_d \xrightarrow{\theta} E' \xrightarrow{\hat{\phi}} E, (z,w) \mapsto \left(\frac{w}{z^2}, \frac{w^2 - d}{z^4}\right).$$

Proof. We construct a homogeneous space corresponding the cohomology class $\{\xi\} \in H^1\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E'[\hat{\phi}]\right)$ following the exposition in [7, §3, pgs. 287 – 296]. Fix $d \in \mathbb{Q}^{\times}$, and consider the functions $z, w \in \overline{\mathbb{Q}}(E')$ defined implicitly by

(2.39)
$$z = \frac{\sqrt[4]{d}}{g(P')}$$
 and $-\frac{z}{w} = \frac{1/\sqrt[4]{d}}{g([-1]P')}$ where $P' = (X, Y)$.

It is easy to check that the point (z, w) is on C'_d as defined above. We show these functions are Galois invariant. To this end, choose $\sigma \in \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, and say that $\sigma(\sqrt[4]{d}) = (-1)^{m/2} \sqrt[4]{d}$. For any integer n we have

$$(2.40) \quad \sigma\left(\frac{\sqrt[4]{d^n}}{g([n]P')}\right) = \frac{(-1)^{mn/2}\sqrt[4]{d^n}}{g([n]P'^{\sigma})} = \frac{(-1)^{mn/2}\sqrt[4]{d^n}}{g([n]P'\oplus T')} = \frac{(-1)^{mn/2}}{e_{\phi}(T',T)}\frac{\sqrt[4]{d^n}}{g([n]P')}$$

where $T' = [n] \xi_{\sigma}$ and T = (0,0). It follows from Proposition 2.3 that $e_{\phi}(T',T) = (-1)^{mn/2}$. Hence the equations defining z and w are Galois invariant, so that z and w themselves are Galois invariant.

This defines a map $E' \to C'_d$ sending $(X, Y) \mapsto (z, w)$ in terms of

(2.41)
$$z = \sqrt[4]{d} \frac{4t(X+A)(X+2A)}{2(X+2A)(2Y+X+A)-tX(X+A)},$$

$$w = \sqrt{d} \frac{2(X+2A)(2Y+X+A)+tX(X+A)}{2(X+2A)(2Y+X+A)-tX(X+A)}.$$

The inverse map $\theta: C'_d \to E'$ – which can be found by considering $g(P') \pm g([-1]P')$ – sends $(z, w) \mapsto (X, Y)$ in terms of

$$(2.42) X = 4 A \frac{w - \sqrt{d}}{\sqrt[4]{d} z - 2(w - \sqrt{d})}, Y = (X + A) \frac{-\sqrt[4]{d} z + t(w + \sqrt{d})}{2\sqrt[4]{d} z}.$$

One checks using the formulas in (2.7) that the composition $\psi' = \hat{\phi} \circ \theta$ is the map $(z, w) \mapsto (w/z^2, (w^2 - d)/z^4)$.

2.5. **Selmer and Shafarevich-Tate Groups.** Combining the exact sequences introduced above, we have the exact diagram

$$0 \longrightarrow \frac{E'(\mathbb{Q})}{\phi(E(\mathbb{Q}))} \longrightarrow S^{(\phi)}(E/\mathbb{Q}) \longrightarrow \operatorname{III}(E/\mathbb{Q})[\phi] \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$0 \longrightarrow \frac{E(\mathbb{Q})}{4E(\mathbb{Q})} \longrightarrow S^{(4)}(E/\mathbb{Q}) \longrightarrow \operatorname{III}(E/\mathbb{Q})[4] \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \phi$$

$$0 \longrightarrow \frac{E(\mathbb{Q})}{\hat{\phi}(E'(\mathbb{Q}))} \stackrel{\delta_E}{\longrightarrow} S^{(\hat{\phi})}(E'/\mathbb{Q}) \longrightarrow \operatorname{III}(E'/\mathbb{Q})[\hat{\phi}] \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$0 \qquad \qquad 0 \qquad \qquad 0$$

where the Selmer and Shafarevich-Tate groups are defined as

$$(2.44) S^{(\phi)}(E/\mathbb{Q}) = \ker \left\{ H^1\left(\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}), E[\phi]\right) \to \prod_{\nu} WC\left(E/\mathbb{Q}_{\nu}\right) \right\},$$

$$\operatorname{III}(E/\mathbb{Q}) = \ker \left\{ WC\left(E/\mathbb{Q}\right) \to \prod_{\nu} WC\left(E/\mathbb{Q}_{\nu}\right) \right\};$$

taking the product over all places ν of \mathbb{Q} . One can compute the rank of $E(\mathbb{Q})$ with the explicit map in Proposition 2.4 – assuming that $\coprod (E'/\mathbb{Q})[\hat{\phi}]$ is trivial.

Proposition 2.5. Let Σ be the finite set of places of \mathbb{Q} consisting of $2, \infty$, and those primes occurring in the factorization of t and $t^2 + 4$. We have

(2.45)
$$S^{(\hat{\phi})}(E'/\mathbb{Q}) \simeq \left\{ d = \pm \prod_{p \in \Sigma} p^{a_p} \in \frac{\mathbb{Q}^{\times}}{(\mathbb{Q}^{\times})^4} \middle| C'_d(\mathbb{Q}_{\nu}) \neq \emptyset \text{ for all } \nu \in \Sigma \right\},$$

$$III(E'/\mathbb{Q})[\hat{\phi}] \simeq \left\{ \{C'_d/\mathbb{Q}\} \in WC(E'/\mathbb{Q})[\hat{\phi}] \middle| C'_d(\mathbb{Q}_{\nu}) \neq \emptyset \text{ for all } \nu \in \Sigma \right\}.$$

Proof. We may identify the Selmer group with a subgroup of $\mathbb{Q}/(\mathbb{Q}^{\times})^4$ by the isomorphism in Proposition 2.3. The subgroup $WC(E'/\mathbb{Q})[\hat{\phi}] \subseteq WC(E'/\mathbb{Q})$ consists of the equivalence classes $\{C'_d/\mathbb{Q}\}$. By definition, a class $\{C'_d/\mathbb{Q}_{\nu}\} \in WC(E'/\mathbb{Q}_{\nu})$ is trivial if and only if $C'_d(\mathbb{Q}_{\nu})$ is nonempty.

2.6. Complete Descent via 4-Isogeny. We collect the results proved thus far into one statement.

Theorem 2.6. Fix $t \in \mathbb{Q}^{\times}$, and let E_t and E'_t denote the elliptic curves

$$(2.46) E_t: v^2 = u^3 + (t^2 + 2) u^2 + u, \quad E'_t: V^2 = U^3 - 2(t^2 - 4) U^2 + (t^2 + 4)^2 U.$$

- (1) There is an isogeny $\phi: E_t \to E'_t$ of degree 4 with kernel generated by (-1,t).
- (2) Upon identifying the Selmer and Shafarevich-Tate groups as in Proposition 2.5, there is an exact sequence

$$(2.47) \quad 0 \longrightarrow \frac{E_t(\mathbb{Q})}{\hat{\phi}(E'_t(\mathbb{Q}))} \stackrel{\delta'}{\longrightarrow} S^{(\hat{\phi})}(E'_t/\mathbb{Q}) \longrightarrow \mathrm{III}(E'_t/\mathbb{Q})[\hat{\phi}] \longrightarrow 0$$

where δ' sends $[m] (-1,t) \mapsto \left(-4\,t^2\right)^m$ and $(u,v) \mapsto u^2 + 2\,(t^2+1)\,u + 1 - 2\,t\,v$ otherwise; while the second map sends $d \mapsto \{C'_{d,t}/\mathbb{Q}\}$ in terms of

(2.48)
$$C'_{d,t}: \quad d\left(w - \frac{z^2}{4t^2}\right)z^2 = \left(w^2 - d\right)^2.$$

(3) The map $\psi': C'_{d,t} \to E_t$ sending

$$(2.49) (z,w) \mapsto \left(\frac{4t^2(w^2-d)^2}{dz^4}, \frac{4t^3(w^2-d)(w^2+d)}{dz^4}\right)$$

satisfies the congruence $\delta'(\psi'(P)) \equiv d \mod (\mathbb{Q}^{\times})^4$.

Proof. The curves E_t and E'_t are related to E and E' by the transformation

(2.50)
$$x = \frac{u+1}{4t^2}$$
, $y = \frac{v-tu}{8t^3}$ and $X = \frac{U-(t^2+4)}{64}$, $Y = \frac{V-4U}{512}$.

The first statement follows from Proposition 2.1. The image of δ in the second statement follows from Proposition 2.3, Proposition 2.4, and the congruence

(2.51)
$$x^2 - y \equiv (u^2 + 2(t^2 + 1)u + 1) - 2tv \mod (\mathbb{Q}^{\times})^4;$$

while the image of the second map follows from Proposition 2.4. The third statement follows from Proposition 2.4 where we write the map ψ' in terms of u and v rather than x and y. From the identity

$$(2.52) (u,v) = \psi'(z,w) \implies (u^2 + 2(t^2 + 1)u + 1) - 2tv = d\left(\frac{2t}{z}\right)^4$$

we find the congruence $(u^2 + 2(t^2 + 1)u + 1) - 2tv \equiv d \mod (\mathbb{Q}^{\times})^4$.

3. Descent via Two-Isogeny

The descent algorithm associated to the rational point of order 4 described in the previous section can be refined to exploit the 2-isogeny coming from the doubling of this point.

3.1. **The 2-Isogenies.** We begin by factoring the 4-isogeny.

Proposition 3.1. Let $\phi: E \to E'$ as in Proposition 2.1, and denote the curve

(3.1)
$$E'': \qquad \mathfrak{v}^2 = \mathfrak{u}(\mathfrak{u} - 4)(\mathfrak{u} + t^2).$$

There exist rational 2-isogenies $\varphi: E \to E''$ and $\eta: E' \to E''$ such that $\phi = \hat{\eta} \circ \varphi$.

Proof. The isogeny ϕ and the "multiplication-by-2" map on E induce the following exact diagram:

We will construct isogenies $\varphi: E \to E''$ and $\eta: E' \to E''$ with kernels

$$(3.3) \quad E[\varphi] = 2\,E[\phi] = \big\{ (-a,0), \; \mathcal{O} \big\} \quad \text{and} \quad E'[\eta] = \phi(E[2]) = \big\{ (-A,0), \; \mathcal{O} \big\}.$$

To this end, assume without loss of generality that $E = E_t$ and $E' = E'_t$ as in Theorem 2.6, where now (u, v) = (0, 0) and (U, V) = (0, 0) are the rational points of order 2. Following the exposition in [7], we define the isogenies

(3.4)
$$\varphi: E \to E'', \qquad (u,v) \mapsto \left(\frac{(u+1)^2}{u}, \frac{1-u^2}{u^2}v\right);$$
$$\eta: E' \to E'', \qquad (U,V) \mapsto \left(\frac{V^2}{4U^2}, \frac{(t^2+4)^2-U^2}{8U^2}V\right);$$

with dual isogenies

$$(3.5) \qquad \begin{aligned} \hat{\varphi}: \quad E'' \to E, \qquad & (\mathfrak{u}, \mathfrak{v}) \mapsto \left(\frac{\mathfrak{v}^2}{4\left(\mathfrak{u} + t^2\right)^2}, \, -\frac{-4\,t^2 + 2\,t^2\,\mathfrak{u} + \mathfrak{u}^2}{8\left(\mathfrak{u} + t^2\right)^2}\,\mathfrak{v}\right); \\ \hat{\eta}: \quad E'' \to E', \qquad & (\mathfrak{u}, \mathfrak{v}) \mapsto \left(\frac{\mathfrak{v}^2}{\mathfrak{u}^2}, \, \frac{-4\,t^2 - \mathfrak{u}^2}{\mathfrak{u}^2}\,\mathfrak{v}\right). \end{aligned}$$

One checks that $\hat{\varphi} \circ \varphi = [2]$ and $\hat{\eta} \circ \eta = [2]$ are the "multiplication-by-2" maps on E, and E', respectively. Moreover, one checks that $\phi = \hat{\eta} \circ \varphi$ and $\hat{\phi} = \hat{\varphi} \circ \eta$.

3.2. Complete Descent via 2-Isogeny. The previous proposition implies the following exact diagram as a refinement of the diagram in (2.43): (3.6)

The first (nontrivial) row in the diagram above comes from the explicit relation

$$(3.7) \quad E''(\mathbb{Q})[\hat{\varphi}] = \left\{ (-t^2, 0), \mathcal{O} \right\} \qquad \text{yet} \qquad (\mathfrak{u}, \mathfrak{v}) = \eta(U, V) \implies \mathfrak{u} = \frac{V^2}{4U^2} > 0;$$

while the second and fourth rows are related to a descent via 2-isogeny. We recall the main results for the latter:

Proposition 3.2. Fix $t \in \mathbb{Q}^{\times}$, denote E_t as in Theorem 2.6, and denote $E_t'' = E''$ as in Proposition 3.1.

(1) There is an exact sequence

$$(3.8) \ 0 \longrightarrow \frac{E_t(\mathbb{Q})}{\hat{\varphi}(E_t''(\mathbb{Q}))} \stackrel{\delta''}{\longrightarrow} S^{(\hat{\varphi})}(E_t''/\mathbb{Q}) \longrightarrow \coprod (E_t''/\mathbb{Q})[\hat{\varphi}] \longrightarrow 0$$

where δ'' sends $[m](0,0) \mapsto (t^2+4)^m$ and $(u,v) \mapsto u$ otherwise; while the second map sends $d \mapsto \{C''_{d,t}/\mathbb{Q}\}$ in terms of

(3.9)
$$C''_{d,t}: \quad dW^2 = d^2 + \left(t^2 + 2\right) dZ^2 + Z^4.$$

(2) The map $\psi': C'_{d,t} \to E_t$ in Theorem 2.6 factors as the composition

$$(3.10) C'_{d,t} \xrightarrow{\eta_*} C''_{d,t} \xrightarrow{\psi''} E_t$$

where η_* and ψ'' are the maps sending

$$(3.11) \ \ (z,w) \mapsto \left(-\frac{d\,z^2}{2\,t\,(w^2-d)}, \ \frac{d\,z^2\,(w^2+d)}{2\,(w^2-d)^2} \right), \qquad (Z,W) \mapsto \left(\frac{d}{Z^2}, \ -\frac{d\,W}{Z^3} \right).$$

In the Proposition above we identify, as in Proposition 2.5, the Selmer and Shafarevich-Tate groups as

$$(3.12) S^{(\hat{\varphi})}(E''/\mathbb{Q}) \simeq \left\{ d = \pm \prod_{p \in \Sigma} p^{a_p} \in \frac{\mathbb{Q}^{\times}}{(\mathbb{Q}^{\times})^2} \middle| C''_d(\mathbb{Q}_{\nu}) \neq \emptyset \text{ for all } \nu \in \Sigma \right\},$$
$$III(E''/\mathbb{Q})[\hat{\varphi}] \simeq \left\{ \{C''_d/\mathbb{Q}\} \in WC(E''/\mathbb{Q})[\hat{\varphi}] \middle| C''_d(\mathbb{Q}_{\nu}) \neq \emptyset \text{ for all } \nu \in \Sigma \right\}.$$

Hence by the canonical exact sequence

$$(3.13) 1 \longrightarrow \frac{(\mathbb{Q}^{\times})^2}{(\mathbb{Q}^{\times})^4} \longrightarrow \frac{\mathbb{Q}^{\times}}{(\mathbb{Q}^{\times})^4} \longrightarrow \frac{\mathbb{Q}^{\times}}{(\mathbb{Q}^{\times})^2} \longrightarrow 1$$

we may think of $S^{(\hat{\phi})}(E'/\mathbb{Q})$ as being a cover of $S^{(\hat{\varphi})}(E''/\mathbb{Q})$.

Proof. The first statement follows from Proposition 3.1 and the results from [7], so we focus on the second statement. Consider the diagram

$$(3.14) C'_{d,t} \xrightarrow{\theta} E'_t \xrightarrow{\hat{\phi}} E_t$$

$$\downarrow^{\eta}$$

$$C''_{d,t} \xrightarrow{\vartheta} E''_t \xrightarrow{\hat{\varphi}} E_t$$

where θ is the isomorphism in (2.42) and ϑ is the isomorphism that sends

(3.15)
$$\vartheta: (Z,W) \mapsto \left(2\frac{Z^2 + \sqrt{d}W + d}{Z^2}, 2\sqrt{d}\frac{(t^2 + 2)Z^2 + 2\sqrt{d}W + 2d}{Z^3}\right);$$

which has the inverse

(3.16)
$$\vartheta^{-1}: \quad (\mathfrak{u},\mathfrak{v}) \mapsto \left(\sqrt{d} \frac{2\mathfrak{v}}{\mathfrak{u}(\mathfrak{u}-4)}, \sqrt{d} \frac{-4t^2 + 2t^2\mathfrak{u} + \mathfrak{u}^2}{\mathfrak{u}(\mathfrak{u}-4)}\right).$$

Define the map $\eta_*: C'_{d,t} \to C''_{d,t}$ by $\eta_* = \vartheta^{-1} \circ \eta \circ \theta$. This gives the composition

(3.17)
$$\psi'' \circ \eta_* = (\hat{\varphi} \circ \vartheta) \circ (\vartheta^{-1} \circ \eta \circ \theta) = (\hat{\varphi} \circ \eta) \circ \theta = \hat{\phi} \circ \theta = \psi'.$$

Moreover, one checks that η_* maps (z, w) as above.

3.3. Modified Descent via 4-Isogeny. We present an effective version of the four-descent by using the two-descent. This is the main result of the paper.

Theorem 3.3. Fix $t \in \mathbb{Q}^{\times}$, and denote the elliptic curves

(3.18)
$$E_{t}: v^{2} = u^{3} + (t^{2} + 2) u^{2} + u,$$

$$E'_{t}: V^{2} = U^{3} - 2(t^{2} - 4) U^{2} + (t^{2} + 4)^{2} U,$$

$$E''_{t}: v^{2} = u^{3} + (t^{2} - 4) u^{2} - 4t^{2} u.$$

(1) There are isogenies $\phi: E_t \to E_t'$ of degree 4 with kernel generated by (-1,t), and $\varphi: E_t \to E_t''$ of degree 2 generated by [2](-1,t) = (0,0).

(2) Upon identifying the Selmer groups as subgroups of quotients of $\mathbb{Q}^{\times}/(\mathbb{Q}^{\times})^4$ and Shafarevich-Tate groups as a collection of homogeneous spaces, there is an exact diagram

$$(3.19) \qquad 0 \longrightarrow \frac{E_t(\mathbb{Q})}{\hat{\phi}(E'_t(\mathbb{Q}))} \xrightarrow{\delta'} S^{(\hat{\phi})}(E'_t/\mathbb{Q}) \longrightarrow \operatorname{III}(E'_t/\mathbb{Q})[\hat{\phi}] \longrightarrow 0$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \eta$$

$$0 \longrightarrow \frac{E_t(\mathbb{Q})}{\hat{\varphi}(E''_t(\mathbb{Q}))} \xrightarrow{\delta''} S^{(\hat{\varphi})}(E''_t/\mathbb{Q}) \longrightarrow \operatorname{III}(E''_t/\mathbb{Q})[\hat{\varphi}] \longrightarrow 0$$

where we have the connecting homomorphisms

(3.20)
$$\delta': \frac{(-1,t)\mapsto -4\,t^2}{(u,v)\mapsto u^2+2\,(t^2+1)\,u+1-2\,t\,v} \qquad \delta'': \qquad \frac{(0,0)\mapsto t^2+4}{(u,v)\mapsto u}$$

while the maps into the Shafarevich-Tate groups send $d \mapsto \{C'_{d,t}/\mathbb{Q}\}$ and $d \mapsto \{C''_{d,t}/\mathbb{Q}\}$ in terms of

(3.21)
$$C'_{d,t}: d\left(w - \frac{z^2}{4t^2}\right)z^2 = \left(w^2 - d\right)^2,$$
$$C''_{d,t}: dW^2 = d^2 + (t^2 + 2) dZ^2 + Z^4.$$

(3) The composition

$$(3.22) \psi': C'_{d,t} \xrightarrow{\eta} C''_{d,t} \xrightarrow{\psi''} E_t$$

in terms of the maps η and ψ'' sending

$$(3.23) \quad (z,w) \mapsto \left(-\frac{d\,z^2}{2\,t\,(w^2-d)}, \, \frac{d\,z^2\,(w^2+d)}{2\,(w^2-d)^2}\right), \qquad (Z,W) \mapsto \left(\frac{d}{Z^2}, \, -\frac{d\,W}{Z^3}\right);$$

$$satisfy \ the \ congruence \ \delta'(\psi'(P)) \equiv d \mod (\mathbb{Q}^\times)^4.$$

Note that we abuse notation and write η instead of η_* .

Proof. This follows directly from Theorem 2.6 and Proposition 3.2. \Box

4. Examples

In this section we assume that the elliptic curve E and its 4-isogenous curve E' have Weierstrass equations as in Proposition 2.1.

4.1. **Example:** $\mathbb{T} \simeq \mathbb{Z}/4\mathbb{Z}$. Consider t=8. Using the ideas in [2] as well as Proposition 2.2 it is easy to see that the torsion subgroup of both $E(\mathbb{Q})$ and $E'(\mathbb{Q})$ is $\mathbb{Z}/4\mathbb{Z}$. In fact, one may use either Cremona's mwrank or the package MAGMA so see that the Mordell-Weil groups are

(4.1)
$$E(\mathbb{Q}) \simeq \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z},$$
$$E'(\mathbb{Q}) \simeq \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z},$$
$$E''(\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}.$$

We consider the diagram in (3.6) in detail.

Performing a 2-descent with mwrank (recall that $\eta: E' \to E''$ and $\hat{\varphi}: E'' \to E$ are 2-isogenies) one computes the quantities

$$(4.2) \qquad \frac{E''(\mathbb{Q})}{\eta\left(E'(\mathbb{Q})\right)} \simeq S^{(\eta)}(E'/\mathbb{Q}) \simeq \{\pm 1, \pm 2\}, \qquad \qquad \mathrm{III}(E'/\mathbb{Q})[\eta] \simeq \{0\};$$

$$\frac{E(\mathbb{Q})}{\hat{\varphi}\left(E''(\mathbb{Q})\right)} \simeq S^{(\hat{\varphi})}(E''/\mathbb{Q}) \simeq \{\pm 1\}, \qquad \qquad \mathrm{III}(E''/\mathbb{Q})[\hat{\varphi}] \simeq \{0\}.$$

Considering the diagram in (3.6) we see that

$$(4.3) \qquad \frac{E(\mathbb{Q})}{\hat{\phi}\left(E'(\mathbb{Q})\right)} \simeq S^{(\hat{\phi})}(E'/\mathbb{Q}) \simeq \frac{\mathbb{Z}}{2\,\mathbb{Z}} \times \frac{\mathbb{Z}}{2\,\mathbb{Z}} \qquad \text{and} \qquad \mathrm{III}(E'/\mathbb{Q})[\hat{\phi}] \simeq \{0\}.$$

Using Theorem 2.6, we may compute generators explicitly by exploiting the homogeneous spaces. The curves E and E' both have conductor $2^4 \cdot 17$ so that they have good reduction away from $\Sigma = \{2, 17, \infty\}$. The homogeneous spaces of interest are of the form

(4.4)
$$C'_d: d(w+az^2)z^2 = (w^2-d)^2, a = -4^{-4}.$$

When d=-1 we find the solution (z,w)=(4,0); this can be predicted from the connecting homomorphism $\delta':(0,0)\mapsto a^{-1}\equiv -1$. When d=4 we find the solution (z,w)=(16,10); this maps to the point of infinite order

(4.5)
$$\left(\frac{w}{z^2}, \frac{w^2 - d}{z^4}\right) = \left(\frac{5}{128}, \frac{3}{2048}\right) \in E(\mathbb{Q}).$$

Hence $S^{(\hat{\phi})}(E'/\mathbb{Q}) \simeq \{\pm 1, \pm 4\}$. A brief search shows that $(-1/2, -3/4) \in E'(\mathbb{Q})$. In fact, one checks that

$$\phi(0,0) = \mathcal{O}, \qquad \phi\left(\frac{5}{128}, \frac{3}{2048}\right) = [2] \left(-\frac{1}{2}, -\frac{3}{4}\right);$$

$$\hat{\phi}(0,0) = [2] (0,0), \qquad \hat{\phi}\left(-\frac{1}{2}, -\frac{3}{4}\right) = [2] \left(\frac{5}{128}, \frac{3}{2048}\right)$$

This shows that

$$(4.7) \qquad \frac{E(\mathbb{Q})}{\hat{\phi}\left(E'(\mathbb{Q})\right)} \simeq \frac{\mathbb{Z}}{2\,\mathbb{Z}} \times \frac{\mathbb{Z}}{2\,\mathbb{Z}} \quad \text{and} \quad \frac{E'(\mathbb{Q})}{\phi\left(E(\mathbb{Q})\right)} \simeq \frac{\mathbb{Z}}{2\,\mathbb{Z}} \times \frac{\mathbb{Z}}{4\,\mathbb{Z}}.$$

4.2. **Example:** $\mathbb{T} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$. Consider t=3/2. One computes that the Mordell-Weil groups are

(4.8)
$$E(\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z},$$
$$E'(\mathbb{Q}) \simeq \mathbb{Z}/8\mathbb{Z},$$
$$E''(\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}.$$

We consider Proposition 2.2 in more detail because $\sqrt{t + \sqrt{t^2 + 4}}$ is rational. Performing a 2-descent with mwrank one computes the quantities

(4.9)
$$\frac{E''(\mathbb{Q})}{\eta(E'(\mathbb{Q}))} \simeq S^{(\eta)}(E'/\mathbb{Q}) \simeq \{\pm 1\}, \qquad \qquad \mathrm{III}(E'/\mathbb{Q})[\eta] \simeq \{0\};$$
$$\frac{E(\mathbb{Q})}{\hat{\varphi}(E''(\mathbb{Q}))} \simeq S^{(\hat{\varphi})}(E''/\mathbb{Q}) \simeq \{\pm 1\}, \qquad \qquad \mathrm{III}(E''/\mathbb{Q})[\hat{\varphi}] \simeq \{0\};$$

and so

$$(4.10) \qquad \frac{E(\mathbb{Q})}{\hat{\phi}\left(E'(\mathbb{Q})\right)} \simeq S^{(\hat{\phi})}(E'/\mathbb{Q}) \simeq \frac{\mathbb{Z}}{2\,\mathbb{Z}} \qquad \text{and} \qquad \mathrm{III}(E'/\mathbb{Q})[\hat{\phi}] \simeq \{0\}.$$

We now compute explicit points. The curves have good reduction away from $\Sigma = \{2, 3, 5, \infty\}$, and the homogeneous spaces of interest are of the form

(4.11)
$$C'_d: d(w+az^2)z^2 = (w^2-d)^2, \quad a = -3^{-2}.$$

When d = -9 we find the solution (z, w) = (3, -3); this maps to the point

$$\left(\frac{w}{z^2}, \frac{w^2 - d}{z^4}\right) = \left(-\frac{1}{3}, \frac{2}{9}\right) \in E(\mathbb{Q}).$$

(Note how we may predict $S^{(\hat{\phi})}(E'/\mathbb{Q}) \simeq \{1, -9\}$ from the connecting homomorphism $\delta': (0,0) \mapsto a^{-1} \equiv -9$.) In fact, the group of rational points $E(\mathbb{Q})$ is generated by (-1/3, 2/9) and (0,0), while the group $E'(\mathbb{Q})$ is generated by (25/64, 125/1024); with orders 2, 4, and 8, respectively. (Compare with the proof of Proposition 2.2 where $\gamma = \sqrt{t + \sqrt{t^2 + 4}} = 2$.) We have $\phi(0,0) = \mathcal{O}$ while (4.13)

$$\phi\left(-\frac{1}{3}, \frac{2}{9}\right) = [4] \left(\frac{25}{64}, -\frac{125}{1024}\right) \quad \text{and} \quad \hat{\phi}\left(\frac{25}{64}, \frac{125}{1024}\right) = \left(-\frac{1}{3}, \frac{2}{9}\right) \oplus (0, 0);$$

this shows that

(4.14)
$$\frac{E(\mathbb{Q})}{\hat{\phi}(E'(\mathbb{Q}))} \simeq \frac{\mathbb{Z}}{2\mathbb{Z}} \quad \text{and} \quad \frac{E'(\mathbb{Q})}{\phi(E(\mathbb{Q}))} \simeq \frac{\mathbb{Z}}{4\mathbb{Z}}.$$

4.3. **Example:** $\mathbb{T} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$. Consider the collection of elliptic curves studied in [2] as discussed in the introduction; each rational r corresponds to $t = (r^4 - 6\,r^2 + 1)\,/\,(2\,r^3 - 2\,r)$. Bounds on the rank can be computed by mwrank and by MAGMA, but these programs do not seem to be able to compute the rank exactly. For these five examples we will consider the non-negative integer R such that

(4.15)
$$E(\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z} \times \mathbb{Z}^{R},$$
$$E'(\mathbb{Q}) \simeq \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}^{R},$$
$$E''(\mathbb{Q}) \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}^{R}.$$

Various data about the curves is summarized in Table 1. We will see below that, assuming the conjectures of Birch and Swinnerton-Dyer, the lower bounds on R are sharp and the Shafarevich-Tate groups are non-trivial.

We focus on size of the Selmer groups. The program mwrank computes the groups $S^{(\eta)}(E'/\mathbb{Q})$ and $S^{(\hat{\varphi})}(E''/\mathbb{Q})$ associated with the 2-isogenies, so it suffices to consider the group $S^{(\hat{\phi})}(E'/\mathbb{Q})$ associated with the 4-isogeny. We may compute the size of the latter Selmer group using the exact diagram in (3.6); the data is collected in Table 2. Consider the image of the torsion subgroup of $E(\mathbb{Q})$ under the connecting homomorphism $\delta': (x,y) \mapsto x^2 - y \mod (\mathbb{Q}^{\times})^4$. Quite explicitly, the torsion subgroup is generated by the points

$$(4.16) \qquad P = \left(\frac{r^2}{(r^2 + 2r - 1)(r^2 - 2r - 1)}, \frac{2r^4}{(r^2 + 2r - 1)^2(r^2 - 2r - 1)^2}\right)$$

$$Q = \left(\frac{r(r+1)(r-1)^2}{(r^2 + 2r - 1)(r^2 - 2r - 1)^2}, \frac{r(r+1)^2(r-1)^3}{(r^2 + 2r - 1)^2(r^2 - 2r - 1)^3}\right)$$

r	t	Set of Bad Primes Σ	Rank Bounds
$\frac{15}{56}$	$-\frac{5651521}{4890480}$	$\{2,3,5,7,41,71,1231,3361,4591,\infty\}$	$2 \le R \le 4$
$\frac{24}{65}$	$-\frac{3580801}{11384880}$	$\{2,3,5,7,13,23,41,89,967,4801,\infty\}$	$2 \le R \le 4$

 $\{2,3,5,7,11,23,29,223,2441,3079,\infty\}$

 $\{2, 3, 5, 7, 11, 19, 23, 79, 113, 281, 7793, \infty\}$

 $\{2, 3, 5, 7, 17, 41, 97, 109, 233, 991, 11593, \infty\}$

11

69

7

88

12

97

4806319

 $\overline{1760880}$

57695201

9480240

80420641

21568920

Table 1. Data for $E_t(\mathbb{Q})$ where $t = (r^4 - 6r^2 + 1)/(2r^3 - 2r)$

TABLE 2. Selmer Group Computations via mwrank

r	$S^{(\eta)}(E'/\mathbb{Q})$	$S^{(\hat{\varphi})}(E''/\mathbb{Q})$	$\#S^{(\hat{\phi})}(E'/\mathbb{Q})$
$\frac{15}{56}$	$\langle -1, 2, 3, 5, 7, 41, 71 \rangle$	$\langle -1, 2, 3, 5, 7, 41, 71 \rangle$	2^{13}
$\frac{24}{65}$	$\langle -1, 2, 3, 5, 13, 41, 89 \rangle$	$\langle -1, 2, 3, 5, 13, 41, 89 \rangle$	2^{13}
$\frac{11}{69}$	$\langle -1, 2, 5, 23, 29, 33 \rangle$	$\langle -1,2,5,23,29,33\rangle$	2^{11}
$\frac{7}{88}$	$\langle -1, 2, 7, 11, 15, 57, 113, 843 \rangle$	$\langle -1,2,7,57,11\cdot 15\rangle$	2^{12}
$\frac{12}{97}$	$\langle -1, 2, 51, 291, 1635, 57965 \rangle$	$\langle -1, 2, 51, 291, 1635 \rangle$	2^{10}

having orders 2 and 8, respectively, and the connecting homomorphism δ' sends

$$P \mapsto -\frac{r^4}{(r^2 + 2r - 1)^2 (r^2 - 2r - 1)^2}, \qquad Q \mapsto \frac{r (r+1)^3 (r-1)^3}{(r^2 + 2r - 1)^2 (r^2 - 2r - 1)^4}.$$

For the $d \in S^{(\hat{\phi})}(E'/\mathbb{Q})$ above, consider the homogeneous spaces

$$(4.18) C'_d: d(w+az^2)z^2 = (w^2-d)^2, a = -\frac{r^2(r+1)^2(r-1)^2}{(r^2+2r-1)^2(r^2-2r-1)^2}.$$

Some generators (x,y) on $E(\mathbb{Q})$ can be found by $\mathtt{mwrank};$ these can be pulled back to a point (z,w) on $C'_d(\mathbb{Q})$ via the mapping $\psi':C'_d\to E$ in Theorem 2.6. We summarize these rational points in Table 3.

Rathbun [5] has remarked that the points in Table 3 correspond to the complete list generators on the elliptic curve if we assume the validity of the Birch and Swinnerton-Dyer conjectures. That is, we consider the order of the vanishing of the L-series in order to compute the rank, as well as the relationship with the residue and regulator of the elliptic curve to compute the generators; the information is summarized in Table 4. (For more information on the BSD conjectures, consult [6, Chapter 4].) Notice that the $\hat{\phi}$ -component of the Shafarevich-Tate group is not

Table 3. Points $(z,w) \in C'_d(\mathbb{Q})$ Corresponding to $d \in S^{(\hat{\phi})}(E'/\mathbb{Q})$

r	d	z	w
$\frac{15}{56}$	$-1 \cdot 5651521^2$	$\frac{5651521}{840}$	5651521
	$2\cdot 8733^2$	$\frac{86033006559349680248647}{367497397145772723536}$	$\frac{3354273204372689787801}{320119683924889132}$
	$41 \cdot 71 \cdot 8617^2$	$\frac{32245866415137}{37085066480}$	$\frac{1001789642807}{3730892}$
	$3 \cdot 5 \cdot 7 \cdot 56515210^2$	$\frac{37902870933931733579}{3188191909720998}$	$\frac{5724171826291316638240}{12960129714313}$
$\frac{24}{65}$	$-1 \cdot 6769^2$	$\frac{155687}{1560}$	6769
0.5	$2 \cdot 15^2$	$\frac{3893880282213551}{2376079357686320}$	$\frac{45660470553}{2504457868}$
	$41 \cdot 89 \cdot 130^2$	$\frac{83760182508961}{72984831998502}$	$\frac{26347553866400}{3333554033}$
	$3\cdot 5\cdot 13\cdot 527982^2$	$\frac{4690464211822333}{4980114669368}$	$\frac{128551710083251455}{15183276431}$
$\frac{11}{69}$	$-1 \cdot 4806319^2$	$\frac{9612638}{759}$	4806319
	$5\cdot 4683^2$	$\frac{68687104829}{107909560}$	$\frac{585885447}{42652}$
	$23 \cdot 2453^2$	$\frac{670894550748541}{5454390952560}$	$\frac{71568676933}{12390148}$
	$2 \cdot 29 \cdot 33 \cdot 215530^2$	$\frac{1001805034739772531298429}{834126406602556183512}$	$\frac{24018346018297942811880275}{3159569721979379483}$
$\frac{7}{88}$	$-1 \cdot 57695201^2$	$\frac{57695201}{616}$	57695201
	$2\cdot 6441^2$	$\frac{5596434497}{2776356}$	$\frac{41962889}{738}$
	$7 \cdot 7658655^2$	$\frac{15608602011772334}{143349694803}$	$\frac{271628289087723675}{2275391981}$
	$11 \cdot 15 \cdot 57 \cdot 135723^2$	$\frac{26233163006176556}{5388839579079}$	$\frac{5056981351335585}{954960053}$
$\frac{12}{97}$	$-1 \cdot 80420641^2$	$\frac{80420641}{1164}$	80420641
	$17 \cdot 97 \cdot 34^2$	$\frac{3173285824541959}{59077245130310}$	$\frac{1304349940207}{558755747}$
	$1635 \cdot 34779^2$	$\frac{115031523259990561}{28757077855714}$	$\frac{31422940575120}{79995877}$

r	Rank	$\#S^{(\hat{\phi})}(E'/\mathbb{Q})$	$\# \mathrm{III}(E'/\mathbb{Q})[\hat{\phi}]$
$\frac{15}{56}$	2	2^{13}	2^6
$\frac{24}{65}$	2	2^{13}	2^6
$\frac{11}{69}$	2	2^{11}	2^4
$\frac{7}{88}$	2	2^{12}	2^5
$\frac{12}{97}$	1	2^{10}	2^7

Table 4. Shafarevich-Tate Group Computations Assuming BSD

always of square order; we give an explanation. For the moment, let $\phi: E \to E'$ be any isogeny. The canonical alternating, non-degenerate, bilinear pairing

$$(4.19) \qquad \qquad \coprod (E'/\mathbb{Q}) \times \coprod (E'/\mathbb{Q}) \longrightarrow \mathbb{Q}/\mathbb{Z}$$

induces a non-degenerate, bilinear pairing

$$(4.20) \qquad \qquad \coprod (E'/\mathbb{Q})[\hat{\phi}] \times \frac{\coprod (E'/\mathbb{Q})}{\phi (\coprod (E/\mathbb{Q}))} \longrightarrow \mathbb{Q}/\mathbb{Z}.$$

(For more information on the Cassels pairing, consult [7, Appendix C, §17].) When ϕ is the "multiplication-by-m" map we have $\coprod (E/\mathbb{Q})/m \coprod (E/\mathbb{Q}) \simeq \coprod (E/\mathbb{Q})[m]$ – assuming that $\coprod (E/\mathbb{Q})$ is finite – so the pairing induces an alternating, non-degenerate, bilinear pairing

$$(4.21) \qquad \qquad \coprod (E/\mathbb{Q})[m] \times \coprod (E/\mathbb{Q})[m] \longrightarrow \mathbb{Q}/\mathbb{Z}.$$

Hence $\coprod (E/\mathbb{Q})[m]$ must have square order. In general, however, ϕ might not be self-dual; that is, upon considering the short exact sequence

$$(4.22) \ 0 \longrightarrow \frac{\mathrm{III}(E/\mathbb{Q})}{\mathrm{III}(E/\mathbb{Q})[\phi]} \stackrel{\phi}{\longrightarrow} \mathrm{III}(E'/\mathbb{Q}) \longrightarrow \frac{\mathrm{III}(E'/\mathbb{Q})}{\phi\left(\mathrm{III}(E/\mathbb{Q})\right)} \longrightarrow 0$$

we might have

$$(4.23) \qquad \# \frac{\mathrm{III}(E/\mathbb{Q})}{\mathrm{III}(E/\mathbb{Q})[\phi]} \neq \# \frac{\mathrm{III}(E'/\mathbb{Q})}{\mathrm{III}(E'/\mathbb{Q})[\hat{\phi}]} \implies \frac{\mathrm{III}(E'/\mathbb{Q})}{\phi \left(\mathrm{III}(E/\mathbb{Q})\right)} \not\simeq \mathrm{III}(E'/\mathbb{Q})[\hat{\phi}].$$

References

- [1] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. J. Symbolic Comput., 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [2] Garikai Campbell and Edray Herber Goins. Heron Triangles, Diophantine Problems, and Elliptic Curves. Submitted, 2003.
- [3] Daniel Sion Kubert. Universal bounds on the torsion of elliptic curves. Proc. London Math. Soc. (3), 33(2):193–237, 1976.
- [4] J. R. Merriman, S. Siksek, and N. P. Smart. Explicit 4-descents on an elliptic curve. Acta Arith., 77(4):385–404, 1996.
- [5] Randall L. Rathbun. Personal communication. October 2003.

- [6] Alice Silverberg. Open questions in arithmetic algebraic geometry. In *Arithmetic algebraic geometry (Park City, UT, 1999)*, volume 9 of *IAS/Park City Math. Ser.*, pages 83–142. Amer. Math. Soc., Providence, RI, 2001.
- [7] Joseph H. Silverman. The arithmetic of elliptic curves. Springer-Verlag, New York-Berlin, 1986.
- [8] Jacques Vélu. Isogénies entre courbes elliptiques. C. R. Acad. Sci. Paris Sér. A-B, 273:A238–A241, 1971.

Department of Mathematics, California Institute of Technology, Pasadena, CA 91125

 $E ext{-}mail\ address: goins@caltech.edu}$