L^p -estimates for Riesz transforms on forms in the Poincaré space

Joaquim Bruna*

Departament de Matemàtiques Universitat Autònoma de Barcelona

Abstract

Using hyperbolic form convolution with doubly isometry-invariant kernels, the explicit expression of the inverse of the de Rham laplacian Δ acting on *m*-forms in the Poincaré space \mathbb{H}^n is found. Also, by means of some estimates for hyperbolic singular integrals, L^p -estimates for the Riesz transforms $\nabla^i \Delta^{-1}$, $i \leq 2$, in a range of *p* depending on *m*, *n* are obtained. Finally, using these, it is shown that Δ defines topological isomorphisms in a scale of Sobolev spaces $H^s_{m,p}(\mathbb{H}^n)$ in case $m \neq \frac{n \pm 1}{2}, \frac{n}{2}$.

Mathematics Subject Classification(2000): 53C21, 58J05, 58J50, 58J70

Key words: Hodge-de Rham laplacian, Sobolev spaces, Riesz transforms, hyperbolic form convolution

1 Statement of results and preliminaries

1.1. The main object of study in this paper is the Hodge-de Rham Laplacian Δ acting on m-forms in the Poincaré hyperbolic space (\mathbb{H}^n, g) . The aim is to prove that Δ defines topological isomorphisms in a range $H^s_{m,p}(\mathbb{H}^n)$ of Sobolev spaces of forms defined as follows. For $0 \leq m \leq n, 1 \leq p < \infty$ and $s \in \mathbb{N}$, the Sobolev space $H^s_{m,p}(\mathbb{H}^n)$ is the completion of the space $\mathcal{D}_m(\mathbb{H}^n)$ of smooth m-forms with compact support with respect the norm

$$\|\eta\|_{p,s} = \sum_{i=0}^{s} \|\nabla^{(i)}\eta\|_{p}$$

Here $\nabla^{(i)}$ means the *i*-th covariant differential of η , and for a covariant tensor α

$$\|\alpha\|_p = \left(\int_{\mathbb{H}^n} |\alpha(x)|^p \, d\mu(x)\right)^{\frac{1}{p}}$$

 $|\alpha|$ being the pointwise norm of α with respect the metric g and $d\mu$ the volume-invariant measure on \mathbb{H}^n given by g. The space $H^s_{m,p}(\mathbb{H}^n)$ can be alternatively defined in terms of weak derivatives. The main result of this paper is:

^{*}supported by projects BFM2002-04072-C02-02 and 2001SGR00172

Theorem A. Δ is a topological isomorphism from $H^{s+2}_{m,p}(\mathbb{H}^n)$ to $H^s_{m,p}$ for $p \in (p_1, p_2)$ with

$$p_1 = \frac{2(n-1)}{n-2+|n-2m|}$$
 $\frac{1}{p_1} + \frac{1}{p_2} = 1$

in case $m \neq \frac{n\pm 1}{2}, \frac{n}{2}$.

In the exceptional case $m = \frac{n\pm 1}{2}$, Δ is one to one but is not a topological isomorphism for any p. For this case we obtain as well some weighted estimates. If $m = \frac{n}{2}$, Δ is known to have a non-trivial kernel. Of course, Sobolev spaces $H^s_{m,p}$ can be considered for non integer s as well, and the same results holds by interpolation.

Notice that the Hodge star operator * establishes an isometry from $H^s_{m,p}(\mathbb{H}^n)$ to $H^s_{n-m,p}(\mathbb{H}^n)$ which commutes with Δ , and this is why the range (p_1, p_2) depends only on |n - 2m|. Notice too that the range (p_1, p_2) always contains p = 2 in the non-critical case |n - 2m| > 1 and that for functions (m = 0), the range of p is $(1, \infty)$ (see comments below). We point out that the range (p_1, p_2) equals $|\frac{1}{p} - \frac{1}{2}| < \frac{\sqrt{\mu}}{n-1}$, where μ denotes the greatest lower bound for the spectrum of Δ in $H^0_{m,2}(\mathbb{H}^n)$, whose value ([D]) is $\mu = \frac{(n-1-2m)^2}{4}$ (for $m < \frac{n}{2}$).

For the Sobolev spaces for p = 2, $H^s_{m,2}(\mathbb{H}^n)$, another proof of the theorem, based in energy methods and valid for an arbitrary complete hyperbolic manifold, is given in [BG]. The motivation for the theorem, as with [BG], comes from mathematical physics, where most operators exhibit Δ as their principal part and results like the above become essential to establish existence and uniqueness theorems.

Our method of proof is simply to construct an explicit inverse L for Δ on $\mathcal{D}_m(\mathbb{H}^n)$ and show that there is a gain of two covariant derivatives

$$||L\eta||_{p,s+2} \le \operatorname{const} ||\eta||_{p,s}.$$

Thus $L\eta$ plays the role of the classical Riesz transform in the Euclidean setting. The most delicate part is of course

$$\|\nabla^{(2)}L\eta\|_p \le \operatorname{const} \|\eta\|_p, \qquad p_1$$

Riesz-type operators such as $\nabla \Delta^{-\frac{1}{2}}$, $\nabla^{(2)} \Delta^{-1}$ have extensively been studied in different contexts, for the case of *functions*. On symmetric spaces, they are bounded in L^p , 1 and of weaktype (1,1). This was shown in [AL] for the first order ones in some spaces, and later extended to $all symmetric spaces in [A]. The <math>L^p$ -boundedness holds as well for higher order Riesz transforms in symmetric spaces, but not generally the weak type (1,1) estimate. In more general contexts, this has been shown in [L1], [L2], [L3], among others. In case of *m*-forms, 0 < m < n, as far as we know, there are much less known results, and is for those that our result is new. In [P1], [P2] some aspects of harmonic analysis of forms are developed; in particular, the exact expression for the heat kernel is given, and it is very likely that from it one can get as well an explicit expression for Δ^{-1} . Strictly speaking, to prove the result an exact expression of Δ^{-1} is not needed, it is enough having estimates for the resolvent both local and at infinity. In [L3], estimates of this kind are obtained and applied to Sobolev-type inequalities for forms, and they might work for this purpose too. However, we feel that our approach, that we next describe, is more elementary and might be interesting in itself.

The de Rham Laplacian Δ is invariant by all isometries φ of \mathbb{H}^n . These form a a group that we denote here by $\operatorname{Iso}(\mathbb{H}^n)$. Denoting by $\varphi^*(\eta)(x) = \eta(\varphi(x))$ the pull-back of a form η by φ , this means that Δ and φ^* commute, for all $\varphi \in \operatorname{Iso}(\mathbb{H}^n)$. Therefore the inverse L of Δ should commute too with $\operatorname{Iso}(\mathbb{H}^n)$. Among all isometries of \mathbb{H}^n , the hyperbolic translations $\operatorname{Tr}(\mathbb{H}^n)$ constitute a (noncommutative) subgroup, in one to one correspondence with \mathbb{H}^n itself. In section 2 we do some harmonic analysis for forms in \mathbb{H}^n and introduce hyperbolic convolution of forms to describe all operators acting on *m*-forms and commuting with $\operatorname{Tr}(\mathbb{H}^n)$. In a second step (subsection 2.2) we characterize the hyperbolic convolution kernels k(x, y) corresponding to operators commuting with the full group $\operatorname{Iso}(\mathbb{H}^n)$.

Once the general expression of an operator commuting with $\operatorname{Iso}(\mathbb{H}^n)$ has been found, we look for our L among these. This corresponds to L having a kernel k(x, y) which is a fundamental solution of Δ in a certain sense, and having the best decay of infinity. This kernel turns out to be unique for $m \neq \frac{n\pm 1}{2}$, $\frac{n}{2}$, we call it the *Riesz kernel for m-forms in* \mathbb{H}^n , it is found in subsection 3.1 and estimated in subsection 3.2. Section 4 is devoted to the proof of the L^p -estimates. Here we use standard techniques in real analysis (Haussdorf-Young inequalities, Schur's lemma, etc.). For the second-order Riesz transform, to show its boundedness in the specified range (p_1, p_2) needs considering some notion of "hyperbolic singular integral". There exist some references dealing with this, e.g. [L4], [I], and giving some criteria for L^p -boundedness that might apply; however, as the singular integral arises locally we have found easier and more elementary to treat it with the classical Euclidean Calderón-Zygmund theory as a local model, and patch it in a suitable way to infinity.

1.2. We collect here several notations and known facts about \mathbb{H}^n . We will use both the unit ball model \mathbb{B}^n with metric $g = 4(1 - |x|^2)^{-2} \sum_i dx^i dx^i$ and the half-space model $\mathbb{R}^n_+ = \{x_n > 0\}$ with metric $g = x_n^{-2} \sum_i dx^i dx^i$. Both models are connected via the Cayley transform $\psi \colon \mathbb{R}^n_+ \to \mathbb{B}^n$ given in coordinates by

$$y_i = \frac{2x_i}{\sum_{i=1}^{n-1} x_i^2 + (x_n+1)^2}, \quad i = 1, \dots, n-1; \qquad \qquad y_n = \frac{\sum_{i=1}^n x_i^2 - 1}{\sum_{i=1}^{n-1} x_i^2 + (x_n+1)^2}.$$

We denote by $e \in \mathbb{H}^n$ the point $(0, 0, \dots, 1) \in \mathbb{R}^n_+$ or $0 \in \mathbb{B}^n$.

The metric g defines a pointwise inner product $(\alpha, \beta)(x)$ between forms at x, for every $x \in \mathbb{H}^n$, and a volume measure $d\mu$. In the ball model $d\mu$ is written $d\mu(x) = 2^n (1 - |x|^2)^{-n} dx^1 \dots dx^n$, and $d\mu(x) = x_n^{-n} dx^1 \dots dx^n$ in the half-space model. We denote by \langle , \rangle the pairing between forms that makes $H^s_{m,2}(\mathbb{H}^n)$ a Hilbert space

$$\langle \alpha, \beta \rangle = \int_{\mathbb{H}^n} (\alpha, \beta)(x) \, d\mu(x) \, .$$

We write $|\alpha|$ and $||\alpha||$ for the pointwise and global norms, respectively, of the form α . In terms of the Hodge star operator * the inner product can be written too

$$\langle \alpha,\beta\rangle = \int_{\mathbb{H}^n} \alpha\wedge\ast\beta\,.$$

The group $\operatorname{Tr}(\mathbb{H}^n)$ of hyperbolic translations is in one to one correspondence $x \mapsto T_x$ with \mathbb{H}^n through the equation $T_x(e) = x$. The equations of $z = T_x y$ are better described in the half-space model by

$$z_i = x_n y_i + x_i$$
, $i = 1, \dots, n-1$; $z_n = x_n y_n$.

It is easily checked that indeed $\operatorname{Tr}(\mathbb{H}^n)$ is a (non-commutative) group. The inverse transformation of T_x will be denoted S_x . Another explicit isometry φ_x mapping e to x, satisfying $\varphi_x^{-1} = \varphi_x$ is given in the ball model by

$$\varphi_x(y) = \frac{(|x|^2 - 1)y + (|y|^2 - 2xy + 1)x}{|x|^2|y|^2 - 2xy + 1}.$$
(1)

Since the isotropy group of 0 is the orthogonal group O(n), the general expression of $\varphi \in \text{Iso}(\mathbb{H}^n)$ is $\varphi = \varphi_x \circ U$, with $x = \varphi(0)$.

The hyperbolic (or geodesic) distance between $x, y \in \mathbb{H}^n$ is written d(x, y). We will rather use the *pseudohyperbolic distance* r = r(x, y), related to d by the formula $d(x, y) = 2 \operatorname{arctanh} r(x, y)$. The explicit expression of $r(x, y)^2$ in the \mathbb{R}^n_+ model and the \mathbb{B}^n model is respectively

$$r^{2} = \frac{|x - y|^{2}}{|x - y|^{2} + 4x_{n}y_{n}}, \quad x, y \in \mathbb{R}^{n}_{+},$$

$$r^{2} = |\varphi_{x}(y)|^{2} = \frac{|x - y|^{2}}{(1 - |x|^{2})(1 - |y|^{2}) + |x - y|^{2}}, \qquad x, y \in \mathbb{B}^{n}.$$
(2)

Associated to the group of translations we have the basis of orthonormal translation-invariant vector fields $X_i(x) = (T_x)_*(X_i(e))$, such that $X_i(e) = \frac{\partial}{\partial x_i}$. They satisfy $X_i(u \circ T_x) = (X_i u) \circ T_x$ for every smooth function u. We will denote by $w^i(x)$ the dual basis of X_i , which accordingly is orthonormal and translation invariant too: $T_x^* w^i = w^i$. Their expression in the \mathbb{R}^n_+ model is simply

$$X_i(x) = x_n \frac{\partial}{\partial x_i}, \qquad w^i(x) = x_n^{-1} dx^i, \quad i = 1, \dots, n$$

Because of their translation-invariance property, the (X_i, w^i) are more suitable than the (X_i, η^i) defined in the ball model \mathbb{B}^n by

$$Y_i(x) = \frac{(1-|x|^2)}{2} \frac{\partial}{\partial x_i}, \qquad \eta^i(x) = 2(1-|x|^2)^{-1} dx^i.$$

For an increasing multindex I of length |I| = m we write $w^I = w^{i_1} \wedge w^{i_2} \wedge \cdots \wedge w^{i_m}$, and similarly dx^I or η^I . The $\{w^I\}_I$ is an orthonormal translation-invariant basis of *m*-forms.

Recall that the de Rham Laplacian is defined $\Delta = d\delta + \delta d$, where δ is the adjoint of d with respecto to \langle , \rangle . Although strictly speaking not needed, the following expression of Δ in w^{I} coordinates will simplify the analysis at some points. If $\alpha = \sum_{I} \alpha_{I} w^{I}$, a computation shows that in case $n \notin J$

$$(\Delta \alpha)_J = \Delta \alpha_J + 2 \sum_{k \in J} X_k \alpha_{Jk} - p(n-p-1)\alpha_J.$$
(3.1)

Here Jk means the multiidex obtained replacing k by n. In case $n \in J$,

$$(\Delta \alpha)_J = \Delta \alpha_J - 2 \sum_{l \notin J} X_l \alpha_{lJ} - (1-p)(p-n)\alpha_J$$
(3.2)

where lJ means the multiindex obtained replacing n by l. For a function f

$$\Delta f = -\sum_{i=1}^{n} X_i^2 f + (n-1)X_n f.$$

In the ball model, with usual coordinates,

$$\Delta f = -\frac{1}{4}(1 - |x|^2)^2 \sum_{i,j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j} + (1 - \frac{n}{2})(1 - |x|^2) \sum x_i \frac{\partial f}{\partial x_i}$$
(3.3)

2 Translation invariant and isometry invariant operators on forms

2.1. We are interested in finding the general expression of an operator acting on *m*-forms, and isometry-invariant. In a first step we consider *translation-invariant operators* acting an *m*-forms; these are described by what we might call *hyperbolic convolution* as follows. Let k(x, y) be a double *m*-form in x, y and define

$$(C_k\alpha)(x) = \int_{\mathbb{H}^n} \alpha(y) \wedge *_y k(x, y) = \langle \alpha, k(x, \cdot) \rangle, \qquad \alpha \in \mathcal{D}_m(\mathbb{H}^n).$$

If T_z is a translation with inverse S_z

$$C_k(T_z^*\alpha)(x) = \int_{\mathbb{H}^n} (T_z^*\alpha)(y) \wedge *_y k(x,y) = \int_{\mathbb{H}^n} \alpha(T_z y) \wedge *_y k(x,y)$$
$$= \int_{\mathbb{H}^n} \alpha(y) \wedge *_y k(x, S_z y)$$
$$T_z^*(C_k\alpha)(x) = C_k \alpha(T_z x) = \int_{\mathbb{H}^n} \alpha(y) \wedge *_y k(T_z x, y).$$

Therefore C_k is translation invariant if k is doubly translation invariant in the sense that

$$k(x,y) = k(S_z x, S_z y) \qquad \forall S_z \,.$$

Using the translation-invariant basis of m-forms w^{I} we see that the general expression of k is

$$k(x,y) = \sum_{I,J} k_{I,J}(x,y) w^{I}(x) \otimes w^{J}(y)$$

with $k_I(x, y)$ doubly-invariant functions, that is, of the form $k_{I,J}(x, y) = a_{I,J}(S_y x)$ for some function (or distribution) $a_{I,J}$. If δ_0 denotes the Delta-mass at e and

$$\delta(x,y) = \sum_{I,J} \delta_0(S_y x) w^I(x) \otimes w^J(y)$$

then formally

$$\alpha(x) = \int_{\mathbb{H}^n} \alpha(y) \wedge *_y \delta(x, y) \,.$$

If P is an operator on m-forms commuting with the T_y, S_y , we will thus have

$$P\alpha(x) = \int_{\mathbb{H}^n} \alpha(y) \wedge *_y P_x(\delta(x,y))$$

and indeed $k(x,y) = P_x(\delta(x,y))$ is formally doubly-invariant. This shows, in loose terms, that the operator C_k of convolution with a doubly translation invariant kernel k gives the general translation-invariant operator acting on m-forms. If

$$k(x,y) = \sum_{I,J} a_{I,J}(S_y x) w^I(x) \otimes w^J(y)$$

and $\alpha(x) = \sum \alpha_I(x) w^I(x)$, then $C_k \alpha$ has in the basis $w^I(x)$ coefficients given by

$$(C_k\alpha)_I(x) = \sum_J \int_{\mathbb{H}^n} a_{I,J}(S_y x) \alpha_J(y) \, d\mu(y) \, .$$

Thus in the basis w^I everything reduces of course to convolution of functions. For a function convolution kernel $a(S_y x)$ and a test function $u \in \mathcal{D}(\mathbb{H}^n)$ we may think in

$$C_a u(x) = \int_{\mathbb{H}^n} u(y) a(S_y x) \, d\mu(y)$$

as an infinite linear combination of inverse translates $a(S_y x)$ of a(x). Since the vector fields X_i commute with translations, it follows that whenever everything makes sense

$$X_i(C_a u) = C_{X_i a} u \,. \tag{4}$$

We point out that this convolution is not commutative, $C_a u$ is in general different from $C_u a$. Correspondingly, $X_i C_a u - C_a X_i u$ is in general not zero; in fact one can easily show ([BG, lemma 3.1]) that these commutators are linear combinations of other convolution operators built from $a(S_y x)$. **2.2.** Let *P* be a generic translation-invariant operator acting on *m*-forms. We have seen in the previous subsection that we can associate to *P* a doubly-translation invariant kernel k(x, y) so that $P = C_k$. By the same argument as before, *P* will be isometry invariant if and only if $k(\varphi x, \varphi y) = k(x, y)$ $\forall \varphi \in \text{Iso}(\mathbb{H}^n)$ in which case we say that *k* is *doubly isometry-invariant*. Working in the ball model and since every $\varphi \in \text{Iso}(\mathbb{H}^n)$ is the composition of a translation with some $U \in O(n)$, the additional requirement on the kernel $k(x, y) = \sum a_{I,J}(S_y x) w^I(x) \otimes w^J(y)$ amounts to k(Ux, U0) = k(x, 0), that is,

$$\sum_{I,J} a_{I,J}(Ux)U^*w^I(x) \otimes U^*w^J(0) = \sum_{I,J} a_{I,J}(x)w^I(x) \otimes w^J(0), \forall U$$

Thus we are interested in describing those k(x,0) —which is a *m*-form at 0 whose coefficients are *m*-forms in *x*— that are doubly invariant by all $U \in O(n)$ in the sense above. Once the k(x,0)having this property are known, $k(x,y) = k(S_yx,0)$ defines the general doubly isometry invariant *m*-form. For m = 0 the k(x,0) are simply the radial functions a(|x|), and $a(|S_yx|) = a(|\varphi_yx|)$ is the general doubly isometry invariant function. For $m \neq 0$ their general expression is not so simple. We find it more convenient to use the usual basis dx^I so we look at k(x,0) in the form

$$k(x,0) = \sum_{|I|=|J|=m} b_{I,J}(x) \, dx^I \otimes dx^J(0) \tag{5}$$

and we must impose $\sum_{I,J} b_{I,J}(Ux) d(Ux)^I \otimes d(Ux)^J(0) = k(x,0) \ \forall U$. For instance

$$\gamma(x,0) = \sum_{i=1}^{n} dx^{i} \otimes dx^{i}(0)$$

is easily seen to be doubly O(n)-invariant, and so is

$$\gamma_m = \frac{1}{m!} \gamma \wedge \dots \wedge \gamma = \sum_{|I|=m} dx^I \otimes dx^I(0)$$

(here we use the symbol \wedge to denote as well the exterior product of double forms defined by $(\alpha_1 \otimes \beta_1) \wedge (\alpha_2 \otimes \beta_2) = (\alpha_1 \wedge \alpha_2) \otimes (\beta_1 \wedge \beta_2)$). Another doubly O(n)-invariant 1-form is

$$\tau(x,0) = \left(\sum_{i=1}^n x_i \, dx^i\right) \otimes \left(\sum_{i=1}^n x_i \, dx^i(0)\right).$$

Lemma 2.1. The double forms γ and τ generate all doubly O(n)-invariant k(x, 0). More precisely, their general expression in the ball model is

$$k(x,0) = A_1(|x|)\gamma_m + A_2(|x|)\tau \wedge \gamma_{m-1} \qquad 0 < m < n$$

$$k(x,0) = A(|x|)\gamma_m, \qquad m = 0, n$$
(6)

Proof. First we prove by induction the following statement S(n): if k(x,0) is a doubly invariant (p,q)-form $\sum_{|I|=p,|J|=q} c_{I,J} dx^I \otimes dx^J(0)$ with constant coefficients, then $k \equiv 0$ if $p \neq q$, or k is diagonal i.e. $k(x,0) = c \sum dx^I \otimes dx^I(0) = c\gamma_n$ if p = q. Of course S(1) is obvious: assuming S(n-1), let

i.e. $k(x,0) = c \sum_{|I|=p}^{n-1} dx^I \otimes dx^I(0) = c\gamma_p$ if p = q. Of course S(1) is obvious; assuming S(n-1), let us break k(x,0) in four pieces, depending on whether $i_1, j_1 = 1$ or not:

$$k = \sum_{i_1=j_1=1} c_{I,J} \, dx^I \otimes dx^J(0) + \sum_{i_1=1,j_1\neq 1} + \sum_{i_1\neq 1,j_1=1} + \sum_{i_1\neq 1,j_1\neq 1} \stackrel{\text{def}}{=} k_1 + k_2 + k_3 + k_4$$

We may write $k_1 = (dx^1 \otimes dx^1(0)) \wedge \widetilde{k_1}, k_2 = (dx^1 \otimes 1) \wedge \widetilde{k_2}, k_3 = (1 \otimes dx^1(0)) \wedge \widetilde{k_3}$, with $\widetilde{k_1}, \widetilde{k_2}, \widetilde{k_3}, k_4$ double forms in the $dx^2, \ldots, dx^n, dx^2(0), \ldots, dx^n(0)$ of bidegrees (p-1, q-1), (p-1, q), (p, q-1) and (p, q), respectively. Imposing that k is doubly invariant by U of the type

$$U = \begin{pmatrix} 1 & 0 & 0 & \cdots \\ 0 & & & \\ 0 & & U_1 & \\ \vdots & & & \\ 0 & & & \end{pmatrix}, \qquad U_1 \in O(n-1)$$
(7)

we see that $\tilde{k_1}$, $\tilde{k_2}$, $\tilde{k_3}$ and k_4 are O(n-1)-invariant. We apply the induction hypothesis: if p = q $\tilde{k_2} = \tilde{k_3} = 0$, and $\tilde{k_1}$, k_4 are diagonal, i.e.

$$k = c_1 \sum_{i_1=1} dx^I \otimes dx^I(0) + c_2 \sum_{i_1 \neq 1} dx^I \otimes dx^I(0) \,. \tag{8}$$

If we use now $U \in O(n)$ permuting the first two axes we see that $c_1 = c_2$ and hence k is diagonal, establishing S(n) in case p = q. If |p - q| > 1 everything is 0. Finally if |p - q| = 1, say p = q + 1, then $\tilde{k_2}$ is diagonal and all others are zero

$$k = c(dx^1 \otimes 1) \wedge \sum_{|J|=q} dx'^J \otimes dx'^J(0)$$

where $x' = (x_2, \dots, x_n)$. If we impose the invariance under the permutation of the first two axes as before, it is clear that k must be zero.

Having proved that S(n) holds for all n, let now k(x,0) be as in (5) doubly O(n)-invariant. Clearly k(x,0) is then determined by its values $k(\vec{r},0)$, where $\vec{r} = (r,0,0,\ldots,0)$. Fixed $r, k(\vec{r},0)$ may be regarded as a double (m,m)-form with constant coefficients, which is invariant by all $U \in O(n)$ fixing \vec{r} , that is, of type (7). We write now the decomposition of $k(\vec{r},0)$ in terms of $\tilde{k}_1(r,0), \tilde{k}_2(r,0), \tilde{k}_3(r,0)$ and $k_4(r,0)$ as before, and applying S(n) we get (8)

$$k(\vec{r},0) = c_1(r) \sum_{i_1=1} dx^I(\vec{r}) \otimes dx^I(0) + c_2(r) \sum_{i_1\neq 1} dx^I(\vec{r}) \otimes dx^I(0)$$

(if m = n the last term is zero and the first is γ_m) which we write

$$= (c_1(r) - c_2(r)) \sum_{\substack{i_1=1\\|I|=m}} dx^I(\vec{r}) \otimes dx^I(0) + c_2(r) \sum_{\substack{|I|=m\\|I|=m}} dx^I(\vec{r}) \otimes dx^I(0)$$

$$= (c_1(r) - c_2(r)) dx^1(\vec{r}) \otimes dx^1(0) \wedge \sum_{\substack{|I|=m-1\\|I|=m-1}} dx^I(\vec{r}) \otimes dx^I(0) + c_2(r) \sum_{\substack{|I|=m\\|I|=m}} dx^I(\vec{r}) \otimes dx^I(0)$$

$$= (c_1(r) - c_2(r)) r^{-2} \tau(\vec{r}, 0) \gamma_{m-1}(\vec{r}, 0) + c_2(r) \gamma_m(\vec{r}, 0).$$

Finally, fixed x we choose U such that $Ux = \overrightarrow{r}$, r = |x|, and use the invariance of k, τ , γ to find (6) with $A_1(r) = c_2(r)$, $A_2(r) = r^{-2}(c_1(r) - c_2(r))$.

To find the general expression of a doubly isometry invariant kernel k(x, y) we must translate k(x, 0) to an arbitrary point: $k(x, y) = k(S_y x, S_y y)$. We may use any isometry mapping y to 0, for instance we may use φ_y given by (1) instead of S_y . We introduce the basic forms α , β , τ and γ

$$\begin{aligned} \alpha &= \alpha(x,y) = \sum_{i} \varphi_{y}^{i}(x) \, d\varphi_{y}^{i}(x) \,, \qquad \beta = \sum_{i} \varphi_{y}^{i}(x) \, d\varphi_{y}^{i}(y) = -\sum_{i} \varphi_{y}^{i}(x) \frac{dy^{i}}{1 - |y|^{2}} \\ \tau &= \alpha \otimes \beta; \gamma(x,y) = \sum_{i=1}^{n} d\varphi_{y}^{i}(x) \otimes d\varphi_{y}^{i}(y) = \frac{-1}{1 - |y|^{2}} \sum_{i=1}^{n} d\varphi_{y}^{i}(x) \otimes dy^{i} = d_{x}\beta \,. \end{aligned}$$

The lemma gives part (a) of the following theorem. Part (b) gives other equivalent general expressions, which are intrinsic, that is, independent of the model of \mathbb{H}^n at use.

Theorem 2.2. (a) The general expression of an (m, m)-form k(x, y) doubly isometry-invariant in \mathbb{H}^n , in the ball model, is

$$\begin{split} k(x,y) &= A_1(|\varphi_y x|)\gamma_m(x,y) + A_2(|\varphi_y x|)\tau(x,y) \wedge \gamma_{m-1}(x,y), \qquad 0 < m < n \\ k(x,y) &= A(|\varphi_y x|)\gamma_m(x,y) \qquad m = 0, n \,. \end{split}$$

(b) Another equivalent expression for 0 < m < n is

$$k(x,y) = B_1(D)(d_x d_y D)^m + B_2(D)(d_x D \otimes d_y D) \wedge (d_x d_y D)^{m-1} =$$

= $(C_1(D)d_x d_y D + C_2(D)d_x D \otimes d_y D)^m$

where D denotes an arbitrary function of the geodesic distance d(x, y). (c) All such k(x, y) are symmetric in $x, y \in \mathbb{H}^n$

Proof. Part (a) has been already proved. For (b) note first that it is enough to consider *one* function of d: we choose $D = r(x, y)^2$ which in the ball model equals $|\varphi_y(x)|^2$. Then $d_x D = 2\alpha$, and using (1), (2) one finds

$$d_y D = 2(1-D) \sum_i \varphi_y^i(x) \frac{d_y^i}{1-|y|^2} = -2(1-D)\beta.$$

This gives $\tau = \alpha \otimes \beta = \frac{-1}{4} \frac{1}{1-D} d_x D \otimes d_y D$, and

$$d_x d_y D = +2d_x D \otimes \beta - 2(1-D)d_x \beta = +4\tau - 2(1-D)\gamma.$$

Therefore $(d_x d_y D)^{m-1}$ and $2^{m-1}(1-D)^{m-1}\gamma_{m-1}$ differ in a term containing τ , and so (b) follows. Part (c) is a consequence of (b).

We will need the expression of the generators τ , γ in terms of the invariant basis w^i . We obtain these using formula (2) for $r^2(x, y)$ in the half-space model. First

$$\alpha = \frac{d_x r^2}{2} = \frac{1 - r^2}{2(|x - y|^2 + 4x_n y_n)} \left(2\sum_{i=1}^{n-1} x_n (x_i - y_i) w^i(x) + (2x_n (x_n - y_n) - |x - y|^2) w^n(x) \right)$$

$$\beta = \frac{d_y r^2}{2(r^2 - 1)} = \frac{-1}{2(|x - y|^2 + 4x_n y_n)} \left(2\sum_{j=1}^n y_n (y_j - x_j) w^j (y) + (2y_n (y_n - x_n) - |x - y|^2) w^n (y) \right)$$

In the following we write $w^{ij} = w^i(x) \otimes w^j(y)$. We have

$$\tau = \alpha \otimes \beta = \frac{1}{4} \frac{1 - r^2}{(|x - y|^2 + 4x_n y_n)^2} \sum_{ij} P_{i,j}(x, y) w^{i,j}$$

where the $P_{ij}(x, y)$ are certain homogeneous polynomials. As we know, everything can be written in terms of $z = S_y x$: for instance

$$1 - r^{2} = \frac{4x_{n}y_{n}}{|x - y|^{2} + 4x_{n}y_{n}} = \frac{4z_{n}}{|z|^{2} + 2z_{n} + 1}$$

and say for i, j < n

$$\frac{P_{ij}}{(|x-y|^2+4x_ny_n)} = \frac{x_ny_n(x_i-y_i)(x_j-y_j)}{(|x-y|^2+4x_ny_n)^2} = \frac{z_nz_iz_j}{(|z|^2+2z_n+1)^2}$$

Therefore we may write

$$\tau = \frac{1 - r^2}{(|z|^2 + 2z_n + 1)^2} \sum_{i,j} p_{i,j}(z) w^{i,j} \,. \tag{9}$$

For $\gamma = d_x \beta$ we obtain a similar expression

$$\frac{4}{1-r^2}\gamma = \sum_{i,j=1}^{n-1} \left(\delta_{ij} - \frac{2(x_i - y_i)(x_j - y_j)}{|x - y|^2 + 4x_n y_n} \right) w^{i,j} + \left(\frac{2\sum_{i=1}^{n-1} |x_i - y_i|^2}{|1 - \frac{2\sum_{i=1}^{n-1} |x_i - y_i|^2}{|x - y|^2 + 4x_n y_n}} \right) w^{n,n} + \sum_{i=1}^{n-1} \frac{2(x_i - y_i)(x_n - y_n)}{|x - y|^2 + 4x_n y_n} (w^{i,n} - w^{n,i}).$$

Again this can be written

$$\gamma = \frac{1 - r^2}{4(|x - y|^2 + 4x_n y_n)} \sum_{i,j} Q_{ij}(x, y) w^{i,j} = \frac{1 - r^2}{(|z|^2 + 2z_n + 1)} \sum_{i,j} q_{ij}(z) w^{i,j}.$$
 (10)

Notice that

$$\frac{p_{ij}(z)}{(|z|^2 + 2z_n + 1)^2} = O(1), \qquad \frac{q_{ij}(z)}{(|z|^2 + 2z_n + 1)} = O(1)$$

and hence

$$|\tau(x,y)| = O(1-r^2), \qquad |\gamma(x,y)| = O(1-r^2).$$
 (11)

3 Riesz forms and Riesz form-potentials in \mathbb{H}^n

3.1. Our next objective is now to find an explicit left-inverse L for Δ on $\mathcal{D}_m(\mathbb{H}^n)$. Since Δ is invariant by all isometries, L should be too. By what has been discussed in section 2, L should have a kernel $k_m(x, y)$,

$$L\eta(x) = \int_{\mathbb{H}^n} \eta(y) \wedge *_y k_m(x,y)$$

doubly invariant by all isometries. Alternatively, notice that if k is some kernel such that

$$\eta(x) = \int_{\mathbb{H}^n} \Delta \eta(y) \wedge *_y k(x, y), \qquad \eta \in \mathcal{D}_m(\mathbb{H}^n)$$
(12)

(which formally exists because $\Delta \eta = 0, \eta \in \mathcal{D}_m(\mathbb{H}^n)$ imply $\eta = 0$) then its average over the unitary group O(n) with respect the normalized left-invariant measure $d\mu(U)$,

$$k_1(x,y) = \int_{O(n)} k_0(Ux,Uy) d\mu(U)$$

still satisfies (12) and it is doubly invariant by O(n). If φ_x is an isometry mapping x to 0, $k_2(x, y) = k_1(\varphi_x x, \varphi_x y)$ is independent of φ_x , satisfies (12) and is doubly invariant by all isometries.

Anyway, we look for a doubly isometry-invariant kernel k_m for which (12) holds, and then consider the operator L defined by k_m as above. Taking for granted by now that this operator Lis well defined on $\mathcal{D}_m(\mathbb{H}^n)$ and maps $\mathcal{D}_m(\mathbb{H}^n)$ into locally integrable *m*-forms, notice that (12) and the symmetry of k_m together imply that L is a right-inverse too, that is, $\Delta L\alpha = \alpha$ for $\alpha \in \mathcal{D}_m(\mathbb{H}^n)$ in the weak sense:

$$\begin{split} \langle \Delta L\alpha, \eta \rangle &= \langle L\alpha, \Delta \eta \rangle = \int_x L\alpha(x) \wedge * \Delta \eta(x) = \\ &= \int_x \left\{ \int_y \alpha(y) \wedge *_y k_m(x, y) \right\} \wedge * \Delta \eta(x) \\ &= \int_y \alpha(y) \wedge *_y \left\{ \int_x k_m(x, y) \wedge * \Delta \eta(x) \right\} = \langle \alpha, \eta \rangle \end{split}$$

We work in the ball model. By Theorem 2.2, $k_m(x, y)$ is of type

$$k_m(x,y) = A(|\varphi_x y|)\gamma_m, \quad m = 0, n$$

$$k_m(x,y) = A_1(|\varphi_x y)\gamma_m + A_2(|\varphi_x y|)\tau \wedge \gamma_{m-1}, \qquad 0 < m < m$$

where $\gamma = \sum_{i} d\varphi_x^i(x) \otimes d\varphi_x^i(y), \tau = \alpha \otimes \beta$ with $\alpha = \sum_{i} \varphi_x^i(y) d\varphi_x^i(x), \beta = \sum_{i} \varphi_x^i(y) d\varphi_x^i(y)$ (notice that we are exchanging x, y, using (c) in Theorem 2.2). Condition (12) implies $\Delta_y k_m(x, y) = 0$ in $y \neq x$ (while $\Delta Lw = w$ implies $\Delta_x k_m(x, y) = 0$ in $x \neq y$). In fact, (12) amounts to requiring $\Delta_y k_m(x, y) = \delta_x$ in a sense to be described below.

3.2. In a first step we look for conditions on the A_1 , A_2 , so that $\Delta_y k_m(x, y) = 0$ in $y \neq x$. A lengthy computation will show that the general harmonic k_m depends on four parameters. By the invariance of k_m , we may assume x = 0, in which case, writing r = |y|,

$$k_m(x,y) = A(r)\gamma_m$$
, $m = 0, n$
 $k_m(0,y) = A_1(r)\gamma_m + A_2(r)\tau \wedge \gamma_{m-1}$

with $\gamma = \sum dx^i(0) \otimes dy^i$, $\tau = \alpha \otimes \beta$, $\alpha = \sum y^i dx^i(0)$, $\beta = rdr$. Since $*_x *_y k_m(x, y)$ is again doubly invariant, it must have an analogous expression with m replaced by n - m. Indeed, it is easily checked that

$$*_{x} *_{y} \gamma_{m} = \frac{m!}{(n-m)!} (1-r^{2})^{2m-n} \gamma_{n-m}$$
$$*_{x} *_{y} (\tau \wedge \gamma_{m-1}) = (m-1)! (1-r^{2})^{2m-n} \left(r^{2} \frac{\gamma_{n-m}}{(n-m)!} - \frac{\tau \wedge \gamma_{n-m-1}}{(n-m-1)!} \right)$$

whence $*_x *_y k_m(x, y) = \frac{m!}{(n-m)!} (1-r^2)^{2m-n} \gamma_{n-m}$ for m = 0, n and for 0 < m < n,

$$*_{x} *_{y} k_{m}(0, y) = \frac{(m-1)!(1-r^{2})^{2m-n}}{(n-m)!} \left[(mA_{1}+r^{2}A_{2})\gamma_{n-m} - (n-m)A_{2}\tau \wedge \gamma_{n-m-1} \right].$$
(13)

Moreover, since * commutes with Δ , it is natural to require as well that $*_x *_y k_m = k_{n-m}$, that is, we may assume from now on that $0 \le m \le n/2$. For m = 0, using (3.3) we find

$$\Delta(A(r)) = \frac{1}{4}(1-r^2) \left[-1(1-r^2)A'' + ((3-n)r + r^{-1}(1-n))A' \right]$$

from which it follows that $A'(r) = c_0(1-r^2)^{n-2}r^{1-n}$ and

$$A(r) = c_1 - c_0 \int_r^1 (1 - s^2)^{n-2} s^{1-n} \, ds \, .$$

We start now computing $\Delta_y k_m(0, y)$ for $0 < m \le n/2$ using that on *m*-forms Δ equals $(-1)^{m+1}(*d*d+(-1)^n d*d*)$. The double form $\Delta_y k_m(x, y)$ is also doubly invariant, and therefore it must have the same expression as k_m with A_1 , A_2 replaced by other functions B_1 , B_2 to be found. In the computations we will use besides (13) the equations

$$d_y \alpha = \gamma, \quad d_y (\tau \wedge \gamma_{m-1}) = -r dr \wedge \gamma_m = -\beta \wedge \gamma_m$$
$$*_x *_y dr \wedge \gamma_m = (-1)^m \frac{m!}{(n-m-1)!} (1-r^2)^{2m+2-n} r^{-1} \alpha \wedge \gamma_{n-m-1}$$

which are easily checked as well. First, $d_y k_m(0,y) = (A'_1 - rA_2)dr \wedge \gamma_m$, so by the equations above

$$*_{x} *_{y} d_{y}k_{m}(0, y) = (-1)^{m} \frac{m!}{(n-m-1)!!} (1-r^{2})^{2m+2-n} (A'_{1}-rA_{2})r^{-1}\alpha \wedge \gamma_{n-m-1} = (14)$$

$$\stackrel{\text{def}}{=} \frac{(-1)^{m}m!}{(n-m-1)!} A_{3}\alpha \wedge \gamma_{n-m-1} ,$$

$$*_{x}d_{y} *_{y} d_{y}k_{m}(0, y) = \frac{(-1)^{m}m!}{(n-m-1)!} (A_{3}\gamma_{n-m} + A'_{3}r^{-1}\tau \wedge \gamma_{n-m-1})$$

$$*_{y} d_{y} *_{y} d_{y}k_{m}(0, y) = (-1)^{m(n-m-1)} *_{y} *_{x}(A_{3}\gamma_{n-m} + A'_{3}r^{-1}\tau \wedge \gamma_{n-m-1}) =$$

$$= (-1)^{m(n-m-1)} \frac{m!}{(n-m-1)!} (1-r^{2})^{n-2n} \left(A_{3}\frac{(n-m)!}{m!} \gamma_{m} + A'_{3}r\frac{(n-m-1)!}{m!} \gamma_{m} \right)$$

$$-A'_{3}r^{-1}\frac{(n-m-1)!}{(m-1)!}\tau \wedge \gamma_{m-1}\Big)$$

= $(-1)^{m(n-m+1)}(1-r^{2})^{n-2m}\left[((n-m)A_{3}+A'_{3}r)\gamma_{m}-mA'_{3}r^{-1}\tau \wedge \gamma_{m-1}\right].$

By the analogous computation, applying d_y to (13)

$$*_{x}d_{y}*_{y}k_{m}(0,y) = \frac{(m-1)!}{(n-m)!} \left[\left[(mA_{1}+r^{2}A_{2})(1-r^{2})^{n-2m} \right]' + (n-m)rA_{2}(1-r^{2})^{n-2m} \right] dr \wedge \gamma_{n-m}$$

$$*_{y} d_{y} *_{y} k_{m}(0, y) = (-1)^{(m+1)(n-m)} (1-r^{2})^{2m+2-n} r^{-1}$$

$$= \left[\left[(mA_{1}+r^{2}A_{2})(1-r^{2})^{n-2m} \right]' + (n-m)rA_{2}(1-r^{2})^{n-2m} \right] \alpha \wedge \gamma_{m-1} = (15)$$

$$\stackrel{\text{def}}{=} (-1)^{(m+1)(n-m)} A_{4}\alpha \wedge \gamma_{m-1}$$

$$d_{y} *_{y} d_{y} *_{y} k_{m}(0, y) = (-1)^{(n-m)(m+1)} (A_{4}'r^{-1}\tau \wedge \gamma_{m-1} + A_{4}\gamma_{m})$$

It follows finally that $\Delta = (-1)^{nm+1} (*d * d + (-1)^n d * d *)$ on k_m equals

$$\Delta_y k_m(0,y) = B_1 \gamma_m + B_2 \tau \wedge \gamma_{m-1}$$

with

$$B_1 = -A_4 - (1 - r^2)^{n-2m} ((n - m)A_3 + A'_3 r)$$

$$B_2 = -A_4 r^{-1} + m(1 - r^2)^{n-2m} A'_3 r^{-1}.$$

Therefore, $\Delta_y k(0, y) = 0$ is equivalent to the system $B_1 = 0$, $B_2 = 0$. It easily follows from this that A_3 satisfies the equation

$$r(1-r^2)A_3'' + \left[(n+1) - r^2(3n+1-4m) \right] A_3' - 2(n-2m)(n-m)rA_3 = 0.$$

Replacing in the equation $B_1 = 0$, A_4 by its expression in terms of A_1 and A_2 , and then A_2 by its expression in terms of A_1 and A_3 , we find that A_1 satisfies the inhomogeneous equation

$$r(1-r^2)A_1'' + \left[(n+1) + (n-1-4m)r^2\right]A_1' + 2m(n-2m)rA_1 =$$

= 2rA_3(1+r^2)(1-r^2)^{n-2m-2}

The change of variables $A_1(r) = G(x)$, $A_3(r) = H(x)$, $x = r^2$, transforms these into the hypergeometric equations

$$x(1-x)H''(x) + \left[\frac{n}{2} + 1 - (\frac{3}{2}n + 1 - 2m)x\right]H'(x) - (\frac{n}{2} - m)(n-m)H = 0$$

$$x(1-x)G''(x) + \left[\frac{n}{2} + 1 - (2m + 1 - \frac{n}{2})x\right]G'(x) - m(m - \frac{n}{2})G = (16)$$

$$= \frac{1}{2}(1+x)(1-x)^{n-2m-2}H(x) \stackrel{\text{def}}{=} f(x)$$

This system is equivalent to $\Delta_y k_m(x, y) = 0$ in $y \neq x$, whence the general doubly-invariant k_m harmonic in $y \neq x$ depends on four parameters. Note that for $m = \frac{n}{2}$ the homogeneous equations are the same and can be solved explicitly: the general solution is $H = as^{-\frac{n}{2}} + b$ and

$$G(x) = cx^{-\frac{n}{2}} + d + \frac{1}{2} \int_{1/2}^{x} t^{-\frac{n}{2}-1} \left\{ \int_{0}^{t} s^{n/2} (1+s)(1-s)^{-3} (as^{-\frac{n}{2}} + b) \, ds \right\} \, dt \,. \tag{17}$$

For $m < \frac{n}{2}$ a fundamental family for the first equation in (16) is given by

$$u_1(x) = x^{-\frac{n}{2}}F(-m, \frac{n}{2} - m, 1 - \frac{n}{2}, x), u_2(x) = F(\frac{n}{2} - m, n - m, \frac{n}{2} + 1, x)$$

The hypergeometric function in u_1 is a polynomial in x of degree m with positive coefficients, 1 + x if m = 1. A fundamental family for the second is given by

$$u_{3}(x) = x^{-\frac{n}{2}}F(m-n, m-\frac{n}{2}, 1-\frac{n}{2}, x) = x^{-\frac{n}{2}}(1-x)^{n+1-2m}F(\frac{n}{2}+1-m, 1-m, 1-\frac{n}{2}, x),$$
$$u_{4}(x) = F(m, m-\frac{n}{2}, 1+\frac{n}{2}, x)$$

The hypergeometric function in u_3 is a polynomial of degree m-1 with positive coefficients (see [E] for all these facts). The wronskian w(x) for this second equation is, by Liouville's formula

$$W(x) = W(x_0) \exp -\int_{x_0}^x \frac{\frac{n}{2} + 1 - (2m + 1 - \frac{n}{2})t}{t(1-t)} dt = c_{mn} x^{-\frac{n}{2} - 1} (1 - x^{n-2m})$$

It follows from this that the parametrization for G is given by

$$G(x) = c(x)u_3(x) + d(x)u_4(x)$$
(18)

where c(x), d(x) satisfy, with $H(x) = au_1(x) + bu_2(x)$,

$$c'(x) = \frac{u_4(x)f(x)}{x(1-x)W(x)} = \frac{1}{2}c_{mn}^{-1}H(x)(1+x)x^{\frac{n}{2}}(1-x)^{-3}u_4(x)$$

$$d'(x) = -\frac{u_3(x)f(x)}{x(1-x)W(x)} = -\frac{1}{2}c_{mn}^{-1}H(x)(1+x)x^{\frac{n}{2}}(1-x)^{-3}u_3(x).$$

Once $A_1(r) = G(r^2)$ and $A_3(r) = H(r^2)$ are known, the kernel $k_m(x, y)$ is completely known, because by the definition of A_3 in (14), $A_2(r) = -(1 - r^2)^{n-2m-2}A_3(r) + r^{-1}A'_1(r) = -(1 - x)^{n-2m-2}H(x) + 2G'(x)$.

The choice a = 0, c(0) = 0 (a = c = 0 in the parametrization (17) for $m = \frac{n}{2}$) gives all doubly invariant $k_m(x, y)$ which are globally harmonic, with no singularity, and they are therefore spanned by the forms corresponding to the choice $G = u_4$ and to the choice a = 0, b = 1, c(0) = 0, d(0) = 0,

$$G(x) = \left\{ \int_0^x (1+t)(1-t)^{-3} t^{n/2} u_2(t) u_4(t) dt \right\} u_3(x) \\ - \left\{ \int_0^x (1+t)(1-t)^{-3} t^{n/2} u_2(t) u_3(t) dt \right\} u_4(x) \,.$$

As a particular case, note that for $m = \frac{n}{2}$, γ_m is harmonic in \mathbb{H}^{2m} , and it is the simplest example of a non-zero harmonic *m*-form in $L^2(\mathbb{H}^{2m})$.

3.3. Besides being harmonic in $y \neq x$, the singularity at y = x most be such that (12) holds. Again, we may assume x = 0; we check this property using *second's Green identity*, whose version for general forms we recall now.

The operator δ being the adjoint of d, one has for a smooth domain $\overline{\Omega} \subset \mathbb{B}^n$ and α, β smooth forms on $\overline{\Omega}$ with degree $\alpha = \text{degree}\beta - 1$

$$\int_{\partial\Omega} \alpha \wedge \ast \beta = \int_{\Omega} d\alpha \wedge \ast \beta - \int_{\Omega} \alpha \wedge \ast \delta \beta \,.$$

Given two *m*-forms η , ω , applying this with $\alpha = \delta \eta$, $\beta = \omega$, next with $\alpha = \omega$, $\beta = d\eta$ and substracting one gets the first Green's identity for *m*-forms

$$\int_{\partial\Omega} (\delta\eta \wedge *\omega - \omega \wedge *d\eta) = \int_{\Omega} (\Delta\eta \wedge *\omega - \delta\eta \wedge *\delta\omega - d\eta \wedge *d\omega)$$

Permuting ω, η and substracting again gives the second Green's identity

$$\int_{\partial\Omega} (\delta\eta \wedge *\omega - \omega \wedge *d\eta - \delta\omega \wedge *\eta + \eta \wedge *d\omega) = \int_{\Omega} (\Delta\eta \wedge *\omega - \Delta\omega \wedge *\eta) \, .$$

We apply this to $\Omega = B(0, R) - B(0, \varepsilon)$ $0 < \varepsilon < R < 1, \eta \in \mathcal{D}_m(\mathbb{H}^n)$ and our $k_m(0, y)$ to get

$$\int_{|y|\geq\varepsilon} \Delta\eta \wedge *_y k_m(0,y) = \int_{|y|=\varepsilon} (k_m \wedge *d\eta + \delta_y k_m \wedge *\eta - \delta\eta \wedge *_y k_m - \eta \wedge *dk_m).$$
(19)

In case m = 0, the terms in $\delta k_m, \delta \eta$ are of course zero; to get a term in $\eta(0)$ on the right when $\varepsilon \to 0$ we need dk_m of the order of ε^{1-n} and k_m of the order of ε^{2-n} in $|y| = \varepsilon$. That makes k_m locally integrable too, and (12) is obtained leting $\varepsilon \to 0$. This means that for m = 0 k is unique and is given by the well-known Green's function

$$A(r) = c_n \int_r^1 (1 - s^2)^{n-2} s^{1-n} \, ds \tag{20}$$

for an appropriate choice of c_n . In case m > 0, again we need $|k_m(0, y)| = o(r^{1-n})$ as $r \to 0$, so that the first and third terms on the right have limit 0 as $\varepsilon \to 0$; then k_m is integrable in y and the integral on the left converges to $\int \Delta \eta \wedge *k_m$. Using the expression for $*dk_m$ in (14), we find

$$\int_{|y|=\varepsilon} \eta \wedge *d_y k_m = \frac{(-1)^{m(n-m+1)} n!}{(n-m-1)!} A_3(\varepsilon) *_x \int_{|y|=\varepsilon} \eta \wedge \alpha \wedge \gamma_{n-m-1}.$$

By Stoke's theorem, and since $\alpha = O(r)$, the last integral equals

$$(-1)^m \int_{|y|<\varepsilon} \eta \wedge \gamma_{n-m} + O(\varepsilon).$$

If $A_3(\varepsilon) = a_0 \varepsilon^{-n} + \dots$, we see that

$$\lim_{\varepsilon} \int_{|y|=\varepsilon} \eta \wedge * d_y k_m = c_n(n-m) \, m! \, a_0 \eta(0) \, .$$

Using (15) for $\delta k_m = (-1)^{n(m+1)+1} * d*$, and proceeding in the same way,

$$\int_{|y|=\varepsilon} \delta_y k_m \wedge *\eta = -A_4(\varepsilon) \int_{|y|=\varepsilon} \alpha \wedge \gamma_{m-1} \wedge *\eta$$
$$= -A_4(\varepsilon) \int_{|y|<\varepsilon} (\gamma_m \wedge *\eta + O(\varepsilon))$$

But by the equation $B_1 = 0$, $A_4(\varepsilon) = -(1 - \varepsilon^2)^{n-2m}((n-m)A_3(\varepsilon) + \varepsilon A'_3(\varepsilon)) = a_0 m \varepsilon^{-n} + O(\varepsilon^{1-n})$, and hence the limit of the above expression is $-c_n m! a_0 m \eta(0)$. Altogether, we conclude that if $A_3(\varepsilon) = a_0 \varepsilon^{-n} + o(\varepsilon^{1-n})$ and $k_m(0, y) = o(r^{1-n})$, one has

$$\int \Delta \eta \wedge *_y k_m(0,y) = -c_n \, n \, m! \, a_0 \, \eta(0)$$

so (12) will hold for an appropriate choice of a_0 . Taking into account the definition of A_3 in (14) and that $|k_m| \simeq |A_1| + r^2 |A_2|$, we see from (17) that if $m = \frac{n}{2}$ this is accomplished by the choice $c = 0, a = a_0$; then $G(x) \sim \log x, A_1(r) \sim \log r, A'_1(r) = O(1/r), A_2(r) = O(r^2)$ if n = 2; if n > 2, $A_1(r) \sim r^{2-n}$ and $A_2 = O(r^{-n})$. For $0 < m < \frac{n}{2}$, in terms of the functions H, G introduced before, this translates to $H(x) \sim a_0 x^{-\frac{n}{2}}, G(x) \sim x^{1-\frac{n}{2}}$. Now look at the general expression of H, G in (18). The condition $H(x) \sim c_0 x^{-\frac{n}{2}}$ fixes $a = a_0$; then near x = 0 c'(x) is bounded and d'(x) behaves like $x^{-\frac{n}{2}}$. Since $u_4(x)$ is bounded, the term $d(x)u_4(x)$ behaves like $x^{1-\frac{n}{2}}$. So, we must normalize c(x) by c(0) = 0, so that c(x) = O(x) and the other term $c(x)u_3(x)$ will behave like $x^{1-\frac{n}{2}}$.

In conclusion, all this discussion shows that the doubly invariant kernels $k_m(x, y)$ satisfying (12) constitute a *two parameter family* described by $H = a_0u_1(x) + bu_2(x)$, c(0) = 0. The two parameters are b and the constant of integration for d(x) in (18). Equivalently, they are obtained by adding to the form corresponding to $H = a_0u_1(x)$, c(0) = 0 and say $d(\frac{1}{2}) = 0$ the general globally smooth one described before.

3.4. In order to produce the best estimates, in a sense we need to choose the best of the kernels k_m . Naturally enough, we choose the k_m having the best behaviour at infinity, x = 1, that is, so that G, H have the best decrease in size as $x \to 1$. In case $m = \frac{n}{2}$, where we already have the normalization c = 0, $a = a_0$, the choice b = -a gives the best growth H(x) = O(1 - x) and $G(x) = O(\log(1 - x))$.

The hypergeometric function u_3 behaves like $(1-x)^{n+1-2m}$ near x = 1 while $u_4(x) = F(m, m - \frac{n}{2}, 1 + \frac{n}{2}, x)$ is bounded because $1 + \frac{n}{2} - m - (m - \frac{n}{2}) = 1 + n - 2m > 0$. Similarly, u_1 is bounded near x = 1; for $u_2(x) = F(\frac{n}{2} - m, n - m, \frac{n}{2} + 1, x)$ we have $\frac{n}{2} + 1 - (\frac{n}{2} - m) - (n - m) = 2m + 1 - n$ and hence it behaves like $(1 - x)^{2m+1-n}$ if 2m < n - 1 and like $\log(1 - x)$ if 2m = n - 1. We use equations (18)

$$c(x) = c_{m,n} \int_0^x H(t)(1+t)t^{n/2}(1-t)^{-3}u_4(t)dt$$
$$d(x) = -c_{m,n} \int_{\frac{1}{2}}^x H(t)(1+t)t^{n/2}(1-t)^{-3}u_3(t)dt + d_0$$

If $b \neq 0$, then $H(t) = a_0 u_1(t) + b u_2(t)$ behaves like $(1-t)^{2m+1-n}$ if 2m < n-1 and like $\log(1-t)$ if 2m = n-1, resulting in $c(x) = O((1-x)^{2m-n-1}, d(x)) = O(\log(1-x))$ if 2m < n-1 and $c(x) = O(((1-x)^{-2}\log(1-x)), d(x)) = O(((1-x)^{-1}\log(1-x)))$ if 2m = n-1. So if $b \neq 0$ one has $G(x) = O(\log(1-x))$ if 2m < n-1 and $G(x) = O(((1-x)^{-1}\log(1-x)))$ if 2m = n-1. If b = 0, then H is bounded, giving $c(x) = O(((1-x)^{-2}))$ and d(x) = O(1) for 2m < n-1, $d(x) = O(\log(1-x))$ for 2m = n-1. In case 2m < n-1, however, we can choose the constant d_0 so that d(1) = 0, and then $d(x) = O((1-x)^{n-2m-1})$. This choice gives $G(x) = O((1-x)^{n-2m-1}$ for 2m < n-1. For 2m = n-1, no choice of d_0 can improve the bound $G(x) = O(\log(1-x))$.

It remains to estimate the growth of $A_2(r)$ near r = 1. Recall that the definition (14) of A_3 translates to $A_2(r) = 2G'(x) - (1-x)^{n-2m-2}H(x)$. Both terms growth like $(1-x)^{n-2m-2}$, but a

cancellation occurs. The functions u_1, u_3 are C^{∞} at 1 and have developments

$$u_3(x) = A(1-x)^{n+1-2m} + O(1-x)^{n+2-2m}$$

$$u'_3(x) = -A(n+1-2m)(1-x)^{n-2m} + O(1-x)^{n+1-2m}$$

$$H(x) = a_0 u_1(x) = B + O(1-x).$$

In $u_4(x) = F(m, m - \frac{n}{2}, 1 + \frac{n}{2}, x)$, $1 + \frac{n}{2} - m - (m - \frac{n}{2}) = n + 1 - 2m \ge 2$, whence u_4 has a finite derivative at 1 and a development

$$u_4(x) = C + D(1-x) + O(1-x)^{1+\varepsilon} \quad \forall \varepsilon < 1, \quad u'_4(x) = O(1).$$

Then $W(x) = u'_3 u_4 - u_3 u'_4 = CA(2m - n - 1)(1 - x)^{n-2m} + \dots$, and so the constant c_{mn} in (18) is CA(2m - n - 1). Then from (18)

$$c'(x) = \frac{B(1-x)^{-3}}{A(2m-n-1)} + O(1-x)^{-2}$$
$$d'(x) = -\frac{B(1-x)^{n-2m-2}}{C(2m-n-1)} + O(1-x)^{n-2m-1}$$

which gives

$$c(x) = \frac{1}{2} \frac{B}{2(2m - n - 1)} (1 - x)^{-2} + O(1 - x)^{-1}$$

$$d(x) = O(1 - x)^{n - 2m - 1}, \ 2m < n - 1, \quad O(\log(1 - x)), 2m = n - 1$$

But $G' = c(x)u'_3(x) + d(x)u'_4(x)$; the second term $d(x)u'_4(x)$ satisfies the required bound while the first $c(x)u'_3(x)$ has a development

$$c(x)u'_{3}(x) = -\frac{1}{2}\frac{B}{A(2m-n-1)}A(n+1-2m)(1-x)^{n-2m-2} + O(1-x)^{n-2m-1}$$
$$= \frac{B}{2}(1-x)^{n-2m-2} + O(1-x)^{n-2m-1}.$$

As $(1-x)^{n-2m-2}H(x) = B(1-x)^{n-2m-2} + O(1-x)^{n-2m-1}$ the bound for A_2 follows for $2m \le n-1$. However, for $m = \frac{n}{2}$, this no longer holds. Indeed, from (17), where c = 0, $a = a_0$, b - a)

$$2G'(x) = x^{-\frac{n}{2}-1} \int_0^x s^{n/2} (1+s)(1-s)^{-3} a(s^{-\frac{n}{2}}-1) \, ds$$

has development

$$2G'(x) = na(1-x)^{-1} + O(\log(1-x))$$

while

$$(1-x)^{-2}a(x^{-\frac{n}{2}}-1) = \frac{n}{2}a(1-x)^{-1} + \dots$$

We point out that all this can be obtained, in loose terms, working directly with the hypergeometric equations relating G, H

$$x(1-x)G''(x) + \left[\frac{n}{2} + 1 - (2m+1-\frac{n}{2})x\right]G'(x) - m(m-\frac{n}{2})G = \frac{1}{2}(1+x)(1-x)^{n-2m-2}H(x)$$

and using asymptotic developments. If $H(x) = h_0 + h_1(1-x) + \dots$ and $G(x) = g_j(1-x)^j + \dots$, identifying the lower order terms in both sides gives,

$$g_j j(j-1-n+2m)(1-x)^{j-1} = h_0(1-x)^{n-2m-2}$$

When $H \equiv 0$, one must have either j = 0 (corresponding to u_4) or j = n - 2m + 1 (corresponding to u_3). For the inhomogeneous equation, if $j \neq 0$, $j \neq n + 1 + 2m$ (that is G contains no contribution from u_3, u_4) one finds j = n - 2m - 1 if 2m < n - 1 and $g_j j = -\frac{h_0}{2}$. Then $2G'(x) = h_0(1-x)^{n-2m-2} + \dots, (1-x)^{n-2m-2}H(x) = h_0(1-x)^{n-2m-2}$, showing cancellation. An analogous argument works if 2m = n - 1, but not for 2m = n.

We summarize the results in this and the previous subsections:

Theorem 3.1. For |n-2m| > 1, there is a unique doubly invariant kernel

$$k_m(x,y) = A_1(|\varphi_x y|)\gamma_m + A_2(|\varphi_x y|)\tau \land \gamma_{m-1}, m \neq 0; k_m(x,y) = A(|\varphi_x y|)\gamma_m, m = 0, m \neq 0$$

for which (12) holds, and satisfying moreover

$$|A_i(r)| = O(1 - r^2)^{|n-2m|-1}$$
 as $r \to 1$.

For $m = \frac{n \pm 1}{2}$, there is a one-parameter family of such kernels satisfying

$$|A_i(r)| = O(\log(1 - r^2))$$

For $m = \frac{n}{2}$, there is a one-parameter family of such kernels satisfying

$$|A_i(r)| = O(1 - r^2)^{-1}$$

In all cases $A_1(r) \sim r^{2-n}$, $A_2(r) \sim r^{-n}$ as $r \to 0$.

For |n-2m| > 1, we call $k_m(x, y)$ the Riesz kernel for m-forms in \mathbb{H}^n , and

$$L\eta(x) = \int_{\mathbb{H}^n} \eta(y) \wedge *_y k_m(x,y)$$

the Riesz potential of η , whenever this is defined. From (11) we see that

$$|k_m(x,y)| = O(1-r^2)^{n-m-1}.$$
(21)

With the notations used before, the function $A_3(r) = H(r^2)$ is bounded with bounded derivatives near r = 1. Then (14) and symmetry imply

$$|d_x k_m(x,y)|, |d_y k_m(x,y)| = O(1-r^2)^{n-m-1}$$
(22)

too. The growth of A_3 also implies $A_4 = O(1 - r^2)^{n-2m}$ because $B_1 \equiv 0$, and then (15) gives as well

$$|\delta_x k_m(x,y)|, \ |\delta_y k_m(x,y)| = O(1-r^2)^{n-m-1}.$$
(23)

By construction, one has $L\Delta \eta = \eta$ for $\eta \in \mathcal{D}_m(\mathbb{H}^n)$. We will need the following generalization of this fact.

Proposition 3.2. If η is a smooth form in \mathbb{H}^n such that

$$|\eta(y)|, |\nabla \eta(y)| = o(1 - |y|^2)^m, \quad y \in \mathbb{B}^n$$

then $L\Delta\eta = \eta$.

Proof. In (19) we would get an extra term

$$\int_{|y|=R} (k_m \wedge *d\eta + \delta k_m \wedge *\eta - \delta \eta \wedge *k_m - \eta \wedge *dk_m) \, dk_m \, d$$

Estimates (21), (22) and (23) imply that with x fixed and $|y| = R \nearrow 1$

$$|k_m|, |\delta k_m|, |dk_m| = O(1 - R^2)^{n - m - 1}$$

Inserting $|\eta(y)|, |\nabla \eta(y)| = o(1 - |y|^2)^m$ we see that this extra term vanishes as $R \nearrow 1$.

4 Proof of the main theorem

4.1. Once the Riesz form $k_m(x, y)$ has been found, our aim is now to prove that the corresponding convolution

$$L_m\eta(x) = \int_{\mathbb{H}^n} \eta(y) \wedge *_y k_m(x,y)$$

satisfies

$$||L_m\eta||_{p,s+2} \le c||\eta||_{p,s} \tag{24}$$

for $m \neq \frac{n\pm 1}{2}$, $\frac{n}{2}$, and p in the range $p_1(m) = \frac{n-1}{n-1-m} , and for a compactly supported$ *m* $-form <math>\eta$ (recall that we are assuming without loss of generality that $m \leq \frac{n}{2}$). Since these are dense in the Sobolev spaces and we already know that $\Delta L_m \eta = L_m \Delta \eta = \eta$, this will prove the theorem for $m \neq \frac{n\pm 1}{2}$, $\frac{n}{2}$. The case $m = \frac{n\pm 1}{2}$ will be commented later.

We work in the translation invariant basis w^{I} . Taking into account formulas (9) and (10) for γ, τ , the Riesz- form is written in the \mathbb{R}^{n}_{+} model

$$k_m(x,y) = \sum_{|I|=|J|=m} a_{I,J}(S_y x) w^I(x) \otimes w^J(y)$$

where each coefficient $a_{I,J}$ has an expression, with $z = S_y x$

$$a_{I,J}(z) = \Psi_{I,J}(r) \frac{p_{I,J}(z)}{(|z|^2 + 2z_n + 1)^{2m}}, \quad r^2 = \frac{1 + |z|^2 - 2z_n}{1 + |z|^2 + 2z_n} = \frac{|x - y|^2}{|x - y|^2 + 4x_n y_n}$$

Here $p_{I,J}(z)$ is a certain polynomial in z_1, \ldots, z_n , $\Psi_{I,J}$ is C^{∞} in (0,1) with $\Psi_{I,J}(r) \sim c_0 r^{2-n}$ as $r \searrow 0$, $\Psi_{I,J}(r) = O(1-r^2)^{n-m-1}$ as $r \nearrow 1$. The term $q_{I,J}(z) = p_{I,J}(z)/(|z|^2 + 2z_n + 1)^{2m}$ is bounded.

If $\eta = \sum_{I} \eta_{I}(y) w^{I}(y)$, the coefficient $(L\eta)_{I}(x)$ of $L\eta$ in the basis w^{I} is a finite linear combination of hyperbolic convolutions

$$(L\eta)_I(x) = \sum_J \int_{\mathbb{H}^n} \Psi_{I,J}(r) q_{I,J}(z) \eta_J(y) \, d\mu(y) \, .$$

By ellipticity of Δ , $L\eta$ is a smooth form. Moreover, since η has compact support, we see from (2) and (21), (22), (23) that, in the ball model,

$$|L\eta(x)|, |d(L\eta)(x)|, |\delta(L\eta)(x)| = O(1 - |x|^2)^{n-m-1}$$

which amounts to

$$|(L\eta)_I(x)|, |X_i(L\eta)_I(x)| = O(1-|x|^2)^{n-m-1}.$$
(25)

We claim that for second-order derivatives we have too

$$|X_j X_i(L\eta)_I(x)| = O(1 - |x|^2)^{n-m-1}, \quad i.e. \quad |\nabla^{(2)}(L\eta)(x)| = O(1 - |x|^2)^{n-m-1}.$$
(26)

Notice that since we already know that $\Delta L\eta = \eta$, from the expression of Δ in the basis w^I given in (3) it follows that it is enough to show that for j < n. We will see below (equation (30) and invariance of the X_i) that each of the functions $a(z) = \Psi_{I,J}(r)q_{I,J}(z)$ satisfies

$$|X_j X_i a(z)| = O(1 - r^2)^{n - m - 1}$$

from which (26) follows as before. In fact, the discussion that follows will show that $|\nabla^{(k)}L\eta(x)| = O(1-|x|^2)^{n-m-1} \quad \forall k.$

We continue the proof of (24). We claim first that it is enough to prove (24) for s = 0. For a smooth form $\eta = \sum \eta_I w^I$ let $X_i \eta$ denote here the *m*-form $X_i \eta = \sum X_i \eta_I w^I$. It is clear from formulas (3) and the commutation properties

$$[X_i, X_j] = 0, \qquad i, j < n, \qquad [X_1, X_i] = X_i, \qquad i < n$$

that for each i there is an operator P_i of order two in the X_1, \ldots, X_n such that

$$X_i \Delta \eta - \Delta(X_i \eta) = P_i(X) \eta \,.$$

Applying this to $L\eta$, which is smooth by the ellipticity of Δ , we get $(X_i - \Delta X_i L)\eta = P_i(X)L\eta$. But $X_i L\eta$ satisfies, by (25) and (26)

$$|X_i L\eta(x)|, |d(X_i L\eta)(x)|, |\delta(X_i L\eta)(x)| = O(1 - |x|^2)^{n-m-1}$$

and hence by Proposition 3.2, $L\Delta = \text{Id on it.}$ We conclude that for all $\eta \in \mathcal{D}_m(\mathbb{H}^n)$

$$(LX_i - X_iL)\eta = LP_i(X)L\eta.$$

Assume that (24) has been proved up to s, so that by density it holds for $\alpha \in H^s_{m,p}(\mathbb{H}^n)$ too, and let γ be a multiindex of length $|\gamma| \leq s$. For $i = 1, \ldots, n$ and $\eta \in \mathcal{D}_m(\mathbb{H}^n)$,

$$X^{\gamma}X_{i}L\eta = X^{\gamma}LX_{i}\eta - X^{\gamma}LP_{i}(X)L\eta$$

so using twice the induction hypothesis

$$\begin{aligned} ||X^{\gamma}X_{i}L\eta||_{p} &\leq \operatorname{const}\left(||X_{i}\eta||_{p,s} + ||P_{i}(X)L\eta||_{p,s}\right) \leq \\ &\leq \operatorname{const}\left(||\eta||_{p,s+1} + ||\eta||_{p,s}\right) \end{aligned}$$

proving (24) for s + 1. Proving (24) for s > 0 means proving

$$||(L\eta)_I||_p$$
, $||X_i(L\eta)_I||_p$, $||X_jX_i(L\eta)_I||_p \le \text{const} ||\eta||_p$.

As before, using that we already know that $\Delta L\eta = \eta$ we see that for the second-order derivatives we may assume j < n. In the following we delete the indexes I, J and denote by $a(z) = \psi(r)Q(z)$ a convolution kernel with ψ, Q as above, and proceed to prove that the convolution

$$(C_a\alpha)(z) = \int_{\mathbb{H}^n} a(S_y x) \alpha(y) \, d\mu(y)$$

satisfies

$$||C_a\alpha||_p, ||X_i(C_a\alpha)||_p, ||X_jX_iC_a(\alpha)||_p \le \text{const} ||\alpha||_p, \qquad p_1 \le p \le p_2$$
(27)

where in the last case we may assume that j < n. The fields X_i are invariant, and therefore $X_i C_a \alpha$, $X_j X_i C_a \alpha$ are obtained, respectively, by convolution with $Z_i a$, $Z_j Z_i a$ (by (4)). Recall that $\psi(r) = O(1 - r^2)^{n-m-1} = O\left(\frac{4z_n}{1 + |z|^2 + 2z_n}\right)^{n-m-1} \text{ as } r \nearrow 1 \text{ and } \psi(r) \sim r^{2-n} \text{ as } r \searrow 0.$

In order to estimate $Z_i a, Z_i Z_j a$, we collect first some auxiliary estimates. We claim that

$$|Z_i Q| \le \operatorname{const}, \qquad |Z_i Z_j Q| \le \operatorname{const}$$

$$|Z_i r| \le \operatorname{const}(1 - r^2), \qquad |Z_i Z_j r| \le \operatorname{const} r^{-1}(1 - r^2).$$
(28)

The first two are routinely checked, for instance, when differentiating the denominator in Q,

$$\left| Z_i \frac{1}{(1+|z|^2+2z_n)^{2m}} \right| = \left| \frac{4mz_n z_1}{(1+|z|^2+2z_n)^{2m+1}} \right| \le \frac{\text{const}}{(1+|z|^2+2z_n)^{2m}} \qquad (i < n)$$

so that the term $p_{I,J}(z)Z_i\left[(1+|z|^2+2z_n)^{-2m}\right]$ will still be bounded. All other terms can be treated similarly. Differentiating $1 - r^2 = \frac{4z_n}{1 + |z|^2 + 2z_n}$ we get

$$Z_{i}r = \frac{1-r^{2}}{2} \frac{z_{i}z_{n}}{r(1+|z|^{2}+2z_{n})}, \qquad Z_{n}r = -\frac{1-r^{2}}{2} \frac{1}{r} \frac{1+|z|^{2}-2z_{n}^{2}}{1+|z|^{2}+2z_{n}}$$

$$Z_{j}Z_{i}r = \frac{1-r^{2}}{2r} \left\{ \frac{\delta_{ij}z_{n}^{2}}{1+|z|^{2}+2z_{n}} - \frac{1+5r^{2}}{2r^{2}} \frac{z_{i}z_{j}z_{n}^{2}}{(1+|z|^{2}+2z_{n})^{2}} \right\}, \qquad i,j < n$$

$$Z_{j}Z_{n}r = \frac{1-r^{2}}{r} \left\{ -\frac{2z_{n}^{2}z_{j}(1+z_{n})}{(1+|z|^{2}+2z_{n})^{2}} + \frac{(1+r^{2})}{4r^{2}} \frac{z_{j}z_{n}(1+|z|^{2}-2z_{n})}{(1+|z|^{2}+2z_{n})^{2}} \right\}, \qquad j < n$$

These imply (28) because

$$|z_i z_n|, 1 + |z|^2 - 2z_n^2 \le (1 + |z|^2 - 2z_n)^{1/2} (1 + |z| + 2z_n)^{1/2} = r(1 + |z|^2 + 2z_n).$$

Now

$$Z_{i}a(z) = \psi'(r)Z_{i}rQ(z) + \psi(r)(Z_{i}Q)(z)$$

$$Z_{j}Z_{i}a(z) = \psi''(r)(Z_{i}r)(Z_{j}r)Q(z) + \psi'(r)(Z_{j}Z_{i}r)Q + \psi'(r)Z_{i}rZ_{j}Q$$

$$+ \psi'(r)Z_{j}rZ_{i}Q + \psi(r)Z_{j}Z_{i}Q.$$
(29)

The estimates (28) imply

$$|a(z)|, |Z_i a(z)|, |Z_j Z_i a(z)| = O(1 - r^2)^{n - m - 1} \quad \text{as} \quad r \nearrow 1$$

$$|a(z)| = O(r^{2 - n}), |Z_i a(z)| = O(r^{1 - n}), |Z_j Z_i a(z)| = O(r^{-n})$$
(30)

We will call a convolution kernel b(z) *m*-admissible if $|b(z)| = O(r^{1-n})$ as $r \searrow 0$ and, moreover, $|b(z)| = O(1-r^2)^{n-m-1}$ as $r \nearrow 1$. We will prove later (Theorem 4.2.) that hyperbolic convolution with *m*-admissible kernels defines a bounded operator in $L^p(\mathbb{H}^n)$ for the range $(p_1(m), p_2(m))$ specified in the statement of the main result. From the estimates (29) we see that *a* and $Z_i a$ are *m*-admissible kernels, and so (27) will be proved for them. As $|Z_j Z_i a(z)| = O(r^{-n})$ has the critical non-integrable singularity at r = 0, $Z_j Z_i a(z)$ is not an *m*-admissible kernel. Notice however from (29), (30) that the last three terms $\psi'(r)Z_irZ_jQ$, $\psi'(r)Z_jrZ_iQ$, $\psi(r)Z_jZ_iQ$ are indeed *m*-admissible. Moreover, the estimate $|Z_iQ| \leq \text{const}$ implies that *Q* is Lipschitz with respect the hyperbolic metric, in particular

$$Q(z) = Q(e) + O(\log \frac{1+r}{1-r}) = Q(e) + O(r)$$

for small r. This means that replacing Q by Q-Q(e) in the first two terms leads to an m-admissible kernel again. All this leaves us with the kernel

$$\psi''(r)Z_irZ_jr + \psi'(r)Z_jZ_ir, \qquad j < n.$$

If $\psi(r) = c_0 r^{2-n} + \dots$, write $\phi(r) = c_0 r^{2-n} (1-r^2)^{n-m-1}$; then the above differs from

$$\phi''(r)Z_irZ_jr + \phi'(r)Z_jZ_ir$$

in an *m*-admissible kernel. By the same reason, we may replace $\phi''(r)$, $\phi'(r)$ respectively by $(r^{2-n})''(1-r^2)^{n-m-1}$, $(r^{2-n})'(1-r^2)^{n-m-1}$, that is to say we must deal with the convolution kernel

$$(1 - r^2)^{n - m - 1} Z_j Z_i(r^{2 - n}). (31)$$

We introduce a class of singular hyperbolic convolution kernels to deal with the later. For this purpose it is more convenient to work in the ball model, so now b is defined in \mathbb{B}^n and r = |z|. We

replace the integrable singularity r^{1-n} by a typical Calderón-Zygmund singularity (see e.g. [S]). Thus, we will call b a *m*-Calderón-Zygmund singular kernel if it has the form

$$b(z) = \Omega(w)r^{-n}(1-r^2)^{n-m-1}, \quad z = rw, \quad w \in S^{n-1}$$

where Ω is say a Lipschitz function on S^{n-1} satisfying the cancellation condition

$$\int_{S^{n-1}} \Omega(w) \, d\sigma(w) = O \,. \tag{32}$$

In Theorem 4.2. below we prove that *m*-Calderón-Zygmund singular kernels define bounded operators in the same range of *p*. With the following proposition, applied to $\phi_2(z) = |z|^{2-n}$ this will end the proof of the main result. The proposition is the analogue of the well-known statement that for ϕ smooth and homogeneous of degree 1-n in \mathbb{R}^n , $\frac{\partial \phi}{\partial x_i}$ defines a Calderón-Zygmund kernel (it is homogeneous of degree -n and the cancellation condition (32) is automatically satisfied, because $\int_{r_1 < |x| < r_2} \frac{\partial \phi}{\partial x_i} dV(x) = \left(\int_{|x|=r_2} - \int_{|x|=r_1}\right) \phi(x) dx^1 \wedge \cdots \wedge dx^i \wedge \cdots \wedge dx^n = 0$).

Proposition 4.1. If ϕ_1, ϕ_2 are homogeneous functions of degree 1-n, 2-n respectively, the kernels $(1-r^2)^{n-m-1}Z_i\phi_1, (1-r^2)^{n-m-1}Z_jZ_i\phi_2$ are sum of (m-1)-admissible and (m-1)-Calderón-Zygmund singular kernels.

Proof. We replace the Z_j by $Y_j = (1 - r^2) \frac{\partial}{\partial z_j}$; we have

$$Y_{i}(\phi_{1}) = (1 - r^{2})\frac{\partial\phi_{1}}{\partial z_{i}} \qquad Y_{i}\phi_{2} = (1 - r^{2})\frac{\partial\phi_{2}}{\partial z_{i}} = (1 - r^{2})O(r^{1 - n})$$
$$Y_{j}Y_{i}\phi_{2} = (1 - r^{2})\frac{\partial^{2}\phi_{2}}{\partial z_{i}\partial z_{j}} - 2(1 - r^{2})z_{j}\frac{\partial\phi_{2}}{\partial z_{i}} = (1 - r^{2})\frac{\partial^{2}\phi_{2}}{\partial z_{i}\partial z_{j}} + (1 - r^{2})O(r^{2 - n})$$

so in all cases we get an extra factor $(1-r^2)$. Besides $\frac{\partial \phi_1}{\partial z_i}$, $\frac{\partial^2 \phi_2}{\partial z_i \partial z_j}$ are, as noted before, homogeneous of degree -n, and satisfy the cancellation condition (32).

4.2. It remains to prove

Theorem 4.2. Both *m*-admissible and *m*-Calderón-Zygmund kernels define by hyperbolic convolution bounded operators in $L^p(\mathbb{H}^n)$ for $\frac{n-1}{n-1-m} , <math>0 \le m < \frac{n-1}{2}$.

We will make use of the following well-known Schur's lemma for boundedness in L^p of an integral operator with positive kernel.

Lemma 4.3. If K(x,y) is a positive kernel in a measure space X and $1 , the operator <math>Kf(x) = \int_X K(x,y)f(y) d\mu(y)$ is bounded in $L^p(\mu)$ if and only if there exists $h \ge 0$ such that

$$\int_X K(x,y)h(y)^q \, d\mu(y) = O(h(x)^q) \,, \qquad x \in X \tag{33}$$

$$\int_{X} K(x, y) h(x)^{p} d\mu(x) = O(h(y)^{p}), \qquad y \in Y.$$
(34)

Here q is the conjugate exponent of p, $\frac{1}{p} + \frac{1}{q} = 1$. If h can be taken $\equiv 1$ that is

$$\sup_{x} \int_{X} K(x,y) \, d\mu(y) \, , \sup_{y} \int_{X} K(x,y) \, d\mu(x) < +\infty$$

then K is bounded in $L^p(\mu)$ for all $p, 1 \leq p \leq \infty$.

Proof. Let us prove Theorem 4.2. If b is m-admissible, $b = b_1 + b_2$ with $b_1(z) = O(r^{1-n})$ for $r \le 1/2$, $b_1(z) = 0$ for r > 1/2, and $b_2(z) = O(1 - r^2)^{n-m-1}$ for all r. We apply to b_1 the second criteria in Lemma 4.3 working in the ball model (recall that $|S_yX| = |\varphi_yx|$ is symmetric in x, y)

$$\int_X b_1(S_y x) \, d\mu(x) \, , \int_X b_1(S_y x) \, d\mu(y) \le c \int_{|S_y x| \le 1/2} |S_y x|^{1-n} d\mu(x)$$
$$= c \int_{|z| \le 1/2} |z|^{1-n} d\mu(z) = \text{const} \int_0^{1/2} \frac{dr}{(1-r^2)^n} < +\infty.$$

We apply to $(1 - r^2)^{n-m-1}$ the criteria of the first part on Lemma 4.3, working this time for convenience in the half-space model, where the kernel is written

$$K(x,y) = (1-r^2)^{n-m-1} = \left(\frac{4z_n}{1+|z|^2+2z_n}\right)^{n-m-1} = \left(\frac{4x_ny_n}{|x-y|^2+4x_ny_n}\right)^{n-m_1}$$

We test $h(y) = y_n^{\alpha}$ in (33) for an exponent α to be chosen, so we need

$$\int_{y_n>0} \frac{y_n^{-k-1+\alpha q} \, dy}{(|x-y|^2 + 4x_n y_n)^{n-k-1}} = O\left(x_n^{\alpha q+k+1-n}\right) \, .$$

We write $|x-y|^2 + 4x_n y_n = |x'-y'|^2 + (x_n+y_n)^2$, where $x' = (x_1, \ldots, x_{n-1}) \in \mathbb{R}^{n-1}$ and analogously for y', and integrate first in y'. One has for 2m < n-1

$$\int_{\mathbb{R}^{n-1}} \frac{dy'}{(|x'-y'|^2 + (x_n + y_n)^2)^{n-m-1}} = c \int_0^\infty \frac{s^{n-2}}{(s^2 + (x_n + y_n)^2)^{n-m-1}}$$
$$= O((x_n + y_n)^{2m+1-n})$$

and so the above becomes

$$\int_0^\infty \frac{y_n^{\alpha q - m - 1} dy_n}{(x_n + y_n)^{n - 1 - 2m}} = O(x_n^{\alpha q + m + 1 - n}).$$

By homogeneity $(y_n = x_n t)$ this reduces to

$$\int_0^\infty \frac{t^{\alpha q - m - 1}}{(1 + t)^{n - 1 - 2m}} = O(1)$$

which holds whenever $m < \alpha q < n - 1 - m$. By symmetry, for (34) we need as well $m < \alpha p < n - 1 - m$. Therefore, a choice of α is possible whenever $m \max\left(\frac{1}{p}, \frac{1}{q}\right) < (n - 1 - n) \min\left(\frac{1}{p}, \frac{1}{q}\right)$ and this gives the range $\frac{n-1}{n-1-m} .$

Consider now a *m*-Calderón-Zygmund kernel $b(z) = \Omega(w)r^{-n}(1-r^2)^{n-m-1}$. Since $|S_yx| = |\varphi_yx|$, we may replace $z = S_yx$ by $z = \varphi_xy$. Using (1) this is given by

$$z = \frac{(x-y)(1-|x|^2) + x|x-y|^2}{A}$$

where we use the notation $A = (1 - |x|^2)(1 - |y|^2) + |x - y|^2$; note that

$$(1 - |x|^2), (1 - |y|^2) \lesssim A^{1/2}$$

Also recall that r = |z| and $Ar^2 = |x - y|^2$. Hence we can write

$$\frac{z}{r} - \frac{x - y}{|x - y|} = \frac{x - y}{|x - y|} \left(\frac{1 - |x|^2}{\sqrt{A}} - 1\right) + x \cdot r.$$

But

$$\frac{1-|x|^2}{\sqrt{A}} - 1 = \frac{(1-|x|^2)^2 - A}{\sqrt{A}((1-|x|^2) + \sqrt{A})} = \frac{(1-|x|^2)O(|x-y|) + O(|x-y|^2)}{A}$$

is O(r). Therefore, modulo an *m*-admissible kernel, we may replace $\Omega(w)$ by $\Omega\left(\frac{x-y}{|x-y|}\right)$. This leaves us with the kernel

$$K = (1 - r^2)^{n - m - 1} \Omega\left(\frac{x - y}{|x - y|}\right) r^{-n} = (1 - r^2)^{n - m - 1} |x - y|^{-n} \Omega\left(\frac{x - y}{|x - y|}\right) A^{\frac{n}{2}}(x, y).$$

Fix p, 1 . Write

$$A^{\frac{n}{2}}(x,y) = (1-|x|^2)^{\frac{n}{p}}(1-|y|^2)^{\frac{n}{q}} + O\left(|x-y|A^{\frac{n-1}{2}}\right)$$

Since $|x - y|^{1-n} A^{\frac{n-1}{2}} = r^{1-n}$, the kernel K differs from

$$(1-r^2)^{n-m-1}|x-y|^{-n}\Omega\left(\frac{x-y}{|x-y|}\right)(1-|x|^2)^{\frac{n}{p}}(1-|y|^2)^{\frac{n}{q}}$$

in a m-admissible kernel, so we keep this one. We write it as the sum of

$$|x-y|^{-n}\Omega\left(\frac{x-y}{|x-y|}\right)(1-|x|^2)^{\frac{n}{p}}(1-|y|^2)^{\frac{n}{q}} = K_1(x,y)$$

and another $K_2(x, y)$ which we estimate by

$$|K_2(x,y)| = O\left(r^2|x-y|^{-n}(1-|x|^2)^{\frac{n}{p}}(1-|y|^2)^{\frac{n}{q}}\right) = O\left(r^{2-n}(1-|x|^2)^{\frac{n}{p}}(1-|y|^2)^{\frac{n}{q}}A^{-\frac{n}{2}}\right).$$

Write K_{Ω} for the (euclidean) Calderón-Zygmund convolution operator with kernel $|x-y|^{-n}\Omega\left(\frac{x-y}{|x-y|}\right)$, which as it is well-known, satisfies an $L^p(dV)$ -estimate. Notice that

$$K_1 f(x) = (1 - |x|^2)^{\frac{n}{p}} K_\Omega \left(f(1 - |y|^2)^{-\frac{n}{p}} \right)$$

and therefore, using the L^p -boundedness of K_{Ω}

$$\int_{\mathbb{B}^n} |K_1 f(x)|^p \, d\mu(x) = \int_{\mathbb{B}^n} \left| K_{\Omega} (f(1-|y|^2)^{-\frac{n}{p}}) \right|^p \, dV(x) \le \int_{\mathbb{B}^n} |f(x)|^p \, d\mu(y) \, .$$

For K_2 , we can ignore the integrable simgularity r^{2-n} and arguing as we just did with K_1 , we need to show that the integral operator

$$K_3 f(x) = \int_{|y| \le 1} \frac{1}{(1 - |x| + |x - y|)^n} f(y) \, dV(y)$$

satisfies $L^p(dV)$ -estimates for all $p, 1 . To see this, just check that the criteria in Lemma 4.3 holds with <math>h(x) = (1 - |x|^2)^{-\frac{1}{pq}}$

Notice that in case m = 0 a *m*-Calderón-Zygmund kernel defines a bounded operator in all $L^{p}(\mathbb{H}^{n}), 1 : this is the right analogue of the euclidian kernels, because <math>(1 - r^{2})^{n-1}$ is the typical growth at infinity of a weak $L^{1}(d\mu)$ function in \mathbb{H}^{n} .

4.3. Finally we make some comments, with no proofs, on the critical case $m = \frac{n-1}{2}$ in the main Theorem. In this case the *m*-admissible and *m*-Calderón-Zygmund operators appearing in $X_j X_i C_a u$, etc. have $(1-r^2)^{\frac{n-1}{2}} \log \frac{1}{1-r^2}$ instead of $(1-r^2)^{n-m-1} = (1-r^2)^{\frac{n-1}{2}}$ as a factor. One can then prove that for $\beta > 0$ and $2 \le p < 2 + \frac{2\beta}{(n-1)}$,

$$||L_p\eta||_{p,2} \le \operatorname{const} \int_{\mathbb{B}^n} |\eta|^p (1-|y|^2)^{-\beta} d\mu(y).$$

The L^p -estimates do not hold in this case for any p, because they do not hold for p = 2 and Δ is self-adjoint.

References

- [BG] J. BRUNA AND J. GIRBAU Mapping properties of the Laplacian in Sobolev spaces of forms on complete hyperbolic manifolds, preprint Departament de Matemàtiques de la UAB, 2002
- [AL] J. P. ANKER AND N. LOHOUÉ Multiplicateurs sur certains espaces symétriques, Amer.
 J. Math. 108 (1986), No.6, 1303–1353
- [A] J. P. ANKER Sharp estimates for some functions of the Laplacian on noncompace symmetric spaces, Duke Math. J. 65 (1992), No.2, 257–297
- [D] H. DONELLY The differential form spectrum of hyperbolic space, Manuscripta Mathematica **33** (1981), 365–385
- [E] A. ERDÉLYI, Higher transcendental functions, vol 1, McGraw Hill, New York, 1953.
- [L1] N. LOHOUÉ Comparison des champs de vecteurs et des puissances du laplacien sur une variété riemannienne à courbure non positive, J. Funct. Anal. 61 (1985), No.2, 164–201

[L2]	N. LOHOUÉ Transformées de Riesz et fonctions de Littlewood-Paley sur les groupes non moyenables, C. R. Acad. Sci. Paris Sér. I Math. 306 (1988), No.7, 327–330.
[L3]	N. LOHOUÉ Estimations asymptotiques des noyaux résolvents du laplacien des formes differentielles sur les espaces symétriques de rang un, de type non compact et applica- tions, C. R. Acad. Sci. Paris Sér. I Math. 307 (1988), No.11, 551–554.
[L4]	N. LOHOUÉ Remarques sur les inégrales singulières sur les variétés à courbure non positive, C. R. Acad. Sci. Paris Sér. I Math. 307 (1988), No.12, 647–649.
[LV]	N. LOHOUÉ AND N. TH. VAROPOULOS Remarques sur les transformées de Riesz sur les groupes de Lie nilpotents, C. R. Acad. Sci. Paris Sér. I Math. 301 (1985), No.11, 559–560.
[I]	 A. D. IONESCU Singular integrals on symmetric spaces of real rank one, Duke Math. J. 114 (2002), No.1, 101–122.
[P1]	E. PEDON Analyse harmonique des formes differentielles sur l'espace hyperbolique réel, I, C. R. Acad. Sci. Paris Sér. I Math. 326 (1998), No.6, 671–676.
[P2]	E. PEDON Analyse harmonique des formes differentielles sur l'espace hyperbolique réel, II, C. R. Acad. Sci. Paris Sér. I Math. 326 (1998), No.7, 781–786.
[S]	E. STEIN, Singular integrals and differentiability properties of functions. Princeton University Press, 1970.

Departament de Matemàtiques Universitat Autònoma de Barcelona Campus de Bellaterra 08193 Cerdanyola del Vallès (Barcelona), Spain bruna@mat.uab.es