## AN EXAMPLE OF $\infty$ -HARMONIC FUNCTION, WHICH IS NOT $C^2$ ON A DENSE SUBSET

## HAYK MIKAYELYAN

ABSTRACT. We show that for certain boundary values McShane-Whitney's minimal-extension-like function is  $\infty$ -harmonic near the boundary and is not  $C^2$  on a dense subset.

Let us consider the strip  $\{(u,v) \in \mathbb{R}^2 : 0 < v < \delta\}$ . The function we are going to construct will be defined in this strip. Take a function  $f \in C^{1,1}(\mathbb{R})$  with  $L_f := \|f'\|_{\infty}$  and  $L'_f := Lip(f')$ . Let us consider an analogue of the minimal extension of McShane and Whitney

$$u(x,d) := \sup_{y \in \mathbb{R}} [f(y) - L|(x,d) - (y,0)|], \tag{1}$$

where  $0 < d < \delta$  and  $L > L_f$ . Note that in order to get the classical minimal extension of McShane and Whitney we have to take  $L = L_f$ .

From now on let us fix the function f, constants  $L > L_f$  and  $\delta > 0$ . We are going to find some conditions on  $\delta > 0$ , which will make our statements to be true. The real number x will be associated with the point  $(x, \delta) \in \Gamma_{\delta} := \{(u, v) \in \mathbb{R}^2 : v = \delta\}$  and the real number y with the point  $(y, 0) \in \Gamma_0$ . In the sequel the values of u on the line  $\Gamma_{\delta}$  will be of our interest and we write u(x) for  $u(x, \delta)$ .

## Proposition 1.

$$u(x) = \sup_{y \in \mathbb{R}} [f(y) - L\sqrt{\delta^2 + (x - y)^2}] = \max_{|y - x| \le D\delta} [f(y) - L\sqrt{\delta^2 + (x - y)^2}], \quad (2)$$

where  $D := \frac{2LL_f}{L^2 - L_f^2}$ .

**Proof.** From the definition of u we have that

$$f(x) - L\delta \le u(x)$$

so it is enough to show that if  $|x-y| > D\delta$  then

$$f(y) - L\sqrt{\delta^2 + (x - y)^2} < f(x) - L\delta.$$

On the other hand from the boundedness of f' we have

$$f(y) - L\sqrt{\delta^2 + (x - y)^2} \le f(x) + L_f|x - y| - L\sqrt{\delta^2 + (x - y)^2}$$

Thus we note that all values of y for which

$$f(x) + L_f|x - y| - L\sqrt{\delta^2 + (x - y)^2} < f(x) - L\delta$$

can be ignored in taking supremum in the definition of u.

Key words and phrases. Infinity-Laplacian.

 $2000\ Mathematics\ Subject\ Classification.\ 35B65,\ 35J70,\ 26B05.$ 




FIGURE 1. Touched by hyperbola

We write

$$L_f|x-y| + L\delta < L\sqrt{\delta^2 + (x-y)^2}$$

and arrive at

$$L_f^2|x-y|^2 + 2LL_f\delta|x-y| + L^2\delta^2 < L^2\delta^2 + L^2|x-y|^2$$

Thus

$$2LL_f\delta < (L^2 - L_f^2)|x - y| <=> |x - y| > D\delta.$$

Let y(x) be one of the points in  $\{|y-x| \leq D\delta\}$ , where the maximum in (2) is achieved

$$u(x) = f(y(x)) - L\sqrt{\delta^2 + (x - y(x))^2}.$$
 (3)

**Lemma 2.** If  $\delta > 0$  is small enough then for every  $x \in \Gamma_{\delta}$  the point y(x) is unique and  $y(x) : \mathbb{R} \to \mathbb{R}$  is a bijective Lipschitz map.

**Proof.** For each  $x \in \Gamma_{\delta}$  consider the function  $g_x(y) := u(x) + L\sqrt{\delta^2 + (x-y)^2}$  defined on  $\Gamma_0$ . The graph of  $g_x$  is a hyperbola and the graph of any other function  $g_{x'}$  can be obtained by a translation. Obviously  $f(y) \le g_x(y)$  on  $\Gamma_0$  and  $g_x(y(x)) = f(y(x))$ . If at every point  $y \in \Gamma_0$  the graph of f can be touched from above by some hyperbola  $g_x(y)$  then we will get the surjectivity of y(x). To obtain this the following will be enough

$$g_x''(y) > L_f', \text{ for all } |y - x| \le D\delta.$$
 (4)

So we arrive at

$$\delta < \frac{L}{L'_f (1 + D^2)^{\frac{3}{2}}}. (5)$$

Note that also uniqueness of y(x) follows from (4); assume we have y(x) and  $\tilde{y}(x)$ , then

$$|L'_f|y(x) - \tilde{y}(x)| < \left| \int_{y(x)}^{\tilde{y}(x)} g''_x(t)dt \right| = |f'(y(x)) - f'(\tilde{y}(x))| \le L'_f|y(x) - \tilde{y}(x)|.$$

We have used here that

$$f'(y(x)) = g'_x(y(x)) = \frac{L(y(x) - x)}{\sqrt{\delta^2 + (y(x) - x)^2}}$$
(6)

(derivatives in y at the point y(x)).

The injectivity of the map y(x) follows from differentiability of f. Assume  $y_0 = y(x) = y(\tilde{x})$ , so we have

$$f(y_0) = g_x(y_0) = g_{\tilde{x}}(y_0).$$

On the other hand  $f(y) \leq \min(g_x(y), g_{\tilde{x}}(y))$  this contradicts differentiability of f at  $y_0$ .

The monotonicity of y(x) can be obtained using same arguments; if  $x < \tilde{x}$  then the 'left' hyperbola  $g_x(y)$  touches the graph of f 'lefter' than the 'right' hyperbola  $g_{\tilde{x}}(y)$ , since both hyperbolas are above the graph of f.

Now we will prove that y(x) is Lipschitz. From (6) it follows that

$$y(x) - x = \frac{\delta f'(y(x))}{\sqrt{L^2 - (f'(y(x)))^2}}.$$
 (7)

Taking Y(x) := y(x) - x we can rewrite this as

$$Y(x) = \frac{\delta f'(Y(x) + x)}{\sqrt{L^2 - (f'(Y(x) + x))^2}} = \delta \Phi(f'(Y(x) + x)), \tag{8}$$

where  $\Phi(t)=\frac{t}{\sqrt{L^2-t^2}}$ . For  $\delta<\frac{(L^2-L_f^2)^{\frac{3}{2}}}{L^2L_f'}$  we can use Banach's fix point theorem and get that this functional equation has unique continuous solution. On the other hand it is not difficult to check that

$$\left| \frac{Y(x_2) - Y(x_1)}{x_2 - x_1} \right| \le \frac{\delta C}{1 - \delta C},$$

where 
$$C = \frac{L^2 L_f^2}{(L^2 - L_f^2)^{\frac{3}{2}}}$$
.

**Corollary 3.** If  $\delta$  is as small as in the previous Lemma, then the function u is  $\infty$ -harmonic in the strip between  $\Gamma_0$  and  $\Gamma_{\delta}$ .

**Proof.** This follows from the fact that if we take the strip with boundary values f on  $\Gamma_0$  and u on  $\Gamma_\delta$  then McShane-Whitney's minimal and maximal solutions will coincide, obviously with u.

Remark 4. We can rewrite (7) in the form

$$x(y) = y - \frac{\delta f'(y)}{\sqrt{L^2 - (f'(y))^2}},\tag{9}$$

where x(y) is the inverse of y(x). This together with (3) gives us the following

$$u(x(y)) = f(y) - \frac{\delta L^2}{\sqrt{L^2 - (f'(y))^2}}$$

Using the recent result of O.Savin that u is  $C^1$ , we conclude that function x(y) is as regular as f', so we cannot expect to have better regularity than Lipschitz.

**Lemma 5.** If  $\delta > 0$  is as small as above and function f is not twice differentiable at  $y_0$ , then the function u is not twice differentiable at  $x_0 := x(y_0)$ .

**Proof.** First note that for all x and y, such that x = x(y) we have

$$u'(x) = f'(y).$$

This can be checked analytically but actually is a trivial geometrical fact; the hyperbola 'slides' in the direction of the growth of f at point y, thus the cone which generates this hyperbola and 'draws' with its peak the graph of u moves in same direction which is the direction of the growth of u at point x = x(y).

Now assume we have two sequences  $y_k \to y_0$  and  $\tilde{y}_k \to y_0$  such that

$$\frac{f'(y_k) - f'(y_0)}{y_k - y_0} \to \underline{f''}(y_0) \text{ and } \frac{f'(\tilde{y}_k) - f'(y_0)}{\tilde{y}_k - y_0} \to \overline{f''}(y_0)$$

and  $\underline{f''}(y_0) < \overline{f''}(y_0)$ . Let us define appropriate sequences on  $\Gamma_\delta$  denoting by  $x_k := x(y_k)$  and by  $\tilde{x}_k := x(\tilde{y}_k)$  and compute the limits of

$$\frac{u'(x_k) - u'(x_0)}{x_k - x_0}$$
 and  $\frac{u'(\tilde{x}_k) - u'(x_0)}{\tilde{x}_k - x_0}$ .

We have

$$\frac{u'(x_k) - u'(x_0)}{x_k - x_0} = \frac{f'(y_k) - f'(y_0)}{y_k - y_0} \frac{y_k - y_0}{x_k - x_0}$$

the first multiplier converges to  $\underline{f''}(y_0)$ , let us compute the limit of the second one. From (9) we get that

$$\frac{x_k - x_0}{y_k - y_0} \to 1 - \delta \Phi'(f'(y_0)) \underline{f''}(y_0),$$

where  $\Phi(t) = \frac{t}{\sqrt{L^2 - t^2}}$ . Thus

$$\frac{u'(x_k) - u'(x_0)}{x_k - x_0} \to \frac{\underline{f''}(y_0)}{1 - \delta \Phi'(f'(y_0))\underline{f''}(y_0)},$$

and analogously

$$\frac{u'(\tilde{x}_k) - u'(x_0)}{\tilde{x}_k - x_0} \to \frac{\overline{f''}(y_0)}{1 - \delta \Phi'(f'(y_0))\overline{f''}(y_0)}.$$

To complete the proof we need to use the monotonicity of the function

$$\frac{t}{1-\delta Ct}, \ -L_f' < t < L_f',$$
 where  $\frac{1}{L} < C < \frac{L^2}{(L^2-L_f^2)^{\frac{3}{2}}}.$ 

We would like to note that if the function f is not  $C^2$  at a point y then u constructed here is not  $C^2$  on the whole line connecting y and x(y). So choosing f to be not twice differentiable on a dense set we can get a function u which is not  $C^2$  on the collection of corresponding line-segments. Note that a similar example is the distance function from a convex set, whose boundary is  $C^1$  and not  $C^2$  on a dense subset. Then the distance function is  $\infty$ -harmonic and is not  $C^2$  on appropriate lines.

Our example has the property of having constant  $|\nabla u|$  on gradient flow curves (lines in our case). It would be interesting to find a general answer to the question:

'What geometry do the gradient flow curves have, on which  $|\nabla u|$  is not constant?'

From Aronsson's results we know that u is not  $C^2$  on such a curve. This is our motivation for the investigation of  $C^2$ -differentiability of  $\infty$ -harmonic functions.

The author has only one item in the list of references. The history and the recent developments of the theory of  $\infty$ -harmonic functions, as well as a complete reference list could be found in that paper.

**Acknowledgment.** The author is grateful to Gunnar Aronsson, Michael Crandall and Arshak Petrosyan for valuable discussions.

## References

[ACJ] G. Aronsson, M. Crandall, P. Juutinen A tour of the theory of absolutely minimizing functions Bull. Amer. Math. Soc. 41 439–505

Hayk Mikayelyan, Mathematisches Institut, Universität Leipzig, Augustusplatz  $10/11,\,04109$  Leipzig, Germany

 $E ext{-}mail\ address: hayk@math.uni-leipzig.de}$