# FORMALITY OF k-CONNECTED SPACES IN 4k + 3 AND 4k + 4 DIMENSIONS

### GIL RAMOS CAVALCANTI

ABSTRACT. Using the concept of s-formality we are able to extend the bounds of a Theorem of Miller and show that a compact k-connected (4k+3)- or (4k+4)-manifold with  $b_{k+1}=1$  is formal. We study k connected n-manifolds, n=4k+3,4k+4, with a hard Lefschetz-like property and prove that in this case if  $b_{k+1}=2$ , then the manifold is formal, while, in 4k+3-dimensions, if  $b_{k+1}=3$  all Massey products vanish. We finish with examples inspired by symplectic geometry and manifolds with special holonomy.

## Introduction

A theorem of Miller [13] states that any compact orientable k-connected manifold of dimension  $d \leq (4k+2)$  is formal. In particular, a compact simply-connected n-manifold is formal if  $n \leq 6$ . Recently, Fernández and Muñoz gave examples of simply-connected nonformal 7- and 8-manifolds [9] and Dranishnikov and Rudyak gave examples of k-connected nonformal manifolds in any dimension greater than 4k+2 [6], therefore proving that Miller's theorem can not be improved without further hypotheses.

Here, we adopt the point of view that for a k-connected manifold, the smaller the  $k + 1^{th}$  Betti number,  $b_{k+1}$ , is, the simpler the topology. Then we establish the biggest value of  $b_{k+1}$  for which one can still assure formality of an n-manifold, n = 4k + 3, 4k + 4:

**Theorem 1.** A compact orientable k-connected manifold of dimension 4k + 3 or 4k + 4 with  $b_{k+1} = 1$  is formal.

One motivation for the study of such manifolds comes from the existence of special geometric structures in 7- and 8-manifold, e.g.,  $G_2$ , Spin(7) and symplectic structures. A compact irreducible Riemannian manifold with holonomy group  $G_2$  or Spin(7) has finite fundamental group and hence its universal cover is compact, simply-connected and has special holonomy. Other examples of manifolds with special holonomy are Kähler manifolds which are formal [4], hence it is conceivable that there is a connexion between the existence of special holonomy metrics and formality.

Another reason to study formality of 8-manifolds comes from symplectic geometry, as this is the only dimension where the question of existence of compact 1-connected nonformal examples is still open.

Date: November 14, 2004.

Researcher supported by CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Ministério da Educação e Cultura), Brazilian Government. Grant 1326/99-6.

One property shared by irreducible  $G_2$ - and Spin(7)-manifolds is that they have a hard Lefschetz-like property. If  $M^n$  is one such manifold, there is a closed (n-4)-form  $\varphi$  for which:

(1) 
$$[\varphi] \cup : H^2(M, \mathbb{R}) \stackrel{\cong}{\to} H^{n-2}(M, \mathbb{R}),$$

is an isomorphism [11]. Of course, this last property is also shared by 2-Lefschetz symplectic 8-manifolds, where, by definition, (1) holds with  $\varphi = \omega^2$ .

We are interested in how (1) can be used to improve on Theorem 1. So we prove that if  $M^n$ , n=4k+3,4k+4, is k-connected and there is  $\varphi \in H^{n-2k-2}(M)$  for which

(2) 
$$\varphi \cup : H^{k+1}(M, \mathbb{R}) \stackrel{\cong}{\to} H^{n-k-1}(M, \mathbb{R})$$

is an isomorphism and  $b_{k+1}(M) = 2$ , then, M is formal. If n = 4k + 3 and  $b_{k+1}(M) = 3$ , then all Massey products on M vanish uniformly. We construct examples showing that our bounds are optimal.

This paper is organized as follows. In the first section we cover background material: introduce minimal models, formality, s-formality and Massey products. The proof that every compact 1-connected manifold has a minimal model is given to fix notation. In Section 2, we prove Theorem 1 by explicitly constructing a minimal model. In Section 3, we prove that the existence of the isomorphism (2) has a 'formalizing tendency' in a k-connected orientable 4k+3-or 4k+4-manifold, as it implies formality for  $b_2=2$  and, in the 4k+3-dimensional case, vanishing of Massey products for  $b_2=3$ . In the last section we study examples from symplectic and Riemannian geometry where our results can be applied.

I would like to thank Nigel Hitchin for his guidance and Marisa Fernández and Yuli Rudyak for their suggestions and encouragement.

# 1. Minimal models and s-formality

In this section we introduce the basic concepts we are interested in: differential graded algebras, minimal models, formality and s-formality. All the results are standard (see [10, 7]). Some proofs are given in order to fix notation.

**Definition.** A differential graded algebra, or DGA for short, is an  $\mathbb{N}$  graded vector space  $\mathcal{A}^{\bullet}$  over a field k, endowed with a product and a differential d satisfying:

(1) The product maps  $\mathcal{A}^i \times \mathcal{A}^j$  to  $\mathcal{A}^{i+j}$  and is graded commutative:

$$a \cdot b = (-1)^{ij}b \cdot a;$$

- (2) The differential has degree 1:  $d: A^i \to A^{i+1}$ ;
- (3)  $d^2 = 0$ ;
- (4) The differential is a derivation: for  $a \in \mathcal{A}^i$  and  $b \in \mathcal{A}^j$

$$d(a \cdot b) = da \cdot b + (-1)^i a \cdot db.$$

We will deal only with DGAs over  $\mathbb{R}$ . A nontrivial example of DGA is the complex of differential forms on a manifold equipped with the exterior derivative. Another example is the cohomology of a manifold, with d=0 as differential. The cohomology of a DGA is defined in the standard way.

**Definition.** A DGA  $(\mathcal{M}, d)$  is *minimal* if it is free as a DGA (i.e. polynomial in even degree and skew symmetric in odd degree) and has generators  $e_1, e_2, \ldots, e_n, \ldots$  such that:

- (1) The degree of the generators form a weakly increasing sequence of positive numbers;
- (2) There are finitely many generators in each degree;
- (3) The differential satisfies  $de_i \in \land \{e_1, \ldots, e_{i-1}\}.$

A minimal model for a differential graded algebra (A, d) is a minimal DGA,  $\mathcal{M}$ , together with a map of differential graded algebras,  $\rho : \mathcal{M} \to \mathcal{A}$ , inducing isomorphism in cohomology. If (A, d) is the complex of differential forms on a manifold M we also refer to the minimal model of  $\mathcal{A}$  as the minimal model of M.

Every simply-connected compact manifold has a minimal model which is unique up to isomorphism. As the cohomology algebra of a manifold with real coefficients is also a DGA we can also construct its minimal model. This minimal model is not necessarily the same as the minimal model for the manifold and in general it carries less information.

**Definition.** A manifold is *formal* if the minimal models for M and for its cohomology algebra are isomorphic, or equivalently, if there is a map of differential graded algebras  $\psi : \mathcal{M} \to H^{\bullet}(M, \mathbb{R})$  inducing isomorphism in cohomology, where  $\mathcal{M}$  is the minimal model of M.

The proof that every 1-connected manifold has a minimal model is done in a constructive way and the fundamental tool is a Hirsch extension. If  $\mathcal{A}$  is a differential algebra, V a finite dimensional vector space and  $d: V \to \ker(d) \cap (\mathcal{A}^{k+1})$  a linear map, we can form the Hirsch extension  $\mathcal{A} \otimes_d (\wedge V)_k$  as follows: the elements of  $\mathcal{A} \otimes_d (\wedge V)_k$  are elements of the tensor product of  $\mathcal{A}$  with the free algebra generated by V in degree k, the differential is defined by linearity plus the product rule  $d(a \otimes v) = da \otimes v + (-1)^{\deg(a)} a \wedge dv \otimes 1$ . We denote  $1 \otimes v$  by v and  $a \otimes 1$  by a and then  $a \otimes v$  becomes simply  $a \wedge v$ .

In doing such an extension one is, amongst other things,

- i) killing cohomology in degree k+1 since the elements in the image of d that were closed in  $\mathcal{A}$ , but not exact, become exact in  $\mathcal{A} \otimes_d (\wedge V)_k$ ;
- ii) creating cohomology in degree k, namely, the new classes are given by the kernel of  $d: V \to \mathcal{A}^{k+1}/\mathrm{Im}\,(d: \mathcal{A}^k \to \mathcal{A}^{k+1})$ .

Therefore, we can always split  $V = B^k \oplus Z^k$ , where the map  $d: B^k \to \mathcal{A}^{k+1}/\mathrm{Im}\,(d: \mathcal{A}^k \to \mathcal{A}^{k+1})$  is an injection and  $Z_k = \ker(d: V \to \mathcal{A}^{k+1})/\mathrm{Im}\,(d: \mathcal{A}^k \to \mathcal{A}^{k+1})$ , although this splitting is not canonical.

**Theorem 2.** (Sullivan [15]) Every compact simply-connected manifold has a minimal model.

*Proof.* The proof is accomplished by constructing the minimal model, which is done by induction. The starting point is a free DGA,  $\mathcal{M}_2$ , generated by  $H^2(M)$  in degree 2 and with vanishing differential. We define  $\rho_2: \mathcal{M}_2^{\bullet} \to \Omega^{\bullet}(M)$  by choosing arbitrary representatives for the cohomology classes in  $H^2(M)$  and extend it to higher symmetric powers of  $H^2(M)$  so that it is a map of algebras.

Now assume we have a free DGA,  $\mathcal{M}_k$ , generated by elements of degree at most k with a map of DGAs  $\rho_k : \mathcal{M}_k^{\bullet} \to \Omega^{\bullet}(M)$  such that  $\rho_k^* : H^i(\mathcal{M}_k) \to H^i(M)$  is an isomorphism for  $i \leq k$ 

and an injection for i = k + 1. Let

$$V^{k+1} = \ker(\rho_k^* : H^{k+2}(\mathcal{M}_k) \to H^{k+2}(M)) \oplus H^{k+1}(M)/\rho_k^* (H^{k+1}(\mathcal{M}_k)).$$

Choosing linearly a representative v for each cohomology class

$$[v] \in \ker(\rho_k^* : H^{k+2}(\mathcal{M}_k) \to H^{k+2}(M)),$$

define  $\mathcal{M}_{k+1}$  to be the Hirsch extension of  $\mathcal{M}_k$  by  $V^{k+1}$ , with d given by:

$$\begin{cases} d[v] = v & \text{if } [v] \in \ker(\rho_k^* : H^{k+2}(\mathcal{M}_k) \to H^{k+2}(M), \\ dv = 0 & \text{if } v \in H^{k+1}(M)/\rho_k^*(H^{k+1}(\mathcal{M}_k)). \end{cases}$$

Finally, we let  $\rho_{k+1}|_{\mathcal{M}_k} = \rho_k$  and define it in  $V^{k+1}$  in the following way. For  $[v] \in \ker(\rho_k^* : H^{k+2}(\mathcal{M}_k) \to H^{k+2}(M))$ , by definition,  $\rho_k(d[v]) = da$ , for some k+1-form a. We let  $\rho_{k+1}([v]) = a$ . For  $v \in H^{k+1}(M)/\rho_k^*(H^{k+1}(\mathcal{M}_k))$ , we choose a closed form b whose cohomology class represents the same element and let  $\rho_{k+1}(v) = b$ . These choices can be made so the  $\rho_{k+1}: V^{k+1} \to \Omega^{k+1}(M)$  is linear. The map  $\rho_{k+1}$  is extended to the rest of  $\mathcal{M}_{k+1}$  by requiring it to be a map of DGAs.

With these choices,  $\mathcal{M}_{k+1}$  is a free DGA generated by elements of degree at most k+1 with a map of DGAs  $\rho_{k+1}: \mathcal{M}_k \to \Omega^{\bullet}(M)$  inducing an isomorphism  $\rho_{k+1}^*: H^i(\mathcal{M}_{k+1}) \to H^i(M)$  for  $i \leq k+1$  and an injection for i=k+2 and the inductive step is complete.

This procedure gives us a family of free DGAs, each obtained by a Hirsch extension from the previous one:

$$\mathcal{M}_2 < \mathcal{M}_3 < \cdots \mathcal{M}_k < \cdots,$$

and maps  $\rho_k: \mathcal{M}_k \to \Omega^{\bullet}(M)$  such that:

- i)  $\rho_k|\mathcal{M}_i = \rho_i$  for  $i \leq k$ , hence induce a map in the limit DGA  $\rho: \mathcal{M} \to \Omega^{\bullet}(M)$ ;
- ii) The induced maps in cohomology are isomorphisms  $\rho^*: H^i(\mathcal{M}_k) \cong H^i(M)$ , for  $i \leq k$  and therefore, in the limit,  $H^i(\mathcal{M}) \cong H^i(M)$  for all i.

Thus the limit  $\mathcal{M}$  of the  $\mathcal{M}_k$  is the minimal model for M.

The main reason to present the proof of this theorem is that it is a constructive one and sometimes properties about the minimal model are more easily obtained using this point of view. For example we have an alternative description of formality (see [7, 10], for a proof).

**Proposition 1.** Let  $V^k$  be the spaces introduced in degree k when constructing the minimal model of a manifold M. M is formal if and only if there is a splitting  $V^k = B^k \oplus Z^k$ , such that

- (1)  $d: B^k \to \mathcal{M}^{k+1}$  is an injection;
- $(2) \ d(Z^k) = 0;$
- (3) If an element of the ideal  $\mathcal{I}(\bigoplus_{k\in\mathbb{N}}B^k)<\mathcal{M}$  is closed, then it is exact.

This characterization of formality allows one to consider weaker versions. Notably, Fernández and Muñoz introduced in [8] the useful concept of s-formality.

**Definition.** A manifold is s-formal if there is a choice of splitting  $V^k = Z^k \oplus B^k$  satisfying (1) and (2) above and such that every closed element in the ideal  $\mathcal{I}(\oplus_{k \leq s} B^k) < \mathcal{M}_s$  is exact in  $\mathcal{M}$ .

Clearly this is a weaker concept, in general, but it is also obvious that if an n-dimensional manifold is n-formal, it is formal. The following result of Fernández and Muñoz shows that sometimes the weaker condition of s-formality implies formality.

**Theorem 3.** (Fernández and Muñoz [8]): If a compact orientable manifold  $M^n$  is s-formal for  $s \ge n/2 - 1$ , then M is formal.

The point of this theorem is that in order to prove formality by constructing the minimal model and finding the splitting as in Proposition 1, it is only necessary to determine the beginning of the minimal model.

So far we have two ways to tell whether a manifold is formal: from the definition of formality and from a (partial) construction of the minimal model and the theorem of Fernández and Muñoz. Next we introduce Massey products, which are an obstruction to formality. The ingredients are  $a_{12}, a_{23}, a_{34} \in \mathcal{A}$  three closed elements such that  $a_{12}a_{23}$  and  $a_{23}a_{34}$  are exact. Then, denoting  $\bar{a} = (-1)^{|a|}a$ , we define

$$\begin{cases} \overline{a_{12}}a_{23} &= da_{13} \\ \overline{a_{23}}a_{34} &= da_{24}. \end{cases}$$

In this case, one can consider the element  $\overline{a_{13}}a_{34} + \overline{a_{12}}a_{24}$ . By the choice of  $a_{13}$  and  $a_{24}$  this element is closed, hence it represents a cohomology class. Observe, however, that  $a_{13}$  and  $a_{24}$  are not well defined and we can change them by any closed element, hence the expression above does not define a unique cohomology class but instead an element in the quotient  $H^{\bullet}(\mathcal{A})/\mathcal{I}([a_1], [a_3])$ .

**Definition.** The Massey product or triple Massey product  $\langle [a_{12}], [a_{23}], [a_{34}] \rangle$ , of the cohomology classes  $[a_{12}], [a_{23}]$  and  $[a_{34}]$  with  $[a_{12}][a_{23}] = [a_{23}][a_{34}] = 0$  is the coset

$$\langle [a_{12}], [a_{23}], [a_{34}] \rangle = [\overline{a_{12}}a_{24} + \overline{a_{13}}a_{34}] + ([a_{12}], [a_{34}]) \in H^{\bullet}(\mathcal{A})/\mathcal{I}([a_{12}], [a_{34}]),$$

where  $a_{13}$  and  $a_{24}$  are defined by (\*). We say that all the Massey products vanish uniformily if there is a fixed set of choices for which all possible Massey products are represented by exact forms.

The importance of Massey products for this work comes from the following result: If M is formal, all the Massey products vanish uniformily.

## 2. Extending Miller's bounds

In this section we prove Theorem 1. The way to prove formality in this case is by constructing the beginning of the minimal model for the manifold and then using the theorem of Fernández and Muñoz (Theorem 3 above).

PROOF OF THEOREM 1. We will only prove the theorem for 4k + 4-manifolds, as the other case is analogous. The proof is accomplished by constructing the minimal model.

As M is k-connected,  $\mathcal{M}_{2k}$  is a free algebra generated by  $H^i(M)$  in degree  $i \leq 2k$  with vanishing differential and  $\rho$  maps linearly each cohomology class to a representative form. The first time we may have to use a nonzero differential (and hence introduce one of the  $B^j$  spaces) is in degree 2k + 1. This will be the case only if the generator  $a \in \mathcal{M}_{k+1}$  satisfies  $a^2 \neq 0$  and

 $\rho(a^2)$  is an exact form. Hence if either  $\rho(a^2)$  is not exact or  $a^2 = 0$ , all the spaces  $B^j$  are trivial for  $j \leq 2k + 1$ , showing that M is 2k + 1-formal and hence, by Theorem 3, formal.

So we only have to consider the case where the cohomology class  $\alpha \in H^{k+1}(M)$  satisfies  $\alpha^2 = 0$  and k+1 is even. In this case, the Hirsch extension in degree 2k+1 is given by

$$V^{2k+1} = H^{2k+1}(M) \oplus \operatorname{span}\{b\},\$$

where d vanishes in  $H^{2k+1}(M)$ ,  $db = a^2$  and  $\rho(b)$  is a form such that  $d\rho(b) = \rho(a^2)$ . With this splitting,  $\mathcal{I}(\bigoplus_{j \leq 2k+1} (B^j)) < \mathcal{M}_{2k+1}$  is just the ideal generated by b and to prove formality using Theorem 3 we have to show that any closed form in this ideal is exact in  $\mathcal{M}$ .

A closed form in this ideal, being the product of b and an element of degree at least k+1, will have degree at least 3k+2. Since M is k-connected, Poincaré duality gives  $H^j(M) = \{0\}$  for 3k+3 < j < 4k+4. If an element in  $\mathcal{I}(b)$  of degree 3k+2 or 3k+3 is closed and nonexact in  $\mathcal{M}$ , Poincaré duality implies that its dual is in either  $\mathcal{M}_{k+2}^{k+2} \cong H^{k+2}(M)$  (in the former case) or in  $\mathcal{M}_{k+1}^{k+1} \cong H^{k+1}(M)$  (in the latter). Either way, from there we can produce a degree 4k+4 closed nonexact element in  $\mathcal{I}(b)$ . So we only have to check that any 4k+4 closed element in  $\mathcal{I}(b)$  is exact.

The only elements in  $\mathcal{I}(b)$  in degree 4k+4 are of the form abv, with  $v \in H^{k+2}(M) \cong V^{k+2}$ , which have derivative  $d(abv) = a^3v$  and hence are not closed if  $v \neq 0$ . This shows that M is (2k+1)-formal and therefore formal.

### 3. Formality of hard Lefschetz manifolds

In this section we study compact orientable k-connected n-manifolds, n = 4k + 3, 4k + 4, for which there is a cohomology class  $\varphi \in H^{n-2k-2}(M)$  inducing an isomorphism

(2) 
$$\varphi \cup : H^{k+1}(M) \stackrel{\cong}{\to} H^{n-k-1}(M).$$

We prove that this property has a 'formalizing tendency' in the following sense.

**Theorem 4.** A compact orientable k-connected manifold  $M^n$ , n = 4k + 3, 4k + 4, satisfying (2) with  $b_{k+1} = 2$  is formal. If n = 4k + 3 and  $b_{k+1} = 3$  all the Massey products vanish uniformly.

*Proof.* We start considering the case  $b_{k+1} = 2$ . The cases n = 4k + 3 and n = 4k + 4 are similar, so we only deal with the latter. Due to (2), the class  $\varphi$  induces a nondegenerate bilinear form on  $H^{k+1}(M)$ . If k is even, this bilinear form is skew, and if k is odd, it is symmetric.

For k odd, the signature of the bilinear form, i.e., the difference between the number of positive and negative eigenvalues, is either 2, 0 or -2. By changing  $\varphi$  to  $-\varphi$ , the case of signature -2 can be transformed into signature 2, hence there are two possibilities to consider. As they are similar, we will only treat the signature 2 case.

Let  $a_1, a_2$  ( $da_1 = da_2 = 0$ ) be generators of  $\mathcal{M}_{k+1} = Sym^{\bullet}H^{k+1}(M)$ , the first nontrivial stage of the minimal model for M. We may further assume that the bilinear form induced by  $\varphi$  is diagonal in the basis  $\{a_1, a_2\}$  and

$$\int \varphi a_i^2 = 1.$$

As with Theorem 1,  $\mathcal{M}^{2k}$  is just the free algebra generated by  $H^j(M)$  in degree  $j \leq k$  with vanishing differential and the first time we may have to introduce one of the  $B^j$  spaces is in

degree 2k + 1, where

$$B^{2k+1} \cong \ker Sym^2H^{k+1}(M) \stackrel{\cup}{\to} H^{2k+2}(M).$$

We know that both  $a_1^2$  and  $a_2^2$  are nonzero in  $H^{2k+2}(M)$ , since by (2) they pair nontrivially with  $\varphi$ . So

$$\dim B^{2k+1} = \dim(\ker Sym^2H^{k+1}(M) \xrightarrow{\cup} H^{2k+2}(M)) \le \dim(Sym^2H^{k+1}(M)) - 1 = 2,$$

and hence there may be at most two generators in degree 2k + 1 in  $\mathcal{M}_{2k+1}$  to kill cohomology classes in  $\mathcal{M}_{2k}$  that are not present in  $H^{2k+2}(M)$ . The case when only one generator is added has a proof very similar to the one of Theorem 1, so we move on to the case when there are two generators  $b_1$  and  $b_2$  added to kill cohomology in degree 2k + 2.

Then

$$db_i = k_{11}^i a_1^2 + k_{12}^i a_1 a_2 + k_{22}^i a_2^2, \qquad i = 1, 2.$$

Multiplying by  $\varphi$  and integrating we get  $k_{11}^i = -k_{22}^i$ . Hence, by re-scaling and taking linear combinations we may assume that  $db_1 = a_1^2 - a_2^2$  and  $db_2 = a_1a_2$ .

Now, if  $c \in \mathcal{I}(b_1, b_2)_{\leq 2k+1}$  is a closed element (again, by Poincaré duality we may assume it has degree n) we can write it as

$$c = (a_1c_{11} + a_2c_{12})b_1 + (a_1c_{21} + a_2c_{22})b_2 + (k_1a_1 + k_2a_2)b_1b_2,$$

where  $c_{ij}$  are closed elements of degree k+2 and  $k_i \in \mathbb{R}$ . The condition dc=0 implies that c=0, thus every closed form in  $(b_1,b_2)_{\leq 2k+1}$  is exact and M is 2k+1-formal and therefore formal.

The case when k is even is easier. Indeed, the argument above can be used again, but with  $Sum^2H^{k+1}(M)$  replaced by  $\wedge^2H^{k+1}(M)$  and hence the nondegeneracy of the pairing implies

$$\dim B^{2k+1} = \dim(\ker \wedge^2 H^{k+1}(M) \xrightarrow{\cup} H^{2k+2}(M)) \le \dim(\wedge^2 H^{k+1}(M)) - 1 = 0,$$

hence M is trivially 2k + 1-formal and therefore formal.

To finish the proof, we consider a 4k + 3-manifold with  $b_{k+1} = 3$  and prove that all triple Massey products vanish if (2) holds. Initially we remark that  $b_{k+1} = 3$  and (2) can not happen if k is even, as there is no nondegenerate skew bilinear form in an odd dimensional vector space.

We also observe that the Massey product  $\langle a, b, c \rangle$  has degree at least 3k + 2 and since  $H^{j}(M) = \{0\}$ , for 3k + 2 < j < 4k + 3, this product will vanish, whenever defined, if its degree is neither 3k + 2 nor 4k + 3. If  $\langle a, b, c \rangle \in H^{4k+3}(M)$ , it lies in the ideal generated by ([a], [c]), so the product also vanishes, therefore the only case left is when a, b and c have degree k + 1 and the product lies in  $H^{3k+2}(M)$ .

This product vanishes trivially if  $c = \lambda a$ , hence we can assume a and c linearly independent. Since  $\varphi ab$  and  $\varphi bc$  vanish, but  $\varphi$  induces a nondegenerate bilinear form, there is  $\beta \in H^{k+1}(M)$  such that  $\varphi b\beta \neq 0$  and we can express the Massey product in the basis  $\{\varphi[a], \varphi[\beta], \varphi[c]\}$ :

$$\langle a, b, c \rangle = -v_1c + av_2 = k_1\varphi[a] + k_2\varphi[\beta] + k_3\varphi[c].$$

where  $ab = dv_1$  and  $bc = dv_2$ . Multiplying the equation above by -[b] and integrating over M we get

$$-k_2 \int \varphi b\beta = \int -v_1 cb + av_2 b = \int -v_1 dv_2 + v_2 dv_1 = \int d(v_1 v_2) = 0.$$

Thus  $k_2 = 0$  and  $\langle a, b, c \rangle = k_1 \varphi[a] + k_3 \varphi[c] \in ([a], [c])$ . So the Massey product vanishes.

Observe that if  $ac = dv_3$ , it is still possible to choose  $v_i$ , i = 1, 2, 3 in such a way that the Massey products  $\langle a, b, c \rangle$ ,  $\langle b, c, a \rangle$  and  $\langle c, a, b \rangle$  vanish simultaneously. Indeed, let us assume a, b and c are linearly independent (the linearly dependent case is similar) and a set of choices of  $v_i$  was made:

$$\begin{cases} \langle a, b, c \rangle = [v_1c - av_2] = k_1\varphi[a] + k_2\varphi[c], \\ \langle b, c, a \rangle = [v_2a - bv_3] = l_1\varphi[b] + l_2\varphi[a], \\ \langle c, a, b \rangle = [v_3b - cv_1] = m_1\varphi[c] + m_2\varphi[b]. \end{cases}$$

Then we can set  $v_1 = \tilde{v}_1 + k_2 \varphi$ ,  $v_2 = \tilde{v}_2 + l_2 \varphi$  and  $v_3 = \tilde{v}_3 + m_2 \varphi$ . With these choices we get

$$\begin{cases} \langle a,b,c\rangle = [\tilde{v}_1c - a\tilde{v}_2] = \tilde{k}\varphi[a], \\ \langle b,c,a\rangle = [\tilde{v}_2a - b\tilde{v}_3] = \tilde{l}\varphi[b], \\ \langle c,a,b\rangle = [\tilde{v}_3b - c\tilde{v}_1] = \tilde{m}\varphi[c]. \end{cases}$$

Adding them up, the left hand side vanishes, giving

$$0 = \tilde{k}\varphi[a] + \tilde{l}\varphi[b] + \tilde{m}\varphi[c].$$

Since  $\varphi[a]$ ,  $\varphi[b]$  and  $\varphi[c]$  are linearly independent,  $\tilde{k}$ ,  $\tilde{l}$  and  $\tilde{m}$  vanish and, with these choices, all the Massey products vanish *simultaneously*, hinting at formality.

# 4. Examples

In this section give simple examples showing that the bounds established in Theorem 1 are sharp. We also apply our results to the blow-up of  $\mathbb{C}P^8$  along a symplectic submanifold and to Kovalev's examples of  $G_2$ -manifolds. We finish with an example of a compact 1-connected 7-manifold which satisfies all known topological restrictions imposed by a  $G_2$  structure, has  $b_2 = 4$  and nonvanishing Massey products. This last example shows that the results of Theorem 4 are sharp in 7 dimensions and that one can not answer the question of formality of  $G_2$ -manifolds using only the currently known topological properties of those.

**Example 1.** In this example we show that the bounds obtained in Theorem 1 are sharp in two ways: we construct nonformal compact k-connected manifolds  $M^n$  with  $b_{k+1} = 2$ , for n = 4k + 3, 4k + 4 and with  $b_{k+1} = 1$  for n > 4k + 4.

We shall start dealing with odd dimensional examples. Let  $k_1 \leq k_2 \in \mathbb{N}$  be two natural numbers such that  $k_1 + k_2$  is even and let V be a  $(k_1 + k_2 + 2)$ -dimensional vector bundle over  $S^{k_1+1} \times S^{k_2+1}$  whose Euler class  $\chi$  is a nonvanishing top degree cohomology class. Let  $X^{2k_1+2k_2+3}$  be the total space of the sphere bundle of V. We claim that X has a nontrivial Massey product. Indeed, using the Gysin sequence for this sphere bundle

$$\cdots \xrightarrow{\chi \cup} H^{j}(S^{k_{1}+1} \times S^{k_{2}+1}) \xrightarrow{\pi^{*}} H^{j}(X) \xrightarrow{\pi_{*}} H^{j-k_{1}-k_{2}-1}(S^{k_{1}+1} \times S^{k_{2}+1}) \xrightarrow{\chi \cup} H^{j+1}(S^{k_{1}+1} \times S^{k_{2}+1}) \rightarrow \cdots$$

we see that  $H^{k_1+1}(X) = \operatorname{span}\{v_1\}$ ,  $H^{k_2+1}(X) = \operatorname{span}\{v_2\}$  and  $H^{k_1+k_2+2}(X) = \{0\}$ , where  $v_i$  are generators for the top degree cohomology of each sphere. Therefore,  $v_1 \cup v_2 = 0$ . If  $\omega_i$  are volume forms pulled back from each  $S^{k_i+1}$ , then  $\omega_1 \wedge \omega_2 = d\xi$ , where

$$\int_{S^{k_1+k_2+1}} \xi = \int_{S^{k_1+1} \times S^{k_2+1}} \chi.$$

Therefore we can compute the Massey product

$$\langle v_1, v_2, v_2 \rangle = -[\xi \wedge \omega_2].$$

This is not exact, as it pairs nontrivially with  $v_1$ ,

$$\int_{X} \xi \wedge \omega_{1} \wedge \omega_{2} = \int_{S^{k_{1}+1} \times S^{k_{2}+1}} \chi \cdot \int_{S^{k_{1}+1}} \omega_{1} \int_{S^{k_{2}+1}} \omega_{2} \neq 0.$$

And this Massey product has no indeterminacy, hence X is not formal.

It is clear that X is  $k_1$ -connected and has dimension  $2k_1 + 2k_2 + 3 \ge 4k_1 + 3$ . If  $k_1 < k_2$ , then  $b_{k_1+1}(X) = 1$ , and if  $k_1 = k_2 = k$ , then  $b_{k+1}(X) = 2$ .

To obtain even dimensional examples we consider  $k_1$  and  $k_2$  as before, but now we take a vector bundle V of rank  $k_1 + k_2 + 2$  over the manifold  $(S^{k_1+1} \times S^{k_2+1}) \# \mathbb{C}P^{k+1}$ , where # indicates connected sum and  $k = \frac{1}{2}(k_1 + k_2)$ . We can further assume that V is trivial outside a small disc. As before, the total space, X, of the sphere bundle of V yields a nonformal  $(2k_1 + 2k_2 + 3)$ -manifold. The total space of a principal circle bundle over X with Chern class equal to the generator of the second cohomology  $H^2(\mathbb{C}P^{k+1},\mathbb{Z})$  will be a nonformal  $k_1$ -connected  $(2k_1 + 2k_2 + 4)$ -manifold.

**Example 2.** Blowing up  $\mathbb{C}P^n$  along a suitable submanifold, Babenko and Taimanov proved in [2] that there are compact 1-connected nonformal symplectic manifolds in any dimension  $2k \geq 10$ . Due to Miller's theorem, such examples do not exist in dimensions 6 or less and the question is still open for 8-manifolds.

Using the techniques of [3], one can show that the blow-up, X, of  $\mathbb{C}P^n$  along any submanifold is always 2-Lefschetz and, if the submanifold is connected,  $b_2(X) = 2$ . Therefore, our results imply the blow-up of  $\mathbb{C}P^8$  along a connected symplectic submanifold is always formal. One can also try to blow up  $\mathbb{C}P^n$  and then take a sequence of Donaldson submanifolds until the result is an 8-manifold, but the manifold obtained this way is also 2-Lefschetz [8] and has  $b_2 = 2$  [5].

**Example 3.** In [12], Kovalev produces a series of examples of 1-connected compact  $G_2$ -manifolds from pairs of Fano 3-folds via twisted connected sums. If  $M^7$  is obtained from the Fanos  $F_1$  and  $F_2$ , he proves that  $b_2(M) \leq \min\{b_2(F_1), b_2(F_2)\} - 1$ . Our results imply that if either of the Fanos involved have  $b_2(F_i) \leq 3$ ,  $M^7$  is formal, while if  $b_2(F_1) = 4$  and  $b_2(F_2) \geq 4$ , the Massey products vanish. Hence the only possibility for one of his examples to have nontrivial Massey products is if it is constructed from 2 Fanos with  $b_2(F_i) \geq 5$ .

According to the classification of Fano 3-folds [14], these have  $b_2 \leq 10$  and if  $b_2 \geq 6$ , the Fano is just the blow-up of  $\mathbb{C}P^3$  in an appropriate number of points. It is easy to follow Kovalev's construction to prove that if one of the summands is  $\mathbb{C}P^3$  with some points blown-up,  $M^7$  is formal. The case where both the Fanos have  $b_2 = 5$  is more difficult, but one still has formality.

In [11], Joyce shows that a compact Riemannian manifold with holonomy  $G_2$  has finite  $\pi_1$ , nonvanishing first Pontryagin class  $p_1$  and, if  $\varphi$  is the closed 3-form determining the structure,

(3) 
$$\int a^2 \varphi < 0 \text{ for } a \in H^2(M) \setminus \{0\} \qquad \text{ and } \qquad \int p_1 \wedge [\varphi] < 0.$$

Using circle bundles we can construct a nonformal manifold satisfying all the topological properties above. The key is Wall's classification of 1-connected spin 6-manifolds.

**Theorem 5.** (Wall [16]) Diffeomorphism classes of oriented 6-manifolds with torsion-free homology and vanishing second Stiefel-Witney class correspond bijectively to isomorphism classes of systems of invariants:

- Two finitely generated free abelian groups  $H^2$ ,  $H^3$ , the latter of even rank;
- A symmetric trilinear map  $\mu: H^2 \times H^2 \times H^2 \to Z$ ;
- A homomorphism  $p_1: H^2 \to Z$ ;
- Subject to: for  $x, y \in H^2$ ,

$$\mu(x,x,y)=\mu(x,y,y) \mod (2),$$
 for  $x\in H^2,$  
$$p_1(x)=4\mu(x,x,x) \mod (24).$$

With appropriate choices for the pairing  $\mu$  and for the Chern class of the principal circle bundle, we can obtain nonformal 7-manifolds satisfying (3). As the base manifold is spin, so will be the total space of the circle bundle. We finish this paper with one example constructed this way.

**Example 4.** We let  $H^2 = \langle \omega, \alpha_1, \alpha_2, \alpha_3, \gamma \rangle$  and define the cup product on  $H^2$  so as to have the following relations

$$\omega \gamma = 0$$
,  $\omega \alpha_i \alpha_j = 2\delta_{ij}$ ,  $\omega^2 \alpha_i = 0$ ,  $\omega^3 = 2$ ,  $\omega \alpha_1 = \gamma \alpha_3$ ,  
 $\alpha_1 \alpha_2 = 0$ ,  $\alpha_2 \alpha_3 = \gamma \alpha_1$  and  $\alpha_3 \omega = \gamma \alpha_1$ .

One set of choices that gives the desired result is the following

$$\omega \alpha_i \alpha_j = 2\delta_{ij} \qquad \alpha_1 \alpha_2^2 = 0 \qquad \alpha_1^2 \alpha_2 = 0 \qquad \alpha_2 \alpha_3^2 = 2 \qquad \alpha_3 \gamma^2 = 0$$

$$\omega \gamma \alpha_i = 0 \qquad \alpha_1 \alpha_2 \alpha_3 = 0 \qquad \alpha_1^2 \alpha_3 = 0 \qquad \alpha_2 \alpha_3 \gamma = 0 \qquad \alpha_3^2 \gamma = 0$$

$$\omega \gamma^2 = 0 \qquad \alpha_1 \alpha_2 \gamma = 0 \qquad \alpha_1^2 \gamma = 0 \qquad \alpha_2 \gamma^2 = 2 \qquad \alpha_3^3 = 0$$

$$\omega^2 \alpha_i = 0 \qquad \alpha_1 \alpha_3^2 = 0 \qquad \alpha_1^3 = 0 \qquad \alpha_2^2 \alpha_3 = 0 \qquad \gamma^3 = 0$$

$$\omega^2 \gamma = 0 \qquad \alpha_1 \alpha_3 \gamma = 2 \qquad \alpha_2^2 \gamma = 2$$

$$\omega^3 = 2 \qquad \alpha_1 \gamma^2 = 0 \qquad \alpha_2^3 = 0$$

With these choices,  $\omega \gamma = 0$  and this is the only 2-cohomology class that pairs trivially with  $\gamma$ .

Now we let M be a simply connected spin 6-manifold with  $H^2(M) = H^2$ , cup product as described above, arbitrary  $H^3$  and first Pontryagin class  $p_1 = 4\omega^2$ . Let  $w, a_i, c_1$  be a set of closed forms representing  $\omega, \alpha_i, \gamma$  and let X be a circle bundle over M with connection form  $\theta$  and first Chern class  $\gamma$  with  $d\theta = c_1$ . Then X is spin, has first Pontryagin class  $p_1 = 4\omega^2$  and has degree 2 and 3 cohomology  $H^2(X) = \langle \omega, \alpha_1, \alpha_2, \alpha_3 \rangle$ ,  $H^3(X) = H^3(M) \oplus \langle [\theta \otimes \omega] \rangle$ . The term  $[\theta \otimes \omega]$  is the one we are concerned about: As  $\gamma \omega = 0$ , in the form level we have  $c_1 \wedge w = d\xi$ , for some 3-form  $\xi$  pulled back from M, hence  $\theta \wedge w - \xi$  is a closed form. This form represents the cohomology class  $[\theta \otimes \omega]$ .

Letting  $\varphi = -\theta \wedge w + \xi$  we see that  $\varphi$  induces a negative definite bilinear form on  $H^2(X)$  as

$$\int_X \varphi \wedge a_i \wedge a_j = \int_M -w \wedge a_i \wedge a_j = -2\delta_{ij},$$

and similarly

$$\int_X \varphi \wedge w \wedge a_j = 0 \quad \text{and} \quad \int_X \varphi \wedge w \wedge w = -2.$$

Also,

$$\int_{X} \varphi p_{1} = \int_{X} -\theta w 4w^{2} = -\int_{M} 4w^{3} = -8.$$

Finally, since  $\gamma$  pulls back to zero in  $H^2(X)$ , we can define the Massey products  $\langle \omega, \alpha_1, \alpha_2 \rangle$ ,  $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ ,  $\langle \alpha_2, \alpha_3, \omega \rangle$  and  $\langle \alpha_3, \omega, \alpha_1 \rangle$ . To prove that X is not formal we compute  $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ :

$$a_1 a_2 = da_{12}, \qquad a_2 a_3 = d(\theta a_1 - a_{23}),$$

where  $a_{ij}$  are pull-backs from the base. So

$$\int_X \langle \alpha_1, \alpha_2, \alpha_3 \rangle \omega = \int_X a_{12} a_3 w - a_1 (\theta a_1 - a_{23}) w = -\int_X \theta w a_1^2 = -\int_M w a_1^2 = -2.$$

Which means that, for these choices, the Massey product pairs nontrivially with the cohomology class  $\omega$  and therefore is a closed nonexact form. One can also check that different choices *keep* the integral above unchanged so the Massey product does not vanish.

Remark. If we deal with the Spin(7) case in a similar fashion, i.e., stripping off the Riemannian structure and working only with the implied topological properties, circle bundles will not provide possible examples of nonformal Spin(7)-manifolds. This is because Spin(7)-manifolds have  $\hat{A}$ -genus 1 and by a result of Atiyah and Hirzebruch [1] if a compact connected Lie group acts differentiably and non-trivially on a compact orientable spin manifold X, then  $\hat{A}(X) = 0$ .

### References

- [1] M. Atiyah and F. Hirzebruch. Spin-manifolds and group actions. In Essays on Topology and Related Topics (Mémoires dédiés à Georges de Rham), pages 18–28. Springer, New York, 1970.
- [2] I. K. Babenko and T. I. A. Massey products in symplectic manifolds. Sb. Math, 191:1107–1146, 2000.
- [3] G. R. Cavalcanti. The Lefschetz property, formality and blowing up in symplectic geometry. math.SG/0403067.
- [4] P. Deligne, P. Griffiths, J. Morgan, and D. Sullivan. *Real homotopy theory of Kähler manifolds*. Invent. Math., 29:245–274, 1975.
- [5] S. Donaldson. Symplectic submanifolds and almost-complex geometry. J. Differential Geometry, 44:666-705, 1996.
- [6] A. N. Dranishnikov and Y. Rudyak. Examples of non-formal closed (k-1)-connected manifolds of dimensions  $\geq 4k-1$ . math.AT/0306299, to appear in Proc. Amer. Math. Soc.
- [7] Y. Félix, S. Halperin, and J.-C. Thomas. *Rational homotopy theory*, volume 205 of *Graduate Texts in Mathematics*. Springer Verlag, 2001.
- [8] M. Fernández and V. Muñoz. Formality of Donaldson submanifolds. math.SG/0211017, to appear in Math. Zeit.
- [9] M. Fernández and V. Muñoz. On non-formal simply connected manifolds. Topology Appl., 135(1-3):111-117, 2004
- [10] P. Griffiths and J. Morgan. Rational Homotopy Theory and Differential Forms, volume 16 of Progress in Mathematics. Birkhäuser, 1981.

- [11] D. Joyce. Compact Manifolds with Special Holonomy. Oxford Mathematical Monographs. Oxford University Press, 2000.
- [12] A. Kovalev. Twisted connected sums and special Riemannian holonomy. J. Reine Angew. Math., 565:125–160, 2003.
- [13] T. J. Miller. On the formality of k-1 connected compact manifolds of dimension less than or equal to 4k-2. Illinois J. Math., 23:253–258, 1979.
- [14] S. Mori and S. Mukai. Classification of Fano 3-folds with  $B_2 \geq 2$ . I. In Algebraic and topological theories (Kinosaki, 1984), pages 496–545. Kinokuniya, Tokyo, 1986.
- [15] D. Sullivan. Infinitesimal computations in topology. Inst. Hautes Études Sci. Publ. Math., 47:269–331, 1978.
- [16] C. T. C. Wall. Classification Problems in Differential Topology. V. On certain 6-manifolds. Invent. Math., 1:355–374, 1966.

MATHEMATICAL INSTITUTE, St. GILES 24-29, OXFORD, OX1 3BN, UK *E-mail address*: gilrc@maths.ox.ac.uk