

Unit Vectors, Morita Equivalence and Endomorphisms

Michael Skeide*

Dipartimento S.E.G.e S.

Università degli Studi del Molise

Via de Sanctis

86100 Campobasso, Italy

E-mail: skeide@math.tu-cottbus.de

Homepage: http://www.math.tu-cottbus.de/INSTITUT/lswas/_skeide.html

December 2004, Revised and extended in March 2005 and January 2006

Abstract

With every (strict or normal) unital endomorphism of the algebra of all adjointable operators on a Hilbert module there is associated a *correspondence* (that is, a Hilbert bimodule) such that the endomorphism can be recovered as amplification of the identity representation with that correspondence. In these notes we show the converse of this statement in the case of strongly full W^* -correspondences by establishing that every W^* -correspondence is Morita equivalent to one that has a unit vector. This, actually, means that every discrete product system of strongly full W^* -correspondences comes from a discrete E_0 -semigroup. (We also show the C^* -analogue of this result in the special case of full C^* -correspondences over a unital C^* -algebras.) Taking into account the duality between a *von Neumann correspondence* (that is, a W^* -correspondence over a von Neumann algebra) and its *commutant*, we furnish a different proof of Hirshberg's recent result that C^* -correspondences (with faithful left action) admit a (faithful) *essential* (that is, nondegenerate) representation on a Hilbert space, and we add that for W^* -correspondences this representation can be chosen normal.

*This work is supported by research funds of the Department S.E.G.e S. of University of Molise.

Let E denote Hilbert module over a C^* -algebra \mathcal{B} and suppose that ϑ is a strict unital endomorphism of the algebra $\mathcal{B}^a(E)$ of all adjointable operators on E .^[1] The theory of strict representations of $\mathcal{B}^a(E)$ according to Muhly, Skeide and Solel [MSS04] associates with ϑ the correspondence $E_\vartheta = E^* \odot_\vartheta E$ ^[2] from \mathcal{B} to \mathcal{B} (or **over** \mathcal{B}). Here E^* is the **dual correspondence** from \mathcal{B} to $\mathcal{B}^a(E)$ of E equipped with module actions $bx^*a := (a^*xb^*)^*$ and inner product $\langle x^*, y^* \rangle := xy^*$, while $E = {}_\vartheta E$ is viewed as a correspondence from $\mathcal{B}^a(E)$ to \mathcal{B} with left action of $\mathcal{B}^a(E)$ **via** ϑ .

The key points in the representation theory are that $E \odot E^* = \mathcal{K}(E)$ as correspondences from $\mathcal{B}^a(E)$ to $\mathcal{B}^a(E)$ ^[3] (via the isomorphism $x \odot y \mapsto xy^*$) and that ${}_\vartheta E$ can also be viewed as a correspondence from $\mathcal{K}(E)$ to \mathcal{B} . (The critical issue, nondegeneracy of the left action of $\mathcal{K}(E)$, follows because ϑ is assumed strict so that a bounded approximate unit of $\mathcal{K}(E)$ formed by finite-rank operators^[4] converges strictly to id_E .) Therefore,

$$E \odot E_\vartheta = E \odot E^* \odot_\vartheta E = \mathcal{K}(E) \odot_\vartheta E = {}_\vartheta E$$

as correspondence from $\mathcal{B}^a(E)$ to \mathcal{B} , that is, $a \odot \text{id}_{E_\vartheta} = \vartheta(a)$ ($a \in \mathcal{B}^a(E)$). An explicit isomorphism is given by

$$x \odot (y^* \odot z) \mapsto \vartheta(xy^*)z. \quad (*)$$

We observe that the **range ideal** $\mathcal{B}_E := \overline{\text{span}}\langle E, E \rangle$ of E and of E_ϑ necessarily coincide. Also, the left action of \mathcal{B}_E on E_ϑ is already nondegenerate. It is not difficult to show that E_ϑ is the unique (up to unique isomorphism) correspondence over \mathcal{B}_E that generates ϑ via an isomorphism $E = E \odot E_\vartheta$ as amplification $\vartheta(a) = a \odot \text{id}_{E_\vartheta}$ and that every other correspondence over \mathcal{B} that does so must contain E_ϑ (in a canonical way); see [MSS04]. In particular, if E is

^[1]The strict topology on $\mathcal{B}^a(E)$ is that inherited from the identification of $\mathcal{B}^a(E) = M(\mathcal{K}(E))$ as the multiplier algebra of the algebra $\mathcal{K}(E)$ of **compact operators** on E , that is, the the closed linear span of the **rank-one operators** $xy^* : z \mapsto x\langle y, z \rangle$ ($x, y \in E$). A bounded linear mapping on $\mathcal{B}^a(E)$ is **strict**, if it is strictly continuous on bounded subsets. That is equivalent to say it is $*$ -strongly continuous on bounded subsets.

^[2]The **tensor product** $E \odot F$ of a correspondence E from \mathcal{A} to \mathcal{B} and a correspondence F from \mathcal{B} to \mathcal{C} is the unique correspondence from \mathcal{A} to \mathcal{C} that is generated by elementary tensors $x \odot y$ with inner product $\langle x \odot y, x' \odot y' \rangle = \langle y, \langle x, x' \rangle y' \rangle$. A **correspondence** from \mathcal{A} to \mathcal{B} is just a Hilbert \mathcal{A} - \mathcal{B} -module, that is, a Hilbert \mathcal{B} -module with a **nondegenerate**(!) representation of \mathcal{A} by adjointable operators on E . The term **correspondence** is now standard for Hilbert bimodules (motivated by the idea of a **correspondence** from a set A to a set B as a generalized mapping). It has notational advantages, but it certainly obstructs a clear capture of other useful meanings of the word ‘correspondence’. Not necessarily standard is the requirement of nondegeneracy for the left action. But the reader will note that everywhere in these note it is indispensable that \mathcal{A} acts a neutral element under tensor product from the left.

^[3]Every closed ideal I in a C^* -algebra \mathcal{B} (in particular, \mathcal{B} itself) can be viewed as a correspondence from \mathcal{B} to \mathcal{B} with the natural bimodule action of \mathcal{B} and with inner product $\langle i, i' \rangle := i^*i'$.

^[4]The algebra of **finite rank operators** is $\mathcal{F}(E) := \text{span}\{xy^* : x, y \in E\}$. (Rieffel [Rie74a] used the term **imprimitivity algebra** of E .)

full^[5], then E_ϑ is the unique correspondence over \mathcal{B} generating ϑ .

The whole construction can be modified easily to a normal unital endomorphism of the W^* -algebra $\mathcal{B}^a(E)$ when E is a W^* -module.^[6] In this case, the tensor product \odot is replaced by the tensor product $\bar{\odot}^s$ of W^* -correspondences^[7]. Then $E_\vartheta = E^* \bar{\odot}^s_\vartheta E$ is a W^* -correspondence over \mathcal{B} such that $E = E \bar{\odot}^s E_\vartheta$ via the isomorphism defined by the same Equation (*) and as a W^* -correspondence over $\overline{\mathcal{B}_E^s}$ it is determined uniquely by ϑ .

It is the goal of these notes to show the converse of the construction of a correspondence from a unital endomorphism in the W^* -case:

Theorem. *Let E be a strongly full W^* -correspondence over a W^* -algebra \mathcal{B} (or, more generally, a W^* -correspondence such that $\overline{\mathcal{B}_E^s}$ acts nondegenerately on E). Then there exists a W^* -module F over \mathcal{B} (with $\overline{\mathcal{B}_F^s} = \overline{\mathcal{B}_E^s}$) and a normal unital endomorphism ϑ of $\mathcal{B}^a(F)$ such that $E = F^* \bar{\odot}^s_\vartheta F$.*

We will refer to this theorem as the *main theorem*. In Section 6 we will show a variant for C^* -correspondence at least in the case when the correspondence is full over a unital C^* -algebra.

As we explain in Section 1, this result implies also that a *discrete* product system (that is, a product system indexed by \mathbb{N}_0) of W^* -correspondences can be derived from an E_0 -semigroup on some $\mathcal{B}^a(F)$. Indeed, we will prove the main theorem by an analysis of the whole E_0 -semigroup $(\vartheta^n)_{n \in \mathbb{N}_0}$ and its relation with product systems of correspondences (Skeide [Ske02]), dilations of CP-semigroups (Bhat and Skeide [BS00]) and Morita equivalence of correspondences (Muhly and Solel [MS00]). In particular, as an intermediate step we will show that a W^* -correspondence E stems from an endomorphism, if and only if the product system $E^{\bar{\odot}^s} = (E^{\bar{\odot}^s n})_{n \in \mathbb{N}_0}$ it generates is Morita equivalent to another one $F^{\bar{\odot}^s} = (F^{\bar{\odot}^s n})_{n \in \mathbb{N}_0}$ that has a unital unit (Theorem 4.10). This theorem holds also, when the product system and the E_0 -semigroup are indexed by \mathbb{R}_+ . In the discrete case existence of a unital unit just means that $F = F_1$ contains a unit vector.

It is a typical feature in the construction of dilations of a single mapping that a dilation of that mapping is, actually, a dilation of the whole semigroup generated by that mapping. In these

^[5]A Hilbert \mathcal{B} -module is *full*, if $\mathcal{B}_E = \mathcal{B}$.

^[6]A W^* -*module* is a Hilbert module E over a W^* -algebra \mathcal{B} that is *self-dual*, i.e., every bounded right-linear mapping $\Phi: E \rightarrow \mathcal{B}$ has the form $\Phi(x) = \langle y, x \rangle$ for a (unique) $y \in E$.

^[7]A W^* -*correspondence* from a W^* -algebra \mathcal{A} to a W^* -algebra \mathcal{B} is a correspondence from \mathcal{A} to \mathcal{B} that is also a W^* -module over \mathcal{B} such that the left action of \mathcal{A} is normal. The tensor product of W^* -correspondences E and F is the unique *minimal self-dual extension* of the correspondence $E \odot F$ according to Paschke [Pas73]. This self-dual extension can be obtained considerably more easily, namely as a strong closure in an operator space, if we pass to von Neumann modules; see Section 7. In this sense, by $\bar{}^s$ we will indicate *strong closure*, although the reader who wishes to do so may think of σ -weak closure with respect to a suitably chosen pre-dual. Also, we will always consider strong limits of nets, but they may conveniently be replaced by (possibly different!) nets that converge σ -weakly.

notes we dilate a correspondence to an endomorphism by, acutally, “dilating” the (discrete) product system generated by the correspondence to an E_0 -semigroup. Therefore, it is has some point to think of E_0 -semigroups as dilations of product systems. So Arveson’s statement in [Arv89] that every **Arveson system** (that is, a product systems of Hilbert spaces or, equivalently, of correspondences over \mathbb{C}) stems from an E_0 -semigroup on $\mathcal{B}(H)$ can be stated as “every Arveson system admits a dilation”.

In Section 5 we proof the main theorem by showing that every (strongly) full W^* -correspondence is Morita equivalent to one that has a unit vector. For the construction of that correspondence with a unit vector we need Lemmata 2.2 and 3.2 which assert that given a full Hilbert module over a unital C^* -algebra or a (strongly) full W^* -module, then a direct sum of a suitable number of copies has a unit vector.

Then, in Section 7, we recall the duality between a *von Neumann correspondence* E (that is, a W^* -correspondence over a von Neumann algebra) and its *commutant* E' as described in Skeide [Ske03a] and show (by a reinterpretation of a result from Muhly and Solel [MS99]) that the commutant E' admits a *nondegenerate normal* representation on a Hilbert space, if and only if E is the correspondence of an endomorphism (Theorem 7.4). Together with our main theorem this reproves a recent result by Hirshberg [Hir05a], who shows by completely different methods that every C^* -correspondence admits such a representation. Additionally, we show that in the case of W^* -correspondences this representation can be chosen normal (left open in [Hir05a]).

In Section 8 we indicate some implications of the main theorem and of Section 7 for the continuous case. In Section 9 we illustrate the abstract constructions explicitly in an example.

A note on footnotes. We apologize for the extensive use of footnotes in the beginning of this introduction. They contain definitions the expert surely will now (maybe, except the convention on nondegenerate left actions) and that would disturb the presentation of the argument too much. However, as we wish to be comprehensive also for the nonexpert in Hilbert modules, we decided to add the basic definitions in form of footnotes.

Acknowledgements. We wish to express our gratitude to P. Muhly for warm hospitality during several stays in Iowa City with many inspiring discussions that influenced this work. We thank I. Hirshberg for intriguing discussions around his article [Hir05a]. We are particularly grateful to B. Solel for pointing out to us two inaccuracies of arguments in an earlier version. Last but not least, we acknowledge also the support by Research Fonds of the Department S.E.G.e S. of University of Molise.

1 Prerequisites on E_0 -semigroups and product systems

In this section we recall the construction from Skeide [Ske02] of a product system from an E_0 -semigroup on $\mathcal{B}^a(E)$ based on existence of a unit vector in E and put it into perspective with the new construction from Muhly, Skeide and Solel [MSS04] as sketched in the introduction. Then we discuss the relation with units, CP-semigroups and the construction of dilations of CP-semigroups as described in Bhat and Skeide [BS00].

Let E be a Hilbert module over a C^* -algebra \mathcal{B} and let $\vartheta = (\vartheta_t)_{t \in \mathbb{S}}$ be a strict E_0 -*semigroup* on $\mathcal{B}^a(E)$, that is, a semigroup of unital endomorphisms ϑ_t of $\mathcal{B}^a(E)$ that are strict. \mathbb{S} is either the set of nonnegative integers $\mathbb{N}_0 = \{0, 1, \dots\}$ or the set of nonnegative reals $\mathbb{R}_+ = [0, \infty)$. (We are mainly interested in the discrete case $\mathbb{S} = \mathbb{N}_0$ but there is no reason to restrict the discussion to that case. The only exception regards existence of *units*. While here we are exploiting the fact that in the discrete case every vector in E_1 generates a unit, in the continuous case units need not exist.)

For every $t > 0$ denote by $E_t := E^* \odot_{\vartheta_t} E$ the correspondence over \mathcal{B} generating the endomorphism ϑ_t via the isomorphism $E = E \odot E_t$ described by (*). We extend the definition to $t = 0$ by putting $E_0 = \mathcal{B}$. (If E is full, then this is automatic.) The E_t form a product system $E^\odot = (E_t)_{t \in \mathbb{S}}$, that is

$$E_s \odot E_t = E_{s+t} \quad (E_r \odot E_s) \odot E_t = E_r \odot (E_s \odot E_t),$$

via

$$\begin{aligned} E_s \odot E_t &= (E^* \odot_{\vartheta_s} E) \odot (E^* \odot_{\vartheta_t} E) \\ &= E^* \odot_{\vartheta_s} (E \odot (E^* \odot_{\vartheta_t} E)) = E^* \odot_{\vartheta_s} (\vartheta_t E) = E^* \odot_{\vartheta_{s+t}} E = E_{s+t}. \end{aligned}$$

In order to indicate that an elementary tensor $x^* \odot y$ is to be understood in $E^* \odot_{\vartheta_t} E$ we shall write $x^* \odot_t y$. We leave it as an exercise to check (using the notation \odot_t appropriately) on elementary tensors that the suggested identification is, indeed, associative.

Suppose now that \mathcal{B} is unital and that E has a **unit vector** ξ (that is, $\langle \xi, \xi \rangle = \mathbf{1}$ so that, in particular, E is full). Then we may identify E_t with the submodule $\vartheta_t(\xi\xi^*)E$ of E when equipped with the left action $bx := \vartheta_t(\xi b \xi^*)x$. The concrete identification $E^* \odot_{\vartheta_t} E \rightarrow \vartheta_t(\xi\xi^*)E$ is

$$x^* \odot_t y \mapsto \vartheta_t(\xi x^*)y \quad \text{with inverse} \quad \vartheta_t(\xi\xi^*)z \mapsto \xi^* \odot_t z,$$

(To see that the inverse is really surjective, write $x^* \odot_t y \in E^* \odot_{\vartheta_t} E$ as $x^* \odot_t y = (\xi^* \xi x^*) \odot_t y = \xi^* \odot_t \vartheta_t(\xi x^*)y$.) The resulting identification $\vartheta_s(\xi\xi^*)E \odot \vartheta_t(\xi\xi^*)E = \vartheta_{s+t}(\xi\xi^*)E$ is

$$\vartheta_s(\xi\xi^*)x \odot \vartheta_t(\xi\xi^*)y \mapsto (\xi^* \odot_s x) \odot (\xi^* \odot_t y) \mapsto \xi^* \odot_{s+t} (\vartheta_t(x\xi^*)y) \mapsto \vartheta_{s+t}(\xi\xi^*)\vartheta_t(x\xi^*)y.$$

This is the way how we determined in Skeide [Ske02] the product system of an E_0 -semigroup (when E has a unit vector) generalizing Bhat's construction in [Bha96] of the Arveson system of an E_0 -semigroup on $\mathcal{B}(H)$ (where existence of a unit vector is not a problem as every nonzero Hilbert space has unit vectors).

By [Ske02] the family $T_t(b) = \langle \xi, \vartheta_t(\xi b \xi^*) \xi \rangle$ of maps on \mathcal{B} defines a semigroup, if and only if the family of projections $p_t = \vartheta_t(\xi \xi^*)$ is increasing. In this case, $\xi_t := \vartheta_t(\xi \xi^*) \xi = \xi$ is contained in all E_t and, moreover, $\xi_s \odot \xi_t = \xi_{s+t}$ and $\xi_0 = \mathbf{1}$, that is, the family $\xi^\odot = (E_t)_{t \in \mathbb{S}}$ defines a **unit** for E^\odot . This unit is **unital**, that is, all ξ_t are unit vectors.

Under the preceding conditions the construction is reversible (see [BS00, Ske02] for details) and this reversion is what we need for the constructive part in the proof of the main theorem. More specifically, let E^\odot be a product system of correspondences over \mathcal{B} and let ξ^\odot be a unital unit for E^\odot . By $\xi_s \odot \text{id}_{E_t}: x_t \mapsto \xi_s \odot x_t$ we define isometric (right linear but, usually, not bilinear) embeddings $E_t \rightarrow E_{s+t}$. These embeddings form an inductive system so that we may construct the (completed) inductive limit $E_\infty = \overline{\lim \text{ind}_{t \rightarrow \infty} E_t}$. The factorization property of the product system E^\odot turns over to E_∞ , that is

$$E_\infty \odot E_t = E_{\infty+t} = E_\infty \quad (E_\infty \odot E_s) \odot E_t = E_\infty \odot (E_s \odot E_t), \quad (1.1)$$

so that $\vartheta_t(a) := a \odot \text{id}_{E_t} \in \mathcal{B}^a(E_\infty \odot E_t) = \mathcal{B}^a(E_\infty)$ defines an E_0 -semigroup on $\mathcal{B}^a(E_\infty)$. (We need to know how to make the intuitive identification $E_\infty = E_\infty \odot E_t$ precise only for the concrete example in Section 9. Here we content ourselves to know that it works, and refer the reader to [BBL04, Section 4.4].) If product system and unit are those constructed in the preceding paragraph give back E^\odot and ξ^\odot from an E_0 -semigroup on $\mathcal{B}^a(E)$ and a unit vector, then we recover E as $E = E_\infty$, if and only if the projections p_t (of the last paragraph) converge to id_E strongly.

1.1 Observation. We summarize: If $E = E_1$ is a correspondence over \mathcal{B} with a unit vector $\xi = \xi_1$, then $E^\odot = (E_n)_{n \in \mathbb{N}_0}$ with $E_n := E^{\odot n}$ is a (discrete) product system and $\xi^\odot = (\xi_n)_{n \in \mathbb{N}_0}$ with $\xi_n := \xi^{\odot n}$ a unital unit. The inductive limit E_∞ over that unit carries a strict E_0 -semigroup $\vartheta = (\vartheta_n)_{n \in \mathbb{N}_0}$ with $\vartheta_n(a) = a \odot \text{id}_{E_n}$ whose product system is E^\odot . In particular, $E = E_1$ occurs as the correspondence of the unital strict endomorphism ϑ_1 of $\mathcal{B}^a(E_\infty)$.

All results have analogues for W^* -modules replacing strict mappings with normal (or σ -weak) mappings and the tensor product of C^* -correspondences with that of W^* -correspondences.

2 Unit vectors in Hilbert modules

In this section we discuss when full Hilbert modules over unital C^* -algebras have unit vectors. In particular, we show that even if there is no unit vector, then a finite direct sum will

admit a unit vector. What we actually need for the proof our main theorem is the version for W^* -modules which we discuss in the following section. But the result in this section is simple, allows to illustrate the problems free from technical ballast, motivates the following section and, finally, has some consequences in proving statements about finitely generated modules that are independent from the rest of the paper. It will play its role in the weaker weaker version of the main theorem, Theorem 6.6, for full correspondences over unital C^* -algebras.

Of course, a Hilbert module E over a unital C^* -algebra \mathcal{B} that is not full cannot have unit vectors. But also if E is full this does not necessarily imply existence of unit vectors.

2.1 Example. Let $\mathcal{B} = \mathbb{C} \oplus M_2 = \begin{pmatrix} \mathbb{C} & 0 \\ 0 & M_2 \end{pmatrix} \subset M_3 = \mathcal{B}(\mathbb{C}^3)$. The M_2 - \mathbb{C} -module $\mathbb{C}^2 = M_{21}$ may be viewed as a correspondence over \mathcal{B} (with operations inherited from $M_3 \supset \begin{pmatrix} 0 & 0 \\ \mathbb{C}^2 & 0 \end{pmatrix}$). Also its dual, the \mathbb{C} - M_2 -module $\mathbb{C}^{2*} = M_{12} =: \mathbb{C}_2$, may be viewed as a correspondence over \mathcal{B} . It is easy to check that $M = \mathbb{C}^2 \oplus \mathbb{C}_2 = \begin{pmatrix} 0 & \mathbb{C}_2 \\ \mathbb{C}^2 & 0 \end{pmatrix}$ is a Morita equivalence (see Section 4) from \mathcal{B} to \mathcal{B} (in particular, M is full) without a unit vector.

Observe that all modules and correspondences in Example 2.1 are W^* -modules, so missing unit vectors are not caused by insufficient closure. The reason why M does not contain a unit vector is because the full Hilbert M_2 -module \mathbb{C}_2 has “not enough space” to allow for sufficiently many orthogonal vectors. (Not two nonzero vectors of this module are orthogonal.) Another way to argue is to observe that every nonzero inner product $\langle x, y \rangle$ is a rank-one operator in $M_2 = \mathcal{B}(\mathbb{C}^2)$ while the identity has rank two. As soon as we create “enough space”, for instance, by taking the direct sum of sufficiently many (in our case two) copies of \mathbb{C}_2 the problem disappears.

In this section we show a lemma asserting that for every full Hilbert module a finite number of copies will be “enough space”. The basic idea is that, if $\langle x, y \rangle = \mathbf{1}$, then by Cauchy-Schwartz inequality $\mathbf{1} = \langle x, y \rangle \langle y, x \rangle \leq \langle x, x \rangle \|y\|^2$ so that $\langle x, x \rangle$ is invertible and $x \sqrt{\langle x, x \rangle}^{-1}$ is a unit vector. Technically, the condition $\langle x, y \rangle = \mathbf{1}$ is realized only approximately and by elements in E^n rather than in E .

2.2 Lemma. *Let E be a full Hilbert module over a unital C^* -algebra. Then there exists $n \in \mathbb{N}$ such that E^n has a unit vector.*

PROOF. E is full, so there exist $x_i^n, y_i^n \in E$ ($n \in \mathbb{N}; i = 1, \dots, n$) such that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \langle x_i^n, y_i^n \rangle = \mathbf{1}.$$

The subset of invertible elements in \mathcal{B} is open. Therefore, for n sufficiently big $\sum_{i=1}^n \langle x_i^n, y_i^n \rangle$ is invertible. Defining the elements $X_n = (x_1^n, \dots, x_n^n)$ and $Y_n = (y_1^n, \dots, y_n^n)$ in E^n we have, thus,

that

$$\langle X_n, Y_n \rangle = \sum_{i=1}^n \langle x_i^n, y_i^n \rangle$$

is invertible. So, also $\langle X_n, Y_n \rangle \langle Y_n, X_n \rangle$ is invertible and, therefore, bounded below by a strictly positive constant. Of course, $\|Y_n\| \neq 0$. By Cauchy-Schwartz inequality also

$$\langle X_n, X_n \rangle \geq \frac{\langle X_n, Y_n \rangle \langle Y_n, X_n \rangle}{\|Y_n\|^2}$$

is bounded below by a strictly positive constant and, therefore, $\langle X_n, X_n \rangle$ is invertible. It follows that $X_n \sqrt{\langle X_n, X_n \rangle^{-1}}$ is a unit vector in E^n . ■

2.3 Corollary. *If E (as before) contains an arbitrary number of mutually orthogonal copies of a full Hilbert submodule (for instance, if E is isomorphic to E^n for some $n \geq 2$), then E has a unit vector.*

Excluding the one-dimensional case, Arveson systems (continuous time) have always infinite-dimensional members. We believe that something similar should be true also for general (continuous time) product systems, in the sense that the product systems E° whose members E_t ($t \in \mathbb{R}_+$) do not fulfill not assumption of the corollary should belong to an unrealistic subclass that can be ruled out and understood more easily. (In Arveson's setting they come from automorphism groups. But we know that one-dimensional product systems can have nontrivial structures. But as their members are \mathcal{B} as right module, they all have unit vectors. The class we wish to rule out cannot have \mathcal{B} as a direct summand.) Example 9.5 tells us that there are serious examples in the discrete case where not one member E_n ($n > 0$) of the product system has a unit vector.

Lemma 2.2 implies that, if $\mathcal{K}(E)$ is unital, then $\mathcal{K}(E) = \mathcal{F}(E)$. (Just apply the lemma to the full Hilbert $\mathcal{K}(E)$ -module E^* . Of course, this may be obtained directly, using again that the invertible elements form an open subset in the C^* -algebra $\mathcal{K}(E)$ and the fact that $\mathcal{F}(E)$ is dense in $\mathcal{K}(E)$.) This has an implication for the theory of finitely generated Hilbert modules.

2.4 Corollary. *If $\mathcal{K}(E)$ is unital, then E is algebraically finitely generated.*

This is some sort of inverse to the well-known fact that an (algebraically) finitely generated Hilbert \mathcal{B} -module is isomorphic to a (complemented) submodule of \mathcal{B}^n for some n . Indeed, at least if \mathcal{B} is unital, then E being a complemented submodule of \mathcal{B}^n implies that the identity $\mathbf{1}_{\mathcal{B}^n}$ in $\mathcal{B}^a(\mathcal{B}^n) = M_n(\mathcal{B}) = \mathcal{K}(\mathcal{B}^n)$ gives rise to a an identity $p_E \mathbf{1}_{\mathcal{B}^n} p_E \in \mathcal{F}(E)$ for E , where p_E denotes the projection onto E .

3 Unit vectors in W^* -modules

In this section we proof the analogue of Lemma 2.2 for W^* -modules. Of course, a W^* -module is a Hilbert module. If it is full then Lemma 2.2 applies. But the good notion of fullness for a W^* -module is that it is **strongly full**, that is, the inner product of the W^* -module generates \mathcal{B} as a W^* -algebra. (Strong fullness is the more useful notion for W^* -modules, because it can always be achieved by restricting \mathcal{B} to the W^* -subalgebra generated by the inner product. Example 3.1 tells us that the same is not true for fullness in the case of W^* -modules.) Usually, for a W^* -module we will just say “full”, while we mention explicitly, when we intend “full in the Hilbert module sense”. It is the assumption of strong fullness for which we want to prove the main theorem, and not the stronger assumption of fullness (that might be not achievable). We thank B. Solel for pointing out to us this gap in the first version of these notes.

We see immediately that for W^* -modules the cardinality of the direct sum in Lemma 2.2 can no longer be kept finite.

3.1 Example. Let H be an infinite-dimensional Hilbert space. Then H^* is a W^* -module over $\mathcal{B}(H)$, that is strongly full but not full as a Hilbert $\mathcal{B}(H)$ -module. (Indeed, the range ideal of H^* in $\mathcal{B}(H)$ is $\mathcal{B}(H)_{H^*} = \mathcal{K}(H) \neq \mathcal{B}(H)$.) For every finite direct sum H^{*n} the inner product $\langle X_n, X_n \rangle$ ($X_n \in H^{*n}$) has rank not higher than n . Therefore, H^{*n} does not admit a unit vector. Only if we consider $\overline{H^{*n}}^s$, the self-dual extension of H^{*n} , where $n = \dim H$, then the vector in $\overline{H^{*n}}^s$ with the components e_i^* ((e_i) some orthonormal basis of H) is a unit vector. But this vector is not in H^{*n} if n is infinite.

The example is in some sense typical. In fact, we constructed a multiple of H^* that contains a unit vector by choosing an orthonormal basis for its dual H . This will also be our strategy for general W^* -modules. A suitable substitute for orthonormal bases are *quasi* orthonormal bases. A **quasi orthonormal basis** in a W^* -module E over \mathcal{B} is a family $(e_i, p_i)_{i \in S}$ where S is some index set (of cardinality n , say), p_i are projections in \mathcal{B} and e_i a elements in E such that

$$\langle e_i, e_j \rangle = \delta_{i,j} p_j \quad \text{and} \quad \sum_{i \in S} e_i e_i^* = \text{id}_E$$

(strong limit in the W^* -algebra $\mathcal{B}^a(E)$ over the finite subsets of S in the case S is not finite; see Footnote [7]). Existence of a quasi orthonormal basis follows by self-duality (Paschke [Pas73]).

3.2 Lemma. *Let E be a (strongly) full W^* -module. Then there exists a cardinal number n such that $\overline{E^n}^s$ has a unit vector.*

PROOF. Let us choose a quasi orthonormal basis $(e_i^*, e_i e_i^*)_{i \in S}$ for the dual $\mathcal{B}^a(E)$ -module E^* . Then

$$\sum_{i \in S} e_i^* e_i = \sum_{i \in S} \langle e_i, e_i \rangle = \text{id}_E.$$

The second sum is, actually, over the elements $\langle e_i, e_i \rangle$ when considered as operator acting from the left on E^* . But, as E is strongly full, the action of \mathcal{B} on E^* is faithful. In particular, the only element in \mathcal{B} having the action id_E is, really, $\mathbf{1} \in \mathcal{B}$. Now, if we put $n = \#S$, then the vector in E^n with components e_i is a unit vector. ■

We note that this idea of choosing a quasi orthonormal basis for E^* (with E full) can be used in order to generalize Bhat's approach to product systems to the case of W^* -modules without a unit vector; see [Ske05b].

3.3 Remark. Of course, if E is countably generated as W^* -module, then in Lemma 3.2 we may choose $n = \#\mathbb{N}$. For this it is not necessary that \mathcal{B} is countably generated. Thus, neither \mathcal{B} nor E need have a separable pre-dual.

4 Morita equivalence for product systems

In this section we review the notions of (strong) Morita equivalence (Rieffel [Rie74b]), Morita equivalence for Hilbert modules (new in these notes) and Morita equivalence for correspondences (Muhly and Solel [MS00]). We put some emphasis on the difference between the C^* -case and the W^* -case that is in part responsible for the fact that we cannot prove our main theorem in the general C^* -case. Then we show that a product system of W^* -correspondences can be derived from an E_0 -semigroup, if and only if it is Morita equivalent to a product system that has a unital unit. In the discrete case this means a W^* -correspondence stems from a unital endomorphism of some $\mathcal{B}^a(E)$, if and only if it is Morita equivalent to a W^* -correspondence that has a unit vector.

For a detailed introduction to *Morita equivalence* (Morita [Mor58]) and its motivation (and its relation with the constructions in Section 7) we refer the reader to Rieffel [Rie74b] (and to Skeide [Ske05a]).

A correspondence M from \mathcal{A} to \mathcal{B} is called a **Morita equivalence** from \mathcal{A} to \mathcal{B} , if it is full and if the canonical mapping from \mathcal{A} into $\mathcal{B}^a(M)$ corestricts to an isomorphism $\mathcal{A} \rightarrow \mathcal{K}(M)$. Clearly, the two conditions can be written also as

$$M^* \odot M = \mathcal{B} \qquad M \odot M^* = \mathcal{A}$$

in each case under the suggested canonical mappings $\mathcal{A} \rightarrow \mathcal{K}(M) = M \odot M^*$ and $M^* \odot M = \mathcal{B}_M \rightarrow \mathcal{B}$. From these equations one concludes easily a couple of facts. Firstly, if M is a Morita equivalence from \mathcal{A} to \mathcal{B} , then M^* is a Morita equivalence from \mathcal{B} to \mathcal{A} . Secondly, the tensor product of Morita equivalences, if it exists, is a Morita equivalence. Thirdly, M and M^* are inverses under tensor product. Two C^* -algebras are called **strongly Morita equivalent**, if they admit a Morita equivalence from one to the other. Strong Morita equivalence is finer than

Morita equivalence, but considerably coarser than isomorphism. Usually, we say just *Morita equivalent* also when we intend *strongly Morita equivalent*.

4.1 Example. All M_n are (strongly) Morita equivalent to \mathbb{C} via the Morita equivalence \mathbb{C}^n . In fact, a representation of M_n factors the representation space H into $\mathbb{C}^n \otimes \mathfrak{H}$ and for every n the dimension of the *multiplicity space* \mathfrak{H} is the parameter that classifies the representations.

In the category of W^* -algebras with W^* -correspondences a correspondence from \mathcal{A} to \mathcal{B} is a *Morita W^* -equivalence*, if M is strongly full and if the canonical mapping $\mathcal{A} \rightarrow \mathcal{B}^a(M)$ is an isomorphism.

4.2 Remark. Clearly, in the W^* -case we have $M \overset{\circ}{\circ} M^* = \mathcal{B}^a(M)$. The fact that Morita equivalence for W^* -algebra relates \mathcal{A} to $\mathcal{B}^a(M)$ while strong Morita equivalence of C^* -algebras relates \mathcal{A} only to $\mathcal{K}(M)$ is one of the reasons why our proof of the main theorem works only in the W^* -case, respectively, runs considerably less smooth the particular C^* -case discussed in Section 6.

4.3 Example. The M_n are W^* -algebras, the \mathbb{C}^n and their duals are W^* -correspondences and all tensor products are tensor products in the W^* -sense.

4.4 Remark. Versions of Examples 4.1 and 4.3 for infinite-dimensional matrices and \mathbb{C} replaced with \mathcal{B} are crucial for the proof of the main theorem. Essentially, we are going to use \mathcal{B}^n as Morita equivalence from $M_n(\mathcal{B})$ to \mathcal{B} . Of course, either we have to pass to strong closures (Section 5) or to a weaker notion of Morita equivalence (Section 6).

4.5 Definition (Muhly and Solel [MS00]). A correspondence E over \mathcal{B} and a correspondence F over \mathcal{C} are (strongly) *Morita equivalent*, if there is a Morita equivalence M from \mathcal{B} to \mathcal{C} such that $E \overset{\circ}{\circ} M = M \overset{\circ}{\circ} F$ (or $E = M \overset{\circ}{\circ} F \overset{\circ}{\circ} M^*$).

We add here:

4.6 Definition. A Hilbert \mathcal{B} -module E and a Hilbert \mathcal{C} -module F are (strongly) *Morita equivalent*, if there is a Morita equivalence M from \mathcal{B} to \mathcal{C} such that $E \overset{\circ}{\circ} M = F$ (or $E = F \overset{\circ}{\circ} M^*$).

Of course, the definitions for the W^* -case are analogue.

Morita equivalence of Hilbert modules and Morita equivalence of correspondences are related by the following crucial proposition. Suppose $\alpha: \mathcal{B}^a(E) \rightarrow \mathcal{B}^a(F)$ is a (bi-)strict isomorphism. By [MSS04] this is the case, if and only if E and F are Morita equivalent where the Morita equivalence M induces α as $\alpha(a) = a \overset{\circ}{\circ} \text{id}_M$.

Now suppose there are two strict unital endomorphisms ϑ and θ on $\mathcal{B}^a(E)$ and $\mathcal{B}^a(F)$, respectively. We may ask whether they are *conjugate*, that is, whether there exists a (bi-)strict isomorphism $\alpha: \mathcal{B}^a(E) \rightarrow \mathcal{B}^a(F)$ such that $\theta = \alpha \circ \vartheta \circ \alpha^{-1}$.

4.7 Proposition. ϑ and θ are conjugate, if and only if there is a Morita equivalence inducing an isomorphism $F = E \odot M$ such that $E_\vartheta \odot M = M \odot F_\theta$, that is, if and only if E and F as well as E_ϑ and F_θ are Morita equivalent by the same Morita equivalence.

The proof consists very much of computations like the second half of the proof Theorem 4.10 below. We leave it as an exercise.

4.8 Remark. Note that in the scalar case $\mathcal{B} = \mathcal{C} = \mathbb{C}$, where \mathbb{C} is the only Morita equivalence over \mathbb{C} , we recover the well-known facts that every normal isomorphism $\alpha: \mathcal{B}(G) \rightarrow \mathcal{B}(H)$ is induced by a unitary $G \rightarrow H$ and that the multiplicity spaces of two endomorphisms conjugate by α must be equal.

Clearly, if $E^\odot = (E_t)_{t \in \mathbb{S}}$ is a product system of correspondences over \mathcal{B} and M is a Morita equivalence from \mathcal{B} to \mathcal{C} , then $M^* \odot E^\odot \odot M := F^\odot = (F_t)_{t \in \mathbb{S}}$ with $F_t := M^* \odot E_t \odot M$ and isomorphisms

$$F_s \odot F_t = M^* \odot E_s \odot M \odot M^* \odot E_t \odot M = M^* \odot E_s \odot E_t \odot M = M^* \odot E_{s+t} \odot M = F_{s+t}$$

is a product system of correspondences over \mathcal{C} .

4.9 Definition. We say E^\odot and F^\odot are (strongly) **Morita equivalent**, if there exists a Morita equivalence M and an **isomorphism** $u^\odot = (u_t)_{t \in \mathbb{S}}: M^* \odot E^\odot \odot M \rightarrow F^\odot$ (that is, the u_t are bilinear unitaries in $M^* \odot E_t \odot M \rightarrow F_t$ such that $u_s \odot u_t = u_{s+t}$ and $u_0 = \text{id}_C$).

The version for W^* -correspondences is analogue.

4.10 Theorem. Let $E^{\bar{\odot}^s} = (E_t)_{t \in \mathbb{S}}$ be a product system of (strongly) full W^* -correspondences E_t over a W^* -algebra \mathcal{B} . Then $E^{\bar{\odot}^s}$ is the product system of a normal E_0 -semigroup $\vartheta = (\vartheta_t)_{t \in \mathbb{S}}$ on $\mathcal{B}^a(E)$ for some W^* -module E over \mathcal{B} , if and only if $E^{\bar{\odot}^s}$ is Morita equivalent to a product system $F^{\bar{\odot}^s}$ of W^* -correspondences over a W^* -algebra \mathcal{C} that contains a unital unit ζ^\odot .

PROOF. “ \implies ”. Suppose $E^{\bar{\odot}^s}$ is the product system of the normal E_0 -semigroup ϑ on the W^* -algebra $\mathcal{C} := \mathcal{B}^a(E)$. Put $F_t := E \bar{\odot}^s E_t \bar{\odot}^s E^*$. As $E \bar{\odot}^s E_t = E$ and $\vartheta_t(a) = a \odot \text{id}_{E_t}$, we find $F_t = \vartheta_t \mathcal{B}^a(E)$ and $a_s \odot a_t = \vartheta_t(a_s) a_t$ is the isomorphism $F_s \bar{\odot}^s F_t = F_{s+t}$. Clearly, $\zeta_t = \text{id}_E \in \mathcal{B}^a(E) = F_t$ defines a unital unit ζ^\odot for $F^{\bar{\odot}^s}$. (One easily verifies that also the inductive limit $F_\infty = \mathcal{B}^a(E) = E \bar{\odot}^s E^*$ constructed from that unit is that obtained from E via the Morita equivalence $M := E^*$ as $F_\infty = E \bar{\odot}^s E^*$.)

“ \impliedby ”. Suppose M is a Morita W^* -equivalence from \mathcal{B} to \mathcal{C} such that $F^{\bar{\odot}^s} := M^* \bar{\odot}^s E^{\bar{\odot}^s} \bar{\odot}^s M$ has a unital unit ζ^\odot . Construct the inductive limit F_∞ with the normal E_0 -semigroup $\theta_t(a) = a \odot \text{id}_{F_t}$ ($a \in \mathcal{B}^a(F_\infty)$) and put $E := F_\infty \bar{\odot}^s M^*$. Then

$$\vartheta_t(a) := \theta_t(a \odot \text{id}_M) \odot \text{id}_{M^*} = a \odot \text{id}_M \odot \text{id}_{F_t} \odot \text{id}_{M^*} = a \odot \text{id}_{E_t}$$

($a \in \mathcal{B}^a(E)$) where

$$\begin{aligned} E &= E \bar{\otimes}^s M \bar{\otimes}^s M^* (= F_\infty \bar{\otimes}^s M^* = F_\infty \bar{\otimes}^s F_t \bar{\otimes}^s M^*) \\ &= E \bar{\otimes}^s M \bar{\otimes}^s F_t \bar{\otimes}^s M^* = E \bar{\otimes}^s E_t \end{aligned}$$

is the induced semigroup on $\mathcal{B}^a(E)$. As F_∞ is full (it contains a unit vector) also $E = F_\infty \bar{\otimes}^s M^*$ is full ($\overline{C_{F_\infty}}^s = C$ acts nondegenerately on M^* so that $\overline{\mathcal{B}_E}^s = \overline{\mathcal{B}_{M^*}}^s = \mathcal{B}$). Therefore, by uniqueness the product systems associated with ϑ gives us back E_t . ■

4.11 Remark. $F_t = {}_{\vartheta_t} \mathcal{B}^a(E)$ is exactly the one-dimensional product system of correspondences over $\mathcal{B}^a(E)$ constructed as in [BS00] from ϑ when considered as CP-semigroup; see Skeide [Ske01a, Example 11.1.3]. We have used the same trick in Skeide [Ske03a], switching between a W^* -correspondence $\mathcal{B}(H) \bar{\otimes}^s \mathfrak{H}$ over $\mathcal{B}(H)$ (note that such an exterior tensor product is the most general form of a W^* -correspondence over $\mathcal{B}(H)$) and the Hilbert space (that is, the correspondence over \mathbb{C}) \mathfrak{H} that corresponds to a member of an Arveson system, in order to solve a problem due to Powers. An extension to the module case will appear in Bhat, Liebscher and Skeide [BLS06]. In the recent preprint [MM04] Marrero and Muhly use the trick to determine the index of a CP-map on M_n .

What happens, if $E \bar{\otimes}^s$ is not full? Suppose it is derived from an E_0 -semigroup on $\mathcal{B}^a(E)$. Then by the extended uniqueness result, the E_t are actually correspondences over $\overline{\mathcal{B}_E}^s$. Therefore, if a product system is derived from a semigroup the range ideal for each E_t cannot depend on t and it must act nondegenerately on each E_t . In other words, either the range ideal is stationary and acts nondegenerately on each E_t so that we may simply pass from \mathcal{B} to that ideal and apply the theorem, or the product system is not derived from an E_0 -semigroup.

4.12 Corollary. *A correspondence not satisfying the condition in brackets in the main theorem cannot be derived from a unital endomorphism.*

We emphasize that this does not mean that for a product system with nonstationary range ideals it was not possible to find isomorphisms $E \odot E_t = E$ so that $\vartheta_t(a) := a \odot \text{id}_{E_t}$ defines a semigroup. (In fact it is easy to describe such examples by *exterior* direct sum constructions. The *exterior direct sum* of a \mathcal{B} -module and a C -module is a $\mathcal{B} \oplus C$ -module.) It just means that the product system associated with ϑ will be smaller.

4.13 Example. A nontrivial example of a product system with range ideals decreasing to $\{0\}$ we obtain as follows. Let $\mathcal{B} = \mathcal{C}_0(0, \infty)$ be the continuous functions on \mathbb{R}_+ that vanish at 0 and at ∞ . Let $s = (s_t)_{t \in \mathbb{R}_+}$ be the usual right shift $[s_t(f)](x) = f(x - t)$ putting $f(x) = 0$ when $x < 0$. Then $E_t := s_t(\mathcal{B})$ with left action $b \cdot s_t(b') = s_t(bb')$ is a product system via

$s_s(b) \odot s_t(b') = s_{s+t}(b)s_t(b') \in s_{s+t}(\mathcal{B})$. Note that this product system does not admit units nor (nonzero) elements that commute with the algebra.

Product systems like in this example are usually derived not from E_0 -semigroups but from E -semigroups, that is, semigroups of not necessarily unital endomorphisms. In fact, the product system is that associated with the shift semigroup s .

5 Proof of the main theorem

We put together the results to form a proof of the main theorem. But before we do that, we explain why even under the assumptions of Lemma 2.2 (resulting into finite direct sums) we would not be able to produce a proof based on strong Morita equivalence that works also in the C^* -case.

Let E be a full correspondence over a unital C^* -algebra \mathcal{B} . By Lemma 2.2 we know that for some $n \in \mathbb{N}$ the correspondence E^n has a unit vector. We observe that $E^n = \mathcal{B}^n \odot E$, where \mathcal{B}^n is a Morita equivalence from $M_n(\mathcal{B})$ to \mathcal{B} . If we could show existence of a unit vector in $M_n(E) = \mathcal{B}^n \odot E \odot \mathcal{B}_n$ where $\mathcal{B}_n := (\mathcal{B}^n)^*$ is the dual of \mathcal{B}^n , then E was Morita equivalent to a correspondence with a unit vector. In this case the “ \Leftarrow ” direction of the proof of Theorem 4.10 works even without strong closure. (The main reason for strong closure is that rarely $\mathcal{B}^a(E) = \mathcal{K}(E)$ so E is rarely a Morita equivalence from $\mathcal{B}^a(E)$ to \mathcal{B}_E .)

Unfortunately, $M_n(E)$ need not have a unit vector. Suppose $n \geq 2$ is minimal. To produce a $\mathbf{1}$ in a place in the diagonal we need n orthogonal vectors, and to produce $\mathbf{1}$ in every of the n places in the diagonal we need n^2 orthogonal vectors. However, the $M_n(\mathcal{B})$ -correspondence $M_n(E)$ still has “space” only for n orthogonal vectors (with suitable inner products). The problem remains, when we use $M_m(E)$ ($m > n$) instead. Still we need mn orthogonal vectors to produce m units in the diagonal, but we have “space” only for m orthogonal vectors. The problem disappears if $m = \infty$, because then we can “slice” $m = \infty$ into n slices still of size $m = \infty$. The problem is now that the sums when calculating inner products of elements in $M_\infty(E)$ (or also in products of elements in $M_\infty(\mathcal{B})$) converge only strongly. (For instance, $\mathbf{1}_\infty \in M_\infty(\mathcal{B})$ is approximated by $\mathbf{1}_m \in M_m(\mathcal{B})$.) This is a second reason why we have to switch to the W^* -case.

In the context of W^* -modules Lemma 3.2 allows for arbitrary cardinalities \aleph . We start by giving a precise meaning to $M_\aleph(\mathcal{B})$ and $M_\aleph(E)$. So let E be a W^* -correspondence over a W^* -algebra \mathcal{B} . Let H be a Hilbert space with an ONB $(e_k)_{k \in S}$ where $\#S = \aleph$. Then we set $M_\aleph(\mathcal{B}) := \mathcal{B}(H) \bar{\otimes}^s \mathcal{B}$ (tensor product of W^* -algebras) and we identify an element $B \in M_\aleph(\mathcal{B})$ with the matrix $(b_{ij})_{i,j \in S}$ where

$$b_{ij} = (e_i^* \otimes \text{id}_{\mathcal{B}})B(e_j \otimes \text{id}_{\mathcal{B}}) \in \mathcal{B}.$$

We put $M_\aleph(E) := \mathcal{B}(H) \bar{\otimes}^s E$, that is, the *exterior tensor product* of W^* -modules; see [Ske01a,

Section 4.3]. We identify an element $X \in M_n(E)$ with the matrix $(x_{ij})_{i,j \in S}$ where

$$x_{ij} = (e_i^* \otimes \text{id}_E)X(e_j \otimes \text{id}_E) \in E.$$

The operations in this correspondence over $M_n(\mathcal{B})$ are

$$\langle X, Y \rangle_{ij} = \sum_k \langle x_{ki}, y_{kj} \rangle \quad (XB)_{ij} = \sum_k x_{ik} b_{kj} \quad (BX)_{ij} = \sum_k b_{ik} x_{kj}, \quad (5.1)$$

where all sums are strong limits (see Footnote [7]). A matrix $X = (x_{ij})$ is an element of $M_n(E)$, if and only if all $\sum_k \langle x_{ki}, x_{kj} \rangle$ exist strongly and define the matrix elements of an element in $M_n(\mathcal{B})$.

Clearly, $M := H \bar{\otimes}^s \mathcal{B} = \overline{\mathcal{B}^n}^s$ is a Morita W^* -equivalence from $M_n(\mathcal{B})$ to \mathcal{B} and

$$M \bar{\otimes}^s E \bar{\otimes}^s M^* = M_n(E).$$

5.1 Corollary. *E and $M_n(E)$ are Morita equivalent W^* -correspondences.*

5.2 Proposition. *Suppose E is (strongly) full and let n be an **infinite** cardinal number not smaller than that granted by Lemma 3.2. Then $M_n(E)$ has a unit vector.*

PROOF. Denote by l the cardinal number from Lemma 3.2 and fix n as stated. Choose sets S, T with $\#S = l, \#T = n$. Let x_ℓ ($\ell \in S$) denote the components of a unit vector in $\overline{E^l}^s$. As n is infinite (by assumption!) and $l \leq n$ so that $ln = n$, we may fix a bijection $\varphi: T \rightarrow S \times T$. Denote by φ_1 and φ_2 the first and the second component, respectively, of φ . Define a matrix $X \in M_n(E)$ by setting

$$x_{ij} = x_{\varphi_1(i)} \delta_{\varphi_2(i), j}.$$

Then

$$\begin{aligned} \langle X, X \rangle_{ij} &= \sum_{k \in T} \langle x_{ki}, x_{kj} \rangle = \sum_{k \in T} \delta_{\varphi_2(k), i} \delta_{\varphi_2(k), j} \langle x_{\varphi_1(k)}, x_{\varphi_1(k)} \rangle \\ &= \sum_{(\ell, k) \in S \times T} \delta_{k, i} \delta_{k, j} \langle x_\ell, x_\ell \rangle = \left(\sum_{k \in T} \delta_{k, i} \delta_{k, j} \right) \left(\sum_{\ell \in S} \langle x_\ell, x_\ell \rangle \right) = \delta_{i, j} \mathbf{1}. \quad \blacksquare \end{aligned}$$

Putting together Corollary 5.1 and Proposition 5.2 with Theorem 4.10 completes the proof of our main theorem.

6 Full C^* -correspondences over unital C^* -algebras

In this section we prove a version of the main theorem for C^* -correspondences that have a full inner product in a unital C^* -algebra. The proof is less streamlined than that for the W^* -case, so we do not develop a complete analogue of the treatment of the W^* -version, also because, partly, it is not possible.

One problem was to have a notion of Morita equivalence that understands a full Hilbert \mathcal{B} -module E as a Morita equivalence from $\mathcal{B}^a(E)$ to \mathcal{B} and not just from $\mathcal{K}(E)$ to \mathcal{B} . In the previous sections the strongly closed versions for W^* -objects did the job. In this section we elaborate a version for *strict* closure (or what is the same for strict or $*$ -strong completion). And we elaborate this *strict* Morita equivalence only for the case, where one of the algebras is $\mathcal{B}^a(\mathcal{B})$. This will allow for the necessary matrix constructions, and Lemma 2.2 will guarantee existence of a unit vector in the matrix modules. There are two reasons why we cannot proof the result for nonunital C^* -algebras. The fact that we have available neither Lemma 2.2 nor Lemma 3.2 is responsible for that we cannot proof the result for nonunital C^* -algebras. The former works only for full Hilbert modules over unital C^* -algebras. The proof of the latter is based on quasi orthonormal bases that, in strict completions, are not available.

6.1 Example. Let $\mathcal{B} = C_0(-2, 2)$ and $I = C_0(-1, 1) \subset \mathcal{B}$ an ideal and define the full Hilbert \mathcal{B} -module $E = \mathcal{B} \oplus I$. Then the strict completion of E , $\mathcal{B}^a(\mathcal{B}, E)$, is the direct sum $C_b(-2, 2) \oplus C_0(-1, 1)$. The only projection in $\mathcal{B}^a(\mathcal{B}) = C_b(-2, 2)$ is $\mathbf{1}$. Every element in $\mathcal{B}^a(\mathcal{B}, E)$ that has unit length leaves a nonzero complement that does not admit a further unit vector. So $\mathcal{B}^a(\mathcal{B}, E)$ has no quasi orthonormal basis.

Let us start with some generalities, however, without discussing (as would be natural) how the definitions fit into the frame of multiplier algebras, double centralizers and strict topology. If E is a Hilbert \mathcal{B} -module, then by the *strict completion* of E we understand the space $\mathcal{B}^a(\mathcal{B}, E)$. If \mathcal{B} is unital, then $\mathcal{B}^a(\mathcal{B}, E)$ is just E where we consider $x \in E$ as the map $b \mapsto xb$. In general, $\mathcal{B}^a(\mathcal{B}, E)$ is Hilbert module over $\mathcal{B}^a(\mathcal{B})$ with inner product $\langle X, X' \rangle = X^*X'$. Further, $\mathcal{B}^a(\mathcal{B}, E)$ has the same operators as E , that is, $\mathcal{B}^a(\mathcal{B}^a(\mathcal{B}, E)) = \mathcal{B}^a(E)$ where an element $a \in \mathcal{B}^a(E)$ acts on $X \in \mathcal{B}^a(\mathcal{B}, E)$ simply by composition aX .

Now we wish to define an appropriate tensor product among such spaces.

6.2 Proposition. *Let E be a Hilbert \mathcal{B} -module and let F be a correspondence from \mathcal{B} to \mathcal{C} . Then:*

1. *The left action of \mathcal{B} on F extends to a (unique and strict) action of $\mathcal{B}^a(\mathcal{B})$. Therefore, also $\mathcal{B}^a(\mathcal{C}, F)$ has a left action of $\mathcal{B}^a(\mathcal{B})$.*
2. *For every $X \in \mathcal{B}^a(\mathcal{B}, E)$ by setting $\eta(X) = X \odot \text{id}_F$ we define a map in $\mathcal{B}^a(\mathcal{B} \odot F, E \odot F) = \mathcal{B}^a(F, E \odot F)$ with adjoint $\eta(X)^* = X^* \odot \text{id}_F$.*
3. *The map $X \odot Y \mapsto \eta(X)Y$ defines an isometry from the tensor product of $\mathcal{B}^a(\mathcal{B}, E)$ and $\mathcal{B}^a(\mathcal{C}, F)$ over \mathcal{B} onto a strictly dense subset of $\mathcal{B}^a(\mathcal{C}, E \odot F)$.*

PROOF. Part 1 is a statement we used already in the representation theory of general $\mathcal{B}^a(E)$ (applied to $\mathcal{B}^a(\mathcal{B})$). See, for instance, [MSS04, Corollary 1.20] for a proof.

Part 2 is general theory of tensor products, but it may also easily deduced from the proof of Part 3.

For Part 3 let us choose a bounded approximate unit $(u_\lambda)_{\lambda \in \Lambda}$ for \mathcal{B} . Then

$$\eta(X)Yc = \lim_\lambda \eta(X)u_\lambda Yc = \lim_\lambda (Xu_\lambda) \odot (Yc) \in E \odot F,$$

where we made use of $u_\lambda y \rightarrow y$ in norm for all $y \in F$. (This follows from Part 1, but may also easily be verified by three epsilons.) So the map is into $\mathcal{B}^a(C, E \odot F)$. Clearly, when restricted to the subset $E \odot F$ of $\mathcal{B}^a(\mathcal{B}, E) \odot \mathcal{B}^a(C, F)$, we obtain all maps of the form $c \mapsto cz$ for $z \in E \odot F$ that form a strictly dense subset of $\mathcal{B}^a(C, E \odot F)$. ■

6.3 Definition. By the *strict tensor product* $\mathcal{B}^a(\mathcal{B}, E) \bar{\odot} \mathcal{B}^a(C, F)$ we understand the space $\mathcal{B}^a(C, E \odot F)$.

The following corollary can be proved as Part 3.

6.4 Corollary. *For every correspondence G from C to \mathcal{D} (that may be viewed also as a correspondence G from $\mathcal{B}^a(C)$ to \mathcal{D} in a unique way) we have*

$$(\mathcal{B}^a(\mathcal{B}, E) \bar{\odot} \mathcal{B}^a(C, F)) \odot G = \mathcal{B}^a(C, E \odot F) \odot G = E \odot F \odot G$$

Clearly, if E is a correspondence from \mathcal{A} to \mathcal{B} , then $\mathcal{B}^a(C, E \odot F)$ is a correspondence from $\mathcal{B}^a(\mathcal{A})$ to $\mathcal{B}^a(C)$. In particular, if E^\odot is a product systems of correspondence over \mathcal{B} , then the family of all $\mathcal{B}^a(\mathcal{B}, E_t)$ form a strict tensor product system of correspondences over $\mathcal{B}^a(\mathcal{B})$. If this product system has a unital unit Ξ^\odot , then we may proceed as in Section 1. So $\Xi_s \odot \text{id}_{E_t}$ defines an inductive system of isometric embeddings $\mathcal{B}^a(\mathcal{B}, E_t) \rightarrow \mathcal{B}^a(\mathcal{B}, E_{s+t})$. From the inductive limit of this system we may extract a Hilbert \mathcal{B} -module

$$E_\infty = (\lim_{t \rightarrow \infty} \text{ind } \mathcal{B}^a(\mathcal{B}, E_t)) \odot \mathcal{B}$$

so that $\lim_{t \rightarrow \infty} \text{ind } \mathcal{B}^a(\mathcal{B}, E_t)$ embeds as a strictly dense subset into $\mathcal{B}^a(\mathcal{B}, E_\infty)$. (Note that \mathcal{B} is a self-inverse Morita equivalence over \mathcal{B} and that the left action of \mathcal{B} extends to a strict left action of $\mathcal{B}^a(\mathcal{B})$ on \mathcal{B} in the canonical way.) $\mathcal{B}^a(\mathcal{B}, E_\infty)$ fulfills

$$\mathcal{B}^a(\mathcal{B}, E_\infty) \bar{\odot} \mathcal{B}^a(\mathcal{B}, E_t) = \mathcal{B}^a(\mathcal{B}, E_\infty) \tag{6.1}$$

and associativity conditions similar to (1.1) so that $\vartheta_t(a) = a \odot \text{id}_{E_t}$ defines an E_0 -semigroup on $\mathcal{B}^a(\mathcal{B}^a(\mathcal{B}, E_\infty))$. But, we know that $\mathcal{B}^a(\mathcal{B}^a(\mathcal{B}, E_\infty))$ is just $\mathcal{B}^a(E_\infty)$. It is easy to show that that this E_0 -semigroup is strict and that its product system is nothing but E^\odot . The following proposition is slightly more general and implies what we just asserted in the special case $M = \mathcal{B}$.

6.5 Proposition. *Let M denote a Morita equivalence from \mathcal{B} to \mathcal{C} (so that M carries a unique and strict extension of its left action to $\mathcal{B}^a(\mathcal{B})$). Put $F_\infty := E_\infty \odot M$. Then*

$$F_\infty = F_\infty \odot (M^* \odot E_t \odot M)$$

(via (6.1) and Corollary 6.4) and $\theta_t(a) := a \odot \text{id}_{M^* \odot E_t \odot M}$ defines a strict E_0 -semigroup on $\mathcal{B}^a(F_\infty)$ whose product system is $F^\odot := M^* \odot E^\odot \odot M$.

PROOF. The isomorphism $F_\infty = F_\infty \odot F_t$ is

$$\begin{aligned} F_\infty &= E_\infty \odot M = \mathcal{B}^a(\mathcal{B}, E_\infty) \odot M = (\mathcal{B}^a(\mathcal{B}, E_\infty) \bar{\odot} \mathcal{B}^a(\mathcal{B}, E_t)) \odot M \\ &= E_\infty \odot E_t \odot M = E_\infty \odot M \odot M^* \odot E_t \odot M = F_\infty \odot F_t. \end{aligned}$$

The remaining statements follow as in the second half of the proof of Theorem 4.10 just the roles of E^\odot and F^\odot have now switched. ■

We are now in a position to proof the analogue of the main theorem in the situation of this section.

6.6 Theorem. *Let E be a full correspondence over a unital C^* -algebra \mathcal{B} . Then there is a (necessarily full) Hilbert \mathcal{B} -module F and a unital endomorphism ϑ of $\mathcal{B}^a(F)$ such that $F_\vartheta = E$.*

PROOF. Denote by $E^\odot = (E^{\odot n})_{n \in \mathbb{N}_0}$ the product system generated by E . We define $M_\infty(\mathcal{B})$ and $M_\infty(E_t)$ as the completions of the spaces of matrices with finitely many nonzero entries in the respective norm topologies and operations like in (5.1). To come to the setting of the preceding proposition we make up a dictionary.

Propositions 6.5	here
\mathcal{B}	$M_\infty(\mathcal{B})$
\mathcal{C}	\mathcal{B}
M	\mathcal{B}^∞
E^\odot	$M_\infty(E^\odot) := (M_\infty(E_n))_{n \in \mathbb{N}_0}$
F^\odot	E^\odot
F_∞	F
θ	ϑ

In order to apply Proposition 6.5 (providing us with the F and the ϑ we seek according to the dictionary) it remains to show that $\mathcal{B}^a(M_\infty(\mathcal{B}), M_\infty(E))$ has a unit vector Ξ (determining a unital unit Ξ^\odot for the whole product system $M_\infty(E^\odot)$ as ingredient). But this can be done as in Proposition 5.2 using, however, the ingredients from Lemma 2.2 (that is, 1 finite so that $n = \#\mathbb{N}$ is sufficient) instead of those from Lemma 3.2. ■

6.7 Remark. If a product system E^\odot consists of correspondences over a nonunital C^* -algebra \mathcal{B} , then it is no problem to pass to the unitization $\widetilde{\mathcal{B}}$ of \mathcal{B} . However, even if E^\odot was full, now it is no longer full so Theorem 6.6 is not applicable.

In this context we would like to mention that the whole basic classification of product systems is based on the existence of units — of continuous units to be precise. A unit, by definition, assumes the value $\mathbf{1} \in E_0 = \mathcal{B}$ at $t = 0$, so that the concept of unit is not defined. Even if we try to admit units in also for nonunital \mathcal{B} , then the continuity conditions would require that $\langle \xi_t, \xi_t \rangle$ converges to some projection p in \mathcal{B} . But there are projectionless C^* -algebras. Anyway, allowing for units not converging to identity at $t = 0$ would considerably weaken the conclusions that may be drawn from existence of units and change completely the classification results from [Ske01b, BBL04, Ske03b]. For instance, if \mathcal{B} is a projectionless C^* -algebra, then the time-ordered product system over \mathcal{B} (consisting of time-ordered Fock modules) would not have a single continuous unit, not even a trivial one. The only solution would be to take $t = 0$ away from the product system. But it is the classification as in [Ske01b, BBL04, Ske03b] which is most similar to Arveson's for Arveson systems.

7 Commutants and nondegenerate representations

In this section we put the result into relation with a recent result by Hirshberg [Hir05a] on existence of nondegenerate (or *essential*) representations of a correspondence over a C^* -algebra. As a consequence, we furnish a different proof of Hirshberg's result. Moreover, our method shows that in the case of a W^* -correspondence the representation can be chosen compatible with the σ -weak topology of that W^* -correspondence, a result that was not accessible by the method in [Hir05a].

Our approach here makes use of a duality between von Neumann correspondences over the von Neumann algebra $\mathcal{B} \subset \mathcal{B}(G)$ and its *commutant*, a von Neumann correspondence over the commutant of \mathcal{B} . By the crucial Theorem 7.4 under the commutant our main theorem applied to a von Neumann correspondence translates into Hirshberg's result for the commutant (and back) in a version von Neumann correspondences. Von Neumann modules (Skeide [Ske00a]) and commutants of von Neumann correspondences (Skeide [Ske03a, Ske05a]) play a crucial role, so we have to give a brief account on these subjects.

Let K be a Hilbert space. A **representation** of a correspondence E over \mathcal{B} on K is a pair (ψ, η) of linear mappings $\psi: \mathcal{B} \rightarrow \mathcal{B}(K)$ and $\eta: E \rightarrow \mathcal{B}(K)$ where ψ is a representation of \mathcal{B} and η is a bimodule map (that is, $\eta(b_1 x b_2) = \psi(b_1) \eta(x) \psi(b_2)$) such that $\eta(x)^* \eta(y) = \psi(\langle x, y \rangle)$. We always assume that ψ is nondegenerate. The representation (ψ, η) is **nondegenerate** (or **essential**), if also η is nondegenerate, that is, if $\overline{\text{span}} \eta(E)K = K$.

Observe that ψ and, therefore, also η are contractions. Moreover, like every representation of

a C^* -algebra on a pre-Hilbert space, the induced representation of $\mathcal{B}^a(E)$ on $\text{span } \eta(E)K$ defined by $a \mapsto (\eta(x)k \mapsto \eta(ax)k)$ is by bounded operators and, therefore, extends to a representation first on $\overline{\text{span}} \eta(E)K$ and then to K (acting as 0 on $\eta(E)K^\perp$). In particular, (ψ, η) extends to a (nondegenerate) representation of the *linking algebra* $\begin{pmatrix} \mathcal{B} & E^* \\ E & \mathcal{B}^a(E) \end{pmatrix}$ on $K \oplus \overline{\text{span}} \eta(E)K$ so that all mappings are even completely contractive.

7.1 Remark. The nomenclature here differs, for instance, from Muhly and Solel [MS98], who call *covariant representation* a pair (ψ, η) of completely contractive mappings without $\eta(x)^*\eta(y) = \psi(\langle x, y \rangle)$. They call a covariant representation *isometric* if also $\eta(x)^*\eta(y) = \psi(\langle x, y \rangle)$ holds, and *fully coisometric* if $\overline{\text{span}} \eta(E)K = K$.

The connection between representations of correspondences and endomorphisms of some $\mathcal{B}^a(E)$ is not direct, but via a duality, called *commutant*, between von Neumann correspondences over \mathcal{B} and von Neumann correspondences over \mathcal{B}' as introduced in Skeide [Ske03a]. (Independently, Muhly and Solel [MS04] have considered a W^* -version and in [MS05] generalized the construction to \mathcal{A} - \mathcal{B} -correspondences.) It works nicely only if we pass to W^* -algebras that are represented concretely as operator algebras acting (nondegenerately) on a Hilbert space, that is, if we pass to *von Neumann algebras*, and to W^* -modules that are concrete spaces of operators between Hilbert spaces, that is, if we pass to *von Neumann modules*; see Skeide [Ske00a].

Let $\mathcal{B} \subset \mathcal{B}(G)$ be a von Neumann algebra (acting nondegenerately on the Hilbert space G). Then every (pre-)Hilbert \mathcal{B} -module E may be identified as a concrete operator \mathcal{B} -submodule of some $\mathcal{B}(G, H)$ (nondegenerate in the sense that $\overline{\text{span}} EG = H$) in the following way. Put $H = E \odot G$. For every $x \in E$ define the mapping $L_x: g \mapsto x \odot g$ in $\mathcal{B}(G, H)$. Then $L_{xb} = L_x b$ and $L_x^* L_y = \langle x, y \rangle$. Identifying x with L_x identifies E as a subspace of $\mathcal{B}(G, H)$. Following [Ske00a], we say E is a *von Neumann \mathcal{B} -module*, if E is strongly closed in $\mathcal{B}(G, H)$.

7.2 Remark. One may show that E is a von Neumann module, if and only if E is self-dual, that is, if and only if E is a W^* -module; see [Ske00a, Ske05c]. The point about von Neumann modules is that it is easier to obtain them (from pre-Hilbert modules over a von Neumann algebra) than W^* -modules. Simply take strong closure. In the sequel, we will learn another possibility that is completely algebraic and parallels the operation of taking the double commutant of an operator algebra in order to obtain a von Neumann algebra; see Remark 7.3.

$\mathcal{B}^a(E)$ embeds into $\mathcal{B}(H)$ via the identification $a = a \odot \text{id}_G$. Clearly, if E is a von Neumann module, then $\mathcal{B}^a(E)$ is a von Neumann subalgebra of $\mathcal{B}(H)$. When E is also a correspondence over \mathcal{B} such that the canonical representation $\rho: \mathcal{B} \rightarrow \mathcal{B}^a(E) \rightarrow \mathcal{B}(H)$ is normal, then we say E is a *von Neumann correspondence*. (So E is a von Neumann correspondence, if and only if it is also a W^* -correspondence.) We refer to ρ as the *Stinespring representation* of \mathcal{B} .

On H there is a second (normal nondegenerate) representation, namely, the so-called **commutant lifting** ρ' of \mathcal{B}' defined as $\rho'(b') = \text{id}_E \odot b'$. It is not difficult to show that the intertwiner space $C_{\mathcal{B}'}(\mathcal{B}(G, H)) := \{x \in \mathcal{B}(G, H) : \rho'(b')x = xb' \ (b' \in \mathcal{B}')\}$ is a von Neumann \mathcal{B} -module (see [Rie74b]) and that E is a von Neumann module, if and only if $E = C_{\mathcal{B}'}(\mathcal{B}(G, H))$ (see [Ske05c]). Less obvious is the converse statement: If ρ' is a normal nondegenerate representation of \mathcal{B}' on a Hilbert space H , then the von Neumann \mathcal{B} -module $E := C_{\mathcal{B}'}(\mathcal{B}(G, H))$ acts nondegenerately on G (see [MS02]), that is, $E \odot G = H$ via $x \odot g = xg$. Clearly, the commutant lifting for that E is the ρ' we started with.

7.3 Remark. If E is only a pre-Hilbert module over the von Neumann algebra \mathcal{B} , then \overline{E}^s is just $C_{\mathcal{B}'}(\mathcal{B}(G, H))$ and provides us with the minimal self-dual extension of E in the sense of Paschke [Pas73]; see [Rie74b, Ske05c]. This is the *double commutant theorem* for von Neumann modules.

Summarizing, we have a one-to-one correspondence between von Neumann \mathcal{B} -modules and representations of \mathcal{B}' and a one-to-one correspondence between von Neumann correspondences E over \mathcal{B} and pairs of representations (ρ, ρ', H) of \mathcal{B} and \mathcal{B}' with mutually commuting range. (See also Connes [Con80], taking into account that the commutant of \mathcal{B} in standard representation is \mathcal{B}^{op} .) In the latter picture of correspondences as two representations nobody prevents us from exchanging the roles of \mathcal{B} and \mathcal{B}' . In that way, we obtain a further von Neumann correspondence, namely

$$E' := C_{\mathcal{B}}(\mathcal{B}(G, H)) := \{x' \in \mathcal{B}(G, H) : \rho(b)x' = x'b \ (b \in \mathcal{B})\},$$

this time over \mathcal{B}' with left action of \mathcal{B}' via ρ' . This duality between E and its **commutant** E' was mentioned in [Ske03a]. See [Ske06] for a careful discussion about “one-to-one” and Skeide [Ske05a] for definitions where the commutant becomes, really, a bijective functor.

We are now in a position to formulate the theorem about the relation between nondegenerate representations and derivability from unital endomorphisms. But first let us recall a few (easy to proof) facts. A von Neumann \mathcal{B} -module is strongly full, if and only if the commutant lifting ρ' is faithful. In turn, this implies that a von Neumann correspondence E over \mathcal{B} is strongly full, if and only if the left action of \mathcal{B}' on the commutant E' defines a faithful representation of \mathcal{B}' on E' . Observe also that a representation (ψ, η) of a correspondence E over \mathcal{B} is faithful (in the sense that both ψ and η are faithful), if and only if ψ is faithful. And if (ψ', η') is a faithful nondegenerate representation of E' , so that left action of \mathcal{B}' on E' is faithful, then necessarily E is strongly full. Finally, note that for a representation (ψ, η) of a W^* -correspondence E being σ -weak, it is necessary and sufficient that ψ is normal. (This is so, because every normal representation ψ of \mathcal{B} on K extends to a unique normal representation of the linking algebra $\begin{pmatrix} \mathcal{B} & E^* \\ E & \mathcal{B}^{\alpha(E)} \end{pmatrix}$ on $\mathcal{B}(K \oplus (E \odot K))$ and η , when considered as mapping $E \rightarrow \mathcal{B}(K, \overline{\text{span}} \eta(E)K)$, is

unitarily equivalent to the restriction of that representation to the corner E .) Therefore, we speak of a **normal** representation (ψ, η) , if ψ is normal.

7.4 Theorem. *Let E be a von Neumann correspondence over a von Neumann algebra $\mathcal{B} \subset \mathcal{B}(G)$ and E' its commutant. Then the following conditions are equivalent.*

1. *E is the correspondence of a normal unital endomorphism ϑ of $\mathcal{B}^a(F)$ for some strongly full von Neumann \mathcal{B} -module F .*
2. *E' admits a faithful normal nondegenerate representation (ψ', η') on a Hilbert space K .*

Moreover, if either of the conditions is fulfilled, then E is strongly full or, equivalently, the left action of \mathcal{B}' on E' is faithful.

PROOF. Let (ρ, ρ', H) be the triple that determines E as $C_{\mathcal{B}'}(\mathcal{B}(G, H))$ and E' as $C_{\mathcal{B}}(\mathcal{B}(G, H))$.

Suppose that F is a strongly full von Neumann \mathcal{B} -module and that ϑ is a normal unital endomorphism of $\mathcal{B}^a(F)$ such that $F = F \bar{\odot}^s E$ and $\vartheta(a) = a \odot \text{id}_E$. (As F is strongly full, E is uniquely determined by these properties and necessarily E is itself strongly full.) Put $K := F \odot G$. As $F = F \bar{\odot}^s E$ we have $K = F \odot E \odot G$. (If the last factor in a tensor product is a Hilbert space, then norm closure is sufficient.) By construction we have $E \odot G = \overline{\text{span}} EG = H = \overline{\text{span}} E'G = E' \odot G$. (Note that $\overline{\text{span}} EG = H = \overline{\text{span}} E'G$ is true equality of Hilbert spaces. The equalities $E \odot G = \overline{\text{span}} EG$ and $E' \odot G = \overline{\text{span}} E'G$ are by canonical isomorphism. The equality $F = F \bar{\odot}^s E$ is by $(*)$ and canonical by a suitable universal property.) We find

$$F \odot G = K = F \odot E' \odot G.$$

There are several ways to understand why $\eta'(x'): y \odot g \mapsto y \odot x' \odot g$ is a well-defined element of $\mathcal{B}(K)$. One is that $\eta'(x') = \text{id}_F \odot x'$ where x' is considered a \mathcal{B} - \mathbb{C} -linear operator from G to $H = E' \odot G$. Let ψ' denote the (normal!) commutant lifting of \mathcal{B}' on $K = F \odot G$. We leave it as an instructive exercise to check that (ψ', η') is a representation of E' on K . Obviously this representation is nondegenerate. It is normal, because ψ' is normal. It is faithful because F is strongly full.

Suppose now that (ψ', η') is a faithful normal nondegenerate representation of E' on K . We put $F := C_{\mathcal{B}'}(\mathcal{B}(G, K))$. As ψ' is faithful, F is strongly full. Again

$$F \odot G = K = F \odot E' \odot G$$

now via $\eta'(x')(y \odot g) \mapsto y \odot x' \odot g$. (Note that the set $\eta'(E')F \odot G$ is total in K , because η' is nondegenerate.) Again we substitute $E' \odot G = H = E \odot G$ so that $F \odot G = F \odot E \odot G$. The action of $b' \in \mathcal{B}'$ on these spaces is the same. To see this we observe, first, that $b'(y \odot x \odot g) = y \odot x \odot b'g = y \odot \rho'(b')(x \odot g)$. Then, writing a typical element of $H = E \odot G$ not as elementary

tensor $x \odot g$ but as elementary tensor $x' \odot g$ and recalling that the action of b' on $x' \odot g$ is just $\rho'(b')$, we find

$$b'(y \odot x' \odot g) = y \odot \rho'(b')(x' \odot g) = y \odot b'x' \odot g = \eta'(b'x')(y \odot g) = \psi'(b')\eta'(x')(y \odot g).$$

As the commutant liftings on $F \odot G$ and on $F \odot E \odot G$ coincide, also the modules F and $F \bar{\odot}^s E$ (being intertwiner spaces for the same commutant lifting) must coincide and $\vartheta(a) = a \odot \text{id}_E$ induces a unital normal endomorphism of $\mathcal{B}^a(E)$. Once again, as F is strongly full, a correspondence E is determined uniquely by these properties, so that $F^* \bar{\odot}^s {}_{\vartheta}F$ gives us back E . ■

7.5 Remark. The theorem has an obvious generalization to product systems; see Definition 8.1 and the discussion that follows that definition.

With this theorem the following two results are corollaries of our main theorem. The first one is a version of Hirshberg's result for von Neumann correspondences (assuring that the faithful nondegenerate representation can be chosen normal what was not accessible to the methods in [Hir05a]). The second is Hirshberg's result for C^* -correspondences now with a completely different proof.

7.6 Theorem. *Every W^* -correspondence over a W^* -algebra with a faithful left action admits a normal faithful nondegenerate representation on a Hilbert space.*

PROOF. In order to pass to von Neumann correspondences choose a faithful (normal nondegenerate) representation of the W^* -algebra on a Hilbert space G and identify the W^* -algebra as von Neumann algebra acting on G . In order to be compatible with the notations in the remainder of these notes, we denote the von Neumann algebra by \mathcal{B}' , its commutant by $\mathcal{B} := \mathcal{B}'$ and the correspondence over \mathcal{B}' by E' . The statement of the theorem is now: E' admits a normal faithful nondegenerate representation. By Theorem 7.4 this is equivalent to that the commutant $E := E''$ of E' is derivable from an endomorphism. As E is strongly full, this is granted by our main theorem. ■

7.7 Theorem. *Every C^* -correspondence over a C^* -algebra with a faithful left action admits a faithful nondegenerate representation on a Hilbert space.*

PROOF. Suppose we choose a faithful nondegenerate representation of the C^* -algebra on a Hilbert space G , so that the correspondence becomes identified as a submodule of $\mathcal{B}(G, H)$ for some Hilbert space H . On H there is a commuting pair of a representation ρ' of the C^* -algebra and a normal representation ρ of its commutant.

So far, we do not yet know whether ρ' extends to a normal representation of the double commutant of the C^* -algebra. (We thank B. Solel for having pointed out to us that gap in the

first version.) The problem disappears, however, if we choose for the faithful representation ρ' from the beginning the universal representation. (In that case, the double commutant is isomorphic to the bidual so that every representation extends to a unique normal representation.)

The intertwiner space for the commutant lifting ρ is a von Neumann correspondence over the bicommutant of the C^* -algebra (namely the strong closure of the original correspondence in the operator space $\mathcal{B}(G, H)$ and also its bicommutant). Now apply Theorem 7.6 to that bicommutant and restrict the resulting representation to the (strongly dense!) original correspondence. By σ -weak continuity also this restriction must act nondegenerately. ■

7.8 Remark. Muhly and Solel [MS99] have constructed from a nondegenerate representation (ψ', η') on K an endomorphism of $\psi'(\mathcal{B}') \subset \mathcal{B}(K)$. Taking into account that this algebra coincides exactly with our $\mathcal{B}^a(F) \subset \mathcal{B}(K)$ puts into perspective the second part of the proof of Theorem 7.4 with the result from [MS99]. In fact, the constructions of the endomorphism are very much the same, except that we have added the construction of F and the interpretation of the algebra on which the endomorphism acts as $\mathcal{B}^a(F)$. This considerably facilitates understanding why everything is well-defined.

7.9 Remark. Theorem 7.4 describes (under the condition that E is full, respectively, E' has faithful left action) a general correspondence between endomorphisms and representations. Recall again that the crucial point in the main theorem is that the endomorphism be unital. In Theorem 7.4 this corresponds exactly to that the representation of the commutant be nondegenerate. It is always easy to construct a nonunital endomorphism of $\mathcal{B}^a(L^2(E^\circ))$ and on the level of the commutants this corresponds exactly to constructing the standard representation of the Cuntz-Pimsner algebra over E' (in the discrete case) or of the spectral algebra over the product system E'^{\otimes^s} (in the continuous case); see Hirshberg [Hir05b]. Note also here the anti-symmetry between E , that acts by tensoring from the right to generate the endomorphism, and E' , that acts by tensoring from the left to define the action of its representation.

Suppose now that $E = H$ is a Hilbert space of dimension $n = 2, 3, \dots, \infty$. Then the commutant H' of H is isomorphic to H and we recover the well-known fact that representations of the Cuntz algebra \mathcal{O}_n correspond to endomorphisms of index n of $\mathcal{B}(K)$, and that nondegenerate representations correspond to unital endomorphisms. Note that the isomorphism $H \cong H'$ is by no means a trivial issue. One may see this by looking at the discrete product systems generated by H and H' , respectively. One is the commutant of the other, but their product system structures are anti-isomorphic. This is the same relation as that between the Bhat system and the Arveson system constructed from an E_0 -semigroup on $\mathcal{B}(K)$; see also Remark 8.4.

Theorem 7.4 can be refined to the case when F is not necessarily full and, consequently, ψ' is not necessarily faithful. The construction of a nondegenerate (ψ', η') in the first part of the proof

works also if we start from the situation $F = F \bar{\odot}^s E$, $\vartheta(a) = a \odot \text{id}_E$. It is not difficult to show that $\overline{\mathcal{B}_F}^s = \overline{\mathcal{B}_E}^s$, if and only if $\ker \psi' = \ker \rho'$. (In fact, the central projection p_F that generates the ideal $\overline{\mathcal{B}_F}^s$ is the “orthogonal complement” of the central projection $\mathbf{1} - p_F$ that generates the ideal $\ker \psi'$; see [Ske06].) The correspondence associated with ϑ is $F^* \bar{\odot}^s \vartheta F = p_F E p_F$. So in order that E is derived from an endomorphism of $\mathcal{B}^a(F)$ we may assume that $\overline{\mathcal{B}_F}^s = \overline{\mathcal{B}_E}^s$ (that is, $\ker \psi' = \ker \rho'$) and that $\overline{\mathcal{B}_E}^s$ acts nondegenerately on E .

Under these circumstances our main theorem and the consequences Theorems 7.6 and 7.7 remain true when we dispense with the requirement that the representation of E' be faithful. However, it is surprisingly tricky to find out what the condition $\overline{\mathcal{B}_E}^s E = E$ means when expressed exclusively in terms of the correspondence E' . A constructive approach can be based on the result from [Ske03a, Theorem 2.3] that $(E \bar{\odot}^s F)' = F' \bar{\odot}^s E'$ for every pair of von Neumann correspondences over \mathcal{B} . (We observe that the condition $\overline{\text{span}}^s \overline{\mathcal{B}_E}^s E = E$ can be rephrased as $\overline{\mathcal{B}_E}^s \bar{\odot}^s E = E$. Applying the mentioned result we obtain $E' \bar{\odot}^s \overline{\mathcal{B}_E}^s = E'$. A careful discussion shows that the commutant of the correspondence $\overline{\mathcal{B}_E}^s$ over \mathcal{B} is just $p_E \mathcal{B}'$ where p_E is the central projection generating $\overline{\mathcal{B}_E}^s$. So the condition means that $x' p_E = x'$ for all $x' \in E'$ or $x' b' = 0$ for all $x' \in E'$ and all $b' \in (\mathbf{1} - p_E) \overline{\mathcal{B}_E}^s = \ker \rho'$.) Here we just state the condition in terms of E' and prove that it does the job.

7.10 Proposition. *Let E' be a von Neumann correspondence over \mathcal{B}' . Then $E := E'$ satisfies $\overline{\mathcal{B}_E}^s E = E$, if and only if E' satisfies the condition*

$$E' \cdot \ker \rho' = \{0\},$$

where $\ker \rho'$ is the kernel of the canonical representation of \mathcal{B}' on E' by left multiplication.

PROOF. Let p_E denote the central projection such that $(\mathbf{1} - p_E) \mathcal{B}' = \ker \rho'$. We know that this means that $p_E \mathcal{B} = \overline{\mathcal{B}_E}^s$. Of course, $E' \cdot \ker \rho' = \{0\}$, if and only if $x'(\mathbf{1} - p_E) = 0$ or $x' = x' p_E = \rho(p_E)x'$ for all x' , that is, if and only if $\rho(p_E)$ acts as identity on $E' \odot G = H = E \odot G$, in other words, if and only if the left action of $\overline{\mathcal{B}_E}^s$ on E is nondegenerate. ■

A way to express this without using $\ker \rho'$ is that for every $b' \in \mathcal{B}'$

$$b' x' = 0 \forall x' \in E' \implies x' b' = 0 \forall x' \in E'. \quad (7.1)$$

We have learned that if E fulfills the condition on \mathcal{B}_E , then it may be considered as a correspondence over $\overline{\mathcal{B}_E}^s$. We find the analogue for E' .

7.11 Corollary. *A von Neumann correspondence E over \mathcal{B} maybe considered a von Neumann correspondence E over $\overline{\mathcal{B}_E}^s$ (that is, $\overline{\mathcal{B}_E}^s$ acts nondegenerately), if and only if its commutant E' may be considered a von Neumann correspondence E over $\mathcal{B}' / \ker \rho'$ (that is, the inner product of E' has values only in $\mathcal{B}' / \ker \rho'$).*

7.12 Corollary. *If E' is a correspondence over a C^* -algebra \mathcal{B}' fulfilling (7.1), then E' admits a nondegenerate representation (ψ', η') . If E' is also a W^* -correspondence, then this representation can be chosen normal.*

8 Remarks on the duality between representations and dilation of product systems

We have shown in our main theorem that every discrete product system of full W^* -correspondences comes from a discrete normal E_0 -semigroup. The following definition (from [MS04], but in a different terminology; see Remark 7.1) extends suitably the definition of a representation of a single correspondence to the definition of a representation of a whole product system.

8.1 Definition. A *representation* of a product system E^\odot of correspondences over a C^* -algebra \mathcal{B} is a pair (ψ, η) where ψ is a nondegenerate representation of \mathcal{B} on a Hilbert space K and $\eta = (\eta_t)_{t \in \mathbb{S}}$ is a family such that each (ψ, η_t) is a representation of E_t on K and such that

$$\eta_{s+t}(x_s \odot y_t) = \eta_s(x_s)\eta_t(y_t).$$

A representation is *nondegenerate*, if every (ψ, η_t) is nondegenerate. In case of product systems of W^* -correspondences we require that ψ (and, therefore, every (ψ, η_t)) is normal.

The main theorem together with Theorem 7.4 states now that the commutant system of a discrete product system has a faithful nondegenerate normal representation. Speaking about a whole product system instead of a single correspondence, Theorem 7.4 remains true (with practically no changes in the proof, appart from a view more indices) for product systems of von Neumann correspondences indexed by \mathbb{N}_0 or \mathbb{R}_+ . We phrase it here.

8.2 Theorem. *Let E^{\odot^s} be a product system of von Neumann correspondences over a von Neumann algebra $\mathcal{B} \subset \mathcal{B}(G)$ and E'^{\odot^s} its commutant. Suppose that all E_t are strongly full or, equivalently, that the left action of \mathcal{B}' is faithful on all E'_t . Then there is a one-to-one correspondence between normal E_0 -semigroups ϑ associated with E^{\odot^s} (acting on a necessarily strongly full von Neumann \mathcal{B} -module) and nondegenerate normal faithful representations (ψ', η') of E'^{\odot^s} .*

PROOF. Just do for every couple ϑ_t and η_t what we did in the proof of Theorem 7.4 for single mappings, and verify the additional conditions. This proceeding reveals also automatically how the product system structure of the commutant of a product system must be defined. ■

8.3 Remark. The theorem has two extensions. The first is to the nonfull case. Here we must require that $\overline{\mathcal{B}_{E_t}^s}$ is stationary for $t > 0$ and acts nondegenerately on all E_t . (Recall from Section

1 that $E_0 := \mathcal{B}$ is defined by hand.) We may also use Corollary 7.11 to phrase an equivalent condition on the E'_t . Dropping strong fullness, on the commutant side this leads to possibly non faithful ψ' where, however, still every η'_t is injective. All this can be proved very simply, by restricting $\overline{\mathcal{B}_{E_t}}^s$ to the smaller algebra $\overline{\mathcal{B}_{E_t}}^s$ (acting nondegenerately on a subspace of G) and its commutant. Then we are in the strongly full case and, after applying Theorem 8.2 the missing part of \mathcal{B} may be added without any problem.

The second extension is to E -semigroups, that is, to semigroups of non necessarily unital endomorphisms. (The definition of the product system associated with an E -semigroup on $\mathcal{B}^a(E)$ is the same. The only difference is that now we do no longer obtain an isomorphism $E \odot E_t \rightarrow E$ but only an isometry onto the subspace $\vartheta_t(\mathbf{1})E$ of E .) On the commutant side this leads to possibly degenerate representations. In this setting we are no longer sure that $\overline{\mathcal{B}_{E_t}}^s$ is stationary for $t > 0$, so we possibly leave also the strongly full case. This time η'_t need no longer be injective. Anyway, also in this case we remain with a one-to-one correspondence of E -semigroups associated with $E^{\odot s}$ and normal representations of $E'^{\odot s}$.

8.4 Remark. In the case of continuous time $t \in \mathbb{R}_+$ (and under further technical conditions product system and E_0 -semigroups must fulfill) Arveson [Arv89] showed that every Arveson system comes from an E_0 -semigroup on $\mathcal{B}(H)$. But a careful analysis shows that the product system Arveson associates with the E_0 -semigroup is actually the commutant of that one we would associate with the E_0 -semigroup as Bhat [Bha96] does. (By accident $\mathbb{C}' = \mathbb{C}$, but, as [Ske03a, Theorem 2.3] shows, the product system and its commutant are anti-isomorphic. Tsirelson [Tsi00] has shown that the two need not be isomorphic.) And what Arveson actually showed is that every Arveson system admits a faithful nondegenerate representation. (Hirshberg's result [Hir05a] is based on ideas from [Arv89].) Therefore, Arveson showed that every Arveson system can be obtained as the commutant of the Bhat system of an E_0 -semigroup on $\mathcal{B}(H)$. After the product system version of Theorem 7.4 (that is inherent in Arveson's work) we obtain the statement that every Bhat system (that is, of course, also an Arveson system) is the Bhat system of an E_0 -semigroup on $\mathcal{B}(H)$.

Liebscher [Lie03] furnished us with a new proof of Arveson's result in which commutants or intertwiners do not play a role. Instead his proof is based on vector states, that is, on unit vectors. We strongly believe that there is a good chance that Lemma 3.2 and its use via Proposition 5.2 and Theorem 4.10 can be a promising starting point to adapt Liebscher's proof to our case. (The use of Theorem 4.10 would not be direct but in two steps. First, we show that the product system is Morita equivalent to one where every member has a unit vector. Then, we apply Liebscher's ideas to show that this second one comes from an E_0 -semigroup and, therefore, is Morita equivalent to one with a unital unit. After that also the original one is Morita equivalent to that with the unit.) But we should also say that there are serious obstacles. For instance, the

possibility to write $\mathcal{B}(H_s \otimes H_t)$ as $\mathcal{B}(H_s) \bar{\otimes}^s \mathcal{B}(H_t)$ plays a crucial role, but a similar identity does not hold for $E_s \odot E_t$. We have some ideas to overcome that difficulty but, for the time being, we are not sure that it will work. It is also possible that the method will work via a deviation to *free product systems* (see Skeide [Ske01b]), replacing tensor products of $\mathcal{B}(H_s)$ and $\mathcal{B}(H_t)$ by (equivalence classes of) reduced free products.

9 An example

In this section we discuss in detail what our constructions assert for the correspondence in Example 2.1. The reader might object that this correspondence is a Morita equivalence and that, therefore, the endomorphism granted by our main theorem is an automorphism. However, the discussion will show that we have to work considerably already in this simple case, if we wish to see explicitly what our construction does. As a side product we find a new interpretation of the Sz.-Nagy-Foias dilation of an isometry to a unitary.

In fact, our construction involves an inductive limit, that is, a highly abstract construction. Understanding the result of such an abstract construction as a concrete space, for instance as a space of functions on a concrete set with values in a Hilbert space or module (like, for instance, the elements of a Fock space or module), is a difficult issue. In the case of commutative algebras, that is, of a classical Markov process, our inductive limit would consist of functions on the abstract Kolmogorov product space, that is, on a projective limit, with all known hard problems when one wishes to capture it concretely. In certain simple cases, namely, when the product systems consists of Fock modules and the unit with respect to which we construct the inductive limit is central, we recognize the inductive limit explicitly as Fock module; see Skeide [Ske01a, Example 11.5.2]. When the unit is not central, but fulfills some technical conditions then quantum stochastic calculus may help to show that the inductive limit must be at least contained in a Fock module; see Goswami and Sinha [GS99] and Skeide [Ske00b].

The general case is perfectly unclear. In fact, we will start not directly with Example 2.1, but with a still simpler case. As algebra we choose $\mathcal{B}(G)$, where G is some Hilbert space, and as correspondence also $\mathcal{B}(G)$, the trivial $\mathcal{B}(G)$ -correspondence. Then $(\mathcal{B}(G)^{\odot n})_{n \in \mathbb{N}_0}$ with $\mathcal{B}(G)^{\odot n} = \mathcal{B}(G)$ via $a_1 \odot \dots \odot a_n = a_1 \dots a_n$ is a product system where $a_n \odot a_m = a_n a_m$ defines the isomorphism $\mathcal{B}(G)^{\odot n} \odot \mathcal{B}(G)^{\odot m} = \mathcal{B}(G)^{\odot(n+m)}$.

Clearly, $\mathcal{B}(G)$ has the unit vector id_H . If we use that unit vector for the inductive limit construction, then we are in the situation of the “if” direction in the proof of Theorem 4.10 (with $\mathcal{B}(G)$ instead of $\mathcal{B}^a(E)$ and with the trivial product system structure instead of a nontrivial one). There is nothing surprising in this case. The induced endomorphism is just the identity. (This is due to the trivial product system structure as compared with the proof of Theorem 4.10 that, in general, leads to true endomorphisms.) Also if we use a unitary $u \in \mathcal{B}(G)$ then practically

nothing changes. Just the identity as induced endomorphism is replaced by conjugation with u .

In general, a unit vector in $\mathcal{B}(G)$ is just an isometry. So what does our construction, if we suppose that $v \in \mathcal{B}(G)$ is a nonunitary isometry? Of course, as $\mathcal{B}(G)$ is a Morita equivalence the induced endomorphism still must be an automorphism. Further, the inductive limit is a von Neumann $\mathcal{B}(G)$ -module so that the algebra of operators on which the automorphism acts is some $\mathcal{B}(H)$ and also an automorphism on that $\mathcal{B}(H)$ must be implemented by a unitary (this time on H).

Let us construct H , following step by step the inductive limit construction described in [BS00] and in Section 1. $\mathcal{B}(G) = \mathcal{B}(G)^{\odot m}$ embeds into $\mathcal{B}(G) = \mathcal{B}(G)^{\odot(n+m)}$ as $v^n \odot \mathcal{B}(G)^{\odot m}$. The inductive limit into which $\mathcal{B}(G)^{\odot m}$ embeds is always the same, but the embedding depends on n . When v is a pure isometry, then we can even say that the image “disappears” (in the strong topology) as n tends to infinity. So to view $\mathcal{B}(G)^{\odot n}$ as $\mathcal{B}(G)$ is not a good idea, if we want to capture the inductive limit $n \rightarrow \infty$.

Instead of “shifting” G to smaller and smaller subspaces $v^n G$ of G , we try to change to a point of view where $v^n G$ is fixed. That makes it necessary to “extend” G into the other direction in each step by exactly the portion that would be missing in vG to G . A precise formulation of this intuitive idea leads to the following construction. Put $G_0 := (vG)^\perp \subset G$ and define a unitary

$$v_0 = v \oplus \text{id}_{G_0} : G \oplus G_0 \longrightarrow vG \oplus G_0 = G.$$

Further, we put $H := G \oplus \bigoplus_{k=1}^{\infty} G_0$ and define a unitary $u \in \mathcal{B}(H)$ as

$$u = v_0^* \oplus \text{id} : G \oplus \bigoplus_{k=1}^{\infty} G_0 \longrightarrow (G \oplus G_0) \oplus \bigoplus_{k=1}^{\infty} G_0 = G \oplus \bigoplus_{k=0}^{\infty} G_0 = G \oplus \bigoplus_{k=1}^{\infty} G_0, \quad (9.1)$$

where in the last step we simply shift a sequence starting with index 0 to one starting with index 1.

The restriction of u^n to G defines an isometric embedding k_n of $\mathcal{B}(G) = \mathcal{B}(G)^{\odot n}$ onto the (right) submodule $\mathcal{B}(G, G \oplus \bigoplus_{k=1}^n G_0)$ of $\mathcal{B}(G, H)$. We claim $\mathcal{B}(G, H)$ is the inductive limit $\lim \text{ind}_{n \rightarrow \infty} \mathcal{B}(G)^{\odot n}$ and k_n are the canonical embeddings. To show this we just check that

$$k_{n+m}(v^n \odot a_m) = k_{n+m}v^n a_m = u^{n+m}v^n a_m = u^n a_m = k_n a_m,$$

what shows that the k_n behave correctly with respect to the inductive system $\mathcal{B}(G)^{\odot m} \rightarrow v^n \odot \mathcal{B}(G)^{\odot m} \subset \mathcal{B}(G)^{\odot(n+m)}$. Of course, $\bigcup_{n=0}^{\infty} k_n \mathcal{B}(G)^{\odot n}$ is strongly dense in $\mathcal{B}(G, H)$.

What is the induced endomorphism ϑ on $\mathcal{B}(H)$? For that goal we have to identify correctly the isomorphism $\mathcal{B}(G, H) \bar{\odot}^s \mathcal{B}(G)^{\odot 1} \rightarrow \mathcal{B}(G, H)$. By [BS00] the isomorphism is fixed as

$$(k_n a_n) \odot a_1 \longmapsto k_{n+1}(a_n \odot a_1)$$

on a strongly dense subset of $\mathcal{B}(G, H)$. In our case it is even sufficient to take $a_1 = \text{id}_G$ so that simply $a_n \odot a_1 = a_n \in \mathcal{B}(G) = \mathcal{B}(G)^{\odot(n+1)}$. So, using the definition of k_n , we find that the

elementary tensor $u_n a_n \odot \text{id}_G$ is sent to $u_{n+1} a_n = u(u^n a_n)$. This means in order to write an element $x \in \mathcal{B}(G, H)$ as $y \odot \text{id}_G \in \mathcal{B}(G, H) \bar{\odot}^s \mathcal{B}(G)^{\odot 1}$, just put $y = u^* x$. If a is in $\mathcal{B}(H) = \mathcal{B}^a(\mathcal{B}(G, H))$, then the action of $\vartheta(a)$ on $y \odot \text{id}_G$ gives by definition $ay \odot \text{id}_G$. To go back from $\mathcal{B}(G, H) \bar{\odot}^s \mathcal{B}(G)^{\odot 1}$ to $\mathcal{B}(G, H)$ we simply apply u to $ay = au^* x$. In the end ϑ is nothing but the automorphism $\vartheta(a) = uau^*$.

Of course, if v is a proper isometry, then G is infinite-dimensional, so that $\mathcal{B}(G) \cong \mathcal{B}(H)$. In other words, the automorphism ϑ of $\mathcal{B}(H)$ is conjugate to the identity automorphism on $\mathcal{B}(G)$, as it should be because the associated correspondence is the same in both cases. However, there is no canonical isomorphism to fix neither between G and H nor between $\mathcal{B}(G)$ and $\mathcal{B}(H)$.

9.1 Remark. Let $p \in \mathcal{B}(H)$ denote the projection onto $G \subset H$. Then $pu^* \upharpoonright G = v$. In fact, $u^* \in \mathcal{B}(H)$ is nothing but the Sz.-Nagy-Foias dilation of an isometry to a unitary.

The whole procedure works under slightly more general circumstances.

9.2 Proposition. *Let \mathcal{B} denote a unital C^* -algebra with a proper isometry $v \in \mathcal{B}$. Then the inductive limit over the trival product system $(\mathcal{B}^{\odot n})_{n \in \mathbb{N}_0}$ with respect to the unit $(v^{\odot n})_{n \in \mathbb{N}_0}$ has the form*

$$F := \mathcal{B} \oplus \bigoplus_{k=1}^{\infty} \mathcal{B}_0 \quad (9.2)$$

where $\mathcal{B}_0 := (\mathbf{1} - vv^*)\mathcal{B}$, and the induced endomorphism ϑ of $\mathcal{B}^a(F)$ is $\vartheta(a) = uau^*$ with u defined in the same way as in (9.1).

It is unclear, whether $F \cong \mathcal{B}$ as right module or whether $\mathcal{B} \cong \mathcal{B}^a(F)$. What we know is that $\mathcal{B} \cong \mathcal{B} \oplus \bigoplus_{k=1}^n \mathcal{B}_0$ for every n , but the simple dimension arguments that helped in the case $\mathcal{B} = \mathcal{B}(G)$ do no longer help to understand the limit $n \rightarrow \infty$.

Now let us come to Example 2.1. So we put $\mathcal{B} = \begin{pmatrix} \mathbb{C} & 0 \\ 0 & M_2 \end{pmatrix} \subset M_3$ and $E = \begin{pmatrix} 0 & \mathbb{C}_2 \\ \mathbb{C}^2 & 0 \end{pmatrix} \subset M_3$. The operations of the correspondence E over \mathcal{B} are those inherited from M_3 . This remains even true for the tensor product:

$$x \odot y = xy \in E \odot E = \mathcal{B}.$$

In particular,

$$E_n := E^{\odot n} = \begin{cases} \mathcal{B} & n \text{ even,} \\ E & n \text{ odd.} \end{cases}$$

Fortunately, the structure of Hilbert \mathcal{B} -modules F is not much more complicated than that of Hilbert spaces and we still can say in advance how automorphisms of $\mathcal{B}^a(F)$ may look like. In particular, we can say when an automorphism is associated with the correspondence E .

Let $p_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and $p_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ denote the two nontrivial central projections in \mathcal{B} . Every Hilbert \mathcal{B} -module F decomposes into the direct sum $F = F_1 \oplus F_2$ with $F_i = F p_i$. The summand F_1 has

inner product in $\begin{pmatrix} \mathbb{C} & 0 \\ 0 & 0 \end{pmatrix}$. We may identify it with a Hilbert space \mathfrak{H}_1 . The summand F_2 has inner product in $\begin{pmatrix} 0 & 0 \\ 0 & M_2 \end{pmatrix}$. Its structure is therefore that of a Hilbert M_2 -module. A short computation shows that

$$F_2 = F_2 \odot M_2 = F_2 \odot \mathbb{C}^2 \odot \mathbb{C}_2 = \mathfrak{H}_2 \odot \mathbb{C}_2 = \mathfrak{H}_2 \otimes \mathbb{C}_2,$$

where we defined the Hilbert space $\mathfrak{H}_2 := F_2 \odot \mathbb{C}^2$ and where we used in the last step that there is no difference between the interior tensor product \odot over \mathbb{C} and the exterior tensor product \otimes .

We note that F is also a W^* -module. Also most tensor products we write down in the sequel are strongly closed if they are norm closed.

An operator a on F cannot mix the components in F_1 and in F_2 . (To see this simply multiply with p_i from the right and use right linearity of a .) Therefore, a decomposes as $a = a_1 \oplus a_2$ where each a_i is an operator on F_i alone. a_1 can be any element in $\mathcal{B}(\mathfrak{H}_1)$, while a_2 must be an element in $\mathcal{B}(\mathfrak{H}_2)$ that acts on $F_2 = \mathfrak{H}_2 \otimes \mathbb{C}_2$ as $a_2 \otimes \text{id}_{\mathbb{C}_2}$. (To see the latter statement we may, for instance, observe that tensoring with \mathbb{C}_2 is an operation of Morita equivalence so that F_2 and \mathfrak{H}_2 , indeed, have the same operators.) We find $\mathcal{B}^a(F) = \mathcal{B}(\mathfrak{H}_1) \oplus \mathcal{B}(\mathfrak{H}_2)$.

It is easy to check that an automorphism of $\mathcal{B}^a(F)$ either sends $\mathcal{B}(\mathfrak{H}_i)$ onto $\mathcal{B}(\mathfrak{H}_i)$ or sends $\mathcal{B}(\mathfrak{H}_1)$ onto $\mathcal{B}(\mathfrak{H}_2)$ and *vice versa*. The first type is simply implemented by two unitaries $u_i \in \mathcal{B}(\mathfrak{H}_i)$. It is, therefore, conjugate to the identity automorphism and the associated correspondence is \mathcal{B} . In order to have the second case necessarily \mathfrak{H}_1 and \mathfrak{H}_2 are isomorphic, to a Hilbert space \mathfrak{H} say, and the action of the automorphism is exchange of the two copies of $\mathcal{B}(\mathfrak{H})$ plus, possibly, an automorphism of the first type that may be absorbed into the identifications of \mathfrak{H}_1 and \mathfrak{H}_2 with \mathfrak{H} . This second case is, thus, simply the flip $\mathcal{F}(a_1 \oplus a_2) = a_2 \oplus a_1$ on $\mathcal{B}(\mathfrak{H}) \oplus \mathcal{B}(\mathfrak{H})$ (up to conjugation with a unitary in $\mathcal{B}(\mathfrak{H}) \oplus \mathcal{B}(\mathfrak{H})$).

We claim that the correspondence associated with the flip is E . We show this by giving an isomorphism from $F \odot E$ to F that implements the flip as $a \mapsto a \odot \text{id}_E$ and appeal to the uniqueness of the correspondence inducing \mathcal{F} . Indeed, one checks easily that

$$\begin{pmatrix} h_1 \\ h_2 \otimes v^* \end{pmatrix} \odot \begin{pmatrix} 0 & v_2^* \\ v_1 & 0 \end{pmatrix} \longmapsto \begin{pmatrix} h_2 \langle v, v_1 \rangle \\ h_1 \otimes v_2^* \end{pmatrix} \quad (h_1, h_2 \in \mathfrak{H}; v, v_1, v_2 \in \mathbb{C}^2)$$

defines a surjective isometry. Moreover, choosing an arbitrary unit vector $e \in \mathbb{C}^2$ we see that $(a_1 \oplus a_2) \odot \text{id}_E$ acting on

$$\begin{pmatrix} h_1 \\ h_2 \otimes v^* \end{pmatrix} = \begin{pmatrix} h_2 \\ h_1 \otimes e^* \end{pmatrix} \odot \begin{pmatrix} 0 & v^* \\ e & 0 \end{pmatrix}$$

gives

$$\begin{pmatrix} a_1 h_2 \\ a_2 h_1 \otimes e^* \end{pmatrix} \odot \begin{pmatrix} 0 & v^* \\ e & 0 \end{pmatrix} = \begin{pmatrix} a_2 h_1 \\ a_1 h_2 \otimes v^* \end{pmatrix} = \mathcal{F}(a_1 \oplus a_2) \begin{pmatrix} h_1 \\ h_2 \otimes v^* \end{pmatrix}.$$

The discussion shows that a Hilbert \mathcal{B} -module F with an endomorphism on $\mathcal{B}^a(F)$ that has E as associated correspondence must have the form $F = \mathfrak{H} \oplus (\mathfrak{H} \otimes \mathbb{C}_2)$ and that the endomorphism is the flip \mathcal{F} on $\mathcal{B}^a(F) = \mathcal{B}(\mathfrak{H}) \oplus \mathcal{B}(\mathfrak{H})$ up to unitary equivalence in $\mathcal{B}^a(F)$. That is, the possible

endomorphisms associated with E are simply classified by the dimension of \mathfrak{H} . We ask now which of them can be obtained by the steps used in the proof of the main theorem.

E does not admit a unit vector. That is why we study this example. But E is full already in the sense of Hilbert modules, so we are in the situation of Lemma 2.2. In fact, E^2 has a unit vector ξ . Visualizing the elements of E^2 as 6×3 -matrices (on which an element $b \in \mathcal{B} \subset M_3$ acts from the left as block diagonal 6×6 -matrix $\begin{pmatrix} b & 0 \\ 0 & b \end{pmatrix}$), we may choose, for instance,

$$\xi = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \xi_1 \oplus \xi_2, \quad \text{where } \xi_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \xi_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \text{ both in } E.$$

As the cardinality in Lemma 2.2 is $l = 2$, the minimal cardinality n in Proposition 5.2 is simply countably infinite, which we denote $n = \infty$. The fact that l is finite allows to choose the bijection φ such that the unit vector $\Xi \in M_\infty(E)$ has the form

$$\Xi = \begin{pmatrix} \xi & & \\ & \xi & \\ & & \ddots \end{pmatrix}.$$

(Recall ξ stands for a 6×3 -matrix, so Ξ is not really block diagonal.)

In order to understand the inductive limit over the product system $(M_\infty(E)^{\otimes^s n})_{n \in \mathbb{N}_0}$ with respect to the unit $(\Xi^{\otimes n})_{n \in \mathbb{N}_0}$ we study first the part coming only from even n and see later on how to incorporate also the odd numbers. (As the inductive limit is increasing, it is not necessary to do it step by step. An arbitrary subsequence, like the even numbers, will be sufficient.) Recall that $M_\infty(E)^{\otimes^s 2n} = M_\infty(E^{\otimes 2n}) = M_\infty(\mathcal{B})$ (as W^* -correspondence over $M_\infty(\mathcal{B})$) and that the tensor product is just matrix multiplication (of matrices with blocks in M_3). Taking into account that

$$\xi_1^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \xi_1 \xi_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad \xi_2^2 = \xi_2 \xi_1 = 0,$$

we find

$$\Xi \odot \Xi = \begin{pmatrix} \xi_1^2 & & & & \\ 0 & & & & \\ \xi_1 \xi_2 & & & & \\ 0 & & & & \\ & \xi_1^2 & & & \\ & 0 & & & \\ & \xi_1 \xi_2 & & & \\ & 0 & & & \\ & & \ddots & & \end{pmatrix}$$

(all blocks in $\mathcal{B} \subset M_3$). We are in the situation of Proposition 9.2 and identifying the right-submodule (that is, the right ideal) $(\mathbf{1} - (\Xi \odot \Xi)(\Xi \odot \Xi)^*)M_\infty(\mathcal{B})$ in $M_\infty(\mathcal{B})$, we could give the result rather explicitly. (In fact, it is similar to the simpler case $\mathcal{B}(H)$, that is, we obtain just a unitary perturbation of the identity endomorphism of $M_\infty(\mathcal{B})$.) But, fortunately, this (tedious) identification is not necessary. For us it is just sufficient to know that the inductive limit contains at least one copy of $M_\infty(\mathcal{B})$, and Equation (9.2) tells us that exactly this is true. (Note that this is independent of the concrete choice of Ξ .)

Indeed, from the inductive limit over $M_\infty(\mathcal{B})$ we obtain the \mathcal{B} -module, in which we are actually interested, by Morita equivalence, that is, by tensoring with $\overline{\mathcal{B}^\infty}^s$ from the right. (See the proof of Theorem 4.10 and the discussion leading to Corollary 5.1.) The fact that the inductive limit contains $M_\infty(\mathcal{B})$ as a direct summand means that the \mathcal{B} -module F we seek contains $M_\infty(\mathcal{B}) \otimes^s \overline{\mathcal{B}^\infty}^s = \overline{\mathcal{B}^\infty}^s$ as a direct summand. This means, however the inductive limit looks like, the module F must be determined by a Hilbert space \mathfrak{H} that is not finite-dimensional. On the other hand, the inductive limit is obtained by a ‘‘countable’’ procedure. Therefore, \mathfrak{H} must be separable. This determines F up to isomorphism.

What we have so far is the \mathcal{B} -module F and the endomorphism ϑ^2 that acts up to unitary equivalence as the identity endomorphism of $\mathcal{B}^a(F)$. What remains is to find ϑ itself, that is, some square root of $\vartheta^2 = \text{id}_{\mathcal{B}^a(F)}$ that essentially (that is, up to unitary equivalence in $\mathcal{B}^a(F)$) flips the two components $\mathcal{B}(\mathfrak{H})$ contained in $\mathcal{B}^a(F)$. We know already that the freedom that remains in addition to the flip is a unitary that may be absorbed into the way how we identify F with $\mathfrak{H} \oplus (\mathfrak{H} \otimes \mathbb{C}_2)$.

We summarize.

9.3 Proposition. *Let F and ϑ be the Hilbert \mathcal{B} -module and ϑ the endomorphism of $\mathcal{B}^a(F)$, respectively, such that the correspondence associated with ϑ is E as constructed in the proof of the main theorem with minimal choices of cardinalities. Then F can be identified with $\mathfrak{H} \oplus (\mathfrak{H} \otimes \mathbb{C}_2)$ in such a way that ϑ is the flip ϖ on $\mathcal{B}^a(F) = \mathcal{B}(\mathfrak{H}) \oplus \mathcal{B}(\mathfrak{H})$, where \mathfrak{H} is an infinite-dimensional separable Hilbert space. Every other Hilbert space doing the job of \mathfrak{H} must be isomorphic to \mathfrak{H} .*

If we do not choose minimal cardinality (in Proposition 5.2), then \mathfrak{H} still has unique dimension, namely, the cardinality \mathfrak{n} .

If we vary, instead, the dimension of \mathfrak{H} in $F := \mathfrak{H} \oplus (\mathfrak{H} \otimes \mathbb{C}_2)$ arbitrarily and look at the flip, then we obtain all endomorphisms (up to unitary equivalence in $\mathcal{B}^a(F)$) that have E as associated correspondence. In particular, our construction via Morita equivalence and inductive limit does not give all endomorphisms that have E as associated correspondence, but only those where \mathfrak{H} is infinite-dimensional.

9.4 Remark. The reader might object that the correspondence E we discuss is a Morita equivalence and, consequently, we are speaking about endomorphisms that are automorphisms. In fact, we do consider it as an interesting problem to deal with general unital endomorphisms of $\mathcal{B}^a(F)$, that is, of $\mathcal{B}(\mathfrak{H}) \oplus \mathcal{B}(\mathfrak{H})$ or, more generally, of $\mathcal{B}(\mathfrak{H}_1) \oplus \mathcal{B}(\mathfrak{H}_2)$. The general form of such endomorphisms is still comparably accessible and it should be easy to see how the associated correspondences look like. However, as our example shows, it will be quite space consuming to work out which of them can be obtained by our inductive limit construction. (We remind the reader of the fact that we gave explicit identifications of the inductive limit construction only in the simpler case $\mathcal{B}(G)$, while the identification of the case E remains only an identification

up to unitary equivalence. (An explicit identification would fill several pages with large block matrices containing lots of zeros.) Additionally, most correspondences associated with a proper endomorphism will have enough “space” (that is, enough “multiplicity”) to admit unit vectors so that the example will be not typical to illustrate the methods of these notes. See, however, Example 9.5.

We briefly discuss the relation with Section 7. \mathcal{B} is a von Neumann subalgebra of $M_3 = \mathcal{B}(\mathbb{C}^3)$ and $H = E \odot \mathbb{C}^3$ is simply \mathbb{C}^3 with the identification by matrix multiplication, that is,

$$\begin{pmatrix} 0 & v_2^* \\ v_1 & 0 \end{pmatrix} \odot \begin{pmatrix} z \\ v \end{pmatrix} = \begin{pmatrix} \langle v_2, v \rangle \\ v_1 z \end{pmatrix} \quad (v, v_1, v_2 \in \mathbb{C}^2; z \in \mathbb{C}).$$

In this identification the representation ρ of \mathcal{B} on $H = \mathbb{C}^3$ is simply the identity representation.

The commutant of \mathcal{B} in M_3 is $\mathcal{B}' = \begin{pmatrix} \mathbb{C} & 0 \\ 0 & \mathbb{C}\mathbf{1} \end{pmatrix}$. The representation ρ' of \mathcal{B}' on H is defined as

$$\begin{aligned} \rho' \left(\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \mathbf{1} \end{pmatrix} \right) \begin{pmatrix} \langle v_2, v \rangle \\ v_1 z \end{pmatrix} &= \rho' \left(\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \mathbf{1} \end{pmatrix} \right) \left(\begin{pmatrix} 0 & v_2^* \\ v_1 & 0 \end{pmatrix} \odot \begin{pmatrix} z \\ v \end{pmatrix} \right) \\ &:= \begin{pmatrix} 0 & v_2^* \\ v_1 & 0 \end{pmatrix} \odot \left(\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \mathbf{1} \end{pmatrix} \begin{pmatrix} z \\ v \end{pmatrix} \right) = \begin{pmatrix} 0 & v_2^* \\ v_1 & 0 \end{pmatrix} \odot \begin{pmatrix} z_1 z \\ z_2 v \end{pmatrix} = \begin{pmatrix} z_2 \langle v_2, v \rangle \\ z_1 v_1 z \end{pmatrix} = \begin{pmatrix} z_2 & 0 \\ 0 & z_1 \mathbf{1} \end{pmatrix} \begin{pmatrix} \langle v_2, v \rangle \\ v_1 z \end{pmatrix}. \end{aligned}$$

So ρ' just flips the coefficients z_1 and z_2 of an element of \mathcal{B}' .

The commutant of E is $E' := C_{\mathcal{B}}(\mathcal{B}(\mathbb{C}^3, H))$. As $H = \mathbb{C}^3$ and the action of \mathcal{B} on H is simply the identity representation, we find $E' = C_{\mathcal{B}}(M_3) = \mathcal{B}'$ as right \mathcal{B}' -module. However, left action of \mathcal{B}' on E' is via the flip. E' has an obvious representation (ψ', η') on \mathbb{C}^2 , namely, $\psi' \left(\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \mathbf{1} \end{pmatrix} \right) = \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \in M_2 = \mathcal{B}(\mathbb{C}^2)$ and $\eta' \left(\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \mathbf{1} \end{pmatrix} \right) = \begin{pmatrix} 0 & z_2 \\ z_1 & 0 \end{pmatrix}$. All other representations are multiples of that representation. It is not difficult to check that the multiplicity space is exactly \mathfrak{H} , when \mathfrak{H} is the Hilbert space that characterizes the endomorphism ϑ related to (ψ', η') by Theorem 7.4.

Of course, it is also possible to switch the roles played by E and by E' , that is, to investigate representations of E and endomorphisms of $\mathcal{B}^a(F')$ (where F' is a Hilbert \mathcal{B}' -module) that have E' as associated correspondence. Also this is not really a typical example because \mathcal{B}' has a unit vector and this unit vector even must be unitary. (This does not mean that we are in the situation of Proposition 9.2. In fact, E' is \mathcal{B}' only as a right module. The left action makes it different from the trivial \mathcal{B}' -correspondence.) The discussion is similar to the discussion for E but simpler. Also here if an endomorphism ϑ' of $\mathcal{B}^a(F')$ has E' as associated correspondence, then $\mathcal{B}^a(F') = \mathcal{B}(\mathfrak{H}) \oplus \mathcal{B}(\mathfrak{H})$ and ϑ' is (conjugate to) the flip. Additionally, if ϑ' comes from the inductive limit construction, then $\mathfrak{H} = \mathbb{C}$, because the unit vector is unitary. The representation of E (as the commutant of E') is simply the identity representation on \mathbb{C}^3 (as a subset of $M_3 = \mathcal{B}(\mathbb{C}^3)$).

9.5 Example. Without running through the explicit construction in the proof of the main theorem, we give an example of a correspondence without unit vector, that comes from a true endomorphism. Moreover, no tensor power of this correspondence admits a unit vector.

We consider the von Neumann algebra $\mathcal{B} = \overline{\bigoplus_{n \in \mathbb{N}} M_n}^s$ acting on $G = \bigoplus_{n \in \mathbb{N}} \mathbb{C}^n$. Recall that $M_{n,m} = \mathbb{C}^n \otimes \mathbb{C}_m$ is a von Neumann correspondence from M_n to M_m (actually, a Morita equivalence) that may also be considered as a correspondence over \mathcal{B} . As E we choose the von Neumann \mathcal{B} -correspondence direct sum

$$E := \mathbb{C} \oplus \overline{\bigoplus_{n \in \mathbb{N}} \mathbb{C}^n \otimes \mathbb{C}_{n+1}}^s.$$

Here \mathcal{B} acts on direct summands of E from either side with that direct summand M_n that fits the correct dimension. That is, M_1 acts from the left on the summands \mathbb{C} and $\mathbb{C}^1 \otimes \mathbb{C}_2 = \mathbb{C}_2$ but from the right only on \mathbb{C} . It is easy to check that

$$E^{\otimes^s m} := \mathbb{C} \oplus \mathbb{C}_2 \oplus \dots \oplus \mathbb{C}_n \oplus \overline{\bigoplus_{n \in \mathbb{N}} \mathbb{C}^n \otimes \mathbb{C}_{n+m}}^s.$$

All $E^{\otimes^s m}$ are strongly full but none of them has a unit vector.

E is not a Morita equivalence, so it must come from a true endomorphism. To understand which endomorphisms could be associated with E , we analyze the general structure of a von Neumann \mathcal{B} -module F and look for which (strongly full) F we can write down an isomorphism $F \otimes^s E = F$. According to the minimal ideals M_n in \mathcal{B} , also F decomposes into a direct sum of von Neumann M_n -modules F_n . Every F_n must have the form $\mathfrak{H}_n \otimes \mathbb{C}_n$ for some Hilbert module. Of course, $\mathcal{B}^a(F) = \overline{\bigoplus_{n \in \mathbb{N}} \mathcal{B}(\mathfrak{H}_n)}^s$. A short computation yields that

$$F \otimes^s E := \mathfrak{H}_1 \oplus \overline{\bigoplus_{n \in \mathbb{N}} \mathfrak{H}_n \otimes \mathbb{C}_{n+1}}^s.$$

Therefore, $F \otimes^s E \cong F$, if and only if $\mathfrak{H}_n = \mathfrak{H}$ for all $n \in \mathbb{N}$. Another computation shows that the endomorphism induced by this isomorphism acts on $\mathcal{B}^a(F)$ as $\vartheta(a_1, a_2, \dots) = (a_1, a_1, a_2, \dots)$. It is nothing but the unitization of the one-sided shift on $\mathcal{B}^a(F)$. As our construction of the inductive limit runs through a countable inductive system of proper isometries, \mathfrak{H} coming from our construction must be infinite-dimensional and separable. Note that in this case F has a unit vector, while if \mathfrak{H} is finite-dimensional, then F fails to have a unit vector.

References

- [Arv89] W. Arveson, *Continuous analogues of Fock space III: Singular states*, J. Operator Theory **22** (1989), 165–205.
- [BBLs04] S.D. Barreto, B.V.R. Bhat, V. Liebscher, and M. Skeide, *Type I product systems of Hilbert modules*, J. Funct. Anal. **212** (2004), 121–181.
- [Bha96] B.V.R. Bhat, *An index theory for quantum dynamical semigroups*, Trans. Amer. Math. Soc. **348** (1996), 561–583.

- [BLS06] B.V.R. Bhat, V. Liebscher, and M. Skeide, (*Tentative title*) *A problem of Powers and the product of spatial product systems*, Preprint, Campobasso, in preparation, 2006.
- [BS00] B.V.R. Bhat and M. Skeide, *Tensor product systems of Hilbert modules and dilations of completely positive semigroups*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **3** (2000), 519–575.
- [Con80] A. Connes, *Correspondences*, His hand-written notes, unpublished, 1980.
- [GS99] D. Goswami and K.B. Sinha, *Hilbert modules and stochastic dilation of a quantum dynamical semigroup on a von Neumann algebra*, *Commun. Math. Phys.* **205** (1999), 377–403.
- [Hir05a] I. Hirshberg, *Essential representations of C^* -correspondences*, *Int. J. Math.* **16** (2005), 765–775.
- [Hir05b] ———, *On the universal property of Pimsner-Toeplitz C^* -algebras and their continuous analogues*, *J. Funct. Anal.* **219** (2005), 21–33.
- [Lie03] V. Liebscher, *Random sets and invariants for (type II) continuous tensor product systems of Hilbert spaces*, Preprint, ArXiv: math.PR/0306365, 2003.
- [MM04] A.E. Marrero and P.S. Muhly, *Cuntz-Pimsner algebras, completely positive maps and Morita equivalence*, Preprint, ArXiv: math.OA/0409511, 2004.
- [Mor58] K. Morita, *Duality for modules and its applications to the theory of rings with minimum condition*, *Tokyo Kyoiku Daigaku* **6** (1958), 83–142.
- [MS98] P.S. Muhly and B. Solel, *Tensor algebras over C^* -correspondences: representations, dilations, and C^* -envelopes*, *J. Funct. Anal.* **158** (1998), 389–457.
- [MS99] ———, *Tensor algebras, induced representations and the Wold decomposition*, *Canad. J. Math.* **51** (1999), 850–880.
- [MS00] ———, *On the Morita equivalence of tensor algebras*, *Proc. London Math. Soc.* **81** (2000), 113–168.
- [MS02] ———, *Quantum Markov processes (correspondences and dilations)*, *Int. J. Math.* **51** (2002), 863–906.
- [MS04] ———, *Hardy algebras, W^* -correspondences and interpolation theory*, *Math. Ann.* **330** (2004), 353–415.
- [MS05] ———, *Duality of W^* -correspondences and applications*, *QP-PQ: Quantum Probability and White Noise Analysis XVIII* (M. Schürmann and U. Franz, eds.), World Scientific, 2005, pp. 396–414.
- [MSS04] P.S. Muhly, M. Skeide, and B. Solel, *Representations of $\mathcal{B}^a(E)$* , Preprint, ArXiv: math.OA/0410607, 2004, To appear in *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*
- [Pas73] W.L. Paschke, *Inner product modules over B^* -algebras*, *Trans. Amer. Math. Soc.* **182** (1973), 443–468.

- [Rie74a] M.A. Rieffel, *Induced representations of C^* -algebras*, Adv. Math. **13** (1974), 176–257.
- [Rie74b] ———, *Morita equivalence for C^* -algebras and W^* -algebras*, J. Pure Appl. Algebra **5** (1974), 51–96.
- [Ske00a] M. Skeide, *Generalized matrix C^* -algebras and representations of Hilbert modules*, Mathematical Proceedings of the Royal Irish Academy **100A** (2000), 11–38.
- [Ske00b] ———, *Quantum stochastic calculus on full Fock modules*, J. Funct. Anal. **173** (2000), 401–452.
- [Ske01a] ———, *Hilbert modules and applications in quantum probability*, Habilitationsschrift, Cottbus, 2001, Available at http://www.math.tu-cottbus.de/INSTITUT/lswas/_skeide.html.
- [Ske01b] ———, *The index of (white) noises and their product systems*, Preprint, Rome, 2001, To appear in Infin. Dimens. Anal. Quantum Probab. Relat. Top.
- [Ske02] ———, *Dilations, product systems and weak dilations*, Math. Notes **71** (2002), 914–923.
- [Ske03a] ———, *Commutants of von Neumann modules, representations of $\mathcal{B}^a(E)$ and other topics related to product systems of Hilbert modules*, Advances in quantum dynamics (G.L. Price, B. M. Baker, P.E.T. Jorgensen, and P.S. Muhly, eds.), Contemporary Mathematics, no. 335, American Mathematical Society, 2003, pp. 253–262.
- [Ske03b] ———, *Dilation theory and continuous tensor product systems of Hilbert modules*, QP-PQ: Quantum Probability and White Noise Analysis XV (W. Freudenberg, ed.), World Scientific, 2003.
- [Ske05a] ———, *Commutants of von Neumann Correspondences and Duality of Eilenberg-Watts Theorems by Rieffel and by Blecher*, Preprint, ArXiv: math.OA/0502241, 2005, To appear in Proceedings of “25th QP-Conference Quantum Probability and Applications”, Bedlewo 2004.
- [Ske05b] ———, *Three ways to representations of $\mathcal{B}^a(E)$* , QP-PQ: Quantum Probability and White Noise Analysis XVIII (M. Schürmann and U. Franz, eds.), World Scientific, 2005, pp. 504–517.
- [Ske05c] ———, *Von Neumann modules, intertwiners and self-duality*, J. Operator Theory **54** (2005), 119–124.
- [Ske06] ———, *Intertwiners, dual quasi orthonormal bases and representations*, in preparation, 2006.
- [Tsi00] B. Tsirelson, *From random sets to continuous tensor products: answers to three questions of W. Arveson*, Preprint, ArXiv: math.FA/0001070, 2000.