

# Holomorphic Functions of Slow Growth on Coverings of Pseudoconvex Domains in Stein Manifolds

Alexander Brudnyi\*

Department of Mathematics and Statistics  
University of Calgary, Calgary  
Canada

## Abstract

We apply the methods developed in [Br1] to study holomorphic functions of slow growth on coverings of pseudoconvex domains in Stein manifolds. In particular, we extend and strengthen certain results of Gromov, Henkin and Shubin [GHS] on holomorphic  $L^2$  functions on coverings of pseudoconvex manifolds in the case of coverings of Stein manifolds.

## 1. Introduction.

Let  $M$  be a complex manifold satisfying

$$M \subset\subset \widetilde{M} \subset N \quad \text{and} \quad \pi_1(M) = \pi_1(N) \quad (1.1)$$

where  $M$  and  $\widetilde{M}$  are open connected subsets of a complex manifold  $N$  and  $\widetilde{M}$  is Stein. (Here  $\pi_1(X)$  stands for the fundamental group of  $X$ .) Condition (1.1) is valid, e.g., for  $M$  a strictly pseudoconvex domain or an analytic polyhedra in a Stein manifold. It implies that the group  $\pi_1(N)$  is finitely generated. In [Br1] we presented a method to construct integral representation formulas for holomorphic functions of slow growth defined on unbranched coverings of  $M$ . Using such formulas we established that some known results for holomorphic functions on  $M$  can be extended to similar results for holomorphic functions of slow growth on coverings of  $M$ . In this paper we continue to study holomorphic functions of slow growth on coverings of  $M$  and apply the methods developed in [Br1] to extend and strengthen certain results of Gromov, Henkin and Shubin [GHS] on holomorphic  $L^2$  functions on coverings of pseudoconvex manifolds in the case of coverings of Stein manifolds.

---

\*Research supported in part by NSERC.

2000 *Mathematics Subject Classification*. Primary 46E15. Secondary 32T15, 32T40.

*Key words and phrases*. Covering, peak point, strictly pseudoconvex sets, Banach vector bundle.

The presentation in this paper is focused on several problems and results formulated in [GHS]. To describe them we, first, recall some definitions.

Let  $M \subset\subset N$  be a domain with a smooth boundary  $bM$  in an  $n$ -dimensional complex manifold  $N$ , that is,

$$M = \{z \in N : \rho(z) < 0\} \quad (1.2)$$

where  $\rho$  is a real-valued function of class  $C^2(\Omega)$  in a neighbourhood  $\Omega$  of the compact set  $\overline{M} := M \cup bM$  such that

$$d\rho(z) \neq 0 \quad \text{for all } z \in bM. \quad (1.3)$$

Let  $z_1, \dots, z_n$  be complex local coordinates in  $N$  near  $z \in bM$ . Then the tangent space  $T_z N$  at  $z$  is identified with  $\mathbb{C}^n$ . By  $T_z^c(bM) \subset T_z N$  we denote the complex tangent space to  $bM$  at  $z$ , i.e.,

$$T_z^c(bM) = \{w = (w_1, \dots, w_n) \in T_z(N) : \sum_{j=1}^n \frac{\partial \rho}{\partial z_j}(z) w_j = 0\}. \quad (1.4)$$

The *Levi form* of  $\rho$  at  $z \in bM$  is a hermitian form on  $T_z^c(bM)$  defined in the local coordinates by the formula

$$L_z(w, \overline{w}) = \sum_{j,k=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \overline{z}_k}(z) w_j \overline{w}_k. \quad (1.5)$$

The manifold  $M$  is called *pseudoconvex* if  $L_z(w, \overline{w}) \geq 0$  for all  $z \in bM$  and  $w \in T_z^c(bM)$ . It is called *strictly pseudoconvex* if  $L_z(w, \overline{w}) > 0$  for all  $z \in bM$  and all  $w \neq 0$ ,  $w \in T_z^c(bM)$ .

Equivalently, strictly pseudoconvex manifolds can be described as the ones which locally, in a neighbourhood of any boundary point, can be presented as strictly convex domains in  $\mathbb{C}^n$ . It is also known (see [C], [R]) that any strictly pseudoconvex manifold admits a proper holomorphic map with connected fibres onto a normal Stein space.

Without loss of generality we may and will assume that  $\pi_1(M) = \pi_1(N)$  for  $M$  as above. Let  $r : N_G \rightarrow N$  be the regular covering of  $N$  with (discrete) transformation group  $G$ . Then  $M_G := r^{-1}(M)$  is a regular covering of  $M$  (with the same transformation group). It is a domain in  $N_G$  with the smooth boundary  $bM_G := r^{-1}(bM)$ . By  $\overline{M}_G := M_G \cup bM_G$  we denote the closure of  $M_G$  in  $N_G$ .

Let  $X$  be a subspace of the space  $\mathcal{O}(M_G)$  of all holomorphic functions on  $M_G$ . A point  $z \in bM_G$  is called a *peak point* for  $X$  if there exists a function  $f \in X$  such that  $f$  is unbounded on  $M_G$  but bounded outside  $U \cap M_G$  for any neighbourhood  $U$  of  $z$  in  $N_G$ .

A point  $z \in bM_G$  is called a *local peak point* for  $X$  if there exists a function  $f \in X$  such that  $f$  is unbounded in  $U \cap M_G$  for any neighbourhood  $U$  of  $z$  in  $N_G$  and there exists a neighbourhood  $U$  of  $z$  in  $N_G$  such that for any neighbourhood  $V$  of  $z$  in  $N_G$  the function  $f$  is bounded on  $U \setminus V$ .

The Oka-Grauert theorem [G1] states that if  $M$  is strictly pseudoconvex and  $bM$  is not empty then every  $z \in bM$  is a peak point for  $\mathcal{O}(M)$ . In general it is not known whether the similar statement is true for boundary points of  $M_G$  with an infinite  $G$ .

Let  $dV_{M_G}$  be the Riemannian volume form on  $M_G$  obtained by a Riemannian metric pulled back from  $N$ . By  $H^2(M_G)$  we denote the Hilbert space of holomorphic functions  $g$  on  $M_G$  with norm

$$\left( \int_{z \in M_G} |g(z)|^2 dV_{M_G}(z) \right)^{1/2}.$$

In [GHS] the von Neumann  $G$ -dimension  $\dim_G$  was used to measure the space  $H^2(M_G)$ . In particular, in [GHS, Theorem 0.2] the following result was proved.

**Theorem A** *If  $M$  is strictly pseudoconvex, then*

- (a)  $\dim_G H^2(M_G) = \infty$  and
- (b) *each point in  $bM_G$  is a local peak point for  $H^2(M_G)$ .*

In [GHS, Theorem 0.5] the similar result was established for a covering  $M_G$  of a pseudoconvex manifold  $M$  with a strictly plurisubharmonic  $G$ -invariant function existing in a neighbourhood of  $bM_G$ . Finally, in [GHS, section 4] the following open problems were formulated.

Suppose that  $M$  is strictly pseudoconvex.

1. Does there exist a finite number of functions in  $H^2(M_G) \cap C(\overline{M}_G)$  which separate all points in  $bM_G$ ?
2. Assume that  $\dim_{\mathbb{C}} M = 2$ . Does there exist  $f \in H^2(M_G) \cap C(\overline{M}_G)$  such that  $f(x) \neq 0$  for all  $x \in bM_G$ ?
3. Is it true that for every  $CR$ -function  $f \in L^2(bM_G) \cap C(bM_G)$  in case  $\dim_{\mathbb{C}} M_G > 1$  there exists  $f' \in H^2(M_G) \cap C(\overline{M}_G)$  such that  $f'|_{bM_G} = f$ ?

Here  $L^2(bM_G)$  is defined similar to  $H^2(M_G)$  with respect to the volume form on  $bM_G$  obtained by a Riemannian metric pulled back from  $N$ . Also, recall that  $f \in C(bM_G)$  is called a  $CR$ -function if for every smooth  $(n, n-2)$ -form  $\omega$  with a compact support one has

$$\int_{bM_G} f \wedge \bar{\partial}\omega = 0.$$

If  $f$  is smooth this is equivalent to the fact that  $f$  is a solution of the tangential  $CR$ -equations:  $\bar{\partial}_b f = 0$  (see, e.g., [KR]).

The present paper deals with the above results and problems in the case of coverings of  $M$  satisfying condition (1.1).

## 2. Formulation of Main Results.

**2.1.** We start with some results related to Problem 2 of the Introduction.

Let  $M$  be a manifold satisfying condition (1.1) and  $M'$  be an unbranched covering of  $M$ . Condition (1.1) implies that there is a covering  $r : N' \rightarrow N$  of  $N$  such that  $M'$  is a domain in  $N'$  (i.e.,  $\pi_1(M') = \pi_1(N')$ ). As above  $\overline{M'}$  denotes the closure of  $M'$  in  $N'$ .

Let  $\phi : N' \rightarrow \mathbb{R}$  be a function uniformly continuous with respect to the path metric induced by a Riemannian metric pulled back from  $N$ .

**Theorem 2.1** *There exist a function  $f_\phi \in \mathcal{O}(M') \cap C(\overline{M'})$  and a constant  $C = C(\phi, M', N')$ <sup>1</sup> such that*

$$|f_\phi(z) - \phi(z)| < C \quad \text{and} \quad |df_\phi(z)| < C \quad \text{for all } z \in M' .$$

(Here the norm  $|\omega(z)|$  of a differential form  $\omega$  at  $z \in M'$  is determined with respect to the Riemannian metric pulled back from  $N$ .)

As a corollary of this result we answer an extended version of Question 2 of the Introduction for coverings of manifolds  $M$  satisfying condition (1.1). Namely, let  $d$  be the path metric on  $N'$  obtained by the pullback of a Riemannian metric defined on  $N$ . Fix a point  $o \in M'$  and set

$$d_o(x) := d(o, x) , \quad x \in N' .$$

From the triangle inequality it follows that the function  $\phi(x) := d_o(x)$ ,  $x \in N'$ , satisfies the hypothesis of Theorem 2.1.

**Corollary 2.2** *Let  $f := f_\phi$  be the function from Theorem 2.1 for  $\phi = d_o$ . Then there exists a constant  $\alpha > 0$  such that  $F = e^{-\alpha f} \in H^2(M') \cap C(\overline{M'})$ .*

(Note that here  $F(x) \neq 0$  for all  $x \in M'$  and there are no restrictions on  $\dim_{\mathbb{C}} M$ .)

**2.2.** In this part we formulate our results related to Theorem A of the Introduction.

Let  $M'$  be an unbranched covering of  $M$  satisfying (1.1). Let  $\psi : M' \rightarrow \mathbb{R}_+$  be a continuous function and  $dV_{M'}$  be the Riemannian volume form on  $M'$  obtained by a Riemannian metric pulled back from  $N$ . For an open set  $D \subset M$  we introduce the Banach space  $H_\psi^p(D')$  of holomorphic functions  $g$  on  $D' := r^{-1}(D) \subset M'$  with norm

$$\left( \int_{z \in M'} |g(z)|^p \psi(z) dV_{M'}(z) \right)^{1/p} .$$

Let  $r : N' \rightarrow N$  be the covering of  $N$  satisfying (1.1) such that  $\pi_1(N') = \pi_1(M')$ . Then  $M' (= r^{-1}(M))$  is a domain in  $N'$ . Suppose that  $\psi : N' \rightarrow \mathbb{R}_+$  is such that  $\log \psi$  is uniformly continuous with respect to the path metric induced by a Riemannian metric pulled back from  $N$ . We set  $\phi := \log \psi$  and consider the holomorphic function  $f_\phi$  from Theorem 2.1. This theorem implies that for  $\tilde{C} := e^C$

$$\frac{1}{\tilde{C}} \psi(z) \leq |e^{f_\phi(z)}| \leq \tilde{C} \psi(z) , \quad z \in \overline{M'} . \quad (2.1)$$

Therefore the following result holds.

**Proposition 2.3** *For any open set  $D \subset M$  and every  $p \in [1, \infty)$  the map  $L_\psi : H_\psi^p(D') \rightarrow H_1^p(D')$ ,  $L_\psi(g) = g \cdot e^{f_\phi/p}$ , is an isomorphism of Banach spaces.  $\square$*

<sup>1</sup>Here and below the notation  $C = C(\alpha, \beta, \gamma, \dots)$  means that the constant depends only on the parameters  $\alpha, \beta, \gamma, \dots$ .

Let us now formulate an extension of Theorem A of the Introduction.

Suppose that  $M$  is a strictly pseudoconvex domain in a complex manifold  $N$  such that  $\pi_1(M) = \pi_1(N)$  and  $N$  is a domain in a Stein manifold. Let  $r : N' \rightarrow N$  be an unbranched covering of  $N$  and  $M' = r^{-1}(M)$  be the corresponding covering of  $M$ . Let  $bM' = r^{-1}(bM)$  be the boundary of  $M'$  in  $N'$ .

**Theorem 2.4** *Each point in  $bM'$  is a peak point for  $\mathcal{O}(M')$  and for every  $H_1^p(M')$ ,  $1 \leq p < \infty$ .*

From Theorem 2.4 and Proposition 2.3 we get (for  $\psi$  as in Proposition 2.3)

**Corollary 2.5** *Each  $z \in bM'$  is a peak point for  $H_\psi^p(M')$ ,  $1 \leq p < \infty$ .  $\square$*

**Remark 2.6** The main ingredient of the proof of Theorem 2.4 is uniform estimates for solutions of certain  $\bar{\partial}$ -equations on  $M'$ . In fact similar estimates are valid on coverings of, so-called, non-degenerate pseudoconvex polyhedrons on Stein manifolds (see [SH] and [He] for their definition). This class contains, in particular, piecewise strictly pseudoconvex domains and non-degenerate analytic polyhedrons on Stein manifolds. Also, every  $M$  from this class satisfies condition (1.1). Let  $M'$  be a covering of such  $M$  and  $z \in bM'$  be such that  $M' \cap U$  is strictly pseudoconvex for a neighbourhood  $U \subset N'$  of  $z$ . Then arguing as in the proof of Theorem 2.4 one obtains that  $z$  is a peak point for  $\mathcal{O}(M')$  and for every  $H_1^p(M')$ ,  $1 \leq p < \infty$ .

**2.3.** In this section we discuss some results related to Problem 3 of the Introduction.

Let  $r : N' \rightarrow N$  be a covering of  $N$  satisfying (1.1). As before we set  $M' = r^{-1}(M) \subset N'$ . Consider a function  $\psi : N' \rightarrow \mathbb{R}_+$  such that  $\log \psi$  is uniformly continuous with respect to the path metric induced by a Riemannian metric pulled back from  $N$ . For such  $\psi$  and every  $x \in M$  we introduce the Banach space  $l_{p,\psi,x}(M')$ ,  $1 \leq p \leq \infty$ , of functions  $g$  on  $r^{-1}(x) \subset M'$  with norm

$$|g|_{p,\psi,x} := \left( \sum_{y \in r^{-1}(x)} |g(y)|^p \psi(y) \right)^{1/p}. \quad (2.2)$$

Next, for an open set  $D \subset M$  we introduce the Banach space  $\mathcal{H}_{p,\psi}(D')$ ,  $1 \leq p \leq \infty$ , of functions  $f$  holomorphic on  $D' := r^{-1}(D) \subset M'$  with norm

$$|f|_{p,\psi}^D := \sup_{x \in D} |f|_{p,\psi,x}. \quad (2.3)$$

Clearly, one has a continuous embedding  $\mathcal{H}_{p,\psi}(D') \hookrightarrow H_\psi^p(D')$ . Let  $U \subset N$  be an open set containing  $\bar{D}$  and  $U' = r^{-1}(U)$ . Then for  $\psi$  as above using the mean value property for plurisubharmonic functions one can easily show that for each  $p \in [1, \infty]$  the restriction  $f \mapsto f|_{D'}$  induces a linear continuous map  $H_\psi^p(U') \rightarrow \mathcal{H}_{p,\psi}(D')$ . Also, for such  $\psi$  from the results proved in [Br1] follow that holomorphic functions from  $\mathcal{H}_{p,\psi}(M')$  separate all points in  $M'$  (for each  $p \in [1, \infty]$ ).

Let us formulate the main result of this section.

Let  $M \subset\subset \tilde{M} \subset N$  be manifolds satisfying condition (1.1) with  $\dim_{\mathbb{C}} M \geq 2$ . Suppose that  $D \subset\subset M$  is an open subset whose boundary  $bD$  is a connected  $C^k$  submanifold of  $M$  ( $1 \leq k \leq \infty$ ). For a covering  $r : N' \rightarrow N$  we set  $D' = r^{-1}(D)$  and  $bD' = r^{-1}(bD)$ .

**Theorem 2.7** For every CR-function  $f \in C^s(bD')$ ,  $0 \leq s \leq k$ , satisfying

$$f|_{r^{-1}(x)} \in l_{p,\psi,x}(M') \quad \text{for all } x \in D \quad \text{and} \quad \sup_{x \in bD} |f|_{p,\psi,x} < \infty$$

there exists a function  $f' \in \mathcal{H}_{p,\psi}(D') \cap C^s(\overline{D'})$  such that  $f'|_{bD'} = f$ .

**Remark 2.8** (1) The converse to this theorem is always true: the restriction of every  $f' \in \mathcal{H}_{p,\psi}(D') \cap C^s(\overline{D'})$  to  $bD'$  is a CR-function satisfying the hypotheses of the theorem.

(2) We will prove also (see (6.6)) that for some  $c = c(M', M, \psi, p)$

$$|f'|_{p,\psi}^D \leq c \sup_{x \in bD} |f|_{p,\psi,x} .$$

As a corollary we obtain an analog of the Hartogs extension theorem. We formulate it for functions of the maximal possible growth for which our method works.

Suppose that  $M$  satisfies (1.1). Let  $D \subset M$  be a domain and  $K \subset\subset D$  be a compact set such that  $U := D \setminus K$  is connected. Consider a covering  $r : M' \rightarrow M$  and set  $D' = r^{-1}(D)$ ,  $U' = r^{-1}(U)$ . By  $d_o$ ,  $o \in M'$ , we denote the distance function on  $M'$  as in Corollary 2.2.

**Corollary 2.9** There exists a constant  $c > 0$  such that for every  $f \in \mathcal{O}(U')$  satisfying for some  $c_2 > 0$  and  $0 < c_1 < c$  the inequality

$$|f(z)| \leq e^{c_2 e^{c_1 d_o(z)}} , \quad z \in U' ,$$

there is  $f' \in \mathcal{O}(D')$  such that

$$|f'(z)| \leq e^{c_3 e^{c_1 d_o(z)}} , \quad z \in D' , \quad \text{and} \quad f'|_{U'} = f ;$$

where  $c_3$  depends on  $c_2, c_1, c, M, M'$  only.

We don't know whether a similar extension result holds for functions  $f$  growing faster than those of the corollary.

**2.4.** Finally, we formulate a result related to Problem 1 of the Introduction. First, we recall some definitions of the theory of flat vector bundles (see, e.g., [O]).

Let  $X$  be a complex manifold and  $\rho : \pi_1(X) \rightarrow GL_k(\mathbb{C})$  be a homomorphism of its fundamental group. We set  $G := \pi_1(X)/\text{Ker } \rho$ . It is well known (see, e.g., Example 3.2 (b) below) that to any such  $\rho$  corresponds a complex flat vector bundle  $E_\rho$  on  $X$  (i.e., a bundle constructed by a locally constant cocycle). We call  $E_\rho$  the *bundle associated with*  $\rho$ . Assume that  $\rho$  is such that  $E_\rho$  is topologically trivial, i.e., is isomorphic in the category of continuous bundles to the bundle  $X \times \mathbb{C}^k$ . Every such  $\rho$  can be obtained as the monodromy of the equation  $dF = \omega F$  on  $X$  where  $\omega$  is a matrix-valued 1-form on  $X$  satisfying  $d\omega - \omega \wedge \omega = 0$ . By  $\mathcal{T}(X)$  we denote the class of quotient groups  $G$  obtained by representations  $\rho$  as above.

Now, let  $r : M_G \rightarrow M$  be the regular covering of  $M$  satisfying condition (1.1) with transformation group  $G$ . Let  $G_1 \subset G$  be a subgroup of a finite index. Then there is a finite covering  $r_1 : M_1 \rightarrow M$  whose fibre is the quotient set  $G/G_1$  such that  $M_G$  is also the regular covering of  $M_1$  with transformation group  $G_1$ .

**Theorem 2.10** *Assume that  $G_1 \in \mathcal{T}(M_1)$ . Then there is a finite number of functions in  $\mathcal{H}_{2,1}(M_G) \cap C(\overline{M}_G)$  which separate all points in  $M_G$ .*

**Remark 2.11** (1) We will see from the proof that the functions in Theorem 2.10 can be taken even from  $\mathcal{H}_{2,\psi}(M_G)$  where  $\psi : M_G \rightarrow \mathbb{R}_+$  has a double exponential growth.

(2) As the group  $G$  in Theorem 2.10 one can take, e.g., a finitely generated free group (see, e.g., [O]) or a polycyclic group (see, e.g., [Ra]). If  $\dim_{\mathbb{C}} M = 1$  then, since  $M$  is homotopically equivalent to a one-dimensional  $CW$ -complex (see, e.g., [GR]), every quotient group  $G$  obtained by a linear representation  $\rho$  belongs to  $\mathcal{T}(M)$ .

### 3. Preliminary Results.

**3.1.** First, we recall some basic facts from the theory of bundles see, e.g., [Hi].

Let  $X$  be a complex analytic space and  $S$  be a complex analytic Lie group with unit  $e \in S$ . Consider an effective holomorphic action of  $S$  on a complex analytic space  $F$ . Here *holomorphic action* means a holomorphic map  $S \times F \rightarrow F$  sending  $s \times f \in S \times F$  to  $sf \in F$  such that  $s_1(s_2f) = (s_1s_2)f$  and  $ef = f$  for any  $f \in F$ . *Efficiency* means that the condition  $sf = f$  for some  $s$  and any  $f$  implies that  $s = e$ .

**Definition 3.1** *A complex analytic space  $W$  together with a holomorphic map (projection)  $\pi : W \rightarrow X$  is called a holomorphic bundle on  $X$  with structure group  $S$  and fibre  $F$ , if there exists a system of coordinate transformations, i.e., if*

(1) *there is an open cover  $\mathcal{U} = \{U_i\}_{i \in I}$  of  $X$  and a family of biholomorphisms  $h_i : \pi^{-1}(U_i) \rightarrow U_i \times F$ , that map “fibres”  $\pi^{-1}(u)$  onto  $u \times F$ ;*

(2) *for any  $i, j \in I$  there are elements  $s_{ij} \in \mathcal{O}(U_i \cap U_j, S)$  such that*

$$(h_i h_j^{-1})(u \times f) = u \times s_{ij}(u)f \quad \text{for any } u \in U_i \cap U_j, f \in F .$$

*A holomorphic bundle  $\pi : W \rightarrow X$  whose fibre is a Banach space  $F$  and the structure group is  $GL(F)$  (the group of linear invertible transformations of  $F$ ) is called a holomorphic Banach vector bundle. A holomorphic section of a holomorphic bundle  $\pi : W \rightarrow X$  is a holomorphic map  $s : X \rightarrow W$  satisfying  $\pi \circ s = id$ .*

We will use the following construction of holomorphic bundles (see, e.g. [Hi, Ch.1]):

Let  $S$  be a complex analytic Lie group and  $\mathcal{U} = \{U_i\}_{i \in I}$  be an open cover of  $X$ . By  $Z_{\mathcal{O}}^1(\mathcal{U}, S)$  we denote the set of holomorphic  $S$ -valued  $\mathcal{U}$ -cocycles. By definition,  $s = \{s_{ij}\} \in Z_{\mathcal{O}}^1(\mathcal{U}, S)$ , where  $s_{ij} \in \mathcal{O}(U_i \cap U_j, S)$  and  $s_{ij}s_{jk} = s_{ik}$  on  $U_i \cap U_j \cap U_k$ . Consider the disjoint union  $\sqcup_{i \in I} U_i \times F$  and for any  $u \in U_i \cap U_j$  identify the point  $u \times f \in U_j \times F$  with  $u \times s_{ij}(u)f \in U_i \times F$ . We obtain a holomorphic bundle  $W_s$  on  $X$  whose projection is induced by the projection  $U_i \times F \rightarrow U_i$ . Moreover, any holomorphic bundle on  $X$  with structure group  $S$  and fibre  $F$  is isomorphic (in the category of holomorphic bundles) to a bundle  $W_s$ .

**Example 3.2 (a)** Let  $M$  be a complex manifold. For any subgroup  $H \subset \pi_1(M)$  consider the unbranched covering  $r : M(H) \rightarrow M$  corresponding to  $H$ . We will describe  $M(H)$  as a holomorphic bundle on  $M$ .

First, assume that  $H \subset \pi_1(M)$  is a normal subgroup. Then  $M(H)$  is a regular covering of  $M$  and the quotient group  $G := \pi_1(M)/H$  acts holomorphically on  $M(H)$  by deck transformations. It is well known that  $M(H)$  in this case can be thought of as a *principle fibre bundle* on  $M$  with fibre  $G$  (here  $G$  is equipped with the discrete topology). Namely, let us consider the map  $R_G(g) : G \rightarrow G$ ,  $g \in G$ , defined by the formula

$$R_G(g)(q) = q \cdot g^{-1}, \quad q \in G.$$

Then for an open cover  $\mathcal{U} = \{U_i\}_{i \in I}$  of  $M$  by sets biholomorphic to open Euclidean balls in some  $\mathbb{C}^n$  there is a locally constant cocycle  $c = \{c_{ij}\} \in Z_{\mathcal{O}}^1(\mathcal{U}, G)$  such that  $M(H)$  is biholomorphic to the quotient space of the disjoint union  $V = \sqcup_{i \in I} U_i \times G$  by the equivalence relation:  $U_i \times G \ni x \times R_G(c_{ij})(q) \sim x \times q \in U_j \times G$ . The identification space is a holomorphic bundle with projection  $r : M(H) \rightarrow M$  induced by the projections  $U_i \times G \rightarrow U_i$ . In particular, when  $H = e$  we obtain the definition of the universal covering  $M_u$  of  $M$ .

Assume now that  $H \subset \pi_1(M)$  is not necessarily normal. Let  $X_H = \pi_1(M)/H$  be the set of cosets with respect to the (left) action of  $H$  on  $\pi_1(M)$  defined by left multiplications. By  $[Hq] \in X_H$  we denote the coset containing  $q \in \pi_1(M)$ . Let  $A(X_H)$  be the group of all homeomorphisms of  $X_H$  (equipped with the discrete topology). We define the homomorphism  $\tau : \pi_1(M) \rightarrow A(X_H)$  by the formula:

$$\tau(g)([Hq]) := [Hqg^{-1}], \quad q \in \pi_1(M).$$

Set  $Q(H) := \pi_1(M)/\text{Ker}(\tau)$  and let  $\tilde{g}$  be the image of  $g \in \pi_1(M)$  in  $Q(H)$ . By  $\tau_{Q(H)} : Q(H) \rightarrow A(X_H)$  we denote the unique homomorphism whose pullback to  $\pi_1(M)$  coincides with  $\tau$ . Consider the action of  $H$  on  $V = \sqcup_{i \in I} U_i \times \pi_1(M)$  induced by the left action of  $H$  on  $\pi_1(M)$  and let  $V_H = \sqcup_{i \in I} U_i \times X_H$  be the corresponding quotient set. Define the equivalence relation  $U_i \times X_H \ni x \times \tau_{Q(H)}(\tilde{c}_{ij})(h) \sim x \times h \in U_j \times X_H$  with the same  $\{c_{ij}\}$  as in the definition of  $M(e)$ . The corresponding quotient space is a holomorphic bundle with fibre  $X_H$  biholomorphic to  $M(H)$ .

**(b)** We retain the notation of example (a). Let  $B$  be a complex Banach space and  $GL(B)$  be the group of invertible bounded linear operators  $B \rightarrow B$ . Consider a homomorphism  $\rho : G \rightarrow GL(B)$ . Without loss of generality we assume that  $\text{Ker}(\rho) = e$ , for otherwise we can pass to the corresponding quotient group. The *holomorphic Banach vector bundle*  $E_\rho \rightarrow M$  associated with  $\rho$  is defined as the quotient of  $\sqcup_{i \in I} U_i \times B$  by the equivalence relation  $U_i \times B \ni x \times \rho(c_{ij})(w) \sim x \times w \in U_j \times B$  for any  $x \in U_i \cap U_j$ . Let us illustrate this construction by an example.

Let  $\phi : X_H \rightarrow \mathbb{R}^+$  ( $X_H := \pi_1(M)/H$ ) be a function satisfying

$$\phi(\tau(h)(x)) \leq c_h \phi(x), \quad x \in X_H, \quad h \in \pi_1(M), \quad (3.1)$$

where  $c_h$  is a constant depending on  $h$ . By  $l_{p,\phi}(X_H)$ ,  $1 \leq p \leq \infty$ , we denote the



Banach space of complex functions  $f$  on  $X_H$  with norm

$$\|f\|_{p,\phi} := \left( \sum_{g \in X_H} |f(g)|^p \phi(g) \right)^{1/p}. \quad (3.2)$$

Then according to (3.1) the map  $\rho$  defined by the formula  $[\rho(g)(f)](x) := f(\tau(g)(x))$ ,  $g \in \pi_1(M)$ ,  $x \in X_H$ , is a homomorphism of  $\pi_1(M)$  into  $GL(l_{p,\phi}(X_H))$ . By  $E_{p,\phi}(X_H)$  we denote the holomorphic Banach vector bundle associated with this  $\rho$ .

**3.2.** We retain the notation of Example 3.2. Let  $r : M' \rightarrow M$  be a covering where  $M' = M(H)$  (i.e.,  $\pi_1(M') = H$ ). Assume that  $M$  satisfies condition (1.1), i.e.,  $M \subset\subset N$  and  $\pi_1(M) = \pi_1(N)$ . Then there is an embedding  $M(H) \hookrightarrow N(H)$ . (Without loss of generality we consider  $M(H)$  as an open subset of  $N(H)$ .) Let  $\{V_i\}_{i \in I}$  be a finite acyclic open cover of  $\overline{M}$  by relatively compact sets. We set  $U_i := V_i \cap M$  and consider the open cover  $\mathcal{U} = \{U_i\}_{i \in I}$  of  $M$ . Then as in Example 3.2 (a) we can define  $M(H)$  by a cocycle  $c = \{c_{ij}\} \in Z_{\mathcal{O}}^1(\mathcal{U}, \pi_1(M))$ .

Further, let  $\psi : N(H) \rightarrow \mathbb{R}_+$  be a function such that  $\log \psi$  is uniformly continuous with respect to the path metric induced by a Riemannian metric pulled back from  $N$ . Fix a point  $z_0 \in M$  and identify  $r^{-1}(z_0)$  with  $z_0 \times X_H$  ( $X_H := \pi_1(M)/H$ ). We define the function  $\phi : X_H \rightarrow \mathbb{R}_+$  by the formula

$$\phi(x) := \psi(z_0, x), \quad x \in X_H.$$

It was proved in [Br1, Lemma 2.3] that  $\phi$  satisfies inequality (3.1). Then the bundle  $E_{p,\phi}(X_H)$  is well defined. By definition, any holomorphic section of this bundle is determined by a family  $\{f_i(z, g)\}_{i \in I}$  of holomorphic functions on  $U_i$  with values in  $l_{p,\phi}(X_H)$  satisfying

$$f_i(z, \tau(c_{ij})(h)) = f_j(z, h) \quad \text{for any } z \in U_i \cap U_j.$$

We introduce the Banach space  $B_{p,\phi}(X_H)$  of *bounded holomorphic sections*  $f = \{f_i\}_{i \in I}$  of  $E_{p,\phi}(X_H)$  with norm

$$\|f\|_{p,\phi} := \sup_{i \in I, z \in U_i} \|f_i(z, \cdot)\|_{p,\phi}. \quad (3.3)$$

(Here  $\|\cdot\|_{p,\phi}$  is the norm on  $l_{p,\phi}(X_H)$ , see (3.2).)

Further, let  $f \in \mathcal{H}_{p,\psi}(M(H))$  (see section 2.2 for the definition). We define the family  $\{f_i\}_{i \in I}$  of functions on  $U_i$  with values in the space of functions on  $X_H$  by the formula

$$f_i(z, g) := f(z, g), \quad z \in U_i, \quad i \in I, \quad g \in X_H. \quad (3.4)$$

It was established in [Br1, Proposition 2.4] that *the correspondence*  $f \mapsto \{f_i\}_{i \in I}$  *determines an isomorphism of Banach spaces*  $D : \mathcal{H}_{p,\psi}(M(H)) \rightarrow B_{p,\phi}(X_H)$ . (Here  $D$  is an isometry for  $\psi \equiv 1$ .)

Next, suppose that  $\{x_n\}_{n \geq 1} \subset M$  converges to  $x \in M$ . Then for sufficiently big  $n$  we can arrange  $r^{-1}(x_n)$  and  $r^{-1}(x)$  in sequences  $\{y_{in}\}_{i \geq 1}$  and  $\{y_i\}_{i \geq 1}$  such that every  $\{y_{in}\}$  converges to  $y_i$  as  $n \rightarrow \infty$ . For such  $n$  we define maps  $\tau_n(x) : r^{-1}(x) \rightarrow r^{-1}(x_n)$  so that  $\tau_n(y_i) = y_{in}$ ,  $i \in \mathbb{N}$ . Below,  $\tau_n^*$  denotes the transpose map generated by  $\tau_n$  on functions defined on  $r^{-1}(x_n)$  and  $r^{-1}(x)$ .

**Definition 3.3** Let  $X \subset M$  be a subset. We say that a function  $f$  on  $r^{-1}(X)$  belongs to the class  $C_{p,\psi}(r^{-1}(X))$  if

- (1)  $f|_{r^{-1}(x)} \in l_{p,\psi,x}(M')$  for all  $x \in X$  and
- (2) for any  $x \in X$  and any sequence  $\{x_n\} \subset X$  converging to  $x$  the sequence of functions  $\{\tau_n^*(f|_{r^{-1}(x_n)})\}$  converges to  $f|_{r^{-1}(x)}$  in the norm of  $l_{p,\psi,x}(M')$ .

By  $C_{p,\psi}^b(r^{-1}(X))$  we denote the Banach space of functions  $f \in C_{p,\psi}(r^{-1}(X))$  with norm

$$\|f\|_{p,\psi}^X := \sup_{x \in X} \|f|_{r^{-1}(x)}\|_{p,\psi,x} . \quad (3.5)$$

Note that if  $X \subset M$  is compact, then  $\|f\|_{p,\psi}^X < \infty$  for every  $f \in C_{p,\psi}(r^{-1}(X))$ .

Comparing with the above definition of  $D$  one determines a similar map for  $C_{p,\psi}^b(r^{-1}(X))$ . This gives an isomorphism  $D : C_{p,\psi}^b(r^{-1}(X)) \rightarrow CB_{p,\phi}^X(X_H)$  where  $CB_{p,\phi}^X(X_H)$  is the Banach space of *bounded continuous sections* of  $E_{p,\phi}(X_H)|_X$  with norm defined as in (3.3).

**3.3.** Most of our proofs are based on Theorem 1.3 of [Br1]. In its proof we use the above isomorphisms  $D$  and Cartan's A and B theorems for coherent Banach vector sheaves (see [B]). Let us formulate this result.

Suppose that  $r : M' \rightarrow M$  is a covering with  $M$  satisfying (1.1). We define  $\psi : M' \rightarrow \mathbb{R}_+$  as in section 3.2. Also, we define  $\mathcal{H}_{p,\psi}(M')$  and  $l_{p,\psi,x}(M')$  as in section 2.3. For Banach spaces  $E$  and  $F$  by  $\mathcal{B}(E, F)$  we denote the space of all linear bounded operators  $E \rightarrow F$  with norm  $\|\cdot\|$ .

**Theorem 3.4** For any  $p \in [1, \infty]$  there is a family  $\{L_z \in \mathcal{B}(l_{p,\psi,z}(M'), \mathcal{H}_{p,\psi}(M'))\}_{z \in M}$  holomorphic in  $z$  such that

$$(L_z h)(x) = h(x) \quad \text{for any } h \in l_{p,\psi,z}(M') \quad \text{and } x \in r^{-1}(z) .$$

Moreover,

$$\sup_{z \in M} \|L_z\| < \infty .$$

The following facts are simple corollaries of this result.

Suppose that  $X \subset M$  and  $f \in C_{p,\psi}^b(r^{-1}(X))$ . We define the function  $F$  on  $X \times M'$  by the formula

$$F(x, z) := (L_x(f|_{r^{-1}(x)}))(z) , \quad x \times z \in X \times M' . \quad (3.6)$$

Then  $F$  is continuous and  $F(x, \cdot) \in \mathcal{H}_{p,\psi}(M')$  for every  $x$ . Moreover, if  $X$  is open and  $f \in \mathcal{H}_{p,\psi}(r^{-1}(X))$ , then  $F \in \mathcal{O}(X \times M')$  and the map  $X \rightarrow \mathcal{H}_{p,\psi}(M')$ ,  $x \mapsto F(x, \cdot)$ , is holomorphic.

We can also express  $F$  in local coordinates. Namely, take  $x \in X$  and let  $U \subset M$  be a neighbourhood of  $x$  biholomorphic to an open Euclidean ball. Then  $r^{-1}(U) = \sqcup_{y \in r^{-1}(x)} V_y$  and there are biholomorphisms  $s_y : U \rightarrow V_y$  such that  $r \circ s_y = id$ . Now, the restriction of  $f \in C_{p,\psi}^b(r^{-1}(X))$  to  $r^{-1}(U \cap X)$  can be written as

$$f(z) = \sum_{y \in r^{-1}(x)} f(z) \chi_y(z) , \quad z \in r^{-1}(U \cap X) , \quad (3.7)$$

where  $\chi_y$  is the characteristic function of  $V_y$ . Let us introduce the functions  $\tilde{f}_y$ ,  $y \in r^{-1}(x)$ , by the formulas

$$\tilde{f}_y(v) = f(s_y(v)) , \quad v \in X \cap U .$$

Then we have

$$f(z) = \sum_{y \in r^{-1}(x)} \tilde{f}_y(v) \chi_y(z) , \quad v = r(z) \in U \cap X . \quad (3.8)$$

Consider the series

$$\sum_{y \in r^{-1}(x)} \tilde{f}_y(v) L_v(\chi_y|_{r^{-1}(v)}) , \quad v \in U \cap X , \quad (3.9)$$

with  $L_v$  as in Theorem 3.4.

**Proposition 3.5** *For  $p \in [1, \infty)$  the series in (3.9) converges in  $\mathcal{H}_{p,\psi}(M')$  to  $F(v, \cdot) := L_v(f|_{r^{-1}(v)})$  uniformly on every compact subset of  $U \cap X$ . If  $p = \infty$  and  $f \in C_{1,1}^b(r^{-1}(X))$  then this series also converges in  $\mathcal{H}_{\infty,\psi}(M')$  to  $F(v, \cdot)$  uniformly on every compact subset of  $U \cap X$ .*

**Proof.** Suppose that  $p \in [1, \infty)$  and  $f \in C_{p,\psi}^b(r^{-1}(X))$ . Let  $C \subset U \cap X$  be a compact subset. By the definition the function  $\Phi : U \cap X \rightarrow l_{p,\psi}(X_H)$ ,  $z \mapsto \tilde{f}(z)$ , is continuous (here we identify  $r^{-1}(x)$  with  $X_H$ ). Thus  $\Phi(C) \subset l_{p,\psi}(X_H)$  is compact. Fix a family  $\{X_i\}_{i \in \mathbb{N}}$  of finite subsets of  $X_H$  such that  $X_i \subset X_{i+1}$  for any  $i$  and  $\cup_{i=1}^{\infty} X_i = X_H$ . Let  $V_i \subset l_{p,\psi}(X_H)$  be a finite-dimensional subspace generated by functions  $\delta_z$  on  $X_H$  with  $z \in X_i$ . Here  $\delta_z(v) = 1$  if  $v = z$  and  $\delta_z(v) = 0$  if  $v \neq z$ . Then  $\cup_{i=1}^{\infty} V_i$  is everywhere dense in  $l_{p,\psi}(X_H)$  (since  $1 \leq p < \infty$ ). This and compactness of  $\Phi(C)$  imply that for any  $\epsilon > 0$  there exists an integer  $l$  such that  $\Phi(C) \subset V_l + B_\epsilon$  where  $B_\epsilon$  is the open ball in  $l_{p,\psi}(X_H)$  centered at 0 of radius  $\epsilon$ . By  $p_l : l_{p,\psi}(X_H) \rightarrow V_l$  we denote the projection sending  $v = \sum_{x \in X_H} v_x \delta_x \in l_{p,\psi}(X_H)$  to  $\sum_{x \in X_l} v_x \delta_x \in V_l$  (here all  $v_x \in \mathbb{C}$ ). Then for  $\epsilon$  as above and every  $v \in \Phi(C)$  we have  $\|v - p_l(v)\|_{p,\phi} < \epsilon$ . From this by (3.8), identifying  $r^{-1}(x)$  with  $X_H$ , we obtain

$$\sup_{v \in C} \|f|_{r^{-1}(v)} - \sum_{y \in X_l} \tilde{f}_y(v) \chi_y|_{r^{-1}(v)}\|_{p,\phi} < \epsilon . \quad (3.10)$$

Thus by the definition of operators  $L_v$  (see Theorem 3.4)

$$\sup_{v \in C} |F(v, \cdot) - \sum_{y \in X_l} \tilde{f}_y(v) L_v(\chi_y|_{r^{-1}(v)})|_{p,\psi}^M < C\epsilon \quad (3.11)$$

for some constant  $C$ . This implies the required uniform convergence for  $p \in [1, \infty)$ .

For  $p = \infty$  and  $f \in C_{1,1}^b(r^{-1}(X))$  we obtain anew that  $\Phi(C) \subset l_{1,1}(X_H)$  is compact. Then in the above notation we easily get  $\|v - p_l(v)\|_{\infty,\phi} < \epsilon$  for any  $v \in \Phi(C)$  (because  $\|\cdot\|_{\infty,\phi} = \|\cdot\|_{\infty,1} \leq \|\cdot\|_{1,1}$ ). Thus (3.10) is also valid for  $p = \infty$ . This gives (3.11) with  $p = \infty$ .  $\square$

## 4. Proofs of Theorem 2.1 and Corollary 2.2.

**Proof of Theorem 2.1.** Let  $M \subset\subset \widetilde{M} \subset N$  be complex manifolds such that  $\pi_1(M) = \pi_1(N)$  and  $\widetilde{M}$  is Stein. Consider an unbranched covering  $r : N' \rightarrow N$  of  $N$  and the corresponding coverings  $M' = r^{-1}(M)$  and  $\widetilde{M}' = r^{-1}(\widetilde{M})$  of  $M$  and  $\widetilde{M}$ . According to Example 3.2 (a)  $\widetilde{M}'$  is defined on an open cover  $\mathcal{U} = \{U_i\}_{i \in I}$  of  $\widetilde{M}$  by sets biholomorphic to open Euclidean balls by a locally constant cocycle  $\{\tilde{c}_{ij}\} \in Z_{\mathcal{O}}^1(\mathcal{U}, Q(H))$ . (Here we retain the notation of Example 3.2 (a) so that  $\widetilde{M}' = \widetilde{M}(H)$ .) Using this construction we identify  $r^{-1}(U_i)$  with  $U_i \times X_H$  ( $X_H = \pi_1(M)/H$ ). Also, we choose some points  $z_i \in U_i$  and assume that diameters of all  $U_i$  in the path metric on  $N$  induced by a Riemannian metric are uniformly bounded by a constant.

Let  $\phi : N' \rightarrow \mathbb{R}$  be a function uniformly continuous with respect to the path metric induced by the Riemannian metric pulled back from  $N$ . For every  $i \in I$  we define a function  $\phi_i : r^{-1}(U_i) \rightarrow \mathbb{R}$  by the formula

$$\phi_i(z, g) := \phi(z_i, g) , \quad z \times g \in r^{-1}(U_i) .$$

Then from the uniform continuity of  $\phi$  and boundedness of  $\text{diam}(U_i)$  for all  $i$  we obtain that there exists a constant  $c$  such that

$$|\phi(v) - \phi_i(v)| \leq c \quad \text{for every } v \in r^{-1}(U_i) , \quad i \in I . \quad (4.1)$$

Define a locally constant cocycle  $\phi_{ij}$  on the open cover  $\{r^{-1}(U_i)\}$  of  $\widetilde{M}'$  by the formula

$$\phi_{ij}(v) = \phi_i(v) - \phi_j(v) \quad \text{for } v \in r^{-1}(U_i \cap U_j) .$$

Then from (4.1) by the triangle inequality we get

$$\sup_{i,j,v} |\phi_{ij}(v)| \leq 2c . \quad (4.2)$$

This inequality implies that rewriting cocycle  $\{\phi_{ij}\}$  in the coordinates on  $\widetilde{M}$  (i.e., taking its direct image  $\{r_*(\phi_{ij})\}$  with respect to  $r$ ) we can regard it as a holomorphic cocycle on the cover  $\mathcal{U}$  with values in the Banach vector bundle  $E_{\infty,1}(X_H)$  with fibre  $l_{\infty,1}(X_H)$  defined on  $\widetilde{M}$  (see Example 3.2 (b)). This correspondence is described in [Br2, Proposition 2.4]. Since  $\mathcal{U}$  is acyclic and  $\widetilde{M}$  is Stein, from the above construction, a version of Cartan's B theorem for coherent Banach sheaves (see [B]), and the classical Leray theorem we obtain as in [Br2] that there are holomorphic functions  $f_i \in \mathcal{O}(r^{-1}(U_i))$  such that

(1) for every compact set  $K \subset U_i$ ,

$$\sup_{y \in r^{-1}(K)} |f_i(y)| < \infty ;$$

(2)

$$f_i(z) - f_j(z) = \phi_{ij}(z) \quad \text{for } z \in r^{-1}(U_i \cap U_j) .$$

Let  $\mathcal{V} = \{V_j\}_{j \in J}$  be a refinement of  $\mathcal{U}$  such that every  $V_j$  is open and relatively compact in some  $U_{i_j}$ . Then condition (1) implies that

$$\sup_{y \in r^{-1}(V_j)} |f_{i_j}(y)| < \infty . \quad (4.3)$$

Finally, define a function  $\tilde{f} \in \mathcal{O}(\tilde{M}')$  by the formula

$$\tilde{f}(z) := \phi_i(z) - f_i(z) , \quad z \in r^{-1}(U_i) . \quad (4.4)$$

Since  $\overline{M} \subset \tilde{M}$  is a compact set, there is a finite subcover of  $\mathcal{V}$  that covers  $\overline{M}$ . From here, (4.3) and (4.1) for the restriction  $f_\phi := \tilde{f}|_{\overline{M}'}$  we obtain (for some  $C$ )

$$|f_\phi(z) - \phi(z)| < C \quad \text{and} \quad |df_\phi(z)| < C \quad \text{for any} \quad z \in M' . \quad (4.5)$$

The proof of the theorem is complete.  $\square$

**Proof of Corollary 2.2.** We retain the notation of the proof of Theorem 2.1.

Let  $d_o := d(o, \cdot)$  be the distance on  $N'$  from a fixed point  $o \in M'$  and let  $f := f_\phi$  be the function from Theorem 2.1 for  $\phi = d_o$ . Consider a finite open cover  $\{U_i\}_{i=1}^l$  of  $\overline{M}$  such that every  $U_i \subset\subset \tilde{M}$  is biholomorphic to an open Euclidean ball. As above we identify  $r^{-1}(U_i) \subset N'$  with  $U_i \times X_H$ . Fix an element  $e \in X_H$  and set  $o_i(z) = (z, e) \in r^{-1}(U_i)$  for every  $z \in U_i$ ,  $1 \leq i \leq l$ , and  $d_{o_i(z)}(v) := d(o_i(z), v)$ ,  $v \in N'$ . Then from compactness of every  $\overline{U_i}$  by the triangle inequality we get

$$|d_{o_i(z)}(v) - d_o(v)| \leq a , \quad 1 \leq i \leq l , \quad (4.6)$$

for some constant  $a$ . By  $B_{z,i}(R)$  we denote the open ball on  $r^{-1}(z)$  of radius  $R$  centered at  $o_i(z)$  with respect to the induced metric  $d|_{r^{-1}(z)}$ . Also, by  $\#A$  we denote the number of elements of  $A$ . Now we prove

**Lemma 4.1** *There is  $k \in \mathbb{N}$  such that*

$$\#B_{z,i}(R) \leq e^{kR} , \quad 1 \leq i \leq l .$$

**Proof.** Let  $\tilde{r} : N_u \rightarrow N$  be the universal covering of  $N$  and  $r' : N_u \rightarrow N'$  be the intermediate covering, i.e.,  $\tilde{r} = r \circ r'$ . We equip  $N_u$  with the path metric  $\tilde{d}$  induced by the Riemannian metric pulled back from  $N$ , the same as in the definition of the metric  $d$  on  $N'$ . Let  $\tilde{o}_i(z) \in \tilde{r}^{-1}(z)$  be such that  $r'(\tilde{o}_i(z)) = o_i(z)$ . By  $\tilde{B}_{z,i}(R)$  we denote the open ball on  $\tilde{r}^{-1}(z)$  of radius  $R$  centered at  $\tilde{o}_i(z)$  with respect to the metric  $\tilde{d}|_{\tilde{r}^{-1}(z)}$ . Let us check that  $r'(\tilde{B}_{z,i}(R)) = B_{z,i}(R)$ .

Indeed, let  $y \in \tilde{B}_{z,i}(R)$  and  $\gamma_y$  be a path joining  $\tilde{o}_i(z)$  and  $y$  in  $N_u$  whose length is less than  $R$  (such a path exists by the definition of  $\tilde{d}$ ). Then  $r'(\gamma_y)$  is a path joining  $o_i(z)$  and  $r'(y)$  in  $N'$ . By the definition of the metrics on  $N_u$  and  $N'$  the length of  $r'(\gamma_y)$  does not exceed the length of  $\gamma_y$ . In particular, it is less than  $R$ . Thus  $d_{o_i(z)}(r'(y)) < R$ , i.e.,  $r'(y) \in B_{z,i}(R)$ . Conversely, let  $w \in B_{z,i}(R)$  and let  $\gamma_w$  be a path in  $N'$  joining  $o_i(z)$  and  $w$  with length less than  $R$ . By the covering homotopy theorem (see, e.g., [Hu, Chapter III]) there is a path  $\tilde{\gamma}_w \subset N_u$  that covers  $\gamma_w$  and joins  $\tilde{o}_i(z)$  with some point  $\tilde{w}$  such that  $r'(\tilde{w}) = w$ . Moreover, by the definition, the

length of  $\tilde{\gamma}_w$  is the same as the length of  $\gamma_w$ . In particular, it is less than  $R$ . Thus  $\tilde{w} \in \tilde{B}_{z,i}(R)$ . This shows that  $r'(\tilde{B}_{z,i}(R)) = B_{z,i}(R)$ . In turn, the latter implies that

$$\#B_{z,i}(R) \leq \#\tilde{B}_{z,i}(R) . \quad (4.7)$$

Next, let  $A$  be a finite set of generators of  $\pi_1(N)$  (recall that condition (1.1) implies that  $\pi_1(N)$  is finitely generated). By  $d_w$  we denote the word metric on  $\pi_1(N)$  with respect to  $A$ . Now, from compactness of every  $\overline{U_i}$  by the Švarc-Milnor lemma (see, e.g., [BH, p.140]) we obtain that there exists a constant  $c$  such that for any  $z \in U_i$ ,  $1 \leq i \leq l$ , and  $g, h \in \pi_1(N)$ ,

$$c^{-1}d_w(g, h) \leq \tilde{d}((z, g), (z, h)) \leq cd_w(g, h) \quad (4.8)$$

(Here we identify  $\tilde{r}^{-1}(U_i)$  with  $U_i \times \pi_1(N)$  as in Example 3.2 (a).) Let  $B_R \subset \pi_1(N)$  be the open ball of radius  $R$  centered at 1 with respect to  $d_w$ . Then there is a natural number  $\tilde{k}$  such that

$$\#B_R \leq e^{\tilde{k}R} \quad \text{for any } R \geq 0 . \quad (4.9)$$

From here, (4.8) and (4.7) we get for  $k := \tilde{k}c$

$$\#B_{z,i}(R) \leq e^{kR} , \quad 1 \leq i \leq l . \quad \square$$

We proceed with the proof of the corollary. Let us define  $\alpha := \frac{k+1}{2}$  and prove that  $F = e^{-\alpha f} \in \mathcal{H}_{2,1}(M') \cap C(\overline{M'})$ . Since  $\mathcal{H}_{2,1}(M') \hookrightarrow H_1^2(M')$  (see section 2.3), this implies the required statement.

Let  $z \in U_i$  for some  $1 \leq i \leq l$ . We will estimate  $|F|_{2,1,z}$  (see (2.2)). By the definition using inequalities (4.5), (4.6) and Lemma 4.1 we obtain

$$\begin{aligned} |F|_{2,1,z}^2 &= \sum_{y \in r^{-1}(z)} |e^{-\alpha f(y)}|^2 \leq e^{2\alpha(a+C)} \cdot \sum_{y \in r^{-1}(z)} e^{-2\alpha d_{o_i(z)}(y)} \leq \\ &e^{2\alpha(a+C)} \cdot \sum_{R=0}^{\infty} e^{-2\alpha R} \cdot \#B_{z,i}(R) \leq e^{2\alpha(a+C)} \cdot \sum_{R=0}^{\infty} e^{(-2\alpha+k)R} = \frac{e^{2\alpha(a+C)+1}}{e-1} . \end{aligned}$$

Therefore

$$|F|_{2,1}^M := \sup_{z \in M} |F|_{2,1,z} \leq \left( \frac{e^{2\alpha(a+C)+1}}{e-1} \right)^{1/2} .$$

This shows that  $F \in \mathcal{H}_{2,1}(M') \cap C(\overline{M'})$ .  $\square$

**Remark 4.2** (1) Using some construction from [Br2] one can prove that the constant  $C$  in Theorem 2.1 for  $\phi = d_o$  (see (4.5)) can be chosen independent of the covering  $r : M' \rightarrow M$ . It depends only on  $M, \overline{M}$  and the Riemannian metric on  $N$ . (2) Consider the holomorphic map  $f : M' \rightarrow \mathbb{C}$  with  $f$  as in Corollary 2.2. Then

$$f(M') \subset S := \{z \in \mathbb{C} : |Im z| < C , -C < Re z < \infty\}$$

where  $C$  is the constant in Theorem 2.1 for  $\phi = d_o$ . Let  $B_t = \{x \in M' : d_o(x) < t\}$  be the open ball in  $M'$  centered at  $o$  of radius  $t$  and  $S_R := \{z \in S : Re z \geq R\}$ . Then

$$f^{-1}(S_R) \subset M' \setminus B_{R-C} \quad \text{for } R > C \quad \text{and} \quad f^{-1}(S \setminus S_R) \subset B_{R+C} .$$

Using such  $f$  one can construct holomorphic functions on  $M'$  decreasing faster than the function  $F$  from Corollary 2.2. Actually, let  $l : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a continuous function monotonically increasing for  $x \geq R_0$ . Consider a holomorphic function  $g$  on  $S$  satisfying

$$\log |g(x + iy)| \geq l(x) \quad \text{for } x \geq R_0 \quad \text{and} \quad \inf_{z \in S} |g(z)| > 0. \quad (4.10)$$

Then one can easily check that the function  $G = g \circ f \in \mathcal{O}(M') \cap C(\overline{M}')$  satisfies

$$|G(z)| \geq e^{l(d_o(z)-C)} \quad \text{for } d_o(z) \geq R_0 + C.$$

In particular,  $H = 1/G \in \mathcal{O}(M') \cap C(\overline{M}')$  satisfies (for some  $c_1 > 0$ )

$$|H(z)| \leq c_1 e^{-l(d_o(z)-C)}, \quad z \in \overline{M}' \setminus B_C. \quad (4.11)$$

Observe that the Harnack inequality for positive harmonic functions implies for  $g$  as in (4.10) (for some positive  $\tilde{c}_1, \tilde{c}_2$ )

$$l(x) \leq \log |g(x)| \leq \tilde{c}_1 e^{\tilde{c}_2 x}, \quad x \geq R_0.$$

This and the properties of  $f$  impose the following restriction on the decay of  $H$ :

$$|H(z)| \geq e^{-\tilde{c}_3 e^{\tilde{c}_2 d_o(z)}}, \quad z \in M'. \quad (4.12)$$

**Example 4.3** As the function  $g$  in (4.10) one can take, e.g.,  $g(z) = e^{z^n}$  for  $n \in \mathbb{N}$  (in this case  $l(x) = (1 - \epsilon)x^n$  for any  $\epsilon > 0$ ), or  $g(z) = e^{C_1 e^{C_2 z}}$  for  $C_1 > 0$  and  $0 < C_2 < \frac{\pi}{2C}$  with  $C$  as above (in this case  $l(x) = C_1 \cos(C_2 C) e^{C_2 x}$ ). For the latter example estimate (4.11) shows that the lower bound (4.12) of the decay of  $H$  is attainable.

## 5. Proof of Theorem 2.4.

Suppose that  $M \subset\subset N$  are domains in a Stein manifold,  $M$  is strictly pseudoconvex and  $\pi_1(M) = \pi_1(N)$ . Using the Remmert embedding theorem (see, e.g., [GR]) we may assume without loss of generality that  $N$  is a domain in a closed complex submanifold of some  $\mathbb{C}^k$ . Let  $r : N' \rightarrow N$  be an unbranched covering of  $N$ . As usual, we set  $M' = r^{-1}(M)$  and  $bM' = r^{-1}(bM)$  where  $bM$  is the boundary of  $M$ . We must show that every point in  $bM'$  is a peak point for  $\mathcal{H}_{p,1}(M')$ ,  $1 \leq p \leq \infty$ . In our proof we use a result on uniform estimates for solutions of certain  $\bar{\partial}$ -equations on  $M'$ . To its formulation we first introduce the corresponding class of  $(0,1)$ -forms on  $M'$ .

Let  $\{V_i\}_{i \in I}$  be a finite acyclic open cover of  $\overline{M}$  by relatively compact complex coordinate systems. We set  $U_i \cap M$  and consider the open cover  $\mathcal{U} = \{U_i\}_{i \in I}$  of  $M$ . Let  $X_H$  be the fibre of  $N'$  with  $H = \pi_1(N')$ . Using the construction of Example 3.2

(a) we identify  $r^{-1}(U_i)$  with  $U_i \times X_H$ . Let  $\omega$  be a  $(0,1)$ -form on  $M'$ . Then in local coordinates on  $r^{-1}(U_i)$  it is presented as

$$\omega(v, x) = \sum_{j=1}^n a_j(v, x) d\bar{v}_j \quad \text{for } v \times x \in U_i \times X_H$$

where  $v = (v_1, \dots, v_n)$  are coordinates on  $U_i$ . Consider every  $a_j$  as a function on  $U_i$  with values in the space of functions on  $X_H$ . We assume that for every  $i \in I$

$$a_j \in C^\infty(U_i, l_{p,1}(X_H)) , \quad 1 \leq j \leq n . \quad (5.1)$$

Then for such an  $\omega$  its direct image  $r_*(\omega)$  is a bounded  $C^\infty$  form with values in the Banach vector bundle  $E_{p,1}(X_H)$  (see section 3.1). Also, we assume that the norm of  $\omega$  defined by the formula

$$\|\omega\| := \sup_{i \in I} \max_{1 \leq j \leq n} |a_j|_{p,1}^{U_i} \quad (5.2)$$

is finite. (Recall that  $|\cdot|_{p,1}^{U_i}$  are norms on  $C_{p,1}(r^{-1}(U_i))$ , see (3.5).)

**Proposition 5.1** *There is a constant  $C > 0$  and for each  $\bar{\partial}$ -closed  $(0,1)$ -form  $\omega$  satisfying (5.1) there is a function  $f \in C^\infty(M') \cap C_{p,1}^b(M')$  such that*

$$\bar{\partial}f = \omega \quad \text{and} \quad |f|_{p,1}^M \leq C\|\omega\| .$$

**Proof.** We apply the operators  $L_z$  from Theorem 3.4 to  $\omega$ . Namely, let us define a form  $\tilde{\omega}$  on  $M$  by the formula

$$\tilde{\omega}(v, z) := \sum_{j=1}^n (L_v a_j(v, \cdot))(z) d\bar{v}_j \quad \text{for } v \times z \in U_i \times M' , \quad i \in I .$$

It is readily seen that  $\tilde{\omega}$  is a bounded  $\bar{\partial}$ -closed  $C^\infty$  form on  $M$  with values in  $\mathcal{H}_{p,1}(M')$ . We define the norm of  $\tilde{\omega}$  by

$$\|\tilde{\omega}\| := \sup_{i \in I, v \in U_i} \max_{1 \leq j \leq n} |L_v a_j(v, \cdot)|_{p,1}^M \quad (5.3)$$

where  $|\cdot|_{p,1}^M$  is norm on  $\mathcal{H}_{p,1}(M')$ . Then according to Theorem 3.4 there is a constant  $c$  (independent of  $\omega$ ) such that

$$\|\tilde{\omega}\| \leq c\|\omega\| . \quad (5.4)$$

Further, we use Lemma 1 from [He]. According to this lemma *there exist a strictly pseudoconvex domain  $W \subset \mathbb{C}^k$  with  $C^2$  boundary such that  $W \cap N = M$  and a holomorphic map  $\pi$  from a neighbourhood  $U(\bar{W})$  of  $\bar{W}$  onto  $U(\bar{W}) \cap N$  such that  $\pi(W) = M$  and  $\pi|_{U(\bar{W}) \cap N}$  is the identity map.*

Using this result we obtain that the pullback  $\pi^*\tilde{\omega}$  with respect to  $\pi$  is a bounded  $\bar{\partial}$ -closed  $C^\infty$  form on  $W$  with values in  $\mathcal{H}_{p,1}(M')$ . Moreover, there is a constant  $c'$  (depending on  $\pi$  and  $W$ ) such that

$$\|\pi^*\tilde{\omega}\| \leq c'\|\tilde{\omega}\| . \quad (5.5)$$



Here for  $\pi^*\tilde{\omega}(w, \cdot) = \sum_{j=1}^k \tilde{a}_j(w, \cdot) d\bar{w}_j$ ,  $w = (w_1, \dots, w_k) \in \mathbb{C}^k$ , we define

$$\|\pi^*\tilde{\omega}\| := \sup_{w \in W} \max_{1 \leq j \leq k} |\tilde{a}_j(w, \cdot)|_{p,1}^M .$$

In [SH] uniform estimates for solutions of  $\bar{\partial}$ -equations on so-called pseudoconvex polyhedra were obtained by means of global integral formulas. This class contains, in particular, strictly pseudoconvex domains with  $C^2$  boundaries. Note that the estimates in [SH] remain valid if one solves Banach-valued  $\bar{\partial}$ -equations. Therefore from the results of [SH] we obtain that *there exists a bounded  $C^\infty$  function  $h$  on  $W$  with values in  $\mathcal{H}_{p,1}(M')$  such that  $\bar{\partial}h = \pi^*\tilde{\omega}$ . Moreover,*

$$\|h\| \leq c'' \|\pi^*\tilde{\omega}\| \tag{5.6}$$

for some  $c''$  (depending on  $W$  only). Here

$$\|h\| := \sup_{w \in W} |h(w, \cdot)|_{p,1}^M .$$

Finally, define a function  $f$  on  $M'$  by the formula

$$f(z) := h(r(z), z) , \quad z \in M' .$$

Using that  $r$  is holomorphic,  $\tilde{\omega}(r(z), z) = \omega(z)$ ,  $z \in M'$ , and  $(\pi^*\tilde{\omega})|_M = \tilde{\omega}$  we easily conclude that  $\bar{\partial}f = \omega$ . By the definition  $f \in C^\infty(M') \cap C_{p,1}^b(M')$  and from (5.4)-(5.6) we have (for some  $C$ )

$$|f|_{p,1}^M \leq C \|\omega\| . \quad \square$$

**Remark 5.2** (1) An analogous to Proposition 5.1 statement is valid for a similar class of bounded  $\bar{\partial}$ -closed  $(0, q)$ -forms on  $M'$ .

(2) Using the main result of [He] and the estimates from [SH] one can show that the result of Proposition 5.1 is valid also for coverings of non-degenerate pseudoconvex polyhedrons on Stein manifolds (see [He] and [SH] for the definition).

We pass to the proof of Theorem 2.4. Take a point  $z \in bM'$  and set  $v = r(z) \in bM$ . Let  $U \subset\subset N$  be a simply connected coordinate neighbourhood of  $v$  and let  $W \subset N'$  be the neighbourhood of  $z$  such that  $r : W \rightarrow U$  is biholomorphic. Since  $M$  is strictly pseudoconvex,  $v$  is a peak point for  $\mathcal{O}(U \cap M)$  for a sufficiently small  $U$ . Moreover, for such  $U$  we can find  $\tilde{f} \in \mathcal{O}(U \cap M)$  with a peak point at  $v$  such that  $\tilde{f} \in L^q(U \cap M)$  for all  $1 \leq q < \infty$  (see [GHS, p.575]). Then  $f := (r^*\tilde{f})|_{W \cap M'}$  has a peak point at  $z$  and  $f \in L^q(W \cap M')$  for all  $1 \leq q < \infty$ . Next, let  $\tilde{\rho} \in C^\infty(U)$  be a cut-off function which equals 1 in a neighbourhood  $O \subset\subset U$  of  $v$  and 0 outside  $U$ . Consider its pullback  $\rho := (r^*\tilde{\rho})|_W \in C^\infty(W)$ . Clearly the  $(0, 1)$ -form  $\omega = \bar{\partial}(\rho f)$  on  $M'$  satisfies conditions of Proposition 5.1. Then this proposition implies that there exists a function  $h \in C^\infty(M') \cap C_{p,1}^b(M')$  such that  $\bar{\partial}h = \omega$ . Finally, consider the function  $h_z := \rho f - h$ . Then  $h_z$  is holomorphic, has a peak point at  $z_0$  and belongs to  $H_1^p(M')$  for  $1 \leq p < \infty$  by the choice of  $f$ . Also, for  $p = \infty$  the function  $h_z$  is bounded outside  $W$ .

The proof of the theorem is complete.  $\square$

## 6. Proofs of Theorem 2.7 and Corollary 2.9.

**Proof of Theorem 2.7.** Let  $M \subset\subset \widetilde{M} \subset N$  be manifolds satisfying condition (1.1) with  $\dim_{\mathbb{C}} M \geq 2$ . Let  $D \subset\subset M$  be an open subset whose boundary  $bD$  is a connected  $C^k$  submanifold of  $M$  ( $1 \leq k \leq \infty$ ). Consider a covering  $r : N' \rightarrow N$  and set  $M' = r^{-1}(M)$ ,  $D' = r^{-1}(D)$  and  $bD' = r^{-1}(bD)$ . Let  $\psi : N' \rightarrow \mathbb{R}_+$  be such that  $\log \psi$  is uniformly continuous with respect to the path metric induced by a Riemannian metric pulled back from  $N$ . Let  $f \in C^s(bD')$ ,  $0 \leq s \leq k$ , be a  $CR$ -function satisfying the hypotheses of Theorem 2.7.

(A) First, we will prove the theorem for  $s = 0$  under the additional assumption

$$f \in C_{p,\psi}(bD') \quad \text{for } p \in [1, \infty) \quad \text{and} \quad f \in C_{1,1}(bD') \quad \text{for } p = \infty. \quad (6.1)$$

(We use here that  $C_{1,1}(bD') \subset C_{\infty,1}(bD') = C_{\infty,\psi}(bD')$ , see Definition 3.3.)

For a  $CR$ -function  $f$  satisfying (6.1) we define a continuous  $\mathcal{H}_{p,\psi}(M')$ -valued function  $F$  on  $bD'$  by the formula

$$F(v) := L_v(f|_{r^{-1}(v)}), \quad v \in bD,$$

where  $L_v$  are operators from Theorem 3.4, see section 3.3.

**Lemma 6.1**  *$F$  is a  $\mathcal{H}_{p,\psi}(M')$ -valued continuous  $CR$ -function.*

**Proof.** Let  $U \subset M$  be a simply connected coordinate neighbourhood of a point  $x \in bD$ . It suffices to check that  $F|_{U \cap bD}$  satisfies the required property. Note that by Proposition 3.5

$$[F(v)](z) = \sum_{y \in r^{-1}(x)} \tilde{f}_y(v) H_y(v, z), \quad v \times z \in (U \cap bD) \times M', \quad (6.2)$$

where  $\tilde{f}_y(v) = f(s_y(v))$ ,  $v \in U \cap bD$ , and  $s_y : U \rightarrow V_y$  is a biholomorphic map onto the connected component  $V_y$  of  $r^{-1}(U)$  containing  $y$ . By the definition of operators  $L_v$  functions  $H_y$  are restrictions to  $bD \times M'$  of some holomorphic functions on  $U \times M'$ . Moreover, Proposition 3.5 implies that the series in (6.2) converges uniformly to  $F$  on every compact subset of  $(U \cap bD) \times M'$ . Next, since  $f|_{V_y \cap bD'}$  is a continuous  $CR$ -function,  $\tilde{f}_y$  is a continuous  $CR$ -function on  $U \cap bD$ . Also, by the definition of  $H_y$ , for a fixed  $z \in M'$  every  $H_y(\cdot, z)$  is a continuous  $CR$ -function on  $U \cap bD$ . Hence,  $\tilde{f}_y \cdot H_y(\cdot, z)$  is a continuous  $CR$ -function on  $U \cap bD$ , as well. Indeed, for each  $(n, n-2)$ -form  $\omega$  with a compact support in  $U$  we have

$$\int_{U \cap bD} \tilde{f}_y(v) \cdot H_y(v, z) \bar{\partial} \omega(v) = \int_{U \cap bD} \tilde{f}_y(v) \bar{\partial} (H_y(v, z) \cdot \omega(v)) = 0$$

because  $\tilde{f}_y$  is  $CR$ . Since the series in (6.2) converges uniformly to  $[F(\cdot)](z)$  on every compact subset of  $(U \cap bD) \times z$ , every  $[F(\cdot)](z)$ ,  $z \in M'$ , is a continuous  $CR$ -function on  $U \cap bD$ . This implies the required statement.  $\square$

Further, since  $[F(v)](z)$  from Lemma 6.1 is holomorphic in  $z \in M'$ , we can expand it in the Taylor series in a complex coordinate neighbourhood  $U_z$ ,

$$[F(v)](w) = \sum_{0 \leq |\alpha| < \infty} F_\alpha(v) w^\alpha, \quad v \in bD. \quad (6.3)$$

Here  $\alpha = (\alpha_1, \dots, \alpha_s) \in (\mathbb{Z}_+)^s$ ,  $|\alpha| = \sum_{i=1}^s \alpha_i$ ,  $w^\alpha = w_1^{\alpha_1} \dots w_n^{\alpha_n}$  and  $w = (w_1, \dots, w_n)$  are coordinates on  $U_z$  such that  $w(z) = 0$ . Now, from Lemma 6.1 follows that each  $F_\alpha$  in (6.3) is a continuous  $CR$ -function on  $bD$ . Then by Theorem 3.14 of Harvey [Ha] for every  $F_\alpha$  there exists a function  $\tilde{F}_\alpha \in \mathcal{O}(D) \cap C(\overline{D})$  such that  $\tilde{F}_\alpha|_{bD} = F_\alpha$ . Also, for a sufficiently small  $U_z$  using estimates of the Cauchy integrals for derivatives of a holomorphic function and compactness of  $bD$  we get from (6.3)

$$M := \sup_{\alpha, v \in bD} |F_\alpha(v)| < \infty .$$

Thus by the maximum modulus principle

$$\sup_{\alpha, v \in \overline{D}} |\tilde{F}_\alpha(v)| = M < \infty .$$

The latter implies that for a sufficiently small  $U_z$  the series

$$\tilde{F}_z(v, w) = \sum_{0 \leq |\alpha| < \infty} \tilde{F}_\alpha(v) w^\alpha , \quad v \times w \in \overline{D} \times U_z ,$$

converges absolutely and uniformly. Hence,  $\tilde{F}_z \in \mathcal{O}(D \times U_z) \cap C(\overline{D} \times U_z)$ . Further, assume that for  $y, z \in M'$  we have  $U_y \cap U_z \neq \emptyset$ . Then for every  $w \in U_y \cap U_z$  and  $v \in bD$

$$\tilde{F}_y(v, w) - \tilde{F}_z(v, w) = [F(v)](w) - [F(v)](w) = 0 .$$

This leads to the identity

$$\tilde{F}_y(\cdot, w) = \tilde{F}_z(\cdot, w) , \quad w \in U_y \cap U_z .$$

Thus we can define a function  $F \in \mathcal{O}(D \times M') \cap C(\overline{D} \times M')$  by the formula

$$\tilde{F}(v, w) := \tilde{F}_z(v, w) , \quad v \times w \in \overline{D} \times U_z . \quad (6.4)$$

**Lemma 6.2**  $\tilde{F}(v, \cdot) \in \mathcal{H}_{p, \psi}(M')$  for any  $v \in D$ .

**Proof.** Observe that the evaluation at  $v \in D$  is a linear continuous functional on the Banach space  $\mathcal{O}(D) \cap C(\overline{D})$  equipped with supremum norm. Identifying  $\mathcal{O}(D) \cap C(\overline{D})$  with its trace space on  $bD$  and using the Hahn-Banach and F. Riesz theorems we have

$$h(v) = \int_{bD} h(\xi) d\mu(\xi) , \quad h \in \mathcal{O}(D) \cap C(\overline{D}) ,$$

where  $\mu$  is a complex regular Borel measure on  $bD$  with the total variation  $Var \mu = 1$ . Thus for every fixed  $w \in M'$  we have

$$\tilde{F}(v, w) = \int_{bD} [F(\xi)](w) d\mu(\xi) .$$

Now, by the definition of the norm on  $\mathcal{H}_{p,\psi}(M')$  using the triangle inequality, the identity  $\tilde{F}(v, \cdot) = F(v)$ ,  $v \in bD$ , and the fact that  $F$  is a continuous  $\mathcal{H}_{p,\psi}(M')$ -valued function on  $bD$  we obtain

$$\begin{aligned} |\tilde{F}(v, \cdot)|_{p,\psi}^M &:= \sup_{z \in M} \left( \sum_{y \in r^{-1}(z)} |\tilde{F}(v, y)|^p \psi(y) \right)^{1/p} \leq \\ &\sup_{z \in M} \left( \sum_{y \in r^{-1}(z)} \left( \int_{bD} |\tilde{F}(\xi, y)| |d\mu(\xi)| \right)^p \psi(y) \right)^{1/p} \leq \\ &\sup_{z \in M} \left( \int_{bD} \left( \sum_{y \in r^{-1}(z)} |[F(\xi)](y)|^p \psi(y) \right)^{1/p} |d\mu(\xi)| \right) \leq \sup_{\xi \in bD} |F(\xi)|_{p,\psi}^M < \infty . \quad \square \end{aligned}$$

Further, set

$$f'(z) := \tilde{F}(r(z), z) , \quad z \in \overline{D'} . \quad (6.5)$$

Then using the inequalities of Lemma 6.2 we get

$$f' \in \mathcal{O}(D') \cap C(\overline{D'}) , \quad f'|_{bD'} = F|_{bD'} = f , \quad \text{and}$$

$$|f'|_{p,\psi,z} := \left( \sum_{y \in r^{-1}(z)} |f'(y)|^p \psi(y) \right)^{1/p} \leq \sup_{\xi \in bD} |F(\xi)|_{p,\psi}^M \leq c |f|_{p,\psi}^{bD} , \quad z \in D ,$$

see (3.5) for the definition of  $|\cdot|_{p,\psi}^{bD}$ . (Here the last inequality follows directly from Theorem 3.4.) The latter implies that  $f'|_{r^{-1}(z)} \in l_{p,\psi,z}(M')$ ,  $z \in D$ , see section 2.3. Thus  $f' \in \mathcal{H}_{p,\psi}(D') \cap C(\overline{D'})$  and

$$|f'|_{p,\psi}^D := \sup_{z \in D} |f'|_{p,\psi,z} \leq c \sup_{z \in bD} |f|_{p,\psi,z} \quad (:= c |f|_{p,\psi}^{bD}) . \quad (6.6)$$

This completes the proof of the theorem for  $s = 0$  under assumption (6.1).

(B) Let us consider the general case of a continuous  $CR$ -function  $f$  on  $bD'$  satisfying

$$f|_{r^{-1}(x)} \in l_{p,\psi,x}(M') \quad \text{for any } x \in D \quad \text{and} \quad m := \sup_{x \in bD} |f|_{p,\psi,x} < \infty . \quad (6.7)$$

According to Remark 4.2 (2) and Example 4.3 there is a constant  $c > 0$  such that for any  $0 < c_1 < c$  and  $c_2 > 0$  there exists a function  $F_{c_1,c_2} \in \mathcal{O}(M') \cap C(\overline{M'})$  satisfying

$$e^{-c_3 e^{c_1 d_o(z)}} \leq |F_{c_1,c_2}(z)| \leq e^{-c_2 e^{c_1 d_o(z)}} \quad \text{for all } z \in \overline{M'} \quad (6.8)$$

with  $c_3$  depending on  $c_2, c_1, c, M, M'$  such that  $c_3 \rightarrow 0$  as  $c_2 \rightarrow 0$ . (Recall that  $d_o, o \in M'$ , is the distance on  $N'$  defined as in Corollary 2.2.) Define a continuous  $CR$ -function  $f_{c_1,c_2}$  by the formula

$$f_{c_1,c_2}(z) := f(z) F_{c_1,c_2}(z) , \quad z \in bD' .$$

**Lemma 6.3**  $f_{c_1, c_2}$  satisfies assumption (6.1).

**Proof.** Note that for any  $l \in \mathbb{N}$  there is a nonnegative  $r$  such that

$$e^{-c_2 e^{c_1 d_o(z)}} < e^{-l d_o(z)} \quad \text{for } d_o(z) > r, z \in \overline{M'}.$$

Let  $U \subset\subset M$  be a neighbourhood of  $\overline{D}$  and  $U' = r^{-1}(U) \subset M'$ . From the above inequality arguing as in the proof of Corollary 2.2 we obtain that  $F_{c_1, c_2} \in \mathcal{H}_{p,1}(U')$  for any  $p \in [1, \infty]$ . Then from [Br, Proposition 2.4] follows that  $F_{c_1, c_2}|_{bD'}$  belongs to  $C_{p,1}(bD')$  for all  $p$ .

Next, take a point  $x \in bD$  and prove that  $f_{c_1, c_1}$  is  $C_{p, \psi}$ -continuous over  $x$ . Let  $U_x \subset M$  be a complex (simply connected) coordinate neighbourhood of  $x$ . We will identify  $r^{-1}(U_x)$  with  $U_x \times X_H$  where  $X_H$  is the fibre of  $r : M' \rightarrow M$ . Consider a sequence  $\{x_n\} \subset U_x \cap bD$  convergent to  $x$ . For  $g \in X_H$  put  $a_n(g) := f(x_n, g)$ ,  $a(g) := f(x, g)$ ,  $b_n(g) := F_{c_1, c_2}(x_n, g)$ ,  $b(g) := F_{c_1, c_2}(x, g)$  and  $c_n := a_n b_n$ ,  $c := ab$ . Then we must check that

$$\lim_{n \rightarrow \infty} |c - c_n|_{p, \psi, x} := \lim_{n \rightarrow \infty} \left( \sum_{g \in X_H} |c(g) - c_n(g)|^p \psi(x, g) \right)^{1/p} = 0.$$

Using the triangle inequality we have

$$|c - c_n|_{p, \psi, x} \leq |(a - a_n)b|_{p, \psi, x} + |a_n(b - b_n)|_{p, \psi, x} := I + II.$$

According to (6.8) for any  $\epsilon > 0$  we can decompose  $b$  in the sum  $b' + b''$  where  $b' = 0$  outside a finite subset  $S_\epsilon \subset X_H$  and  $b'' = 0$  on  $S_\epsilon$  such that  $|b''(g)| < \epsilon$  for all  $g$ . Note also that  $|b'(g)| \leq 1$  for all  $g$ . Also, (6.7) and uniform continuity of  $\log \psi$  on the compact set  $bD$  imply that  $|a - a_n|_{p, \psi, x} \leq km$  for some  $k > 0$ . Finally, by continuity of  $f$  on  $bD'$  we can find a number  $N$  such that for any  $n \geq N$  we have  $|(a - a_n)\chi_\epsilon|_{p, \psi, x} < \epsilon$ , where  $\chi_\epsilon$  is the characteristic function of  $S_\epsilon$ . Using all these facts we get for  $n \geq N$

$$I \leq |(a - a_n)\chi_\epsilon b'|_{p, \psi, x} + |(a - a_n)b''|_{p, \psi, x} \leq \epsilon + km\epsilon = (1 + km)\epsilon.$$

To estimate  $II$  observe that from (6.7) and uniform continuity of  $\log \psi$  follow that for each  $g \in X_H$

$$(|a_n(g)|^p \psi(x, g))^{1/p} \leq |a_n|_{p, \psi, x} \leq k'|a_n|_{p, \psi, x_n} \leq k'm$$

(for some  $k'$ ). Moreover, since  $F_{c_1, c_2}|_{bD'} \in C_{p,1}(bD')$ , there is an integer  $N'$  such that for any  $n \geq N'$  we have  $|b - b_n|_{p, 1, x} < \epsilon$ . These two inequalities yield for  $n \geq N'$

$$II \leq \sup_{g \in X_H} (|a_n(g)|^p \psi(x, g))^{1/p} |b - b_n|_{p, 1, x} \leq k'm\epsilon.$$

Combining the estimates for  $I$  and  $II$  we obtain that

$$\lim_{n \rightarrow \infty} |c - c_n|_{p, \psi, x} = 0.$$

This is equivalent to  $C_{p,\psi}$ -continuity of  $f_{c_1,c_2}$  over  $x$ .

Similarly one can check that if  $p = \infty$  then  $f_{c_1,c_2}$  belongs to  $C_{1,1}(bD')$ . We leave it as an exercise to the readers.  $\square$

Let us finish the proof of the theorem. According to Lemma 6.3 and the case (A) there is a function  $f'_{c_1,c_2} \in \mathcal{O}(D') \cap C(\overline{D'})$  such that  $f'_{c_1,c_2}|_{bD'} = f_{c_1,c_2}$ . Note also that since  $|F_{c_1,c_2}| \leq 1$  inequality (6.6) yields

$$|f'_{c_1,c_2}|_{p,\psi}^D \leq c \sup_{z \in bD} |f|_{p,\psi,z} := cm .$$

Consider  $f' := f_{c_1,c_2}/F_{c_1,c_2}$ . Then  $f' \in \mathcal{O}(D') \cap C(\overline{D'})$  and  $f'|_{bD'} = f$ . The uniqueness property for holomorphic functions implies that  $f'$  does not depend on  $c_1$  and  $c_2$ . Since  $F_{c_1,c_2}$  converges uniformly on compact subsets of  $M'$  to 1 as  $c_2 \rightarrow 0$  from the last inequality we get

$$|f'|_{p,\psi}^D \leq c \sup_{z \in bD} |f|_{p,\psi,z} .$$

Therefore  $f' \in \mathcal{H}_{p,\psi}(D') \cap C(\overline{D'})$ .

The proof of the theorem for  $s = 0$  is complete. If, in addition,  $f \in C^s(bD)$  for  $1 \leq s \leq k$ , then in fact the extended function  $f' \in C^s(\overline{D'})$  (see, e.g., Theorem 3.14 in [Ha], and the discussion that follows it).  $\square$

**Proof of Corollary 2.9.** Let us consider the function  $F_{c_1,c_2}$  from (6.8). Suppose that  $f \in \mathcal{O}(U')$  satisfies the hypotheses of Corollary 2.9 with  $c_1, c_2$  and  $c$  as in the definition of  $F_{c_1,c_2}$ . Then  $f_{c_1,c_2} := fF_{c_1,c_2} \in H^\infty(U')$  with the norm bounded by 1. Further, by the hypotheses we can find a connected  $C^\infty$  compact submanifold  $bS \subset U$  that bounds a domain  $S$  containing  $K$ . By Theorem 2.7 (applied to  $S' := r^{-1}(S)$  and  $bS' := r^{-1}(bS)$ ) the function  $f_{c_1,c_2}$  admits an extension  $f'_{c_1,c_2} \in H^\infty(D')$ . Since  $f'_{c_1,c_2} = f_{c_1,c_2}$  on  $U'$  and  $h(z) := \sup_{y \in r^{-1}(z)} |f'_{c_1,c_2}(y)|$  is a continuous plurisubharmonic function on  $D$ ,

$$|f'_{c_1,c_2}|_\infty^D = \sup_{z \in U'} |f(z)| \leq 1 .$$

Then the function  $f' := f'_{c_1,c_2}/F_{c_1,c_2} \in \mathcal{O}(D')$  extends  $f$  and satisfies

$$|f'(z)| \leq e^{c_3 e^{c_1 d_0(z)}} , \quad z \in D' ,$$

with  $c_3$  as in (6.8).  $\square$

## 7. Proof of Theorem 2.10.

Let  $r : M_G \rightarrow M$  be the regular covering of  $M$  satisfying condition (1.1) (for some  $\widetilde{M}$  and  $N$ ) with transformation group  $G$ . Let  $G_1 \subset G$  be a subgroup of a finite index and let  $r_1 : M_1 \rightarrow M$  be the covering with fibre  $G/G_1$ . Then there are coverings  $N_1$  and  $\widetilde{M}_1$  of  $N$  and  $\widetilde{M}$  with fibre  $G/G_1$  such that  $M_1 \subset \subset \widetilde{M}_1 \subset N_1$ . Clearly this triple satisfies condition (1.1), as well. Thus without loss of generality we may assume that  $M := M_1$ ,  $G := G_1$ ,  $\widetilde{M} := \widetilde{M}_1$  and  $N := N_1$ , and so  $G \in \mathcal{T}(M)$ . The latter means that  $G$  admits a linear representation  $\rho$  into  $GL_k(\mathbb{C})$  and that the flat vector bundle  $E_\rho$  on  $N$  associated with  $\rho$  is topologically trivial. Then the

restriction  $E_\rho|_{\tilde{M}}$  is topologically trivial. Since  $\tilde{M}$  is Stein, according to the Oka-Grauert principle (see [G2]),  $E_\rho|_{\tilde{M}}$  is holomorphically trivial. In particular,  $\rho$  can be obtained as the monodromy of an equation  $dF = \omega F$  on  $\tilde{M}$  where  $\omega$  is a matrix-valued holomorphic 1-form on  $\tilde{M}$  satisfying  $d\omega - \omega \wedge \omega = 0$ . Let  $\tilde{\omega} = r^*\omega$  be the pullback of  $\omega$  on  $\tilde{M}_G = r^{-1}(\tilde{M})$ . (Here  $r : N_G \rightarrow N$  is the regular covering of  $N$  with the transformation group  $G$  so that  $M_G$  and  $\tilde{M}_G$  are domains in  $N_G$ .) Then there exists a function  $\tilde{F} \in \mathcal{O}(\tilde{M}_G, GL_k(\mathbb{C}))$  such that  $d\tilde{F} = \tilde{\omega}\tilde{F}$ . This follows from the fact that the monodromy of the last equation is the restriction of  $\rho$  to  $\pi_1(M_G)$  and so it is trivial (since  $\pi_1(M_G) \subset Ker \rho$ ). Note that  $\tilde{F}$  can be obtained by Picard iteration applied to  $\tilde{\omega}$ . Since  $\overline{M}$  is a compact subset of  $\tilde{M}$  (and so  $\omega|_M$  is bounded), the Picard iteration produces for some positive  $c = c(M, \omega)$  the estimate

$$\|\tilde{F}(z)\|_2 \leq e^{cd_o(z)}, \quad z \in M_G, \quad (7.1)$$

where  $\|\cdot\|_2$  is the  $l_2$ -norm on  $GL_k(\mathbb{C})$  and  $o \in M_G$ . (Here as before  $d_o$  is the distance from  $o$  in the path metric induced by a Riemannian metric pulled back from  $N$ .) Moreover, for every  $z \in \tilde{M}_G$  there exists a matrix  $C_z \in GL_k(\mathbb{C})$  such that  $\tilde{F}(gz) = C_z^{-1}\rho(g)\tilde{F}(z)$  for any  $g \in G$ . These are standard facts of the theory of flat connections. In particular, from the last identity we derive easily that  $\tilde{F}$  separates points in every orbit of the action of  $G$  on  $\tilde{M}_G$ .

Next, let  $f$  be the function from Corollary 2.2. Then by (7.1) we get (for some  $c_1 = c_1(f, \alpha)$ )

$$\|e^{-\alpha f(z)}\tilde{F}(z)\|_2 \leq c_1 e^{(c-\alpha)d_o(z)}, \quad z \in M_G.$$

From here arguing as in the proof of Corollary 2.2 we deduce that for a sufficiently big  $\alpha$  all entries of the matrix  $e^{-\alpha f} \cdot \tilde{F}$  belong to  $\mathcal{H}_{2,1}(M_G) \cap C(\overline{M}_G)$ . Now the family consisting of these entries and the function  $e^{-\alpha f}$  separate all points in any orbit of the action of  $G$  on  $\tilde{M}_G$ . For otherwise, there are  $x, y \in \tilde{M}_G$ ,  $y = gx$ ,  $g \neq 1$ ,  $g \in G$ , such that  $e^{-\alpha f(x)}\tilde{F}(x) = e^{-\alpha f(y)}\tilde{F}(y)$  and  $e^{-\alpha f(x)} = e^{-\alpha f(y)}$ . But this implies that  $\tilde{F}(x) = \tilde{F}(y)$ , a contradiction. Finally, since  $M \subset \subset \tilde{M}$  and  $\tilde{M}$  is Stein, by the Remmert embedding theorem there are holomorphic functions  $h_1, \dots, h_l$  from  $H^\infty(M) \cap C(\overline{M})$  that separate all points in  $\overline{M}$ . We set  $\tilde{h}_i = e^{-\alpha f} r^* h_i$ ,  $1 \leq i \leq l$ . Then by the definition  $\tilde{h}_i \in \mathcal{H}_{2,1}(M_G) \cap C(\overline{M}_G)$  and so the family consisting of all  $\tilde{h}_i$ , entries of  $e^{-\alpha f}\tilde{F}$  and  $e^{-\alpha f}$  separates all points in  $\overline{M}_G$ .

The proof of Theorem 2.10 is complete.  $\square$

## References

- [B] L. Bungart, On analytic fibre bundles I. Holomorphic fibre bundles with infinite dimensional fibres. *Topology*, **7** (1) (1968), 55-68.
- [BH] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*. Springer, 1999.
- [Bo] S. Bochner, Analytic and meromorphic continuation by means of Green's formula. *Ann. of Math.*, **44** (1943), 652-673.

- [Br1] A. Brudnyi, Integral representations of holomorphic functions on coverings of pseudoconvex domains in Stein manifolds. Preprint (2005), 18 pp.
- [Br2] A. Brudnyi, Projections in the space  $H^\infty$  and the corona theorem for subdomains of coverings of finite bordered Riemann surfaces. Ark. Mat., **42** (2004), 31-59.
- [C] H. Cartan, Sur les fonctions de plusieurs variables complexes. Les espaces analytiques. Proc. Intern. Congress Mathematicians Edinburgh 1958, Cambridge Univ. Press, 1960, 33-52.
- [G1] H. Grauert, On Levi's problem and the imbedding of real-analytic manifolds. Ann. of Math., **68** (1958), 460-472.
- [G2] H. Grauert, Analytische Faserungen über holomorph-vollständigen Räumen. Math. Ann., **135**, no. 3 (1958), 263-273.
- [GHS] M. Gromov, G. Henkin and M. Shubin, Holomorphic  $L^2$  functions on coverings of pseudoconvex manifolds. GAFA, Vol. 8 (1998), 552-585.
- [GR] H. Grauert and R. Remmert, Theorie der Steinschen Räume. Springer, New York, 1977.
- [Ha] R. Harvey, Holomorphic chains and their boundaries. Several complex variables (Proc. Sympos. Pure Math., Vol. XXX, Part 1, Williams Coll., Williamstown, Mass., 1975), 309-382. Amer. Math. Soc., Providence, R. I., 1977.
- [He] D. Heunemann, Uniform estimates for the  $\bar{\partial}$ -equation on pseudoconvex polyhedra on Stein manifolds. Math. Nachr., **114** (1983), 191-196.
- [Hi] F. Hirzebruch, Topological methods in Algebraic Geometry. Springer-Verlag, New York, 1966.
- [Hu] S.-T. Hu, Homotopy theory. Academic Press, new York, 1959.
- [KR] J. J. Kohn and H. Rossi, On the extension of holomorphic functions from the boundary of a complex manifold. Ann. of Math. (2), **81** (1965), 451-472.
- [O] A. L. Onishchik, Certain concepts and applications of non-Abelian cohomology theory. Trudy Moskov. Mat. Obshch., **17** (1967), 45-88.
- [Ra] M. S. Raghunathan, Discrete subgroups of Lie groups. Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 68, Springer Verlag, New York, 1972.
- [R] R. Remmert, Sur les espaces analytiques holomorphiquement séparables et holomorphiquement convexes. C. R. Acad. Sci. Paris, **243** (1956), 118-121.
- [SH] A. G. Sergeev and G. M. Henkin, Uniform estimates of the solutions of the  $\bar{\partial}$ -equation in pseudoconvex polyhedra. (Russian) Mat. Sb. (N.S.) **112 (154)** (1980), no. 4 (8), 522-567.