

CHERN OBSTRUCTIONS FOR COLLECTIONS OF 1-FORMS ON SINGULAR VARIETIES

W. EBELING AND S. M. GUSEIN-ZADE

Dedicated to Jean-Paul Brasselet on the occasion of his 60th birthday

ABSTRACT. We introduce a certain index of a collection of germs of 1-forms on a germ of a singular variety which is a generalization of the local Euler obstruction corresponding to Chern numbers different from the top one.

INTRODUCTION

The aim of this paper is to bring together some ideas of [3] and [6]. A germ of a vector field or of a 1-form on the complex affine space \mathbb{C}^n at the origin not vanishing in a punctured neighbourhood of it has a topological invariant — the Poincaré–Hopf index. The sum of the Poincaré–Hopf indices of the singular points of a vector field on a compact complex manifold is equal to the Euler characteristic of the manifold. There are several generalizations of this notion to vector fields and/or to 1-forms on complex analytic varieties with singularities (isolated or not) started by M.-H. Schwartz: [11, 4, 8, 1, 12, 5, 3, ...]. For the case of an isolated complete intersection singularity there is defined an index which is sometimes called the GSV index: [8, 12, 5]. Another generalization which makes sense not only for isolated complete intersection singularities and also not only for varieties with isolated singularities is the so called local Euler obstruction: [2, 3] (its analogue for 1-forms was considered in [7]). One can say that in some sense all these invariants correspond to the Euler characteristic, which, for a compact complex analytic manifold M^n , coincides with the top Chern number $\langle c_n(M), [M] \rangle$.

A generalization of the GSV-index corresponding to other Chern numbers (different from the top one) was introduced and studied in [6]. It is defined for a collection of germs of 1-forms on an isolated

1991 *Mathematics Subject Classification.* 14B05, 58A10, 55S99.

Key words and phrases. singular variety, 1-form, index.

Partially supported by the DFG-programme "Global methods in complex geometry" (Eb 102/4–3), grants RFBR–04–01–00762, NSh–1972.2003.1.

complete intersection singularity. For a collection of 1-forms on a projective complex complete intersection with isolated singularities, the sum of these indices of the singular points is equal to plus-minus the corresponding Chern number of a smoothing of the variety.

Here we define and study an index of a collection of germs of 1-forms on a germ of a singular variety which is an analogue of the local Euler obstruction corresponding to a Chern number different from the top one.

1. SPECIAL POINTS OF 1-FORMS

Let $(X^n, 0) \subset (\mathbb{C}^N, 0)$ be the germ of a purely n -dimensional reduced complex analytic variety at the origin (generally speaking with a non-isolated singularity). Let $\mathbf{k} = \{k_i\}$, $i = 1, \dots, s$, be a fixed partition of n (i.e., k_i are positive integers, $\sum_{i=1}^s k_i = n$). Let $\{\omega_j^{(i)}\}$ ($i = 1, \dots, s$, $j = 1, \dots, n - k_i + 1$) be a collection of germs of 1-forms on $(\mathbb{C}^N, 0)$ (not necessarily complex analytic; it suffices that the forms $\omega_j^{(i)}$ are complex linear functions continuously depending on a point of \mathbb{C}^N). Let $\varepsilon > 0$ be small enough so that there is a representative X of the germ $(X, 0)$ and representatives $\omega_j^{(i)}$ of the germs of 1-forms inside the ball $B_\varepsilon(0) \subset \mathbb{C}^N$.

Definition: A point $P \in X$ is called a *special* point of the collection $\{\omega_j^{(i)}\}$ of 1-forms on the variety X if there exists a sequence $\{P_m\}$ of points from the non-singular part X_{reg} of the variety X converging to P such that the sequence $T_{P_m}X_{\text{reg}}$ of the tangent spaces at the points P_m has a limit L as $m \rightarrow \infty$ (in the Grassmann manifold of n -dimensional vector subspaces of \mathbb{C}^N) and the restrictions of the 1-forms $\omega_1^{(i)}, \dots, \omega_{n-k_i+1}^{(i)}$ to the subspace $L \subset T_P\mathbb{C}^N$ are linearly dependent for each $i = 1, \dots, s$.

Definition: The collection $\{\omega_j^{(i)}\}$ of 1-forms has an *isolated special point* on the germ $(X, 0)$ if it has no special points on X in a punctured neighbourhood of the origin.

Remarks. 1. If the 1-forms $\omega_j^{(i)}$ are complex analytic, the property to have an isolated special point is a condition on the classes of these 1-forms in the module

$$\Omega_{X,0}^1 = \Omega_{\mathbb{C}^N,0}^1 / \{f \cdot \Omega_{\mathbb{C}^N,0}^1 + df \cdot \mathcal{O}_{\mathbb{C}^N,0} \mid f \in \mathcal{I}_X\}$$

of germs of 1-forms on the variety X (\mathcal{I}_X is the ideal of germs of holomorphic functions vanishing on X).

2. For the case $s = 1$ (and therefore $k_1 = n$), i.e. for one 1-form ω , there exists a notion of a *singular* point of the 1-form ω on X (see, e.g., [7]). It is defined in terms of a Whitney stratification of the variety X . A point $x \in X$ is a *singular* point of the 1-form ω on the variety X if the restriction of the 1-form ω to the stratum of X containing x is equal to zero at the point x . (One should consider points of all zero-dimensional strata as singular ones.) One can easily see that a special point of the 1-form ω on the variety X is singular, but not vice versa. (E.g. the origin is a singular point of the 1-form dx on the cone $\{x^2 + y^2 + z^2 = 0\}$, but not a special one.) On a smooth variety these two notions coincide.

The notion of a *non-degenerate* special (singular) point of a collection of germs of 1-forms on a smooth variety was introduced in [6]. The index of a non-degenerate point of a collection of germs of holomorphic 1-forms is equal to 1.

Let

$$\mathcal{L}^{\mathbf{k}} = \prod_{i=1}^s \prod_{j=1}^{n-k_i+1} \mathbb{C}_{ij}^{N^*}$$

be the space of collections of linear functions on \mathbb{C}^N (i.e. of 1-forms with constant coefficients).

Proposition 1. *There exists an open and dense subset $U \subset \mathcal{L}^{\mathbf{k}}$ such that each collection $\{\ell_j^{(i)}\} \in U$ has only isolated special points on X and, moreover, all these points belong to the smooth part X_{reg} of the variety X and are non-degenerate.*

Proof. Let $Y \subset X \times \mathcal{L}^{\mathbf{k}}$ be the closure of the set of pairs $(x, \{\ell_j^{(i)}\})$ where $x \in X_{\text{reg}}$ and the restrictions of the linear functions $\ell_1^{(i)}, \dots, \ell_{n-k_i+1}^{(i)}$ to the tangent space $T_x X_{\text{reg}}$ are linearly dependent for each $i = 1, \dots, s$. Let $\pi : Y \rightarrow \mathcal{L}^{\mathbf{k}}$ be the projection to the second factor. One has $\text{codim } Y = \sum_{i=1}^s k_i = n$ and therefore $\dim Y = \dim \mathcal{L}^{\mathbf{k}}$. Moreover, $Y \setminus (X_{\text{reg}} \times \mathcal{L}^{\mathbf{k}})$ is a proper subvariety of Y and therefore its dimension is strictly smaller than $\dim \mathcal{L}^{\mathbf{k}}$. A generic point of the space $\mathcal{L}^{\mathbf{k}}$ is a regular value of the map π which means that it has only finitely many preimages, all of them belong to $X_{\text{reg}} \times \mathcal{L}^{\mathbf{k}}$ and the map π is non-degenerate at them. This implies the statement. \square

Corollary 1. *Let $\{\omega_j^{(i)}\}$ be a collection of 1-forms on X with an isolated special point at the origin. Then there exists a deformation $\{\tilde{\omega}_j^{(i)}\}$ of the collection $\{\omega_j^{(i)}\}$ whose special points lie in X_{reg} and are non-degenerate.*

Moreover, as such a deformation one can use $\{\omega_j^{(i)} + \lambda \ell_j^{(i)}\}$ with a generic collection $\{\ell_j^{(i)}\} \in \mathcal{L}^{\mathbf{k}}$, $\lambda \neq 0$ small enough.

Corollary 2. *The set of collections of holomorphic 1-forms with a non-isolated special point at the origin has infinite codimension in the space of all holomorphic collections.*

2. LOCAL CHERN OBSTRUCTIONS

Let $\{\omega_j^{(i)}\}$ be a collection of germs of 1-forms on $(X, 0)$ with an isolated special point at the origin. Let $\nu : \widehat{X} \rightarrow X$ be the Nash transformation of the variety $X \subset B_\varepsilon(0)$ defined as follows. Let $G(n, N)$ be the Grassmann manifold of n -dimensional vector subspaces of \mathbb{C}^N . There is a natural map $\sigma : X_{\text{reg}} \rightarrow B_\varepsilon(0) \times G(n, N)$ which sends a point $x \in X_{\text{reg}}$ to $(x, T_x X_{\text{reg}})$. The Nash transform \widehat{X} of the variety X is the closure of the image $\text{Im } \sigma$ of the map σ in $B_\varepsilon(0) \times G(n, N)$, ν is the natural projection. The Nash bundle \widehat{T} over \widehat{X} is a vector bundle of rank n which is the pullback of the tautological bundle on the Grassmann manifold $G(n, N)$. There is a natural lifting of the Nash transformation to a bundle map from the Nash bundle \widehat{T} to the restriction of the tangent bundle $T\mathbb{C}^N$ of \mathbb{C}^N to X . This is an isomorphism of \widehat{T} and $TX_{\text{reg}} \subset T\mathbb{C}^N$ over the non-singular part X_{reg} of X .

The collection of 1-forms $\{\omega_j^{(i)}\}$ gives rise to a section $\widehat{\omega}$ of the bundle

$$\widehat{\mathbb{T}} = \bigoplus_{i=1}^s \bigoplus_{j=1}^{n-k_i+1} \widehat{T}_{i,j}^*$$

where $\widehat{T}_{i,j}^*$ are copies of the dual Nash bundle \widehat{T}^* over the Nash transform \widehat{X} numbered by indices i and j . Let $\widehat{\mathbb{D}} \subset \widehat{\mathbb{T}}$ be the set of pairs $(x, \{\alpha_j^{(i)}\})$ where $x \in \widehat{X}$ and the collection $\{\alpha_j^{(i)}\}$ of elements of \widehat{T}_x^* (i.e. of linear functions on \widehat{T}_x) is such that $\alpha_1^{(i)}, \dots, \alpha_{n-k_i+1}^{(i)}$ are linearly dependent for each $i = 1, \dots, s$. The image of the section $\widehat{\omega}$ does not intersect $\widehat{\mathbb{D}}$ outside of the preimage $\nu^{-1}(0) \subset \widehat{X}$ of the origin. The map $\widehat{\mathbb{T}} \setminus \widehat{\mathbb{D}} \rightarrow \widehat{X}$ is a fibre bundle. The fibre $W_x = \widehat{\mathbb{T}}_x \setminus \widehat{\mathbb{D}}_x$ of it is $(2n-2)$ -connected, its homology group $H_{2n-1}(W_x; \mathbb{Z})$ is isomorphic to \mathbb{Z} and has a natural generator: see, e.g., [6]. The latter fact implies that the fibre bundle $\widehat{\mathbb{T}} \setminus \widehat{\mathbb{D}} \rightarrow \widehat{X}$ is homotopically simple in dimension $2n-1$, i.e. the fundamental group $\pi_1(\widehat{X})$ of the base acts trivially on the homotopy group $\pi_{2n-1}(W_x)$ of the fibre, the last one being isomorphic to the homology group $H_{2n-1}(W_x; \mathbb{Z})$: see, e.g., [14].

Definition: The *local Chern obstruction* $\text{Ch}_{X,0} \{\omega_j^{(i)}\}$ of the collections of germs of 1-forms $\{\omega_j^{(i)}\}$ on $(X, 0)$ at the origin is the (primary, and in fact the only) obstruction to extend the section $\widehat{\omega}$ of the fibre bundle $\widehat{\mathbb{T}} \setminus \widehat{\mathbb{D}} \rightarrow \widehat{X}$ from the preimage of a neighbourhood of the sphere $S_\varepsilon = \partial B_\varepsilon$ to \widehat{X} , more precisely its value (as an element of $H^{2n}(\nu^{-1}(X \cap B_\varepsilon), \nu^{-1}(X \cap S_\varepsilon); \mathbb{Z})$) on the fundamental class of the pair $(\nu^{-1}(X \cap B_\varepsilon), \nu^{-1}(X \cap S_\varepsilon))$.

The definition of the local Chern obstruction $\text{Ch}_{X,0} \{\omega_j^{(i)}\}$ can be reformulated in the following way. Let $\mathcal{D}_X^{\mathbf{k}} \subset \mathbb{C}^N \times \mathcal{L}^{\mathbf{k}}$ be the closure of the set of pairs $(x, \{\ell_j^{(i)}\})$ such that $x \in X_{\text{reg}}$ and the restrictions of the linear functions $\ell_1^{(i)}, \dots, \ell_{n-k_i+1}^{(i)}$ to $T_x X_{\text{reg}} \subset \mathbb{C}^N$ are linearly dependent for each $i = 1, \dots, s$. (For $s = 1$, $\mathbf{k} = \{n\}$, $\mathcal{D}_X^{\mathbf{k}}$ is the (non-projectivized) conormal space of X [15].) The collection $\{\omega_j^{(i)}\}$ of germs of 1-forms on $(\mathbb{C}^N, 0)$ defines a section $\tilde{\omega}$ of the (trivial) fibre bundle $\mathbb{C}^N \times \mathcal{L}^{\mathbf{k}} \rightarrow \mathbb{C}^N$. Then

$$\text{Ch}_{X,0} \{\omega_j^{(i)}\} = (\tilde{\omega}(\mathbb{C}^N) \circ \mathcal{D}_X^{\mathbf{k}})_0$$

where $(\cdot \circ \cdot)_0$ is the intersection number at the origin in $\mathbb{C}^N \times \mathcal{L}^{\mathbf{k}}$. This description can be considered as a generalization of an expression of the local Euler obstruction as a microlocal intersection number defined in [9], see also [10, Sections 5.0.3 and 5.2.1].

Remarks. 1. On a smooth manifold X the local Chern obstruction $\text{Ch}_{X,0} \{\omega_j^{(i)}\}$ coincides with the index $\text{ind}_{X,0} \{\omega_j^{(i)}\}$ of the collection $\{\omega_j^{(i)}\}$ defined in [6].

2. The local Euler obstruction is defined for vector fields as well as for 1-forms. One can see that vector fields are not well adapted to a definition of the local Chern obstruction. A more or less direct version of the definition above for vector fields demands to consider vector fields on a singular variety $X \subset \mathbb{C}^N$ to be sections $v = v(x)$ of $T\mathbb{C}^N|_X$ such that $v(x) \in T_x X \subset T_x \mathbb{C}^N$ ($\dim T_x X$ is not constant). (Traditionally vector fields tangent to smooth strata of the variety X are considered.) There exist only continuous (non-trivial, i.e. with $s > 1$) collections of such vector fields "on X " with isolated special points, but not holomorphic ones.

3. The definition of the local Chern obstruction $\text{Ch}_{X,0} \{\omega_j^{(i)}\}$ may also be formulated in terms of a collection $\{\omega_j^{(i)}\}$ of germs of 1-forms with values in vector spaces L_i of dimensions $n - k_i + 1$. Therefore (via differentials) it is also defined for a collection $\{f^{(i)}\}$ of germs of maps

$f^{(i)} : (\mathbb{C}^N, 0) \rightarrow (\mathbb{C}^{n-k_i+1}, 0)$ (just as the Euler obstruction is defined for a germ of a function).

Being a (primary) obstruction, the local Chern obstruction satisfies the law of conservation of number, i.e. if a collection of 1-forms $\{\tilde{\omega}_j^{(i)}\}$ is a deformation of the collection $\{\omega_j^{(i)}\}$ and has isolated special points on X , then

$$\text{Ch}_{X,0} \{\omega_j^{(i)}\} = \sum \text{Ch}_{X,Q} \{\tilde{\omega}_j^{(i)}\}$$

where the sum on the right hand side is over all special points Q of the collection $\{\tilde{\omega}_j^{(i)}\}$ on X in a neighbourhood of the origin. With Corollary 1 this implies the following statements.

Proposition 2. *The local Chern obstruction $\text{Ch}_{X,0} \{\omega_j^{(i)}\}$ of a collection $\{\omega_j^{(i)}\}$ of germs of holomorphic 1-forms is equal to the number of special points on X of a generic (holomorphic) deformation of the collection.*

This statement is an analogue of Proposition 2.3 in [13].

Proposition 3. *If a collection $\{\omega_j^{(i)}\}$ of 1-forms on a compact (say, projective) variety X has only isolated special points, then the sum of the local Chern obstructions of the collection $\{\omega_j^{(i)}\}$ at these points does not depend on the collection and therefore is an invariant of the variety.*

It is reasonable to consider this sum as $((-1)^n$ times) the corresponding Chern number of the singular variety X .

Let $(X, 0)$ be an isolated complete intersection singularity. As it was mentioned above, a collection of germs of 1-forms $\{\omega_j^{(i)}\}$ on $(X, 0)$ with an isolated special point at the origin has an index $\text{ind}_{X,0} \{\omega_j^{(i)}\}$ which is an analogue of the GSV-index of a vector field: [5]. The fact that both the Chern obstruction and the index satisfy the law of conservation of number and they coincide on a smooth manifold yields the following statement.

Proposition 4. *For a collection $\{\omega_j^{(i)}\}$ of germs of 1-forms on an isolated complete intersection singularity $(X, 0)$ the difference*

$$\text{ind}_{X,0} \{\omega_j^{(i)}\} - \text{Ch}_{X,0} \{\omega_j^{(i)}\}$$

does not depend on the collection and therefore is an invariant of the germ of the variety.

Since, by Proposition 1, $\text{Ch}_{X,0} \{\ell_j^{(i)}\} = 0$ for a generic collection $\{\ell_j^{(i)}\}$ of linear functions on \mathbb{C}^N , one has the following statement.

Corollary 3. *One has*

$$\mathrm{Ch}_{X,0} \{\omega_j^{(i)}\} = \mathrm{ind}_{X,0} \{\omega_j^{(i)}\} - \mathrm{ind}_{X,0} \{\ell_j^{(i)}\}$$

for a generic collection $\{\ell_j^{(i)}\}$ of linear functions on \mathbb{C}^N .

REFERENCES

- [1] Ch. Bonatti, X. Gómez-Mont: The index of holomorphic vector fields on singular varieties I. *Astérisque* **222**, 9–35 (1994).
- [2] J.-P. Brasselet, Lê Dũng Tráng, J. Seade: Euler obstruction and indices of vector fields. *Topology* **39**, 1193–1208 (2000).
- [3] J.-P. Brasselet, D. Massey, A. J. Parameswaran, J. Seade: Euler obstruction and defects of functions on singular varieties. *J. London Math. Soc. (2)* **70**, 59–76 (2004).
- [4] J.-P. Brasselet, M.-H. Schwartz: Sur les classes de Chern d'un ensemble analytique complexe. In: *Caractéristique d'Euler-Poincaré*, *Astérisque* **82–83**, 93–147 (1981).
- [5] W. Ebeling, S. M. Gusein-Zade: Indices of 1-forms on an isolated complete intersection singularity. *Moscow Math. J.* **3**, 439–455 (2003).
- [6] W. Ebeling, S. M. Gusein-Zade: Indices of vector fields or 1-forms and characteristic numbers. *math.AG/0303330*. *Bull. London Math. Soc.* (to appear).
- [7] W. Ebeling, S. M. Gusein-Zade: Radial index and Euler obstruction of a 1-form on a singular variety. *math.AG/0402388*. *Geom. Dedicata* (to appear).
- [8] X. Gómez-Mont, J. Seade, A. Verjovsky: The index of a holomorphic flow with an isolated singularity. *Math. Ann.* **291**, 737–751 (1991).
- [9] M. Kashiwara, P. Schapira: *Sheaves on Manifolds*. Springer-Verlag, 1990.
- [10] J. Schürmann: *Topology of Singular Spaces and Constructible Sheaves*. Birkhäuser, 2003.
- [11] M.-H. Schwartz: Classes caractéristiques définies par une stratification d'une variété analytique complexe. *C. R. Acad. Sci. Paris Sér. I Math.* **260**, 3262–3264, 3535–3537 (1965).
- [12] J. A. Seade, T. Suwa: A residue formula for the index of a holomorphic flow. *Math. Ann.* **304**, 621–634 (1996).
- [13] J. Seade, M. Tibăr, A. Verjovsky: Milnor numbers and Euler obstruction. *math.CV/0311109*.
- [14] N. Steenrod: *The Topology of Fibre Bundles*. Princeton Math. Series, Vol. **14**, Princeton University Press, Princeton, N. J., 1951.
- [15] B. Teissier: Variétés polaires. II. Multiplicités polaires, sections planes, et conditions de Whitney. In: *Algebraic geometry (La Rábida, 1981)*, *Lecture Notes in Math.*, Vol. **961**, Springer, Berlin, 1982, pp. 314–491.

UNIVERSITÄT HANNOVER, INSTITUT FÜR MATHEMATIK, POSTFACH 6009, D-30060 HANNOVER, GERMANY.

E-mail address: `ebeling@math.uni-hannover.de`

MOSCOW STATE UNIVERSITY, FACULTY OF MECHANICS AND MATHEMATICS, MOSCOW, 119992, RUSSIA.

E-mail address: `sabir@mccme.ru`