

A note on Primes in Short Intervals

Tsz Ho Chan

August 17, 2018

Abstract

Instead of a strong quantitative form of the Hardy-Littlewood prime k -tuple conjecture, one can assume an average form of it and still obtains the same distribution result on $\psi(x+h) - \psi(x)$ by Montgomery and Soundararajan [1].

1 Introduction

Let $\Lambda(n)$ be von Mangoldt lambda function, $\mu(n)$ be the Möbius function and $\phi(n)$ be the Euler's phi function. Let $e(\theta) = e^{2\pi i\theta}$, $\epsilon > 0$ and $\psi(x) = \sum_{n \leq x} \Lambda(n)$. In [1], Montgomery and Soundararajan studied the distribution of primes in short intervals

$$M_K(N; H) := \sum_{n=1}^N (\psi(n+H) - \psi(n) - H)^K \quad (1)$$

under a strong quantitative form of the Hardy-Littlewood prime k -tuple conjecture:

Conjecture 1.

$$\sum_{n \leq x} \prod_{i=1}^k \Lambda(n + d_i) = \mathfrak{S}(\mathcal{D})x + E_k(x; \mathcal{D}) \quad (2)$$

holds with

$$E_k(x; \mathcal{D}) \ll_{\epsilon, K} N^{1/2+\epsilon}$$

uniformly for $1 \leq k \leq K$, $0 \leq x \leq N$, and distinct d_i satisfying $1 \leq d_i \leq H$. Here $\mathcal{D} = \{d_1, d_2, \dots, d_k\}$, and

$$\mathfrak{S}(\mathcal{D}) = \sum_{\substack{q_1, \dots, q_k \\ 1 \leq q_i < \infty}} \left(\prod_{i=1}^k \frac{\mu(q_i)}{\phi(q_i)} \right) \sum_{\substack{a_1, \dots, a_k \\ 1 \leq a_i \leq q_i \\ (a_i, q_i) = 1 \\ \sum a_i / q_i \in \mathbb{Z}}} e\left(\sum_{i=1}^k \frac{a_i d_i}{q_i} \right)$$

is the singular series as in equation (2) of [1].

They proved

Theorem 1. *Under Conjecture 1,*

$$M_K(N; H) = \mu_k H^{K/2} \int_1^N (\log x/H + B)^{K/2} dx \\ + O\left(N(\log N)^{K/2} H^{K/2} \left(\frac{H}{\log N}\right)^{-1/(8K)} + H^K N^{1/2+\epsilon}\right)$$

uniformly for $\log N \leq H \leq N^{1/K}$, where $\mu_k = 1 \cdot 3 \cdots (k-1)$ if k is even, and $\mu_k = 0$ if k is odd; $B = 1 - C_0 - \log 2\pi$ and C_0 denotes Euler's constant.

The first author of [1] suggested to the present author that Theorem 1 is probably true under an average form of the Hardy-Littlewood prime k -tuple conjecture, namely:

Conjecture 2. *For $x \geq H$,*

$$\sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} E_k(x, \mathcal{D})^2 \ll_k x^{1+\epsilon} H^k.$$

Our goal in this paper is to prove Theorem 1 under Conjecture 2.

Acknowledgement The author would like to thank Prof. Hugh Montgomery for suggesting the problem and the American Institute of Mathematics for provision both financially and spatially.

2 Proof of Theorem 1

Lemma 1. *With $\Lambda_0(n) = \Lambda(n) - 1$ and $|A|$ denoting the size of a set A ,*

$$\sum_{n \leq x} \prod_{i=1}^k \Lambda_0(n + d_i) = \mathfrak{S}_0(\mathcal{D})x + O\left(\sum_{\mathcal{J} \subset \{1, 2, \dots, k\}} |E_{|\mathcal{J}|}(x; \mathcal{D}_{\mathcal{J}})|\right)$$

where

$$\mathfrak{S}_0(\mathcal{D}) = \sum_{\substack{q_1, \dots, q_k \\ 1 < q_i < \infty}} \left(\prod_{i=1}^k \frac{\mu(q_i)}{\phi(q_i)} \right) \sum_{\substack{a_1, \dots, a_k \\ 1 \leq a_i \leq q_i \\ (a_i, q_i) = 1 \\ \sum a_i / q_i \in \mathbb{Z}}} e\left(\sum_{i=1}^k \frac{a_i d_i}{q_i}\right)$$

and $\mathcal{D}_{\mathcal{J}} = \{d_j\}_{j \in \mathcal{J}}$.

Proof: The left hand side

$$\begin{aligned}
&= \sum_{n \leq x} \prod_{i=1}^k (\Lambda(n + d_i) - 1) = \sum_{n \leq x} \sum_{\mathcal{J} \subset \{1, 2, \dots, k\}} (-1)^{k-|\mathcal{J}|} \prod_{i \in \mathcal{J}} \Lambda(n + d_i) \\
&= \sum_{\mathcal{J} \subset \{1, 2, \dots, k\}} (-1)^{k-|\mathcal{J}|} \sum_{n \leq x} \prod_{i \in \mathcal{J}} \Lambda(n + d_i) \\
&= \sum_{\mathcal{J} \subset \{1, 2, \dots, k\}} (-1)^{k-|\mathcal{J}|} (\mathfrak{S}(\mathcal{D}_{\mathcal{J}})x + E_{|\mathcal{J}|}(x; \mathcal{D}_{\mathcal{J}})) \\
&= \mathfrak{S}_0(\mathcal{D})x + O\left(\sum_{\mathcal{J} \subset \{1, 2, \dots, k\}} |E_{|\mathcal{J}|}(x; \mathcal{D}_{\mathcal{J}})|\right)
\end{aligned}$$

by (2) and the identity $\mathfrak{S}_0(\mathcal{D}) = \sum_{\mathcal{J} \subset \mathcal{D}} (-1)^{|\mathcal{J}|} \mathfrak{S}(\mathcal{J})$ (see equation (5) of [1]).

Proof of Theorem 1 under Conjecture 2: Following [1], we expand (1) and have

$$\begin{aligned}
M_K(N; H) &= \sum_{k=1}^K \frac{1}{k!} \sum_{\substack{1 \leq M_1, \dots, M_k \\ \sum M_i = K}} \binom{K}{M_1 \cdots M_k} \\
&\quad \times \sum_{\substack{m_1, \dots, m_k \\ 0 \leq m_i < M_i}} \prod_{i=1}^k (-1)^{M_i-1-m_i} \binom{M_i-1}{m_i} L_k(\mathbf{m}),
\end{aligned} \tag{3}$$

where

$$L_k(\mathbf{m}) := \sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} \sum_{n=1}^N \prod_{i=1}^k \Lambda_{m_i}(n + d_i) \tag{4}$$

and $\Lambda_m(n) := \Lambda(n)^m \Lambda_0(n)$. To estimate $L_k(\mathbf{m})$, one needs to distinguish between those i for which $m_i = 0$ and those for which $m_i > 0$. Following [1], we set $\mathcal{K} = \{1, \dots, k\}$ and introduce

$$\mathcal{H} = \{i \in \mathcal{K} : m_i \geq 1\}, \mathcal{J} = \{i \in \mathcal{K} : m_i = 0\}, \mathcal{J} \subset \mathcal{K}.$$

Then

$$\begin{aligned}
& \sum_{n \leq x} \prod_{i \in \mathcal{J}} \Lambda_0(n + d_i) \prod_{i \in \mathcal{H}} \Lambda(n + d_i) \\
&= \sum_{n \leq x} \sum_{\substack{\mathcal{J} \\ \mathcal{J} \subset \mathcal{J} \subset \mathcal{K}}} \prod_{i \in \mathcal{J}} \Lambda_0(n + d_i) = \sum_{\substack{\mathcal{J} \\ \mathcal{J} \subset \mathcal{J} \subset \mathcal{K}}} \sum_{n \leq x} \prod_{i \in \mathcal{J}} \Lambda_0(n + d_i) \\
&= \sum_{\substack{\mathcal{J} \\ \mathcal{J} \subset \mathcal{J} \subset \mathcal{K}}} \left[\mathfrak{S}_0(\mathcal{D}_{\mathcal{J}})x + O\left(\sum_{\mathcal{J}' \subset \mathcal{J}} |E_{|\mathcal{J}'|}(x; \mathcal{D}_{\mathcal{J}'})|\right) \right] \\
&= x \sum_{\substack{\mathcal{J} \\ \mathcal{J} \subset \mathcal{J} \subset \mathcal{K}}} \mathfrak{S}_0(\mathcal{D}_{\mathcal{J}}) + O_k\left(\sum_{\mathcal{K}' \subset \mathcal{K}} |E_{|\mathcal{K}'|}(x; \mathcal{D}_{\mathcal{K}'})|\right)
\end{aligned}$$

by Lemma 1. We write the above as $f(x) = cx + E_{x, \mathcal{K}}$. In general,

$$\begin{aligned}
& \int_{1^-}^X g(x) df(x) \\
&= g(X)f(X) - \int_{1^-}^X f(x)g'(x)dx \\
&= g(X)[cX + E_{X, \mathcal{K}}] - \int_{1^-}^X (cx + E_{x, \mathcal{K}})g'(x)dx \\
&= c \int_{1^-}^X g(x)dx + O\left(E_{X, \mathcal{K}}|g(X)| + \int_{1^-}^X E_{x, \mathcal{K}}|g'(x)|dx\right).
\end{aligned}$$

Thus, by integration by parts,

$$\begin{aligned}
& \sum_{n=1}^N \left(\prod_{i \in \mathcal{J}} \Lambda_0(n + d_i) \right) \left(\prod_{i \in \mathcal{H}} \Lambda(n + d_i) (\log(n + d_i))^{m_i - 1} (\log(n + d_i) - 1) \right) \\
&= \int_{1^-}^N \prod_{i \in \mathcal{H}} (\log(x + d_i))^{m_i - 1} (\log(x + d_i) - 1) df(x) \\
&= c \int_1^N \prod_{i \in \mathcal{H}} (\log(x + d_i))^{m_i - 1} (\log(x + d_i) - 1) dx + O\left(E_{N, \mathcal{K}} \log^K N \right. \\
&\quad \left. + \int_{1^-}^N E_{x, \mathcal{K}} \frac{\log^K(x + H)}{x} dx\right).
\end{aligned}$$

This is the analogue of equation (65) in [1]. Now, note that

$$\Lambda_m(n) = \Lambda(n) (\log n)^{m-1} (\log n - 1)$$

when n is prime. We have, by following the argument in [1],

$$\begin{aligned}
& \sum_{n=1}^N \prod_{i=1}^k \Lambda_{m_i}(n + d_i) \\
&= \left(\sum_{\mathcal{J} \subset \mathcal{J} \subset \mathcal{K}} \mathfrak{S}_0(\mathcal{D}_{\mathcal{J}}) \right) (I_{\mathbf{m}}(N) + O(H(\log N)^{K-k})) + O(N^{1/2+\epsilon}) \\
&+ O\left(E_{N, \mathcal{K}} \log^K N + \int_{1^-}^N E_{x, \mathcal{K}} \frac{\log^K(x+H)}{x} dx \right).
\end{aligned} \tag{5}$$

where

$$I_{\mathbf{m}}(N) := \int_1^N \prod_{i \in \mathcal{J}} ((\log x)^{m_i-1} (\log x - 1)) dx.$$

Putting (5) into (4),

$$\begin{aligned}
L_k(\mathbf{m}) &= I_{\mathbf{m}}(N) \sum_{\mathcal{J} \subset \mathcal{J} \subset \mathcal{K}} \sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} \mathfrak{S}_0(\mathcal{D}_{\mathcal{J}}) + O(H^k N^{1/2+\epsilon}) \\
&+ O\left(\sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} E_{N, \mathcal{K}} \log^K N + \int_{1^-}^N \sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} E_{x, \mathcal{K}} \frac{\log^K x}{x} dx \right).
\end{aligned} \tag{6}$$

Now, we use Conjecture 2. By Cauchy-Schwarz inequality,

$$\sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} |E_k(x, \mathcal{D})| \ll_k x^{1/2+\epsilon} H^k.$$

In particular,

$$\begin{aligned}
\sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} E_{x, \mathcal{K}} &= \sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} \sum_{\mathcal{K}' \subset \mathcal{K}} |E_{|\mathcal{K}'|}(x; \mathcal{D}_{\mathcal{K}'})| \\
&\ll_k \sum_{j=0}^k \sum_{\substack{d_1, \dots, d_j \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} |E_j(x; \mathcal{D})| \ll_k x^{1/2+\epsilon} H^k
\end{aligned}$$

Applying this to (6), the second error term is $\ll N^{1/2+\epsilon} H^k$ while the third error

term is

$$\begin{aligned}
&= \int_H^N \sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} E_{x, \mathcal{X}} \frac{\log^K(x+H)}{x} dx + \int_{1^-}^H \sum_{\substack{d_1, \dots, d_k \\ 1 \leq d_i \leq H \\ d_i \text{ distinct}}} E_{x, \mathcal{X}} \frac{\log^K(x+H)}{x} dx \\
&\ll_k N^{1/2+\epsilon} H^k + \int_1^H H^{k+1} \frac{\log^K(x+H)}{x} dx \ll_K N^{1/2+\epsilon} H^k
\end{aligned}$$

as $H \leq N^{1/2}$. Hence, (6) has an error $O(H^k N^{1/2+\epsilon})$ and the rest of the proof in [1] follows. Therefore, we have Theorem 1 under Conjecture 2.

References

- [1] H.L. Montgomery and K. Soundararajan, *Primes in Short Intervals*, Commun. Math. Phys. **252** (2004), 589-617.

Tsz Ho Chan
American Institute of Mathematics
360 Portage Avenue
Palo Alto, CA 94306
U.S.A.
thchan@aimath.org