ON THE UNIQUENESS OF THE FOLIATION OF SPHERES OF CONSTANT MEAN CURVATURE IN ASYMPTOTICALLY FLAT 3-MANIFOLDS

JIE QING AND GANG TIAN

ABSTRACT. In this note we study constant mean curvature surfaces in asymptotically flat 3-manifolds. We prove that, outside a given compact subset in an asymptotically flat 3-manifold with positive mass, stable spheres of given constant mean curvature are unique. Therefore we are able to conclude that there is a unique foliation of stable spheres of constant mean curvature in an asymptotically flat 3-manifold with positive mass.

1. Introduction

In the description of isolated gravitational system in General relativity a spacelike time-slice has the structure of a complete Riemannian 3-manifold with an asymptotically flat end. Such asymptotically flat end is diffeomorphic to $R^3 \setminus B_1(0)$ and the metric on it asymptotically approaches the Euclidean metric near the infinity:

$$g_{ij} = (1 + \frac{2m}{r})\delta_{ij} + O(r^{-2}),$$

where r is the Euclidean distance in R^3 . The constant m can be interpreted as the total mass of the isolated system and is referred to as ADM mass in literature [ADM]. It has also been established in [B] that with reasonable conditions ADM mass can be geometrically defined independent of the choices of coordinate system at infinity.

Often it is better to consider an asymptotically flat end as a perturbation of the static time-slice of the Schwarzchild space-time. Let us start with a precise definition of asymptotically flat 3-manifolds adopted from [HY] for our discussions in this note as follows:

¹After we posted our first version of this paper, we got some feedbacks indirectly about our main result compared to previous results on this topic. In this new version we stated our main result more precisely and gave more specific references for a clearer comparison of our main result to the previous ones.

Definition 1.1. A complete Riemannian 3-manifold (M, g) is said to be an asymptotically flat 3-manifold with mass m if there is a compact domain K of M such that $M \setminus K$ is diffeomorphic to $R^3 \setminus B_1(0)$ and the metric g in this coordinate system is given as

$$g_{ij}(x) = (1 + \frac{m}{2|x|})^4 \delta_{ij} + T_{ij}(x),$$

for all $x \in \mathbb{R}^3 \setminus B_1(0)$ with a constant C such that

(1.1)
$$|\partial^l T_{ij}|(x) \le C|x|^{-2-l}, \quad 1 \le l \le 4,$$

where ∂ denotes partial derivatives with respect to the Euclidean coordinates.

The existence of a unique foliation of spheres of constant mean curvature near the end in an asymptotically flat manifold is very important question. Among many applications, the unique foliation of spheres of constant mean curvature can be used to construct a geometrically canonical coordinate system at the infinity of asymptotically flat end. It can also be used to define a geometric center of mass for an isolated gravitational system (cf. [HY]). The existence of such unique foliation of spheres of constant mean curvature at the asymptotically flat end is also helpful to the study of Penrose inequality regarding the mass (cf. [Br]). In this note we show that indeed outside a given compact subset in an asymptotically flat 3-manifold with positive mass there is a unique foliation of stable spheres of constant mean curvature. Our main theorem² is

Theorem 1.1. Suppose (M,g) is an asymptotically flat 3-manifold with positive mass. Then there exists a compact domain K such that stable spheres of given constant mean curvature which separates the infinity from the compact domain K are unique. Hence there exists a unique foliation of stable spheres of constant mean curvature outside the compact domain K in M.

The existence of a foliation of stable spheres of constant mean curvature near asymptotically flat ends was established by Huisken and Yau in [HY] (also see [Ye]). Some uniqueness results with additional assumptions were also proven in [Br] [HY] [Ye]. The major difficulty of establishing the uniqueness of spheres of given constant

²The uniqueness problem addressed here was referred as the global uniqueness of stable CHC surfaces in [HY]. Their result on this global uniqueness was stated in Theorem 5.1 in [HY]. They proved that for $q > \frac{1}{2}$, if H sufficiently small, there is a unique stable constant mean curvature surface of mean curvature H outside $B_{H-q}(0)$. It has been a long-standing question whether stable constant mean curvature surfaces are unique outside a fixed compact subset. In the paragraph after Theorem 5.1 on page 301, Huisken and Yau stated: "it is an open question whether stable constant mean curvature surfaces are actually completely unique outside a fixed compact subset." Our main theorem gives an affirmative answer to this question.

mean curvature is that possible drifting of the spheres of constant mean curvature presents a hurdle to any useful global a priori estimates on the curvature. As a matter of fact, the uniqueness is known if one assumes no drifting (cf. [HY] [Ye]). Moreover, it was proven in [HY] that, if the drifting was somehow mild, then the uniqueness holds (cf. Theorem 5.1 on page 301 in [HY]).

Our main technical contributions can be summarized as follows: First, as a sharp contrast to the Euclidean space, similar to (5.13) in [HY], we find the following scale invariant integral which detects the nonzero mass. Suppose that N is a surface of constant mean curvature in an asymptotically flat end $(R^3 \setminus B_1(0), g)$ with positive mass m. Then

(1.2)
$$\frac{1}{8\pi} \int_{N} \frac{H}{|x|} \nu \cdot b d\sigma + \frac{1}{4\pi} \int_{N} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} d\sigma \le C m^{-1} r_{0}^{-1},$$

where $d\sigma$ is induced from the Euclidean metric, C>0 is some constant, b is any vector in R^3 , ν is the unit out-going normal vector of N in R^3 with respect to the Euclidean metric, and

$$(1.3) r_0 = \min\{|x| : x \in N \subset R^3 \setminus B_1(0)\},\$$

provided that

where $d\mu$ is induced from g. Secondly³, we are able to obtain estimates (cf. Corollary 4.4 and Corollary 4.5 in Section 4), which are beyond one individual scale in the blow-down analysis, via an asymptotic analysis used in an early work of us [QT]. The blow-down for a surface N of constant mean curvature H with the scale H is defined as,

(1.5)
$$\tilde{N} = \{\frac{1}{2}Hx : x \in N \subset \mathbb{R}^3 \setminus B_1(0)\} \subset \mathbb{R}^3.$$

The use of the asymptotic analysis introduced in Section 4 is the key which allows us to obtain some finer estimates and untangle the problem that uniform roundness

³In [HY], a global estimate was sought after (cf. Lemma 5.6 in [HY]), with a compromise to assume that the inner radius is not smaller than H^{-q} for $q > \frac{1}{2}$. They stated in the paragraph after Theorem 5.1 (page 21, [HY]) that their assumption on inner radius "seems to be optimal from a technical point of view". While in this paper we do different estimates in three different scales. Particularly we establish some decay estimate for the intermediate scales by using an asymptotic analysis developed in [QT].

and non-drifting of spheres of constant mean curvature hinge on each other. More precisely, to eliminate the possible drifting, one carefully calculates the two integrals in left-hand side of (1.3) for \tilde{N} ,

(1.6)
$$\frac{1}{4\pi} \int_{\tilde{N}} \frac{1}{|x|} \nu \cdot b d\sigma + \frac{1}{4\pi} \int_{\tilde{N}} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma$$

with some particular choice of b. If drifting happened, then the rescaled surface \tilde{N} would approach the origin. Then one evaluates the integrals over three different regions: 1) the part of \tilde{N} that is any fixed distance away from the origin; 2) the part of \tilde{N} that is near the origin in the scale of Hr_0 ; 3) the transition between the above two. We will employ Corollary 4.5 in Section 4 to show the integrals on third region contribute something negligible. Consequently we are able to prove that the drifting of stable spheres of constant mean curvature does not happen at all in an asymptotically flat 3-manifold with positive mass. Then using the early existence and uniqueness results in [HY] and [Ye], for instance, Theorem 5.1 in [HY], we may conclude our main theorem.

It is worthwhile to note that the uniqueness of spheres of given constant mean curvature outside the horizon in the Schwarzchild space is an interesting open problem. In his thesis [Br], Bray proved the coordinate spheres are the unique minimizing surfaces of given constant mean curvature outside the horizon in Scwarzchild space, in an attempt to prove the Penrose inequality regarding the mass by the foliation of constant mean curvature surfaces. Theorem 1.1 in the above particularly implies that the coordinate spheres are the only stable sphere of constant mean curvature near the infinity of the Schwarzchild space which separates the infinity from the horizon.

The paper is organized as follows: In Section 2 we will obtain the curvature estimates based on the Simons' identity and the smallness of the integral of the traceless part of the second fundamental form. In Section 3 we introduce the blowdown analysis in all scales. In Section 4 we recall the asymptotic analysis from [QT] and prove a technical proposition. Finally in Section 5 we introduce a sense of the center of mass and prove our main theorem.

2. Curvature estimates

First let us recall the Simons' identity [SSY] [Sj] for a hypersurface N in a Riemannian manifold (M, g) (cf. Lemma 1.3 in [HY]):

(2.1)
$$\Delta h_{ij} = \nabla_i \nabla_j H + H h_{ik} h_{jk} - |A|^2 h_{ij} + H R_{3i3j} - h_{ij} R_{3k3k} + h_{jk} R_{klil} + h_{ik} R_{klil} - 2h_{lk} R_{iljk} + \nabla_i R_{3kik} + \nabla_k R_{3ijk}$$

where $A = (h_{ij})$ is the second fundamental form for N in M, H = TrA is mean curvature, and R_{ijkl} and ∇R_{ijkl} are curvature and covariant derivatives of curvature for (M, g). When N is a constant mean curvature hypersurface, we rather like to rewrite it as an equation for the traceless part \mathring{A} of A, i.e. $\mathring{A} = A - \frac{1}{2}H$.

$$\Delta \mathring{A}_{ij} = H \mathring{A}_{ik} \mathring{A}_{jk} - \frac{1}{2} H |\mathring{A}|^2 \delta_{ij} - (|\mathring{A}|^2 + \frac{1}{2} H^2) \mathring{A}_{ij}$$

$$+ H R_{3i3j} - \frac{1}{2} H R_{3k3k} \delta_{ij} - \mathring{A}_{ij} R_{3k3k}$$

$$+ \mathring{A}_{jk} R_{klil} + \mathring{A}_{ik} R_{kljl} - 2 \mathring{A}_{lk} R_{likj}$$

$$+ \nabla_j R_{3kik} - \nabla_k R_{3ikj}.$$

Lemma 2.1. Suppose that N is a constant mean curvature surface in an asymptotically flat end $(R^3 \setminus B_1(0), g)$. Then

(2.3)
$$-|\mathring{A}|\Delta|\mathring{A}| \leq |\mathring{A}|^4 + CH|\mathring{A}|^3 + CH^2|\mathring{A}|^2 C|\mathring{A}|^2|x|^{-3} + CH|\mathring{A}||x|^{-3} + C|\mathring{A}||x|^{-4}.$$

Note that, in an asymptotically flat end (cf. Definition 1.1 in Section 1),

$$(2.4) |R_{ijkl}| \le C|x|^{-3}, |\nabla R_{ijkl}| \le C|x|^{-4}.$$

We refer readers to [HY] for the calculations of curvature of the Schwarzschild space and asymptotically flat ends.

Lemma 2.2. Suppose that N is a constant mean curvature surface in an asymptotically flat end $(R^3 \setminus B_1(0), g)$. Then $\int_N H_e^2 d\sigma$ is bounded if and only if $\int_N H^2 d\mu$ is bounded, provided that r_0 is sufficiently large.

Proof. First one may calculate

(2.5)
$$H_e = \left(1 + \frac{m}{2r}\right)^2 H + 2\left(1 + \frac{m}{2r}\right)^{-1} \frac{m}{r^3} x \cdot \nu + O(r^{-3}),$$

where H_e is the mean curvature of $N \subset \mathbb{R}^3$ with respect to the Euclidean metric (cf. Lemma 1.4 in [HY]). Hence

$$H_e^2 = H^2 + O(r^{-1})H^2 + O(r^{-2})H + O(r^{-3}).$$

Following Lemma 5.2 in [HY] and the fact that g is quasi-isometric to the Euclidean metric $|dx|^2$, we have:

$$\begin{split} \int_N H_e^2 d\sigma & \leq C \int_N H_e^2 d\mu \leq C \int_N H^2 d\mu + C (\int_N H^2 d\mu)^{\frac{1}{2}} (\int_N r^{-4} d\mu)^{\frac{1}{2}} + C \int_N r^{-3} d\mu \\ & \leq C \int_N H^2 d\mu \end{split}$$

and

$$(1-Cr_0^{-1})\int_N H^2 d\mu \leq C\int_N H_e^2 d\sigma.$$

Thus the lemma is proved.

Therefore, following Lemma 1 in [Si], we have

Lemma 2.3. Suppose that N is a constant mean curvature surface in an asymptotically flat end $(R^3 \setminus B_1(0), g)$ with $r_0(N)$ sufficiently large, and that

$$\int_{N} H^{2} d\mu \le C.$$

Then

$$C_1 H^{-1} \le diam(N) \le C_2 H^{-1}$$
.

We would like to point out that, if the surface N separates the infinity from the compact part, i.e. the origin is inside $N \subset \mathbb{R}^3$, then the above lemma implies

$$(2.6) C_1 H^{-1} \le r_1(N) \le C_2 H^{-1},$$

where the outer radius $r_1(N)$ is defined as

$$r_1(N) = \max\{|x| : x \in N \subset R^3 \setminus B_1(0)\}.$$

Based on Michael and Simon [MS], one has the following Sobolev inequality (cf. Lemma 5.6 in [HY]).

Lemma 2.4. Suppose that N is a constant mean curvature surface in an asymptotically flat end $(R^3 \setminus B_1(0), g)$ with $r_0(N)$ sufficiently large, and that

$$\int_{N} H^{2} d\mu \le C.$$

Then

(2.7)
$$(\int_{N} f^{2} d\mu)^{\frac{1}{2}} \leq C(\int_{N} |\nabla f| d\mu + \int_{N} H|f| d\mu).$$

Now we are ready to state and prove the main curvature estimates:

Theorem 2.5. Suppose that $(R^3 \setminus B_1(0), g)$ is an asymptotically flat end. Then there exist positive numbers σ_0 , ϵ_0 and δ_0 such that for any constant mean curvature surface in the end, which separates the infinity from the compact part, we have

(2.8)
$$|\mathring{A}|^{2}(x) \leq C|x|^{-2} \int_{B_{\delta_{0}|x|}(x)} |\mathring{A}|^{2} d\mu + C|x|^{-4},$$

provided that

$$\int_{N} |\mathring{A}|^2 d\mu \le \epsilon_0$$

and $r_0(N) \ge \sigma_0$. And the corresponding a priori estimates for all covariant derivatives of curvature also hold consequently.

Proof. Recall that

$$-|\mathring{A}|\Delta|\mathring{A}| \le |\mathring{A}|^4 + CH|\mathring{A}|^3 + CH^2|\mathring{A}|^2$$
$$C(|\mathring{A}|^2|x|^{-3} + CH|\mathring{A}||x|^{-3} + C|\mathring{A}||x|^{-4}).$$

Multiply the two sides with ϕ^3 , where ϕ is an appropriate cutoff function of small support, and integrate,

$$\int_{N} -\phi^{3} |\mathring{A}| \Delta |\mathring{A}| d\mu \leq \int_{N} \phi^{3} |\mathring{A}|^{4} d\mu + C \int_{N} H \phi^{3} |\mathring{A}|^{3} d\mu + C \int_{N} H^{2} \phi^{3} |\mathring{A}|^{2} d\mu + C r_{0}^{-1} \int_{N} \phi^{3} (|\mathring{A}|^{2} |x|^{-2} + CH |\mathring{A}| |x|^{-2} + C |\mathring{A}| |x|^{-3}) d\mu$$

where

$$\begin{split} \int_N -\phi^3 |\mathring{A}| \Delta |\mathring{A}| d\mu &= \int_N \nabla (\phi^3 |\mathring{A}|) \nabla |\mathring{A}| d\mu \\ &= \int_N \phi^2 \nabla (\phi |\mathring{A}|) \nabla |\mathring{A}| + \int_N 2\phi^2 |\mathring{A}| \nabla \phi \nabla |\mathring{A}| d\mu \\ &= \int_N \phi |\nabla (\phi |\mathring{A}|)|^2 d\mu + \int_N \phi |\mathring{A}| \nabla (\phi |\mathring{A}|) \nabla \phi d\mu + \int_N 2\phi^2 |\mathring{A}| \nabla \phi \nabla |\mathring{A}| d\mu \\ &\geq \frac{3}{4} \int_N \phi |\nabla (\phi |\mathring{A}|)|^2 d\mu - C \int_N \phi |\mathring{A}|^2 |\nabla \phi|^2 d\mu + \int_N 2\phi^2 |\mathring{A}| \nabla \phi \nabla |\mathring{A}| d\mu \\ &\geq \frac{1}{2} \int_N \phi |\nabla (\phi |\mathring{A}|)|^2 d\mu - C \int_N \phi |\mathring{A}|^2 |\nabla \phi|^2 d\mu \\ &\int_N \phi^3 |\mathring{A}|^4 d\mu \leq (\int_{\operatorname{supp}(\phi)} |\mathring{A}|^2 d\mu)^{\frac{1}{2}} (\int_N (\phi |\mathring{A}|)^6 d\mu)^{\frac{1}{2}} \end{split}$$

and

$$\int_N H\phi^3 |\mathring{A}|^3 d\mu \leq (\int_{\mathrm{supp}(\phi)} H^2 d\mu)^{\frac{1}{2}} (\int_N (\phi |\mathring{A}|)^6 d\mu)^{\frac{1}{2}}.$$

For other terms

$$\int_{N} H^{2} \phi^{3} |\mathring{A}|^{2} d\mu \leq C r_{M}^{-2} \int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu \leq C |x_{0}|^{-2} \int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu,$$

$$\int_{N} \phi^{3} |x|^{-2} |\mathring{A}|^{2} d\mu \leq C |x_{0}|^{-2} \int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu,$$

$$\int_{N} \phi^{3} H |x|^{-2} |\mathring{A}| d\mu \leq C |x_{0}|^{-2} (\int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu)^{\frac{1}{2}},$$

and

$$\int_{N} \phi^{2} |x|^{-3} |\mathring{A}| d\mu \le C|x_{0}|^{-2} \left(\int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu \right)^{\frac{1}{2}}.$$

Note that, for a given point x_0 , we may choose the cutoff function ϕ so that it has the suppose of a disk of radius, say, $\delta_0|x_0|$ (δ_0 to be determined). Now, combining all terms, we have

$$\int_{N} \phi |\nabla(\phi|\mathring{A}|)|^{2} d\mu \leq 2 \left(\int_{N} |\mathring{A}|^{2} d\mu\right)^{\frac{1}{2}} \left(\int_{N} (\phi|\mathring{A}|)^{6} d\mu\right)^{\frac{1}{2}} + C \left(\int_{\text{supp}(\phi)} H^{2} d\mu\right)^{\frac{1}{2}} \left(\int_{N} (\phi|\mathring{A}|)^{6} d\mu\right)^{\frac{1}{2}} + C |x_{0}|^{-2} \left(\int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu\right)^{\frac{1}{2}}.$$

Applying the Sobolev inequality with $f = \phi^3 g^3$ where $g = |\mathring{A}|$, we have

$$\begin{split} &(\int_{N}(\phi g)^{6}d\mu)^{\frac{1}{2}} \leq C(3\int_{N}(\phi g)^{2}|\nabla(\phi g)|d\mu + \int_{N}H(\phi g)^{3}) \\ &\leq C(\int_{N}\phi^{3}g^{4}d\mu)^{\frac{1}{2}}(\int_{N}|\nabla(\phi g)|^{2}\phi d\mu)^{\frac{1}{2}} + (\int_{\mathrm{supp}(\phi)}H^{2}d\mu)^{\frac{1}{2}}(\int_{N}(\phi g)^{6}d\mu)^{\frac{1}{2}} \\ &\leq C(\int_{N}g^{2}d\mu)^{\frac{1}{2}}(\int_{N}(\phi g)^{6}d\mu)^{\frac{1}{2}} + C(\int_{\mathrm{supp}(\phi)}H^{2}d\mu)^{\frac{1}{2}}(\int_{N}(\phi g)^{6}d\mu)^{\frac{1}{2}} \\ &\quad + C\int_{N}|\nabla(\phi g)|^{2}\phi d\mu. \end{split}$$

Thus

$$(2.9) \qquad (\int_{N} (\phi |\mathring{A}|)^{6} d\mu)^{\frac{1}{2}} \leq C|x_{0}|^{-2} (\int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu)^{\frac{1}{2}},$$

which implies

(2.10)
$$\int_{N} (\phi |\mathring{A}|)^{4} d\mu \leq C |x_{0}|^{-2} \int_{\text{supp}(\phi)} |\mathring{A}|^{2} d\mu.$$

Note that we have chosen δ_0 small enough so that

$$C\int_{\operatorname{supp}(\phi)} H^2 d\mu \le \frac{1}{8}$$

and $\int_N |\mathring{A}|^2 d\mu \leq \epsilon_0$, where ϵ_0 is small enough so that

$$C\int_{N} |\mathring{A}|^2 d\mu \le \frac{1}{8}.$$

Now we proceed to get the point-wise estimates. First, if we take $f=u^2$ in the Sobolev inequality, then

$$\begin{split} (\int_N u^4 d\mu)^{\frac{1}{2}} & \leq C(2 \int_N |u| |\nabla u| d\mu + \int_N H u^2 d\mu) \\ & \leq C(\int_N u^2 d\mu)^{\frac{1}{2}} (\int_N |\nabla u|^2 d\mu)^{\frac{1}{2}} + C(\int_{\mathrm{supp}(u)} H^2 d\mu)^{\frac{1}{2}} (\int_N u^4 d\mu)^{\frac{1}{2}}. \end{split}$$

When u has the support as the cutoff function ϕ , we have

$$(2.11) \qquad (\int_N u^4 d\mu)^{\frac{1}{2}} \le C(\int_N u^2 d\mu)^{\frac{1}{2}} (\int_N |\nabla u|^2 d\mu)^{\frac{1}{2}}.$$

To finish the point-wise estimates we use the following rather standard estimate:

Lemma 2.6. Suppose that a nonnegative function v in L^2 solves

$$(2.12) -\Delta v \le fv + h$$

on $B_{2R}(x_0)$, where

$$\int_{B_{2R}(x_0)} f^2 d\mu \le CR^{-2}$$

and $h \in L^2(B_{2R}(x_0))$. And Suppose that

$$\left(\int_{N} u^{4} d\mu\right)^{\frac{1}{2}} \leq C\left(\int_{N} u^{2} d\mu\right)^{\frac{1}{2}} \left(\int_{N} |\nabla u|^{2} d\mu\right)^{\frac{1}{2}}$$

holds for all u with support inside $B_{2R}(x_0)$. Then

$$\sup_{B_R(x_0)} v \le CR^{-1} \|v\|_{L^2(B_{2R}(x_0))} + CR \|h\|_{L^2(B_{2R}(x_0))}.$$

Proof. We will simply use the Moser iteration method. For convenience, we may rescale so that we are working on B_2 . The correct scales would be

$$v_R(x) = v(Rx), f_R(x) = R^2 f(Rx), \text{ and } h_R = R^2 h(Rx).$$

Let $k = ||h||_{L^2(B_2)}$ and $\bar{v} = v + k$. Multiply the equation with $\phi^2 \bar{v}^{p-1}$ on the both sides

$$\int |\nabla (\phi \bar{v}^{\frac{p}{2}})|^2 \le \frac{2}{p} \int f \phi^2 \bar{v}^p + \int h \phi^2 \bar{v}^{p-1} + C \int |\nabla \phi|^2 \bar{v}^p.$$

Set $\bar{f} = \frac{h}{k} + f$ we have

$$\int |\nabla (\phi \bar{v}^{\frac{p}{2}})|^2 \le \frac{2}{p} \int \bar{f} \phi^2 \bar{v}^p + C \int |\nabla \phi|^2 \bar{v}^p.$$

Note that $\|\bar{f}\|_{L^2(B_2)} \leq 1 + \|f\|_{L^2(B_2)}$. By the assumed Sobolev inequality, we

$$\left(\int (\phi \bar{v}^{\frac{p}{2}})^4\right)^{\frac{1}{2}} \le C(p \int |\bar{f}|\phi^2 \bar{v}^p)^{\frac{1}{2}} \left(\int \phi^2 \bar{v}^p\right)^{\frac{1}{2}} + C(\int |\nabla \phi|^2 v^p)^{\frac{1}{2}} \left(\int \phi^2 v^p\right)^{\frac{1}{2}}.$$

To handle the first term, we apply Hólder inequality

$$\begin{split} (p\int |\bar{f}|\phi^2\bar{v}^p)^{\frac{1}{2}} (\int \phi^2\bar{v}^p)^{\frac{1}{2}} &\leq p^{\frac{1}{2}} (\int |\bar{f}|^2)^{\frac{1}{4}} (\int \phi^2\bar{v}^p)^{\frac{1}{2}} (\int (\phi\bar{v}^{\frac{p}{2}})^4)^{\frac{1}{4}} \\ &\leq \frac{1}{2C} (\int (\phi\bar{v}^{\frac{p}{2}})^4)^{\frac{1}{2}} + Cp (\int |\bar{f}|^2)^{\frac{1}{2}} \int \phi^2\bar{v}^p. \end{split}$$

Hence

$$\left(\int (\phi \bar{v}^{\frac{p}{2}})^4\right)^{\frac{1}{2}} \le C(p\|\bar{f}\|_{L^2} \int \phi^2 \bar{v}^p + \int |\nabla \phi|^2 \bar{v}^p).$$

Now, for $i = 1, 2, \ldots$, let $p = 2^i$ and

$$\phi = \begin{cases} 1 & \forall x \in B_{1+2^{-i}} \\ 0 & \forall x \notin B_{1+2^{-i+1}} \end{cases}$$

Then

$$(\int_{B_{1+2^{-i}}} \bar{v}^{2^{i+1}})^{2^{-i-1}} \leq C^{2^{-i}} 2^{i2^{-i}} (\int_{B_{1+2^{-i+1}}} \bar{v}^{2^i})^{2^{-i}}.$$

Thus

$$\sup_{B_1} v \le \sup_{B_1} \bar{v} \le C^{\sum_{i=1} 2^{-i}} 2^{\sum_{i=1} i 2^{-i}} (\int_{B_2} \bar{v}^2)^{\frac{1}{2}} \le C(\|v\|_{L^2(B_2)} + \|h\|_{L^2(B_2)}),$$

whose scaled version gives the lemma.

To get curvature estimates, we write the equation in such way as (2.12) that we may apply the above lemma for

$$f = C(|\mathring{A}|^2 + H|\mathring{A}| + H^2 + r^{-3})$$
 and $h = C(Hr^{-3} + r^{-4})$,

in the light of (2.9) and (2.10).

3. Blow-down analysis

In order to understand a surface of constant mean curvature N in an asymptotically flat end $(R^3 \setminus B_1(0), g)$, we will need to blow down the surface in different scales. We first consider, the blow-down by the scale H,

(3.1)
$$\tilde{N} = \frac{1}{2}HN = \{\frac{1}{2}Hx : x \in N\}.$$

Suppose that there is a sequence of constant mean curvature surfaces $\{N_i\}$ such that

(3.2)
$$\lim_{i \to \infty} r_0(N_i) = \infty \text{ and } \lim_{i \to \infty} \int_{N_i} H^2 d\mu = 16\pi.$$

Then, by a similar argument to the proof of Lemma 2.2 in the previous section, we have

(3.3)
$$\lim_{i \to \infty} \int_{N_i} H_e^2 d\sigma = 16\pi.$$

Hence, by the curvature estimates established in the previous section combining the proof of Theorem 1 in [Si], we have

Lemma 3.1. Suppose that $\{N_i\}$ is a sequence of constant mean curvature surfaces in a given asymptotically flat end $(R^3 \setminus B_1(0), g)$ and that

$$\lim_{i \to \infty} r_0(N_i) = \infty \text{ and } \lim_{i \to \infty} \int_{N_i} H^2 d\mu = 16\pi.$$

And suppose that N_i separates the infinity from the compact part. Then, there is a subsequence of $\{\tilde{N}_i\}$ which converges in Gromov-Hausdorff distance to a round sphere $S_1^2(a)$ of radius 1 and centered at $a \in R^3$. Moreover, the convergence is in C^{∞} sense away from the origin.

From the above lemma, the difficulty will be to study the possibility of having the origin lying on the sphere $S^2(a)$, that is,

(3.5)
$$\lim_{i \to \infty} r_0(N_i) = \infty, \text{ and } \lim_{i \to \infty} r_0(N_i)H(N_i) = 0.$$

Then, in the light of the curvature estimates we obtained in the previous section, we may use the smaller scale $r_0(N_i)$ to blow down the surface

(3.6)
$$\hat{N} = r_0(N)^{-1}N = \{r_0^{-1}x : x \in N\}.$$

Lemma 3.2. Suppose that $\{N_i\}$ is a sequence of constant mean curvature surfaces in a given asymptotically flat end $(R^3 \setminus B_1(0), g)$ and that

$$\lim_{i \to \infty} r_0(N_i) = \infty \ and \ \lim_{i \to \infty} \int_{N_i} H^2 d\mu = 16\pi.$$

And suppose that

$$\lim_{i \to \infty} r_0(N_i)H(N_i) = 0.$$

Then there is a subsequence of $\{\hat{N}_i\}$ converges to a 2-plane at distance 1 from the origin. Moreover the convergence is in C^{∞} in any compact set of R^3 .

As one would expect, the real difficulty is to understand the behavior of the surfaces N_i in the scales between $r_0(N_i)$ and $H^{-1}(N_i)$. To start we consider the intermediate scales r_i such that

(3.7)
$$\lim_{i \to \infty} \frac{r_0(N_i)}{r_i} = 0 \text{ and } \lim_{i \to \infty} r_i H(N_i) = 0$$

and blow down the surfaces

(3.8)
$$\bar{N}_i = r_i^{-1} N = \{ r_i^{-1} x : x \in N \}.$$

Lemma 3.3. Suppose that $\{N_i\}$ is a sequence of constant mean curvature surfaces in a given asymptotically flat end $(R^3 \setminus B_1(0), g)$ and that

$$\lim_{i \to \infty} r_0(N_i) = \infty \ and \ \lim_{i \to \infty} \int_{N_i} H^2 d\mu = 16\pi.$$

And suppose that $\{r_i\}$ are such that

$$\lim_{i \to \infty} \frac{r_0(N_i)}{r_i} = 0 \text{ and } \lim_{i \to \infty} r_i H(N_i) = 0.$$

Then there is a subsequence of $\{\bar{N}_i\}$ converges to a 2-plane at the origin in Gromov-Hausdorff distance. Moreover the convergence is C^{∞} in any compact subset away from the origin.

4. Asymptotic analysis

In this section we would like to apply the asymptotic analysis used in [QT] to obtain some estimate that holds over the whole transition region between the scales $r_0(N_i)$ and $r_1(N_i)$. But first, let us revise Proposition 2.1 in [QT] as follows. Let us denote

$$||u||_i^2 = \int_{[(i-1)L,iL]\times S^1} |u|^2 dt d\theta.$$

Lemma 4.1. Suppose $u \in W^{1,2}(\Sigma, \mathbb{R}^k)$ satisfies

$$\Delta u + A \cdot \nabla u + B \cdot u = h$$
 in Σ .

where $\Sigma = [0, 3L] \times S^1$. And suppose that L is given and large. Then there exists a positive number δ_0 such that, if

$$||h||_{L^2(\Sigma)} \le \delta_0 \max_{1 \le i \le 3} \{||u||_i\}$$

and

$$||A||_{L^{\infty}(\Sigma)} \le \delta_0, \quad ||B||_{L^{\infty}(\Sigma)} \le \delta_0,$$

then.

- (a). $||u||_3 \le e^{-\frac{1}{2}L} ||u||_2$ implies $||u||_2 < e^{-\frac{1}{2}L} ||u||_1$,
- (b). $||u||_1 \le e^{-\frac{1}{2}L}||u||_2$ implies $||u||_2 < e^{-\frac{1}{2}L}||u||_3$, and
- (c). If both $\int_{L\times S^1} u d\theta$ and $\int_{2L\times S^1} u d\theta \le \delta_0 \max_{1\le i\le 3} \{\|u\|_i\}$, then either $\|u\|_2 < e^{-\frac{1}{2}L} \|u\|_1$ or $\|u\|_2 < e^{-\frac{1}{2}L} \|u\|_3$.

Proof. We refer to the proof of Proposition 2.1 in [QT] for more details. In the proof by contradiction argument one needs to make sure that the sequence of normalized u_k converges to a non-zero harmonic function u and the non-zero harmonic function u violates one of (a)-(c). Interior elliptic estimates give the strong convergence in the middle section $I_2 = [L, 2L] \times S^1$, which implies that u is not trivially zero. Because, with the assumption of the proof by contradiction, the middle one is the largest. Finally u indeed induces a contradiction due to the Fatou lemma.

We would like to point out that Proposition 2.1 in [QT] is overstated since it is not correct for l > 3. But, in the proof of Proposition 3.1 in [QT], where Corollary 2.2 is used, one may replace the shifting cylinder with length 3L instead of 5L. The proof still works the same, which is, one push to the direction of growth the cylinder of length 3L when Corollary 2.2 in [QT] applies and it gives the estimates regardless of where one is stopped applying Corollary 2.2.

Given a surface N in \mathbb{R}^3 , recall from, for example, (8.5) in [Ka], that

$$(4.2) \Delta \nu + |\nabla \nu|^2 \nu = \nabla H_e$$

where ν is the Gauss map from $N \longrightarrow S^2$. For the constant mean curvature surfaces in the asymptotically flat end $(R^3 \setminus B_1(0), g)$,

$$(4.3) |\nabla H_e|(x) \le C|x|^{-3}.$$

Therefore we consider that the Gauss map of the constant mean curvature surfaces in the asymptotically flat end $(R^3 \setminus B_1(0), g)$ is an almost harmonic map. Hence we are in a situation which is very similar to that in [QT]. We will refer readers to [QT] for rather elementary yet involved analysis since the proof we present here is some modifications from the proof in [QT]. We will not carry the indices for the surfaces N_i if it does not cause any confusion. Set

$$(4.4) A_{r_1,r_2} = \{x \in N : r_1 \le |x| \le r_2\}.$$

 A_{r_1,r_2}^0 stands for the standard annulus in R^2 . We are concerned with the behavior of ν on the part $A_{Kr_0(N),sH^{-1}(N)}$ of N where K will be fixed large and s will be fixed small. The first difference from [QT] is that, while we had a fixed domain in [QT], we need the following lemma in order to be in the position to use Lemma 4.1 in the above.

Lemma 4.2. Suppose that N is a constant mean curvature surfaces in a given asymptotically flat end $(R^3 \setminus B_1(0), g)$. Then, for any $\epsilon > 0$ and L fixed, there are ϵ_0 , s and K such that, if

$$\int_{N} |\mathring{A}|^2 d\sigma \le \epsilon_0$$

and $Kr_0(N) < r < sH^{-1}(N)$, then $(r^{-1}A_{r,e^{4L}r}, r^{-2}g_e)$ may be represented as $(A_{1,e^{4L}}^0, \bar{g})$ where

(4.5)
$$\|\bar{g} - |dx|^2 \|_{C^1(A^0_{1,e^{4L}})} \le \epsilon.$$

In other words, in the cylindrical coordinates $(S^1 \times [\log r, 4L + \log r], \bar{g}_c)$,

This is a consequence of Lemma 3.3 in the previous section. Another difference from [QT] is that, we are considering maps with tension fields possibly blowing up at point but no energy concentration, while in [QT] we were considering almost harmonic maps with concentration of energy but tension fields uniformly bounded in L^2 . In cylindrical coordinates, the tension fields

(4.7)
$$|\tau(\nu)| = r^2 |\nabla H_e| \le Cr^{-1} = Ce^{-t}$$

for $t \in [\log(Kr_0), \log(sH^{-1})]$. Thus,

(4.8)
$$\int_{S^1 \times [t,t+L]} |\tau(\nu)|^2 dt d\theta \le Ce^{-2t}$$

which decays as needed in the arguments in [QT]. But to get the growth (or decay) of the energy along the cylinder we first can only have the estimate (3.8) in [QT]. Then we need to use the Hopf differential

$$\Phi = |\partial_t \nu|^2 - |\partial_\theta \nu|^2 - 2\sqrt{-1}\partial_t \nu \cdot \partial_\theta \nu$$

and the stationary property, in complex variable $z = t + \sqrt{-1}\theta$,

$$\bar{\partial}\Phi = \partial\nu \cdot \tau(\nu)$$

to bound $\int |\partial_t \nu|^2$ by $\int |\partial_\theta \nu|^2$ (cf. [QT] [DT]) as follows:

$$\int_{S^1 \times [t,t+L]} |\partial_t \nu|^2 dt d\theta \le \int_{S^1 \times [t,t+L]} |\Phi| dt d\theta + \int_{S^1 \times [t,t+L]} |\partial_\theta \nu|^2 dt d\theta.$$

By the elliptic estimates (cf. [DT]), we have

$$\int_{S^1\times[t,t+L]}|\Phi|dtd\theta\leq \int_{N\cap B^c_r}|\Phi|dtd\theta\leq C(\int_{N\cap B^c_r}|\nabla\nu|^2d\sigma)^{\frac{1}{2}}(\int_{N\cap B^c_r}|\tau(\nu)|^2d\sigma)^{\frac{1}{2}},$$

where $N \cap B_r^c$ is the part of N which is outside of B_r and is a disk since N is a sphere topologically. Hence, we have

(4.10)
$$\int_{S^1 \times [t, t+L]} |\Phi| dt d\theta \le C \left(\int_{N \cap B_r^c} |\tau(\nu)|^2 d\sigma \right)^{\frac{1}{2}} \le C e^{-t}.$$

Notice that in [QT] we instead used the fact that the tension fields is uniformly bounded in L^2 inside B_{δ} (cf. lines between (3.8) and (3.9) in [QT]). The rest of the proof of Proposition 3.1 in [QT] works with little modifications. Thus we have

Proposition 4.3. Suppose that $\{N_i\}$ is a sequence of constant mean curvature surfaces in a given asymptotically flat end $(R^3 \setminus B_1(0), g)$ and that

$$\lim_{i\to\infty} r_0(N_i) = \infty \ \ and \ \lim_{i\to\infty} \int_{N_i} H^2 d\mu = 16\pi.$$

And suppose that

$$\lim_{i \to \infty} r_0(N_i)H(N_i) = 0.$$

Then there exist a large number K, a small number s and i_0 such that, when $i \geq i_0$,

(4.11)
$$\max_{I_j} |\nabla \nu| \le C \left(\int_{B_{sH^{-1}(N_i)} \cap N_i} |\nabla \nu|^2 d\sigma + r_0^{-1} \right) \left(e^{-\frac{1}{4}jL} + e^{-\frac{1}{4}(n_i - j)L} \right),$$

where

$$I_j = S^1 \times [\log(Kr_0(N_i)) + jL, \log(Kr_0(N_i)) + (j+1)L]$$

and

$$j \in [0, n_i]$$
 and $\log(Kr_0(N_i)) + (n_i + 1)L = \log(sH(N_i)^{-1}).$

This finer analysis improves our understanding of the blow-downs that we discussed in the previous section. Namely,

Corollary 4.4. Assume the same conditions as Proposition 4.3. Then the limit plane in Lemma 3.2 and the limit plane in Lemma 3.3 are all orthogonal to the vector a. In fact, we may choose s small and i large enough so that,

$$|\nu(x) + a| \le \epsilon$$

for all $x \in N_i$ and $|x| \le sH^{-1}(N_i)$.

And we have

Corollary 4.5. Asume the same condition as Proposition 4.3. Let $\nu_i = \nu(p_i)$ for some $p_i \in I_{\frac{n_i}{2}}$. Then

(4.12)
$$\leq C \frac{1}{1 - e^{-\frac{1}{4}L}} \left(\int_{B_{sH^{-1}(N_i)} \cap N_i} |\nabla \nu|^2 d\sigma + r_0^{-1} \right) \left(e^{-\frac{1}{4}jL} + e^{-\frac{1}{8}n_i L} \right)$$

for $j \in [0, \frac{1}{2}n_i]$ and

$$(4.13) \qquad \max_{I_j} |\nu - \nu_i|$$

$$\leq C \frac{1}{1 - e^{-\frac{1}{4}L}} \left(\int_{B_{sH^{-1}(N_i)} \cap N_i} |\nabla \nu|^2 d\sigma + r_0^{-1} \right) \left(e^{-\frac{1}{8}n_i L} + e^{-\frac{1}{4}(n_i - j)L} \right)$$

for $j \in [\frac{1}{2}n_i, n_i]$.

The two corollary above will be the key for us to calculate the integrals in next section to prove our main theorem.

5. Center of mass

First let us recall that, for any embedded surface N in \mathbb{R}^3 and any given vector $b \in \mathbb{R}^3$,

$$\int_{N} H_{e}\nu \cdot bd\sigma = 0.$$

One may consider this as the first variation of the area of surface $N_t = N + tb \subset R^3$. On the other hand, if N is a constant mean curvature surface in the asymptotically flat end $(R^3 \setminus B_1(0), g)$, then

(5.2)
$$\int_{N} H\nu \cdot bd\sigma = H \int_{N} \nu \cdot bd\sigma = 0.$$

Since the flux is zero across any surface for a given constant velocity b. Thus, for any constant mean curvature surface in the asymptotically flat end,

(5.3)
$$\int_{N} (H_e - H)\nu \cdot bd\sigma = 0.$$

One may calculate and find

(5.4)
$$H_e - H = m\left(\frac{H}{|x|} + 2\frac{\nu \cdot x}{|x|^3}\right) + O(|x|^{-2})H + O(|x|^{-3}).$$

Lemma 5.1. Suppose N is a surface of constant mean curvature in the asymptotically flat end with positive mass $m \neq 0$. And suppose that

$$\int_N H^2 d\mu < \infty$$

and $r_0(N)$ is sufficiently large. Then for any given b and for some C>0,

(5.5)
$$\frac{1}{8\pi} \int_{N} \frac{H}{|x|} \nu \cdot b d\sigma + \frac{1}{4\pi} \int_{N} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} d\sigma \le C m^{-1} r_{0}^{-1}.$$

Proof. Simply multiply $\nu \cdot b$ to the both sides of (5.4) and integrate over the surface N, we have,

$$\frac{1}{8\pi} \int_N (H_e - H)\nu \cdot bd\sigma = \frac{m}{8\pi} \int_N \frac{H}{|x|} \nu \cdot bd\sigma + \frac{m}{4\pi} \int_N \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma + O(r_0^{-1}).$$

Here we used the Lemma 5.2 in [HY]. Then the lemma is proved due to (5.3).

Now, we are ready to state and prove our main theorem in this note as follows:

Theorem 5.2. Suppose that $\{N_i\}$ is a sequence of spheres of constant mean curvature in a given asymptotically flat end with positive mass $m \neq 0$ and that

$$\lim_{i \to \infty} r_0(N_i) = \infty \ and \ \lim_{i \to \infty} \int_{N_i} H^2 d\sigma = 16\pi.$$

And suppose that N_i separates the infinity from the compact part. Then

(5.6)
$$\lim_{i \to \infty} \frac{r_0(N_i)}{r_1(N_i)} = 1.$$

Proof. We may apply Lemma 3.1 for the blow-down

$$\tilde{N} = \frac{1}{2}HN = \{\frac{1}{2}Hx : x \in N\}.$$

If the surfaces \tilde{N}_i stay away from the origin, i.e.

$$0 < C \le H_i^{-1} r_0(N_i)$$

for some positive constants C, then a subsequence of \tilde{N}_i converges to a sphere $S^2(a)$ radius 1 and centered at $a \in R^3$ in C^{∞} by the curvature estimates Theorem 2.5 in Section 2. Also notice that (2.6) implies that the blow-down surfaces \tilde{N}_i always stay within a bounded region in R^3 . On one hand, by (5.5) in Lemme 5.1, we have

(5.7)
$$\frac{1}{4\pi} \int_{S^2(a)} \frac{\nu \cdot b}{|x|} d\sigma + \frac{1}{4\pi} \int_{S^2(a)} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma = 0,$$

for any b. On the other hand, if take $b = -\frac{a}{|a|}$, then

(5.8)
$$\frac{1}{4\pi} \int_{S^2(a)} \frac{\nu \cdot b}{|x|} d\sigma + \frac{1}{4\pi} \int_{S^2(a)} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma = |a|$$

due to an explicit calculation when the origin is inside. Therefore

$$a = 0$$
 and $\lim_{i \to \infty} \frac{1}{2} r_0(N_i) H(N_i) = \lim_{i \to \infty} \frac{r_0(N_i)}{r_1(N_i)} = 1.$

To conclude this is all that can happen we need only to exclude the case when

(5.9)
$$\lim_{i \to \infty} H_i^{-1} r_0(N_i) = 0.$$

Assume otherwise, according to Lemma 3.1, the blow-down sequence \tilde{N}_i converges to a unit round sphere $S^2(a)$ centered at $a \in R^3$ with |a| = 1 in Hausdorff topology. We will take $b = -\frac{a}{|a|}$. From Lemma 5.1, we know

(5.10)
$$\lim_{i \to \infty} \left(\int_{\tilde{N}_{\epsilon}} \frac{\nu \cdot b}{|x|} d\sigma + \int_{\tilde{N}_{\epsilon}} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma \right) = 0.$$

But, we claim, on the other hand,

(5.11)
$$\lim_{i \to \infty} \left(\int_{\tilde{N}_i} \frac{\nu \cdot b}{|x|} d\sigma + \int_{\tilde{N}_i} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma \right) = 4\pi$$

which gives us the contradiction. First, we have from explicit calculations

(5.12)
$$\int_{S^{2}(a)} \frac{\nu \cdot b}{|x|} d\sigma = \frac{4}{3}\pi, \ \int_{S^{2}(a)} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} d\sigma = \frac{2}{3}\pi.$$

The first term in (5.11) is an easy term because the uniform integrability

(5.13)
$$\lim_{i \to \infty} \int_{\tilde{N}_i} \frac{\nu \cdot b}{|x|} d\sigma = \int_{S^2(a)} \frac{\nu \cdot b}{|x|} d\sigma = \frac{4}{3}\pi.$$

To deal with the second term in (5.11), we break up the integral into three parts. For any fixed small number s > 0 and large number K > 0,

(5.14)
$$\int_{\tilde{N}_{i}} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} d\sigma = \int_{\tilde{N}_{i} \cap B_{s}^{c}(0)} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} d\sigma + \int_{\tilde{N}_{i} \cap (B_{s}(0) \setminus B_{KHr_{0}}(0))} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} d\sigma + \int_{\tilde{N}_{i} \cap B_{KHr_{0}}(0)} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} d\sigma.$$

Then

$$(5.15) \qquad \lim_{i \to \infty} \int_{\tilde{N}_i \cap B_c^c(0)} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma = \int_{S^2(a) \cap B_c^c} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma$$

and

$$\lim_{i\to\infty}\int_{\tilde{N}_i\bigcap\,B_{KH(N_i)r_0(N_i)}(0)}\frac{(\nu\cdot x)(\nu\cdot b)}{|x|^3}d\sigma=\int_{P\bigcap\,B_K(0)}\frac{(\nu\cdot x)(\nu\cdot b)}{|x|^3}d\sigma,$$

where P is the limit plane in Lemma 3.2. By Corollary 4.4, we know

$$\int_{P} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma = 2\pi.$$

due to a simple calculation. Notice that

$$\int_{\tilde{N}_i} \frac{\nu \cdot x}{|x|^3} d\sigma = 4\pi$$

for any i and

$$\int_{S^2(a)} \frac{\nu \cdot x}{|x|^3} d\sigma = 2\pi$$

because the origin is on the sphere $S^2(a)$. Since

(5.18)
$$\lim_{i \to \infty} \int_{\tilde{N}_i \cap B_c^c(0)} \frac{\nu \cdot x}{|x|^3} d\sigma = \int_{S^2(a) \cap B_c^c(0)} \frac{\nu \cdot x}{|x|^3} d\sigma,$$

(5.19)
$$\lim_{i \to \infty} \int_{\tilde{N}_i \cap B_{KH_{r_0}}(0)} \frac{\nu \cdot x}{|x|^3} d\sigma = \int_{P \cap B_K(0)} \frac{\nu \cdot x}{|x|^3} d\sigma$$

and

$$\int_{P} \frac{\nu \cdot x}{|x|^3} d\sigma = 2\pi,$$

we know

(5.21)
$$\lim_{i \to \infty, s \to 0, K \to \infty} \int_{\tilde{N}_i \cap (B_s^c(0) \setminus B_{KH_{r_0}}(0))} \frac{\nu \cdot x}{|x|^3} d\sigma = 0.$$

Now we are ready to handle the difficult term: the integral over the transition region in (5.14). Our goal is to show that

(5.22)
$$\lim_{i \to \infty, s \to 0, K \to \infty} \int_{\tilde{N}_i \cap (B_s(0) \setminus B_{KHr_0}(0))} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^3} d\sigma = 0.$$

The key point is to use Corollary 4.5 to prove (5.22) from (5.21). Let ν_i be chosen as in Corollary 4.5. Then

(5.23)
$$\int_{\tilde{N}_{i} \cap (B_{s} \setminus B_{KHr_{0}})} \frac{(\nu \cdot x)(\nu \cdot b)}{|x|^{3}} =$$

$$(\nu_{i} \cdot b) \int_{\tilde{N}_{i} \cap (B_{s} \setminus B_{KHr_{0}})} \frac{\nu \cdot x}{|x|^{3}} + \int_{\tilde{N}_{i} \cap (B_{s} \setminus B_{KHr_{0}})} \frac{(\nu \cdot x)((\nu - \nu_{i}) \cdot b)}{|x|^{3}}.$$

Hence we only need to deal with the second term on the right side of the above (5.23). We are better now to use the cylindrical coordinates used in Section 4.

$$\int_{\tilde{N}_{i} \cap (B_{s}(0) \setminus B_{KHr_{0}}(0))} \frac{(\nu \cdot x)((\nu - \nu_{i}) \cdot b)}{|x|^{3}} d\sigma$$

$$= \sum_{j=1}^{n_{i}} \int_{I_{j}} \frac{(\nu \cdot x)((\nu - \nu_{i}) \cdot b)}{|x|^{3}} A(t) d\theta dt$$

$$\leq C \sum_{j=1}^{n_{i}} L \max_{I_{j}} |\nu - \nu_{i}|$$

$$= C \sum_{j=1}^{n_{i}/2} L \max_{I_{j}} |\nu - \nu_{i}| + C \sum_{j=n_{i}/2+1}^{n_{i}/2} L \max_{I_{j}} |\nu - \nu_{i}|.$$

iFrom (4.12) and (4.13), we have

$$\int_{\tilde{N}_{i} \cap (B_{s}(0) \setminus B_{KHr_{0}}(0))} \frac{(\nu \cdot x)((\nu - \nu_{i}) \cdot b)}{|x|^{3}} d\sigma$$

$$\leq C \eta \left(\sum_{j=1}^{n_{i}/2} (e^{-\frac{1}{4}Lj} + e^{-\frac{1}{8}Ln_{i}}) + \sum_{j=1}^{n_{i}/2} (e^{-\frac{1}{8}Ln_{i}} + e^{-\frac{1}{4}(n_{i}-j)L}) \right)$$

$$\leq C \eta (n_{i}e^{-\frac{1}{8}n_{i}L} + 2),$$

where

$$\eta = \int_{B_{sH^{-1}(N_i)} \cap N_i} |\nabla \nu|^2 d\sigma + r_0^{-1}$$

and η can be arbitrarily small as long as $s \to 0$ and $r_0(N_i) \to \infty$. Thus (5.22) is proved and the proof of the theorem is completed.

Corollary 5.3. Suppose $(R^3 \setminus B_1(0), g)$ is an asymptotically flat end with positive mass. Then there exist a large number K > 0 and a small number $\epsilon > 0$ such that, for any $H < \epsilon$, there exists a unique stable spheres N of constant mean curvature H with $N \subset R^3 \setminus B_K(0)$ and separates the infinity from the compact part. Hence there exists a unique foliation of stable spheres of constant mean curvature near the infinity.

Proof. In the light of Proposition 5.3 in [HY] we know

(5.26)
$$\lim_{i \to \infty} \int_{N_i} H^2 d\mu = 16\pi,$$

provided each N_i is a stable sphere of constant mean curvature. Thus Corollary 5.3 follows from Theorem 5.2 in the above, Theorem 4.1 in [HY] and the proof of Theorem 5.1 in [HY].

References:

- [ADM] R. Amowitt, S. Deser and C. Misner, Coordinate invariant and energy expression in general relativity, Phys. Rev. 122 (1966) 997-1006.
 - [B] R. Barnik, The mass of an asymptotically flat manifold, Comm. Pure. Appl. Math. 39 (1986) 661-693.
 - [Br] H. Bray, The Penrose inequality in general relativity and volume comparison theorem involving scalar cutvature, Stanford Thesis 1997.
 - [MS] J. Michael and L. Simon, Sobolev and mean-value inequalities on general submanifolds of \mathbb{R}^n , Comm. Pure. Appl. Math. 26 (1973) 361-379.
 - [HY] G. Huisken and S. T. Yau, Definition of center of mass for isolated physical systems and unique foliations by stable spheres of constant mean curvature, Invent. Math. 124 (1996) 281-311.
 - [Ka] K. Kenmotsu, "Surfaces with constant mean curvature", Tanslations of Math. Monographs, v. 221, AMS, 2003.
 - [QT] Jie Qing and Gang Tian, Bubbling of the heat flow for harmonic maps from surfaces, Comm. Pure and Appl. Math., Vol. L (1997), 295-310.
 - [SSY] R. Schoen, L. Simon and S.T. Yau, Curvature estimates for minimal hypersurfaces, Acta Math. 134 (1975) 275-288.
 - [Si] L. Simon, Existence of Willmore surface, Proc. Centre for Math. Anal. 10 (1985) 187-216.

- [Sj] J. Simons, Minimal Varieties in Riemannian manifolds, Ann. of Math. 88 (1968) 62-105.
- [Ye] R. Ye, Foliation by constant mean curvature spheres on asymptotically flat manifolds, Geom. Anal. and the Calculus of Variations, 369-383, Intern. Press, Cambridge, MA, 1996.

JIE QING, DEPT. OF MATH., UC, SANTA CRUZ, SANTA CRUZ, CA 95064 $E\text{-}mail\ address:}$ qing@ucsc.edu

GANG TIAN, DEPT. OF MATH., PRINCETON UNIVERSITY, PRINCETON, NJ 08544 *E-mail address*: tian@math.princeton.edu