DIRECT IMAGE OF LOGARITHMIC COMPLEXES AND INFINITESIMAL INVARIANTS OF CYCLES

MORIHIKO SAITO

ABSTRACT. We show that the direct image of the filtered logarithmic de Rham complex is a direct sum of filtered logarithmic complexes with coefficients in variations of Hodge structures, using a generalization of the decomposition theorem of Beilinson, Bernstein and Deligne to the case of filtered *D*-modules. The advantage of using the logarithmic complexes is that we have the strictness of the Hodge filtration by Deligne after taking the cohomology group in the projective case. As a corollary, we get the total infinitesimal invariant of a (higher) cycle in a direct sum of the cohomology of filtered logarithmic complexes with coefficients, and this is essentially equivalent to the cohomology class of the cycle.

Introduction

Let X, S be complex manifolds or smooth algebraic varieties over a field of characteristic zero. Let $f: X \to S$ be a projective morphism, and D be a divisor on S such that f is smooth over $S \setminus D$. We have a filtered locally free \mathcal{O} -module (V^i, F) on $S \setminus D$ underlying a variation of Hodge structure whose fiber V_s^i at $s \in S \setminus D$ is the cohomology of the fiber $H^i(X_s, \mathbb{C})$. If D is a divisor with normal crossings on S, let \widetilde{V}^i denote the Deligne extension [7] of V^i such that the the eigenvalues of the residue of the connection are contained in [0, 1). The Hodge filtration F is naturally extended to \widetilde{V}^i by [25]. We have the logarithmic de Rham complex

$$\mathrm{DR}_{\mathrm{log}}(\widetilde{V}^i) = \Omega_S^{\bullet}(\log D) \otimes_{\mathcal{O}} \widetilde{V}^i,$$

which has the Hodge filtration F^p defined by $\Omega^j_S(\log D) \otimes_{\mathcal{O}} F^{p-j}\widetilde{V}^i$. In general, V^i can be extended to a regular holonomic \mathcal{D}_S -module M^i on which a local defining equation of D acts bijectively. By [23], M^i and hence the de Rham complex $\mathrm{DR}(M^i)$ have the Hodge filtration F. If $Y := f^*D$ is a divisor with normal crossings on X, then $\Omega^{\bullet}_X(\log Y)$ has the Hodge filtration F defined by the truncation σ (see [8]) as usual, i.e. $F^p\Omega^{\bullet}_X(\log Y) = \Omega^{\bullet \geq p}_X(\log Y)$.

Theorem 1. Assume $Y = f^*D$ is a divisor with normal crossings. There is an increasing split filtration L on the filtered complex $\mathbf{R}f_*(\Omega_X^{\bullet}(\log Y), F)$ such that we have noncanonical and canonical isomorphisms in the filtered derived category:

$$\mathbf{R}f_*(\Omega_X^{\bullet}(\log Y), F) \simeq \bigoplus_{i \in \mathbb{Z}} (\mathrm{DR}(M^i), F)[-i],$$
$$\mathrm{Gr}_i^L \mathbf{R}f_*(\Omega_X^{\bullet}(\log Y), F) = (\mathrm{DR}(M^i), F)[-i].$$

Date: Aug. 28, 2005, v.4.

If D is a divisor with normal crossings, we have also

$$\mathbf{R} f_*(\Omega_X^{\bullet}(\log Y), F) \simeq \bigoplus_{i \in \mathbb{Z}} (\mathrm{DR}_{\log}(\widetilde{V}^i), F)[-i],$$
$$\mathrm{Gr}_i^L \mathbf{R} f_*(\Omega_X^{\bullet}(\log Y), F) = (\mathrm{DR}_{\log}(\widetilde{V}^i), F)[-i].$$

This follows from the decomposition theorem (see [2]) extended to the case of the direct image of (\mathcal{O}_X, F) as a filtered \mathcal{D} -module, see [22]. Note that Hodge modules do not appear in the last statement if D is a divisor with normal crossings. The assertion becomes more complicated in the non logarithmic case, see Remark (i) in (2.5). A splitting of the filtration L is given by choosing the first noncanonical isomorphism in the filtered decomposition theorem, see (1.4.2). A canonical choice of the splitting is given by choosing an relatively ample class, see [9].

Let $\operatorname{CH}^p(X \setminus Y, n)$ be Bloch's higher Chow group, see [3]. In the analytic case, we assume for simplicity that $f:(X,Y) \to (S,D)$ is the base change of a projective morphism of smooth complex algebraic varieties $f':(X',Y') \to (S',D')$ by an open embedding of complex manifolds $S \to S'_{an}$, and an element of $\operatorname{CH}^p(X \setminus Y,n)$ is the restriction of an element of $\operatorname{CH}^p(X' \setminus Y',n)$ to $X \setminus Y$. If n=0, we may assume that it is the restriction of an analytic cycle of codimension p on X. From Theorem 1, we can deduce

Corollary 1. With the above notation and assumption, let $\xi \in CH^p(X \setminus Y, n)$. Then, choosing a splitting of the filtration L in Theorem 1 (or more precisely, choosing the first noncanonical isomorphism in the filtered decomposition theorem (1.4.2)), we have the total infinitesimal invariant

$$\delta_{S,D}(\xi) = (\delta_{S,D}^{i}(\xi)) \in \bigoplus_{i \geq 0} \mathbf{H}^{i}(S, F^{p} \mathrm{DR}(M^{2p-n-i})),$$

$$(resp. \ \overline{\delta}_{S,D}(\xi) = (\overline{\delta}_{S,D}^{i}(\xi)) \in \bigoplus_{i \geq 0} \mathbf{H}^{i}(S, \mathrm{Gr}_{F}^{p} \mathrm{DR}(M^{2p-n-i})),)$$

where $\delta^i_{S,D}(\xi)$ (resp. $\overline{\delta}^i_{S,D}(\xi)$) is independent of the choice of a splitting if the $\delta^j_{S,D}(\xi)$ (resp. $\overline{\delta}^j_{S,D}(\xi)$) vanish for j < i. In the case D is a divisor with normal crossings, the assertion holds with $DR(M^{2p-n-i})$ replaced by $DR_{log}(\widetilde{V}^{2p-n-i})$.

This shows that the infinitesimal invariants in [14], [13], [27], [5], [1], [24] can be defined naturally in the cohomology of filtered logarithmic complexes with coefficients in variations of Hodge structures if D is a divisor with normal crossings, see (2.4) for the compatibility with [1]. Note that if S is Stein or affine, then $\mathbf{H}^i(S, F^p\mathrm{DR}_{\log}(\widetilde{V}^q))$ is the i-th cohomology group of the complex whose j-th component is $\Gamma(S, \Omega_S^j(\log D) \otimes_{\mathcal{O}} F^{p-j}\widetilde{V}^q)$. If D is empty, then an inductive definition of $\delta_{S,D}^i(\xi)$, $\overline{\delta}_{S,D}^i(\xi)$ was given by Shuji Saito [24] using the filtered Leray spectral sequence together with the E_2 -degeneration argument in [6]. He also showed that the infinitesimal invariants depend only on the cohomology class of the cycle. If S is projective, then it follows from [8] that the total infinitesimal invariant $(\delta_{S,D}^i(\xi))$ is equivalent to the cycle class of ξ in $H_{\mathrm{DR}}^{2p-n}(X \setminus Y)$ by the strictness of the Hodge filtration, and the filtration L comes from the Leray filtration on the cohomology of $X \setminus Y$, see Remark (iii) in (2.5).

Corollary 1 is useful to study the behavior of the infinitesimal invariants near the boundary of the variety. If D is empty, let $\delta_S^i(\xi)$ denote $\delta_{S,D}^i(\xi)$. We can define $\delta_{DR,S}^i(\xi)$ as in [19] by omitting F^p before DR in Corollary 1 where $D = \emptyset$.

Corollary 2. Assume S is projective. Let $U = S \setminus D$. Then for each $i \geq 0$, $\delta_{S,D}^{i}(\xi)$, $\overline{\delta}_{S,D}^{i}(\xi)$, $\delta_{U}^{i}(\xi)$ and $\delta_{\mathrm{DR},U}^{i}(\xi)$ are equivalent to each other, i.e. one of them vanishes if and only if the others do.

Indeed, $(\delta_{\mathrm{DR},U}^i(\xi))$ is determined by $(\delta_U^i(\xi))$, and $(\delta_U^i(\xi))$ by $(\delta_{S,D}^i(\xi))$. Moreover, $(\delta_{S,D}^i(\xi))$ is equivalent to $(\delta_{\mathrm{DR},U}^i(\xi))$ by the strictness of the Hodge filtration [8] applied to (X,Y) together with Theorem 1, see (2.3). For the relation with $\overline{\delta}_{S,D}^i(\xi)$, see (2.1). Note that the equivalence between $\delta_U^i(\xi)$ and $\delta_{\mathrm{DR},U}^i(\xi)$ in the case of algebraic cycles (i.e. n=0) was first found by J.D. Lewis and Shuji Saito in [19] (assuming a conjecture of Brylinski and Zucker and the Hodge conjecture and using an L^2 -argument). The above arguments seem to be closely related with their question, see also Remark (i) in (2.5) below.

As another corollary of Theorem 1 we have

Corollary 3. Assume f induces an isomorphism over $S \setminus D$, and $Y = f^*D$ is a divisor with normal crossings on X. Then

$$R^i f_* \Omega_X^p(\log Y) = 0$$
 if $i + p > \dim X$.

This follows immediately from Theorem 1 since $M^i = 0$ for $i \neq 0$. Corollary 3 is an analogue of the vanishing theorem of Kodaira-Nakano. However, this does not hold for a non logarithmic complex (e.g. if f is a blow-up with a point center). This corollary was inspired by a question of A. Dimca.

I would like to thank Dimca, Lewis and Shuji Saito for good questions and useful suggestions.

In Section 1, we prove Theorem 1 after reviewing some basic facts on filtered differential complexes. In Section 2 we explain the application of Theorem 1 to the infinitesimal invariants of (higher) cycles. In Section 3 we give some examples using Lefschetz pencils.

1. Direct image of logarithmic complexes

1.1. Filtered differential complexes. Let X be a complex manifold or a smooth algebraic variety over a field of characteristic zero. Let $D^bF(\mathcal{D}_X)$ (resp. $D^bF(\mathcal{D}_X)^r$) be the bounded derived category of filtered left (resp. right) \mathcal{D}_X -modules. Let $D^bF(\mathcal{O}_X, \mathrm{Diff})$ be the bounded derived category of filtered differential complexes (L, F) where F is exhaustive and locally bounded below (i.e. $F_p = 0$ for $p \ll 0$ locally on X), see [22], 2.2. We have an equivalence of categories

(1.1.1)
$$DR^{-1}: D^b F(\mathcal{O}_X, Diff) \to D^b F(\mathcal{D}_X)^r,$$

whose quasi-inverse is given by the de Rham functor DR^r for right \mathcal{D} -modules, see (1.2) below. Recall that, for a filtered \mathcal{O}_X -module (L, F), the associated filtered right \mathcal{D} -module $DR^{-1}(L, F)$ is defined by

(1.1.2)
$$\mathrm{DR}^{-1}(L,F) = (L,F) \otimes_{\mathcal{O}} (\mathcal{D},F),$$

and the morphisms $(L, F) \to (L', F)$ in $MF(\mathcal{O}_X, \text{Diff})$ correspond bijectively to the morphisms of filtered \mathcal{D} -modules $DR^{-1}(L, F) \to DR^{-1}(L', F)$. More precisely, the condition on $(L, F) \to (L', F)$ is that the composition

$$F_pL \to L \to L' \to L'/F_qL'$$

is a differential operator of order $\leq p-q-1$. The proof of (1.1.1) can be reduced to the canonical filtered quasi-isomorphism for a filtered right \mathcal{D} -module (M, F)

$$DR^{-1} \circ DR^r(M, F) \to (M, F),$$

which follows from a calculation of a Koszul complex.

Note that the direct image f_* of filtered differential complexes is defined by the sheaf-theoretic direct image $\mathbf{R}f_*$, and this direct image is compatible with the direct image f_* of filtered \mathcal{D} -modules via (1.1.1), see [22], 2.3. So we get

$$(1.1.3) \mathbf{R}f_* = \mathrm{DR}^r \circ f_* \circ \mathrm{DR}^{-1} : D^b F(\mathcal{O}_X, \mathrm{Diff}) \to D^b F(\mathcal{O}_S, \mathrm{Diff}),$$

where we use DR^r for right \mathcal{D} -modules (otherwise there is a shift of complex).

1.2. De Rham complex. The de Rham complex $DR^r(M, F)$ of a filtered right \mathcal{D} -module (M, F) is defined by

$$(1.2.1) (DRr(M, F))i = \bigwedge^{-i} \Theta_X \otimes_{\mathcal{O}} (M, F[-i]) for i \le 0.$$

Here $(F[-i])_p = F_{p+i}$ in a compatible way with $(F[-i])^p = F^{p-i}$ and $F_p = F^{-p}$. Recall that the filtered right \mathcal{D} -module associated with a filtered left \mathcal{D} -module (M, F) is defined by

$$(1.2.2) (M,F)^r := (\Omega_X^{\dim X}, F) \otimes_{\mathcal{O}} (M,F),$$

where $\operatorname{Gr}_p^F \Omega_X^{\dim X} = 0$ for $p \neq -\dim X$. This induces an equivalence of categories between the left and right \mathcal{D} -modules. The usual de Rham complex $\operatorname{DR}(M,F)$ for a left \mathcal{D} -module is defined by

$$(1.2.3) (DR(M, F))^i = \Omega_X^i \otimes_{\mathcal{O}} (M, F[-i]) for i \ge 0,$$

and this is compatible with (1.2.1) via (1.2.2) up to a shift of complex, i.e.

(1.2.4)
$$\operatorname{DR}(M, F) = \operatorname{DR}^{r}(M, F)^{r}[-\dim X].$$

1.3. Logarithmic complex. Let X be as in (1.1), and Y be a divisor with normal crossings on X. Let (V, F) be a filtered locally free \mathcal{O} -module underlying a polarizable variation of Hodge structure on $X \setminus Y$. Let (\tilde{V}, F) be the Deligne extension of (V, F) to X such that the eigenvalues of the residue of the connection are contained in [0, 1). Then we have the filtered logarithmic de Rham complex $\mathrm{DR}_{\log}(\tilde{V}, F)$ such that F^p of its i-th component is

$$\Omega_X^i(\log Y) \otimes F^{p-i}\widetilde{V}.$$

If
$$(M, F) = (\mathcal{O}_X, F)$$
 with $\operatorname{Gr}_p^F \mathcal{O}_X = 0$ for $p \neq 0$, then $\operatorname{DR}_{\operatorname{log}}(\mathcal{O}_X, F) = (\Omega_X^{\bullet}(\log X), F).$

Let $\widetilde{V}(*Y)$ be the localization of \widetilde{V} by a local defining equation of Y. This is a regular holonomic left \mathcal{D}_X -module underlying a mixed Hodge module, and has the Hodge filtration F which is generated by the Hodge filtration F on \widetilde{V} , i.e.

$$F_p\widetilde{V}(*Y) = \sum_{\nu} \partial^{\nu} F^{-p+|\nu|} \widetilde{V},$$

where $F_p = F^{-p}$ and $\partial^{\nu} = \prod_i \partial_i^{\nu_i}$ with $\partial_i = \partial/\partial x_i$. Here (x_1, \dots, x_n) is a local coordinate system such that Y is contained in $\{x_1 \cdots x_n = 0\}$. By [23], 3.11, we have a filtered quasi-isomorphism

(1.3.1)
$$\operatorname{DR}_{\log}(\widetilde{V}, F) \xrightarrow{\sim} \operatorname{DR}(\widetilde{V}(*Y), F).$$

This generalizes the filtered quasi-isomorphism in [7]

$$(1.3.2) \qquad (\Omega_X^{\bullet}(\log Y), F) \xrightarrow{\sim} \mathrm{DR}(\mathcal{O}_X(*Y), F).$$

Note that the direct image of the filtered \mathcal{D}_X -module $(\tilde{V}(*Y), F)$ by $X \to pt$ in the case X projective (or proper algebraic) is given by the cohomology group of the de Rham complex $\mathrm{DR}(\tilde{V}(*Y), F)$ (up to a shift of complex) by definition, and the Hodge filtration F on the direct image is strict by the theory of Hodge modules. So we get

(1.3.3)
$$F^{p}\mathbf{H}^{i}(X \setminus Y, \mathrm{DR}(V)) := F^{p}\mathbf{H}^{i}(X, \mathrm{DR}(\widetilde{V}(*Y)))$$
$$= \mathbf{H}^{i}(X, F^{p}\mathrm{DR}(\widetilde{V}(*Y))) = \mathbf{H}^{i}(X, F^{p}\mathrm{DR}_{\mathrm{log}}(\widetilde{V})).$$

1.4. Decomposition theorem. Let $f: X \to S$ be a projective morphism of complex manifolds or smooth algebraic varieties over a field of characteristic zero. Then the decomposition theorem of Beilinson, Bernstein and Deligne [2] is extended to the case of Hodge modules ([22], [23]), and we have noncanonical and canonical isomorphisms

(1.4.1)
$$f_*(\mathcal{O}_X, F) \simeq \bigoplus_{j \in \mathbb{Z}} \mathcal{H}^j f_*(\mathcal{O}_X, F)[-j] \quad \text{in } D^b F(\mathcal{D}_S),$$
$$\mathcal{H}^j f_*(\mathcal{O}_X, F) = \bigoplus_{Z \subset S} (M_Z^j, F) \quad \text{in } MF(\mathcal{D}_S),$$

where Z are irreducible closed analytic or algebraic subsets of S, and (M_Z^j, F) are filtered \mathcal{D}_S -modules underlying a pure Hodge module of weight $j + \dim X$ and with strict support Z, i.e. M_Z^j has no nontrivial sub nor quotient module whose support is strictly smaller than Z. (Here $MF(\mathcal{D}_S)$ denotes the category of filtered left \mathcal{D}_S -modules.) Indeed, the second canonical isomorphism follows from the strict support decomposition which is part of the definition of pure Hodge modules, see [22], 5.1.6. The first noncanonical isomorphism follows from the strictness of the Hodge filtration and the relative hard Lefschetz theorem for the direct image (see [22], 5.3.1) using the E_2 -degeneration argument in [6] together with the equivalence of categories $D^bF(\mathcal{D}_S) \simeq D^bG(\mathcal{B}_S)$. Here $\mathcal{B}_S = \bigoplus_{i \in \mathbb{N}} F_i\mathcal{D}_S$ and $D^bG(\mathcal{B}_S)$ is the derived category of bounded complexes of graded left \mathcal{B}_S -modules M_{\bullet}^{\bullet} such that $M_i^j = 0$ for $i \ll 0$ or $|j| \gg 0$, see [22], 2.1.12. We need a derived category associated

to some abelian category in order to apply the argument in [6] (see also [9]). In the algebraic case, we can also apply [6] to the derived category of mixed Hodge modules on S and it is also possible to use [23], 4.5.4 to show the first noncanonical isomorphism.

If f is smooth over the complement of a divisor $D \subset S$ and $Y := f^*D$ is a divisor with normal crossings, then the filtered direct image $f_*(\mathcal{O}_X(*Y), F)$ is strict (see [23], 2.15), and we have noncanonical and canonical isomorphisms

(1.4.2)
$$f_*(\mathcal{O}_X(*Y), F) \simeq \bigoplus_{j \in \mathbb{Z}} \mathcal{H}^j f_*(\mathcal{O}_X(*Y), F)[-j] \quad \text{in } D^b F(\mathcal{D}_S),$$
$$\mathcal{H}^j f_*(\mathcal{O}_X(*Y), F) = (M_S^j(*D), F) \quad \text{in } MF(\mathcal{D}_S).$$

Here $(M_S^j(*D), F)$ is the 'localization' of (M_S^j, F) along D which is the direct image of $(M_S^j, F)|_U$ by the open embedding $U := S \setminus D \to S$ in the category of filtered \mathcal{D} -modules underlying mixed Hodge modules. (By the Riemann-Hilbert correspondence, this gives the direct image in the category of complexes with constructible cohomology because D is a divisor.) The Hodge filtration F on the direct image is determined by using the V-filtration of Kashiwara and Malgrange, and $(M_S^j(*D), F)$ is the unique extension of $(M_S^j, F)|_U$ which underlies a mixed Hodge module on S and whose underlying \mathcal{D}_S -module is the direct image in the category of regular holonomic \mathcal{D}_S -modules, see [23], 2.11. So the second canonical isomorphism follows because the left-hand side satisfies these conditions. (Note that (M_Z^j, F) for $Z \neq S$ vanishes by the localization, because $Z \subset D$ if $(M_Z^j, F) \neq 0$.) The first noncanonical isomorphism follows from the strictness of the Hodge filtration and the relative hard Lefschetz theorem by the same argument as above.

1.5. Proof of Theorem 1. Let $r = \dim X - \dim S$. By (1.1.3), (1.3.2) and (1.4.2), we have isomorphisms

(1.5.1)
$$\mathbf{R} f_*(\Omega_X^{\bullet}(\log Y), F) = \mathrm{DR}^r \circ f_* \circ \mathrm{DR}^{-1}(\Omega_X^{\bullet}(\log Y), F)$$
$$= \mathrm{DR}^r \circ f_*(\mathcal{O}_X(*Y), F)[-\dim X]$$
$$\simeq \bigoplus_{i \in \mathbb{Z}} \mathrm{DR}(M_S^i(*D), F)[-r - i],$$

where the shift of complex by r follows from the difference of the de Rham complex for left and right \mathcal{D} -modules. Furthermore, letting L be the filtration induced by τ on the complex of filtered \mathcal{D}_S -modules $f_*(\mathcal{O}_X(*Y), F)[-r]$, we have a canonical isomorphism

(1.5.2)
$$\operatorname{Gr}_{i}^{L} f_{*}(\mathcal{O}_{X}(*Y), F)[-r] = (M_{S}^{i-r}(*D), F)[-i],$$

and the first assertion follows by setting $M^i = M_S^{i-r}(*D)$. The second assertion follows from the first by (1.3.1). This competes the proof of Theorem 1.

2. Infinitesimal invariants of cycles

2.1. Cycle classes. Let X be a complex manifold, and $\mathcal{C}^{\bullet,\bullet}$ denote the double complex of vector spaces of currents on X. The associated single complex is denoted by \mathcal{C}^{\bullet} . Let F be the Hodge filtration by the first index of $\mathcal{C}^{\bullet,\bullet}$ (using the truncation

 σ in [8]). Let ξ be an analytic cycle of codimension p on X. Then it is well known that ξ defines a closed current in $F^p\mathcal{C}^{2p}$ by integrating the restrictions of C^{∞} forms with compact supports on X to the smooth part of the support of ξ (and using a triangulization or a resolution of singularities of the cycle). So we have a cycle class of ξ in $H^{2p}(X, F^p\Omega_X^{\bullet})$.

Assume X is a smooth algebraic variety over a field k of characteristic zero. Then the last assertion still holds (where Ω_X^{\bullet} means $\Omega_{X/k}^{\bullet}$), see [11]. Moreover, for the higher Chow groups, we have the cycle map (see [4], [10], [12], [15], [16])

$$cl: \mathrm{CH}^p(X,n) \to F^p H^{2p-n}_{\mathrm{DR}}(X),$$

where the Hodge filtration F is defined by using a smooth compactification of X whose complement is a divisor with normal crossings, see [8]. This cycle map is essentially equivalent to the cycle map to $\operatorname{Gr}_F^p H^{2p-n}_{\operatorname{DR}}(X)$ because we can reduce to the case $k=\mathbb{C}$ where we have the cycle map

$$cl: CH^p(X, n) \to Hom_{MHS}(\mathbb{Q}, H^{2p-n}(X, \mathbb{Q})(p)),$$

and morphisms of mixed Hodge structures are strictly compatible with the Hodge filtration F.

2.2. Proof of Corollary 1. By (2.1) ξ has the cycle class in

$$H^{2p-n}(X, F^p\Omega_X^{\bullet}(\log Y)).$$

By theorem 1, this gives the total infinitesimal invariant

$$\delta_{S,D}(\xi) = (\delta_{S,D}^{2p-n-i}(\xi)) \in \bigoplus_{i \in \mathbb{Z}} \mathbf{H}^{2p-n-i}(S, F^p \mathrm{DR}(M^i)),$$

and similarly for $\overline{\delta}_{S,D}(\xi)$. So the assertion follows.

2.3. Proof of Corollary 2. Choosing the first noncanonical isomorphism in the filtered decomposition theorem (1.4.2), we get canonical morphisms compatible with the direct sum decompositions

$$\bigoplus_{i\geq 0} \mathbf{H}^{i}(S, F^{p}\mathrm{DR}(M^{q-i})) \to \bigoplus_{i\geq 0} \mathbf{H}^{i}(S\setminus D, F^{p}\mathrm{DR}(M^{q-i}))$$
$$\to \bigoplus_{i\geq 0} \mathbf{H}^{i}(S\setminus D, \mathrm{DR}(M^{q-i})),$$

and these are identified with the canonical morphisms

$$\mathbf{H}^{q}(X, F^{p}\Omega_{X}^{\bullet}(\log Y)) \to \mathbf{H}^{q}(X \setminus Y, F^{p}\Omega_{X \setminus Y}^{\bullet})$$
$$\to \mathbf{H}^{q}(X \setminus Y, \Omega_{X \setminus Y}^{\bullet}).$$

By Deligne [8], the composition of the last two morphisms is injective because of the strictness of the Hodge filtration, see also (1.3). So we get the equivalence of $\delta_{S,D}^{i}(\xi)$, $\delta_{U}^{i}(\xi)$, $\delta_{\mathrm{DR},U}^{i}(\xi)$. The equivalence with $\overline{\delta}_{S,D}^{i}(\xi)$ follows from (2.1).

2.4. Compatibility with the definition in [1]. When D is empty, the infinitesimal invariants are defined in [1] by using the extension groups of filtered \mathcal{D} -modules together with the forgetful functor from the category of mixed Hodge modules to that of filtered \mathcal{D} -modules. Its compatibility with the definition in this paper follows from the equivalence of categories (1.1.1) and the compatibility of the direct image functors (1.1.3).

Note that for $(L, F) \in D^b F(\mathcal{O}_X, \text{Diff})$ in the notation of (1.1), we have a canonical isomorphism

(2.4.1)
$$\operatorname{Ext}^{i}((\Omega_{X}^{\bullet}, F), (L, F)) = \mathbf{H}^{i}(X, F_{0}L),$$

where the extension group is taken in $D^bF(\mathcal{O}_X, \text{Diff})$. Indeed, the left-hand side is canonically isomorphic to

$$\operatorname{Ext}^{i}(\operatorname{DR}^{-1}(\Omega_{X}^{\bullet}, F), \operatorname{DR}^{-1}(L, F))$$

$$= \mathbf{H}^{i}(X, F_{0}\mathcal{H}om_{\mathcal{D}}(\operatorname{DR}(\mathcal{D}_{X}, F), \operatorname{DR}^{-1}(L, F))),$$

$$= \mathbf{H}^{i}(X, F_{0}\operatorname{DR}^{r}\operatorname{DR}^{-1}(L)),$$

and the last group is isomorphic to the right-hand side of (2.4.1) which is independent of a representative of (L, F). If X is projective, then this assertion follows also from the adjoint relation for filtered \mathcal{D} -modules.

If X is smooth projective and Y is a divisor with normal crossings, then the cycle class can be defined in

$$\operatorname{Ext}^{2p}((\Omega_X^{\bullet}, F), \Omega_X^{\bullet}(\log Y), F[p])) = \mathbf{H}^{2p}(X, F^{p}\Omega_X^{\bullet}(\log Y))$$
$$= F^{p}\mathbf{H}^{2p}(X, \Omega_X^{\bullet}(\log Y)).$$

2.5. Remarks. (i) If we use (1.4.1) instead of (1.4.2) we get an analogue of Theorem 1 for non logarithmic complexes. However, the assertion becomes more complicated, and we get noncanonical and canonical isomorphisms

(2.5.1)
$$\mathbf{R} f_*(\Omega_X^{\bullet}, F) \simeq \bigoplus_{i \in \mathbb{Z}, Z \subset S} (\mathrm{DR}(M_Z^{i-r}), F)[-i].$$
$$\mathrm{Gr}_i^L \mathbf{R} f_*(\Omega_X^{\bullet}, F) = \bigoplus_{Z \subset S} (\mathrm{DR}(M_Z^{i-r}), F)[-i].$$

This implies an analogue of Corollary 1. If D is a divisor with normal crossings, we have a filtered quasi-isomorphism for Z = S

$$(2.5.2) (DR_{\log}(\widetilde{M}_S^{i-r}), F) \xrightarrow{\sim} (DR(M_S^{i-r}), F),$$

where $\mathrm{DR}_{\mathrm{log}}(\widetilde{M}_S^{i-r})$ is the intersection of $\mathrm{DR}(M_S^{i-r})$ with $\mathrm{DR}_{\mathrm{log}}(\widetilde{V}_S^i)$. This seems to be related with a question of Lewis and Shuji Saito, see also [19].

- (ii) If $\dim S = 1$, we can inductively define the infinitesimal invariants in Corollary 1 by an argument similar to [24] using [26].
- (iii) Assume S is projective and D is a divisor with normal crossings. Then the Leray filtration for $X \to S \to pt$ is given by the truncation τ on the complex of filtered \mathcal{D}_S -modules $f_*(\mathcal{O}_X(*Y), F)$, and gives the Leray filtration on the cohomology of $X \setminus Y$ (induced by the truncation τ as in [8]). Indeed, the graded pieces $\mathcal{H}^j f_*(\mathcal{O}_X(*Y), F)$ of the filtration τ on S coincide with $(\tilde{V}^{j+r}(*D), F)$, and give the open direct images by $U \to S$ of the graded pieces (V^{j+r}, F) of the filtration τ on U as filtered \mathcal{D} -modules underlying mixed Hodge modules. Note that the morphism $U \to S$ is open affine so that the direct image preserves regular holonomic \mathcal{D} -modules.

3. Examples

3.1. Lefschetz pencils. Let Y be a smooth irreducible projective variety of dimension n embedded in a projective space \mathcal{P} over \mathbb{C} . We assume that $Y \neq \mathcal{P}$ and Y is not contained in a hyperplane of \mathcal{P} so that the hyperplane sections of Y are parametrized by the dual projective spaces \mathcal{P}^{\vee} . Let $D \subset \mathcal{P}^{\vee}$ denote the discriminant. This is the image of a projective bundle over Y (consisting of hyperplanes tangent to Y), and hence D is irreducible. At a smooth point of D, the corresponding hyperplane section of Y has only one ordinary double point. We assume that the associated vanishing cycle is not zero in the cohomology of general hyperplane section X. This is equivalent to the non surjectivity of $H^{n-1}(Y) \to H^{n-1}(X)$.

A Lefschetz pencil of Y is a line \mathbb{P}^1 in \mathcal{P} intersecting the discriminant D at smooth points of D (corresponding to hyperplane sections having only one ordinary double point). We have a projective morphism $\pi: \widetilde{Y} \to \mathbb{P}^1$ such that $\widetilde{Y}_t := \pi^{-1}(t)$ is the hyperplane section corresponding $t \in \mathbb{P}^1 \subset \mathcal{P}$ and \widetilde{Y} is the blow-up of Y along a smooth closed subvariety Z of codimension 2 which is the intersection of \widetilde{Y}_t for any (or two of) $t \in \mathbb{P}^1$.

A Lefschetz pencil of hypersurface sections of degree d is defined by replacing the embedding of Y using $\mathcal{O}_Y(d)$ so that a hyperplane section corresponds to a hypersurface section of degree d. Here $\mathcal{O}_Y(d)$ for an integer d denote the invertible sheaf induced by that on \mathcal{P} as usual.

3.2. Hypersurfaces containing a subvariety. Let Y, \mathcal{P} be as in (3.2). Let E be a closed subvariety (which is not necessarily irreducible nor reduced). Let

$$E_{\{i\}} = \{x \in E : \dim T_x E = i\}.$$

Let \mathcal{I}_E be the ideal sheaf of E in Y. Let δ be a positive integer such that $\mathcal{I}_E(\delta)$ is generated by global sections. By [18], [20] (or [21]) we have the following

(3.2.1) If dim $Y > \max\{\dim E_{\{i\}} + i\}$ and $d \ge \delta$, then there is a smooth hypersurface section of degree d containing E.

We have furthermore

(3.2.2) If dim $Y > \max\{\dim E_{\{i\}} + i\} + 1$ and $d \ge \delta + 1$, then there is a Lefschetz pencil of hypersurface sections of degree d containing E.

Indeed, we have a pencil such that \widetilde{Y}_t has at most isolated singularities, because \widetilde{Y}_t is smooth near the center Z which is the intersection of generic two hypersurfaces sections containing E, and hence is smooth, see [18], [20] (or [21]). Note that a local equation of \widetilde{Y}_t near Z is given by f - tg if t is identified with an appropriate affine coordinate of \mathbb{P}^1 where f, g are global sections of $\mathcal{I}_E(d)$ corresponding to smooth hypersurface sections.

To get only ordinary double points, note first that the parameter space of the hypersurfaces containing E is a linear subspace of \mathcal{P}^{\vee} . So it is enough to show that this linear subspace contains a point of the discriminant D corresponding to an ordinary double point. Thus we have to show that an isolated singularity

can be deformed to ordinary double points by replacing the corresponding section $h \in \Gamma(Y, \mathcal{I}_E(d))$ with $h + \sum_i t_i g_i$ where $g_i \in \Gamma(Y, \mathcal{I}_E(d))$ and the $t_i \in \mathbb{C}$ are general with sufficiently small absolute values. Since $d \geq \delta + 1$, we see that $\Gamma(Y, \mathcal{I}_E(d))$ generates the 1-jets at each point of the complement of E. So the assertion follows from the fact that for a function with an isolated singularity f, the singularities of $\{f + \sum_i t_i x_i = 0\}$ are ordinary double points if t_1, \ldots, t_n are general, where t_1, \ldots, t_n are local coordinates. (Note that t_n has an ordinary double point if and only if the morphism defined by $(\partial f/\partial x_1, \ldots, \partial f/\partial x_n)$ is locally biholomorphic at this point.)

3.3. Construction. For Y, \mathcal{P} be as in (3.1), let $i_{Y,\mathcal{P}}: Y \to \mathcal{P}$ denote the inclusion. Assume

(3.3.1)
$$i_{Y,\mathcal{P}}^*: H^j(\mathcal{P}) \to H^j(Y)$$
 is surjective for any $j \neq \dim Y$,

where cohomology has coefficients in any field of characteristic zero. This condition is satisfied if Y is a complete intersection.

Let E_1 , E_2 be m-dimensional irreducible closed subvarieties of Y such that

$$E_1 \cap E_2 = \emptyset$$
, $\deg E_1 = \deg E_2$.

Here dim Y = n = 2m + s + 1 with $m \ge 0$, $s \ge 1$. Let $E = E_1 \cup E_2$. With the notation of (3.2), assume

(3.3.2)
$$d > \delta, \quad \dim Y > \max\{\dim E_{\{i\}} + i\} + s,$$

(3.3.3)
$$i_{X^{(j)},Y}^*: H^{n-j}(Y) \to H^{n-j}(X^{(j)})$$
 is not surjective for $j \leq s$,

where $X^{(j)}$ is a general complete intersection of multi degree (d, \ldots, d) and of codimension j in Y. (This is equivalent to the condition that the vanishing cycles for a hypersurface $X^{(j)}$ of $X^{(j-1)}$ are nonzero.)

Let X be a general hypersurface of degree d in Y containing E, see (3.2.1). Let L denote the intersection of X with a general linear subspace of codimension m+s in the projective space. Then $[E_a]$ (a=1,2) and $c[L\cap X]$ have the same cohomology class in $H^{2m+2s}(X)$ for some $c \in \mathbb{Q}$, because dim $H^{2m+2s}(X) = 1$ by the weak and hard Lefschetz theorems together with (3.3.1). Let

$$\xi_a = [E_a] - c[L \cap X] \in CH^{m+s}(X)_{\mathbb{Q}} \ (a = 1, 2).$$

These are homologous to zero. It may be expected that one of them is non torsion, generalizing an assertion in [24]. More precisely, let S be a smooth affine rational variety defined over a finitely generated subfield k of \mathbb{C} and parametrizing the smooth hypersurfaces of degree d containing E as above so that there is the universal family $\mathcal{X} \to S$ defined over k (see [2], [28]). Assume X corresponds to a geometric generic point of S with respect to k, i.e. X is the geometric generic fiber for some embedding $k(S) \to \mathbb{C}$. Let

$$\xi_{a,\mathcal{X}} = [E_a \times_k S] - c[L]_{\mathcal{X}} \in \mathrm{CH}^{m+s}(\mathcal{X})_{\mathbb{Q}},$$

where $[L]_{\mathcal{X}}$ is the pull-back of [L] by $\mathcal{X} \to Y$. Since the local system $\{H^{2m+2s-j}(\mathcal{X}_s)\}$ on S is constant for j < s and S is smooth affine rational, we see that $\delta_S^j(\xi_{a,\mathcal{X}}) = 0$ for j < s. Then it may be expected that $\delta_S^s(\xi_{a,\mathcal{X}}) \neq 0$ for one of a, where S can be

replaced by any non empty open subvariety. We can show this for s = 1 as follows. (For s > 1, it may be necessary to assume further conditions on d, etc.)

3.4. Case s=1. Consider a Lefschetz pencil $\pi: \widetilde{Y} \to \mathbb{P}^1$ such that $\widetilde{Y}_t := \pi^{-1}(t)$ for $t \in \mathbb{P}^1$ is a hypersurface of degree d in Y containing E. Here \widetilde{Y} is the blow-up of Y along a smooth closed subvariety Z, and Z is the intersection of \widetilde{Y}_t for any $t \in \mathbb{P}^1$. Note that \widetilde{Y}_t has an ordinary double point for $t \in \Lambda \subset \mathbb{P}^1$, where Λ denotes the discriminant, see (3.2.2).

Since Z has codimension 2 in Y, we have the isomorphism

$$(3.4.1) Hn(\widetilde{Y}) = Hn(Y) \oplus Hn-2(Z),$$

so that the cycle class of $[E_a \times \mathbb{P}^1] - c[L]_{\widetilde{Y}} \in \operatorname{CH}^{m+1}(\widetilde{Y})_{\mathbb{Q}}$ in $H^n(\widetilde{Y})$ is identified with the difference of the cycle class $cl_Z(E_a) \in H^{n-2}(Z)$ and the cycle class of L in $H^n(Y)$. Indeed, the injection $H^{n-2}(Z) \to H^n(\widetilde{Y})$ in the above direct sum decomposition is defined by using the projection $Z \times \mathbb{P}^1 \to Z$ and the closed embedding $Z \times \mathbb{P}^1 \to \widetilde{Y}$, and the injection $H^n(Y) \to H^n(\widetilde{Y})$ is the pull-back by $\widetilde{Y} \to Y$, see [17].

By assumption, one of the $cl_Z(E_a)$ is not contained in the non primitive part, i.e. not a multiple of the cohomology class of the intersection of general hyperplane sections. Indeed, if both are contained in the non primitive part, then $cl_Z(E_1) = cl_Z(E_2)$ and this implies the vanishing of the self intersection number $E_a \cdot E_a$ in Z.

We will show that the cycle class of $[E_a \times \mathbb{P}^1] - c[L]_{\widetilde{Y}}$ does not vanish in the cohomology of $\pi^{-1}(U)$ for any non empty open subvariety of \mathbb{P}^1 , in other words, it does not belong to the image of $\bigoplus_{t \in \Lambda'} H^n_{\widetilde{Y}_t}(\widetilde{Y})$ where Λ' is any finite subset of \mathbb{P}^1 containing Λ . (Note that the condition for the Lefschetz pencil is generic, and for any proper closed subvariety of the parameter space, there is a Lefschetz pencil whose corresponding line is not contained in this subvariety.)

Thus the assertion is reduced to that $\dim H^n_{\widetilde{Y}_t}(\widetilde{Y})$ is independent of $t \in \mathbb{P}^1$ because this implies that the image of $H^n_{\widetilde{Y}_t}(\widetilde{Y}) \to H^n(\widetilde{Y})$ is independent of t. (Note that the Gysin morphism $H^{n-2}(\widetilde{Y}_t) \to H^n(\widetilde{Y})$ for a general t can be identified with the direct sum of the Gysin morphism $H^{n-2}(\widetilde{Y}_t) \to H^n(Y)$ and the restriction morphism $H^{n-2}(\widetilde{Y}_t) \to H^{n-2}(Z)$ up to a sign, and the image of the last morphism is the non primitive part by the weak Lefschetz theorem.) By duality, this is equivalent to that $R^n\pi_*\mathbb{Q}_{\widetilde{Y}}$ is a local system on \mathbb{P}^1 . Then it follows from the assumption that the vanishing cycles are nonzero, see (3.3.3).

References

- [1] Asakura, M., Motives and algebraic de Rham cohomology, in The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), CRM Proc. Lecture Notes, 24, Amer. Math. Soc., Providence, RI, 2000, pp. 133–154.
- [2] Beilinson, A., Bernstein, J. and Deligne, P., Faisceaux pervers, Astérisque, vol. 100, Soc. Math. France, Paris, 1982.
- [3] Bloch, S., Algebraic cycles and higher K-theory, Advances in Math., 61 (1986), 267–304.

- [4] Bloch, S., Algebraic cycles and the Beilinson conjectures, Contemporary Math. 58 (1) (1986), 65–79.
- [5] Collino, A., Griffiths' infinitesimal invariant and higher K-theory on hyperelliptic Jacobians, J. Alg. Geom. 6 (1997), 393–415.
- [6] Deligne, P., Théorème de Lefschetz et critères de dégénérescence de suites spectrales, Inst. Hautes Etudes Sci. Publ. Math. 35 (1968), 259–278.
- [7] Deligne, P., Equation différentielle à points singuliers réguliers, Lect. Notes in Math. vol. 163, Springer, Berlin, 1970.
- [8] Deligne, P., Théorie de Hodge I, Actes Congrès Intern. Math., 1970, vol. 1, 425-430; II, Publ. Math. IHES, 40 (1971), 5-57; III, ibid., 44 (1974), 5-77.
- [9] Deligne, P., Décompositions dans la catégorie dérivée, in Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., 55, Part 1, Amer. Math. Soc., Providence, RI, 1994, pp. 115–128.
- [10] Deninger, C. and Scholl, A., The Beilinson conjectures, in Proceedings Cambridge Math. Soc. (eds. Coats and Taylor) 153 (1992), 173–209.
- [11] El Zein, F., Complexe dualisant et applications à la classe fondamentale d'un cycle, Bull. Soc. Math. France Mém. No. 58 (1978)
- [12] Esnault, H. and Viehweg, E., Deligne-Beilinson cohomology, in Beilinson's conjectures on Special Values of *L*-functions, Academic Press, Boston, 1988, pp. 43–92.
- [13] Green, M.L., Griffiths' infinitesimal invariant and the Abel-Jacobi map, J. Differential Geom. 29 (1989), 545–555.
- [14] Griffiths, P.A., Infinitesimal variations of Hodge structure, III, Determinantal varieties and the infinitesimal invariant of normal functions, Compositio Math. 50 (1983), 267–324.
- [15] Jannsen, U., Deligne homology, Hodge-*D*-conjecture, and motives, in Beilinson's conjectures on Special Values of *L*-functions, Academic Press, Boston, 1988, pp. 305–372.
- [16] Jannsen, U., Mixed motives and algebraic K-theory, Lect. Notes in Math., vol. 1400, Springer, Berlin, 1990.
- [17] Katz, N., Etude cohomologique des pinceaux de Lefschetz, in Lect. Notes in Math., vol. 340, Springer Berlin, 1973, pp. 254–327.
- [18] Kleiman, S. and Altman, A., Bertini theorems for hypersurface sections containing a subscheme, Comm. Algebra 7 (1979), 775–790.
- [19] Lewis, J.D. and Saito, S., preprint.
- [20] Otwinowska, A, Monodromie d'une famille d'hypersurfaces, preprint (math.AG/0403151).
- [21] Otwinowska, A. and Saito, M., Monodromy of a family of hypersurfaces containing a given subvariety, preprint (math.AG/0404469).
- [22] Saito, M., Modules de Hodge polarisables, Publ. RIMS, Kyoto Univ. 24 (1988), 849–995.
- [23] Saito, M., Mixed Hodge Modules, Publ. RIMS Kyoto Univ. 26 (1990), 221–333.
- [24] Saito, S., Higher normal functions and Griffiths groups, J. Algebraic Geom. 11 (2002), 161–201.
- [25] Schmid, W., Variation of Hodge structure: the singularities of the period mapping, Inv. Math. 22 (1973), 211–319.
- [26] Steenbrink, J.H.M., Limits of Hodge structures, Inv. Math. 31 (1975/76), no. 3, 229–257.
- [27] Voisin, C., Variations de structure de Hodge et zéro-cycles sur les surfaces générales, Math. Ann. 299 (1994), 77–103.
- [28] Voisin, C., Transcendental methods in the study of algebraic cycles, in Lect. Notes in Math. vol. 1594, pp. 153–222.

RIMS KYOTO UNIVERSITY, KYOTO 606-8502 JAPAN *E-mail address*: msaito@kurims.kyoto-u.ac.jp