# ON THE CRITICAL POINTS OF THE $E_k$ FUNCTIONALS IN KÄHLER GEOMETRY

VALENTINO TOSATTI

ABSTRACT. We prove that a Kähler metric in the anticanonical class which is a critical point of the functional  $E_k$  and has nonnegative Ricci curvature, is necessarily Kähler-Einstein. This partially answers a question of X.X. Chen.

# 1. INTRODUCTION

One of the major problems in Kähler geometry is the study of extremal metrics, especially Kähler-Einstein metrics or constant scalar curvature Kähler metrics (cscK). Let  $(M, \omega)$  be a compact Kähler manifold of complex dimension n, and assume the first Chern class of M to be negative or zero. Then by the celebrated work of Yau [13] (see also Aubin [1]) we know that M admits a Kähler-Einstein metric. The remaining case  $c_1(M) > 0$  is more difficult and still largely open. There is a conjecture of Yau [14] that the existence of a Kähler-Einstein metric should be equivalent to the stability of the manifold in some algebro-geometric sense. The behaviour of the Mabuchi energy [6] plays a central role in this study [12]. X.X. Chen and G. Tian in [4] have introduced a family of functionals  $E_k$   $(0 \le k \le n)$ which generalize the Mabuchi energy  $\nu_{\omega} = E_0$ , and used  $E_1$  in their study of the Kähler-Ricci flow. They can also be defined in terms of the Deligne pairing [9], and they should possibly be related to stability. Bando and Mabuchi showed [2] that Kähler-Einstein metrics realize the minimum of the Mabuchi energy, and recently Song and Weinkove [10] showed that Kähler-Einstein metrics realize the minimum of the  $E_k$ , on the space of metrics with nonnegative Ricci curvature in the anticanonical class for  $k \ge 2$ , and on the whole anticanonical class for k = 1. Pali [8] also investigated the question of the lower boundedness of  $E_1$ .

For the moment we drop the assumption  $c_1(M) > 0$ . Let  $P(M, \omega)$  be the set of Kähler potentials for  $\omega$ , i.e. the set of all smooth real functions  $\phi$  such that

$$\omega_\phi := \omega + \sqrt{-1}\partial\overline{\partial}\phi$$

is positive definite. For  $\phi \in P(M, \omega)$  let  $\phi_t$  be a smooth path in  $P(M, \omega)$  with  $\phi_0 = 0, \phi_1 = \phi$ , and define

$$E_{k,\omega}(\phi) = \frac{k+1}{V} \int_0^1 \int_M (\Delta_{\phi_t} \dot{\phi}_t) \operatorname{Ric}(\omega_{\phi_t})^k \wedge \omega_{\phi_t}^{n-k} dt - \frac{n-k}{V} \int_0^1 \int_M \dot{\phi}_t \left(\operatorname{Ric}(\omega_{\phi_t})^{k+1} - \mu_k \, \omega_{\phi_t}^{k+1}\right) \wedge \omega_{\phi_t}^{n-k-1} dt,$$

2000 Mathematics Subject Classification. Primary 32Q20, 58E11.

The author is supported by a Harvard Mathematics Department grant.

where  $\dot{\phi}_t = \frac{\partial}{\partial t} \phi_t$ ,  $V = \int_M \omega^n$  is the volume of M, and

$$\mu_k = \frac{1}{V} \int_M \operatorname{Ric}(\omega)^{k+1} \wedge \omega^{n-k-1} = (2\pi)^{k+1} \frac{[c_1(M)]^{k+1} \smile [\omega]^{n-k-1}}{[\omega]^n}$$

is a constant that depends only on the cohomology classes  $[c_1(M)]$  and  $[\omega]$ . It can be checked that these functionals do not depend on the choice of the path. The functional  $E_{0,\omega}$  is equal to the Mabuchi energy  $\nu_{\omega}$ . With a simple integration by parts we can rewrite these functionals as

$$E_{k,\omega}(\phi) = \frac{1}{V} \int_0^1 \int_M \dot{\phi}_t \left[ (k+1) \binom{n}{k}^{-1} (\Delta_{\phi_t}(\sigma_k(\omega_{\phi_t})) - \sigma_{k+1}(\omega_{\phi_t})) + (n-k)\mu_k \right] \omega_{\phi_t}^n dt,$$

where  $\sigma_k(\omega)$  is given by

$$(\omega + t\operatorname{Ric}(\omega))^n = \left(\sum_{k=0}^n \sigma_k(\omega)t^k\right)\omega^n.$$

If at a point p we pick normal coordinates in which the Ricci tensor is diagonal with entries  $\lambda_1, \ldots, \lambda_n$  then  $\sigma_k(\omega)$  at p is the k-th elementary symmetric function in the  $\lambda_i$ ,

$$\sigma_k(\omega) = \sum_{1 \le i_1 < \dots < i_k \le n} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_k}.$$

For example we have  $\sigma_0(\omega) = 1$ ,  $\sigma_1(\omega) = R$  the scalar curvature of  $\omega$ ,  $\sigma_2(\omega) = \frac{1}{2}(R^2 - |\text{Ric}|^2)$ . From this we see that the critical points of  $E_{k,\omega}$  are the functions  $\phi$  such that  $\omega_{\phi}$  satisfies

(1) 
$$\sigma_{k+1}(\omega_{\phi}) - \triangle_{\phi}\sigma_{k}(\omega_{\phi}) = \binom{n}{k+1}\mu_{k}$$

For k = 0 we see that the critical metrics are precisely cscK metrics. The critical equation for k = 1 is

(2) 
$$R^2 - |\operatorname{Ric}|^2 - 2\triangle R = n(n-1)\mu_1$$

From now on assume that  $c_1(M) > 0$ , that  $[\omega] = 2\pi [c_1(M)]$  so that  $\mu_k = 1$  for all k. Under these hypotheses Chen and Tian [4] have shown that critical metrics for  $E_n$  with positive Ricci curvature must be necessarily Kähler-Einstein. In a recent paper [3] X.X. Chen asked the following:

**Question** (X.X. Chen). Is a critical metric for  $E_1$  in the anticanonical class necessarily Kähler-Einstein?

Our result is a partial answer to this question, for all k, generalizing Chen and Tian:

**Theorem 1.** Fix  $0 \le k \le n$  and assume  $\omega$  is a Kähler metric in the anticanonical class which is critical for  $E_k$  and has nonnegative Ricci curvature. Then  $\omega$  is Kähler-Einstein. If k = 1 then the same holds if  $\omega$  satisfies only R > -n.

*Remark.* G. Maschler [7] has proved that a critical point of  $E_n$  in the anticanonical class is always Kähler-Einstein.

#### KÄHLER GEOMETRY

## 2. The proof

In case k = 0 the critical metrics are cscK and it is well known that in the anticanonical class they are necessarily Kähler-Einstein, even if the Ricci curvature is not positive (see for example [11, Prop. 2.12]). Next we assume that 0 < k < n, that  $\operatorname{Ric}(\omega) > 0$  and we use the minimum principle. Let  $p \in M$  be a point where  $\sigma_k(\omega)$  achieves its minimum, and pick normal coordinates around p in which the Ricci tensor is diagonal with *positive* entries  $\lambda_1, \ldots, \lambda_n$ . Notice that all the  $\sigma_k(\omega)$  are positive. Then using the critical equation (1) we have that

$$\sigma_{k+1}(\omega)(p) \geq \binom{n}{k+1}.$$

Define the normalized symmetric functions as

$$\Sigma_k(\omega) := \frac{\sigma_k(\omega)}{\binom{n}{k}},$$

so that at p we have  $\Sigma_{k+1}(\omega) \ge 1$ . Since the  $\lambda_i$ 's are positive, a classical inequality (see [5, Sect. 2.22, Thm. 51]) says that for every  $0 \le k \le n-1$  we have

(3) 
$$\Sigma_k^{k+1}(\omega) \ge \Sigma_{k+1}^k(\omega),$$

and this holds everywhere on M. Hence we have  $\Sigma_k(\omega)(p) \ge 1$  and since p is the minimum point of  $\Sigma_k(\omega)$ , we have  $\Sigma_k(\omega) \ge 1$  everywhere. Also using (3) repeatedly we have that

$$\Sigma_1(\omega) \ge \Sigma_2^{\frac{1}{2}}(\omega) \ge \dots \ge \Sigma_k^{\frac{1}{k}}(\omega) \ge 1$$

everywhere on M. Now by hypothesis  $\omega \in 2\pi c_1(M)$  so there exists a smooth real function f such that  $\operatorname{Ric}(\omega) = \omega + \sqrt{-1}\partial\overline{\partial}f$ , and taking the trace of this with respect to  $\omega$  we get  $R = n + \Delta f$ , which can be rewritten as  $\Sigma_1(\omega) = 1 + \frac{\Delta f}{n}$ . Hence  $\Delta f \geq 0$  from which it follows easily that f is constant, and so  $\operatorname{Ric}(\omega) = \omega$  and  $\omega$  is Kähler-Einstein.

If now we only assume that  $\operatorname{Ric}(\omega) \geq 0$ , we have that  $\lambda_i \geq 0$ . Using the minimum principle as before, we see that  $\Sigma_{k+1}(\omega)(p) \geq 1$ , so it is not zero. From this it follows that  $\Sigma_j(\omega)(p) > 0$  for  $0 \leq j \leq k$ , and it is very easy to check that in this case the inequality (3) still holds and the reasoning proceeds as before.

If k = n then the critical equation becomes  $\Delta \sigma_n(\omega) = 0$ . Hence  $\sigma_n(\omega) = 1 = \Sigma_n(\omega)$ , and as before  $\Sigma_1(\omega) \ge \Sigma_n^{\frac{1}{n}}(\omega) = 1$  and so  $\omega$  is Kähler-Einstein. G. Maschler called such metrics *central Kähler metrics* and he proved [7] that the hypothesis Ric  $\ge 0$  can be removed, using Demailly's holomorphic Morse inequalities.

If k = 1 then the critical equation is (2) with  $\mu_1 = 1$ . In this case the inequality (3) holds for arbitrary  $\lambda_i$ , and takes the form  $|\text{Ric}|^2 \geq \frac{R^2}{n}$  (which follows from  $|\text{Ric} - \frac{R}{n}g|^2 \geq 0$ ). With the same argument as above we get at the minimum p of R,  $R^2(p) \geq n^2$ . Since we are assuming R > -n, we must have  $R(p) \geq n$ , so  $R \geq n$  everywhere and  $\omega$  is Kähler-Einstein.

## 3. Further remarks

**Question.** Is Theorem 1 still true if we do not assume the Ricci curvature to be nonnegative (at least for k = 1)?

Notice that this holds for k = 0, n.

#### VALENTINO TOSATTI

*Remark.* Further evidence for the validity of Chen's question is provided by this recent result of Song and Weinkove [10]: assuming M has no nonzero holomorphic vector fields, the functional  $E_1$  is proper in the sense of Tian [12] if and only if there exists a Kähler-Einstein metric. This leads one to expect that Kähler-Einstein metrics and critical points of  $E_1$  coincide.

Acknowledgements. The author thanks Jian Song and Ben Weinkove for many useful discussions, and his advisor Professor S.-T. Yau for his constant support.

# References

- T. Aubin Equations du type Monge-Ampère sur les variétés Kähleriennes compactes, Bull. Sci. Math. 102 (1976), 119-121.
- S. Bando, T. Mabuchi Uniqueness of Einstein Kähler metrics modulo connected group actions, Adv. Stud. Pure Math. 10 (1987), 11-40.
- [3] X.X. Chen On the lower bound of energy functional E<sub>1</sub> (I) a stability theorem on the Kähler-Ricci flow, J. Geom. Anal. 16 (2006) no. 1, 23-38.
- [4] X.X. Chen, G. Tian Ricci flow on Kähler-Einstein surfaces, Invent. Math. 147 (2002), no. 3, 487-544.
- [5] G.H. Hardy, J.E. Littlewood, G. Pólya Inequalities, Cambridge University Press, 1952.
- [6] T. Mabuchi K-energy maps integrating Futaki invariants, Tôhoku Math. J. 38 (1986), 575-593.
- [7] G. Maschler Central Kähler metrics, Trans. Amer. Math. Soc. 355 (2003), no.6, 2161-2182.
- [8] N. Pali A consequence of a lower bound of the K-energy, Int. Math. Res. Not. 2005, no. 50, 3081-3090.
- [9] D.H. Phong, J. Sturm The Futaki invariant and the Mabuchi energy of a complete intersection, Comm. Anal. Geom. 12 (2004), no. 1-2, 323-343.
- [10] J. Song, B. Weinkove Energy functionals and canonical Kähler metrics, to appear in Duke Math. J.
- [11] G. Tian Canonical metrics in Kähler geometry, Birkhähuser, 2000.
- [12] G. Tian Kähler-Einstein metrics with positive scalar curvature, Invent. Math. 137 (1997), 1-37.
- [13] S.-T. Yau On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I, Comm. Pure Appl. Math. 31 (1978), 339-411.
- [14] S.-T. Yau Open problems in geometry, Proc. Sympos. Pure Math. 54 (1993), 1-28 (problem 65).

Department of Mathematics, Harvard University, 1 Oxford St, Cambridge MA 02138  $E\text{-}mail\ address:\ \texttt{tosattiQmath.harvard.edu}$ 

4