arXiv:math/0506136v2 [math.GT] 12 Jan 2007

CONNECTED COMPONENTS OF THE STRATA OF THE MODULI SPACES OF QUADRATIC DIFFERENTIALS

ERWAN LANNEAU

ABSTRACT. In two fundamental classical papers, Masur [Ma1] and Veech [Ve1] have independently proved that the Teichmüller geodesic flow acts ergodically on each connected component of each stratum of the moduli space of quadratic differentials. It is therefore interesting to have a classification of the ergodic components. Veech has proved that these strata are not necessarily connected. In a recent work [KoZo], Kontsevich and Zorich have completely classified the components in the particular case where the quadratic differentials are given by the global square of Abelian differentials.

Here we are interested in the complementary case. In a previous paper [La1], we have described some particular component, namely the *hyperelliptic* connected components, and showed that some strata are non-connected. In this paper, we give the general classification theorem: up to four exceptional cases in low genera, the strata of meromorphic quadratic differentials are either connected, or have exactly two connected components. In this last case, one component is hyperelliptic, the other not. This result was announced in the paper [La1].

Our proof is based on a new approach of the so-called Jenkins-Strebel differential. We will present and use the notion of *generalized permutations*.

RÉSUMÉ. Dans des travaux maintenant classiques, Masur [Ma1] et Veech [Ve1] ont démontré indépendamment que le flot géodésique de Teichüller est ergodique sur chaque composante connexe de chaque strate de l'espace des modules des différentielles quadratiques. Il devient dès lors intéressant d'avoir une description de ces composantes ergodiques. Veech a montré que ces strates ne sont pas nécessairement connexes. Dans un article récent Kontsevich et Zorich [KoZo] donnent une description complète des composantes dans le cas particulier où les différentielles quadratiques sont données par le carré de différentielles abéliennes.

Dans cet article nous considérons le cas complémentaire. Dans un précédent article [La1] nous montrions que les strates ne sont pas forcément connexes. Nous donnions une série de strates non-connexes possédant des composantes connexes *hyperelliptiques*. Dans cet article, nous démontrons le théorème général annoncé dans [La1] : excepté quatre cas particuliers en petits genres, les strates de l'espace des modules des différentielles quadratiques ont au plus deux composantes connexes. Les cas de non-connexité étant décrits exactement par [La1] : une composante est hyperelliptique, l'autre non.

Notre preuve repose principalement sur une nouvelle approche des différentielles quadratiques de type Jenkins-Strebel à savoir la notion de *permutations généralisées*.

1. INTRODUCTION

The moduli space of compact connected Riemann surfaces S of genus g endowed with an integrable meromorphic quadratic differential q is a disjoint union $\mathcal{H}_q \sqcup \mathcal{Q}_q$, where the

Date: October 15, 2006.

²⁰⁰⁰ Mathematics Subject Classification. Primary: 32G15. Secondary: 30F30, 57R30, 37D40.

Key words and phrases. Quadratic differentials, moduli space, measured foliations, Teichümuller geodesic flow.

isomorphism class of (S, q) belongs to \mathcal{H}_g if and only if q is the (global) square of a holomorphic Abelian differential. It can be identified with the cotangent bundle of the moduli space \mathcal{M}_g of compact connected smooth complex curves (see for instance [HuMa]). It carries a natural flow, called the Teichmüller geodesic flow, see for instance [Ma1, Ve1]. It has a natural stratification, whose strata are denoted by $\mathcal{H}(k_1, \dots, k_n) = \mathcal{H}_g(k_1, \dots, k_n)$ contained in \mathcal{H}_g and $\mathcal{Q}(k_1, \dots, k_n) = \mathcal{Q}_g(k_1, \dots, k_n)$ contained in \mathcal{Q}_g , where $k_1, \dots, k - n$ is the (unordered) list of multiplicities of the zeroes and poles of the quadratic differentials. It is well known that the flow preserves this stratification and that each stratum carries a complex algebraic orbifold structure of complex dimension $2g + n - \varepsilon$ (here $\varepsilon = 1$ or -1 depending respectively of the strata of Abelian differential or quadratic differentials). Masur and Smillie [MaSm] proved that all of these strata (corresponding to the multiplicities satisfying the Gauss-Bonnet condition), except four particular cases in low genera, are non-empty.

The aim of this paper is motivated by a fundamental theorem, independently proved by Masur and Veech [Ma1], [Ve1], which asserts that the Teichmüller geodesic flow acts ergodically on each connected component of each stratum (with respect to a finite measure equivalent to the Lebesgue measure).

Kontsevich and Zorich [KoZo] have recently described the set of connected components for the strata in \mathcal{H}_g . In [La1], using a construction developed in [KoZo] (hyperelliptic components), we showed that some strata in \mathcal{Q}_g are non-connected. More precisely, we presented three series of one discrete-parameter strata which are non connected; those strata have a connected component consisting of hyperelliptic curves equipped with an "hyperelliptic differential". This component is called an *hyperelliptic component*.

In this paper, we describe the set of connected components of any stratum of Q_g . The general case stabilizes at genus 5 and corresponds to Theorem 1.1. Theorem 1.2 gives the remaining cases.

Theorem 1.1. Let us fix $g \ge 5$. Each stratum of the moduli space Q_g having an hyperelliptic connected component has exactly two connected components: one is hyperelliptic — the other not; the detailed list is given in [La1].

Any other stratum of the moduli space \mathcal{Q}_g of quadratic differentials is connected.

In small genera, there are some exceptional cases coming from the geometry of genus one and genus two surfaces (respectively elliptic and hyperelliptic curves). There are also 4 mysterious cases which appear.

Theorem 1.2. Let us fix $g \leq 4$. The components of the strata of the moduli space Q_g fall in the following description:

- In genera 0 and 1, any stratum is connected.
- In genus 2, there are two non-connected strata. For these two, one component is hyperelliptic, the other not. Any other stratum of Q_2 is connected.
- In genera 3 and 4, each stratum with an hyperelliptic connected component has exactly two connected components: one is hyperelliptic, the other not.
- There are 4 sporadic strata in genus 3 and 4 which are non-connected and which do not possess an hyperelliptic component.
- Any other stratum of Q_3 and Q_4 is connected.

1.1. Precise formulation of the statements. In order to establish notations and to give a precise statement, we review basic notions concerning moduli spaces, Abelian differentials and quadratic differentials. There is an abundant literature on this subject; for more details and proofs see for instance [DoHu], [EsMaZo], [FaLaPo], [HuMa] [Ko], [KoZo], [Ma1], [St],[Th],[Ve1],[Ve2], [Ve3],... For a nice survey see [MaTa] or [Zo2].

1.1.1. Background. For $g \geq 1$, we define the moduli space of Abelian differentials \mathcal{H}_g as the moduli space of pairs (\mathcal{S}, ω) where \mathcal{S} is a genus g (closed connected) Riemann surface and $\omega \in \Omega(\mathcal{S})$ a non-zero holomorphic 1-form defined on \mathcal{S} . The term moduli spaces means that we identify the points (\mathcal{S}, ω) and (\mathcal{S}', ω') if there exists an analytic isomorphism $f : \mathcal{S} \to \mathcal{S}'$ such that $f^*\omega' = \omega$.

For $g \geq 0$, we also define the moduli space of quadratic differentials \mathcal{Q}_g which are not the global square of Abelian differentials as the moduli space of pairs (\mathcal{S}, q) where \mathcal{S} is a genus g Riemann surface and q a non-zero meromorphic quadratic differential defined on \mathcal{S} such that q is not the global square of any Abelian differential. In addition, we assume that q has at most simple poles, if any. This last condition guaranties that the area of \mathcal{S} in terms of the metric determined by q is finite:

$$\int_{\mathcal{S}} |q| < \infty.$$

We will denote by $\mathcal{H}(k_1, \ldots, k_n)$ the subset of \mathcal{H}_g consisting of (classes of) pairs (\mathcal{S}, ω) such that ω possesses exactly n zeroes on \mathcal{S} with multiplicities (k_1, \ldots, k_n) . We will also denote by $\mathcal{Q}(k_1, \ldots, k_n)$ the subset of \mathcal{Q}_g consisting of pairs (\mathcal{S}, q) such that q possesses exactly n singularities on \mathcal{S} with multiplicities (k_1, \ldots, k_n) , $k_i \geq -1$.

Note that the Gauss-Bonnet formula implies that the sum of the multiplicities $\sum k_i$ equals 2g-2 in the case of $\mathcal{H}(k_1,\ldots,k_n)$ and 4g-4 in the case of $\mathcal{Q}(k_1,\ldots,k_n)$. In Section 2.1.1, we will present Thurston's approach to these surfaces via the theory of measured foliations.

From these definitions, it is a well known part of the Teichmüller theory that these spaces are (Hausdorff) complex analytic, and in fact algebraic, spaces (see [DoHu] for a nice description of the stratum Q(1, ..., 1); see also [HuMa], [Ko], [Ve2]). Basically, one can see that as follows. We first concentrate on the strata of the moduli spaces \mathcal{H}_q .

Let (S, ω^2) be a representative of an element in $\mathcal{H}(k_1, \ldots, k_n)$, S its underlying topological surface, and P_1, \ldots, P_n its singular points. Let us denote by hol = $\operatorname{hol}_{(S,\omega)}$ the group morphism $H_1(S, \{P_1, \ldots, P_n\}, \mathbb{Z}) \to \mathbb{C}$ defined by $\operatorname{hol}([\gamma]) = \int_{\gamma} \omega$ for every 1-cycle γ in S relative to $\{P_1, \ldots, P_n\}$. Fix a basis $(\gamma_1, \ldots, \gamma_{2g+n-1})$ of the free abelian group $H_1(S, \{P_1, \ldots, P_n\}, \mathbb{Z})$. Any other element of $\mathcal{H}(k_1, \ldots, k_n)$ will be represented by an element having the same underlying surface and the same singular points. With these notations, the map

$$\Phi = \begin{pmatrix} \mathcal{H}(k_1, k_2, \dots, k_n) & \longrightarrow & H^1(S, \{P_1, \dots, P_n\}, \mathbb{C}) \\ \mathcal{S}' & \longmapsto & (\gamma_1, \dots, \gamma_{2g+n-1}) \mapsto (hol_{\mathcal{S}'}(\gamma_1), \dots, hol_{\mathcal{S}'}(\gamma_{2g+n-1})) \end{pmatrix}$$

is named the *period map* and is a local homeomorphism in a neighbourhood of (\mathcal{S}, ω^2) . Therefore we get a locally one-to-one correspondence between the corresponding stratum of \mathcal{H}_g and an open domain in the vector space $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C}) \simeq \mathbb{C}^{2g+n-1}$. The change of coordinates are affine maps outside the singularities of $\mathcal{H}(k_1, k_2, \ldots, k_n)$ and produces after a study of the singularities a differentiable orbifold structure on the strata of \mathcal{H}_g .

Let us now consider the case of a stratum of the moduli space \mathcal{Q}_g . For every $(\mathcal{S}, q) \in \mathcal{Q}_g$, consider the canonical double cover $\pi : \hat{\mathcal{S}} \to \mathcal{S}$ such that $\pi^* q = \omega^2$ for some holomorphic Abelian differential ω on $\hat{\mathcal{S}}$ (see for instance [La2]). As above, we consider the period map between a neighborhood of the point $(\hat{\mathcal{S}}, \omega)$ and an open subset of $H^1(\hat{\mathcal{S}}, \{\hat{P}_1, \ldots, \hat{P}_n\}; \mathbb{C})$. The covering involution $\tau : \hat{\mathcal{S}} \to \hat{\mathcal{S}}$ induces an involutive linear map on this cohomology vector

space. Therefore, this vector space decomposes into two eigenspaces for τ^* , say E_{-1} and E_{+1} , with eigenvectors -1 and +1. Abelian differentials in E_{-1} are precisely those which arise from quadratic differentials on S by pull-back by π . Hence we obtain a one-to-one correspondence between a neighborhood of any point in the corresponding strata of Q_g and an open domain of $E_{-1} \simeq \mathbb{C}^{2g+n-2}$.

Next we recall the construction of a measure μ on each stratum. For that, the tangent space to \mathcal{H}_q (respectively \mathcal{Q}_q) at each point contains a lattice:

$$H^{1}(\mathcal{S}, \{P_{1}, \ldots, P_{n}\}; \mathbb{Z}) \oplus i \cdot H^{1}(\mathcal{S}, \{P_{1}, \ldots, P_{n}\}; \mathbb{Z}) \subset H^{1}(\mathcal{S}, \{P_{1}, \ldots, P_{n}\}; \mathbb{C}).$$

We define the function $A : \mathcal{H}_g \to \mathbb{R}_+$ by the formula $A(\mathcal{S}, \omega) = \frac{i}{2} \int_{\mathcal{S}} \omega \wedge \overline{\omega}$. This is the area of \mathcal{S} in terms of the flat metric associated to ω .

The group $\operatorname{SL}_2(\mathbb{R})$ acts by linear transformations with constant coefficients on the pair of real-valued 1-forms ($\operatorname{Re}(\omega), \operatorname{Im}(\omega)$). In the local affine coordinates, this action is the action of $\operatorname{SL}_2(\mathbb{R})$ on the coefficient field of the cohomology vector space $H^1(S, \{P_1, \ldots, P_n\}; \mathbb{C})$. From this description, it is clear that the subgroup $\operatorname{SL}_2(\mathbb{R})$ preserves the measure μ and the function A.

On the hypersurface $\mathcal{H}_g^{(1)} = A^{-1}(1)$, we define the induced measure by the formula

$$\mu^{(1)} = \frac{\mu}{dA}.$$

Recall that each stratum carries a complex algebraic orbifold structure modeled on the first relative cohomology group (see for instance [Ko]). The dimensions are respectively given by

$$\dim_{\mathbb{C}} \mathcal{H}(k_1, \dots, k_n) = 2g + n - 1 \quad \text{where} \quad k_1 + \dots + k_n = 2g - 2$$
$$\dim_{\mathbb{C}} \mathcal{Q}(k_1, \dots, k_n) = 2g + n - 2 \quad \text{where} \quad k_1 + \dots + k_n = 4g - 4.$$

The action of the 1-parameter subgroup of diagonal matrices $g_t := \text{diag}(e^{t/2}, e^{-t/2})$ presents a particular interest for our purpose. It gives a measure-preserving flow for $\mu^{(1)}$, preserving each stratum. This flow is known as the *Teichmüller geodesic flow*. Note that orbits under g_t project to Teichmüller geodesics on the moduli space of Riemann surfaces \mathcal{M}_g . The next fundamental result motives our study (see [Ma1], [Ve1]).

Theorem (Masur, Veech). The Teichmüller geodesic flow acts ergodically on each connected component of each stratum of $\mathcal{H}_g^{(1)}$ with respect to the measure $\mu^{(1)}$, which is finite and in the Lebesgue class.

A direct corollary of the finiteness of the measure $\mu^{(1)}$ on any stratum of $\mathcal{H}_g^{(1)}$ is a proof of the conjecture of Keane [Ke]: almost all intervals exchange transformations are uniquely ergodic.

1.1.2. Topology of the Moduli Space. Following the theorem of Masur and Veech, we are interested in the classification of the connected components of the strata of $\mathcal{H}_g \sqcup \mathcal{Q}_g$. Veech and Arnoux discovered, by direct calculations in terms of *Rauzy classes*, that some strata are non-connected. They have proved that $\mathcal{H}(2)$ is connected, $\mathcal{H}(4)$ have 2 connected components and $\mathcal{H}(6)$ have 3 connected components.

Recently, in the context of moduli space of Abelian differential \mathcal{H}_g , Kontsevich and Zorich (see [KoZo]) obtained the following complete description.

Theorem (Kontsevich, Zorich). Let $g \ge 4$ be any integer. The topology of any stratum of \mathcal{H}_q is given by the following list:

- The strata $\mathcal{H}(2g-2)$ and $\mathcal{H}(2k,2k)$, for any $k \geq 2$, have three connected components.
- Any other stratum $\mathcal{H}(2k_1,\ldots,2k_n)$, for any $k_i \geq 1$, has two connected components.
- The stratum $\mathcal{H}(2k-1, 2k-1)$, for any $k \geq 2$, has two connected components.

Any other stratum of Abelian differentials on a surface of genus $g \ge 4$ is non-empty and connected.

The description of connected components for strata of genera $1 \leq g \leq 3$ is similar to the previous one with some exceptions; we do not present the result here (see [KoZo]). Roughly speaking, Kontsevich and Zorich use two invariants to obtain this classification: the parity of the spin structure and the hyperellipticity.

In [La2], we prove that the first invariant extends trivially to the moduli space Q_g . However, the second produces non-trivial values. In [La1], we classify all strata for which this second one invariant produces non-trivial values. In order to present our statement, we will recall briefly this construction in the coming Section.

In this paper, we will show that this (hyperelliptic) invariant is complete in genera $g \ge 5$: it classifies precisely the components of the strata of \mathcal{Q}_g . For small genera, we obtain a similar result with 4 additional mysterious components.

Remark 1.3. As a direct corollary of our result and Kontsevich-Zorich's theorem, we draw that the ergodic components of the Teichmüller geodesic flow are given by an explicit list. In particular, if $n \ge 5$, for any $g \ge 0$, any stratum of $\mathcal{H} \sqcup \mathcal{Q}_g$ with n singularities is connected.

Remark 1.4. This paper achieves the classification of connected components of the strata of the moduli spaces $\mathcal{H} \sqcup \mathcal{Q}_g$ announced in the vast program in [Ko]. Nevertheless a more precise description of the strata is not completely understood, even for the simplest "non-trivial" case: $\mathcal{H}(2)$.

Conjecture (Kontsevich). Each connected component of the strata of the moduli space $\mathcal{H} \sqcup \mathcal{Q}_g$ is a $K(\pi, 1)$, where π is a group commensurable with some mapping class group.

1.1.3. *Hyperelliptic components*. We will need the following statement. A proof can be found in [KoZo]; see also [La1].

Proposition 1.5. Any stratum of the moduli space $Q(k_1, \ldots, k_n)$ in genus 0 is connected.

Let S_g be a (compact, connected) Riemann surface endowed with a (integrable, meromorphic) quadratic differential q_0 which is not the global square of any Abelian differential. Let (k_1, \ldots, k_n) be its singularity pattern. We do not exclude the case when some of k_i are equal to zero: by convention this means that we have some marked points. Sometimes We shall use the "exponential" notation to denote multiple singularities (simple poles), for example $Q(-1^5, 1) := Q(-1, -1, -1, -1, -1, 1)$.

Let $\pi : \tilde{S}_{\tilde{g}} \to S_g$ be a (ramified) covering such that the image of any ramification point of π is a marked point, or a zero, or a pole of the quadratic differential q_0 . Fix the *combinatorial type* of the covering π : the degree of the covering, the number of critical fibers and the ramification index of the points in every critical fiber. Consider the induced quadratic differential π^*q_0 on $\tilde{S}_{\tilde{g}}$; let $(\tilde{k}_1, \ldots, \tilde{k}_m)$ be its singularity pattern.

Deforming slightly the initial point $(\mathcal{S}_g, q_0) \in \mathcal{Q}(k_1, \ldots, k_n)$, we get a ramified covering over the deformed Riemann surface with the same combinatorial type as the covering π .

The induced quadratic differential $\pi^* q$ has the same singularity pattern $(\tilde{k}_1, \ldots, \tilde{k}_m)$ as $\pi^* q_0$. Thus we obtain a map from the stratum $\mathcal{Q}(k_1, \ldots, k_n)$ to the stratum $\mathcal{Q}(\tilde{k}_1, \ldots, \tilde{k}_m)$. We will denote denote this map by Π . In [La1], we prove that Π is an immersion.

Now we recall an example of this construction for strata having four singularities. Consider a meromorphic quadratic differential q on $\mathbb{C}P(1)$ having the singularity pattern $(2(g-k) - 3, 2k + 1, -1^{2g+2})$, where $k \ge -1$, $g \ge 1$ and $g - k \ge 2$. Consider a ramified double covering π over $\mathbb{C}P(1)$ having ramification points over the 2g + 2 poles of q, and no other ramification points. We obtain a hyperelliptic Riemann surface \tilde{S} of genus g with a quadratic differential π^*q on it. A straightforward computation shows that the induced quadratic differential π^*q has the singularity pattern (2(g-k) - 3, 2(g-k) - 3, 2k + 1, 2k + 1). Thus we get a map

$$Q(2(g-k)-3, 2k+1, -1^{2g+2}) \rightarrow Q(2(g-k)-3, 2(g-k)-3, 2k+1, 2k+1),$$

where $k \ge -1, g \ge 1$ and $g - k \ge 2$. Computing the dimensions of the strata, we get

$$\dim_{\mathbb{C}} \mathcal{Q}(2(g-k)-3,2k+1,-1^{2g+2}) = 2 \cdot 0 + (2g+4) - 2 = 2g+2$$
$$\dim_{\mathbb{C}} \mathcal{Q}(2(g-k)-3,2(g-k)-3,2k+1,2k+1) = 2g+4 - 2 = 2g+2$$

Since the dimension of the strata coincide, and the mapping is an immersion, we obtain an open set inside the stratum Q(2(g-k)-3, 2(g-k)-3, 2k+1, 2k+1). This remark, together with Proposition 1.5 motivate the following definition. We call the *hyperelliptic* connected component $Q^{hyp}(2(g-k)-3, 2(g-k)-3, 2k+1, 2k+1)$ of the stratum $Q^{hyp}(2(g-k)-3, 2(g-k)-3, 2k+1, 2k+1)$, consisting of quadratic differentials on hyperelliptic curves of genus g such that the quotient produces a quadratic differential with singularities data $Q(2(g-k)-3, 2k+1, -1)^{2g+2})$.

In [La1], we classify all components of this type. This motivates the following definition.

Definition. By *hyperelliptic components* we call the following connected components of the following strata of quadratic differentials.

- (1) The connected component $Q^{hyp}(2(g-k)-3, 2(g-k)-3, 2k+1, 2k+1)$ of the stratum Q(2(g-k)-3, 2(g-k)-3, 2k+1, 2k+1) consisting of quadratic differentials of the stratum $Q(2(g-k)-3, 2k+1, -1^{2g+2})$ where $k \ge -1$, $g \ge 1$, $g-k \ge 2$. The corresponding ramified double covering has ramification points over the 2g+2 poles.
- (2) The connected component $\mathcal{Q}^{hyp}(2(g-k)-3,2(g-k)-3,4k+2)$ of the stratum $\mathcal{Q}(2(g-k)-3,2(g-k)-3,4k+2)$ consisting of quadratic differentials of the stratum $\mathcal{Q}(2(g-k)-3,2k,-1^{2g+1})$ where $k \geq 0, g \geq 1$ and $g-k \geq 1$. The corresponding ramified double covering has ramification points over the 2g+1 poles and over the zero of degree 2k.
- (3) The connected component $\mathcal{Q}^{hyp}(4(g-k)-6,4k+2)$ of the stratum $\mathcal{Q}(4(g-k)-6,4k+2)$ consisting of quadratic differentials of the stratum $\mathcal{Q}(2(g-k)-4,2k,-1^{2g})$ where $k \ge 0, g \ge 2$ and $g-k \ge 2$.

The corresponding double ramified covering has ramification points over all the singularities.

Remark 1.6. Hyperelliptic connected components of type (1) were first discovered by Kontsevich.

1.1.4. *Main result*. We are finally in a position to give a precise statement of our result.

Theorem 1.1. Let $g \ge 5$ be any integer. Let us consider the following families \mathcal{F}_i , i = 2, 3, 4, of strata inside the moduli space of quadratic differentials \mathcal{Q}_q .

Then any stratum listed above has exactly two connected components: one is hyperelliptic — the other not.

Any other stratum of the moduli space \mathcal{Q}_g is non-empty and connected.

In small genera, some components are missing compared to the general case. The complete description is given by the following theorem.

Theorem 1.2. Let $g \leq 4$ be any non-negative integer. The components of the strata of the moduli space Q_q fall into the following description.

- In genera 0 and 1, any stratum is non-empty and connected, except Q(0) and Q(-1,1) which are empty.
- In genus 2, there are two non-connected strata: $\mathcal{Q}(-1, -1, 6)$ and $\mathcal{Q}(-1, -1, 3, 3)$; they have two components, one component is hyperelliptic the other not. Any other stratum of \mathcal{Q}_2 is connected.
- In genera 3 and 4, each stratum possessing an hyperelliptic connected component has exactly two connected components: one is hyperelliptic, the other not.
- There are 4 sporadic strata:

$$Q_3(-1,9), Q_3(-1,3,6), Q_3(-1,3,3,3), Q_4(12)$$

which have exactly two connected components.

• Any other stratum of Q_3 and Q_4 is connected.

Remark 1.7. The fact that the four sporadic components are not connected is due to Zorich (see [Zo1]). Zorich proved this result by a direct calculation in terms of *extended Rauzy classes*. It would be interesting to have an algebraic-geometric proof of the non-connectedness of these strata.

Remark 1.8. Using our approach of quadratic differentials, we get a new proof of a result of Masur and Smillie [MaSm] concerning the fact that the strata $\mathcal{Q}(0)$, $\mathcal{Q}(-1,1)$, $\mathcal{Q}(1,3)$ and $\mathcal{Q}(4)$ are empty.

1.2. Outline of the proof. The proof involves the dynamic and topology of measured foliations. We will use the well known fact that quadratic differentials q on S and pair of transverse measured foliations on S define the same objects (see [HuMa]). We will say that a component C_1 of a strata of Q_g is *adjacent* to a component C_2 if $C_2 \subset \overline{C_1}$ where the closure is taken inside the whole space Q_g .

1 (Claim A). Let \mathcal{C} be any component of any stratum $\mathcal{Q}_g(k_1, \ldots, k_n)$ with $n \geq 2$ and $g \geq 3$. We assume that \mathcal{C} is neither a hyperelliptic component nor a sporadic component $\mathcal{Q}^{irr}(-1,9)$, $\mathcal{Q}^{irr}(-1,3,6)$ or $\mathcal{Q}^{irr}(-1,3,3,3)$ (to be defined later on). Then \mathcal{C} is adjacent to a component of the stratum $\mathcal{Q}(4g-4)$.

2. We will use the following corollary of a theorem of Konsevich ([KoZo]): The number of connected components of a stratum of Q_g , which are adjacent to a component of the stratum Q(4g-4), is bounded by the number of components of Q(4g-4). In addition, for a given

stratum, there is at most one hyperelliptic or one sporadic component.

These two remarks, together with Claim A, provide us with a bound on the number of components of any stratum in terms of the number r_g of components of $\mathcal{Q}(4g-4)$. More precisely, the number of components of a statum containing a hyperelliptic component (or a sporadic component) is bounded upper by $r_g + 1$. The number of components of a statum containing nor a hyperelliptic component neither a sporadic component is bounded upper by r_g .

3. The list of stratum which possess a hyperelliptic component has been given in [La1]. In particular, these strata are non connected for $g \ge 3$. Hence the proof of Theorem 1.1 is reduced to the proof that the equality $r_g = 1$ holds for any $g \ge 5$.

The genera 3 and 4 are similar with some additional cases (sporadic strata). Theorem 1.2 follows from the equality $r_3 = 1$ and the inequality $r_4 \leq 2$ that we prove in Section 6. The small genera cases g = 1, 2 are considered separately; this is done in Section 8.

Therefore the proof of our result is reduced to the one of Claim A and to the computation of r_q .

We will say that a saddle connection γ on (S, q) has *multiplicity one* (see [MaZo]) if one can collapse γ to a point to get a new *closed* half-translation surface S'. In particular, this condition is satisfied if γ is "small" with respect to the other saddle connections in the direction of γ . More precisely if

(1.1)
$$|\gamma| < \frac{1}{2} |\eta|$$
, for any saddle connection η on $S \setminus \gamma$ in the direction $\overrightarrow{\gamma}$

then γ has multiplicity one. We will give a useful criterion on half-translation surfaces to have such an inequality.

4. Let $(S,q) \in C$ and γ be a multiplicity one saddle connection of (S,q). Then γ can be collapsed to a point to get a new half-translation surface (S',q'). The component C' containing (S',q') is contained in \overline{C} .

5. Let us assume that S is decomposed into a single cylinder for the horizontal flow. It means that any two regular horizontal geodesics are closed and homologous: they form a family which fill the surface into a metric cylinder. The boundaries of this cylinder consist of a set of saddle connections and separatrix loops. The arrangement of these separatrices is described by a "generalized permutation" π (see Section 3). We will note $(S, q) = S(\pi)$ for a one cylinder surface.

In Section 5 we give a *combinatorial criterion* (namely irreducibility) on the combinatorics of π so that there exists a saddle connection satisfying Equation (1.1); In particular this produces a multiplicity one saddle connection.

6. Let $\mathcal{S}(\pi) \in \mathcal{C}$ be a point in an arbitrary component (such points are dense in each stratum of the moduli space \mathcal{Q}_g). We consider the set of surfaces $\mathcal{S}(\pi_s) := h_s \cdot \mathcal{S}(\pi)$ where $h_s = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix}$.

If $S(\pi_s)$ has a multiplicity one saddle connection for any s, we are done thanks to 4. Otherwise the generalized permutation π_s must obey to some combinatorial conditions. Repeating that for any s we get that either $S(\pi)$ has a multiplicity one saddle connection or π is an exceptional (or hyperelliptic) permutation. The last case implies that C is an exceptional (or hyperelliptic) component contradicting the assumptions of Claim A.

7. The proof of the connectedness of $\mathcal{Q}(4g-4)$ (i.e. of the equality $r_g = 1$) for $g \geq 5$ is done by induction on the genus g of the surfaces. We first show by a direct computation that the minimal stratum in genus 5 is connected. We then use the surgery "Bubbling a handle" for the step of the induction. Precisely, we find in each component a surface with a cylinder filled by closed geodesic such that the boundary is formed by two single multiplicity one saddle connections. Therefore one can "erase" this cylinder to obtain a closed (g-1) half-translation surface.

The paper is organized in the following manner. In Section 2 we remind key results concerning the geometry of quadratic differentials. Then Sections 3–5 are devoted to the notion of generalized permutations. We develop this useful notion and give some relations between the combinatorics of π and the dynamic of the measured foliation on $S(\pi)$. Finally in Section 6 and Section 7 we prove respectively the two points 7 and 6.

1.3. Remarks and Acknowledgments. This paper is the last part of my PhD, the first two parts being the papers [La1] and [La2]. I would like to thank Pascal Hubert and Anton Zorich for encouraging me to write this paper.

I also thank Pascal Hubert, Maxim Kontsevich, Howard Masur, Anton Zorich and the anonymous referee for helpful comments on preliminary versions of this manuscript.

I thank the Institut de Mathématiques in Luminy, the Max-Planck-Institute for Mathematics in Bonn and the Centre de Physique Théorique in Marseille for excellent welcome and working conditions.

2. Preliminaries and preparatory material

In order to establish notations and preparatory material, we review basic notions concerning quadratic differentials versus half-translation surfaces. These surfaces have been considered and studied by numerous authors in various guises, see say [HuMa], [KeMaSm], [St] for more details; See also [EsMaZo] and [MaZo] for recent related developments on surgeries about half-translation surfaces.

2.1. Flat metrics.

2.1.1. Flat surfaces and geodesics. A half-translation surface is a (compact, connexe, real) genus g surface equipped with a flat metric (with isolated conical singularities) such that the holonomy group belongs to $\{\pm Id\}$. Here holonomy means that the parallel transport of a vector along a small loop going around a conical point brings the vector back to itself or to its negative. This implies that all cone angles are integer multiples of π . An equivalent definition is the following. A half-translation surface is a triple $(\mathcal{S}, \mathcal{U}, \Sigma)$ such that \mathcal{S} is a topological compact connected surface, Σ is a finite subset of \mathcal{S} (whose elements are called singularities) and $\mathcal{U} = (U_i, z_i)$ is an atlas of $\mathcal{S} \setminus \Sigma$ such that the transition maps $z_j \circ z_i^{-1}$: $z_i(U_i \cap U_j) \to z_j(U_i \cap U_j)$ are translations or half-translation: $z_i = \pm z_j + const$. This implies that the holonomy belongs to $\{\pm Id\}$.

Therefore, we get on S a flat metric with conical singularities located in Σ (possibly not all Σ). We also get a *quadratic differential* defined locally in the coordinates z_i by the formula $q = dz_i^2$. This form extends to the points of Σ to zeroes, poles or marked points (see [MaTa]). We will sometimes use the notation (S, q) or simply S.

Remark 2.1. The holonomy is trivial if and only if all transition functions are translations or equivalently the quadratic differentials q is the square of an Abelian differential. We will then say that S is a translation surface.

Hence the half-translation structure defines on $S \setminus \Sigma$ a Riemannian structure; we therefore have notions of directional foliation, geodesic, length, angle, measure, etc.

Convention.

- the *singularities* are the zeroes and poles of q.
- leaves of the directional foliation are called geodesics.
- geodesics meeting singularities are called *separatrices*. A geodesic emanating from a singularity and going back to the same is called a *separatrix loop*. A geodesic connecting two different singularities is called a *saddle connection*.
- a geodesic not passing through a singularity is called *regular*.

One can see the equivalence between half-translation surface and quadratic differential as follows. We start with the first definition. If we cut a flat surface S successively along an appropriate collection of saddle connections, we can decompose it into polygons contained in \mathbb{C} . We may then view S as a union of polygons with sides ordered by pairs consisting of parallels sides of the same length. The surface S is then isometric to the polygons where we identify these pair by translations or half-translations (translation post-composed with a central symmetry). Note that we have endowed each polygon with a complex coordinate. By construction, the transition functions in these complex coordinates z have the form

$$z \mapsto \pm z + const.$$

Thus any flat surface with the conical singularities removed is endowed with a natural complex structure. Moreover, consider a holomorphic quadratic differential $q = dz^2$ on every polygon. Since $dz^2 = d(\pm z + const)^2$ we obtain a globally well defined holomorphic quadratic differential on S'. It is a direct calculation to check that the complex structure and the quadratic differential can be extended to the singularities; the quadratic differential q extends to a (possibly meromorphic) form on S with zeroes or simple poles at every conical point. Note that when all transitions function are only translations, the quadratic differential q can be globally written as $q = \omega^2$, where ω is an Abelian differential. In this case the corresponding foliation is oriented.

Conversely, given a pair (S, q), and a point $P \in S$ such that $q(P) \neq 0$, the integral $\int_{z_0}^z \sqrt{q}$ produces a local coordinate z near P such that $q = dz^2$. Thus $|dz|^2$ defines a flat metric on S. At a singularity of multiplicity $k \geq -1$ the total angle we get is $(k + 2)\pi$. Remark that for regular point of q (k = 0), one get regular point of the metric.

2.1.2. $SL_2(\mathbb{R})$ -action. Given any matrix $A \in SL_2(\mathbb{R})$, we can post-compose the local coordinate of the charts of our translation atlas on (\mathcal{S}, q) by A. One easily checks that this gives a new half-translation surface, denoted by $A \cdot (\mathcal{S}, q)$. In local coordinates, this gives

$$A \cdot (\mathcal{S}, dz^2) = (\mathcal{S}, (dAz)^2).$$

Hence this produces an $SL_2(\mathbb{R})$ -action on these half-translation surfaces.

2.1.3. *Cylinders.* Closed regular geodesics appear in families of parallel geodesics of the same length. Such parallel closed geodesic, typically, do not filled the surface, but only a cylindrical subset. Each boundary component of such a cylinder is comprised by saddle connections.

Generically, each boundary component of a cylinder filled with closed regular geodesics is a single closed saddle connection. The converse, however, is false. A closed saddle connection does not necessarily bound a cylinder of regular closed geodesics. In fact, it bounds such a cylinder if and only if the angle at the singularities between the outgoing and incoming segments is exactly π . One calls such a cylinder a *simple* cylinder (see Section 6). See also Figure 7.

2.1.4. Adjacency. Let \mathcal{C} , \mathcal{C}' be two connected components of the moduli space \mathcal{Q}_g . We will say that \mathcal{C}' is adjacent to \mathcal{C} if $\mathcal{C} \subset \overline{\mathcal{C}'}$ (the closure being taken inside the whole space \mathcal{Q}_g).

2.1.5. *Example*. We end this section with an example of half-translation surface with Figure 3. Identifying pairs of sides of the polygon by isometries, we get a half-translation surface of genus g = 1. Note the form dz is not globally defined but dz^2 is; therefore the holonomy is exactly $\{\pm Id\}$.

2.2. **Ĥomologous saddle connections.** Let S be a half-translation surface. We denote by $\pi : \hat{S} \to S$ the standard orientating double covering so that $\pi^*q = \hat{\omega}^2$ (see [La2]). Let τ be the induced involution of the covering. Let γ be a *compact separatrix* on S. We consider γ^+ and γ^- the two lifts of γ by π . We choose an orientation of γ . According to this choice, we define

$$\hat{\gamma} = [\gamma^+] - [\gamma^-]$$

 $(\hat{\gamma} \text{ is well defined up to a sign})$. If P_i denote the singularities of q on S and \hat{P}_i the singularities of ω on \hat{S} , let $H_1^+(\hat{S}, \{\hat{P}_i\}, \mathbb{C})$ be the first homological group invariant with respect to the involution τ and $H_1^-(\hat{S}, \{\hat{P}_i\}, \mathbb{C})$ the first homological group anti-invariant with respect to the involution τ . We therefore get

$$\hat{\gamma} \in H_1^-/\pm$$

2.2.1. \hat{H} omologous saddle connections. Following Masur and Zorich [MaZo], we say that two compact separatrices γ and η are \hat{h} omologous if their corresponding loops $\hat{\gamma}$ and $\hat{\eta}$ on \hat{S} are proportional inside H_1^- .

Note that this definition does not depend of the choice of the orientation of the geodesics neither the choice of a direction on S. Moreover, γ and η are not supposed to be homeomorphic to a circle. For instance, one can have a saddle connection (homeomorphic to a segment) \hat{h} omologous to a separatrix loop (homeomorphic to a circle); see Figure 3. The following proposition gives a necessary condition for two compact separatrices to be \hat{h} omologous.

Proposition 2.2 (Masur, Zorich). Let us assume that γ and η are two compact separatrices. If γ and η are homologous then they are parallel and their lengths are equal or differ by a factor two:

$$\frac{|\gamma|}{|\eta|} \in \left\{1, 2, \frac{1}{2}\right\}$$

Example 2.3. In Figure 3, one can check that the vertical saddle connection $\gamma(\pi)$ and the vertical separatrix loop η are homologous. More precisely, we have $\widehat{\gamma(\pi)} = \hat{\eta}$.

2.2.2. Multiplicity. Let γ be a compact separatrix on S. We will say that γ has multiplicity n, and we will note mult $(\gamma) = n$, if there exists exactly n different classes $[\eta] \in H_1$ where η is a compact separatrices homologous to γ .

We will say that a simple cylinder has *multiplicity* n if the multiplicity of its boundary (which can be represented by a compact separatrix) is n.

Lemma 2.4. Let (S,q) be any half-translation surface in any hyperelliptic connected component of the following type:

$$\begin{array}{l} \mathcal{Q}^{hyp}(4(g-k)-6,4k+2) \\ \mathcal{Q}^{hyp}(2(g-k)-3,2(g-k)-3,4k+2) \\ \mathcal{Q}^{hyp}(2(g-k)-3,2(g-k)-3,2k+1,2k+1) \end{array}$$

For each of these strata, let γ be a saddle connection on (\mathcal{S}, q) between (respectively)

- the two zeroes of degree 4(g-k) 6 and 4k + 2.
- one of the two zeroes of degree 2(g-k) 3 and the other zero of degree 4k + 2.
- one of the two zeroes of degree 2(g k) 3 and one of the two zeroes of degree 2k + 1, 2k + 1.

Then γ has multiplicity at least two.

Proof of the lemma. The surface S is hyperelliptic so it is equipped with an hyperelliptic involution, say τ . Take a saddle connection γ as indicated in the assumptions. By construction [La1], $\gamma_2 := \tau(\gamma) \neq \gamma$ thus we obtain an other saddle connection in the same direction of γ (and of the same length). Always by construction we get that $\hat{\gamma} = \hat{\gamma}_2$. Therefore γ_2 is homologous to γ and $[gamma] \neq \gamma_2$ hence $\operatorname{mult}(\gamma) \leq 2$. The lemma is proven.

2.3. Surgeries. The principal ingredient of our proof is to decrease the complexity of a stratum. By complexity, we mean the genus g of the surfaces and the number n of the singularities. The next two subsections describe how one can increase the complexity. In Section 5 we will present some results to get the converse. For details and proofs of the two next sections we refer to [EsMaZo], [KoZo] and [MaZo].

2.3.1. Breaking up a singularity. Let (S, q) be a half-translation surface and $P \in S$ a singularity of q. Let $k\pi$ be the conical angle around P with $k \leq -1$. Choose any partition of k into two non-zero integers k_1, k_2 with $k_i \geq -1$. We recall the well known construction to obtain a new half-translation structure q' on S with the same singularities pattern as q except at the point P; The new half-translation surface will possess two singularities P_{k_1} and P_{k_2} of multiplicities k_1 and k_2 . Moreover S' will possess a multiplicity one saddle connection between P_{k_1} et P_{k_2} . Here we detail the case k odd and the case k_1, k_2 even.

Consider a small geodesic neighborhood of P, that is an ε -"polydisc" constructed from k + 2 half Euclidean discs of radii ε glued in their centers; Figure 1 (see also also Figure 4 and Figure 5 in [La1]).

Now, for ε small enough, there is no critical geodesic passing through this polydisc, other than the k + 2 emanating rays from P. Let us remove this polydisc and change continuously parameters in the following manner. We break up the singularity P into two singularities of conical angle $(k_1 + 2)\pi$ and $(k_2 + 2)\pi$. This is possible by the choice of k_i (see Figure 1 for details). Now one can re-glue this polydisc on the surface S to obtain the desired new half-translation structure on S.




FIGURE 1. Breaking up a zero of order 3 into two zeroes of orders 1 and 2 correspondingly. Note that the surgery is local: we do not change the flat metric outside of the neighborhood of the zero.

Remark 2.5. The new geodesic for q' on S which connects P_{k_1} and P_{k_2} has multiplicity one. Indeed one can choose its length arbitrary small without changing the others lengths thus Proposition 2.2 applies.

If \mathcal{C} denotes the component which contains (\mathcal{S}, q) and \mathcal{C}' denotes the component which contains (\mathcal{S}, q') then \mathcal{C}' is adjacent to \mathcal{C} .

The construction we have presented is local: we do not change the metric outside the ε -polydisc. In the case where the parameters k_1 and k_2 are odd, the construction is global. This is the *parallelogram construction* (see [MaZo]).

Theorem 2.6 (Masur, Zorich). Let $(S,q) \in Q(k_1, \ldots, k_n)$ be a point. Let us also assume that there exists a multiplicity one saddle connection on S between the singularities $P_1 \in S$ of order k_1 and $P_2 \in S$ of order k_2 . We will make that additional assumption that $\{k_1, k_2\} \neq \{-1, -1\}$.

Then there exists a point $(S',q') \in Q(k_1 + k_2, k_3, ..., k_n)$ such that one can break up the singularity $P'_{k_1+k_2} \in S'$ (for q') into two singularities to obtain the initial half-translation surface (S,q).

Proof of Theorem 2.6. Here we address the proof in cases of k_1 , k_2 even or $k_1 + k_2$ odd. Assume that γ is a multiplicity one saddle connection between P_1 and P_2 . As usual we will assume that γ is vertical. Using the geodesic flow, we contract it to a short segment of length δ . Choose any ε with $\frac{\delta}{2} < \varepsilon$. Now consider an ε -polydisc $D(\varepsilon)$ of these two points as indicated in Figure 1. The assumption on the multiplicity of γ implies that one can and do choose ε and δ small enough so that there are no critical (vertical) geodesics inside $D(\varepsilon)$ other than γ and the $k_1 + 2$ and $k_1 + 2$ verticals emanating from P_1 and P_2 . We can apply above construction to replace this polydisc by a new one, where one has glued the two singularities together. The new surface (\mathcal{S}, q') we have constructed satisfies the conclusions of the theorem: on can break up the singularity $P = P_1 = P_2$ into two to obtain our initial surface (\mathcal{S}, q) . The theorem is proven.

From the previous construction we easily gets the following useful result.

Corollary 2.7. Let S be a flat surface in the stratum $Q(k_1 + k_2, k_3, ..., k_n)$. Let us assume that $k_1 + k_2 \neq k_i$ for all i = 3, ..., n. Then all surfaces obtained from S by breaking up the

singularity of multiplicity $k_1 + k_2$ with discrete parameters k_1, k_2 (and arbitrary continuous parameters) belong to the same connected component of the stratum $Q(k_1, k_2, k_3, \ldots, k_n)$.

In terms of adjacency: If $C_0 \subseteq Q(k_1+k_2,\ldots,k_n)$ is a component and $C_1, C_2 \subset Q(k_1,k_2,\ldots,k_n)$ are two components such that $\mathcal{C}_0 \subset \overline{\mathcal{C}_1} \cap \overline{\mathcal{C}_2}$, then $\mathcal{C}_1 = \mathcal{C}_2$.

2.3.2. "bubbling a handle". Let (S,q) be a half-translation surface and $P \in S$ a singularity of q. Let us break up the singularity P into two singularities $P_1, P_2 \in \mathcal{S}$ (see previous section). One gets a new half-translation structure, say q_1 on \mathcal{S} , and a closed saddle connection γ (of length δ) between P_1 and P_2 . Let us cut this surface along γ . We obtain a surface with some boundaries components. We identify the two points P_1 , P_2 on this surface to obtain a surface \mathcal{S}_1 with a boundary component isometric to the union of two circles, each of length δ . Then let us glue a straight metric cylinder with the following parameters. The height and the twist are chosen arbitrarily, and of weight (circumference) is δ . The new surface (\mathcal{S}', q') we get is a $(\text{genus}(\mathcal{S})+1)$ half-translation surface. The angle between the new handle is k_1+2 (or k_2+2 if we consider the complementary angle). See Figure 7 for an example.

Let (\mathcal{S}, q) be any half-translation surface. Assume that the singularities data of \mathcal{S} are either (4g-3,-1) or (4g-4). Let s be any non-negative integer. Thanks to Corollary 2.7, the surfaces (\mathcal{S}', q') constructed by the surgery "bubbling a handle" at the unique zero of q in \mathcal{S} with arbitrary continuous parameters (height, twist and width of the cylinder) and fix discrete parameter s represent quadratic differentials belonging in the same connected component of the stratum. This motivates the following definition.

Definition 2.8. Let (\mathcal{S},q) be any half-translation surface, with singularities pattern (4g -(3, -1) or (4g - 4) and let s be any non-negative integer. We will denote by $\mathcal{S} \oplus s$ a surface constructed by the surgery "bubbling a handle" at the unique zero of q in S with arbitrary continuous parameters (height, twist and width of the cylinder) and discrete parameter s. If \mathcal{C} denote the component containing (\mathcal{S},q) then we denote the component containing (\mathcal{S}',q') by $\mathcal{C} \oplus s$. In other terms we obtain the two well defined mappings.

for any
$$g \ge 2$$
, $\oplus = \begin{pmatrix} \pi_0(\mathcal{Q}(4g-3,-1)) \times \mathbb{N} & \longrightarrow & \pi_0(\mathcal{Q}(4g+1,-1)) \\ (\mathcal{C},s) & \longmapsto & \mathcal{C} \oplus s \end{pmatrix}$
for any $g \ge 3$, $\oplus = \begin{pmatrix} \pi_0(\mathcal{Q}(4g-4)) \times \mathbb{N} & \longrightarrow & \pi_0(\mathcal{Q}(4g)) \\ (\mathcal{C},s) & \longmapsto & \mathcal{C} \oplus s \end{pmatrix}$

Here $\pi_0(E)$ denotes the set of connected components of the topological space E. Note also that the discrete parameter s corresponding to the angle between the two new sectors can be chosen modulo 2q ($s \in \{1, \ldots, 2q\}$).

Proposition 2.9. Let C be any connected component of any above stratum, namely $\mathcal{Q}(-1, 4g)$ 3) or $\mathcal{Q}(4g-4)$. Then the following statements hold.

- $\begin{array}{ll} (1) \quad \mathcal{C} \oplus s_1 \oplus s_2 = \mathcal{C} \oplus s_2 \oplus s_1 & \text{for any } s_1, \ s_2. \\ (2) \quad \mathcal{C} \oplus s_1 \oplus s_2 = \mathcal{C} \oplus (s_2 2) \oplus (s_1 + 2) & \text{for any } s_2 \geq 3. \\ (3) \quad \mathcal{C} \oplus s_1 \oplus s_2 = \mathcal{C} \oplus (s_2 4) \oplus s_1 & \text{for any } s_2 s_1 \geq 4. \end{array}$

Proof of Proposition 2.9. The proof uses the description of quadratic differentials in terms of separatrices diagrams (see [KoZo]) and ribbons graphs. For such diagrams, an element of $\mathcal{Q}(4q-4)$ (respectively of $\mathcal{Q}(-1, 4q-3)$) is presented by a measured foliation with 4q-4+2(respectively 4g - 3 + 2) leaves emanating from the singularity. Gluing an handle of angle $s\pi$ consists topologically to glue two news sectors with an angle of $s\pi$. We thus get measured foliation with 4g - 4 + 2 + 4 (respectively 4g - 3 + 2 + 4) leaves around the new singularity. Geometrically it is easy to see that the two surgeries consisting to first glue the handle of angle $s_1\pi$ or to first glue the handle of angle $s_2\pi$ produces surfaces belonging to the same component. This is the first point of the proposition.

The second statement is also clear in terms of diagrams. This is illustrated in Figure 2.




FIGURE 2. Gluing two handles attached on a singularity. Gluing first the "white" handle of angle $s_1\pi$ and then the "grey" handle of angle $s_2\pi$ is equivalent to glue first the "grey" handle of angle $(s_2 - 2)\pi$ and then the "white" handle of angle $(s_1 + 2)\pi$.

The last statements follows from the same considerations. The proposition is proven. \Box

2.4. Jenkins-Strebel surfaces. An important class of flat metrics is given by the so-called Jenkins-Strebel differentials. We first explain these particular metrics and then we give a density result.

Let us denote by $\Gamma(q)$ the critical graph of (S, q) induced by the horizontal foliation, that is the union of all separatrices in the horizontal direction. It is easy to see that Γ is compact if and only if the horizontal measured foliation of q is completely periodic; this means that any horizontal geodesic is closed. We then call such a form a *Jenkins-Strebel differential*.

Recall that locally, a stratum of the moduli space Q_g is modeled by the first cohomology group with coefficients in \mathbb{C} :

$$H^{1}(\mathcal{S}, \{P_{1}, \dots, P_{n}\}; \mathbb{Q}) \oplus i \cdot H^{1}(\mathcal{S}, \{P_{1}, \dots, P_{n}\}; \mathbb{Q}) \subset \subset H^{1}(\mathcal{S}, \{P_{1}, \dots, P_{n}\}; \mathbb{C}) = H^{1}(\mathcal{S}, \{P_{1}, \dots, P_{n}\}; \mathbb{R} \oplus i\mathbb{R})$$

Taking forms such that the absolute and relative periods are in $\mathbb{Q} \oplus i\mathbb{Q}$, we get arithmetic surfaces: the orientating covering is itself a ramified covering over the standard two torus. Therefore, up to normalize the direction to the horizontal direction, we get that Jenkins-Strebel differentials are dense in each stratum (see [DoHu], [HuMa], [KoZo] and [St]).

Remark 2.10. In [DoHu], Douady and Hubbard proved a stronger result: the Jenkins-Strebel differentials are dense on each Riemann surface, not just in a stratum.

The complement of $\Gamma(q)$ in S is a disjoint union of *maximal* periodic components for the horizontal foliation. These components are isometric to metric straight cylinders, foliated by regular horizontal leaves. A simple computation with the Euler characteristic, using the cylinders decomposition, shows that the number of the cylinders is bounded by 3g - 3.

In [Ma2] Masur proved that the set of Jenkins-Strebel differentials with exactly r cylinders (for any $1 \le r \le 3g-3$) is dense inside the principal stratum $\mathcal{Q}(1, \ldots, 1)$ of genus g. Recently, Kontsevich and Zorich [KoZo] have obtained a similar result. They have proved that the set of Jenkins-Strebel differentials with exactly one cylinder is dense inside any stratum of \mathcal{H}_g . Here we extend their proof to the case of strata of \mathcal{Q}_q .

Theorem 2.11. The set of quadratic differentials, such that the horizontal foliation is completely periodic and decomposes the surface into a unique straight metric cylinder, forms a dense subset of any connected component of any stratum of Q_g .

Proof of Theorem 2.11. We will only prove the existence of such surfaces on each connected component of the moduli space Q_g . For the density result, we refer to the Kontsevich-Zorich's proof. Note that in this paper, we will only need the existence result.

Let (S,q) be a point. Thanks to previous discussion we may assume that the surface (S,q) is an arithmetic surface: its orientating double covering \hat{S} covers the standard torus $\mathbb{T}^2 = \mathbb{C}/\mathbb{Z}^2$. The vertical foliation on S is completely periodic and decomposes the surface into many vertical cylinders C_i .

Let us construct a closed regular curve γ transverse to this foliation. The surface \hat{S} is a ramified covering $\pi : \hat{S} \to \mathbb{T}^2$. Obviously the directional flow on \hat{S} projects to the directional flow on the two-torus \mathbb{T}^2 . Let us consider a foliation on this torus in the direction $\theta = 1/b$ with b arbitrary large. The lift of this foliation allows us to obtain a closed regular geodesic γ on \hat{S} , and thus a closed geodesic on S. This leaf is transverse to the vertical foliation determined by q and γ does not contain any singularity of q. In addition one can choose γ in such a way that its length with respect to the metric defined by q is arbitrary large; Indeed the length of γ is greater than $\sqrt{1+b^2}$.

The closed loop γ cuts the boundaries of cylinders C_i many times (this means that γ cuts the set of vertical saddle connections and separatrix loops of q). By construction $\partial C_i \setminus \gamma$ is a disjoint union of vertical intervals. One can and do choose γ long enough so that each components of $\partial C_i \setminus \gamma$ contains at most one singularity of q. Now we will change slightly the transverse structure (in the vertical direction) to obtain a periodic horizontal foliation with only one cylinder. We will do that without changing the structure in the direction of γ .

We cut the surface along the vertical critical graph $\Gamma(q)$ of q and also along γ . We obtain a finite union of parallelograms R_i . Up to the $\mathrm{SL}_2(\mathbb{R})$ -action one may assume that γ is vertical. The set of horizontal sides of \mathbb{R}_i is a part of γ and the set of vertical sides of \mathbb{R}_i is a part $\Gamma(q)$. By construction in each vertical side of R_i there is at most one singularity of q.

Let us construct a new foliation as follows. We conserve all horizontal parameters and we change vertical parameters in the following way: we declare that the length of any vertical side of R_i is 1 for all *i*. In addition, if there is a singularity located on a vertical side, we declare that it is located at the middle of this side. With our above considerations, there is no contradiction. Finally we obtain a new set of parallelograms R'_i endowed with the natural metric dz^2 .

Let (S', q') be the flat surface constructed from the new rectangles R'_i with the corresponding identifications of vertical and horizontal sides given by gluing described above. We obtain a new half-translation structure q' on our surface.

The surfaces (S, q) and (S', q') carry the same topology. By construction the vertical critical graphs $\Gamma(q)$ and $\Gamma(q')$ coincide. We just have change absolute and relative periods of the form q. The subvariety of quadratic differentials sharing the same vertical foliation is connected and depends continuously on the suitable of deformations of the vertical foliation

(see [HuMa] and [Ve1]). Thus it implies that the two points (S, q) and (S', q') belong in the same connected component. It is easy to check that in the horizontal direction on S' for q' the foliation is completely periodic and decomposes the surface into a single cylinder.

3. Generalized permutations

In this section, we propose a natural way to encode Jenkins-Strebel differentials; namely we will introduce the notion of *generalized permutations*.

In this section (S, q) will denote a surface which is decomposed into a unique metric cylinder into the horizontal direction. Note that the holonomy of (S, q) is not assumed to be nontrivial.

3.1. Combinatorics of (S, q). Let $\Gamma(q)$ be the horizontal critical graph of (S, q). The complement $S \setminus \Gamma(q)$ is a metric cylinder Cyl(S). The horizontal saddle connections are labelled by $\gamma_1, \ldots, \gamma_k$.

One can encode the sequence of saddle connections contained in the bottom and in the top of $Cyl(\mathcal{S})$, and ordered following the cyclic ordering of the boundary of the cylinder, by a sequence of labels into the following manner. Each saddle connection γ_i is presented twice on the boundaries of $Cyl(\mathcal{S})$. Let us denote by γ_i^1 and γ_i^2 these two copies. Hence the top (respectively the bottom) of $Cyl(\mathcal{S})$ is a sequence of γ_i^{ϵ} where $i \in \{1, \ldots, k\}$ and $\epsilon = 1, 2$. Roughly speaking a generalized permutation π is a table with two lines encoding the sequence of labels of the saddle connections. The precise definition is the following.

Definition 3.1. Let r, l be any non-negative integers with the same parity. A generalized permutation π (of type (r, l)) is an involution, without fixed points, of the set $\{1, \ldots, r, r + 1, \ldots, r + l\}$.

Through this paper, we will represent a generalized permutation by a table.

Example 3.2. The generalized permutation (of type (7,5)) given by $\pi(1) = 10$, $\pi(2) = 12$, $\pi(3) = 5$, $\pi(4) = 7$, $\pi(5) = 3$, $\pi(6) = 11$, $\pi(7) = 4$, $\pi(8) = 9$, $\pi(9) = 8$, $\pi(10) = 1$, $\pi(11) = 6$, $\pi(12) = 2$ is represented by the table

in a natural way.

The term "generalized" is justified by the fact that a classical permutation π_1 of the symmetric group S_k is a generalized permutation with l = r = k and

$$\pi(i) = \begin{cases} k + \pi_1^{-1}(i), & \text{for } i \le k \\ \pi_1(i-k), & \text{for } i > k \end{cases}$$

In the present paper, we are interested by the quadratic differentials which are not the global square of any Abelian differential. Thus, in order to avoid "true" permutations, we require the following technical condition

(3.1) there exist
$$i_0 \leq r$$
 and $j_0 \geq r+1$ such that $\pi(i_0) \leq r$ and $\pi(j_0) \geq r+1$

3.2. Admissible vectors. Let π denote a generalized permutation of type (r, l).

Definition 3.3. We say that $\lambda \in \mathbb{R}^{r+l}_+$ is an admissible vector (for π) if

$$\begin{cases} \lambda_i = \lambda_{\pi(i)} \text{ for all } i = 1, \dots, r + \\ \sum_{i=1}^r \lambda_i = \sum_{j=1}^l \lambda_{r+j} \end{cases}$$

Note that for the "true" permutations π the vector

$$(\lambda_1,\ldots,\lambda_r,\lambda_{\pi(1)},\ldots,\lambda_{\pi(r)})$$

is admissible for any $\lambda_i > 0$.

3.3. Suspension over a generalized permutation. Let π be a generalized permutation and let λ be any admissible vector for π . We denote by w (width of the cylinder) the quantity

$$w := \sum_{i=1}^{r} \lambda_i = \sum_{j=1}^{l} \lambda_{r+j}$$

Let $\mathcal{C} = [0, w] \times [0, 1]$ be an Euclidean cylinder endowed with the form dz^2 . Let us consider the partition of the top (respectively bottom) of \mathcal{C} into horizontal intervals of length λ_i for $i = 1, \ldots, r$ (respectively i = r+1, dots, r+l). Now we identify the horizontal interval labeled i with the horizontal interval $\pi(i)$ for all i into the following manner. If the two intervals are presented twice on a side, we identify them by a centrally symmetry and otherwise we identify them by a translation.

The resulting space is a Riemann surface, denoted by $S(\pi, \lambda)$, endowed with a natural quadratic differential $q = dz^2$. We call this half-translation surface (S, q) the suspension over the element (π, λ) .




FIGURE 3. A suspension over a permutation π with an appropriate admissible vector λ . The resulting point $(\mathcal{S}(\pi, \lambda), dz^2)$ belongs to the stratum $\mathcal{Q}(-1, -1, 2)$. The black bullets correspond to the poles and the white bullets to the unique zero of the differential q on $\mathcal{S}(\pi, \lambda)$.

Notation. The surface $S = S(\pi, \lambda)$ decomposes into a single cylinder in the horizontal direction. By construction there exists in the vertical direction a compact separatrix on this surface. We will denote it by $\gamma(\pi) \subset S$.

Remark 3.4. If π a true permutation then the surface $S = S(\pi, \lambda)$ is a translation surface. The converse is true.

Lemma 3.5. The surfaces $S(\pi, \lambda)$ with fixed parameter π and arbitrary parameters λ belong to the same connected component.

Proof of the lemma. The lengths λ_i of the horizontal intervals correspond to the absolute and to the relative periods of the corresponding form q on S. Thus the lemma is a direct consequence of the local description of the orbifoldic structure of the strata in terms of the cohomological coordinates.

This construction implies a simple but important fact: we can encode the set of connected components using generalized permutations. Given a permutation, it determines completely the type of the singularities and hence a stratum. In addition, above lemma shows that it also determines the connected component of the stratum as well. The set of generalized permutations, for a fix stratum is obviously finite. Thus it gives an independent proof of a theorem of Veech [Ve1].

Theorem (Veech). The set of connected components of a stratum $Q(k_1, \ldots, k_n)$ of the moduli space Q_g of meromorphic quadratic differentials is finite.

3.4. Cyclic order and horocyclic flow. The class of Jenkins-Strebel quadratic differentials is stable under the horocyclic flow $h_s = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix}$. More precisely the corresponding suspended surfaces

$$h_s \cdot \mathcal{S}(\pi, \lambda) = \mathcal{S}(\pi', \lambda')$$

are related into the following way. Elements of π' corresponds to the elements of π with a "rotating" of the first line and the second line. We will say that π is equivalent to π' and we will note $\pi \sim \pi'$. For example the permutation of Example 3.2 is equivalent to the following one

Note that this relation preserves the stratum and also the connected component.




FIGURE 4. Action of the horocyclic flow h_s on a half-translation surface.

4. Representative elements

In this section, we give a "bestiary" of half-translations surfaces in terms of generalized permutations (see previous Section 3).

4.1. Hyperelliptic connected components. Let r, l be two non-negative integers. Let k = r+l. We construct the generalized "hyperelliptic" permutation $\Pi_1(r, l)$ of type (k+2, k+2) into the following way.

$$\Pi_1(r,l) = \begin{pmatrix} 0_1 & 1 & \dots & r & 0_1 & r+1 & \dots & r+l \\ r+l & \dots & r+1 & 0_2 & r & \dots & 1 & 0_2 \end{pmatrix}$$

A direct computation of the type of conical angles identifies the stratum which contain surfaces $S(\Pi_1(r,l),\lambda)$. It depends of the parity of the two integers r and l. We easily establish the next lemma.

Lemma 4.1. Let (S, q) be the half-translation surface given by $S(\Pi_1(r, l), \lambda)$ for any admissible vector λ . If r and l are odd then q has two singularities. If r and l have different parities then q has three singularities. If r and l are even then q has four singularities. The following table gives the type of the singularities in terms of r and l:

| r | l | stratum which contains (\mathcal{S},q) |
|--------|------------|--|
| 2k + 1 | 2(g-k) - 3 | ${\cal Q}(\ 4k+2 \ , \ 4(g-k)-6 \)$ |
| 2k + 2 | 2(g-k) - 3 | $\mathcal{Q}(\ 2k+1 \ , \ 2k+1 \ , \ 4(g-k)-6 \)$ |
| 2k + 2 | 2(g-k)-2 | $\mathcal{Q}(2k+1, 2k+1, 2(g-k)-3, 2(g-k)-3)$ |

According to [La1], each above stratum contains an hyperelliptic connected component. We have the following lemma.

Lemma 4.2. For any λ and any integers r, l, the surface $S(\Pi_1(r, l), \lambda)$ belongs to the hyperelliptic connected component (of the corresponding stratum).

Proof of Lemma 4.2. Here we present the proof of the first case; that is r and l are odd. The other cases are similar and left to the reader. Take r = 2k + 1 and l = 2(g - k) - 3. We consider the rectangle

$$\mathcal{R} = \left[-rac{r+l}{2} - 1, rac{r+l}{2} + 1 \right] \times \left[-1, 1 \right]$$

Let $\tau : \mathcal{R} \to \mathcal{R}$ be the involution of \mathcal{R} given by $\tau(x, y) = (-x, -y)$. The combinatorics of Π_1 implies that τ induces a *global* involution of the surface $\mathcal{S} = \mathcal{S}(\Pi_1(r, l), \lambda)$; we still denote it by τ . By Lemma 4.1 the surface (\mathcal{S}, q) belongs to the stratum $\mathcal{Q}(4k+2, 4(g-k)-6)$ therefore the Gauss-Bonnet formula implies that \mathcal{S} has genus g.

Recall that the hyperelliptic component of this stratum is, by definition, the image of the map

$$Q(2k, 2(g-k) - 4, -1^{2g}) \to Q(4k+2, 4(g-k) - 6)$$

 $(\mathbb{C}P(1), q_0) \mapsto (\mathcal{S}_0, \pi^* q_0)$

where $\pi : S_0 \to \mathbb{C}P(1)$ is a double ramified covering. The locus of ramification being the zeroes and poles of q_0 .

In order to prove that (\mathcal{S}, q) belongs to the hyperelliptic component, we have to construct a double ramified covering $\pi : \mathcal{S} \to \mathbb{P}^1$ and a quadratic differential q_0 on the sphere such that $\pi^* q_0 = q$. Let us count the number of fixed points of the map τ :

- there are r + l fixed points of τ on the horizontal sides of R located at the middle of the intervals (precisely at the middle of separatrix loops).
- there is a fixed point located at the middle of the vertical side.
- there is a fixed point at (0,0).

• there are 2 fixed points which corresponds to the two zeroes of q.

Therefore the total number of fixed points of τ on S_0 are

$$r + l + 1 + 1 + 2 = 2k + 1 + 2(g - k) - 3 + 4 = 2g + 2$$

The Riemann-Hurwitz formula implies that the genus of $S/(x \sim \tau(x))$ is zero. Let us consider the projection map

$$\pi: \mathcal{S} \to \mathcal{S} / (x \sim \tau(x)) \simeq \mathbb{C}P(1).$$

It is easy to check that above covering gives the desired map. Lemma 4.2 is proven. \Box

Remark 4.3. The permutation $\Pi_2(r, l)$ of type (2r, 2l) given by:

$$\Pi_2(r,l) = \left(\begin{array}{ccccc} 1 & \cdots & r & 1 & \cdots & r \\ r+1 & \cdots & r+l & r+1 & \cdots & r+l \end{array}\right)$$

furnishes also a representative element for hyperelliptic connected components.

Through the proof of Theorem 7.2 on adjacency of strata, we will get a characterization of hyperelliptic components. We will obtain the nice description.

Theorem 4.4. Let (S,q) be any point into any hyperelliptic component. Let us assume that S is decomposed into a unique metric cylinder for the horizontal flow. Then there exist $i \in \{1,2\}$, non-negative integers r, l, an admissible vector λ and $s \in \mathbb{R}$ such that

$$(\mathcal{S},q) = h_s \cdot \mathcal{S}(\Pi_i(r,l),\lambda)$$

4.2. Irreducible connected components. Here we give representative elements of the sporadic components discussed in Theorem 1.2: we will denote them by the *irreducible* connected components.

Definition 4.5. The irreducible connected components are defined to be the components containing the elements $S(\pi, \lambda)$ given by the next table.

| irreducible components | Representatives elements | | |
|-------------------------------|--|--|--|
| $\mathcal{Q}^{irr}(-1,9)$ | $\left(\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$ | | |
| $\mathcal{Q}^{irr}(-1,3,6)$ | $\left(\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$ | | |
| $\mathcal{Q}^{irr}(-1,3,3,3)$ | $\left(\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$ | | |
| $\mathcal{Q}^{irr,I}(12)$ | $\left(\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$ | | |
| $Q^{irr,II}(12)$ | $\left(\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$ | | |

5. Dynamical properties of $\mathcal{S}(\pi,\lambda)$ versus combinatorics of π

An important part of our proof is to find surfaces with multiplicity one saddle connection. For that we will give a combinatorial condition such that Proposition 2.2 applies.

5.1. Irreducibility. There exists "bad" permutations π such that, for any λ , the saddle connection $\gamma(\pi)$ has multiplicity at least two on $\mathcal{S}(\pi, \lambda)$. More precisely the two next cases can occur.

 $\forall \lambda, \exists$ saddle connection η such that $|\eta| = |\gamma(\pi_1)|$ on $\mathcal{S}(\pi_1, \lambda)$

and

$$\forall \lambda, \exists$$
 saddle connection η such that $|\eta| = 2|\gamma(\pi_2)|$ on $\mathcal{S}(\pi_2, \lambda)$

The first class of "bad" permutations will lead to the notion of weak reducibility. The second class will lead to the $Irred_2$ notion.

5.1.1. Weak irreducibility. We say that π (of type (r, l)) is weakly reducible if there exist $1 \leq i_0 < r$ and $r+1 \leq j_0 such that at least one of the following two conditions holds.$

- $\pi(\{1,\ldots,i_0\}) = \{r+1,\ldots,j_0\}$ or $\pi(\{i_0+1,\ldots,r\}) = \{r+j_0+1,\ldots,p\}$
- each $1 \leq i \leq r$ with $1 \leq \pi(i) \leq r$ satisfies $i \leq i_0$ and $\pi(i) > i_0$. All other $i \leq i_0$ with $\pi(i) \geq r+1$ satisfy $\pi(i) \leq j_0$. each $r+1 \leq j$ with $r+1 \leq \pi(j)$ satisfies $j \leq j_0$ and $\pi(j) > j_0$. All other $r+1 \leq j \leq j_0$ with $\pi(j) \leq r$ satisfy $\pi(j) \leq i_0$.

We will say that π is weakly irreducible if π is not weakly reducible.

Lemma 5.1. Let $S = S(\pi, \lambda)$ be a half-translation surface. Then π is weakly irreducible if and only if there exists a full Lebesgue measure set of admissible vectors λ such that any vertical separatrix η (different from $\gamma(\pi)$) on S satisfies $|\eta| \ge 2 \cdot |\gamma(\pi)|$.

The proof is obvious.

Notation. In order to clarify the situation, for a weakly reducible permutation π , we will denote by a vertical segment the position of the corresponding elements i_0 , j_0 . For instance the permutation

$$\left(\begin{array}{rrrrr|r}1 & 2 & 3 & 4 & 3 & 5\\6 & 1 & 2 & 6 & 5 & 4\end{array}\right)$$

is weakly reducible with corresponding $i_0 = 3$ and $j_0 = 9$.

5.1.2. The condition $Irred_2$. This is a technical condition and we first prefer to address the relating result to this notion. The definition will then become clear.

Proposition 5.2. Let $S = S(\pi, \lambda)$ be a half-translation surface. Let us assume that π satisfies the condition Irred₂. Then there exists a full Lebesgue measure set of admissible vectors λ such that any vertical separatrix η on S satisfies $|\eta| \neq 2 \cdot |\gamma(\pi)|$.

Definition 5.3. We say that π does not satisfy the condition Irred₂ if there exists a decomposition of π (up to exchange lines) into the following way (in terms of table).

$$\pi = \left(\begin{array}{cc} Y_1' & Y_1'' & Y_1''' \\ Y_2' & 0 & Y_2''' \\ \end{array}\right) \left(\begin{array}{cc} Y_1'' & Y_1''' \\ Y_2'' & 0 & Y_2''' \\ \end{array}\right)$$

with

$$\begin{cases} \forall i \in Y_1' \Rightarrow \pi(i) \in Y_1''' \sqcup Y_2' \\ \forall i \in Y_1'' \Rightarrow \pi(i) \in Y_1'' \sqcup Y_2'' \end{cases} \quad \text{and} \quad \begin{cases} \forall j \in Y_2' \Rightarrow \pi(j) \in Y_2''' \sqcup Y_1' \\ \forall j \in Y_2'' \Rightarrow \pi(j) \in Y_2'' \sqcup Y_1'' \end{cases}$$

Example 5.4. The permutation $\begin{pmatrix} 1 & 2 & 2 & 3 & 1 \\ 0 & 0 & \end{pmatrix}$ does not satisfy the condition Irred₂. Indeed, $Y'_1 = \{1\}, Y''_1 = \{2, 2, 3, 3\}, Y'''_1 = \{1\}$ and $Y'_2 = Y''_2 = Y''_2 = \emptyset$ is a decomposition. The permutation of Example 3.2 satisfies the condition Irred₂.

Proof of Proposition 5.2. As usual we will assume that $\gamma(\pi)$ has length 1. Recall that an admissible vector λ satisfy a linear equation given by Equation (3.3). Let us consider the set E of admissible vectors λ such that the entries λ_i defines a codimension one subspace. This is obviously a full Lebesgue measure set. Let us choose any $\lambda \in E$. We will prove that if there exists a vertical separatrix of length 2, then the above decomposition of Irred₂ occurs.

Let us assume that one has found a vertical separatrix $\eta \in S(\pi, \lambda)$ of length 2. A straightforward computation shows that one of the two cases presented by Figure 5 and Figure 6 have to occurs.




FIGURE 5. A separatrix of length 2 (here the canonical separatrix $\gamma(\pi)$ has length 1). The two corresponding horizontal intervals of length λ_{j_0+r} and λ_{i_1} , numbered by j_0 and i_1 , are glued by a translation.




FIGURE 6. A separatrix of length 2 (here the canonical separatrix $\gamma(\pi)$ has length 1). The two corresponding horizontal intervals of length λ_{j_0+r} and λ_{j_1+r} , numbered by j_0 and j_1 , are glued by a central symmetry.

Euclidean elementary geometry on the half-translation surface $\mathcal S$ on Figure 5 gives

$$\sum_{i=1}^{i_0} \lambda_i = \sum_{j=1}^{j_0} \lambda_{j+r} - \varepsilon \quad \text{and} \quad \sum_{i=1}^{i_1} \lambda_i - \varepsilon = \sum_{j=1}^{j_1} \lambda_{j+r}$$

Adding these two formulas and recalling that $\lambda_{i_1} = \lambda_{j_0+r}$, we get

(5.1)
$$2\sum_{i=1}^{i_0} \lambda_i + \sum_{i=i_0+1}^{i_1-1} \lambda_i = 2\sum_{j=1}^{j_0-1} \lambda_{j+r} + \sum_{j=j_0}^{j_1} \lambda_{j+r}$$

The same argument on Figure 6 produces similar equalities. More precisely

$$\sum_{i=1}^{i_0-1} \lambda_i = \sum_{j=1}^{j_0-1} \lambda_{j+r} + \varepsilon \quad \text{and} \quad \sum_{i=1}^{i_1} \lambda_i = \sum_{j=1}^{j_1} \lambda_{j+r} - \varepsilon$$

Adding these two formulas and recalling that $\lambda_{j_0+r} = \lambda_{j_1+r}$, we get the new one:

(5.2)
$$2\sum_{i=1}^{i_0-1} \lambda_i + \sum_{i=i_0}^{i_1} \lambda_i = 2\sum_{j=1}^{j_0-1} \lambda_{j+r} + \sum_{j=j_0+1}^{j_1-1} \lambda_{j+r} + 2\lambda_{j_0+r}$$

Recall that by assumption on λ , there exists exactly one rational relation between λ_i . This is the following one.

(5.3)
$$\sum_{i=1}^{r} \lambda_i = \sum_{j=1}^{l} \lambda_{r+j}$$

It is easy to check that Equation (5.1) can not occur: on the right part there is a term λ_{j_0+r} and on the left part, the corresponding term does not appear.

Therefore only the second case arise. Comparing the coefficients of λ_i , and forcing the terms to cancel (using Equation (5.3)) this leads to the fact that π does not satisfy Irred₂; with corresponding sets

$$\begin{array}{ll} Y_1' = (1, \dots, i_0) & Y_1'' = (i_0 + 1, \dots, i_1 - 1) & Y_1''' = (i_1, \dots, r) \\ Y_2' = (1, \dots, j_0 - 1) & Y_2'' = (j_0 + 1, \dots, j_1 - 1) & Y_2''' = (j_1 + 1, \dots, l) \end{array}$$

Therefore if λ belongs to E and π satisfies Irred₂, any vertical separatrix η has length different from 2. Proposition 5.2 is proven.

5.1.3. Irreducibility.

Definition 5.5. We say that π is *irreducible* if π is weakly irreducible and satisfies the condition Irred₂.

Proposition 5.6. Let us consider a surface $S(\pi, \lambda)$ with an irreducible permutation π . Then there exists a full Lebesgue measure set of admissible vectors λ such that any vertical separatrix $\eta \subset S(\pi, \lambda)$, different from $\gamma(\pi)$, satisfies $|\eta| \geq 3 \cdot |\gamma(\pi)|$.

Proof of Proposition 5.6. The proof is obvious using Lemma 5.1 and Proposition 5.2: the length of any separatrix is a positive integer different from 1 and 2 for a full Lebesgue measure set. \Box

Remark 5.7. Proposition 5.6 is related to Keane's i.d.o.c property. More precisely, For "true" permutations, weak irreducibility coincides with the classical definition. This means that $\pi\{1,\ldots,k\} \neq \{1,\ldots,k\}$ for any $k = 1,\ldots,r-1$. The Keane's property asserts that, for irreducible permutations, for almost all λ , any separatrix $\eta \neq \gamma(\pi)$ has infinite length. It will be interesting to have an analogous result for generalized permutations.

5.2. Irreducibility and weak irreducibility.

Definition 5.8. Let π be a type (r, l) generalized permutation. We say that π satisfies the Condition (*) if there exists only one element $i_0 \leq r$ (respectively $j_0 \geq r+1$) such that $\pi(i_0) \leq r$ (respectively $\pi(j_0) \geq r+1$).

We end this section by the following obvious lemma.

24

Lemma 5.9. Under condition (*), weak irreducibility implies irreducibility.

5.3. Irreducibility and "Breaking up a singularity". Let π be a generalized permutation and λ any admissible vector. Recall (Section 3) that for a horizontal separatrix β , we denote by β^1 and β^2 the two corresponding intervals on the cylinder Cyl(S).

Proposition 5.10. Let π be a generalized permutation and $S = S(\pi, \lambda)$ the suspended flat surface associated to λ . If π is irreducible then there exists a full Lebesgue measure set of λ such that $\gamma(\pi) \in S(\pi, \lambda)$ has multiplicity one.

Proof. It follows from Proposition 5.6 and properties of homologous separatrix detailed in Proposition 2.2. \Box

Proposition 5.11. Let π be a generalized permutation and $S = S(\pi, \lambda)$ the suspended flat surface associated to λ . We denote by β and η any two horizontal separatrices. Assume that one of the two following holds.

- β^1 and β^2 are located in two different horizontal boundaries of the cylinder $Cyl(\mathcal{S})$.
- all intervals β^1 , β^2 and η^1 , η^2 are located in the same horizontal boundary of $Cyl(\mathcal{S})$.

Then there exists a full Lebesgue measure set of λ such that β has multiplicity one.

Proof of Proposition 5.11. In the first case there is no condition on the horizontal parameter $|\beta^1| = |\beta^2| = |\beta|$ (see Equation (5.3). So one can and do choose the length of β in the flat metric, arbitrary small with respect to the other length of horizontal separatrices. Therefore Proposition 2.2 applies.

In the second case there is only one linear relation on the length of γ ; This is Equation (3.3). Therefore one gets:

$$|\beta^{1}| + |\beta^{2}| + |\eta^{1}| + |\eta^{2}| + \dots = \dots$$

In the left part of this equality, the terms β^i , η^i survives. In particular, we can choose $|\beta^1| + |\beta^2| + |\eta^1| + |\eta^2|$ arbitrary small and hence $|\beta^1| = |\beta^2| = |\beta|$ arbitrary small with respect to the length of other horizontal separatrices. We are done.

As direct corollary, one gets the following useful result.

Theorem 5.12. Let us assume that either $\gamma(\pi)$ or β (in above propositions) is a saddle connection connecting two different singularities; not two poles. Let us also assume that one of the assumptions of above Proposition 5.11 holds. Then the surface $S(\pi, \lambda)$ is obtained from the surgery "breaking up a singularity" on a surface in a lower dimensional stratum for a full Lebesgue measure set of λ .

5.4. Irreducibility and "Bubbling a handle".

Notation. Let π be a generalized permutation of the set $\{1, \ldots, r+l\}$ satisfying the condition $\pi(1) = r + 1$. We will denote by $\hat{\pi}$ the *restricted* generalized permutation of the set $\{\hat{1}, 2, \ldots, r, \hat{r+1}, r+2, r+l\}$ (where \hat{i} means that we forgot the element i). In terms of table, this gives

$$\pi = \left(\begin{array}{cc} 0 & A \\ 0 & B \end{array}\right) \qquad \text{and} \qquad \hat{\pi} = \left(\begin{array}{c} A \\ B \end{array}\right)$$

Clearly, the surface $S(\pi, \lambda)$, with π as above, possesses a simple cylinder in the vertical direction (see Figure 7 and Section 2.1.1). This cylinder is filled by regular vertical closed geodesics; each boundary component is a single vertical separatrix.

Using Theorem 5.12, one deduces the following useful result.




FIGURE 7. On the left, the figure represents a flat surface $\mathcal{S}(\pi, \lambda)$ with π satisfying $\pi(1) = r+1$. In the vertical direction, one can easily see that there is a simple cylinder. The boundary component of this cylinder is $\gamma(\pi) \sqcup \beta$. On the figure presented on the right, we have represented the diagram of the vertical foliation of $q = dz^2$. In this example, the angle of this cylinder is 4π (or 6π is we consider the complementary angle). If the generalized permutation $\hat{\pi}$ is irreducible, one can choose horizontal parameters so that $\gamma(\pi)$ has multiplicity one. In this case this surface is obtained from a surface in genus q-1, where $g = \text{genus}(\mathcal{S})$, by "bubbling a handle".

Theorem 5.13. Let $S(\pi, \lambda) \in Q_q(4g-4)$ be a point. Let us assume that $\pi(1) = r+1$. If $\hat{\pi}$ is irreducible then the surface $\mathcal S$ is obtained from the surgery "bubbling a handle" on a surface in a the stratum $\mathcal{Q}_{q-1}(4g-8)$ for a full Lebesgue measure set of λ .

Proof of Theorem 5.13. In the vertical direction, the surface S has a simple cylinder, for any λ . Let us remove it. We obtain a half-translation surface with boundaries. Each boundary component is a single geodesic circle: $\gamma(\pi)$ and one other say β . By construction, they have the same length. Let us remove the singularity and glue these two geodesics sequents together. We obtain a closed half-translation surface S' of genus g-1. The induced quadratic differential has two singularities say P_1 and P_2 of multiplicities k_1 , k_2 depending of the conical angle between $\gamma(\pi)$ and β . We denote this angle by $k\pi$. By definition

$$k_1 = k - 2$$
 and $k_2 = 4g - 6 - k$.

By assumptions $\hat{\pi}$ is irreducible. Thus Theorem 5.12 implies that one can choose λ in such a way that $\gamma(\pi)$ has multiplicity one. Applying Theorem 2.6 we can collapse $\gamma(\pi)$ to a point. Therefore we obtain a closed flat surface S'' of genus q-1 with a unique singularity. By construction, "bubbling a handle" at the unique zero of \mathcal{S}'' with the appropriate angle $k\pi$, we get the surface \mathcal{S} . The theorem is proven.

We end this section with the following lemma.

Lemma 5.14. The three following statements hold.

- (1) The component $\mathcal{Q}^{irr}(-1,3,6)$ is adjacent to the component $\mathcal{Q}^{irr}(-1,9)$.
- (2) The component $\mathcal{Q}^{irr}(-1,3,3,3)$ is adjacent to the component $\mathcal{Q}^{irr}(-1,3,6)$. (3) $\mathcal{Q}^{irr,I}(12) = \mathcal{Q}(8) \oplus 2$ and $\mathcal{Q}^{irr,II}(12) = \mathcal{Q}(8) \oplus 6$.
- (4) $Q^{irr}(-1,9) = Q(-1,5) \oplus 3.$

Proof of the Lemma 5.14. We first prove the two first assertions. Let $\mathcal{S}(\pi, \lambda) \in \mathcal{Q}^{irr}(-1, 3, 6)$ be a point with π as in Definition 4.5. Observing the horizontal foliation, it is easy to see that there exists a multiplicity one saddle connection connecting the two zeroes. Then the first assertion deduces from Proposition 2.6. One proves the second assertion in the same manner. Now let us concentrate with the third assertion. Let $S_1 = S(\pi_1, \lambda), S_2 = S(\pi_2, \lambda)$ be two representative elements of $Q^{irr,I}(12), Q^{irr,II}(12)$ with π_1, π_2 as in Definition 4.5 and $\lambda = (1, \ldots, 1)$.

We detail the first case, that is S_1 ; the second case is similar. The vertical foliation on S_1 is completely periodic and decomposes the surface into two cylinders (see Figure 8). One of




FIGURE 8. A vertical decomposition on a half-translation surface representing the component $Q^{irr,I}(12)$.

the cylinder is a simple cylinder and the angle between the separatrix loops which form the boundary is 2π . Hence $S_1 = Q(8) \oplus 2$. The proof of $S_2 = Q(8) \oplus 6$ is similar and left to the reader.

The last statement is a direct verification and left to the reader. The lemma is proven. \Box

We have now all necessary tools to prove Main Theorem 1.1 and Main Theorem 1.2.

6. The minimal stratum

This section is devoted to the so-called *minimal* stratum $\mathcal{Q}(4g-4)$ in case $g \geq 3$. We will prove this stratum is connected except for genus 4; for that we will show there exist at most two connected components.

The proof is done by induction on the genus of the surfaces. The step of induction is given by Theorem 6.3. The initialization of the induction is reduced to the proof of the connectedness of the stratum $Q_5(16)$ which we establish by a direct argument.

Theorem 6.1. Any connected component of the stratum Q(4g-4) is described by the following list.

- The stratum Q(8) is connected.
- The stratum Q(12) possesses at most two connected components corresponding to $Q^{irr,I}(12)$ and $Q^{irr,II}(12)$.
- Any other stratum $\mathcal{Q}(4g-4)$, in genus $g \geq 5$, is non-empty and connected.

Remark 6.2. The stratum $\mathcal{Q}(4g-4)$ is empty for $g \leq 2$ (see [MaSm]). Zorich (see [Zo1]) has proved that the stratum $\mathcal{Q}(12)$ is non-connected. The proof uses the Extended Rauzy classes.

6.1. Step of induction.

Theorem 6.3. Let C be a connected component of the stratum Q(4g - 4) in genus $g \ge 4$. Then there exist a half-translation surface $(S,q) \in Q(4g-8)$ and $s \in \mathbb{N}^*$ such that the surgery "bubbling a handle" at the unique singularity of q in S (with discrete parameter s and arbitrary

continuous parameters) produces surfaces belonging to the component C. In other words the map

$$\oplus : \pi_0(\mathcal{Q}_{g-1}(4g-8)) \times \mathbb{N}^* \to \pi_0(\mathcal{Q}_g(4g-4))$$

$$(\mathcal{C}', s) \mapsto \mathcal{C}' \oplus s$$

is onto for any $g \geq 4$.

Remark 6.4. Geometrically the previous statement is equivalent to find a half-translation surface in C with a multiplicity one simple cylinder. In order to do that we will use Theorem 5.13. Kontsevich and Zorich have obtained a similar result in the particular case of Abelian differentials. Their proof uses Rauzy classes. Here we first give an independent (geometric) proof of their result. We then give the proof in full generality.

Hence Theorem 6.3 is equivalent to the following.

Theorem 6.5. Let C be a connected component of the stratum Q(4g - 4) in genus $g \ge 4$. Then there exists $S(\pi, \lambda) \in C$ with $\hat{\pi}$ irreducible.

Proof of Theorem 6.3 for Abelian differentials. Let $C \subset \mathcal{H}(2g-2)$ be a connected component in genus $g \geq 2$. In order to find a surface with a multiplicity one simple cylinder in C we will use Section 5.4. Therefore it is sufficient to find a surface $(S, \omega) = S(\pi, \lambda) \in C$ so that $\hat{\pi}$ is irreducible.

Let π be a "true" permutation in the symmetric group S_r . We will assume that the surface $\mathcal{S}(\pi,\lambda) \in \mathcal{C}$ has no marked point that is $\pi(i+1) \neq \pi(i) + 1$ for all $i = 1, \ldots, r$ with the "dummy" condition $\pi(r+1) := \pi(1)$. We will show there exists a permutation π_1 in the class of π for the cyclic order with $\pi_1(1) = 1$ and such that $\hat{\pi}_1$ is irreducible. This will prove the theorem; Indeed the surface $\mathcal{S}(\pi_1, \lambda) \in \mathcal{C}$.

The differential ω has a unique singularity thus one can assume, using the cyclic order, that $\pi(1) = 1$. If $\hat{\pi}$ is irreducible then the theorem holds with $\pi_1 = \pi$. Otherwise let us assume that the restricted permutation $\hat{\pi}$ is reducible. Then by definition, there exists $2 \leq i_0 < r$ such that

$$\pi(\{2,\ldots,i_0\}) = \{2,\ldots,i_0\}.$$

Let us consider the following new set: $\pi(i_0 + 1, \ldots, r) = (A_1 \ r \ A_2)$. With these notations, we have

$$\begin{pmatrix} 1 & 2 & \dots & i_0 & i_0 + 1 & \dots & r \\ 1 & \dots & \dots & \pi(i_0) & A_1 & r & A_2 \end{pmatrix} = \pi \sim \pi_1 = = \begin{pmatrix} r & 1 & 2 & \dots & i_0 & i_0 + 1 & \dots & r - 1 \\ r & A_2 & 1 & \pi(2) & \dots & \dots & \pi(i_0) & A_1 \end{pmatrix}.$$

It is easy to see that $A_2 \neq \emptyset$: otherwise the corresponding surface $S(\pi_1, \lambda_1)$ will possess a marked point (see above). In particular $S(\pi, \lambda)$ possesses also a marked point which is a contradiction. Thus if $\hat{\pi}_1$ is reducible, it is easy to see that the corresponding invariant set (after renumbering elements) $\pi_1(\{2, \ldots, i'_0\}) = \{2, \ldots, i'_0\}$ will satisfy the condition $i'_0 \geq i_0 + 1 > i_0$. The set $\{0, \ldots, r\}$ is finite, thus the theorem follows by repeating finitely many times this process. Theorem 6.3 for Abelian differentials is proven.

To clarify the situation, we decompose the proof into several steps. We first prove the theorem in a weaker version: we add the additional assumption that there exists a surface $S(\pi, \lambda) \in C$

such that π satisfies the condition (*) (see Section 5.2). This corresponds to Proposition 6.6. We then prove Lemma 6.9 and Lemma 6.10 to get this additional condition.

Proposition 6.6. Let π be a generalized permutation such that $S(\pi, \lambda) \in Q(4g - 4)$ with $g \geq 4$. Let us assume that π satisfies the condition (*). Then there exists $\pi_1 \sim \pi$ such that $\hat{\pi}_1$ is irreducible.

We first prove the proposition for four particular permutations. This corresponds to Lemma 6.8.

Notation. A generalized permutation π is an ordered partition of $X = \{0, \ldots, r+l\}$ into two ordered lists, $X = Y_1 \sqcup Y_2$. In the present paper we shall always consider only those generalized permutations, for which each of Y_1 , Y_2 contains at least one entry of multiplicity two. The permutation satisfies the condition (*) so that is each set Y_1 , Y_2 contains *exactly* one entry of multiplicity two. Up to re-labeling, one can assume for the next that the two particular elements are $1 \in Y_1$ and $2 \in Y_2$. Finally, up to cyclic order, one can always put π into the form $\pi = \begin{pmatrix} 0 & A & C \\ 0 & B & D \end{pmatrix}$.

Note that if $\hat{\pi}$ is reducible, it involves one of the two following decomposition cases:

- (1) $\pi(A) = B \text{ or } \pi(C) = D$
- For all $i \in A$, with $i \neq 1$ one have $\pi(i) \in B$; $1 \in C$, $\pi(1) \in C$.
- (2) For all $j \in B$, with $j \neq 2$ one have $\pi(j) \in D$, $1 \in \mathbb{C}$, $\pi(1) \in \mathbb{C}$. For all $j \in B$, with $j \neq 2$ one have $\pi(j) \in A$; $2 \in D$, $\pi(2) \in D$.

In addition, we assume that this decomposition is minimal: we do not have a decomposition into smaller sets A', B', C', D'. This condition, in case of "true" permutation, is equivalent to ask that i_0 is minimal (see the proof versus Abelian differentials).

Lemma 6.7. If $\hat{\pi}$ involves reducibility of type (1) then one can find $\pi_1 \sim \pi$ such that $\hat{\pi}_1$ is irreducible.

Proof. Imitating the proof of Theorem 6.3 versus Abelian differentials, one can easily see that if $\hat{\pi}$ involves reducibility of type (1) then one can find $\pi_1 \sim \pi$ with $\hat{\pi}_1$ irreducible.

Lemma 6.8. Let π be a generalized permutation satisfying the condition (*). Assume also that one can put π into one of the four following forms:

 $\begin{pmatrix} 0 & A & 1 & C & | & 3 & 1 \\ 0 & B & 2 & D & | & 3 & 2 \end{pmatrix} \quad \begin{pmatrix} 0 & A & 1 & C & | & 1 \\ 0 & B & 2 & D & | & 2 \end{pmatrix} \quad \begin{pmatrix} 0 & A & 1 & C & | & 3 & 1 \\ 0 & B & 2 & D & | & 2 & 3 \end{pmatrix} \quad \begin{pmatrix} 0 & A & 1 & C & | & 4 & 3 & 1 \\ 0 & B & 2 & D & | & 4 & 2 & 3 \end{pmatrix}$

Let us assume that $\hat{\pi}$ is reducible (with corresponding marked invariant minimal sets) and $S(\pi, \lambda) \in Q(4g - 4)$ with $g \ge 4$. Then there exists $\pi_1 \sim \pi$ such that $\hat{\pi}_1$ is irreducible.

Proof of Lemma 6.8. Here we address the proof for the first class of permutations; the others being completely similar. By assumption on minimality of the decomposition, at least one of the two sets C or D is non-empty. Up to a permutation of the lines, we assume $C \neq \emptyset$. Up to re-labeling, let us denote $C := (C \ 4)$ with $\pi(4) \in B \sqcup D$. Depending the value of $\pi(4)$, we also put $B := (B_1 \ 4 \ B_2)$ or $D := (D_1 \ 4 \ D_2)$. Therefore π is equivalent to

 $\pi_B = \begin{pmatrix} 4 & 3 & 1 & 0 & A & 1 & C \\ 4 & B_2 & 2 & D & 3 & 2 & 0 & B_1 \end{pmatrix} \text{ if } \pi(4) \in B \quad \text{or } \pi_D = \begin{pmatrix} 4 & 3 & 1 & 0 & A & 1 & C \\ 4 & D_2 & 3 & 2 & 0 & B & 2 & D_1 \end{pmatrix} \text{ if } \pi(4) \in D$

This involves the two following cases:

Case 1. If B_2 or D are non-empty, the restricted permutation $\hat{\pi}_B$ is eventually reducible but in this case it involves decomposition of type (1); thus the lemma follows from Lemma 6.7. Therefore let us assume that $B_2 = D = \emptyset$. If B_1 is empty then the surface $S(\pi, \lambda)$ belongs to the stratum Q(8) and hence it has genus 3 which is a contradiction. Thus up to re-labeling,

one can put $B_1 := (B \ 5)$ with $\pi(5) \in A \sqcup C$. As above depending the value of $\pi(5)$, we also put $A := (A_1 \ 5 \ A_2)$ or $C := (C_1 \ 5 \ C_2)$. Therefore π is equivalent to

$$\pi_A = \begin{pmatrix} 5 & A_2 & 1 & C & 4 & 3 & 1 & 0 & A_1 \\ 5 & 4 & 2 & 3 & 2 & 0 & B \end{pmatrix} \text{ if } \pi(5) \in A \quad \text{or } \pi_C = \begin{pmatrix} 5 & C_2 & 4 & 3 & 1 & 0 & A & 1 & C_1 \\ 5 & 4 & 2 & 3 & 2 & 0 & B \end{pmatrix} \text{ if } \pi(5) \in C$$

Now we easily see that each of these permutations $\hat{\pi}_A$ and $\hat{\pi}_C$ is eventually reducible but it then involves decomposition of Type (1). Thus Lemma 6.7 applies.

Case 2. The discussion of this case is completely similar to the above one, depending the dichotomy $\pi(D_2) \subseteq A$ or not. We do not give the complete details here. The Lemma 6.8 is proven.

Proof of Proposition 6.6. Let π be a generalized permutation, satisfying condition (*), such that $S(\pi, \lambda) \in \mathcal{Q}(4g-4)$ with $g \geq 4$. If $\hat{\pi}$ is reducible then, according to previous notation and Lemma 6.7, one can assume that decomposition of Type (2) arises. We have $\pi = \begin{pmatrix} 0 & A & C \\ 0 & B & D \end{pmatrix}$. with $1 \in A$ and $\pi(1) \in C$, $2 \in B$ and $\pi(2) \in D$. Let us introduce the following new notations: $C = (C_1 \ 1 \ C_2)$ and $D = (D_1 \ 2 \ D_2)$. Then we put π into the following form $\pi = \begin{pmatrix} 0 & A & C \\ 0 & B & D \end{pmatrix}$.

Claim. Either there exists $\pi_1 \sim \pi$ with $\hat{\pi}_1$ irreducible or π can be put to one of the two forms (with $\pi(1) \in C$ and $\pi(2) \in D$):

$$\begin{pmatrix} 0 & A & C & 1 \\ 0 & B & D & 2 \end{pmatrix} \quad or \quad \begin{pmatrix} 0 & A & C & 1 & 3 \\ 0 & B & D & 3 & 2 \end{pmatrix}$$

Proof of the Claim. Obviously, if C_1 and C_2 are empty, the first form arises. Thus, by symmetry, let us assume $C_2 \neq \emptyset$. Up to re-labeling, we put $C_2 := (C_2 \ 3)$ with $\pi(3) \in D_1 \sqcup D_2$. As usual, depending the value of $\pi(3)$ we put $D_1 := (D'_1 \ 3 \ D''_1)$ or $D_2 := (D'_2 \ 3 \ D''_2)$. Thus, each case involves the two new permutations in the class of π :

$$\pi_{D_1} = \begin{pmatrix} 3 & 0 & A & C_1 & 1 & C_2 \\ 3 & D_1'' & 2 & D_2 & 0 & B & D_1' \end{pmatrix} \quad \text{or} \quad \pi_{D_2} = \begin{pmatrix} 3 & 0 & A & C_1 & 1 & C_2 \\ 3 & D_2'' & 0 & B & D_1 & 2 & D_2' \end{pmatrix}$$

For the first permutation: If $D_1'' = D_2 = \emptyset$ then the permutation is of the second class of the lemma thus we are done. Otherwise it is easy to see that the restricted permutation $\hat{\pi}_{D_1}$ is eventually reducible but then the invariant set is larger than A and we are done by repeating finitely many times this process.

For the second permutation: We have $D_2'' \neq \emptyset$. As above, the restricted permutation $\hat{\pi}_{D_2}$ is eventually reducible but then the new invariant set is larger than A and we are also done by repeating finitely many times this process. The claim is proven.

Now we will consider the two lists C, D of the claim. Each permutation of the claim produces two new classes of permutations. Here we do not give the details but the algorithm is completely similar to the one described above. Namely, the following holds:

Claim. Either there exists $\pi_1 \sim \pi$ with $\hat{\pi}_1$ irreducible or π can be put to one of the four forms:

 $\begin{pmatrix} 0 & A & 1 \\ 0 & B & 2 \end{pmatrix} \quad \begin{pmatrix} 0 & A & 3 & 1 \\ 0 & B & 3 & 2 \end{pmatrix} \quad \begin{pmatrix} 0 & A & 3 & 1 \\ 0 & B & 2 & 3 \end{pmatrix} \quad \begin{pmatrix} 0 & A & 4 & 3 & 1 \\ 0 & B & 4 & 2 & 3 \end{pmatrix}$

Now, let us remark that each of the permutations of the previous claim falls in the list of Lemma 6.8, and for those, we have already proved the proposition. Proposition 6.6 is proven. $\hfill \Box$

Lemma 6.9. Let C_0 be a connected component of the minimal stratum Q(4g-4). Then there exist two sequences of connected components

• $\mathcal{C}_i \subset \mathcal{Q}(4g-4)$ for $i=0,\ldots,m \ (m \geq 1)$


• $C^j \subset Q(k_j, 4g - 4 - k_j)$ for $j = 1, ..., m, k_j \ge 1$ and a flat surface $S = S(\pi, \lambda) \in C_m$ such that:

 $C_i \cup C_{i+1} \subset \overline{C^{i+1}}, \ \forall i = 0, \dots, m-1$ and π satisfies the condition (*).

Proof of Lemma 6.9. Let $S(\pi, \lambda) \in C_0$ be a point. If π satisfies the condition (*) we are done. Otherwise let us denote η_1, \ldots, η_m the set of horizontal separatrices such that η_i^1 and η_i^2 belong inside a same boundary component of the cylinder Cyl(S). By assumption $m \geq 3$. Let us "break up" the unique zero into two zeroes to obtain a new surface, say S'_1 , which belong to a component C^1 . By Proposition 5.11 and the fact that $m \geq 3$, the saddle connection η_1 on S'_1 has multiplicity one so one can it to a point. Thanks to this process we get a new surface S_1 in the minimal stratum, which belong to a component C_1 (eventually different from C_0). By construction we have

$$C_0 \cup C_1 \subset \overline{\mathcal{C}^1}$$

Repeating inductively this process on saddle connection η_i for i = 2, ..., m-1 we obtain the following diagram



with $C_i \cup C_{i+1} \subset \overline{C^{i+1}}$.

Finally, on the surface $S_m(\pi_m, \lambda_m) \in C_m$, by construction there exists a single pair of horizontal separatrices η_m and η_{m-1} such that η_i^1 and η_i^2 belong to the same boundary component of the cylinder Cyl(S). In other word π_m satisfies the condition (*). Lemma 6.9 is proven. \Box

Lemma 6.10. Let $C_0 \subset Q(4g-4)$ be a connected component. Let us assume there exist two components $C_1 \subset Q(4g-4)$ and $C^1 \subset Q(k, 4g-4-k)$ such that

 $\mathcal{C}_0 \cup \mathcal{C}_1 \subset \overline{\mathcal{C}^1}$

Let us also assume there exists a flat surface $(S,q) \in C_1$ with a multiplicity one simple cylinder. Then there also exists a flat surface $(S',q') \in C_0$ with a multiplicity one simple cylinder.

Proof of Lemma 6.10. The proof is obvious using description of surfaces in terms of separatrices diagrams. \Box

Corollary 6.11. For each connected component C of Q(4g - 4) with $g \ge 4$, there exists a surface $S(\pi, \lambda) \in C$ such that π satisfies the condition (*).

Proof. It follows from Lemma 6.9 and Lemma 6.10.

Proof of Theorem 6.3. It follows obviously from Corollary 6.11 and Proposition 6.6. \Box

6.2. Connectedness of the minimal stratum. We are ready to prove Theorem 6.1. First we directly show that Q(4g - 4) is connected for g = 3, 5 and has at most two components for g = 4. We then prove the theorem inductively on g.

Lemma 6.12.

The stratum Q(8) is connected. The stratum Q(12) has at most two connected components.

Proof of Lemma 6.12. See Lemma A.1 and Lemma A.2 in the appendix.

Lemma 6.13. The stratum Q(16) is connected.

Proof of Lemma 6.13. Thanks to previous lemma let $C_0 = \mathcal{Q}(8)$ be the unique connected component of the stratum $\mathcal{Q}(8)$. Let $\mathcal{C}_1 = \mathcal{C}_0 \oplus 2 \oplus 2$ be the component of $\mathcal{Q}_5(16)$ obtaining by bubbling two handles on a surface of \mathcal{C}_0 with an angle of 2π . Recall that

$$\mathcal{C}_0 \oplus 2 = \mathcal{Q}^{irr,I}(12)$$
 and $\mathcal{C}_0 \oplus 6 = \mathcal{Q}^{irr,II}(12)$

Let \mathcal{C}' be any component of $\mathcal{Q}(16)$. By Theorem 6.3, there exists s_0 such that $\mathcal{C}' = \mathcal{C} \oplus s_0$ where \mathcal{C} is a component of $\mathcal{Q}(12) = \mathcal{Q}^{irr,I}(12) \cup \mathcal{Q}^{irr,II}(12)$. Using the properties of the map \oplus (Proposition 2.9), we directly get the relations

$$\mathcal{C}_0 \oplus 2 \oplus s = \mathcal{C}_0 \oplus 6 \oplus s = \mathcal{C}_1$$
 for any $s = 1, \dots, 8$.

Thus $\mathcal{C}' = \mathcal{C} \oplus s_0 = \mathcal{C}_1$ which proves the lemma.

Proof of Theorem 6.1. We process by induction, initialization being given by Lemma 6.13. Let us fix g > 5. Let us assume that $\mathcal{Q}(4g'-4)$ is connected for all genera $5 \leq g' < g$. We have to show that $\mathcal{Q}(4g-4)$ is connected. We denote by $\mathcal{C}_{g-1} = \mathcal{Q}(4(g-1)-4)$ the unique connected component of this stratum. We also define $\mathcal{C}_g \subseteq \mathcal{Q}(4g-4)$ by

$$\mathcal{C}_g = \mathcal{C}_{g-1} \oplus 1$$

By Theorem 6.3 the map

$$\oplus : \pi_0(\mathcal{Q}_{g-1}(4g-8)) \times \mathbb{N}^* \to \pi_0(\mathcal{Q}_g(4g-4))$$
$$(\mathcal{C}', s) \mapsto \mathcal{C}' \oplus s$$

is onto. But the stratum $\mathcal{Q}(4(g-1)-4)$ is connected and s can be chosen in $\{1, \ldots, 2g-2\}$ (up to consider the complementary angle). Thus we obtain a (onto) map

$$\oplus : \{1, \dots, 2g - 2\} \to \pi_0(\mathcal{Q}_g(4g - 4)) \\ s \mapsto \mathcal{C}_{g-1} \oplus s$$

In order to end the proof, it remains to show that

(6.1)
$$C_g = C_{g-1} \oplus s$$
 for any s.

Now always by Theorem 6.3, there exists r_0 such that

$$\mathcal{C}_{g-1} = \mathcal{C}_{g-2} \oplus r_0$$
 with $\mathcal{C}_{g-2} \subseteq \mathcal{Q}(4(g-2)-8)$

(this stratum is non-empty because $g-2 \ge 4$). But recalling that the stratum $\mathcal{Q}(4(g-1)-4)$ $(g-1 \ge 5)$ is connected, we also have

$$\mathcal{C}_{g-2} \oplus s = \mathcal{C}_{g-1}$$
 for any s

Using properties of the map \oplus , this yields

$$\mathcal{C}_g = \mathcal{C}_{g-1} \oplus 1 = (\mathcal{C}_{g-2} \oplus s) \oplus 1 = (\mathcal{C}_{g-2} \oplus 1) \oplus s = \mathcal{C}_{g-1} \oplus s \quad \text{for any } s$$

Thus we get the desired relation (6.1). Theorem 6.1 is proven.

32

7. Adjacency of the strata

Theorem 7.1. Let $C \subset Q_g(k_1, \ldots, k_n)$ be a connected component with any $g \geq 3$, $n \geq 2$. Let us make the additional assumption that C is neither an hyperelliptic component nor an irreducible component. Then there exists a connected component $C_0 \subset Q(4g-4)$ such that $C_0 \subset \overline{C}$.

We will deduce Theorem 7.1 from the following one.

Theorem 7.2. Let $C \subset Q_g(k_1, \ldots, k_n)$ be a connected component with $g \ge 1$ and $n \ge 2$. Let us assume that C is none of one of the following:

- the hyperelliptic connected component $\mathcal{Q}^{hyp}(4(g-k)-6,4k+2)$ or $\mathcal{Q}^{hyp}(-1,-1,4g-2)$
- a component of $\mathcal{Q}(-1,5)$
- the irreducible component $Q^{irr}(-1,9)$

Then there exists a flat surface $(S,q) \in Q_g$ in a lower dimensional stratum and a surgery "breaking up a singularity" at a singularity of q on S such that the resulting surface (S',q') belongs to C.

Remark 7.3. Surprisingly, the answer is quite difficult and we find that some components, in small genera, are not hyperelliptic neither adjacent to the minimal stratum. Note also that the corresponding statement for Abelian differentials is trivial.

7.1. Link with our main result. Following the description of the topology of the strata of the moduli space in local coordinates (see [KoZo]), one has

Theorem 7.4 (Kontsevich). For any $(S,q) \in \mathcal{Q}(4g-4)$ with $g \geq 3$ there exists a small open set U(S,q) of (S,q) in the whole space \mathcal{Q}_g such that

$$\mathcal{Q}(k_1,\ldots,k_n)\cap U(\mathcal{S},q)$$

is non-empty and connected for any k_i with $\sum k_i = 4g - 4$.

Combining this result with Theorem 7.1 we get an upper bound for the number of connected components of any stratum. Let us denote by r_g the number of connected components of the minimal stratum $\mathcal{Q}(4g-4)$. Then for any $g \geq 3$, Theorem 7.1 and Theorem 7.4 together give

 $1 \leq \# \left\{ \begin{array}{l} \text{components of a statum which do not contain} \\ \text{a hyperelliptic component neither an exceptional component} \right\} \leq r_g$ $2 \leq \# \{ \text{components of statum which contain a hyperelliptic component} \} \leq r_g + 1$ $1 \leq \# \{ \text{components of statum which contain an exceptional component} \} \leq r_g + 1$

In Section 6 we have proved that $r_q = 1$ for $g \ge 5$ which implies main Theorem 1.1.

7.2. Strategy of the proof of Theorem 7.2. Let \mathcal{C} be a connected component of the stratum $\mathcal{Q}_g(k_1, \ldots, k_n)$ with $g \ge 1$ and $n \ge 2$. According to Theorem 2.6 one has to construct a surface $(\mathcal{S}, q) \in \mathcal{C}$ with a multiplicity one separatrix between two *different singularities*, not two poles. In order to do that we will use the criterion given by Proposition 5.11 and Proposition 5.10.

Let $(S(\pi, \lambda), q)$ be a point in C. Either Proposition 5.11 applies and then we have done or assumptions of the proposition does not hold. In this last case we get some restrictions on the combinatorics of π .

Now let us consider the vertical foliation on $(\mathcal{S}(\pi, \lambda), q)$. As above either Proposition 5.10 applies and we have done or assumptions of the proposition again does not hold. In this last

case we get also new restrictions on the combinatorics of π . In particular a simple computation shows that the permutation π is completely determined. It corresponds to a "hyperelliptic" or "irreducible" permutation. These two cases are avoid; indeed the component C is not hyperelliptic neither irreducible.

The proof is decomposed into several cases. Recall that n denote the number of singularities. First we consider the general case $n \ge 4$. We then prove the case n = 3. Finally we conclude with the holomorphic case n = 2 and the meromorphic case n = 2: the stratum $\mathcal{Q}(-1, 4g-3)$.

Remark 7.5. The corresponding statement of Theorem 7.2 versus Abelian differentials is trivial. Indeed Proposition 5.11 applies directly (all transitions functions are given by translations).

7.3. Proof of Theorem 7.2 in case $n \ge 4$. Let \mathcal{C} be a connected component and $\mathcal{S}(\pi, \lambda) \in \mathcal{C}$. We denote the horizontal cylinder of \mathcal{S} by $Cyl(\mathcal{S})$. The boundaries components of $Cyl(\mathcal{S})$ are denoted by I and J. One associates to each saddle connection η two intervals η^1 , η^2 on $I \sqcup J$.

Claim 1. There exists a saddle connection α between a zero P_1 and another singularity P_2 .

Proof of the claim. The assumption $g \ge 1$ implies there exists at least a zero P_1 . If there exists a saddle connection attached to this zero we are done. Otherwise any separatrix emanating from P_1 is actually a separatrix loop.

Exchanging the role of I and J if necessary, the previous assertion means that any saddle connection located in I is a separatrix loop emanating from P_1 (see Figure 9a).

We claim there exists a separatrix attached to a zero, say $P'_1 \neq P_1$, located on J. Otherwise it will mean that any separatrix in J is attached to a pole. The only possibility to obtain that is presented in Figure 9b. In particular the stratum is $\mathcal{Q}(k, -1, -1)$ contradicting the assumption $n \geq 4$. If there exists no saddle connection attached to P'_1 then any separatrix




FIGURE 9.

located on J is a separatrix loop emanating from P'_1 . It implies that the surface has only two zeroes. This is again a contradiction and the claim is proven.

Let α be a saddle connection between the zero P_1 and the singularity P_2 (we assume that $\alpha^1 \subset I$). If $\alpha^2 \subset J$ then Proposition 5.11 implies that α has multiplicity one and the theorem is proven. Hence let us assume that

$$\alpha^1, \ \alpha^2 \subset I$$

(we refer to Figure 10a).

The complement of α^1, α^2 in I has two connected components. Following the figure, we will refer to these two components as I_1, I_2 . The interval I_i is a union of separatrices.

COMPONENTS OF THE STRATA OF THE MODULI SPACES




FIGURE 10.

Claim 2. Either the theorem is proved or any separatrix τ , with $\tau^1 \subset I_i$, is a separatrix loop attached to P_i . In addition $\tau^2 \subset J$ (eventually $I_2 = \emptyset$).

Proof of the claim. Obviously one has: Either there exists a saddle connection τ between two singularities (not two poles) with $\tau^1 \subset I_1$ or any separatrix of I_1 is a separatrix loop attached to P_1 .

The first case is the conclusion of the theorem. Thus let us consider the second case. Let τ be a separatrix with $\tau^1 \subset I_1$. Then $\tau^2 \subset I \sqcup J$. Recalling that α^1 , $\alpha^2 \subset I$ and applying Proposition 5.11 we get that if $\tau^2 \subset I$ then α has multiplicity one and therefore the theorem is again proved. Otherwise $\tau^2 \subset J$ giving the second conclusion of the claim.

Repeating the same argument for separatrices τ with $\tau^1 \subset I_2$, we get the claim.

Let us assume that the second conclusion of Claim 2 holds. The singularity P_1 is a zero hence there exists a separatrix loop β attached to P_1 with $\alpha_1 \subset I_1$. Thanks to Claim 2 $\alpha^2 \subset J$ (see Figure 10b). In particular there exists a saddle connection η between P_1 and a singularity $P_3 \neq P_1$ (possibly $P_3 = P_2$) in such a way that $\eta^1 \subset J$. If $\eta^2 \subset I$ then thanks to Proposition 5.11, η has multiplicity one and the theorem is proved. Thus one can and do assume $\eta^2 \subset J$. The complement of η^1, η^2 in J has two connected components. Following the figure, we will refer to these two components as J_1, J_3 . The interval J_i is a union of separatrices.

Claim 3. Either the theorem is proved or any separatrix τ , with $\tau^1 \subset J_i$, is a separatrix loop attached to P_i . In addition $\tau^2 \subset I$.

Proof of the claim. The proof parallels the one of previous Claim 2.

Finally thanks the previous claims, either the theorem is proved or S can be putted into the form prescribed by Figure 10c. The last case implies that S has at most three singularities: P_1, P_2, P contradicting $n \ge 4$. Therefore Theorem 7.2 in case $n \ge 4$ is proven.

7.4. Proof of Theorem 7.2 in case n = 3. The proof is similar to the case $n \ge 4$ with a refinement. We first prove the following lemma.

Lemma 7.6. Let $C \subset Q_g(k_1, k_2, k_3)$ be a connected component with $g \ge 1$. Then there exists $(S(\pi, \lambda), q) \in C$ a half-translation surface such that one of the two following holds.

- $(S(\pi, \lambda), q)$ has a multiplicity one saddle connection between two different singularities (not two poles)
- The singularity pattern is exactly $(k_1, k_2, k_3) = (-1, -1, 4g 2)$ and the combinatorics of π has the form

$$\pi = \begin{pmatrix} 1 & \dots & r & 0_1 & 0_1 \\ 0_2 & 0_2 & \pi(1) & \dots & \pi(r) \end{pmatrix} \quad or \quad \pi = \begin{pmatrix} A \\ 0 & 0 \end{pmatrix} \text{ with } \pi(A) = A$$

Proof of Lemma 7.6. Let $S(\pi, \lambda)$ be a point in C. Imitating the proof of Claim 1 one gets that either there exists a saddle connection α between a zero and another singularity or one can put the surface S into the form prescribed by Figure 9b. The last case produces a surface $S(\pi, \lambda)$ belonging to the stratum Q(k, -1, -1) with $\pi = \begin{pmatrix} A \\ 0 & 0 \end{pmatrix}$ and $\pi(A) = A$; and so the lemma is proven.

Therefore let us assume S possesses a saddle connection α between a zero and another singularity. Let us again apply conclusions of Claim 2 and Claim 3. This leads to the following dichotomy. Either S possesses a multiplicity one saddle connection, and so the lemma is proven, or S can put put into the form prescribed by Figure 10c.

Finally one can assume the following situation: Any separatrix τ belonging to I_1 or J_1 is a separatrix loop attached to P_1 . In addition $\tau^1 \subset I_1$ and $\tau^2 \subset J_1$. Moreover n = 3 thus $P_3 \neq P_2$.

If P_2 is not a pole then there exists a separatrix loop τ attached to P_2 with $\tau^1 \subset I_2$. If $\tau^2 \subset J$ then $\tau^2 \subset J_3$ implying $P_2 = P_3$ which is contradiction. Therefore $\tau^2 \subset I$. Now Proposition 5.11 implies (indeed $\alpha^1, \alpha^2 \subset I$) that α has multiplicity one and so the lemma is proven.

The same proof holds if P_3 is not a pole.

Therefore let us assume that P_2 and P_3 are two poles. Then the stratum containing S is Q(k, -1, -1) and the permutation π has the combinatorics $\begin{pmatrix} 1 & \dots & r & 0_1 & 0_1 \\ 0_2 & 0_2 & \pi(1) & \dots & \pi(r) \end{pmatrix}$. The lemma is proven.

Therefore in order to prove Theorem 7.2 in case n = 3 we have to analyse the combinatorics of two permutations given in Lemma 7.6.

Lemma. Let $\pi = \begin{pmatrix} 2 & \dots & r & 0_1 & 0_1 & 1 \\ 0_2 & \pi(1) & \pi(2) & \dots & \pi(r) & 0_2 \end{pmatrix}$. Assume π is different (up to the cyclic order) from $\Pi_1(r, 0)$. Then there exists $\pi' \sim \pi$ such that π' is irreducible and $\gamma(\pi')$ is a saddle connection between the zero and a pole.

Proof of the lemma. First of all, let us remark that the separatrix $\gamma(\pi)$ is actually a saddle connection. Let us also remark that π satisfies the condition (*), therefore by Lemma 5.9, weakly irreducibility implies irreducibility for any permutation in the class of π .

The above generalized permutation π is reducible if and only if $\pi(r) = 1$. One observes that

$$\pi = \left(\begin{smallmatrix} 2 & \dots & r & 0_1 & 0_1 & 1 \\ 0_2 & \pi(1) & \pi(2) & \dots & \pi(r) & 0_2 \end{smallmatrix}\right) \sim \pi' = \left(\begin{smallmatrix} 3 & \dots & r & 0_1 & 0_1 & 1 & 2 \\ 0_2 & \pi(1) & \pi(2) & \dots & \pi(r-1) & 1 & 0_2 \end{smallmatrix}\right).$$

It is easy to check that π' is irreducible if and only if $\pi(r-1) = 2$.

Therefore repeating finitely many times this process, one shows that either there exists an irreducible permutation π' in the class of π or

$$\pi(i) = r - i + 1 \qquad \text{for all } i = 1, \dots r.$$

The last equations mean that $\pi \sim \Pi_1(r, 0)$ which is a contradiction. The lemma is proven. \Box

Lemma. Let $\pi = \begin{pmatrix} A \\ 0 & 0 \end{pmatrix}$ with $\pi(A) = A$. Assume π is different (up to the cyclic order) from $\Pi_2(r, 1)$. Then there exists $\pi' \sim \pi$ such that π' is irreducible and $\gamma(\pi')$ is a saddle connection between the zero and a pole.

Proof. It is easy to see that for each permutation π' in the class of π , the separatrix $\gamma(\pi')$ is actually a saddle connection between the zero and a pole. If π is weakly reducible then by definition, we have $\pi = \begin{pmatrix} B & C \\ 0 & 0 \end{pmatrix}$ with $\pi(B) = C$. Let us decompose the two lists B, C into the following way: $B = (1 \ B_2)$ and $C = (C_1 \ 1 \ C_2)$. With these notations, $\pi \sim \begin{pmatrix} 1 & C_2 \ 1 & B_2 \ 0 \end{pmatrix}$. This permutation is weakly reducible if and only if $C_1 = \emptyset$.

One can repeat finitely many times this process, with $B := B_2$ and $C := C_2$ for the new sets, to get the following statement. Either there exists a weakly irreducible permutation $\pi' \sim \pi$ or $B = C = (1 \ 2 \ \dots \ r)$. The last case means that $\pi = \Pi_2(r, 1)$ which is a contradiction. Therefore we have proved that there exists a weakly irreducible $\pi' \sim \pi$. Form this permutation it is easy to check that there exists $\pi'' \sim \pi'$ irreducible. Hence $\pi'' \sim \pi$ and the lemma is proven.

Now we are ready to prove Theorem 7.2 in case n = 3. Let \mathcal{C} be any component of any stratum $\mathcal{Q}_g(k_1, k_2, k_3)$. Applying Lemma 7.6 combining with the two above lemma one gets the following dichotomy. There exits $\mathcal{S} \in \mathcal{C}$ such that either \mathcal{S} possesses a multiplicity one saddle connection between two singularities (not two poles), and so the theorem is proved, or $\mathcal{S} = \mathcal{S}(\pi, \lambda)$ with $\pi = \prod_1(r, 0)$ or $\pi = \prod_2(r, 1)$. These last two permutations correspond to hyperelliptic permutations and in particular it implies that \mathcal{C} is itself a hyperelliptic component of $\mathcal{Q}(-1, -1, 4g - 2)$ which is a contradiction. The theorem is proven.

7.5. Proof of Theorem 7.2 in holomorphic case n = 2. We first prove the following lemma.

Lemma 7.7. Let $C \subset Q_g(k_1, k_2)$ be a connected component with any k_1 , $k_2 > 0$ and $g \ge 2$. Then there exists $(S(\pi, \lambda), q) \in C$ a half-translation surface such that one of the two following holds.

- $(\mathcal{S}(\pi,\lambda),q)$ has a multiplicity one saddle connection between the two zeroes.
- The combinatorics of π has the form

$$\pi = \begin{pmatrix} 0_1 & 1 & \dots & r & 0_1 & r+1 & \dots & r+l \\ 0_2 & \sigma_1(1) & \dots & \sigma_1(r) & 0_2 & \sigma_2(r+1) & \dots & \sigma_2(r+l) \end{pmatrix} \text{ or } \pi = \begin{pmatrix} A \\ B \end{pmatrix} \text{ with } \pi(A) = A, \ \pi(B) = B$$

where σ_1, σ_2 are "true" permutations of the sets $\{1, \ldots, r\}$ and $\{r+1, \ldots, r+l\}$.

Proof of Lemma 7.7. The proof is similar to the one of Lemma 7.7. It parallels the proof of Claim 1, Claim 2 and Claim 3. \Box

Therefore in order to prove Theorem 7.2 in holomorphic case n = 2 we have to analyse the two permutations of Lemma 7.7.

Lemma. Let $\pi = \begin{pmatrix} 0_1 & 1 & \dots & r & 0_1 & r+1 & \dots & r+l \\ \sigma_1(r) & 0_2 & \sigma_2(r+1) & \dots & \sigma_2(r+l) & 0_2 & \sigma_1(1) & \dots & \sigma_1(r-1) \end{pmatrix}$. Let us assume that $\pi \neq \Pi_1(r,l)$. Then there exists $\pi' \sim \pi$ such that π' is irreducible and $\gamma(\pi')$ is a saddle connection between the two zeroes.

Proof of the lemma. First let us remark that the separatrix $\gamma(\pi)$ on $\mathcal{S}(\pi, \lambda)$ is actually a saddle connection. Note also that π satisfies the Condition (*); Thus the permutation π is irreducible if and only if π is weakly irreducible.

One checks directly that π is reducible if and only if $\sigma_1(r) = 1$. In this case one can put π into the form

$$\pi = \begin{pmatrix} 0_1 & 1 & \dots & r & 0_1 & r+1 & \dots & r+l \\ \sigma_1(r) & 0_2 & \sigma_2(r+1) & \dots & \sigma_2(r+l) & 0_2 & \sigma_1(1) & \dots & \sigma_1(r-1) \end{pmatrix} \sim \pi' = \\ = \begin{pmatrix} 0_1 & 1 & 2 & \dots & r & 0_1 & r+1 & \dots & r+l \\ \sigma_1(r-1) & 1 & 0_2 & \sigma_2(r+1) & \dots & \sigma_2(r+l) & 0_2 & \sigma_1(1) & \dots & \sigma_1(r-2) \end{pmatrix}.$$

Repeating this process to π' this yields to the following dichotomy. Either there exists an irreducible permutation π' with $\pi' \sim \pi$, and so the lemma is proved, or

$$\begin{cases} \sigma_1(i) = r - i + 1 & \text{for } i = 1, \dots, r \\ \sigma_2(j) = 2r + l - j + 1 & \text{for } j = r + 1, \dots, r + l \end{cases}$$

These last equalities mean that $\pi = \Pi_1(r, l)$ which is a contradiction. The lemma is proven.

Lemma. Let $\pi = \begin{pmatrix} A \\ B \end{pmatrix}$ with $\pi(A) = A$, $\pi(B) = B$. Let us also assume that $\pi \neq \Pi_2(r,l)$. Then there exists $\pi' \sim \pi$ such that π' is irreducible and $\gamma(\pi')$ is a saddle connection between the two zeroes.

Proof. The proof is completely similar to the previous proof and left to the reader. \Box

Now we are ready to prove Theorem 7.2 in holomorphic case n = 2. Let C be any component of any stratum $Q_g(k_1, k_2)$. Applying Lemma 7.7 combining with the two above lemma one gets the following dichotomy. There exits $S \in C$ such that either S possesses a multiplicity one saddle connection between two zeroes, and so the theorem is proved, or $S = S(\pi, \lambda)$ with $\pi =$ $\Pi_1(r, l)$ or $\pi = \Pi_2(r, l)$. These last two permutations correspond to hyperelliptic permutations and in particular it implies that C is itself a hyperelliptic component of Q(4(g-k)-6, 4k+2)(where r = 2k + 1 and l = 2(g-k) - 3) which is a contradiction. The theorem is proven.

7.6. Proof of Theorem 7.2 in case $\mathcal{Q}(-1, 4g - 3)$. The proof is decomposed in two steps. We recall that a simple cylinder is a metric cylinder with boundary components consisting of two *single* homologous separatrices. A simple cylinder has multiplicity one if its boundary has multiplicity one.

Proposition 7.8. Let $C \subset Q(-1, 4g - 3)$ with $g \geq 3$. Then there exists $S(\pi, \lambda) \in C$ a half-translation surface such that one of the two followings holds.

- S has a multiplicity one saddle connection between the zero and the pole.
- S has a multiplicity one simple cylinder (i.e. $C = C' \oplus s$ with $s \in \{1, \ldots, 2g\}$ and $C' \subset Q(-1, 4(g-1) 3)$ is component).

Proposition 7.9. The following assertions hold.

(1) For any $s \in \{1, 2, 4\}$, there exists a flat surface $(S, q) \in Q(-1, 5) \oplus s$ with a multiplicity one saddle connection.

- (2) Any connected component of $\mathcal{Q}(-1, 13)$ possesses a half-translation surface equipped with a multiplicity one saddle connection.
- (3) If (S,q) has a multiplicity one saddle connection then $(S,q) \oplus s$ has also a multiplicity one saddle connection by bubbling a handle.

We address the proof of the second proposition in the appendix (see Proposition A.4). Let us first show how these two propositions give the theorem and then let us prove the first proposition.

Proof of Theorem 7.2. It is done by induction on g. The theorem is already proven in case g = 4 (Proposition 7.9, (2)). Let g > 4 be any integer and let $\mathcal{C} \subset \mathcal{Q}(-1, 4g - 3)$ be a connected component. One has to prove that \mathcal{C} possesses a surface with a multiplicity one saddle connection.

Thanks to Proposition 7.8, either \mathcal{C} possesses a surface with a multiplicity one saddle connection, and then the theorem is proved, or \mathcal{C} contains a surface with a multiplicity one simple cylinder. In the last eventuality we have $\mathcal{C} = \mathcal{C}' \oplus s$ where $\mathcal{C}' \subset \mathcal{Q}(-1, 4(g-1)-3) = \mathcal{Q}(-1, 4g'-3)$ is a connected component. We have $4 \leq g' < g$ thus by assumption \mathcal{C}' contains a surface \mathcal{S}' with a multiplicity one saddle connection. Then we conclude by Proposition 7.9, (3): $\mathcal{S}' \oplus s \in \mathcal{C}$ is the required surface and the theorem is proven.

It remains to prove the theorem for genus g = 3 case. Again form Proposition 7.8, if $\mathcal{C} \subset \mathcal{Q}(-1,9)$ has no surface with multiplicity one saddle connection then \mathcal{C} has the form $\mathcal{Q}(-1,5)\oplus s$ with s = 1, 2, 3, 4. Moreover Lemma 5.14, (4) implies $\mathcal{Q}(-1,9)^{irr} = \mathcal{Q}(-1,5)\oplus 3$. Therefore the theorem follows from Proposition 7.9, (1) which examines cases s = 1, 2, 4. The theorem is proven.

Proof of Proposition 7.8. Let $C \subset Q(-1, 4g - 3)$ be a connected component with $g \geq 3$ and $S(\pi, \lambda) \in C$. The surface S has a pole and a unique zero P. Hence one can put S into the form prescribed by Figure 11. Let α be the horizontal saddle connection between the pole




FIGURE 11. A half-translation surface with a unique zero and a unique pole.

and the zero.

Claim. Either α has multiplicity one or $\mathcal{S}(\pi, \lambda) = \mathcal{S}(\pi', \lambda')$ with $\pi' = \begin{pmatrix} 0 & 1 & \dots & r & 0 \\ A & \cdots & m & 0 \end{pmatrix}$ and $\pi'(\{1, \dots, r\}) \subset A$.

Proof of the claim. Apply Proposition 5.11.

Therefore thanks to above claim let us assume that $\mathcal{S}(\pi, \lambda) \in \mathcal{C}$ with π a permutation given by previous claim. We define $\sigma \sim \pi$ by $\sigma = \begin{pmatrix} 1 & 2 & \cdots & r & 0 & 0 \\ 1 & A & \cdots & r & 0 & 0 \end{pmatrix}$ and $\sigma(\{1, \ldots, r\}) \subset A$. Note that the vertical direction on $\mathcal{S}(\sigma, \lambda)$ decomposes the surface into (at least) one simple cylinder (see also

Figure 7). The reducibility of $\hat{\sigma}$ involves one of the two followings possible decompositions.

$$Type(1) \quad \sigma = \begin{pmatrix} 1 & 2 & \dots & k & | & k+1 & \dots & r & 0 & 0 \\ 1 & A_1 & k & A_2 & | & A' & 0 \end{pmatrix} \quad \text{with } \sigma(\{1, \dots, r\}) = A_1 \sqcup A_2$$
$$Type(2) \quad \sigma = \begin{pmatrix} 1 & 2 & \dots & r & 0 & | & 0 \\ 1 & A & | & A' & 0 \end{pmatrix} \quad \text{with } \sigma(A') \subset A$$

Claim. If $\hat{\sigma}$ involves reducibility of Type (1) then there exists $\sigma_1 \sim \sigma$ such that $\hat{\sigma}_1$ is irreducible.

Proof of the claim. The proof parallels the one of Theorem 6.3 versus Abelian differentials (Section 6.1). \Box

For reducible permutations σ of Type (2), let us denote $A = (A_1 \ k \ A_2)$ with $\sigma(A_2) \subseteq A'$ and $2 \leq k \leq r$ (eventually $A_2 = \emptyset$).

Claim. If $\hat{\sigma}$ involves reducibility of Type (2) then either there exists $\sigma_1 \sim \sigma$ such that $\hat{\sigma}_1$ is irreducible or k = r.

Proof of the claim. With above notations, if $k \neq r$ then one has

$$\sigma = \begin{pmatrix} 1 & 2 & \dots & k & \dots & r & 0 & 0 \\ 1 & A_1 & k & A_2 & & A' \end{pmatrix} \sim \sigma_1 = \begin{pmatrix} k & k+1 & \dots & r & 0 & 0 & 1 & 2 & \dots & k-1 \\ k & A_2 & A' & 1 & A_1 & & & \end{pmatrix}.$$

If $\hat{\sigma}_1$ is irreducible then the claim is proven. Otherwise it is easy to see that $\hat{\sigma}_1$ involves reducibility of Type (1). Thus previous claim applies.

Finally the last case to consider is the following one: $\hat{\sigma}$ is reducible with decomposition of Type (2) and k = r. Reporting these data into the permutation π one gets $\pi = \begin{pmatrix} 0 & 1 & \dots & r & 0 \\ r & A_2 & A' & 1 & A_1 \end{pmatrix}$ with $\pi(A_2) \subseteq A'$.

Claim. The vertical separatrix $\gamma(\pi)$ on $S(\pi, \lambda)$ is a saddle connection. Moreover the above permutation π is reducible if and only if

$$\pi = \begin{pmatrix} 0 & 1 & \dots & r & 0 \\ r & A' & 1 & \pi(2) & \dots & \pi(r-1) & A'' \end{pmatrix}$$

with $A'' = \pi(A')$.

Proof of the claim. Let $\pi = \begin{pmatrix} 0 & 1 & \dots & r & 0 \\ r & A_2 & A' & 1 & A_1 \end{pmatrix}$ with $\pi(A_2) \subseteq A'$. A direct observation shows that if π is reducible then $A_2 = \emptyset$. The same approach shows that if π is reducible then $A_1 = \pi(2) \dots \pi(r-1) A''$ with $\pi(A') = A''$. The claim is proven.

The conclusion of previous claims is the following one. We have proved that C contains a surface $S(\pi, \lambda)$ such that either S satisfied the conclusions of the proposition or

$$\pi = \begin{pmatrix} 0 & 1 & \dots & r & 0 \\ r & A' & 1 & \pi(2) & \dots & \pi(r-1) & A'' \end{pmatrix} \quad \text{and} \ \pi(A') = A''$$

Then we have to study the two last cases $r \ge 2$ and r = 1. This corresponds to the two next lemma which end the proof of Proposition 7.8.

Lemma. Let $r \ge 2$ be any integer. Let $\pi = \begin{pmatrix} 0 & 1 & \dots & r & 0 \\ r & A' & 1 & \pi(2) & \dots & \pi(r-1) & A'' \end{pmatrix}$ be a permutation with $\pi(A') = A''$. We also assume that $S(\pi, \lambda) \in \mathcal{Q}(-1, 4g - 3)$ with $g \ge 3$.

Then there exists a permutation $\pi' \sim \pi$ such that $S(\pi', \lambda)$ has a multiplicity one saddle connection.

Proof of the lemma. Let us assume that $r \ge 3$, then $r-1 \ne 1$. Let us introduce some notations to clarify the situation. Let C stand for $(\pi(2) \ldots \pi(r-1))$. One decomposes C into the following way: $C = (C_1 \ r - 1 \ C_2)$. Equipped with these notations: $\pi = \begin{pmatrix} 0 \ 1 \ \cdots \ r-1 \ r \ 0 \\ r \ A' \ 1 \ C_1 \ r-1 \ C_2 \ A'' \end{pmatrix} \sim$ $\sigma = \begin{pmatrix} 0 \ 1 \ \cdots \ r-1 \ r \ 0 \\ r-1 \ C_2 \ A'' \ r \ A' \ 1 \ C_1 \end{pmatrix}$. A direct observation shows that σ is irreducible and $\gamma(sigma)$ is a saddle connection. The lemma then follows from Theorem 5.12.

Now let us assume that r = 2. The list A'' is non-empty. Let us denote $A'' = (3 \ B)$ with $\pi(3) \in A'$. We also use $A' = (B_1 \ 3 \ B_2)$. Thus $\pi = \begin{pmatrix} 0 & 1 & 2 & 0 \\ 2 & A' & 1 & A'' \end{pmatrix} \sim \sigma = \begin{pmatrix} 0 & 1 & 2 & 0 \\ B & 2 & B_1 & 3 & B_2 & 1 & 3 \end{pmatrix}$. This permutation is reducible if and only if $B = B_1 = B_2 = \emptyset$. This case consists of $\pi = \begin{pmatrix} 0 & 1 & 2 & 0 \\ B & 2 & B_1 & 3 & B_2 & 1 & 3 \end{pmatrix}$ and then $S(\pi, \lambda) \in Q(-1, 5)$ contradicting $g \geq 3$. Hence σ is irreducible and the lemma again follows from Theorem 5.12.

Next lemma studies r = 1 case.

Lemma. Let $\pi = \begin{pmatrix} 0 & 1 & 0 \\ A' & 1 & A'' \end{pmatrix}$ be a permutation with $\pi(A') = A''$. Then one of the following assertions holds.

- There exists $\pi' \sim \pi$ such that $\mathcal{S}(\pi', \lambda)$ has a multiplicity one saddle connection.
- The combinatorics of π is given by $A' = A'' = (2 \ 3 \ \dots \ l)$. Moreover there exists $\pi' \sim \pi$ and an admissible vector λ_0 such that $S(\pi', \lambda_0)$ has a multiplicity one simple cylinder.

Proof of the lemma. Let π the permutation $\pi = \begin{pmatrix} 0 & 1 & 0 \\ A' & 1 & A'' \end{pmatrix}$. The list A' is non-empty thus let us introduce $A' = (2 \ A'_2)$ with $\pi(2) \in A''$. Hence we can also introduce lists A''_1 , A''_2 such that $A'' = (A''_1 \ 2 \ A''_2)$. According to these notations, one has $\pi = \begin{pmatrix} 0 & 1 & 0 \\ 2 & A'_2 \ 1 & A''_1 \ 2 & A''_2 \end{pmatrix} \sim \pi' = \begin{pmatrix} 0 & 1 & 0 \\ 2 & A''_2 \ 1 & A''_1 \ 2 & A''_2 \end{pmatrix}$. A straightforward computation shows that the permutation π' is reducible if and only if $A''_1 = \emptyset$. Thus, repeating inductively this process with $A' := A'_2$ and $A'' := A''_2$, we get either $S(\pi', \lambda)$ has a multiplicity one saddle connection or

$$\pi \sim \pi' = \begin{pmatrix} 0 & 1 & 0 \\ 3 & 4 & \dots & l & 2 & 3 & 4 & \dots & l & 1 & 2 \end{pmatrix}$$

In order to finish the proof of the lemma, we have to present an admissible vector λ_0 such that the vertical foliation on $S(\pi', \lambda_0)$ is completely periodic and decomposes the surface with at least a multiplicity one simple cylinder. Let us consider the admissible vector (for π'):

(7.1)
$$\lambda_0 = \left((l-1)\alpha \; ; \; \alpha \; ; \; (l-1)\alpha \; ; \; \underline{\alpha \; ; \; \dots \; ; \; \alpha}_{2l-1 \; \text{times}} \right) \quad \text{for any } \alpha \in \mathbb{R}^+$$

It is easy to see that the vertical foliation on $S(\pi', \lambda_0)$ decomposes the surface into g - 1 cylinders (see Figure 12). One checks that the vertical cylinder corresponding to the horizontal interval numbered l is a multiplicity one simple cylinder (see the figure). In Figure 12, we present a complete description for the surface given by the case r = 5 (i.e. g = 3). This completes the proof of the lemma.

Now we can deduce from Theorem 7.2 our main result, namely Theorem 7.1.

7.7. Proof of Theorem 7.1. We first establish




FIGURE 12. A half-translation surface $S' = S(\pi'\lambda_0)$ suspended over the permutation π' and the admissible vector λ_0 (see Equation (7.1). The vertical foliation decomposes S' in two cylinders C_1 and C_2 . The cylinder C_2 is a simple cylinder: the conical angle of its boundary η is π . Moreover the length of η can be chosen arbitrary small with respect to other vertical parameters: C_2 has multiplicity one.

Proposition 7.10 (Reformulation of Theorem 7.2 in holomorphic case n = 2).

 $C \subseteq Q(k_1, k_2), \ k_i > 0, \ is \ hyperelliptic \iff$ $\forall \ (S, q) \in C, \ any \ saddle \ connection \ on \ S \ has \ multiplicity \ at \ least \ 2$

Through the proof of Theorem 7.2 in case n = 3 and n = 4 one gets a similar characterization of hyperelliptic components (compare with Lemma 2.4).

Proposition 7.11. Hyperelliptic components of strata with 3 and 4 singularities are characterized by the following:

(1) $C \subseteq Q(k_1, k_2, k_3)$ is hyperelliptic $\iff \exists i_0 \neq i_1 \in \{1, 2, 3\}, \forall (S, q) \in C$, any saddle connection on S between P_{k_i} and P_{k_j} , with $i \in \{i_0, i_1\}$ and $j \notin \{i_0, i_1\}$ has multiplicity at least 2. (2) $C \subseteq Q(k_1, k_2, k_3, k_4)$ is hyperelliptic $\iff \exists i_0 \neq i_1 \in \{1, 2, 3, 4\}$ $\forall (S,q) \in C$, any saddle connection on S between P_{k_i} and P_{k_j} , with $i \in \{i_0, i_1\}$ and $j \notin \{i_0, i_1\}$, has multiplicity at least 2.

Proposition 7.12. Let C be a connected component of a $Q(k_1, \ldots, k_n)$ with $n \ge 5$. Then for each $i \ne j$, there exists a half-translation surface $S \in C$ with a multiplicity one saddle connection between two singularities $P_{k_i}, P_{k_j} \in S$ of multiplicities k_i and k_j .

Proof of Theorem 7.1. We discuss the theorem following the different values of n (the number of singularities). First let us note that if we prove the theorem in case n = 2, 3, 4, 5 then the theorem follows for any $n \ge 6$ because the locus of hyperelliptic and irreducible components is located on strata with 2, 3 and 4 singularities.

The theorem in case n = 2 corresponds to Theorem 7.2 which is already proved.

Thus let us assume that n = 3. Let $\mathcal{C} \subseteq \mathcal{Q}(k_1, k_2, k_3)$ be a non-hyperelliptic component. Recall that all k_i are non-zero. Up to re-organize k_i , one can assume $k_1 \leq k_2 \leq k_3$. Proposition 7.11 with $i_0 = 1$, $i_1 = 2$ gives a half-translation surface $(\mathcal{S}, q) \in \mathcal{C}$ with a multiplicity one saddle connection between $P_{k_3} \in \mathcal{S}$ and $P_{k_i} \in \mathcal{S}$ for $i \in \{1, 2\}$. In other terms there exists a component \mathcal{C}' of the strata $\mathcal{Q}(k_1 + k_3, k_2) \sqcup \mathcal{Q}(k_2 + k_3, k_1)$ such that $\mathcal{C}' \subset \overline{\mathcal{C}}$. There is three possibilities: \mathcal{C}' is hyperelliptic, irreducible or "regular" (neither hyperelliptic or irreducible). If \mathcal{C}' is regular then Theorem 7.2 leads to the result. Indeed on can find $\mathcal{C}_0 \subseteq \mathcal{Q}(4g - 4)$ a component with $\mathcal{C}_0 \subset \overline{\mathcal{C}'}$. But $\overline{\mathcal{C}'} \subset \overline{\mathcal{C}}$ therefore $\mathcal{C}_0 \subset \overline{\mathcal{C}}$ and we are done.

Assume that \mathcal{C}' is hyperelliptic. Applying Proposition B.1, one can connect the hyperelliptic component \mathcal{C}' to another non-hyperelliptic component \mathcal{C}'' of $\mathcal{Q}(k_i + k_3, k_j)$ passing through the stratum $\mathcal{Q}(k_1, k_2, k_3)$. Therefore there exists a component $\mathcal{C}_2 \subseteq \mathcal{Q}(k_1, k_2, k_3)$ such that $\mathcal{C}' \sqcup \mathcal{C}'' \subset \overline{\mathcal{C}}_2$. Thus $\mathcal{C}' \subset \overline{\mathcal{C}} \sqcup \overline{\mathcal{C}}_2$. The assumption $k_1 \leq k_2 \leq k_3$ implies $k_i + k_3 \neq k_j$ thus Corollary 2.7 implies $\mathcal{C}_2 = \mathcal{C}$. Therefore \mathcal{C} is adjacent to a non-hyperelliptic component of $\mathcal{Q}(k_i + k_3, k_j)$ and we are again done.

Finally let us assume that C' is irreducible; that is $C' = Q^{irr}(-1,9)$. Recall that the component $C \neq Q^{irr}(-1,3,6)$ therefore the theorem follows from Proposition B.2 and Corollary 2.7.

The proofs for cases n = 4, 5 are similar to the above discussion. Theorem 7.1 is proven. \Box

8. Proof of Main Theorem 1.1 and Main Theorem 1.2

Proof of Main Theorem 1.1. As mentioned in Section 7.1, Main Theorem 1.1 follows from Theorem 7.1 on adjacency of strata and Theorem 6.1 on classification of minimal strata. \Box

Proof of Main Theorem 1.2. Theorem 1.2 is already proved for genus 0 case (see Proposition 1.5). The genus 3 case is also proved (see Theorem 7.1 and Section 7.1).

Thus let us consider genera 1, 2 cases. One can first prove, by a direct argument on finitely many strata, Theorem 1.2 for g = 1, 2 and $n \leq 5$ cases. Indeed the number of such a strata of Q_2 and of Q_1 with n = 2, 3, 4, 5 is finite. We can check, using Rauzy classes, that the theorem holds. In particular we get that any stratum of Q_2 with n = 5 is connected and any stratum of Q_1 with n = 5 is also connected.

Now let us prove that any stratum of \mathcal{Q}_1 with $n \ge 6$ is also connected. Let $\mathcal{C}_1, \mathcal{C}_2$ be two components of $\mathcal{Q}(k_1, \ldots, k_6)$. Let us assume that $k_1 \le \cdots \le k_6$. Then $k_5 + k_6 \ne k_i$ for any $i = 1, \ldots, 4$. By Proposition 7.12 there exists two components \mathcal{C}_0 and \mathcal{C}'_0 of $\mathcal{Q}(k_1, \ldots, k_4, k_5 + k_6)$ with $\mathcal{C}_0 \subset \overline{\mathcal{C}_1}$ and $\mathcal{C}'_0 \subset \overline{\mathcal{C}_2}$. Any stratum of \mathcal{Q}_1 with n = 5 being connected, one has $\mathcal{C}_0 = \mathcal{C}'_0$

hence $C_0 \subset \overline{C_1} \cup \overline{C_2}$. Therefore Corollary 2.7 applies and $C_1 = C_2$; so that any stratum of Q_1 with n = 6 is connected. Above argument extends to any stratum of Q_1 with $n \ge 6$. Theorem 1.2 in case g = 1 is therefore proven. The same argument holds for the genus 2 case.

We now prove Theorem 1.2 in case g = 4. We have already proved that any component of a stratum of \mathcal{Q}_4 which is not irreducible neither hyperelliptic is adjacent to $\mathcal{Q}^{irr,I}(12)$ or $\mathcal{Q}^{irr,II}(12)$ (Theorem 7.1). In order to prove the theorem it suffices to show that those components are in fact adjacent to $\mathcal{Q}^{irr,I}(12)$. By Theorem 7.2, any non-irreducible or nonhyperelliptic component is adjacent to a non-irreducible or non-hyperelliptic component of a stratum of \mathcal{Q}_4 with n = 2. Let \mathcal{C} be any non-irreducible or non-hyperelliptic component of $\mathcal{Q}(k_1, k_2)$ with $k_1 + k_2 = 12$. One has to prove that \mathcal{C} is adjacent to $\mathcal{Q}^{irr,I}(12)$. By Theorem 7.1, \mathcal{C} is adjacent to $\mathcal{Q}^{irr,I}(12)$ or $\mathcal{Q}^{irr,II}(12)$. Let us assume that second case holds. By Proposition B.3 there exists $\mathcal{C}' \subset \mathcal{Q}(k_1, k_2)$ such that $\mathcal{Q}^{irr,II}(12) \subset \overline{\mathcal{C}'}$ and $\mathcal{Q}^{irr,I}(12) \subset \overline{\mathcal{C}'}$. Therefore $\mathcal{Q}^{irr,II}(12) \subset \overline{\mathcal{C}'} \cup \overline{\mathcal{C}}$ and Corollary 2.7 implies $\mathcal{C} = \mathcal{C}'$. Theorem 1.2 is proven. \Box

Appendix A. Connectedness of particular strata

This section is devoted to the computation of the set of connected components of some strata in low dimensional case.

Lemma A.1. The stratum Q(8) is connected.

Proof. Let $(S(\pi, \lambda), q)$ be a genus three half-translation surface, with a single singularity. The conical angle around this singularity is 10π . We will show that the possibilities for the combinatorics of the gluing maps of the set of horizontal separatrix loops is very restrictive.

Recall that to each permutation π one associates two lines of the table representing π (see 3). There are obviously two possibilities for these two lines: either the numbers of elements are equal or they are different. One defines \mathcal{A}_1 the set of generalized permutations corresponding to the first case. We also define \mathcal{A}_2 the set of generalized permutations corresponding to the second case.

By assumption $(S,q) \in \mathcal{Q}(8)$, hence there are exactly five separatrices loops. Therefore the number of elements of lines for permutations in \mathcal{A}_1 is (5,5) and for permutations in \mathcal{A}_2 is (6,4). A straightforward computation shows the two sets \mathcal{A}_1 , \mathcal{A}_2 are very simple. More precisely, up to cyclic order, one has $\#\mathcal{A}_1 = 4$ and $\#\mathcal{A}_2 = 3$.

up to cyclic order:
$$\mathcal{A}_1 = \left\{ \begin{pmatrix} 5 & 3 & 5 & 2 & 4 \\ 1 & 2 & 1 & 3 & 4 \end{pmatrix}, \begin{pmatrix} 5 & 4 & 5 & 2 & 3 \\ 1 & 2 & 1 & 3 & 4 \end{pmatrix}, \begin{pmatrix} 5 & 4 & 5 & 3 & 2 \\ 1 & 2 & 1 & 3 & 4 \end{pmatrix}, \begin{pmatrix} 5 & 3 & 5 & 3 & 4 \\ 1 & 2 & 1 & 3 & 4 \end{pmatrix} \right\}$$

up to cyclic order: $\mathcal{A}_2 = \left\{ \begin{pmatrix} 5 & 2 & 5 & 3 & 4 & 2 \\ 1 & 3 & 1 & 4 & 2 \end{pmatrix}, \begin{pmatrix} 3 & 5 & 4 & 2 & 3 & 4 & 2 \\ 1 & 3 & 1 & 4 & 2 & 2 \end{pmatrix} \right\}$

This proves that the stratum $\mathcal{Q}(8)$ has at most 7 connected components. Now, let us consider surfaces $\mathcal{S}(\pi, \lambda_2)$ with $\pi \in \mathcal{A}_2$ and $\lambda_2 = (1, 1, 1, 1, 1, 1, 2, 1, 2, 1)$. A direct verification shows that the vertical foliation on surfaces $\mathcal{S} = \mathcal{S}(\pi, \lambda_2)$ decomposes \mathcal{S} into a single cylinder. Therefore, we get a permutation encoding this cylinder. One can check that this permutation belongs to the set \mathcal{A}_1 . Thus procedure connects permutations of \mathcal{A}_2 to permutations of \mathcal{A}_1 .

Lemma A.2. The stratum Q(12) has at most two connected components.

Proof. One has to show that $\mathcal{Q}(12) = \mathcal{Q}^{irr,I}(12) \cup \mathcal{Q}^{irr,II}(12)$. Thanks to previous lemma, let \mathcal{C}_0 be the unique connected component of the stratum $\mathcal{Q}(8)$. Theorem 6.3 implies that any component \mathcal{C} of the stratum $\mathcal{Q}(12)$ has the following form.

(A.1)
$$\mathcal{C} = \mathcal{C}_0 \oplus s \text{ with } s = 1, \dots, 6 = 2g$$

We now recall the construction of the two irreducible components $\mathcal{Q}^{irr,I}(12)$, $\mathcal{Q}^{irr,II}(12)$. Let $\pi_1 = \begin{pmatrix} 1 & 2 & 3 & 4 & 2 & 5 & 6 \\ 1 & 4 & 5 & 7 & 6 & 7 & 3 \end{pmatrix}$ and $\pi_2 = \begin{pmatrix} 1 & 2 & 3 & 4 & 3 & 5 & 6 \\ 1 & 5 & 7 & 4 & 2 & 6 & 7 \end{pmatrix}$ be two permutations. Let $\mathcal{S}_i = \mathcal{S}(\pi_i, \lambda_0)$ be the suspended flat surfaces with admissible vector $\lambda_0 = (1, 1, 1, 1, 1, 1, 1)$. By definition $\mathcal{Q}^{irr,I}(12)$ is the component containing \mathcal{S}_1 and $\mathcal{Q}^{irr,II}(12)$ is the component containing \mathcal{S}_2 . Let us recall (see Lemma 5.14) that the vertical foliation on \mathcal{S}_i produces a multiplicity one simple cylinder which gives:

$$\mathcal{Q}^{irr,I}(12) = \mathcal{C}_0 \oplus 2$$
$$\mathcal{Q}^{irr,II}(12) = \mathcal{C}_0 \oplus 6$$

We will show:

$$\mathcal{C}_0 \oplus 4 = \mathcal{C}_0 \oplus 1 = \mathcal{C}_0 \oplus 5 = \mathcal{C}_0 \oplus 2 = \mathcal{Q}^{irr,I}(12)$$

$$\mathcal{C}_0 \oplus 3 = \mathcal{C}_0 \oplus 6 = \mathcal{Q}^{irr,II}(12)$$

Combining with Equation (A.1) this will give the lemma.

Let $C_0 \oplus s$, $s = 1, \ldots, 6$ be any component of Q(12).

case s=3.

Let us consider the new permutation $\pi_2 \sim \pi'_2 = \begin{pmatrix} 5 & 6 & 1 & 2 & 3 & 4 & 3 \\ 5 & 7 & 4 & 2 & 6 & 7 & 1 \end{pmatrix}$. Obviously, $\hat{\pi}'_2$ is irreducible thus one gets a (vertical) multiplicity one simple cylinder on $\mathcal{S}(\pi'_2, \lambda_0)$. This cylinder has angle 3π . In other words $\mathcal{C}_0 \oplus 3 = \mathcal{Q}^{irr,II}(12)$.

case s=4.

Let us consider the new permutation $\pi_1 \sim \pi'_1 = \left(\frac{5}{5} \frac{6}{7} \frac{1}{6} \frac{2}{7} \frac{3}{3} \frac{4}{1} \frac{2}{4}\right)$. Obviously, $\hat{\pi}'_1$ is irreducible thus one gets a (vertical) multiplicity one simple cylinder on $\mathcal{S}(\pi'_1, \lambda_0)$. This cylinder has angle 4π or in other words $\mathcal{C}_0 \oplus 4 = \mathcal{Q}^{irr, I}(12)$.

cases s=1 and s=5.

Now let us consider the new permutation $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 5 \\ 1 & 4 & 7 & 3 & 7 & 2 & 6 \end{pmatrix}$. The permutation $\hat{\sigma}$ is irreducible therefore the surface $S(\sigma, \lambda_0)$ has a (vertical) multiplicity one simple cylinder. This cylinder has angle 4π , therefore this surface $S(\sigma, \lambda_0)$ belongs to component $Q^{irr,I}(12)$.

Let us consider the new permutation $\sigma \sim \sigma' = \begin{pmatrix} 3 & 4 & 5 & 6 & 5 & 1 & 2 \\ 3 & 7 & 2 & 6 & 1 & 4 & 7 \end{pmatrix}$. Obviously, $\hat{\sigma}'$ is irreducible thus one gets a (vertical) multiplicity one simple cylinder on $\mathcal{S}(\sigma', \lambda_0)$. This cylinder has angle π . In other words $\mathcal{C}_0 \oplus 1 = \mathcal{Q}^{irr, I}(12)$.

Let us consider the new permutation $\sigma \sim \sigma'' = \begin{pmatrix} 2 & 3 & 4 & 5 & 6 & 5 & 1 \\ 2 & 6 & 1 & 4 & 7 & 3 & 7 \end{pmatrix}$. Obviously, $\hat{\sigma}''$ is irreducible thus one gets a (vertical) multiplicity one simple cylinder on $\mathcal{S}(\sigma'', \lambda_0)$. This cylinder has angle 5π or in other words $\mathcal{C}_0 \oplus 5 = \mathcal{Q}^{irr, I}(12)$.

The lemma is proven.

Lemma A.3. The stratum $\mathcal{Q}(-1,5)$ is connected.

Proof of Lemma A.3. Let $S(\pi, \lambda) \in Q(-1, 5)$ be any surface. One can directly check that (up to cyclic order), the combinatorics of the permutation π has only two possibilities.

$$\pi_1 = \begin{pmatrix} 0 & 0 & 1 & 2 \\ 1 & 3 & 2 & 3 \end{pmatrix} \quad \text{or} \quad \pi_2 = \begin{pmatrix} 0 & 1 & 0 \\ 2 & 3 & 2 & 1 & 3 \end{pmatrix}$$

Equipped with the two admissible vectors

$$\lambda_1 = (1, 1, 2, 1, 2, 1, 1, 1)$$
 and $\lambda_2 = (2, 1, 2, 1, 1, 1, 1, 1),$

we obtain two flat surfaces $S_i = S(\pi_i, \lambda_i)$, i = 1, 2. A direct computation shows the vertical foliation on S_2 produces the surface S_1 . The lemma is proven.

Proposition A.4. The following assertions hold:

- (1) For any $s \in \{1, 2, 4\}$, there exists a flat surface $(S, q) \in Q(-1, 5) \oplus s$ with a multiplicity one saddle connection.
- (2) Any connected component of $\mathcal{Q}(-1, 13)$ has a half-translation surface with a multiplicity one saddle connection.
- (3) If (S,q) has a multiplicity one saddle connection then $(S,q) \oplus s$ has also a multiplicity one saddle connection by bubbling a handle.

Proof of Proposition A.4. We consider separately the three cases.

Proof of the first point We will present three surfaces in $\mathcal{Q}(-1,5) \oplus s$ (for s = 1, 2, 4) with a multiplicity one saddle connection. For that we will construct surfaces in $\mathcal{Q}(-1,9)$ with a multiplicity one simple cylinder of angle $\pi, 2\pi$ and 4π .

Thus let $\pi_1 = \begin{pmatrix} 3 & 4 & 0 & 0 & 1 & 2 \\ 3 & 5 & 2 & 1 & 4 & 5 \end{pmatrix}$ and $\pi_2 = \begin{pmatrix} 2 & 3 & 4 & 0 & 0 & 1 \\ 2 & 4 & 5 & 1 & 3 & 5 \end{pmatrix}$ be two permutations. The corresponding surfaces belong to $\mathcal{Q}(-1,9)$. The surfaces $\mathcal{S}_1 = \mathcal{S}(\pi_1,\lambda)$ and $\mathcal{S}_2 = \mathcal{S}(\pi_2,\lambda)$ possess a (vertical) multiplicity one simple cylinder. One can check that $\hat{\pi}_i$ is irreducible for i = 1, 2 thus, by Theorem 5.13, each of the surfaces \mathcal{S}_1 and \mathcal{S}_2 possesses a multiplicity one simple cylinder. Thanks to a direct computation this cylinder has angle π for i = 1 and 2π for i = 2. In other terms this proves $\mathcal{S}_1 \in \mathcal{Q}(-1, 5) \oplus 1$ and $\mathcal{S}_2 \in \mathcal{Q}(-1, 5) \oplus 2$.

Let us show that $\mathcal{Q}(-1,5) \oplus 4 = \mathcal{Q}(-1,5) \oplus 1$. One has $\pi_1 \sim \pi_3 = \begin{pmatrix} 1 & 2 & 3 & 4 & 0 & 0 \\ 1 & 4 & 5 & 3 & 5 & 2 \end{pmatrix}$. The permutation $\hat{\pi}_3$ is irreducible and the angle of the cylinder on $\mathcal{S}(\pi_3, \lambda)$ is 4π . Then surface $\mathcal{S}(\pi_3, \lambda)$ belongs to $\mathcal{Q}(-1,5) \oplus 4$ and $\mathcal{Q}(-1,5) \oplus 1$.

It remains to end the proof to find a multiplicity one saddle connection on each surface S_1 , S_2 . This is done with the following remark. Let us consider the permutations

$$\pi_1 = \begin{pmatrix} 3 & 4 & 0 & 0 & 1 & 2 \\ 3 & 5 & 2 & 1 & 4 & 5 \end{pmatrix} \sim \pi'_1 = \begin{pmatrix} 0 & 1 & 2 & 3 & 4 & 0 \\ 1 & 4 & 5 & 3 & 5 & 2 \end{pmatrix} \quad \text{and} \quad \pi_2 = \begin{pmatrix} 2 & 3 & 4 & 0 & 0 & 1 \\ 2 & 4 & 5 & 1 & 3 & 5 \end{pmatrix} \sim \pi'_2 = \begin{pmatrix} 0 & 1 & 2 & 3 & 4 & 0 \\ 4 & 5 & 1 & 3 & 5 & 2 \end{pmatrix}.$$

For i = 1, 2 the vertical separatrix loop $\gamma(\pi'_i)$ is a saddle connection and π'_i is irreducible. Therefore Proposition 5.10 applies. Hence first point of the proposition is proven.

Proof of the second point

One has to prove that each component of $\mathcal{Q}(-1,13)$ has a half-translation surface with a multiplicity one saddle connection. Thanks to Proposition 7.8 we are reduced to consider components of the form $\mathcal{C} \oplus s$ with $\mathcal{C} \subset \mathcal{Q}(-1,9)$. The previous proof shows that each component of $\mathcal{Q}(-1,9)$ (different form $\mathcal{Q}^{irr}(-1,9)$) possesses a surface with a multiplicity one saddle connection. Finally one has to consider components of the form

$$\mathcal{Q}^{irr}(-1,9) \oplus s$$
 with $s = 1, \dots, 6$

Let us recall that $\mathcal{Q}^{irr}(-1,9) = \mathcal{Q}(-1,5) \oplus 3$. Moreover any component $\mathcal{Q}(-1,5) \oplus s$, with s = 1, 2, 4, 5, possesses a flat surface with a multiplicity one saddle connection. Using properties of the map \oplus (see Proposition 2.9), this yields to $\mathcal{Q}(-1,9)^{irr} \oplus s = \mathcal{Q}(-1,5) \oplus s \oplus 3$ which proves the proposition for s = 1, 2, 4, 5.

The case s = 6 is reduced to the case s = 3.

$$\mathcal{Q}(-1,9)^{irr} \oplus 6 = \mathcal{Q}(-1,5) \oplus 6 \oplus 3 = \mathcal{Q}(-1,9)^{irr} \oplus 3.$$

We finish the proof of the second statement using the second property of the map \oplus .

$$\mathcal{Q}(-1,5) \oplus 3 \oplus 3 = \mathcal{Q}(-1,5) \oplus 1 \oplus 5$$

Proof of the third point

The proof is obvious.

We finish this section with an independent proof of a theorem of Masur and Smillie [MaSm]. The original proof uses algebraic geometry. Here we only use combinatorics of generalized permutations.

Theorem (Masur, Smillie). The following strata

$$\mathcal{Q}(\emptyset), \ \mathcal{Q}(1,-1) \ (in \ genus \ g=1) \quad and \quad \mathcal{Q}(4), \ \mathcal{Q}(1,3) \ (in \ genus \ g=2)$$

are empty.

Proof of the theorem. Let us assume that the stratum $\mathcal{Q}(4)$ is non-empty. Thus there exists a genus two half-translation surface with a single zero. Applying "breaking up a singularity" to this zero into two zeroes of order tow we get a point inside the stratum $\mathcal{Q}(2,2)$ (see [La1] or [MaZo], [EsMaZo]). By construction, this new surface has a multiplicity one saddle connection so it belongs to a non-hyperelliptic component of $\mathcal{Q}(2,2)$. Now we will prove that this stratum is connected and equals to its hyperelliptic component which leads to a contradiction.

Let us consider a point $\mathcal{S}(\pi,\lambda) \in \mathcal{Q}(2,2)$. As usual, a direction computation shows that one can put (up to cyclic order) π into one of the two following forms $\begin{pmatrix} 1 & 2 & 1 & 3 \\ 4 & 3 & 4 & 2 \end{pmatrix}$ or $\begin{pmatrix} 1 & 2 & 1 & 2 \\ 3 & 4 & 3 & 4 \end{pmatrix}$. According to Lemma 4.2, each of these two permutations gives rise to surfaces into the component $\mathcal{Q}^{hyp}(2,2)$. Therefore $\mathcal{Q}(2,2) = \mathcal{Q}^{hyp}(2,2)$ is connected and hence the stratum $\mathcal{Q}(4) = \emptyset.$

Using the same approach, we prove that $\mathcal{Q}(1,-1)$ and $\mathcal{Q}(1,3)$ are empty (one considers respectively the connected hyperelliptic strata $\mathcal{Q}(-1, -1, 2)$ and $\mathcal{Q}(1, 1, 2)$).

The stratum $\mathcal{Q}(\emptyset)$ is empty because the quotient of a quadratic differential q by dz^2 is a holomorphic function on the torus thus constant. Therefore $q = \omega^2$ which is a contradiction.

Appendix B. Deformations of hyperelliptic and irreducible components

Proposition B.1. Let $(S,q) \in \mathcal{Q}^{hyp}(4(g-k)-6,4k+2)$ be a point with $0 \ge k \ge g-2$ and $g \ge 2$. Let (k_1, k_2) be a positive partition of 4k + 2. Then there exists a continuous path $\rho: [0,1] \longrightarrow \mathcal{Q}_g$ of the interval [0,1] into the whole moduli space \mathcal{Q}_g such that:

- $\rho(0) = (\mathcal{S}, q)$
- $\rho(t) \in \mathcal{Q}(4(g-k) 6, k_1, k_2) \quad \forall \ 0 < t < 1.$ $\rho(1) \in \mathcal{Q}(4(g-k) 6, 4k + 2) \setminus \mathcal{Q}^{hyp}(4(g-k) 6, 4k + 2).$

Proof. We remark that it is sufficient to prove the proposition for a particular point of the component $\mathcal{Q}^{hyp}(4(g-k)-6,4k+2)$. We first claim:

Claim. Let us fix r = 2k + 1 and l = 2(g - k) - 3. Let a be any integer with $2 \le a \le r + 1$. Let us consider the generalized permutation

$$\Pi_1(r,l,a) = \begin{pmatrix} 0_1 & 0_3 & 1 & \dots & r & 0_1 & r+1 & \dots & r+l \\ r+l & \dots & r+1 & 0_2 & r & \dots & a & 0_3 & a-1 & \dots & 1 & 0_2 \end{pmatrix}.$$

Then for any admissible vector λ one has $\mathcal{S}(\Pi_1(r,l,a),\lambda) \in \mathcal{Q}(a-2, 4k+4-a, 4(g-k)-3)$. Moreover, the horizontal saddle connection labeled 0_3 has multiplicity one. The resulting surface \hat{S} obtained by shrinking this saddle connection to a point is $\mathcal{S}(\Pi_1(r,l),\lambda)$ which belongs to the hyperelliptic component.

Proof of the claim. A straightforward calculation of the angle of the conical singularities located at end-points of the intervals.

The proposition follows from above claim taking $a = k_1 + 2$.

One can easily have similar results on deformations of hyperelliptic components of other strata. Here we present a similar result concerning irreducible component. The proof is just based on the deformations of an adequate generalized permutation.

Proposition B.2. Let $(S,q) \in Q^{irr}(-1,9)$ be a point. Let (k_1,k_2) be any pairs in the list $\{(-1, 10), (1, 8), (2, 7), (3, 6), (4, 5)\}$. Then there exists a continuous path $\rho : [0, 1] \longrightarrow Q_3$ of the interval [0,1] into the whole moduli space Q_3 such that:

•
$$\rho(0) = (\mathcal{S},$$

- $\begin{array}{l} \bullet \ \rho(0) = (\mathcal{S},q) \\ \bullet \ \rho(t) \in \mathcal{Q}(-1,k_1,k_2) \quad \forall \ 0 < t < 1. \\ \bullet \ \rho(1) \in \mathcal{Q}(-1,9) \ \backslash \ \mathcal{Q}^{irr}(-1,9). \end{array}$

Proof. The proof is similar to the previous one, deforming the permutation

representing the irreducible component of $\mathcal{Q}(-1,9)$.

Proposition B.3. Let $(S,q) \in Q^{irr,II}(12)$ be a point. Let (k_1,k_2) be any pair in the list $\{(-1, 13), (1, 11), (2, 10), (3, 9), (4, 8), (5, 7), (6, 6)\}.$

Then there exists a continuous path $\rho: [0,1] \longrightarrow \mathcal{Q}_4$ of the interval [0,1] into the whole moduli space Q_4 such that:

$$\begin{aligned} \bullet \ \rho(0) &= (\mathcal{S}, q) \\ \bullet \ \rho(t) \in \mathcal{Q}(k_1, k_2) \qquad \forall \ 0 < t < 1. \\ \bullet \ \rho(1) \in \mathcal{Q}^{irr, I}(12) \end{aligned}$$

Proof. The proof parallels the one of Proposition B.1 deforming the permutation $\begin{pmatrix} 1 & 2 & 3 & 4 & 3 & 5 & 6 \\ 1 & 5 & 7 & 4 & 2 & 6 & 7 \end{pmatrix}$.

References

[DoHu] A. Douady, J. Hubbard, On the density of Strebel differentials, Invent. Math., 30, (1975), 175–179.

- [EsMaZo] A. Eskin, H. Masur, A. Zorich: Moduli spaces of Abelian differentials: the principal boundary, counting problems and the Siegel Veech constants, Publ. Math. Inst. Hautes Études Sci. 97, (2003), 61 - 179
- [FaLaPo] A. Fathi, F. Laudenbach, V. Poenaru: Travaux de Thurston sur les surfaces, Astérisque 66-67, (1979).
- [HuMa] J. Hubbard, H. Masur: Quadratic differentials and foliations, Acta Math. 142, (1979), 221–274.

- [Ke] M. Keane. Interval exchange transformations, Math. Z. 141 (1975), 25–31.
- [KeMaSm] S. Kerckhoff, H. Masur, J. Smillie. Ergodicity of Billiard Flows and Quadratic Differentials. Ann. of Math., 124, (1986), 293–311.
- [Ko] M.Kontsevich, Lyapunov exponents and Hodge theory. "The mathematical beauty of physics" (Saclay, 1996), (in Honor of C. Itzykson), Adv. Ser. Math. Phys. 24, World Sci. Publishing, River Edge, NJ, (1997), 318–332.
- [KoZo] M. Kontsevich, A. Zorich: Connected components of the moduli spaces of Abelian differentials with prescribed singularities, Invent. Math. 153 no. 3, (2003), 631–678.
- [La1] E. Lanneau, Hyperelliptic components of the moduli spaces of quadratic differentials with prescribed singularities, Comment. Math. Helv. 79 no. 3, (2004), 471–501.
- [La2] E. Lanneau, Parity of the Spin Structure Defined by a Quadratic Differential, Geom. Topol. 8 no. 12, (2004), 511–538.
- [La3] E. Lanneau, Classification of connected components of the strata of the moduli spaces of quadratic differentials with prescribed singularities, Thèse de Doctorat, Université de Rennes 1, (Décembre 2003).
- [Ma1] H. Masur, Interval exchange transformations and measured foliations, Ann of Math. 115 (1982) 169–200.
- [Ma2] H. Masur, The Jenkins-Strebel differentials with one cylinder are dense, Comment. Math. Helv. 54 (1979) 179–184.
- [MaSm] H. Masur, J. Smillie, Quadratic differentials with prescribed singularities and pseudo-Anosov diffeomorphisms, Comment. Math. Helv. 68 no. 2, (1993), 289–307.
- [MaTa] H. Masur, S. Tabachnikov, Rational billiards and flat structures, Handbook of dynamical systems, Vol. 1A, North-Holland, Amsterdam, (2002), 1015–1089.
- [MaZo] H. Masur, A. Zorich, Moduli Spaces of quadratic differentials: The Principal Boundary, Counting Problems and the Siegel-Veech Constants, submitted, (2004).
- [Ra] G. Rauzy, Echanges d'intervalles et transformations induites Acta Arith., 34 (1979), 315–328.
- [St] K. Strebel, *Quadratic differentials*, Springer-Verlag, (1984).
- [Th] W.P. Thurston, On the geometry and dynamics of diffeomorphisms of surfaces Bull. Amer. Math. Soc., 19 (1988), 417–431.
- [Ve1] W.A. Veech, Gauss measures for transformations on the space of interval exchange maps, Ann. of Math. 115, (1982), 201–242.
- [Ve2] W.A. Veech, Moduli spaces of quadratic differentials, Journal d'Analyse Math. 55, (1990), 117–171.
- [Ve3] W. Veech, The Teichmuller geodesic flow, Ann. of Math. **124** (1986), 441–530.
- [Zo1] A. Zorich, Explicit construction of a representative of any extended Rauzy class, to appear, 1–24.
- [Zo2] A. Zorich, *Flat Surfaces*, in collection "Frontiers in Number Theory, Physics and Geometry. Volume
 1: On random matrices, zeta functions and dynamical systems", P. Cartier; B. Julia; P. Moussa;
 P. Vanhove (Editors), Springer-Verlag, Berlin, (2006), 439-586.

Centre de Physique Théorique de Marseille (CPT), UMR CNRS 6207 Université du Sud Toulon-Var and

Fédération de Recherches des Unités de Mathématiques de Marseille

LUMINY, CASE 907, F-13288 MARSEILLE CEDEX 9, FRANCE

E-mail address: lanneau@cpt.univ-mrs.fr