# A CHARACTERISTIC NUMBER OF HAMILTONIAN BUNDLES OVER $S^2$

ANDRÉS VIÑA

ABSTRACT. Each loop  $\psi$  in the group  $\operatorname{Ham}(M)$  of Hamiltonian diffeomorphisms of a symplectic manifold M determines a fibration E on  $S^2$ , whose coupling class [5] is denoted by c. If VTE is the vertical tangent bundle of E, we relate the characteristic number  $\int_E c_1(VTE)c^n$  with the Maslov index of the linearized flow  $\psi_{t*}$  and the Chern class  $c_1(TM)$ . We give the value of this characteristic number for loops of Hamiltonian symplectomorphisms of Hirzebruch surfaces.

MSC 2000: 53D05, 57S05

# 1. INTRODUCTION

A loop  $\psi: S^1 \to \operatorname{Ham}(M, \omega)$  in the group of Hamiltonian diffeomorphisms [7] of a symplectic manifold  $(M^{2n}, \omega)$  can be considered as a clutching function of a Hamiltonian fibration  $E \xrightarrow{\pi} S^2$  with fibre M. The total space E supports the coupling class  $c \in H^2(E, \mathbb{R})$ ; this is the unique class such that  $c^{n+1} = 0$ , and  $i_p^*(c)$ is the cohomology class of the symplectic structure on the fibre  $\pi^{-1}(p)$ , where  $i_p$  is the inclusion of  $\pi^{-1}(p)$  in E [5]. Furthermore one can consider on E the first Chern class  $c_1(VTE)$  of the vertical tangent bundle of E. These canonical cohomology classes on E determine the characteristic number (see [6])

(1.1) 
$$I_{\psi} = \int_{E} c_1(VTE) c^n,$$

which depends only on the homotopy class of  $\psi$ . Since I is an  $\mathbb{R}$ -valued group homomorphism on  $\pi_1(\operatorname{Ham}(M,\omega))$ , the non vanishing of I implies that the group  $\pi_1(\operatorname{Ham}(M,\omega))$  is infinite. That is, I can be used to detect the infinitude of the corresponding homotopy group. Furthermore I calibrates the Hofer's norm  $\nu$  on  $\pi_1(\operatorname{Ham}(M,\omega))$  in the sense that  $\nu(\psi) \geq C|I_{\psi}|$ , for all  $\psi$ , where C is a positive constant [9].

I is a generalization of the mixed action-Maslov homomorphism introduced by Polterovich [8] for monotone manifolds, that is, when  $[\omega] = ac_1(TM)$  and a > 0. The value of this mixed action-Maslov homomorphism on a loop  $\psi$  is, in many cases, easy to calculate, since it is a linear combination of the symplectic action around any orbit  $\{\psi_t(x_0)\}_t$  and the Maslov index of the linearized flow  $(\psi_t)_*$  along this orbit. By contrast, I is defined for Hamiltonian loops in general manifolds (non necessarily monotone), and its value is mostly not so easy to determine from the definition.

Key words and phrases. Hamiltonian Diffeomorphisms, Symplectic Fibrations.

This work has been partially supported by Ministerio de Ciencia y Tecnología, grant MAT2003-09243-C02-00.

#### ANDRÉS VIÑA

Our purpose in this note is to obtain an explicit expression for  $I_{\psi}$ , which can be used to calculate its value. More precisely, when the bundle TM admits local symplectic trivializations whose domains are fixed by the diffeomorphisms  $\psi_t$ , we deduce a formula for  $I_{\psi}$  in which appear a contribution related to the Maslov indices of the linearized flow  $\psi_{t*}$  in the trivializations, and a second one in which are involved transition functions of the bundle  $\det(TM)$ . The second contribution is related with the Chern class  $c_1(TM)$  in the following sense. Using the expression of  $c_1(M)$  in terms of the transition functions of TM determined by the trivializations,  $\langle c_1(M)[\omega]^{n-1}, M \rangle$  can be written as a sum  $\sum_j \int_{R_j} \sigma_j$ , where  $\sigma_j$  is a 2n-1form (see (3.16)). It turns out that the second contribution is equal to this sum "weighted" by a multiple of the Hamiltonian  $f_t$  which generates  $\psi$ ; more concretely, that contribution is  $-n \sum_j \int dt \int_{R_j} (f_t \circ \psi_t) \sigma_j$ .

Let  $(M, \omega, f)$  be an integrable system such that the points where the integrals of motion are dependent form a set P which is union of codimension 2 submanifolds of M, and such that  $M \setminus P$  is invariant under  $\psi_t$  and on it there exist action-angle coordinates. Furthermore we assume that there are  $\psi_t$  invariant Darboux charts which cover P. Then the expression of  $I_{\psi}$  in this atlas reduces to the aforesaid second contribution; that is,  $I_{\psi} = -n \sum_j \int_{R_j} f \sigma_j$ .

The paper is organized as follows. In Section 2 we recall the construction of the coupling class c following [9]. Section 3 is concerned with the proof of the mentioned expression for  $I_{\psi}$ . First we express  $\langle c_1(M)[\omega]^{n-1}, M \rangle$  as the sum  $\sum_j \int_{R_j} \sigma_j$  of integrals of 2n-1 forms, and next we use this result to prove the formula for the invariant  $I_{\psi}$ . In Section 4 we check and apply the formulae obtained in Section 3. Using these formulae, we calculate  $I_{\psi}$ , when  $\psi$  is the loop in  $\operatorname{Ham}(S^2)$  generated by the 1-turn rotation of  $S^2$  around the z-axis. The result  $I_{\psi} = 0$  agrees with the fact that  $\pi_1(\operatorname{Ham}(S^2)) = \mathbb{Z}_2$  and I is a group homomorphism on  $\operatorname{Ham}(M)$ . We also prove that I on  $\pi_1(\operatorname{Ham}(\mathbb{T}^{2n}))$  vanishes identically. When n = 1 this result is consistent with the fact that  $\pi_1(\operatorname{Ham}(\mathbb{T}^2)) = 0$ . Finally we determine the value of I on the loops generated by action of  $\mathbb{T}^2$  on a general symplectic Hirzebruch surface (see Theorem 8).

I thank Dusa McDuff for explaining me properties of the Maslov index of the linearized flow, and Eva Miranda for clarifying me some points relative to actionangle variables.

### 2. The coupling class

Let  $(M, \omega)$  be a compact connected symplectic 2*n*-manifold. Let  $\psi : S^1 = \mathbb{R}/\mathbb{Z} \to$ Ham $(M, \omega)$  be a loop in the group Ham $(M, \omega)$  at id. By  $X_t$  is denoted the timedependent vector field generated by  $\psi_t$  and  $f_t$  is the normalized time-dependent Hamiltonian; that is,

$$\frac{d\psi_t}{dt} = X_t \circ \psi_t, \quad \iota_{X_t} \omega = -df_t, \quad \int_M f_t \omega^n = 0.$$

Given  $\epsilon$ , with  $0 < \epsilon < \pi/2$ , we set

$$D_{+}^{2} := \{ p \in S^{2} \mid 0 \le \theta(p) < \pi/2 + \epsilon \}$$
$$D_{-}^{2} := \{ p \in S^{2} \mid \pi/2 - \epsilon < \theta(p) \le \pi \},\$$

where  $\theta \in [0, \pi]$  is the polar angle from the z-axis.

Next we construct the Hamiltonian bundle E over  $S^2$  determined by  $\psi$ . First of all we extend  $\psi$  to a map defined on  $F := D^2_+ \cap D^2_-$  by putting  $\psi(\theta, \phi) = \psi_t$ , with  $t = \phi/2\pi$ , with  $\phi$  the spherical azimuth angle. We set

$$E = [(D_{+}^{2} \times M) \cup (D_{-}^{2} \times M)] / \simeq, \text{ where}$$
$$(+, p, x) \simeq (-, p', y) \text{ iff } \begin{cases} p = p' \in F, \\ y = \psi_{t}^{-1}(x), \ t = \phi(p)/2\pi. \end{cases}$$

In this way  $M \hookrightarrow E \xrightarrow{\pi} S^2$  is a Hamiltonian bundle over  $S^2$ .

We assume that  $D_{\pm}^2$  are endowed with the orientations induced by the usual one of  $S^2$  (that is, the orientation of  $S^2$  as border of the unit ball). We suppose that  $S^1$  is oriented by  $dt = d\phi/2\pi$ , that is,  $S^1$  is oriented as  $\partial D_+$ . In *E* one considers the orientation induced by the one defined on  $D_+^2 \times M$  by  $d\theta \wedge d\phi \wedge \omega^n$ .

Let  $\alpha$  be a monotone smooth map  $\alpha : [\pi/2 - \epsilon, \pi] \to [0, 1]$ , with  $\alpha(\theta) = 1$  for  $\theta \in [\pi/2 - \epsilon, \pi/2 + \epsilon]$  and  $\alpha(\theta) = 0$  for  $\theta$  near  $\pi$ . Now we consider the 2-form (see [9])

(2.1) 
$$\tau = \begin{cases} \omega, \text{ on } D_+^2 \times M\\ \omega + d(\alpha(f_t \circ \psi_t)) \wedge dt, \text{ on } D_-^2 \times M. \end{cases}$$

As  $\alpha$  vanishes near  $\pi$ ,  $\tau$  is well-defined on  $D^2_- \times M$ ; moreover on  $F \times M \subset D^2_- \times M$ ,  $\tau$  reduces to  $\omega + d(f_t \circ \psi_t) \wedge dt$ . If we denote by h the map

$$h: F \times M \subset D^2_- \times M \to F \times M \subset D^2_+ \times M$$

given by  $h(p,x) = (p,\psi_t(x))$ , with  $t = \phi(p)/2\pi$ , then taking into account that  $h_*(\frac{\partial}{\partial t}) = \frac{\partial}{\partial t} + X_t \circ \psi_t$ , it follows from  $\iota_{X_t}\omega = -df_t$  that  $h^*\omega = \omega + d(f_t \circ \psi_t) \wedge dt$ . So one has the following Proposition

**Proposition 1.**  $\tau$  defines a closed 2-form on E.

Moreover the cohomology class  $[\tau] \in H^2(E, \mathbb{R})$  restricted to each fibre coincides with  $[\omega]$ . On the other hand

$$\int_E \tau^{n+1} = (n+1) \int_{D^2_- \times M} (f_t \circ \psi_t) \alpha'(\theta) d\theta \wedge dt \wedge \omega^n.$$

From the normalization condition for  $f_t$  it follows that  $\int_E \tau^{n+1} = 0$ . Hence  $[\tau]$  is the coupling class c of the fibration E [5] [7].

3. The characteristic number  $I_{\psi}$ .

Denoting  $TM = \{v_x \in T_xM \mid x \in M\}$ , we put

$$VTE = \left[ (D_+^2 \times TM) \cup (D_-^2 \times TM) \right] / \simeq,$$

with

(

$$(+, p, v_x) \simeq (-, p', v'_{x'})$$
 iff  $p = p', x' = \psi_t^{-1}(x), v'_{x'} = (\psi_t^{-1})_*(v_x)$ 

where  $t = \phi(p)/2\pi$ . So VTE is a vector bundle over E; by construction it is the vertical tangent bundle of E.

Let  $(U; X_1, \ldots, X_{2n})$  be a symplectic trivialization of TM on  $U \subset M$ , and  $(V; Y_1, \ldots, Y_{2n})$  be a symplectic trivialization on  $V \subset M$ . We put

(3.1) 
$$U_{\pm} := \{ [\pm, p, x] \mid p \in D^2_+, x \in U \}$$

and similarly for  $V_{\pm}$ . Denoting  $x_t := \psi_t^{-1}(x)$  one has

$$U_{+} \cap U_{-} = \{ [+, p, x] \mid p \in F, x \in U, x_{t} \in U \}$$
$$V_{+} \cap V_{-} = \{ [+, p, x] \mid p \in F, x \in V, x_{t} \in V \}$$
$$V_{-} \cap U_{-} = \{ [-, p, x] \mid p \in D_{-}^{2}, x \in V \cap U \}$$
$$U_{+} \cap V_{+} = \{ [+, p, x] \mid p \in D_{+}^{2}, x \in V \cap U \}$$

The corresponding transition functions of VTE are

$$g_{U_{-}U_{+}}([+, p, x]) = A(t, x) \in Sp(2n, \mathbb{R}), \text{ with } \psi_{t^{*}}^{-1}(X_{i}(x)) = \sum_{k} A^{k}{}_{i}(t, x) X_{k}(x_{t})$$
$$g_{V_{-}V_{+}}([+, p, x]) = B(t, x) \in Sp(2n, \mathbb{R}), \text{ with } \psi_{t^{*}}^{-1}(Y_{i}(x)) = \sum_{k} B^{k}{}_{i}(t, x) Y_{k}(x_{t})$$
$$g_{U_{-}V_{-}}([-, p, x]) = R(x) = g_{U_{+}V_{+}}([+, p, x]), \text{ with } Y_{i}(x) = \sum_{k} R^{k}{}_{i}(x) X_{k}(x).$$

We denote by  $\rho$  the usual map  $\rho: Sp(2n, \mathbb{R}) \to U(1)$  which restricts to the determinant map on U(n) [10], then  $l_{ab} := \rho \circ g_{ab}$  is a transition function for det(VTE). We also use the following notation, the matrices in  $Sp(2n, \mathbb{R})$  are denoted with capital letters and its images by  $\rho$  will be denoted by the corresponding small letters; that is,

(3.2) 
$$a(t,x) := \rho(A(t,x)), \quad b(t,x) := \rho(B(t,x)), \quad r_{UV}(x) := \rho(R(x)).$$

If  $\psi_t(U) \subset U$  for all t, given  $x \in U$ , the winding number of the map  $t \in S^1 \mapsto a^{-1}(t,x) \in U(1)$  is the integer

(3.3) 
$$\frac{i}{2\pi} \int_0^1 a^{-1}(t,x) \frac{\partial a}{\partial t}(t,x) dt.$$

This integer is independent of the point  $x \in U$ , it will be denoted  $J_U$ . The number  $J_U$  is the Maslov index in U of the linearized flow  $\psi_{t*}$ . Analogously, if  $\psi_t(V) \subset V$  for all t we have the integer

(3.4) 
$$J_V = \frac{i}{2\pi} \int_0^1 b^{-1}(t,x) \frac{\partial b}{\partial t}(t,x) dt,$$

x being any point of V; this is the Maslov index in V of  $\psi_{t*}$ .

As a previous step to compute  $I_{\psi}$  we shall prove the following Lemma, in which the value  $\langle c_1(M)[\omega]^{n-1}, [M] \rangle$  is expressed in terms of transition functions of det(TM).

**Lemma 2.** Let  $\{B_1, \ldots, B_m\}$  be a set of trivializations of TM, such that its domains cover M. Then

(3.5) 
$$\langle c_1(TM)[\omega]^{n-1}, [M] \rangle = \frac{-i}{2\pi} \sum_{i < k} \int_{A_{ik}} d(\log s_{ik}) \wedge \omega^{n-1},$$

 $s_{ik}$  being the corresponding transition function of det(TM) and

(3.6) 
$$A_{ik} = \left(\partial B_i \setminus \bigcup_{r < k} B_r\right) \cap B_k.$$

*Proof.*  $c_1(M)$  is represented on  $B_a$  by the 2-form

$$\frac{-i}{2\pi} \sum_{c} d(\varphi_c d \log s_{ac}),$$

where  $\{\varphi_c\}$  is a partition of unity subordinate to the covering  $\{B_1, \ldots, B_m\}$ . If m = 2

$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \frac{-i}{2\pi} \int_{B_1} d(\varphi_2 d \log s_{12}) \wedge \omega^{n-1} + \frac{-i}{2\pi} \int_{B_2 \setminus B_1} d(\varphi_1 d \log s_{21}) \wedge \omega^{n-1}$$

By Stokes' theorem

(3.7) 
$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \int_{\partial B_1} \varphi_2 L_{12} + \int_{\partial (B_2 \setminus B_1)} \varphi_1 L_{21}$$

where

$$L_{jk} := (-i/2\pi)d \log s_{jk} \wedge \omega^{n-1}.$$

Since  $\partial(B_2 \setminus B_1) \cap B_1 = \emptyset$ ,  $\varphi_1$  vanishes on  $\partial(B_2 \setminus B_1)$  and the last integral in (3.7) is zero.

As  $\varphi_2$  is 1 on  $\partial B_1$ , we have

$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \int_{\partial B_1} L_{12}.$$

In this case  $\partial B_1 \subset B_2$ , so  $\partial B_1 = A_{12}$ , and the Lemma is proved when m = 2. If m = 3

(3.8) 
$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \int_{\partial B_1} \left( \varphi_2 L_{12} + \varphi_3 L_{13} \right) + \int_{\partial (B_2 \setminus B_1)} \left( \varphi_1 L_{21} + \varphi_3 L_{23} \right)$$
  
(3.9)  $+ \int \left( \varphi_1 L_{31} + \varphi_2 L_{32} \right).$ 

(3.3) 
$$+ \int_{\partial(B_3 \setminus (B_1 \cup B_2))} (\varphi_1 L_{31} + \varphi_2 L_{32}).$$
  
As  $\partial(B_2 \setminus (B_1 \cup B_2))$  and the interior of  $B_1 \cup B_2$  are disjoint sets.

As  $\partial(B_3 \setminus (B_1 \cup B_2))$  and the interior of  $B_1 \cup B_2$  are disjoint sets,  $\varphi_1$  and  $\varphi_2$  vanish on  $\partial(B_3 \setminus (B_1 \cup B_2))$ , and the integral in (3.9) is zero. Analogously  $\partial(B_2 \setminus B_1)$  and support of  $\varphi_1$  are disjoint so

(3.10) 
$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \int_{\partial B_1} \left( \varphi_2 L_{12} + \varphi_3 L_{13} \right) + \int_{\partial (B_2 \setminus B_1)} \varphi_3 L_{23}$$

On the other hand  $\partial B_1 = A + D$ , with  $A := \partial B_1 \setminus B_2$  (oriented as  $\partial B_1$ ) and  $D := (\partial B_1 \setminus B_2) \cap B_3$  (see Figure 1). Moreover  $\partial (B_2 \setminus B_1) = -A + C$  with  $C := (\partial B_2 \setminus B_1) \cap B_3$  (oriented as  $\partial B_2$ ).

Since  $C \cap (B_1 \cup B_2) = \emptyset$ , then  $\varphi_3|_C = 1$ ; thus

(3.11) 
$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \int_{A+D} \left( \varphi_2 L_{12} + \varphi_3 L_{13} \right) + \int_{-A} \varphi_3 L_{23} + \int_{A_{23}} L_{23}.$$

The last integral in (3.11) is just the term in (3.5) with i = 2, k = 3.

Since  $\varphi_j|_D = 0$ , for j = 1, 2, then  $\varphi_3|_D = 1$ . As A and support of  $\varphi_1$  are disjoint sets, then  $(\varphi_2 + \varphi_3)|_A = 1$ . It follows from these facts together with the cocycle condition  $L_{13} + L_{32} = L_{12}$  that

(3.12) 
$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \int_A L_{12} + \int_D L_{13} + \int_{A_{23}} L_{23}.$$

On the other hand  $A_{12} = (\partial B_1 \setminus B_1) \cap B_2 = A$ . Similarly  $A_{13} = D$ . Therefore (3.12) is the formula given in the statement of Lemma when m = 3.



FIGURE 1.  $A = \partial B_1 \cap B_2$ ,  $D = (\partial B_1 \setminus B_2) \cap B_3$  and  $C = (\partial B_2 \setminus B_1) \cap B_3$ .

The preceding arguments can be generalized to any m

$$\langle c_1(TM)[\omega]^{n-1}, [M] \rangle = \int_{\partial B_1} \sum_{j \neq 1} \varphi_j L_{1j} + \dots + \int_{\partial (B_{m-1} \setminus \bigcup_{r < m-1} B_r)} \sum_{j \neq m-1} \varphi_j L_{m-1,j}$$

$$(3.14) \qquad \qquad + \int_{\partial (B_m \setminus \bigcup_{r < m} B_r)} \sum_{j \neq m} \varphi_j L_{m-1,j}.$$

For any j = 1, ..., m-1 support of  $\varphi_j$  and  $\partial (B_m \setminus \bigcup_{r < m} B_r)$  are disjoint sets. Thus the integral (3.14) is zero (as in the cases m = 2, 3). We decompose

$$\partial(B_{m-1} \setminus \cup_{r < m-1} B_r) = E + G,$$

with

$$E := \left(\partial B_{m-1} \setminus \bigcup_{r < m-1} B_r\right) \cap B_m.$$

Then  $\varphi_j|_E = 0$  for all  $j \neq m$  and  $\varphi_m|_E = 1$ ; thus

(3.15) 
$$\int_{\partial(B_{m-1}\setminus\cup_{r< m-1}B_r)} \sum_{j\neq m-1} \varphi_j L_{m-1,j} = \int_G + \int_{A_{m-1,m}} L_{m-1,m}.$$

The last integral in (3.15) is the term in (3.5) which corresponds to i = m-1, k = m. An analogous, but more tedious, calculation to the one for the case m = 3 allows to identify in (3.13) the remainder terms of (3.5).

Lemma 2 gives a way for expressing  $\langle c_1(TM)[\omega]^{n-1}, [M] \rangle$  as a sum of integrals of 2n-1 differential forms on 2n-1 chains. The righthand side of (3.5) can be written schematically

(3.16) 
$$\sum_{j} \int_{R_j} \sigma_j.$$

In next Theorem we use this expression to give an explicit formula for  $I_{\psi}$  in terms of transition functions of det(TM) and Maslov indices of  $\psi_{t*}$ .

**Theorem 3.** If  $\{B_1, \ldots, B_m\}$  is a set of symplectic trivializations for TM which covers M, and such that  $\psi_t(B_j) = B_j$ , for all t and all j, then

(3.17) 
$$I_{\psi} = \sum_{i=1}^{m} J_i \int_{B_i \setminus \bigcup_{j < i} B_j} \omega^n + \sum_{i < k} N_{ik},$$

where

$$N_{ik} = n \frac{i}{2\pi} \int_0^1 dt \int_{A_{ik}} (f_t \circ \psi_t) (d \log r_{ik}) \wedge \omega^{n-1},$$

 $A_{ik} = (\partial B_i \setminus \bigcup_{r < k} B_r) \cap B_k$ ,  $J_i$  is the Maslov index of  $(\psi_t)_*$  in the trivialization  $B_i$ and  $r_{ik}$  the corresponding transition function of det(TM).

*Proof.* Using the notation (3.1) we put

(3.18) 
$$O_{2\mathbf{a}-\mathbf{1}} := (B_a)_-, \quad O_{2\mathbf{a}} := (B_a)_+.$$

Then  $\{O_{\mathbf{c}} | c = 1, ..., 2m\}$  is a covering for E. We shall denote by  $l_{\mathbf{bc}}$  the respective transition functions for det (VTE). If we set  $U := B_1, V := B_2$ , one has by (3.2)

$$l_{12} = a(t, x), \ l_{13} = r_{UV}(x), \ l_{34} = b(t, x)$$

We can determine  $I_{\psi} = \langle c_1(VTE) c^n, [E] \rangle$  applying the result given in Lemma 2 to the set  $\{O_c\}$  of trivializations of VTE. That is,

(3.19) 
$$I_{\psi} = \sum_{\mathbf{a} < \mathbf{b}} \mathcal{T}_{\mathbf{a}\mathbf{b}}, \text{ where } \mathcal{T}_{\mathbf{a}\mathbf{b}} = \frac{-i}{2\pi} \int_{A_{\mathbf{a}\mathbf{b}}} d \log l_{\mathbf{a}\mathbf{b}} \wedge \tau^n.$$

It follows from (3.18) and (2.1) that  $\tau$  is equal to  $\omega$  on  $A_{ab}$  unless **a** and **b** are both odd; in this case  $\tau = \omega + d(\alpha(f_t \circ \psi_t)) \wedge dt$ .

We will calculate the summand  $\mathcal{T}_{12}$  in (3.19). The set  $A_{12} = \partial O_1 \cap O_2 = \partial U_- \cap U_+$ , and

$$\partial U_{-} = \{ [+, p, x] \mid p \in \partial D_{-}^{2}, \ x \in U \} \cup \{ [-, p, x] \mid p \in D_{-}^{2}, \ x \in \partial U \}.$$

 $\operatorname{So}$ 

$$A_{12} = \{ [+, p, x] \mid p \in \partial D_{-}^2, x \in U \}.$$

Taking into account (3.3), (3.2) together with the fact that orientations of  $S^1$  and  $\partial D^2_-$  are opposite , we deduce

$$\mathcal{T}_{12} = \frac{-i}{2\pi} \int_U \left( \int_{-S^1} a^{-1}(t,x) \frac{\partial a(t,x)}{\partial t} dt \right) \omega^n = J_U \int_U \omega^n.$$

Next we consider the term  $\mathcal{T}_{34}$ . The integration domain is

$$A_{\mathbf{34}} = \left(\partial V_- \setminus (U_- \cup U_+)\right) \cap V_+ = \{[+, p, x] \mid p \in \partial D_-^2, x \in V \setminus U\}$$

Hence

$$\mathcal{T}_{\mathbf{34}} = J_V \int_{V \setminus U} \omega^n.$$

In general,

$$A_{2\mathbf{j}-\mathbf{1},2\mathbf{j}} = \left(\partial B_{j-} \setminus \bigcup_{r < j} (B_{r+} \cup B_{r-})\right) \cap B_{j+}$$
$$= \{[+, p, x] \mid p \in \partial D^2_{-}, x \in B_j \setminus \bigcup_{r < j} B_r\}.$$

Hence the term in (3.19) with  $\mathbf{a} = 2\mathbf{j} - \mathbf{1}$ ,  $\mathbf{b} = 2\mathbf{j}$  gives a contribution to  $I_{\psi}$  equal to

(3.20) 
$$J_{B_j} \int_{B_j \setminus U_{r < j} B_r} \omega^n$$

Now we analyze  $\mathcal{T}_{13}$ .

$$A_{13} = \{ [-, p, x] \mid p \in D_{-}^{2}, \ x \in \partial U \cap V \}.$$

 $D_{-}^{2}$  is oriented by the form  $d\theta \wedge dt$ , and  $\partial U \cap V$  is oriented with the orientation of  $\partial U$ . Hence

(3.21) 
$$\mathcal{T}_{\mathbf{13}} = \frac{-i}{2\pi} \int_{A_{\mathbf{13}}} d \log r_{UV} \wedge \left(\omega + d(\alpha(f_t \circ \psi_t)) \wedge dt\right)^n$$
$$= \frac{-ni}{2\pi} \int_{A_{\mathbf{13}}} d \log r_{UV} (f_t \circ \psi_t)) \alpha'(\theta) d\theta \wedge dt \wedge \omega^{n-1}$$
$$= \frac{+ni}{2\pi} \int_0^1 dt \int_{\partial U \cap V} (f_t \circ \psi_t) d \log r_{UV} \wedge \omega^{n-1}.$$

In general, if j < k

(3.22) 
$$\mathcal{T}_{2\mathbf{j}-1,2\mathbf{k}-1} = \frac{ni}{2\pi} \int_0^1 dt \int_{A_{jk}} (f_t \circ \psi_t) d \log r_{jk} \wedge \omega^{n-1},$$

where  $A_{jk}$  is the set defined in Lemma 2.

On the other hand

$$A_{\mathbf{14}} = \left(\partial U_{-} \setminus (U_{+} \cup V_{-})\right) \cap V_{+} = \{[-, p, x] \mid p \in D_{-}^{2}, x \in \partial U \setminus V\} \cap V_{+} = \emptyset.$$

Thus  $\mathcal{T}_{14} = 0$ . In general, for j < k the integration domain  $A_{2j-1,2k}$  is of the form

$$(\partial B_{j-} \setminus \cup \cdot) \cap B_{k+}.$$

In the union  $\cup \cdot$  appear the sets  $B_{k-}$  and  $B_{j+}$ , hence

$$A_{2j-1,2k} \subset (\partial B_{j-} \setminus (B_{j+} \cup B_{k-})) \cap B_{k+},$$

and this set is empty by the same reason that  $A_{14} = \emptyset$ . Therefore  $\mathcal{T}_{2j-1,2k} = 0$ , for any j < k.

The set  $A_{23}$  is

$$A_{23} = (\partial U_+ \setminus U_-) \cap V_- = \{ [+, p, x] \mid p \in F, x \in \partial U \setminus V \}.$$

As  $d \log l_{23} \wedge \omega^n$  does not contain  $d\theta$ , the term  $\mathcal{T}_{23}$  vanishes. In general, if j < k

$$A_{2j,2k-1} = (\partial B_{j+} \setminus \cup \cdot) \cap B_{k-} \subset (\partial B_{j+} \setminus B_{j-}) \cap B_{k-}$$
$$= \{ [+, p, x] \mid p \in F, x \in \partial B_j \setminus B_k \}.$$

Then  $\mathcal{T}_{2j,2k-1}$  vanishes by the same reason that  $\mathcal{T}_{23} = 0$ .

Analogous arguments as the ones explained in the preceding paragraph show that  $\mathcal{T}_{2j,2k} = 0$ , for any j < k.

So, apart from the terms  $\mathcal{T}_{ab}$  considered in (3.20) and in (3.22), the remainder summands in (3.19) are zero. The theorem follows from (3.20) and (3.22).

8

From the definition of product in  $\pi_1(\text{Ham}(M, \omega))$  by juxtaposition of paths and under the hypotheses of Theorem 3 is obvious that

$$I: \pi_1(\operatorname{Ham}(M,\omega)) \to \mathbb{R}$$

is a group homomorphism. This fact has been proved in [6] for the general case.

**Corollary 4.** If U and V are symplectic trivializations of TM, with  $\psi_t(U) = U$ ,  $\psi_t(V) = V$ , for all t and  $U \cup V = M$  and  $\int_{S^1} (f_t \circ \psi_t) dt$  is a constant k on  $\partial U \cap V$ , then

$$I_{\psi} = J_U \int_U \omega^n + J_V \int_{V \setminus U} \omega^n - nk \langle c_1(TM)[\omega]^{n-1}, M \rangle.$$

**Corollary 5.** If TM is trivial on  $U := M \setminus \{q\}$ , where q is a point of M fixed by  $\psi_t$  for all t, then

$$I_{\psi} = J_U \int_M \omega^n - n \Big( \int_{S^1} f_t(q) dt \Big) \langle c_1(TM)[\omega]^{n-1}, M \rangle.$$

Now we analyze the expression for  $I_{\psi}$  given in Theorem 3 in case of *integrable systems*. Let f be the normalized Hamiltonian which generates the loop  $\psi$ . We assume that  $(M, \omega, f)$  is completely integrable, with  $f_1 = f, f_2, \ldots, f_n$  integrals of motion. We suppose that  $df_1, \ldots, df_n$  are independent at the points of  $M \setminus P =: V$ , where P is a finite union of 2n - 2 dimensional submanifolds of M. We suppose that on V are defined action-angle coordinates. We put

$$Q := \{x \in P \mid \dim \operatorname{Span} \left( df_1(x), \dots, df_n(x) \right) = n - 1 \}.$$

By  $Q_1, \ldots, Q_k$  are denoted the connected components of Q, and let  $V_j$  be a tubular neighborhood of  $Q_j$  in M, invariant under  $\psi_t$  for all t. We assume that on  $V_j$  is defined a symplectic trivialization of TM. Then, for each j one can choose a family of tubular neighborhoods  $\{V_{jb} \subset V_j\}_{b=1,2...}$ , such that

$$\lim_{b \to \infty} \int_{V_{jb}} \omega^n = 0.$$

Lemma 2 applied to the covering  $\{V, V_{jb}\}_{j=1,...,k}$  of  $V \cup Q$  gives

$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \frac{-i}{2\pi} \sum_{j=1}^k \int_{\partial V \cap V_{jb}} d\log r_{VV_{jb}} + \epsilon(b),$$

where  $\epsilon(b)$  goes to 0 as  $b \to \infty$ .

Hence

(3.23) 
$$\langle c_1(M)[\omega]^{n-1}, [M] \rangle = \sum_{j=1}^k z_j,$$

with

(3.24) 
$$z_j := \frac{-i}{2\pi} \sum \int_{\partial V \cap V_{jb}} d \log r_{VV_{jb}} \wedge \omega^{n-1}.$$

**Proposition 6.** Let  $(M, \omega, f, f_2, \dots, f_n)$  be an integrable in which the preceding hypotheses hold, then

$$I_{\psi} = \sum_{j=1}^{k} z'_j,$$

where  $z'_j$  is obtained from the corresponding  $z_j$  by inserting the factor -nf in the integrand of (3.24).

*Proof.* The Maslov index  $J_V = 0$  because of the particular form of the flow equations in action-angle coordinates. On the other hand

$$\int_{V_{jb}\setminus(V\cup\dots)}\omega^n=0$$

Thus the Proposition follows from Theorem 3, together with (3.23) and (3.24).

Similar arguments to the ones involved in this Proposition are used in Section 4 for studying the invariant I in Hirzebruch surfaces.

### 4. Examples.

The invariant I when the manifold is the 2-sphere.

Let  $\psi_t$  be the rotation in  $\mathbb{R}^3$  around  $\vec{e}_3$  of angle  $2\pi t$  with  $t \in [0, 1]$ . Then  $\psi_t$  determines a Hamiltonian symplectomorphism of  $(S^2, \omega_{area})$ . In fact, the isotopy  $\{\psi_t\}$  is generated by the vector field  $\frac{\partial}{\partial \phi}$ , and the function f on  $S^2$  defined by  $f(\theta, \phi) = -2\pi \cos \theta = -2\pi z$  is the corresponding normalized Hamiltonian.

 $TS^2$  can be trivialized on  $U = D^2_+$ , and on  $V = D^2_-$ . Moreover  $\partial U \cap V$  is the parallel  $\theta = \pi/2 + \epsilon$ . On  $\partial U \cap V$  the function  $f \circ \psi_t$  takes the value  $2\pi \sin \epsilon$ .

$$\int_U \omega = 2\pi (1-k'), \quad \int_{V \setminus U} \omega = 2\pi (1+k'),$$

with  $k' := \cos(\pi/2 + \epsilon)$ .

Furthermore the north pole n and the south pole s are fixed points of the isotopy  $\psi_t$ . The rotation  $\psi_t$  transforms the basis  $\vec{e_1}$ ,  $\vec{e_2}$  of  $T_n S^2$  in

 $(\cos 2\pi t \,\vec{e_1} + \sin 2\pi t \,\vec{e_2}, \, -\sin 2\pi t \,\vec{e_1} + \cos 2\pi t \,\vec{e_2}).$ 

So  $J_U$  is the winding number of the map

$$t \in [0, 1] \to e^{2\pi t i} \in U(1);$$

That is,  $J_U = +1$ .

Similarly, by considering the oriented basis  $\vec{e}_2$ ,  $\vec{e}_1$  of  $T_s S^2$  it turns out that the Maslov index  $J_V$  of  $\psi_t$  is -1.

By Corollary 4

$$I_{\psi} = 2\pi(1-k') - 2\pi(1+k') - (-2\pi k')\langle c_1(TS^2), S^2 \rangle = 0.$$

Corollary 5 can also be applied to determine  $I_{\psi}$ . One takes  $U := S^2 \setminus \{s\}$ . As  $f(s) = -2\pi(-1)$ , we obtain again

$$I_{\psi} = +4\pi - 2\pi \langle c_1(TS^2), S^2 \rangle = 0.$$

Using formula (??) we can determine  $I_{\psi}$  again. Now V is  $S^2 \setminus \{n, s\}$ ,  $U_1$  is a small polar cap at n and  $U_2$  the symmetric one at s. By the symmetry

$$\int_{\partial U_1 \cap V} d \log r_{U_1 V} = \int_{\partial U_2 \cap V} d \log r_{U_2 V},$$

so  $y_1 = y_2$ . As f(n) = -f(s), then  $I_{\psi} = 0$ .

This result was expected, because  $\pi_1(\text{Ham}(S^2))$  is isomorphic to  $\mathbb{Z}_2$  (see [9]) and I is a group homomorphism.

The invariant I for Hamiltonian loops in  $\mathbb{T}^{2n}$ .

We identify the torus  $\mathbb{T}^{2n}$  with  $\mathbb{R}^{2n}/\mathbb{Z}^{2n}$ , and we suppose that  $\mathbb{T}^{2n}$  is equipped with the standard symplectic form  $\omega_0$ . If  $\psi_t$  is a Hamiltonian isotopy of  $\mathbb{T}^{2n}$ , it can be written in the form

$$\psi_t(x^1, \dots, x^{2n}) = \left(x^1 + \alpha^1(t, x^i), \dots, x^{2n} + \alpha^{2n}(t, x^i)\right),\$$

where the function  $\alpha^j$ , for j = 1, ..., 2n, is periodic of period 1 in each variable:  $t, x^1, ..., x^{2n}$ . The vector fields  $\{\frac{\partial}{\partial x^i}\}$  give a symplectic trivialization of the tangent bundle. In this case the right hand side of (3.17) has only one term. The matrix of  $(\psi_t)_*$  with respect to  $\{\frac{\partial}{\partial x^i}\}$  is

(4.1) 
$$\left(\delta_i^j + \frac{\partial \alpha^j}{\partial x^i}\right) \in Sp(2n, \mathbb{R}).$$

First, let us assume that each  $\alpha^j$  is a separate variables function; that is,  $\alpha^j(t, x^i) = f^j(t)u^j(x^i)$ . Since  $\alpha^1$  takes the same value at symmetric points on opposite faces of the cube  $I^{2n}$ , there is a point  $p_1 \in I^{2n}$  such

$$\frac{\partial u^1}{\partial x^j}(p_1) = 0,$$

for all j. Hence the first row of the matrix (4.1) at the point  $p_1$  is  $(1, 0, \ldots, 0)$ ; that is, the matrix of  $(\psi_t)_*(p_1)$  is independent of  $f^1$  and thus the Maslov index of  $\{(\psi_t)_*(p_1)\}_t$  does not depend on  $f^1$ . From (3.17) it follows that  $I_{\psi}$  is independent of  $f^1$ . The independence of  $I_{\psi}$  with respect to  $f^i$  is proved in a similar way. Thus in order to determine  $I_{\psi}$  we can assume that  $f^i = 0$  for all i, but in this case  $I_{\psi} = 0$ obviously.

If  $\alpha^j$  is sum of two separate variables functions

$$\alpha^j(t,x^i)=f^j(t)u^j(x^i)+g^j(t)v^j(x^i),$$

we take a point  $q_1 \in I^{2n}$ , such that  $\frac{\partial v^1}{\partial x^j}(q_1) = 0$ , for all j. Then  $I_{\psi}$  is independent of  $g^1$ . The above reasoning gives  $I_{\psi} = 0$  in this case as well.

By the Fourier theory, the original  $C^{\infty}$  periodic function  $\alpha^{j}$  can be approximated (in the uniform  $C^{k}$ -norm) by a sum of separated functions of the form  $\sum f_{a}(t)u_{a}(x^{i})$ , where  $f_{a}$  and  $u_{a}$  are 1-periodic. As  $I_{\psi}$  depends only on the homotopy class of  $\psi$ , we conclude that  $I_{\psi} = 0$  for a general Hamiltonian loop.

**Proposition 7.** The invariant I is identically zero on  $\pi_1(Ham(\mathbb{T}^{2n}, \omega_0))$ .

This result when n = 1 is consistent with the fact that  $\pi_1(\operatorname{Ham}(\mathbb{T}^2)) = 0$  (see [9])

Application to Hirzebruch surfaces.

Given 3 numbers  $k, \tau, \mu$ , with  $k \in \mathbb{Z}_{>0}, \tau, \mu \in \mathbb{R}_{>0}$  and  $k\mu < \tau$ , the triple  $(k, \tau, \mu)$  determine a Hirzebruch surface  $M_{k,\tau,\mu}$  [3]. This manifold is the quotient

$$\{z \in \mathbb{C}^4 : k|z_1|^2 + |z_2|^2 + |z_4|^2 = \tau/\pi, |z_1|^2 + |z_3|^2 = \mu/\pi\}/\mathbb{T}^2,$$

where the  $\mathbb{T}^2$ -action is given by

$$(a,b) \cdot (z_1, z_2, z_3, z_4) = (a^k b z_1, a z_2, b z_3, a z_4),$$

for  $(a, b) \in \mathbb{T}^2$ . The map

$$[z_1, z_2, z_3, z_4] \mapsto ([z_2 : z_4], [z_2^k z_3 : z_4^k z_3 : z_1])$$

# ANDRÉS VIÑA

allows us to represent  $M_{k,\tau,\mu}$  as a submanifold of  $\mathbb{C}P^1 \times \mathbb{C}P^2$ . On the other hand the usual symplectic structures on  $\mathbb{C}P^1$  and  $\mathbb{C}P^2$  induce a symplectic form  $\omega$  on  $M_{k,\tau,\mu}$ , and the following  $\mathbb{T}^2$ -action on  $\mathbb{C}P^1 \times \mathbb{C}P^2$ 

 $(a,b)([u_0:u_1],[x_0:x_1:x_2]) = ([au_0:u_1],[a^kx_0:x_1:bx_2])$ 

gives rise to a toric structure on  $M_{k,\tau,\mu}$ . In terms of the Delzant construction  $(M_{k,\tau,\mu}, \omega)$  is associated to the trapezoid in  $(\mathbb{R}^2)^*$  whose not oblique edges are  $\tau, \mu, \lambda := \tau - k\mu$ , [4] (see Figure 2). Moreover  $\lambda$  is the value that the symplectic form  $\omega$  takes on the exceptional divisor,  $\{[z] \in M \mid z_3 = 0\}$ , of  $M := M_{k,\tau,\mu}$ . And  $\omega$  takes the value  $\mu$  on the class of the fibre in the fibration  $M \to \mathbb{C}P^1$ .



FIGURE 2. Delzant polytope associated to M.

Since M is a toric manifold, the  $\mathbb{T}^2$ -action define symplectomorphisms of M. More precisely, let  $\psi_t$  the diffeomorphism of M defined by

(4.2) 
$$\psi_t[z_1, z_2, z_3, z_4] = [z_1 e^{2\pi i t}, z_2, z_3, z_4]$$

 $\psi = \{\psi_t : t \in [0,1]\}$  is a loop of Hamiltonian symplectomorphisms of  $(M,\omega)$ . Similarly we have

(4.3) 
$$\tilde{\psi}_t[z_1, z_2, z_3, z_4] = [z_1, z_2 e^{2\pi i t}, z_3, z_4],$$

and the corresponding loop  $\tilde{\psi}$  in  $\operatorname{Ham}(M, \omega)$ .

Using Theorem 3 we shall calculate the values of  $I_{\psi}$  and  $I_{\tilde{\psi}}$  in terms of  $\lambda$ ,  $\tau$  and k. The result is stated in Theorem 8 below. The most laborious point in the proof of the following Theorem is to obtain Darboux charts for M which give rise to simple transition functions for det(TM).

**Theorem 8.** Let  $\psi$  and  $\tilde{\psi}$  be the loops of symplectomorphisms of the Hirzebruch surface  $(M_{k,\tau,\mu}, \omega)$ , defined by (4.2) and (4.3) respectively, then

$$I_{\psi} = \frac{2k\mu^2}{3} \left( 1 - \frac{\mu}{2\lambda + k\mu} \right), \text{ and } I_{\tilde{\psi}} = \frac{-k^2\mu^2}{3} \left( 1 - \frac{\mu}{2\lambda + k\mu} \right).$$

 $\lambda$  being  $\tau - k\mu$ .

*Proof.* We will define a Darboux atlas on M. First we consider the following covering for M

$$U_1 = \{ [z] \in M : z_3 \neq 0 \neq z_4 \}, \ U_2 = \{ [z] \in M : z_1 \neq 0 \neq z_4 \}$$

 $U_3 = \{[z] \in M : z_1 \neq 0 \neq z_2\}, \ U_4 = \{[z] \in M : z_2 \neq 0 \neq z_3\}.$ We set  $z_j = \rho_j e^{i\theta_j}$ , with  $\rho_j = |z_j|$ , and on  $U_1$  introduce the coordinates  $(x_1, y_1, a_1, b_1)$  by the formulae

 $x_1 + iy_1 = \rho_1 e^{i\varphi_1}, \ a_1 + ib_1 = \rho_2 e^{i\varphi_2}, \ \varphi_1 = \theta_1 - \theta_3 - k\theta_4, \ \varphi_2 = \theta_2 - \theta_4.$ 

Then  $\omega$  on  $U_1$  can be written  $\omega = dx_1 \wedge dy_1 + da_1 \wedge db_1$ .

On  $U_2$  we consider the Darboux coordinates  $(x_2, y_2, a_2, b_2)$ , with

 $x_2 + iy_2 = \rho_3 e^{i\xi_3}, \ a_2 + ib_2 = \rho_2 e^{i\xi_2}, \ \xi_2 = \theta_2 - \theta_4, \ \xi_3 = \theta_3 - \theta_1 + k\theta_4.$ On  $U_3$  we put

 $x + i x = e^{i\chi_3}$   $x + i h = e^{i\chi_4}$ 

 $x_3 + iy_3 = \rho_3 e^{i\chi_3}, \quad a_3 + ib_3 = \rho_4 e^{i\chi_4}, \quad \chi_3 = \theta_3 - \theta_1 + k\theta_2, \quad \chi_4 = \theta_4 - \theta_2,$ 

and  $\omega = dx_3 \wedge dy_3 + da_3 \wedge db_3$ .

Finally, on  $U_4$  we set

$$x_4 + iy_4 = \rho_1 e^{i\zeta_1}, \quad a_4 + ib_4 = \rho_4 e^{i\zeta_4}, \quad \zeta_1 = \theta_1 - \theta_3 - k\theta_2, \quad \zeta_4 = \theta_4 - \theta_2,$$

and  $\omega = dx_4 \wedge dy_4 + da_4 \wedge db_4$ .

The normalized Hamiltonian function for  $\psi_t$  is  $f = \pi \rho_1^2 - \kappa$ , where  $\kappa$  is a constant determined by the condition  $\int_M f\omega^2 = 0$ . Straightforward calculations give

$$\int_{M} \omega^{2} = \mu (2\tau - k\mu), \text{ and } \int_{M} \pi \rho_{1}^{2} \omega^{2} = \frac{\mu^{2}}{3} (3\tau - 2k\mu).$$

So

(4.4) 
$$\kappa = \frac{\mu}{3} \left( \frac{3\lambda + k\mu}{2\lambda + k\mu} \right).$$

It is not easy to determine the transition function of det(TM) that corresponds to the coordinate transformation  $(x_i, y_i, a_i, b_i) \rightarrow (x_j, y_j, a_j, b_j)$ ; that is why we will introduce polar coordinate on subsets of the domains  $U_j$ .

Given  $0 < \epsilon \ll 1$ , for j = 1, 2, 3, 4 we put

$$B_j = \{ [z] \in U_j : |z_j| < 2\epsilon \}$$
 and  $B_0 = \{ [z] \in M : |z_j| > \epsilon \text{ for all } j \}.$ 

On  $B_0$  are well-defined the coordinates  $(\frac{\rho_1^2}{2}, \varphi_1, \frac{\rho_2^2}{2}, \varphi_2)$ , and in this coordinates

$$\omega = d\left(\frac{\rho_1^2}{2}\right) \wedge d\varphi_1 + d\left(\frac{\rho_2^2}{2}\right) \wedge d\varphi_2.$$

On  $B_j$  (j = 1, 2, 3, 4) we consider the Darboux coordinates  $(x_j, y_j, a_j, b_j)$  defined above. Then  $B_0, B_1, B_2, B_3, B_4$  is a Darboux atlas for M. We assume that M is endowed with the orientation given by  $\omega^2$ . This orientation agrees on  $B_0$  with the one defined by  $d\rho_1^2 \wedge d\varphi_1 \wedge d\rho_2^2 \wedge d\varphi_2$ .

It is evident that  $\psi_t(B_i) = B_i$ , for i = 0, 1, 2, 3, 4. Since  $\psi_t$  on  $B_0$  is simply the translation  $\varphi_1 \to \varphi_1 + 2\pi t$  of the variable  $\varphi_1$ , the Maslov index  $J_0$  of  $\psi$  in the trivialization defined on  $B_0$  vanishes.

As  $B_j$  (for j = 1, 2, 3, 4) has "infinitesimal size" and  $J_0 = 0$ , the expression for  $I_{\psi}$  of Theorem 3 can be written

(4.5) 
$$I_{\psi} = \sum_{i < k} N_{ik} + O(\epsilon)$$

Since  $I_{\psi}$  is obviously independent of the coordinates, it follows from (4.5) that  $N_{ik}$  is independent, up to order  $\epsilon$ , of the chosen Darboux coordinates in  $B_j$ , for j = 1, 2, 3, 4. Moreover  $N_{ik}$  with  $0 \neq i < k$  is also of order  $\epsilon$ .

On the other hand, if we substitute  $B_i$  by

$$B'_j = \{ [z] \in B_j : |z_r| > \epsilon, \ r \neq j \}$$

in the definition of  $N_{ik}$  (see Theorem 3) the new  $N_{ik}$  differs from the old one in a quantity of order  $\epsilon$ . As on  $B'_1$  the variable  $\rho_2 \neq 0$ , we can consider the Darboux coordinates

$$(x_1, y_1, \frac{\rho_2^2}{2}, \varphi_2)$$

on  $B'_1$ . Since  $\rho_3 \neq 0$  on  $B'_2$  we take the coordinates  $(a_2, b_2, \frac{\rho_3^2}{2}, \xi_3)$  on  $B'_2$ . Similarly we will adopt the following coordinates:  $(x_3, y_3, \frac{\rho_4^2}{2}, \chi_4)$  on  $B'_3$  and  $(\frac{\rho_1^2}{2}, \zeta_1, a_4, b_4)$  on  $B'_4$ .

Taking into account the preceding arguments

(4.6) 
$$I_{\psi} = \sum_{j=1}^{4} N'_{0j} + O(\epsilon)$$

where

(4.7) 
$$N'_{0j} = \frac{i}{\pi} \int_{A'_{0j}} f d \log r_{0j} \wedge \omega$$

and

$$A'_{0j} = \{ [z] \in M : |z_r| > \epsilon, \text{ for all } r \neq j \text{ and } |z_j| = \epsilon \}.$$

The submanifold  $A'_{0j}$  is oriented as a subset of  $\partial B_0$ ; that is, with the orientation induced by the one of  $B_0$ .

Next we determine the value of  $N'_{01}$ . To know the transition function  $r_{01}$  one needs the Jacobian matrix R of the transformation

$$(x_1, y_1, \frac{\rho_2^2}{2}, \varphi_2) \to (\frac{\rho_1^2}{2}, \varphi_1, \frac{\rho_2^2}{2}, \varphi_2)$$

in the points of  $A'_{01}$ ; with  $\rho_1^2 = x_1^2 + y_1^2$ ,  $\varphi_1 = \tan^{-1}(y_1/x_1)$ . The non trivial block of R is the diagonal one

$$\begin{pmatrix} x_1 & y_1 \\ r & s \end{pmatrix},$$

with  $r = -y_1(x_1^2 + y_1^2)^{-1}$  and  $s = x_1(x_1^2 + y_1^2)^{-1}$ . The non-real eigenvalues of *R* are

$$\lambda_{\pm} = \frac{x_1 + s}{2} \pm \frac{i\sqrt{4 - (s + x_1)^2}}{2}.$$

On  $A'_{01}$  these non-real eigenvalues occur when  $(s + x_1)^2 < 2$ , that is, if  $|\cos \varphi_1| < 2\epsilon(\epsilon^2 + 1)^{-1} =: \delta$ . If  $y_1 > 0$  then  $\lambda_-$  of the first kind (see [10]) and  $\lambda_+$  is of the first kind, if  $y_1 < 0$ .

Hence, on  $A'_{01}$ ,

$$\rho(R) = \begin{cases} \lambda_+ |\lambda_+|^{-1} = x + iy, & \text{if } |\cos \varphi_1| < \delta \text{ and } y_1 < 0; \\ \lambda_- |\lambda_-|^{-1} = x - iy, & \text{if } |\cos \varphi_1| < \delta \text{ and } y_1 > 0; \\ \pm 1, & \text{otherwise.} \end{cases}$$

where  $x = \delta^{-1} \cos \varphi_1$ , and  $y = \sqrt{1 - x^2}$ .

14

If we put  $\rho(R) = e^{i\gamma}$ , then  $\cos \gamma = \delta^{-1} \cos \varphi_1$  (when  $|\cos \varphi_1| < \delta$ ), and

$$\sin \gamma = \begin{cases} -\sqrt{1 - \cos^2 \gamma}, & \text{if } \sin \varphi_1 > 0; \\ \sqrt{1 - \cos^2 \gamma}, & \text{if } \sin \varphi_1 < 0. \end{cases}$$

So when  $\varphi_1$  runs anticlockwise from 0 to  $2\pi$ ,  $\gamma$  goes round clockwise the circumference; that is,  $\gamma = h(\varphi_1)$ , where h is a function such that

(4.8) 
$$h(0) = 2\pi$$
, and  $h(2\pi) = 0$ .

As  $r_{01} = \rho(R)$ , then  $d \log r_{01} = idh$ .

On  $A'_{10}$  the form  $\omega$  reduces to  $(1/2)d\rho_2^2 \wedge d\varphi_2$ . From (4.7) one deduces

(4.9) 
$$N'_{01} = \frac{i}{2\pi} \int_{A'_{01}} if dh \wedge d\rho_2^2 \wedge d\varphi_2.$$

On the other hand according to the convention about orientations,  $\{[z] : |z_1| = \epsilon\}$  as subset of  $\partial B_0$  is oriented by  $-d\varphi_1 \wedge d\rho_2^2 \wedge d\varphi_2$ . And on  $A'_{01}$  the Hamiltonian function  $f = -\kappa + O(\epsilon)$ . Then it follows from (4.9) together with (4.8)

(4.10) 
$$N_{01}' = 2\tau\kappa + O(\epsilon)$$

The contributions  $N'_{02}$ ,  $N'_{03}$ ,  $N'_{04}$  to  $I_{\psi}$  can be calculated in a similar way. One obtains the following results up to addends of order  $\epsilon$ :

(4.11) 
$$N'_{02} = 2\mu\kappa - \mu^2, \ N'_{03} = 2\lambda(\kappa - \mu), \ N'_{04} = \mu(2\kappa - \mu).$$

As  $I_{\psi}$  is independent of  $\epsilon$ , it follows from (4.6), (4.10), (4.11) and (4.4)

$$I_{\psi} = \frac{2k\mu^2}{3} \left( 1 - \frac{\mu}{2\lambda + k\mu} \right).$$

Next we consider the loop  $\tilde{\psi}$ ; the corresponding normalized Hamiltonian function is  $\tilde{f} = \pi \rho_2^2 - \tilde{\kappa}$ , where

(4.12) 
$$\tilde{\kappa} = \frac{3\lambda^2 + 3k\lambda\mu + k^2\mu^2}{3(2\lambda + k\mu)}$$

As in the preceding case

(4.13) 
$$I_{\tilde{\psi}} = \sum_{j=1}^{4} \tilde{N}'_{0j} + O(\epsilon)$$

where

$$\tilde{N}'_{0j} = \frac{i}{\pi} \int_{A'_{0j}} \tilde{f}d \, \log r_{0j} \wedge \omega.$$

The expression for  $\tilde{N}'_{01}$  can be obtained from (4.9) substituting f for  $\tilde{f}$ ; so

(4.14) 
$$N'_{01} = \tau(2\tilde{\kappa} - \tau) + O(\epsilon)$$

Analogous calculations give the following values for the  $\tilde{N}'_{0j}{}^{\prime}{\rm s},$  up to summands of order  $\epsilon$ 

(4.15)  $\tilde{N}'_{02} = 2\mu\tilde{\kappa}, \quad \tilde{N}'_{03} = \lambda(2\tilde{\kappa} - \lambda), \quad \tilde{N}'_{04} = \mu(2\tilde{\kappa} - k\mu - 2\lambda).$ 

From (4.12), (4.13), (4.14) and (4.15) it follows the value for  $I_{\tilde{\psi}}$  given in the statement of Theorem.

15

## ANDRÉS VIÑA

*Remark.* In [2] is proved that  $\pi_1(\text{Ham}(M)) = \mathbb{Z}$  when k = 1, therefore the quotient of  $I_{\psi}$  by  $I_{\psi'}$ , for arbitrary Hamiltonian loops of symplectomorphisms, is a rational number. For the particular loops considered in Theorem 8 the quotient  $I_{\psi}/I_{\psi}$  equals -k/2, so Theorem 8 is consistent with the result of Abreu and McDuff.

#### References

- [1] Abraham, R. and Marsden, J. Foundations of mechanics. Addison-Wesley, Reading. 1985
- [2] Abreu, M. and McDuff, D. Topology of symplectomorphism groups of rational ruled surfaces. J. Amer. Math. Soc. 13, 971-1009 (2000).
- [3] Audin, M. Torus Actions on symplectic manifolds. Birkhäuser, Basel, 2004
- [4] Guillemin, V. Moment maps and combinatorial invariants of Hamiltonian  $T^n\mbox{-spaces.}$  Birkhäuser, Boston, 1994
- [5] Guillemin, V., Lerman, L. and Sternberg, S. Symplectic fibrations and multiplicity diagrams. Cambridge U.P., Cambridge, 1996
- [6] Lalonde, F., McDuff, D. and Polterovich, L. Topological rigidity of Hamiltonian loops and quantum homology. Invent. Math 135, 369-385 (1999)
- [7] McDuff, D. and Salamon, D., Introduction to symplectic topology, Clarenton Press, Oxford, 1998
- [8] Polterovich, L., Hamiltonian loops and Arnold's principle. Amer. Math. Soc. Transl. (2) 180, 181-187 (1997)
- [9] Polterovich, L., The geometry of the group of symplectic diffeomorphisms, Birkhäuser, Basel, 2001
- [10] Salamon, D. and Zehnder, E. Morse theory for periodic solutions of Hamiltonian systems and the Maslov index. Comm. Pure Appl. Math XLV, 1303-1360 (1992)

Departamento de Física. Universidad de Oviedo. Avda Calvo Sotelo. 33007 Oviedo. Spain.

*E-mail address*: vina@uniovi.es