

# EXTENSIONS OF THE REPRESENTATION MODULES OF A PRIME ORDER GROUP

V. A. BOVDI, V. P. RUDKO

*Dedicated to Professor Kálmán Györy on his 65th birthday*

ABSTRACT. For the ring  $R$  of integers of a ramified extension of the field of  $p$ -adic numbers and a cyclic group  $G$  of prime order  $p$  we study the extensions of the additive groups of  $R$ -representations modules of  $G$  by the group  $G$ .

Let  $\mathfrak{F}$  be the field of fractions of a principal ideal domain  $R$ ,  $F$  a field which contains  $R$ , let  $G$  be a finite group and  $\Gamma$  a matrix  $R$ -representation of  $G$ . Let  $M$  be an  $RG$ -module, which affords the  $R$ -representation  $\Gamma$  of  $G$ , and  $FM = F \otimes_R M$  the smallest linear space over  $F$  which contains  $M$  and  $\widehat{M} = FM^+/M$ , the factor group of the additive group of the space  $FM$  by the additive group of  $M$ . Clearly, the group  $\widehat{M}$  and the space  $FM$  are  $RG$ -modules. Put  $\widehat{F} = F^+/R$ .

Let  $\mathfrak{f} : G \rightarrow \widehat{M}$  be a 1-cocycle of  $G$  with value in  $\widehat{M}$ , i.e.

$$\mathfrak{f}(xy) = x\mathfrak{f}(y) + \mathfrak{f}(x), \quad (x, y \in G).$$

Define  $[g, x]$  by  $\begin{pmatrix} g & x \\ 0 & 1 \end{pmatrix}$  and set

$$\mathfrak{C}\mathfrak{r}\mathfrak{n}\mathfrak{s}(G, M, \mathfrak{f}) = \{ [g, x] \mid g \in G, \quad x \in \mathfrak{f}(g) \},$$

where  $x$  runs over the cosets  $\mathfrak{f}(g) \in \widehat{M}$  for any  $g \in G$ .

Clearly,  $\mathfrak{C}\mathfrak{r}\mathfrak{n}\mathfrak{s}(G, M, \mathfrak{f})$  is a group, where the multiplication is the usual matrix multiplication. Of course  $K_1 = \{[e, x] \mid e \text{ is the unit element of } G, \quad x \in \mathfrak{f}(e)\}$  is a normal subgroup of  $\mathfrak{C}\mathfrak{r}\mathfrak{n}\mathfrak{s}(G, M, \mathfrak{f})$  such that  $K_1 \cong M^+$  and  $\mathfrak{C}\mathfrak{r}\mathfrak{n}\mathfrak{s}(G, M, \mathfrak{f})/K_1 \cong G$ . The group  $\mathfrak{C}\mathfrak{r}\mathfrak{n}\mathfrak{s}(G, M, \mathfrak{f})$  is an extension of the additive group of the  $RG$ -module  $M$  by  $G$ .

We are using the terminology of the theory of group representations [1].

A 1-cocycle  $\mathfrak{f} : G \rightarrow \widehat{M}$  is called *coboundary*, if there exists an  $x \in FM$  such that  $\mathfrak{f}(g) = (g - 1)x + M$  for every  $g \in G$ . The 1-cocycles  $\mathfrak{f}_1 : G \rightarrow \widehat{M}$  and

---

1991 *Mathematics Subject Classification*. Primary 16U60, 16W10.

The research was supported by OTKA No.T 037202 and No.T 038059

$f_2 : G \rightarrow \widehat{M}$  are called *cohomologous* if  $f_1 - f_2$  is a coboundary. Let  $H^1(G, \widehat{M})$  be the first cohomology group. Clearly, each element of  $H^1(G, \widehat{M})$  defines a class of equivalence of groups.

If the 1-cocycles  $f_1, f_2$  are cohomologous, then  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f_1)$  and  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f_2)$  are isomorphic. This isomorphism is called *equivalence* and these groups are called equivalent. In particular, the group  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f)$  is split (i.e.  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f) = M \rtimes G$ ) if and only if  $f$  is coboundary.

The *dimension* of the group  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f)$  is called the  $R$ -rank of the  $R$ -module  $M$ . (Note that  $M$  is a free  $R$ -module of finite rank.) The group  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f)$  is called *irreducible (indecomposable)*, if  $M$  is an irreducible (indecomposable)  $RG$ -module and the 1-cocycle  $f$  is not cohomologous to zero.

The group  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f)$  is *non-split*, if the 1-cocycle  $f$  defines a nonzero element of  $H^1(G, \widehat{M})$ .

Note that the properties of the group  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, f)$  were studied in [5,6,8], in the cases when  $R$  is either the ring of rational integers  $\mathbb{Z}$ , or the  $p$ -adic integers  $\mathbb{Z}_p$ , or the localization  $\mathbb{Z}_{(p)}$  of  $\mathbb{Z}$  at  $p$ .

Let  $G = \langle a \mid a^p = 1 \rangle$  be the cyclic group of prime order  $p$ ,  $R$  the ring of integers of the ramified finite extension  $T$  of the field of  $p$ -adic numbers. We calculate the group  $H^1(G, \widehat{M})$  for some module  $M$  of an indecomposable  $R$ -representation of  $G$ .

Let  $\Phi_p(x) = x^{p-1} + \dots + x + 1$  be a cyclotomic polynomial of degree  $p$  and let  $\eta(x)$  be a divisor of  $\Phi_p(x)$  over the field  $\mathfrak{F}$  with  $\deg(\eta(x)) < p - 1$  (provided that such nontrivial polynomial exists).

**Lemma 1.** *Let  $M_1$  and  $M_2$  be  $RG$ -modules which afford an  $R$ -representation  $\Gamma$  of  $G = \langle a \mid a^p = 1 \rangle$ .*

(i) *If  $M_1 \cong M_2$  then  $H^1(G, \widehat{M}_1) \cong H^1(G, \widehat{M}_2)$ .*

(ii) *If the matrix  $\Gamma(a)$  does not have 1 as eigenvalue, then  $H^1(G, \widehat{M}_1)$  is trivial.*

*Proof.* See [1].  $\square$

**Theorem 1.** *Let  $G = \langle a \mid a^p = 1 \rangle$  and  $M_\eta = \eta(a)RG$ . Then the  $RG$ -module  $M_\eta$  is indecomposable and*

$$H^1(G, \widehat{M}_\eta) \cong R/(\eta(1)R),$$

where  $R/(\eta(1)R)$  is the additive group of the factor ring of  $R$  by the ideal  $\eta(1)R$ .

*Proof.* Let  $t \in R$  be a prime element and  $\overline{R} = R/(tR)$ . Then in  $\overline{R}$  we have that

$$x^p - 1 = (x - 1)^p; \quad \eta(x) = (x - 1)^n; \quad (x^p - 1)\eta^{-1}(x) = (x - 1)^{p-n}, \quad (1)$$

where  $n = \deg(\eta(x))$ .

Put  $\eta_1(x) = (x^p - 1)\eta^{-1}(x)$ . Then  $M_\eta$  and  $RG/(\eta_1(a)RG)$  are isomorphic as  $RG$ -modules. If  $\overline{M}_\eta = M_\eta/(tM_\eta)$ , then by (1) follows that  $\overline{M}_\eta$  is a root subspace of the linear operator  $a$  over  $\overline{R}$ . It is easy to see that  $\overline{M}_\eta$  is not decomposable into a direct sum of invariant subspaces. It follows that  $M_\eta$  is an indecomposable

$RG$ -module. Clearly  $FM_\eta = \eta(a)FG = \eta(a)F + (a-1)FM_\eta$  and the group  $\widehat{M}_\eta = FM_\eta^+/M_\eta$  is isomorphic to a direct sum of groups  $\eta(a)(F^+/R) + (a-1)\widehat{M}_\eta$ . This means that in the class of 1-cocycles there is a cocycle  $\mathfrak{f} : G \rightarrow \widehat{M}_\eta$  such that

$$\mathfrak{f}(a) = \lambda\eta(a) + M_\eta, \quad (\lambda \in F).$$

Moreover, from  $\mathfrak{f}(a^p) = 0$  (in  $\widehat{M}_\eta$ ) it follows that if  $\omega = a^{p-1} + \dots + a + 1$ , then

$$\omega \cdot \mathfrak{f}(a) = \lambda \cdot \eta(1)\omega \in M_\eta$$

if and only if  $\lambda\eta(1) \in R$ . Therefore,  $H^1(G, \widehat{M}_\eta)$  is isomorphic to the subgroup  $\{\lambda + R \mid \lambda \in F, \lambda \cdot \eta(1) \in R\}$  of  $F/R$  and

$$\{\lambda + R \in F/R \mid \lambda \cdot \eta(1) \in R\} \cong R/(\eta(1)R).$$

□

**Corollary 1.** *Let  $\alpha \in R$  and  $\eta(1)R = t^s R$ , where  $t$  is a prime element of  $R$ . Put*

$$K_\alpha(G, M_\eta) = \left\langle \begin{pmatrix} e & m \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} a & \alpha t^{-s} \eta(a) \\ 0 & 1 \end{pmatrix} \mid m \in M_\eta \right\rangle,$$

where  $\alpha$  runs over the representative elements of the cosets of  $R/(t^s R)$ . Up to equivalence, the groups  $K_\alpha(G, M_\eta)$  give all extensions of the additive group of the  $RG$ -module  $M_\eta$  by the group  $G$ .

Suppose  $p = t^d \theta$ , where  $d > 1$  is the ramification index and  $\theta$  is a unit in  $R$ . Set

$$\mathfrak{X}_{ji} = t^j RG + (a-1)^i RG, \quad (1 \leq j < d, 1 \leq i < p).$$

**Theorem 2.** *The module  $\mathfrak{X}_{ji}$  is an  $RG$ -module affording an indecomposable  $R$ -representation of  $G$  and*

$$H^1(G, \widehat{\mathfrak{X}}_{ji}) \cong R/(t^{d-j}R).$$

*Proof.* Suppose that the  $RG$ -module  $\mathfrak{X}_{ji}$  is decomposable into a direct sum of  $RG$ -submodules. Then  $t^j = u_1 + u_2$ , where  $u_1, u_2$  are nonzero elements of  $RG$  with  $u_1 u_2 = 0$ . Thus  $e_1 = t^{-j} u_1$  is an idempotent. Since the trace  $tr(e_1)$  of  $e_1$  is a rational number (see Theorem 3.5, [4] p.21) of the form  $rp^{-1}$  ( $1 \leq r \leq p$ ), we get  $t^j r p^{-1} \in R$ , which is impossible for  $j < d$ . This contradiction proves the indecomposability of  $\mathfrak{X}_{ji}$ . Clearly  $F\mathfrak{X}_{ji} = FG = F + (a-1)FG$ . Therefore, in each class of 1-cocycles there is a cocycle  $\mathfrak{f} : G \rightarrow \widehat{\mathfrak{X}}_{ji}$  such that  $\mathfrak{f}(a) = \lambda + \mathfrak{X}_{ji}$ , where  $\lambda \in F$  with  $\lambda\omega \in \mathfrak{X}_{ji}$ . It follows that  $\lambda p = t^j \alpha$ , where  $\alpha \in R$  and

$$H^1(G, \widehat{\mathfrak{X}}_{ji}) \cong \{\lambda + R \mid \lambda \in F, \lambda t^{d-j} \in R\}$$

is a subgroup of  $F/R$ .

□

Set

$$K_\alpha(G, \mathfrak{X}_{ji}) = \left\langle \begin{pmatrix} e & m \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} a & \alpha t^{j-d} \\ 0 & 1 \end{pmatrix} \mid m \in \mathfrak{X}_{ji} \right\rangle,$$

where  $\alpha$  runs over the representative elements of the cosets of  $R/(t^{d-j}R)$ .

**Corollary 2.** *The groups  $K_\alpha(G, \mathfrak{X}_{ji})$  give all extensions of the additive group of the  $RG$ -module  $\mathfrak{X}_{ji}$  by  $G$ .*

**Lemma 2.** *The set  $\{\mathfrak{X}_{ji} \mid j = 1, \dots, d-1; i = 1, \dots, \frac{p-1}{2}\}$  consists of pairwise non-isomorphic modules.*

*Proof.* Let us consider an indecomposable  $\overline{RG}$ -module  $V_i = \overline{RG}/((a-1)^i \overline{RG})$ , where  $\overline{R} = R/(tR)$  and  $1 \leq i \leq p$ . It is easy to check that the elements

$$u_1 = t^j, \dots, u_i = t^j(a-1)^{i-1}, \quad u_{i+1} = (a-1)^i, \dots, u_p = (a-1)^{p-1} \quad (2)$$

form an  $R$ -basis in  $\mathfrak{X}_{ji}$  and

$$\Phi_p(x) - (x-1)^{p-1} = p\theta(x), \quad (3)$$

where  $\theta(x) \in \mathbb{Z}[x]$ ,  $\deg(\theta(x)) \leq p-2$ . Note that since  $\theta(1) = 1$ , it follows that  $\theta(a)$  is a unit in the group ring  $RG$ . Using the identity

$$xy - 1 = (x-1)(y-1) + (x-1) + (y-1),$$

from (3) we obtain that

$$(a-1)^p = p(a-1) \cdot (\alpha_0 + \alpha_1(a-1) + \dots + \alpha_{p-2}(a-1)^{p-2}), \quad (4)$$

where  $\alpha_0, \alpha_1, \dots, \alpha_{p-2} \in \mathbb{Z}$ . Since  $p = t^d \theta = t(t^{d-1} \theta)$ , from (4) we get

$$(a-1)^p = (a-1)u_p = tm, \quad (5)$$

where  $m \in \mathfrak{X}_{ji}$ . According to (2)  $(a-1)u_i = t^j u_{i+1}$ , and from (5) we obtain that the  $RG$ -module  $\overline{\mathfrak{X}}_{ji} = \mathfrak{X}_{ji}/(t\mathfrak{X}_{ji})$  is isomorphic to a direct sum  $V_i \oplus V_{p-i}$  of indecomposable  $\overline{RG}$ -modules, so by Theorem 2 and Lemma 1 the proof is complete.  $\square$

Let  $n > 1$  be the degree of a divisor of  $\Phi_p(x)$ , which is irreducible over  $R$ . We consider the following  $RG$ -modules:

$$\mathfrak{U}_{ji} = t^j(a-1)RG + (a-1)^{s+1}RG, \quad (1 \leq j < d, \quad 1 \leq s < n).$$

It is easy to check that the  $RG$ -module  $\mathfrak{U}_{ji}$  satisfies the condition (ii) of Lemma 1, so  $H^1(G, \widehat{\mathfrak{U}}_{ji}) = 0$ .

Let  $\mathfrak{Z}_{js}$  be a submodule of the free module  $RG^{(2)} = \{(x, y) \mid x, y \in RG\}$  of rank 2, which consists of the solutions  $(x, y)$  of the equality

$$t^j(a-1)x + (a-1)^{s+1}y = 0. \quad (6)$$

**Lemma 3.** *Let  $\omega = \Phi_p(a)$  and set  $u_1 = [0, \omega]$ ,  $u_2 = [(a-1)^s, -t^j]$  and  $u_3 = [t^{-j}(\omega - (a-1)^{p-1}), (a-1)^{p-s-1}]$ . Then  $\mathfrak{Z}_{j_s}$  is an  $RG$ -module generated by  $u_1, u_2, u_3$ .*

*Proof.* Clearly,  $u_1, u_2, u_3 \in \mathfrak{Z}_{j_s}$ . Let  $u = [x, y]$  be an arbitrary element of  $\mathfrak{Z}_{j_s}$ . If  $x = 0$  then  $u = u_1$ . Suppose  $x \neq 0$ . By subtraction of the elements of  $RGu_3$  from  $u$  we obtain that  $y = \gamma_0 + \gamma_1(a-1) + \cdots + \gamma_{p-s-2}(a-1)^{p-s-2}$  ( $\gamma_r \in R$ ). By (6)

$$t^j(a-1)x + (\gamma_0 + \gamma_1(a-1) + \cdots + \gamma_{p-s-2}(a-1)^{p-s-2}) \cdot (a-1)^{s+1} = 0,$$

which is possible if and only if  $\gamma_0 \equiv \cdots \equiv \gamma_{p-s-2} \equiv 0 \pmod{t^j}$ . Now, since  $u$  is an element of  $RGu_2$ , we obtain that  $y = 0$ . Then  $t^j(a-1)x = 0$  which implies  $x = \alpha\omega$  ( $\alpha \in R$ ) and  $u = \alpha(t^j u_3 - (a-1)^{p-s-1} u_2)$ . □

**Theorem 3.** *The  $RG$ -module  $\mathfrak{Z}_{j_s}$  is indecomposable. Moreover,*

$$H^1(G, \widehat{\mathfrak{Z}}_{j_s}) \cong R/(t^d R) \oplus R/(t^{d-j} R)$$

and the  $RG$ -modules  $\mathfrak{X}_{j_s}$  are pairwise non-isomorphic.

*Proof.* It is easy to see that

$$u_1 = t^j(a-1), \dots, u_{i-1} = t^j(a-1)^{i-1}, \quad u_i = (a-1)^i, \dots, u_{p-1} = (a-1)^{p-1}$$

form an  $R$ -basis in the  $RG$ -module  $\mathfrak{U}_{j_s}$  and

$$\overline{\mathfrak{U}}_{j_s} = \mathfrak{U}_{j_s}/(t\mathfrak{U}_{j_s}) \cong V_s \oplus V_{p-s-1}.$$

Since  $s < n$ , it follows that the  $RG$ -module  $\mathfrak{U}_{j_s}$  is indecomposable. Moreover, it follows that the  $RG$ -modules  $\mathfrak{U}_{j_s}$  are pairwise non-isomorphic and  $RG$ -modules  $\mathfrak{Z}_{j_s}$ ,  $\mathfrak{U}_{j_s}$  and  $RG^2$  form an exact sequence

$$0 \rightarrow \mathfrak{Z}_{j_s} \rightarrow RG^{(2)} \rightarrow \mathfrak{U}_{j_s} \rightarrow 0.$$

Therefore,  $\mathfrak{Z}_{j_s}$  is the kernel of a minimal projective covering of the indecomposable  $RG$ -module  $\mathfrak{U}_{j_s}$ , so  $\mathfrak{Z}_{j_s}$  is also indecomposable.

**Lemma 4.** *Let  $\widehat{\mathfrak{Z}}_{j_s} = (F\mathfrak{Z}_{j_s})^+/\mathfrak{Z}_{j_s}$ ,  $\widehat{F} = F^+/R$  and  $M = (a-1)\widehat{\mathfrak{Z}}_{j_s}$ . Then*

$$\widehat{\mathfrak{Z}}_{j_s}/M = \widehat{F}\nu_1 + \widehat{F}\nu_2,$$

where  $\nu_1 = [0, \omega] + M$ ,  $\nu_2 = [\omega, 0] + M$  and  $a\nu_1 = \nu_1$ ,  $a\nu_2 = \nu_2$ .

*Proof.* Clearly,  $ax = x$  ( $x \in \widehat{\mathfrak{Z}}_{j_s}/M$ ) and  $\widehat{F}\nu_1 = \widehat{F}[0, \omega] + M \in \widehat{\mathfrak{Z}}_{j_s}/M$ . Moreover,

$$\begin{aligned} \omega\widehat{F}u_3 + M &= \widehat{F}[t^{-1}p\omega, 0] + M = \widehat{F}(tp^{-1})[t^{-1}p\omega, 0] + M \\ &= \widehat{F}[\omega, 0] + M. \end{aligned}$$

By analogy

$$\omega \widehat{F}u_2 + M = \widehat{F}[0, -t^j \omega] + M = \widehat{F}[0, \omega] = \widehat{F}\nu_1.$$

Therefore  $\widehat{\mathfrak{Z}}_{js}/M = \widehat{F}\nu_1 + \widehat{F}\nu_2$ .  $\square$

From Lemma 4 it follows that each class of 1-cocycles of the group  $G$  with values in the group  $\widehat{\mathfrak{Z}}_{js} = (F\mathfrak{Z}_{js})^+/\mathfrak{Z}_{js}$  contains a 1-cocycle  $\mathfrak{f}$  such that

$$\mathfrak{f}(a) = \alpha[0, \omega] + \beta[\omega, 0] + \mathfrak{Z}_{js},$$

where  $\alpha, \beta \in F$  and  $\omega(\alpha[0, \omega] + \beta[\omega, 0]) \in \mathfrak{Z}_{js}$ . This condition holds if and only if  $\alpha p, \beta p \in R$ . Moreover

$$\alpha[0, \omega] + \beta[\omega, 0] \in \mathfrak{Z}_{js} + (a-1)\widehat{\mathfrak{Z}}_{js}$$

if and only if  $\alpha \in R$  and  $\beta \in t^{-j}R$ . Using properties of the 1-cocycle  $\mathfrak{f}$  it is easy to show that the two 1-cocycles  $\mathfrak{f}_j$  ( $j = 1, 2$ ):

$$\mathfrak{f}_1(a) = \alpha_1[0, \omega] + \beta_1[\omega, 0] + \mathfrak{Z}_{js}, \quad \mathfrak{f}_2(a) = \alpha_2[0, \omega] + \beta_2[\omega, 0] + \mathfrak{Z}_{js}$$

are cohomologous if and only if

$$p\alpha_1 \equiv p\alpha_2 \pmod{t^d} \quad \text{and} \quad p\beta_1 \equiv p\beta_2 \pmod{t^{d-j}},$$

where  $\alpha_j, \beta_j \in F$ ,  $p\alpha_j, p\beta_j \in R$ . Note that  $p = t^d\theta$ .

It follows that the map  $\mathfrak{f} \mapsto (p\alpha + t^d R, p\beta + t^{d-j} R)$  gives the isomorphism

$$H^1(G, \widehat{\mathfrak{Z}}_{js}) \cong R/(t^d R) \oplus R/(t^{d-j} R).$$

Therefore, according to (ii) of Lemma 1, the  $RG$ -modules  $\mathfrak{Z}_{js}$  ( $1 \leq j < d$ ) are pairwise non-isomorphic.  $\square$

Now, using the description of 1-cocycles it is easy to prove the following

**Corollary 3.** *Put*

$$K_{\alpha, \beta}(G, \mathfrak{Z}_{js}) = \left\langle \begin{pmatrix} e & m \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} a & \alpha t^{-d}[0, \omega] + \beta t^{-d}[\omega, 0] \\ 0 & 1 \end{pmatrix} \mid m \in \mathfrak{Z}_{js} \right\rangle,$$

where  $\alpha$  and  $\beta$  independently run over the representative elements of the cosets  $R/(t^d R)$  and  $R/(t^{d-j} R)$ , respectively. Up to equivalence, the groups  $K_{\alpha, \beta}(G, \mathfrak{Z}_{js})$  give all extensions of the additive group of the  $RG$ -module  $\mathfrak{Z}_{js}$  by the group  $G$ .

If  $R$  is the quadratic extension of the ring of  $p$ -adic integers, then the  $R$ -representations of  $G$  were described by P.M. Gudivok (see [7]).

Finally, we have the following result

**Theorem 4.** *Let  $\Phi_p(x)$  be decomposable into the product of at least two irreducible polynomials over  $R$ . Then the dimensions of the non-split indecomposable groups  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, \mathfrak{f})$  are unbounded.*

*Proof.* Let  $\Phi_p(x) = \eta_1(x) \cdots \eta_k(x)$  ( $k > 2$ ) be a decomposition into a product of polynomials irreducible over  $R$  and suppose that

$$\eta_1(x) = x^n - \alpha_{n-1}x^{n-1} - \cdots - \alpha_1x - \alpha_0 \in R[x].$$

Note that  $\deg(\eta_1(x)) = \deg(\eta_2(x)) = \cdots = \deg(\eta_k(x)) = n$  and  $kn = p - 1$ .

We will use the technic of the integral representation of finite groups, which was developed by S.D. Berman and P.M. Gudivok in [2, 3, 7].

Let  $\varepsilon$  be a primitive  $p$ th root of unity such that  $\eta_1(\varepsilon) = 0$  and let  $r_j$  be a natural number, such that  $\varepsilon_j = \varepsilon^{r_j}$  is a root of the polynomial  $\eta_j(x)$ , where  $r_1 = 1$  and

$j = 1, \dots, k$ . Let  $\tilde{\varepsilon} = \begin{pmatrix} 0 & \cdots & 0 & \alpha_0 \\ 1 & \cdots & 0 & \alpha_1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & \alpha_{n-1} \end{pmatrix}$  be the comparing matrix of  $\eta_1(x)$ .

The following  $R$ -representations of  $G = \langle a \mid a^p = 1 \rangle$  are irreducible:

$$\delta_0 : a \mapsto 1; \quad \delta_1 : a \mapsto \tilde{\varepsilon}; \quad \delta_j : a \mapsto \tilde{\varepsilon}_j = \tilde{\varepsilon}_j^{r_j}, \quad (j = 2, \dots, k).$$

Note that the module which affords representation  $\delta_1$  is  $R[\varepsilon]$  with  $R$ -basis  $1, \varepsilon, \dots, \varepsilon^{n-1}$ .

Let  $m \in \mathbb{N}$ . Define the following  $R$ -representation of  $G = \langle a \rangle$  of degree  $(3n+1)m$ :

$$\Gamma_m : a \mapsto \begin{pmatrix} \Delta_{1m}(a) & U_m(a) \\ 0 & \Delta_{2m}(a) \end{pmatrix},$$

where

$$\Delta_{1m}(a) = \delta_0^{(m)}(a) + \delta_1^{(m)}(a) = \begin{pmatrix} E_m \otimes \delta_0(a) & 0 \\ 0 & E_m \otimes \delta_1(a) \end{pmatrix};$$

$$\Delta_{2m}(a) = \delta_2^{(m)}(a) + \delta_3^{(m)}(a) = \begin{pmatrix} E_m \otimes \delta_2(a) & 0 \\ 0 & E_m \otimes \delta_3(a) \end{pmatrix};$$

$$U_m(a) = \begin{pmatrix} E_m \otimes u & J_m(1) \otimes u \\ E_m \otimes \bar{u} & E_m \otimes \bar{u} \end{pmatrix};$$

$u = (0, 0, \dots, 0, 1)$  defines a nonzero element of  $\text{Ext}(\delta_0, \delta_j)$ ;

$J_m(\lambda)$  is a Jordan block of degree  $m$  with  $\lambda$  in the main diagonal;

$\bar{u}$  is a matrix in which the first row is  $(0, \dots, 0, 1)$  and all other rows are zero. The matrix  $\bar{u}$  defines a nonzero element of the group  $\text{Ext}(\delta_1, \delta_j)$ , where  $j = 2, 3$ ;

$E_m$  is the unity matrix of degree  $m$ .

**Lemma 5.** (see [2,3])  $\Gamma_m$  is an indecomposable  $R$ -representation of  $G$ .

Let  $\mathfrak{W}_m = R^l$  be a module of  $l$ -dimension vectors over  $R$  affording the  $R$ -representation  $\Gamma_m$ . Put  $\widehat{F} = F^+/R$ ,  $\widehat{\mathfrak{W}}_m = F\mathfrak{W}_m^+/\mathfrak{W}_m$ . Clearly  $\widehat{F}^l \cong \widehat{\mathfrak{W}}_m$ . Define  $\tau : F \rightarrow F^n$  by

$$\tau(w) = w(\alpha_0, \alpha_0 + \alpha_1, \alpha_0 + \alpha_1 + \alpha_2, \dots, \alpha_0 + \dots + \alpha_{n-2}, 1), \quad (7)$$

where the  $\alpha_j$  are coefficients of  $\eta_1(x)$  and  $w \in F$ .

**Lemma 6.** (i) Each 1-cocycle of  $G = \langle a \mid a^p = 1 \rangle$  at  $\widehat{\mathfrak{W}}_m$  is cohomologous to a cocycle  $\mathfrak{f}$ , such that

$$\mathfrak{f}(a) = (X, 0, \dots, 0) + \mathfrak{W}_m,$$

where  $X \in F^m$  and  $pX = 0$  in  $\widehat{F}^m$  (i.e.  $pX \in R^m$ ).

(ii) Let  $z \in F^n$  such that  $(\tilde{\varepsilon} - E_n)z = 0$  in  $\widehat{F}^n$ . Then  $z = \tau(w) \pmod{R^n}$ , with  $w \in F$  such that  $\eta_1(1)w = 0$  in  $\widehat{F}$ .

(iii) If  $V = R/(\frac{p}{\eta(1)}R)$  is the residual of ring  $R$  by the ideal  $(\frac{p}{\eta(1)}R)$ , then  $H^1(G, \widehat{\mathfrak{W}}_n) \cong V^m$ .

*Proof.* (i) follows from (ii) of Lemma 1. (ii) is easy to check.

(iii) by (i) we can put  $\mathfrak{f}(a) = (X, 0, 0, 0)$  and  $\mathfrak{g}(a) = (Y, 0, 0, 0)$ , where  $X = (x_1, \dots, x_n)$ ,  $Y = (y_1, \dots, y_n)$  and  $pX = pY = 0$ . Note that all the equalities considered here are understood modulo the group  $R$ . Suppose that these 1-cocycles are cohomologous and  $Z \in F^l$  is such that

$$(\Gamma_m(a) - E_l)Z + \mathfrak{f}(a) = \mathfrak{g}(a). \quad (8)$$

Put  $Z = (Z_1, Z_2, Z_3, Z_4)$ , where  $Z_1 \in F^m$  and  $Z_2, Z_3, Z_4$  are  $m$ -dimensional vectors, with  $i$ -components belong to  $F^n$ , and denoted by  $Z_2^i, Z_3^i$  and  $Z_4^i$ , respectively. By (8) we get

$$(E_m \otimes u)Z_3 + (J_m \otimes u)Z_4 + X = Y, \quad (9)$$

$$(E_m \otimes (\tilde{\varepsilon} - E_n))Z_2 + (E_m \otimes \bar{u})(Z_3 + Z_4) = 0, \quad (10)$$

$$(E_m \otimes (\tilde{\varepsilon}_2 - E_n))Z_3 = 0, \quad (E_m \otimes (\tilde{\varepsilon}_3 - E_n))Z_4 = 0. \quad (11)$$

From (11) and by (ii) we have

$$Z_3 = (\tau(v_1), \dots, \tau(v_m)), \quad Z_4 = (\tau(u_1), \dots, \tau(u_m)), \quad (12)$$

where  $u_j, v_j \in F$ ,  $\tau$  is from (7) and

$$\eta_1(1)u_j = \eta_1(1)v_j = 0. \quad (13)$$

Clearly, the equality (10) consists of  $m$  matrix equalities of the form

$$(\tilde{\varepsilon} - E_n)Z_2^i + \bar{u}\tau(w) = 0, \quad (14)$$



where  $Z_2^i \in F^n$  is the  $i$ th component of  $Z_2$ ,  $i = 1, \dots, m$  and  $w \in F$ . Since  $u\tau(w) = w$  and  $\bar{u}\tau(w) = (w, 0, \dots, 0)$ , when all the rows of (14) are added together we obtain

$$-\eta(1)Z_2^n + w = 0, \quad (15)$$

where  $Z_2^n$  is the last component of the vector  $Z_2$ . According to (12) and (15), (10) gives the equalities

$$-\eta(1)z_j + v_j + u_j = 0, \quad (j = 1, \dots, m) \quad (16)$$

where  $z_j$  are some components of  $Z_2$ . From (9)

$$\begin{aligned} v_j + u_j + u_{j+1} + x_j &= y_j, & (j = 1, \dots, m-1) \\ v_m + u_m + x_m &= y_m, \end{aligned} \quad (17)$$

where  $X = (x_1, \dots, x_n)$ ,  $Y = (y_1, \dots, y_n)$  and  $pX = pY = 0$ . Multiplying (17) by  $\eta_1(1)$  and using (16) we obtain for the components of  $X$  and  $Y$

$$\eta_1(1)x_j = \eta_1(1)y_j \quad (j = 1, \dots, m). \quad (18)$$

Therefore, if the 1-cocycles  $\mathfrak{f}$  and  $\mathfrak{g}$  are cohomologous then (18) holds.

Conversely, suppose that (18) holds. Then it is not difficult to construct vectors  $Z_2, Z_3, Z_4$  that satisfy (9) and (10), which is equivalent to (8), i.e. the 1-cocycles  $\mathfrak{f}$  and  $\mathfrak{g}$  are cohomologous. It follows that by going from a cocycle to an element of the cohomology group, we need to change each component in  $X$  by  $\beta = \alpha \cdot p^{-1}$  modulo the group  $R$ , where  $\alpha \in R$ . Moreover, if  $\eta_1 \cdot \beta \in R$ , then must change  $\beta$  to 0.  $\square$

**Theorem 5.** *Let  $\varepsilon \in R$ , where  $\varepsilon^p = 1$  and  $p > 2$ . Then the description of the non-split indecomposable groups  $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, \mathfrak{f})$  is a wild type problem.*

*Proof.* For arbitrary matrices  $A, B \in M(m, R)$  the map

$$\Gamma_{A,B} : a \mapsto \begin{pmatrix} E & 0 & E & A & E \\ \varepsilon E & E & E & B & \\ & \varepsilon^2 E & 0 & 0 & \\ & & \varepsilon^3 E & 0 & \\ & & & \varepsilon^4 E & \end{pmatrix}$$

is an  $R$ -representation of  $G$  of degree  $l = 5m$ . The  $R$ -representations  $\Gamma_{A,B}$  and  $\Gamma_{A_1,B_1}$  are  $R$ -equivalent if and only if

$$C^{-1}AC \equiv A_1 \pmod{(1-\varepsilon)}, \quad C^{-1}BC \equiv B_1 \pmod{(1-\varepsilon)}$$

for some invertible matrix  $C$ . It follows that the description of the  $R$ -representations  $\Gamma_{A,B}$  of  $G$  is a wild type problem.

For the module affording the representation  $\Gamma_{A,B}$  of  $G$  we put  $R^l$ . Let  $X$  be an  $m$ -dimensional vector over  $F$  with  $pX \in R^m$ . Then there is a 1-cocycle  $\mathfrak{f}_X : G \rightarrow \widehat{R}^l$ , such that  $\mathfrak{f}_X(a) = (X, 0, \dots, 0) + R^l$ . The 1-cocycles  $\mathfrak{f}_X$  and  $\mathfrak{f}_Y$  are cohomologous if and only if

$$(1-\varepsilon)(X-Y) \in R^m.$$

Putting  $X = (p^{-1}, 0, \dots, 0)$  we obtain that  $H^1(G, \widehat{R}^l) \neq 0$ .  $\square$

## REFERENCES

1. Benson, D. J., *Representations and cohomology. Cohomology of groups and modules*, Cambridge Studies in Advanced Math., 31. Cambridge University Press, Cambridge, 1998.
2. Berman, S.D., *Representations of finite groups over an arbitrary field and over rings of integers.*, Izv. Akad. Nauk SSSR Ser. Mat. **30** (1966), 69–132. (Russian)
3. Berman, S.D.; Gudivok, P.M., *Indecomposable representation of finite group over the ring  $p$ -adic integers (Russian)*, Izvestia AN USSR **28(4)** (1964), 875–910.
4. Bovdi, A.A., *Group rings (Russian)*, Kiev UMK BO, 1988, pp. 156.
5. Bovdi, V.A., Gudivok, P.M., Rudko, V.P., *Torsion free groups with indecomposable holonomy group I*, J. Group Theory **5** (2002), 75–96.
6. Bovdi, V.A., Gudivok, P.M., Rudko, V.P., *Torsion-free crystallographic groups with indecomposable holonomy group.II*, J. Group Theory **7(4)** (2004), 555–569.
7. Gudivok, P.M., *Representation of finite groups over quadratic fields*, Doklady AN USSR **5** (2002), 75–96.
8. Kopcha, G.M., *Non-split extension of the indecomposable module of  $p$ -integer representation of the cyclic group of order  $p^2$  (Ukrainian)*, Uzhgorod University Scientific Herald. Math. Ser. **5** (2000), 49–56.

V.A. Bovdi

Institute of Mathematics, University of Debrecen

P.O. Box 12, H-4010 Debrecen, Hungary

Institute of Mathematics and Informatics, College of Nyíregyháza

Sóstói út 31/b, H-4410 Nyíregyháza, Hungary

E-MAIL: VBOVDI@MATH.KLTE.HU

V.P. Rudko

DEPARTMENT OF ALGEBRA, UNIVERSITY OF UZHGOROD

88 000, UZHGOROD, UKRAINE

E-MAIL: MATH1@UNIV.UZHGOROD.UA