

COHOMOLOGICAL DIMENSION WITH RESPECT TO NONABELIAN GROUPS

ATISH MITRA

ABSTRACT. Cencelj and Dranishnikov showed [1, 2, 3] that for certain nilpotent groups G , $K(G_{ab}, 1) \in \text{AE}(X)$ is equivalent to $K(G, 1) \in \text{AE}(X)$ for any compacta X (here G_{ab} is the abelianization of G). We examine the same problem for solvable groups. We also give an elementary proof of this fact for any nilpotent group and any 2-dimensional metric space.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Nilpotent Groups	3
4. Finite Solvable Groups	4
5. Extension Properties of the Pontryagin Disk	7
6. Infinite Solvable Groups	8
References	9

1. INTRODUCTION

We define cohomological dimension $\dim_G X$ of a topological space X with respect to an abelian group G as the largest number n such that there exists a closed subset $A \subset X$ with $\check{H}^n(X, A; G) \neq 0$. Several authors have worked on developing this theory for various classes of spaces like compacta and metrizable spaces. For surveys of the theory the reader is referred to [5], [8].

We use Kuratowski's notation $X \tau M$ if every map from a closed subset of X to M extends over X . The following theorem allows us to formulate an equivalent definition of cohomological dimension of compacta with respect to abelian groups in terms of extension of maps to Eilenberg-MacLane spaces.

Theorem 1.1. [6] *For any compactum X and abelian group G the following conditions are equivalent:*

$$(1) \dim_G X \leq n$$

Date: June 28, 2005 .

1991 Mathematics Subject Classification. 16B50, 18D35, 54C56 .

Key words and phrases. cohomological dimension, nilpotent group, solvable group.

- (2) $\check{H}^{n+1}(X, A; G) = 0$ for all closed $A \subset X$
- (3) $X\tau K(G, n)$

Trying to extend the notion of cohomological dimension to non-abelian groups G , one notices that the corresponding Eilenberg-MacLane spaces are defined only for $n = 1$. Dranishnikov and Repovš considered the case of perfect groups in [7]. Later Cencelj and Dranishnikov studied this notion in a series of three papers [1, 2, 3].

Generalizing Dranishnikov's theorem about extension of maps to simply-connected complexes [5], they obtained the following result.

Theorem 1.2. *For any nilpotent CW-complex M and finite-dimensional metric compactum X , the following are equivalent:*

- (1) $X\tau M$
- (2) $X\tau SP^\infty M$
- (3) $\dim_{H_i(M)} X \leq i$ for every $i > 0$
- (4) $\dim_{\pi_i(M)} X \leq i$ for every $i > 0$

Recall that $SP^i X = X^i / (\Sigma_i)$, where Σ_i is the symmetric group on i letters. If X is pointed, there is a natural embedding $SP^i X \rightarrow SP^{i+1} X$, and $SP^\infty X$ is defined as direct limit of the $SP^i X$.

In course of the proof of the above theorem, they obtained the following theorem.

Theorem 1.3. *For a nilpotent group N and every metric compactum X the following equivalence holds : $\dim_N X \leq 1 \Leftrightarrow \dim_{N_{ab}} X \leq 1$ provided N has one of the following properties:*

- (1) N is a torsion group.
- (2) for every prime p s.t. $Tor_p N \neq 1$
 - (a) N is not p -divisible, or
 - (b) $Tor_p N_{ab} \neq 0$.

In view of Theorem 1.3 it is worthwhile studying the following problem.

Question 1.4. *Let \mathcal{C} be a class of spaces. Describe groups for which the following are equivalent for any $X \in \mathcal{C}$?*

- (1) $X\tau K(G, 1)$
- (2) $X\tau K(G_{ab}, 1)$

In section 2 we list well known results which we use in the sequel. In section 3 we show that if \mathcal{C} is the class of all 2-dimensional metrizable spaces, then nilpotent groups satisfy question 1. In section 3 we show that if \mathcal{C} is the class of all 2-dimensional compacta, then finite solvable groups satisfy question 1.4. Finally we give an example showing that infinite solvable groups do not satisfy question 1.4 when if \mathcal{C} is the class of all 2-dimensional compacta.

2. PRELIMINARIES

We collect below some classical results which we will use in the sequel.

Proposition 2.1. *Let A be closed in X . A map $f : A \rightarrow Y$ has an extension to X iff some homotopic map $g : A \rightarrow Y$ has an extension.*

Recall that a map $f : X \rightarrow Y$ between CW complexes is called cellular if $f(X^{(n)}) \subset Y^{(n)}$ for all n .

Theorem 2.2 (Cellular Approximation Theorem). [11] *Every map $f : X \rightarrow Y$ between CW complexes is homotopic to a cellular map.*

Theorem 2.3 (Seifert - Van Kampen). *Let X_1, X_2 be path connected open subsets of X , and the intersection $X_0 = X_1 \cap X_2$ be path connected. Then the following commutative diagram of fundamental groups (based at some $x_0 \in X_0$) where all maps are induced by inclusions is a pushout diagram:*

$$\begin{array}{ccc} \pi_1(X_0) & \xrightarrow{i_1} & \pi_1(X_1) \\ i_2 \downarrow & & \downarrow j_1 \\ \pi_1(X_2) & \xrightarrow{j_2} & \pi_1(X) \end{array}$$

Theorem 2.4 (Bockstein exact sequence). [8] *If $1 \rightarrow G \rightarrow E \rightarrow \Pi \rightarrow 1$ is a short exact sequence of abelian groups, then there is a natural exact sequence*

$$\dots \rightarrow \check{H}^n(X, A; G) \rightarrow \check{H}^n(X, A; E) \rightarrow \check{H}^n(X, A; \Pi) \rightarrow \check{H}^{n+1}(X, A; G) \rightarrow \dots \quad (2.1)$$

for any paracompact space X and its closed subspace A . Here $\check{H}^n(X, A; G)$ is the n -th Čech cohomology group of the pair (X, A) with coefficients in G .

The following theorem due to Cenceli and Dranishnikov gives one of the implications of question 1 for any group G

Theorem 2.5. [1] *Let X be a compactum and G a group. Then $X\tau K(G, 1) \Rightarrow X\tau K(G_{ab}, 1)$.*

Proof: From [5], theorem 6 it follows that $X\tau SP^\infty K(G, 1)$. We use results from [4], [10] to see that $X\tau K(H_1(K(G, 1)), 1)$, which is equivalent to $X\tau K(G_{ab}, 1)$.

3. NILPOTENT GROUPS

We present a generalization of theorem 1.3 for 2-dimensional metrizable spaces.

Theorem 3.1. *Let G be a nilpotent group and X be a 2-dimensional metric space.*

Then $X\tau K(G, 1) \Leftrightarrow X\tau K(G_{ab}, 1)$.

Recall that the lower central series of a group G is $G = \gamma_1 G \geq \gamma_2 G \geq \dots$, where $\gamma_{i+1} G = [\gamma_i G, g]$. Notice that $\gamma_i G / \gamma_{i+1} G$ lies in the center of $G / \gamma_{i+1} G$ and that the nilpotent class of G equals the length of the lower central series of G . The following lemma shows that the first lower central factor G_{ab} exerts a strong influence on the subsequent lower central factors.

Lemma 3.2 (Robinson). *Let G be a group and let $F_i = \gamma_i G / \gamma_{i+1} G$. Then the mapping $a(\gamma_{i+1} G) \otimes g[G, G] \mapsto [a, g](\gamma_{i+2} G)$ is a well defined module epimorphism from $F_i \otimes_{\mathbb{Z}} G_{ab}$ to F_{i+1} .*

Proof: Let $g \in G$ and $a \in G_i$, and consider the function $(a(\gamma_{i+1} G), g[G, G]) \mapsto [a, g](\gamma_{i+2} G)$. Check that the mapping is well-defined and bilinear. By the universal property of tensor products, we have an induced homomorphism. It is easily checked that it is an epimorphism.

The above lemma gives the following useful corollary.

Corollary 3.3. *The \mathcal{P} be a group theoretical property which is inherited by images of tensor products (of abelian groups) and by extensions. If G is a nilpotent group such that G_{ab} has \mathcal{P} , then G has \mathcal{P} .*

Proof: Let $F_i = \gamma_i G / \gamma_{i+1} G$, and note that $F_1 = G_{ab}$. Suppose F_i has \mathcal{P} : then the previous lemma implies that F_{i+1} has \mathcal{P} . As G is nilpotent, some $\gamma_{c+1} = 1$. Since \mathcal{P} is closed under extensions, G has \mathcal{P} .

Proof of theorem 3.1: Let \mathcal{P} be the group theoretical property of "belonging to \mathcal{G} ". We know that if $1 \rightarrow G \rightarrow E \rightarrow \Pi \rightarrow 1$ is a short exact sequence and $X\tau K(G, 1)$ and $X\tau K(\Pi, 1)$, then $X\tau K(E, 1)$ [8]. We also know that if X is metrizable, then $X\tau K(G, 1) \Rightarrow X\tau K(G \otimes H, 1)$ for any abelian groups G, H [8]. From the Bockstein exact sequence we see that if X is 2-dimensional, G abelian and $X\tau K(G, 1)$, then $X\tau\phi(K(G, 1))$ for any homomorphism ϕ . Apply the previous corollary.

Remark 3.4. The theorem can be improved to the case where X has cohomological dimension 2.

4. FINITE SOLVABLE GROUPS

Our effort to answer question 1 for solvable groups gives the following theorem.

Theorem 4.1. *Let G be a finite solvable group and X be a 2-dimensional metrizable space. Then $X\tau K(G, 1) \Leftrightarrow X\tau K(G_{ab}, 1)$.*

For proving the above theorem, we need some lemmas.

Proposition 4.2. *Let X be a metrizable space and K_1, K_2 be subcomplexes of the CW complex $K_1 \cup K_2$. Then:*

- (1) *If K_1, K_2 and $K_0 = K_1 \cap K_2$ are absolute extensors of X , then so is $K_1 \cup K_2$.*
- (2) *If $K_1 \cup K_2$ and $K_0 = K_1 \cap K_2$ are absolute extensors of X , then so are K_1, K_2 .*

Proof: (1) Let $f : A \rightarrow K_1 \cup K_2$ be a map from a closed subset A of X . Define $C_i = f^{-1}(K_i)$ and note that $A = C_1 \cup C_2$ and $f(C_1 \cap C_2) \subset K_1 \cap K_2$. As the closure of any one of the sets $C_1 - C_2$ and $C_2 - C_1$ misses the other (and as X is completely normal), we can find U open in X such that $C_1 - C_2 \subset U \subset \bar{U} \subset X - (C_2 - C_1)$. Define $D_1 = \bar{U} \cup (C_1 \cap C_2)$ and $D_2 = (X - U) \cup (C_1 \cap C_2)$. Then we have $D_1 \cap A = C_1$, $D_2 \cap A = C_2$ and $D_1 \cup D_2 = X$. As $C_1 \cap C_2$ is closed in $D_1 \cap D_2$ (which is closed in X) and as $K_1 \cap K_2$ is absolute extensor of X , we can extend $f|_{C_1 \cap C_2}$ to $\bar{f} : D_1 \cap D_2 \rightarrow K_1 \cap K_2$. Now we have a map from $C_1 \cup (D_1 \cap D_2)$ to K_1 and a map from $C_2 \cup (D_1 \cap D_2)$ to K_2 , both of which we can extend and then paste the resulting maps to get the desired extension.

(2) Start with $f : A \rightarrow K_1$ and extend to $\bar{f} : X \rightarrow K_1 \cup K_2$. Calling $C_i = (\bar{f})^{-1}(K_i)$, we have $X = C_1 \cup C_2$, $A \subset C_1$ and $\bar{f}(C_1 \cap C_2) \subset K_1 \cap K_2$. Extend $\bar{f}|_{C_1 \cap C_2}$ to $\tilde{f} : C_2 \rightarrow K_1 \cap K_2$. Paste $\bar{f}|_{C_1}$ and $\tilde{f}|_{C_2}$ to get the desired extension.

Definition 4.3. Let G_1, G_2 and A be groups, and $f_i : A \rightarrow G_i$ be homomorphisms. Then the amalgamated product $G_1 *_A G_2$ is defined as $\frac{G_1 * G_2}{\langle f_1(a) = f_2(a), a \in A \rangle}$ (Here $\langle \rangle$ denotes the normal closure). Notice that the amalgamated product is a pushout of the following diagram:

$$\begin{array}{ccc} A & \xrightarrow{f_1} & G_1 \\ f_2 \downarrow & & \\ & & G_2 \end{array}$$

Proposition 4.4. *Let X be a 2-dimensional metrizable space and define $\mathcal{H}_X = \{G : X\tau K(G, 1)\}$.*

- (1) *If G_1, G_2 and A are in \mathcal{H}_X , then so is $G_1 *_A G_2$.*
- (2) *If $G_1 *_A G_2$ and A are in \mathcal{H}_X , then so are G_1 and G_2 .*

Proof: Note that a homomorphism $f : A \rightarrow G$ induces a map $f_* : K(A, 1)^{(2)} \rightarrow K(G, 1)^{(2)}$: we start by sending the generators to the corresponding loops in $K(G, 1)^{(1)}$, and then use the relators of A to extend the map to the 2-skeleton. Let M_{f_*} be the mapping cylinder of f_* : then we have $K(A, 1)^{(2)} \hookrightarrow M_{f_*}$. Apply the previous theorem with $K_1 = M_{f_*1}$, $K_2 = M_{f_*2}$ and $K_0 = K(A, 1)^{(2)}$.

We can now prove the following lemma, which we use later to see that any finite group with solvability index 2 is in \mathcal{G} .

Lemma 4.5. *Let X be a 2-dimensional metrizable space with $\dim_{\mathbb{Z}/p} X = 1$ and let K be a CW complex. Let $\alpha \in \pi_1(K)$ with $\alpha^p = 1$, for some prime p . Then $X\tau K \Leftrightarrow X\tau K \cup_{\alpha} D^2$, where D^2 is a 2-cell.*

Proof: Note that $\alpha : S^1 \rightarrow K$ induces a homomorphism $\alpha_* : \mathbb{Z} \rightarrow \pi_1(K)$ with $\alpha_*(1)^p = 1$, i.e. $\alpha_*(p) = 1$. Then $\pi_1(K \cup_{\alpha} D^2)$ is the pushout of

$$\begin{array}{ccc} \mathbb{Z} & \xrightarrow{\alpha_1} & \pi_1(K) \\ \downarrow & & \\ \{1\} & & \end{array}$$

But in this case it is also the pushout of

$$\begin{array}{ccc} \mathbb{Z}/p & \xrightarrow{\alpha_1|} & \pi_1(K) \\ \downarrow & & \\ \{1\} & & \end{array}$$

Then we can use the previous theorem.

The previous lemma has the following generalization:

Lemma 4.6. *Let X be a 2-dimensional metrizable space, $\mathcal{P} = \{p : \dim_{\mathbb{Z}/p} X = 1\}$ and let K be a CW complex. Let $\alpha_i \in \pi_1(K)$ with $\alpha_i^{p_i^{m_i}} = 1$, for some prime $p_i \in \mathcal{P}$ and positive integer m_i . Then $X\tau K \Leftrightarrow X\tau K \cup_{\alpha_i} D_i^2$, where D_i^2 is a 2-cell.*

Proof of theorem 4.1:

$A = G_{ab}$ is a finite abelian group and hence of the form $\mathbb{Z}/q_1^{n_1} \oplus \cdots \oplus \mathbb{Z}/q_l^{n_l}$. Using the previous lemma, we can "kill" all elements of order products of q_i : we create the quotient of G by the normal subgroups generated by those elements. As G is finite, we can repeat this process a finitely many times to get a solvable group \tilde{G} such that $X\tau K(G, 1) \Leftrightarrow X\tau K(\tilde{G}, 1)$ for 2-dimensional compacta X .

We claim that \tilde{G} has trivial abelianization: in the exact sequence $1 \rightarrow [\tilde{G}, \tilde{G}] \rightarrow \tilde{G} \rightarrow \tilde{G}_{ab} \rightarrow 1$ let y be any generator of \tilde{G}_{ab} and consider its preimage y' . As $y^q = 1$ for some $q \in \{q^1, \dots, q^l\}$, y'^q is mapped to 1 in \tilde{G}_{ab} , and hence $y'^q = k$ for some element $k \in [\tilde{G}, \tilde{G}]$. Every element in $[\tilde{G}, \tilde{G}]$ has order relatively prime to the $q_i(s)$, so suppose k has order p , relatively prime to the $q_i(s)$. Thus we have $y'^{pq} = 1$ which implies $y'^p = 1$, which in turn implies $y^p = 1$. Thus $y = 1$, and our claim is proved.

Remark 4.7. The author thinks that the above theorem is true for any solvable group with all elements of finite order. There are a few details which are being worked on. This paper is a preliminary version before the "22nd annual workshop in geometric topology, Colorado College" and the Bedlewo conference.

5. EXTENSION PROPERTIES OF THE PONTRYAGIN DISK

We will use for the sequel a 2-dimensional compactum called the mod-2 Pontryagin disk $\mathbb{P}D^2$, which is a variant of a classical construction of L.S. Pontryagin [12].

The space $\mathbb{P}D^2$ is constructed as the inverse limit of an inverse sequence of spaces. As the first stage of the construction, we start with the usual 2-cell embedded in \mathbb{R}^3 and take some triangulation of it with mesh less than 1. To get the $k+1$ th stage from the k th stage we take each 2-simplex, remove its interior and glue a disk along the degree 2 map, then give a triangulation of mesh less than $1/(k+1)$ to the resulting space. For the bonding map from $k+1$ th stage to the k th stage, we send the interior 2-simplices of the attached disks to the barycenter of the removed 2-simplices.

The following theorem is very useful in constructing compacta with special extension properties. For explanation of terms and proof see [7]. Note that $\text{mesh}\{\lambda_i\} \rightarrow 0$ means that for every k , $\lim_{i \rightarrow \infty} \{\text{mesh}(q_k^{k+i}(\lambda_{k+i}))\} = 0$, which is not the same as $\lim_{i \rightarrow \infty} \text{mesh}\lambda_i = 0$.

Theorem 5.1. *Suppose that K is a countable CW complex and that X is a compactum such that $X = \lim_{\leftarrow} (X_i, f_i^{i+1})$, where (L_i, f_i^{i+1}) is a K -resolvable inverse system of compact polyhedra L_i with triangulations λ_i such that $\text{mesh}\{\lambda_i\} \rightarrow 0$. Then $X \tau K$.*

Theorem 5.2. *Let $\mathbb{P}D^2$ be the Pontryagin disk. Then $\mathbb{P}D^2 \tau K(\mathbb{Z}/2, 1)$.*

Proof: Let $A \subset \mathbb{P}D^2$ be a closed and consider the map $f : A \rightarrow K(\mathbb{Z}/2, 1)$. Represent $(\mathbb{P}D^2, A) = \lim_{\leftarrow} \{((\mathbb{P}D^2)_i, A_i), q_i^{i+1}\}$ in the usual way as the inverse limit of polyhedral pairs (all lying in I^∞). Like in the preceding theorem, find i_0 such that f can be extended over A_{i_0} to give a map $\overline{f}_{i_0} : A_{i_0} \rightarrow K(\mathbb{Z}/2, 1)$ with $\overline{f}_{i_0} \circ (q_{i_0}^\infty) \simeq f$. Extend \overline{f}_{i_0} over the 1-skeleton of $((\mathbb{P}D^2)_i)$ and get an induced extension upto homotopy $\overline{G} : A \cup (\mathbb{P}D^2)^{(1)} \rightarrow K(\mathbb{Z}/2, 1)$. Pick any 2-simplex $\sigma_j \in ((\mathbb{P}D^2)_{i_0})^{(1)} - (A_{i_0})^{(1)}$ and consider $\gamma_{\sigma_j} = g_{i_0}(\partial\sigma_j) \in \mathbb{Z}/2$. Consider $(q_{i_0}^\infty)^{-1}(\sigma_j) \subset \mathbb{P}D^2$ and note that we have an extension (of f) over $A \cup (q_{i_0}^\infty)^{-1}(\partial\sigma_j)$. We want to extend over $A \cup (q_{i_0}^\infty)^{-1}(\sigma_j)$: if γ_{σ_j} is trivial there is no obstruction. If not, find the smallest index $i_1 > i_0$ such that $(p_{i_0}^{i_1})^{-1}$ has an $(\mathbb{R}P)^2$ (needs explanation / rephrasing). Clearly the extension over $A \cup (q_{i_0}^\infty)^{-1}$ is possible. Do the same for all such 2-simplices σ_j .

6. INFINITE SOLVABLE GROUPS

Theorem 6.1. *Let $\mathbb{P}D^2$ be the Pontryagin disk and Γ be a torsion free group with abelianization being a non-trivial 2-group. Then $K(\Gamma_{ab}, 1)$ is an absolute extensor of $\mathbb{P}D^2$, but $K(\Gamma, 1)$ is not an absolute extensor of $\mathbb{P}D^2$.*

Proof: Let σ be any 2-simplex in the first stage of construction of $\mathbb{P}D^2$ and identity $\partial\sigma$ and its image under in X under $(f_0^\infty)^{-1}\partial\sigma$. Consider the map $f : \partial\sigma \rightarrow K(\Gamma, 1)$ that sends $\partial\sigma$ to the generator x along the identity map $S^1 \rightarrow S^1$.

We claim that this map does not extend: suppose there is an extension $g : \mathbb{P}D^2 \rightarrow K(\Gamma, 1)$. Then there is some integer i_0 such that there is an extension up to homotopy $g_{i_0} : \mathbb{P}D^2_{i_0} \rightarrow K(\Gamma, 1)$. In that case, x could be expressed as products of elements of Γ , each of order 2. But Γ is torsion-free, and we have a contradiction.

We give 2 examples of groups satisfying theorem 6.1, with Γ_{ab} being a 2-group.

Example 6.2. Consider the following group, which was first studied by Hirsch:

$$\Gamma_1 = \langle x, y, z \mid x^z = x^{-1}, y^z = y^{-1}, [x, y] = z^4 \rangle.$$

This example of Hirsch is a torsion free polycyclic group which is not poly infinite cyclic. The following lemma shows that G is not nilpotent.

Lemma 6.3 (Robinson, 5.2.20). *A finitely generated torsion-free nilpotent group has a central series with infinite cyclic factors.*

The author thanks Prof. Thomas Farrell for pointing out the next example.

Example 6.4. It is known [14]theorem 3.5.5 that there are just 6 affine diffeomorphism classes of compact connected orientable flat 3-dimensional Riemannian manifolds, represented by manifolds \mathbb{R}^3/G , where G is one of 6 given groups. One of them is our next example

$$\Gamma_2 = \langle \alpha, \beta, \gamma x, y, z \mid \gamma\beta\alpha = xy, \alpha^2 = x, \alpha y \alpha^{-1} = y^{-1}, \alpha z \alpha^{-1} = z^{-1}, \\ \beta x \beta^{-1} = x^{-1}, \beta^2 = y, \beta z \beta^{-1} = z^{-1}, \gamma x \gamma^{-1} = x^{-1}, \\ \gamma y \gamma^{-1} = y^{-1}, \gamma^2 = z \rangle.$$

It's abelianization is $\mathbb{Z}/4 \oplus \mathbb{Z}/4$. It is torsion free and solvable but not nilpotent.

Remark 6.5. Even solvable groups with some elements of infinite order may not be enough to serve as an example of groups in theorem 6.1.

As an example, consider D_∞ , which is defined to be the semi-direct product $\mathbb{Z} \rtimes_\theta \mathbb{Z}/2$, where $\theta : \mathbb{Z}/2 \rightarrow \text{Aut}(\mathbb{Z})$ is the "t⁻¹" map. A presentation of D_∞ is $\langle x, y \mid y^2 = 1, yxy^{-1} = x^{-1} \rangle$. Note that the relations imply $(yx)^2 = 1$,

so $D_\infty = \mathbb{Z}/2 * \mathbb{Z}/2$ and therefore, the abelianization is $\mathbb{Z}/2 \oplus \mathbb{Z}/2$. The commutator is $\langle x \rangle = \mathbb{Z}$.

As $D_\infty = \mathbb{Z}/2 * \mathbb{Z}/2$, $\mathbb{P}D^2\tau K(D_\infty, 1)$.

REFERENCES

- [1] M.Cenceli and A.N.Dranishnikov, *Extension of maps to nilpotent spaces*, Canadian Mathematical Bulletin, vol. 44(3), 2001 pp.266-269.
- [2] M.Cenceli and A.N.Dranishnikov, *Extension of maps to nilpotent spaces - II*, Topology Appl. 124(2002), no. 1, pp.77-83.
- [3] M.Cenceli and A.N.Dranishnikov, *Extension of maps to nilpotent spaces - III*, Topology Appl. ***.
- [4] A.Dold and R. Thom, *Quasifaserungen und unendliche symmetrische produkte*, Ann. of Math.(2), vol.67 (1958), no **, pp.239-281.
- [5] A.N.Dranishnikov, *Extension of mappings to CW complexes*, Math. USSR Sbornik, vol.74 (1993), no1, pp.47-56.
- [6] A.N.Dranishnikov, *Cohomological Dimension Theory of Compact Metric Spaces*, preprint (1999).
- [7] A.N.Dranishnikov and Dusan Repovs, *Cohomological dimension with respect to perfect groups*, Topology Appl. 74(1996), pp.123-140.
- [8] J. Dydak, *Cohomological dimension and metrizability spaces*, Trans. of the Amer. Math.Soc.,337(1993),219-234.
- [9] J. Dydak, *Cohomological dimension theory*, Handbook of Geometric Topology, Elsevier Science B.V., 2002.
- [10] , *Homotopy Theory and Duality*, Gordon and Breach, New York, 1965.
- [11] A. Lundell and S. Weingram, *The topology of CW complexes*, Van Nostrand Reinhold Company, 1969.
- [12] L.S. Pontrjagin, *Sur Une hypothese fondamentale de la dimension*, C.R. Acad. Sci. 190(1930), 1105-1107.
- [13] D.J.S. Robinson, *Introduction to the theory of groups*, GTM, Springer-Verlag
- [14] J.A. Wolf, *Spaces of Constant Curvature*, McGraw-Hill Book Co.,1967.

UNIVERSITY OF TENNESSEE, KNOXVILLE, TN 37996, USA
E-mail address: ajmitra@math.utk.edu