

**ON CONTINUOUS BUT DIFFERENTIABLE NOWHERE  
SOLUTIONS IN THE MOMENT PROBLEM**

MIGUEL TIERZ

Since the times of Chebyshev and, especially, after Stieltjes seminal memoir [9], the moment problem has always been a subject of remarkable interest. Its notorious influence and appearance in many areas is well-known [1, 7, 8].

Let us briefly present the fundamentals of the moment problem. Let  $I \subseteq \mathbb{R}$  be an interval. For a positive measure  $\mu$  on  $I$ , the  $n$ th moment is defined as  $\int_I x^n d\mu(x)$ , provided the integral exists. Consider now the set  $\{\gamma_n\}_{n=0}^\infty$ , an infinite but countable sequence of real numbers. Then, the moment problem on  $I$  consists in the solution of the following questions:

0.0.1. *Does there exist a positive measure on  $I$  with moments  $\{\gamma_n\}_{n=0}^\infty$  ?*

If so:

0.0.2. *Is this positive measure uniquely determined by the moments  $\{\gamma_n\}_{n=0}^\infty$  ?*

One may also want to consider a third question, in case the previous one is answered in the negative:

0.0.3. *How can all the positive measures on  $I$ , with moments  $\{\gamma_n\}_{n=0}^\infty$ , be described ?*

This last question is the main focus here. The simplest way to show the non-uniqueness feature is to recall Stieltjes famous computation<sup>1</sup> [9]:

$$(0.1) \quad \int_0^\infty x^n e^{-k^2 \log^2 x} \sin\left(2\pi \frac{\log x}{\log q}\right) dx = 0, \quad n \in \mathbb{Z},$$

where  $q = e^{-1/2k^2}$ . Thus, independently of  $\lambda$ , we have:

$$(0.2) \quad \begin{aligned} & \frac{1}{\sqrt{\pi}} \int_0^\infty x^n e^{-k^2 \log^2 x} \left(1 + \lambda \sin\left(2\pi \frac{\log x}{\log q}\right)\right) dx \\ &= \frac{1}{\sqrt{\pi}} \int_0^\infty x^n e^{-k^2 \log^2 x} dx = q^{(n+1)^2/2}. \end{aligned}$$

Our main point is almost trivial but apparently absent in the literature: instead of (0.1), one can more generally write:

$$(0.3) \quad \int_0^\infty x^n e^{-k^2 \log^2 x} \sin\left(2\pi j \frac{\log x}{\log q}\right) dx = 0, \quad n, j \in \mathbb{Z},$$

which means that one can construct the Fourier series:

$$(0.4) \quad f(x) = \frac{1}{\sqrt{\pi}} e^{-k^2 \log^2 x} \left(1 + \lambda \sum_{n=1}^m a_n \sin\left(2\pi b_n \frac{\log x}{\log q}\right)\right),$$

<sup>1</sup>Stieltjes considered  $k = 1$ .

and the integer moments keep their (log-normal) value (0.2). Now, it is manifest that a suitable choice of  $a_n$  and  $b_n$  and the case  $m \rightarrow \infty$  lead to functions that are continuous but differentiable nowhere. The sum-integral exchange in the moment computation of (0.4) (term by term integration) is, by virtue of Lebesgue dominated convergence theorem, not a problem in the  $m \rightarrow \infty$  case, as soon as the series is convergent.

One can get a more compact and elegant description by employing a more general characterization of the non-uniqueness, first found by Chihara in the seventies [3] and more recently exploited in [4]. It says that the moments remain unchanged if  $f(x) = \frac{1}{\sqrt{\pi}} k e^{-k^2 \log^2 x}$  is modified in the following way:

$$(0.5) \quad f(x) \rightarrow f(x)g(x), \text{ with } g(x) = g(qx).$$

This  $q$ -periodic function  $g(x)$  is well known in the  $q$ -calculus literature and is the most general solution of  $D_g g(x) = 0$  where  $D_g g(x) = \frac{g(x) - g(qx)}{(q-1)x}$  denotes the usual  $q$ -derivative [5]. Interestingly enough, this  $q$ -periodic function has been studied in the context of fractal geometry [5]. It turns out that nowhere differentiable functions are perfectly well-behaved under the  $q$ -derivative so, while in  $q$ -calculus a  $q$ -periodic function essentially plays the role of a constant, it may easily be differentiable nowhere under the ordinary derivative.

The property (0.5) was found employing the functional equation satisfied by the log-normal function,  $f(xq) = \sqrt{q} x f(x)$ . This functional equation actually corresponds to the  $q$ -Pearson equation associated to the Stieltjes-Wigert orthogonal polynomials. It can be easily shown that the same argument in [4] can be identically applied to any orthogonal polynomials system that is an indeterminate moment problem (all within the  $q$ -deformed Askey scheme [6]). Thus, the property (0.5), also holds for all of them and hence, the existence of differentiable nowhere solutions appears in the same way. A detailed account of this and other related results will appear elsewhere [10].

**Remark 0.1.** *To conclude, we would like to make a remark of historical nature. In addition to Stieltjes seminal memoir, another relevant mathematical document is the correspondence between Hermite and Stieltjes [2]. There are two well-known letters that are still often quoted.*

*One is a letter from Hermite to Stieltjes (number 351, dated 22 October 1892) which shows the admiration and respect for the results of Stieltjes. This was a reply to previous correspondence from Stieltjes, where he described his discoveries to him. Among these results, the one we have discussed here. In his own words:*

*"L'existence de ces fonctions  $\varphi(u)$  qui, sans être nulles, son telles que*

$$\int_0^\infty u^k \varphi(u) du = 0 \quad (k = 0, 1, 2, 3, \dots)$$

*me paraît très remarquable."*

*The second one, dated 20 May (1893), is even more well-known, and is the one where Hermite emphatically expresses his dislike for the then recently introduced continuous but differentiable nowhere functions.*

Therefore, it seems interesting that the same results that impressed Hermite, actually contained, as we have shown, mathematics that he despised so much. Namely, continuous but differentiable nowhere functions.

## REFERENCES

- [1] N.I. Akhiezer, *The Classical Moment Problem*, Hafner Publishing Co., New York, 1965;
- [2] B. Baillaud and H. Bourget, *Correspondance d'Hermite et Stieltjes I, II*, Gauthiers-Villars, Paris 1905.
- [3] T.S. Chihara, On generalized Stieltjes-Wigert and related orthogonal polynomials, *J. Comp. Appl. Math.* 5(4) (1979), 291-297
- [4] J.S. Christiansen, The moment problem associated with the Stieltjes-Wigert polynomials, *J. Math. Anal. Appl.* **277** (2003), 218-245
- [5] A. Erzan and J.-P. Eckmann, q-analysis of fractal sets, *Phys. Rev. Lett.* 78 (1997); 3245-3248
- [6] R. Koekoek and R. F. Swarttouw : *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue*. Delft University of Technology, Faculty of Information Technology and Systems, Department of Technical Mathematics and Informatics, Report no. 98-17, 1998. Available on-line at: <http://aw.twi.tudelft.nl/~koekoek/askey.html>
- [7] J.A. Shohat and J.D. Tamarkin, *The problem of moments*, Amer. Math. Soc. Surveys, No. 1 (1943).
- [8] B. Simon, The Classical Moment Problem as a Self-Adjoint Finite Difference Operator, *Adv. Math.* 137 (1998) 82-203.
- [9] T. Stieltjes, Recherches sur les fractions continues, *Anns, Sci. Fac. Univ. Toulouse* (1894-1895), 8, J1-J122; 9, A5-A47. English version in T.J. Stieltjes , *Collected Papers*, G. van Dijk (ed.), Vol II, Springer, Berlin 1993.
- [10] M. Tierz, In preparation

INSTITUT D'ESTUDIS ESPACIALS DE CATALUNYA (IEEC/CSIC). CAMPUS UAB, FAC CIENCIES,  
TORRE C5-PARELL-2A PLANTA, E-08193 BELLATERRA (BARCELONA) SPAIN  
*E-mail address:* tierz@ieec.fcr.es