Stein's Method and Random Character Ratios Running head: Stein's Method and Random Character Ratios

Version of 8/13/05

By Jason Fulman

University of Pittsburgh

Department of Mathematics, 301 Thackeray Hall, Pittsburgh PA 15260

email: fulman@math.pitt.edu

Abstract: Stein's method is used to prove limit theorems for random character ratios. Tools are developed for four types of structures: finite groups, Gelfand pairs, twisted Gelfand pairs, and association schemes. As one example an error term is obtained for a central limit theorem of Kerov on the spectrum of the Cayley graph of the symmetric group generated by i-cycles, or equivalently for the character ratio of a Plancherel distributed representation on an i-cycle. Other main examples include an error term for a central limit theorem of Ivanov on character ratios of random projective representations of the symmetric group, and a new central limit theorem for the spectrum of certain graphs whose vertices are the set of perfect matchings on 2n symbols. The error terms in the resulting limit theorems are typically $O(n^{-1/4})$ or better. The results are obtained with remarkably little information: a character formula for a single representation close to the trivial representation and estimates on two step transition probabilities of a random walk. Although the limit theorems stated in this paper are all for the case of normal approximation, many of the tools developed are quite general. Indeed, both the construction of an exchangeable pair used for Stein's method and lemmas computing certain moments are useful for arbitrary distributional approximation.

2000 Mathematics Subject Classification: Primary 05E10, Secondary 60C05.

Key words and phrases: Stein's method, normal approximation, Gelfand pair, character ratio, symmetric group, Plancherel measure, association scheme.

1. INTRODUCTION

Given a fixed $m \times m$ matrix, it is natural to study the distribution of its eigenvalues, where each eigenvalue is chosen with probability $\frac{1}{m}$. As a motivating example around which discussion can be centered, consider the $n! \times n!$ transition matrix for random walk on the symmetric group S_n , where the generating set consists of all i -cycles. Diaconis and Shahshahani [\[DSh\]](#page-50-0) proved that the eigenvalues of this matrix are the numbers $\frac{\chi^{\lambda}_{(i,1^{n-i})}}{dim(\lambda)}$ $\frac{d^{(i,1^{n-i})}}{dim(\lambda)}$ occurring with multiplicity $dim(\lambda)^2$. Here λ parameterizes an irreducible representation of the symmetric group, $\chi^{\lambda}_{(i,1^{n-i})}$ is the corresponding character value on *i*-cycles, and $dim(\lambda)$ is the dimension of the irreducible representation.

Since $\sum_{|\lambda|=n} dim(\lambda)^2 = n!$, the eigenvalue $\frac{\lambda^{\lambda}_{(i,1^{n-i})}}{dim(\lambda)}$ $\frac{\partial f(i,1^{n-1})}{\partial \lim(\lambda)}$ is chosen with probability $dim(\lambda)^2$ $\frac{n(\lambda)}{n!}$.

The probability measure on irreducible representations of S_n which picks the representation corresponding to λ with probability $\frac{dim(\lambda)^2}{n!}$ is known as the Plancherel measure of the symmetric group. Kerov [\[K1\]](#page-51-0) proved that if $i \geq 2$ is fixed, and λ is random from the Plancherel measure of the symmetric group, then the random variable $\sqrt{\binom{n}{i}(i-1)!}\chi^{\lambda}_{(i,1^{n-i})}$ $\frac{\partial^{(i)}(i,1^{n-i})}{\partial (in(\lambda))}$ is asymptotically normal as $n \to \infty$. Kerov's proof used the method of moments and difficult combinatorics; a beautiful exposition of his work is the paper [\[IO\]](#page-50-1). Hora [\[Ho1\]](#page-50-2) gave another proof of Kerov's result, also using the method of moments, but with somewhat simpler combinatorics based on the fact that the kth moment of the character ratio $\frac{\chi^{\lambda}_{(i,1^{n-i})}}{dim(\lambda)}$ $\frac{\partial^{(i,1^{n-1})}}{\partial^2 m(\lambda)}$ is equal to the probability that the walk generated by the set of i-cycles is at the identity after k steps. In very recent work, Sniady [\[Sn1\]](#page-51-1), [\[Sn2\]](#page-51-2) uses the genus expansion of random matrix theory to give another method of moments proof of Kerov's result.

A more probabilistic approach to Kerov's result for the case $i = 2$ was given in [\[F1\]](#page-50-3), where Stein's method was used to obtain the first error term in Kerov's central limit theorem; an error term of $O(n^{-1/4})$ was proved and an error term of $O(n^{-1/2})$ was conjectured. This error term was later improved to $O(n^{-s})$ for any $s < 1/2$ using martingale theory [\[F4\]](#page-50-4). More recently, a proof of the $O(n^{-1/2})$ conjecture appears in [\[ShSu\]](#page-51-3), using a new refinement of Stein's method. For other proofs of the $O(n^{-1/2})$ bound, see [\[CF\]](#page-50-5) for an argument with a different refinement of Stein's method, and [\[F3\]](#page-50-6) for an argument using Bolthausen's variation of Stein's method. All of these results, it should be emphasized, were only for the case $i = 2$. However even in the simple setting of $i = 2$, random character ratios arise in work on the moduli space of curves [\[EO\]](#page-50-7).

Before proceeding further, it should be mentioned that familiarity with Stein's method is not necessary to read this paper. Indeed, Section [2](#page-4-0) gives a very brief introduction to normal approximation by Stein's method. It presents the bare minimum needed to understand this paper, but gives a few pointers to the literature for further reading.

Section [3](#page-5-0) of this paper generalizes the set-up of Kerov's central limit theorem to the case when G is a finite group and the generating set consists of a single conjugacy class C. Quite general results are proved, elucidating our early work on this problem [\[F1\]](#page-50-3) which required different information than the current treatment. As a new application, it is shown that for any fixed $i \geq 2$, one obtains an error term $O(n^{-1/4})$ in Kerov's central limit theorem. The approach presented here is appealing because it uses only the most elementary ingredients, namely a well known character formula for the irreducible representation parameterized by $\lambda = (n-1,1)$ on all elements of S_n , and estimates on the two step transition probabilities of the random walk on S_n generated by C.

Section [4](#page-14-0) proves theorems in the setting of Gelfand pairs. If G is a group and K is a subgroup such that the induced representation 1_K^G is multiplicity free, the pair (G, K) is called a Gelfand pair. Stein's method is used to prove a central limit theorem for random spherical functions of the pair (G, K) . As an application, a new central limit theorem with error term $O(n^{-1/4})$ is obtained for the spectrum of certain random walks on the set of perfect matchings of $2n$ symbols. Equivalently, a central limit theorem is obtained for certain statistics under the $Jack₂$ measure on partitions, which is an interesting object [\[O2\]](#page-51-4). This application is an exact analog of Kerov's central limit theorem, and complements results in the paper [\[F2\]](#page-50-8), which obtained an analog of Kerov's central limit only for $i = 2$, but for the entire Jack_{α} family of measures on partitions, where there is not a known corresponding Gelfand pair. As in the group case, only the simplest ingredients are needed for the proof: a formula for a single spherical function (chosen to be as close to the trivial spherical function as possible) and estimates on the two step transition probabilities of a random walk.

Section [5](#page-26-0) focuses on twisted Gelfand pairs. This terminology, introduced in [\[Stm\]](#page-51-5), refers to a triple (G, K, ϕ) , where ϕ is a linear character of K such that $Ind_K^G(\phi)$ is multiplicity free. An error term is obtained for a central limit theorem of Ivanov [\[I\]](#page-50-9) on character ratios of random projective representations of the symmetric group. There is a close parallel to earlier sections, but there are major differences in argumentation. One striking example of this is that if one attempts a straightforward adaptation of the Markov chain used in the construction of an exchangeable pair in Section [4,](#page-14-0) the resulting "Markov chain" can have negative transition probabilities. However an interesting combinatorial argument shows that only the holding "probabilities" can be negative. Since Stein's method works by analyzing how a statistic changes under small perturbations, holding probabilities are not so important, and by rescaling one obtains a genuine Markov chain and so a legitimate construction of an exchangeable pair.

Section [6](#page-41-0) develops limit theorems for the spectrum of an adjacency matrix of an association scheme. The arguments are analogous to those in previous sections. This is not surprising since experts in the area will realize that Gelfand pairs and association schemes are both generalizations of the finite group case. However the perspective and examples are quite different, and it would confuse rather than clarify matters to present Kerov's central limit theorem as a result about Gelfand pairs or association schemes. As an example, the Hamming association scheme is treated. One obtains a central limit theorem for the spectrum of the Hamming graph, or equivalently for values of q-Krawtchouk polynomials.

Having outlined the contents of this article, it is useful to give four further reasons why we believe the results to be interesting. First, as mentioned in the abstract, many of the results here are applicable for distributional approximations other than normal approximation. Both the construction of an exchangeable pair and the computation of certain moments will be quite useful once the corresponding versions of Stein's method are developed. For an illustration of this point in the context of the spectrum of generalizations of the Bernoulli-Laplace diffusion model, see [\[CF\]](#page-50-5). Second, the problem of studying the spectrum of random walks on G/K where (G, K) is a Gelfand pair, is of ongoing interest, even in the case when G and K are finite groups. In particular if G, K are finite classical groups, this leads to interesting questions in number theory [\[Te\]](#page-51-6), and we are working on using the exchangeable pair constructed in this paper to obtain information. Third, Plancherel measure (which arises in the group case), Jack measure (which arises in the Gelfand pair case), and shifted Plancherel measure (which arise in the twisted Gelfand pair case), are all objects of interest to researchers in random matrix theory [\[AlD\]](#page-49-0), [\[BO1\]](#page-50-10), [\[BOO\]](#page-50-11), [\[De\]](#page-50-12), [\[J\]](#page-50-13), [\[Mat\]](#page-51-7), [\[O1\]](#page-51-8), [\[O2\]](#page-51-4), [\[TW\]](#page-51-9). As is evident from these papers, there are many interesting statistics under these measures, and the method of constructing exchangeable pairs in this paper should be useful for studying them by Stein's method. Fourth, the examples in this paper will be a useful testing grounds for results on Stein's method. For example the refinement of Stein's method in [\[ShSu\]](#page-51-3) arose from trying to obtain an $O(n^{-1/2})$ error term for the $i = 2$ case of Kerov's central limit theorem.

To close the introduction, some deficiencies with this work should be noted. First, the method of this paper does not seem to be applicable for studying the spectrum of arbitrary reversible Markov chains. In all of the examples presented here, there is an abundance of symmetry, and the list of eigenvalues of the matrix of interest is itself an eigenvector for another natural operator. Thus the method of moments, which typically doesn't give error terms, may be more versatile if one's only purpose is to obtain a limit theorem. The examples in the papers [\[Bi\]](#page-49-1), [\[Ho2\]](#page-50-14), [\[HHO\]](#page-50-15) (which are promising candidates for Stein's method as well) illustrate the use of the method of moments to study the spectrum of random walks. Second, it is not clear that the results in this paper are useful for bounding the convergence rates of the Markov chains whose spectrum is studied. Such bounds, reviewed in Chapter 3 of Diaconis [\[Di\]](#page-50-16), usually require information about the edge of the spectrum rather than the bulk of the spectrum. Third, the error terms, while good, are not perfect. As is well known to workers in normal approximation, the transition from an $O(n^{-1/4})$ bound to an $O(n^{-1/2})$ bound is often quite difficult. We conjecture that $O(n^{-1/2})$ bounds hold for Theorems [3.15,](#page-14-1) [4.20,](#page-26-1) and [5.2.](#page-27-0) To prove this will almost certainly require much finer combinatorial information, along the lines of that developed in [\[IO\]](#page-50-1) (which in turn relies on [\[IK\]](#page-50-17)), and even this is only for the special case of Plancherel measure of the symmetric group. The point of the current paper is to obtain good bounds with minimal information and effort.

2. Stein's Method for Normal Approximation

In this section we briefly review Stein's method for normal approximation, using the method of exchangeable pairs [\[Stn1\]](#page-51-10). One can also use couplings to prove normal approximations by Stein's method (see [\[Re\]](#page-51-11) for a survey), but the exchangeable pairs approach is effective for our purposes. For a survey discussing both exchangeable pairs and couplings, the paper [\[RiRo2\]](#page-51-12) can be consulted.

Two random variables W, W' on a state space X are called exchangeable if for all $w_1, w_2, \mathbb{P}(W = w_1, W' = w_2)$ is equal to $\mathbb{P}(W = w_2, W' = w_1)$. As is typical in probability theory, let $\mathbb{E}(A|B)$ denote the expected value of A given B . The following result of Stein uses an exchangeable pair (W, W') to prove a central limit theorem for W.

Theorem 2.1. ([\[Stn1\]](#page-51-10)) Let (W, W') be an exchangeable pair of real random variables such that $\mathbb{E}(W'|W) = (1 - a)W$ with $0 < a < 1$. Then for all real x_0 ,

$$
\mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx
$$

$$
\le \frac{\sqrt{Var(\mathbb{E}[(W'-W)^2|W])}}{a} + (2\pi)^{-\frac{1}{4}} \sqrt{\frac{1}{a} \mathbb{E}[W'-W]^3}.
$$

In order to apply Theorem [2.1](#page-4-1) to study a statistic W , one needs an exchangeable pair (W, W') . The usual way of doing this is to use Markov chain theory. A Markov chain K (with chance of going from x to y denoted by $K(x, y)$ on a finite set X is called reversible with respect to a probability distribution π if $\pi(x)K(x, y) = \pi(y)K(y, x)$ for all x, y . This condition implies that π is a stationary distribution for K. It is straightforward to check that if K is reversible with respect to π , then one obtains an exchangeable pair (W, W') as follows: choose $x \in X$ from π , then obtain x' by taking one step from x according to K, and set $(W, W') = (W(x), W(x'))$.

To apply Theorem [2.1,](#page-4-1) one needs the exchangeable pair arising from K to satisfy $\mathbb{E}(W'|W) = (1-a)W$, and also one must obtain useful bounds on $Var(\mathbb{E}[(W'-W)^2|W])$ and $\mathbb{E}|W-W'|^3$. A main contribution of this paper is to provide a number of algebraically natural examples where this can be done. As the reader will see, the construction of K in our examples can be subtle.

A drawback with Theorem [2.1](#page-4-1) is that in many problems of interest, it gives a convergence rate of $O(n^{-1/4})$ rather than $O(n^{-1/2})$. When $|W'-W|$ is bounded, the following variation often gives the correct rate. A similar result is in [\[RiRo1\]](#page-51-13).

Theorem 2.2. ($[\text{ShSu}]$) Let (W, W') be an exchangeable pair of real random variables such that $\mathbb{E}(W'|W) = (1 - a)W$ with $0 < a < 1$. Suppose that $|W' - W| \leq A$ for some constant A. Then for all real x_0 ,

$$
\left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right|
$$

$$
\le \frac{\sqrt{Var(\mathbb{E}[(W'-W)^2|W])}}{a} + .41 \frac{A^3}{a} + 1.5A.
$$

In many of our examples, $|W' - W|$ is unbounded. Thus as mentioned in the introduction, significantly more work may be needed to go beyond the $O(n^{-1/4})$ rate.

The following general lemmas are helpful in working with the error bounds in either Theorem [2.1](#page-4-1) or Theorem [2.2.](#page-4-2)

Lemma 2.3. Let (W, W') be an exchangeable pair of random variables such that $\mathbb{E}(W'|W) = (1 - a)W$ and $\mathbb{E}(W^2) = 1$. Then $\mathbb{E}(W' - W)^2 = 2a$.

Proof. Since W and W' have the same distribution,

$$
\mathbb{E}(W'-W)^2 = \mathbb{E}(\mathbb{E}(W'-W)^2|W)
$$

= $\mathbb{E}((W')^2) + \mathbb{E}(W^2) - 2\mathbb{E}(W\mathbb{E}(W'|W))$
= $2\mathbb{E}(W^2) - 2\mathbb{E}(W\mathbb{E}(W'|W))$
= $2\mathbb{E}(W^2) - 2(1-a)\mathbb{E}(W^2)$
= $2a$.

 \Box

Lemma [2.4](#page-5-1) is a well known inequality (already used in the monograph [\[Stn1\]](#page-51-10)) and useful because often the right hand side is easier to compute or bound than the left hand side. To make this paper as self-contained as possible, we include a proof. Here x is an element of the state space X .

Lemma 2.4.

$$
Var(\mathbb{E}[(W'-W)^2|W]) \leq Var(\mathbb{E}[(W'-W)^2|x]).
$$

Proof. Jensen's inequality states that if g is a convex function, and Z a random variable, then $g(\mathbb{E}(Z)) \leq \mathbb{E}(g(Z))$. There is also a conditional version of Jensen's inequality (Section 4.1 of $[Du]$) which states that for any σ subalgebra F of the σ -algebra of all subsets of X,

$$
\mathbb{E}(g(\mathbb{E}(Z|F))) \leq \mathbb{E}(g(Z)).
$$

The lemma follows by setting $g(t) = t^2$, $Z = \mathbb{E}((W'-W)^2|x)$, and letting F be the σ -algebra generated by the level sets of W.

3. Finite groups

This section uses Stein's method to study the spectrum of random walk on a finite group G , where the generating set is a conjugacy class C which satisfies $C = C^{-1}$. As will be explained in Subsection [3.1,](#page-6-0) by [\[DSh\]](#page-50-0) this is

equivalent to studying the distribution of the character ratio $\frac{\chi^{\lambda}(C)}{dim(\lambda)}$ where λ is chosen from the Plancherel measure of the group G.

The organization of this section is as follows. Subsection [3.1](#page-6-0) recalls the necessary background from representation theory. Subsection [3.2](#page-8-0) then defines a Markov chain on the set of irreducible representations of G , and uses it to construct an exchangeable pair. It then shows that the moment computations needed for Stein's method can be carried out provided that one knows the character values of a single nontrivial irreducible representation of G (which to be useful should be close to the trivial representation) and that one can estimate the two-step transition probabilities of the random walk on G generated by C . This leads to a general central limit theorem. Subsection [3.3](#page-13-0) applies the theory to the symmetric group S_n with C the conjugacy class of i -cycles, where i is fixed and n is large.

3.1. Background from representation theory. We recall facts from the representation theory of finite groups, referring the reader to [\[Sa\]](#page-51-14) or [\[Se\]](#page-51-15) for more details. In what follows, χ denotes a character of the finite group G , $dim(\chi)$ is the dimension of the corresponding representation, and $Irr(G)$ is the set of all irreducible characters of G. Also \overline{z} denotes complex conjugate of a number z.

Lemma 3.1. Let χ be an irreducible representation of G. Then $\chi(C^{-1}) =$ $\overline{\chi(C)}$. Thus if $C = C^{-1}$, then $\chi(C)$ is real.

Next we recall the orthogonality relations for irreducible characters of G.

Lemma 3.2. Let ν and χ be irreducible characters of a finite group G. Then

$$
\frac{1}{|G|} \sum_{g \in G} \nu(g) \overline{\chi(g)} = \delta_{\nu,\chi}.
$$

Lemma 3.3. Let C be the conjugacy class of G containing the element g . Then for $g \in G$,

$$
\sum_{\chi \in Irr(G)} \chi(g)\overline{\chi(h)}
$$

is equal to $\frac{|G|}{|C|}$ if h, g are conjugate and is 0 otherwise.

Lemma [3.4](#page-6-1) while known, is perhaps not well known, and since analogous results will be needed in later sections, a proof along the lines of one in [\[HSS\]](#page-50-19) is included.

Lemma 3.4. Let G be a finite group with conjugacy classes C_1, \dots, C_t . Let C_k be the conjugacy class of an element $w \in G$. Then the number of $m\text{-tuples } (g_1, \dots, g_m) \in G^m \text{ such that } g_j \in C_{i_j} \text{ and } g_1 \dots g_m = w \text{ is }$

$$
\prod_{j=1}^m |C_{i_j}| \sum_{\chi \in Irr(G)} \frac{dim(\chi)^2}{|G|} \frac{\chi(C_{i_1})}{dim(\chi)} \cdots \frac{\chi(C_{i_m})}{dim(\chi)} \frac{\overline{\chi(C_k)}}{dim(\chi)}.
$$

Proof. Identify each class C_i with its corresponding class sum in the complex group algebra $\mathbb{C}G$. If $\chi^{(1)}, \cdots, \chi^{(t)}$ are the irreducible complex characters of G , then the elements

$$
E_s = \frac{dim(\chi^{(s)})}{|G|} \sum_{j=1}^t \overline{\chi_j}^{(s)} C_j \quad (1 \le s \le t)
$$

are a complete set of orthogonal idempotents for the center of $\mathbb{C}G$, where $\chi^{(s)}_i$ $j^{(s)}_j$ denotes the value of $\chi^{(s)}$ at any $g \in C_j$. Lemma [3.3](#page-6-2) implies that

$$
C_j = |C_j| \sum_{s=1}^{t} \frac{\chi_j^{(s)}}{\dim(\chi^{(s)})} E_s.
$$

Since the E_j 's are orthogonal idempotents (i.e. $E_rE_s = \delta_{r,s}E_r$), it follows that

$$
C_{i_1} C_{i_2} \cdots C_{i_m} = |C_{i_1}| \cdots |C_{i_m}| \sum_{s=1}^t \frac{\chi_{i_1}^{(s)} \cdots \chi_{i_m}^{(s)}}{\dim(\chi^{(s)})^m} E_s
$$

$$
= \frac{|C_{i_1}| \cdots |C_{i_m}|}{|G|} \sum_{k=1}^t C_k \sum_{s=1}^t \frac{\chi_{i_1}^{(s)} \cdots \chi_{i_m}^{(s)} \overline{\chi_k}^{(s)}}{\dim(\chi^{s})^{m-1}},
$$

as desired. \Box

As a corollary one obtains the following result.

Corollary 3.5. ([\[DSh\]](#page-50-0)) Suppose that C is a conjugacy class satisfying $C =$ C^{-1} . Then the eigenvalues of the random walk on G with generating set C are indexed by $\chi \in Irr(G)$ and are the numbers $\frac{\chi(C)}{dim(\chi)}$, occurring with multiplicity $dim(\chi)^2$.

Proof. If M is the $|G| \times |G|$ transition matrix for the random walk, the chance of being at the identity after k steps is the trace of M^k divided by |G|. Thus Lemma [3.4](#page-6-1) implies that for all $k \geq 0$, the trace of M^k is equal to

$$
\sum_{\chi \in Irr(G)} dim(\chi)^2 \left(\frac{\chi(C)}{dim(\chi)}\right)^k.
$$

Since this holds for all $k \geq 0$, the result follows.

As mentioned in the introduction, the Plancherel measure of G is the probability measure on $Irr(G)$ which chooses each χ with probability $\frac{dim(\chi)^2}{|G|}$ $\frac{n(\chi)^{-}}{|G|}$. So Corollary [3.5](#page-7-0) says that the eigenvalues of the random walk on G generated by C are the "character ratios" $\frac{\chi(C)}{dim(\chi)}$ occurring with multiplicity proportional to the Plancherel probability of χ .

$$
\overline{}
$$

3.2. Central limit theorems for character ratios. The goal of this subsection is to prove a central limit theorem for the random variable W defined by $W(\lambda) = \frac{|C|^{1/2} \chi^{\lambda}(C)}{dim(\lambda)}$ $\frac{d^{1/2}\chi^{\Lambda}(C)}{dim(\lambda)}$, where C is a fixed conjugacy class such that $C = C^{-1}$ and λ is random from the Plancherel measure of $G.$

For this purpose, it is useful to construct a Markov chain on the set of irreducible representations of G as follows. First, fix a non-trivial irreducible representation τ whose character is real valued. This gives a Markov chain L_{τ} by defining the probability of transitioning from λ to ρ as

$$
L_{\tau}(\rho,\lambda) := \frac{dim(\rho)}{dim(\lambda)dim(\tau)} \frac{1}{|G|} \sum_{g} \chi^{\lambda}(g) \chi^{\tau}(g) \overline{\chi^{\rho}(g)}.
$$

It is worth remarking that since Stein's method is in the spirit of Taylor approximation, τ should typically be chosen to be as close as possible to the trivial representation.

Lemma [3.6](#page-8-1) verifies that L_{τ} is a Markov chain which is reversible with respect to Plancherel measure.

Lemma 3.6. The transition probabilities of L_{τ} are real and non-negative and sum to 1. Moreover the chain L_{τ} is reversible with respect to the Plancherel measure of G.

Proof. The transition probabilities of L_{τ} are real and non-negative since

$$
\frac{1}{|G|} \sum_{g} \chi^{\lambda}(g) \chi^{\tau}(g) \overline{\chi^{\rho}(g)}
$$

is the multiplicity of ρ in the tensor product of λ and τ . Letting id denote the identity, it follows from Lemma [3.3](#page-6-2) that

$$
\sum_{\rho} L_{\tau}(\lambda, \rho) = \frac{1}{\dim(\lambda)\dim(\tau)|G|} \sum_{g} \chi^{\lambda}(g) \chi^{\tau}(g) \sum_{\rho} \chi^{\rho}(id) \overline{\chi^{\rho}(g)} = 1.
$$

For the reversibility assertion, the fact that χ^{τ} and the transition probabilities of L_{τ} are both real valued implies that

$$
\frac{\dim(\lambda)^2}{|G|}L_{\tau}(\lambda,\rho)=\frac{\dim(\rho)^2}{|G|}L_{\tau}(\rho,\lambda).
$$

An exchangeable pair (W, W') is now constructed from the chain L_{τ} in the standard way. First choose λ from the Plancherel measure of G, then choose ρ with probability $L_{\tau}(\lambda, \rho)$, and finally let $(W, W') = (W(\lambda), W(\rho))$. The remaining results in this subsection show that the exchangeable pair (W, W') has desirable properties.

Lemma 3.7. $\mathbb{E}(W'|W) = \left(\frac{\chi^\tau(C)}{dim(\tau)}\right)$ $dim(\tau)$ $\big)W.$ *Proof.* From the definition of W' ,

$$
\mathbb{E}(W'|\lambda) = |C|^{1/2} \sum_{\rho} \frac{dim(\rho)}{dim(\lambda)dim(\tau)} \frac{1}{|G|} \sum_{g} \chi^{\lambda}(g) \chi^{\tau}(g) \overline{\chi^{\rho}(g)} \frac{\chi^{\rho}(C)}{dim(\rho)}
$$

$$
= \frac{|C|^{1/2}}{|G|} \sum_{g} \frac{\chi^{\lambda}(g)}{dim(\lambda)} \frac{\chi^{\tau}(g)}{dim(\tau)} \sum_{\rho} \chi^{\rho}(C) \overline{\chi^{\rho}(g)}
$$

$$
= \left(\frac{\chi^{\tau}(C)}{dim(\tau)}\right) W(\lambda),
$$

where the last step is Lemma [3.3.](#page-6-2) The result follows since this depends on λ only through W.

Corollary [3.8](#page-9-0) is not needed in what follows, but is worth recording.

Corollary 3.8. The eigenvalues of L_{τ} are $\frac{\chi^{\tau}(C)}{dim(\tau)}$ $\frac{\chi^2(C)}{dim(\tau)}$ as C ranges over conjugacy classes of G. The functions $\psi_C(\lambda) = \frac{|C|^{1/2} \chi^{\lambda}(C)}{dim(\lambda)}$ $\frac{d^{1/2}X^{n}(C)}{dim(\lambda)}$ are a basis of eigenvectors of L_{τ} , orthonormal with respect to the inner product

$$
\langle f_1, f_2 \rangle = \sum_{\lambda} f_1(\lambda) \overline{f_2(\lambda)} \frac{dim(\lambda)^2}{|G|}.
$$

Proof. The proof of Lemma [3.7](#page-8-2) shows that ψ_C is an eigenvector of L_{τ} with eigenvalue $\frac{\chi^{\tau}(C)}{dim(\tau)}$ $\frac{\chi(C)}{dim(\tau)}$. The orthonormality assertion follows from Lemma [3.3,](#page-6-2) and the basis assertion follows since the number of conjugacy classes of G is equal to the number of irreducible representations of G .

Lemma 3.9.
$$
\mathbb{E}(W'-W)^2=2\left(1-\frac{\chi^{\tau}(C)}{dim(\tau)}\right).
$$

Proof. This is immediate from Lemma [2.3](#page-5-2) and [3.7.](#page-8-2)

$$
\Box
$$

For the remainder of this subsection, if K is a conjugacy class of $G, p_m(K)$ will denote the probability that the random walk generated by C is in K after m steps.

Lemma 3.10.

$$
\mathbb{E}((W')^{2}|\lambda) = \frac{|C|}{\dim(\lambda)\dim(\tau)} \sum_{K} p_{2}(K)\chi^{\lambda}(K)\chi^{\tau}(K)
$$

where the sum is over all conjugacy classes K of G.

Proof.

$$
\mathbb{E}((W')^{2}|\lambda)
$$
\n
$$
= \frac{1}{\dim(\lambda)\dim(\tau)|G|} \sum_{\rho} \dim(\rho) \sum_{g} \chi^{\lambda}(g) \chi^{\tau}(g) \overline{\chi^{\rho}(g)} \left(\frac{|C|^{1/2} \chi^{\rho}(C)}{\dim(\rho)}\right)^{2}
$$
\n
$$
= \frac{|C|}{\dim(\lambda)\dim(\tau)|G|} \sum_{g} \chi^{\lambda}(g) \chi^{\tau}(g) \sum_{\rho} \dim(\rho) \overline{\chi^{\rho}(g)} \left(\frac{\chi^{\rho}(C)}{\dim(\rho)}\right)^{2}.
$$

The result now follows from Lemma [3.4.](#page-6-1) \Box

Lemma [3.11](#page-10-0) writes $Var([\mathbb{E}(W'-W)^2|\lambda])$ as a sum of positive quantities. Lemma 3.11.

$$
Var([\mathbb{E}(W'-W)^2|\lambda]) = |C|^2 \sum_{K \neq id} \frac{p_2(K)^2}{|K|} \left(\frac{\chi^{\tau}(K)}{dim(\tau)} + 1 - \frac{2\chi^{\tau}(C)}{dim(\tau)}\right)^2,
$$

where K ranges over all non-identity conjugacy classes of G.

Proof. By Lemmas [3.10,](#page-9-1) [3.9,](#page-9-2) and [3.7,](#page-8-2)

$$
Var([\mathbb{E}(W'-W)^2|\lambda])
$$

= $\mathbb{E}(\mathbb{E}((W'-W)^2|\lambda)^2) - 4\left(1 - \frac{\chi^{\tau}(C)}{\dim(\tau)}\right)^2$
= $|C|^2 \mathbb{E}\left[\frac{\sum_K p_2(K)\chi^{\lambda}(K)\chi^{\tau}(K)}{\dim(\lambda)\dim(\tau)} + \left(1 - \frac{2\chi^{\tau}(C)}{\dim(\tau)}\right)\left(\frac{\chi^{\lambda}(C)}{\dim(\lambda)}\right)^2\right]^2$
- $4\left(1 - \frac{\chi^{\tau}(C)}{\dim(\tau)}\right)^2$
= $T_1 + T_2 + T_3 - 4\left(1 - \frac{\chi^{\tau}(C)}{\dim(\tau)}\right)^2$

where

$$
T_1 = \frac{|C|^2}{\dim(\tau)^2} \mathbb{E}\left(\sum_K \frac{p_2(K)\chi^{\lambda}(K)\chi^{\tau}(K)}{\dim(\lambda)}\right)^2
$$

$$
T_2 = \frac{2|C|^2}{\dim(\tau)} \left(1 - \frac{2\chi^{\tau}(C)}{\dim(\tau)}\right) \sum_{K} p_2(K)\chi^{\tau}(K) \mathbb{E}\left[\frac{\chi^{\lambda}(K)}{\dim(\lambda)} \left(\frac{\chi^{\lambda}(C)}{\dim(\lambda)}\right)^2\right].
$$

$$
T_3 = |C|^2 \left(1 - \frac{2\chi^{\tau}(C)}{\dim(\tau)}\right)^2 \mathbb{E}\left(\frac{\chi^{\lambda}(C)}{\dim(\lambda)}\right)^4.
$$

Fortunately, these terms can be simplified. By Lemma [3.10,](#page-9-1) the expression inside parentheses in T_1 is real. Since χ^{τ} is real valued, Lemma [3.3](#page-6-2) implies that

$$
T_1 = |C|^2 \sum_K \frac{p_2(K)^2}{|K|} \frac{\chi^{\tau}(K)^2}{\dim(\tau)^2}.
$$

Since $\chi^{\lambda}(C)$ is real, Lemma [3.4](#page-6-1) implies that

$$
T_2 = 2|C|^2 \left(1 - \frac{2\chi^{\tau}(C)}{\dim(\tau)}\right) \sum_{K} \frac{p_2(K)^2}{|K|} \frac{\chi^{\tau}(K)}{\dim(\tau)}
$$

and since $C = C^{-1}$,

$$
T_3 = |C|^2 \left(1 - \frac{2\chi^{\tau}(C)}{\dim(\tau)}\right)^2 p_4(id)
$$

= $|C|^2 \left(1 - \frac{2\chi^{\tau}(C)}{\dim(\tau)}\right)^2 \sum_K \frac{p_2(K)^2}{|K|}.$

Thus one can write

$$
T_1 + T_2 + T_3 = |C|^2 \sum_K \frac{p_2(K)^2}{|K|} \left(\frac{\chi^{\tau}(K)}{\dim(\tau)} + 1 - \frac{2\chi^{\tau}(C)}{\dim(\tau)} \right)^2
$$

result follows since $p_2(id) = \frac{1}{|K|}$.

and the result follows since $p_2(id) = \frac{1}{|K|}$

Lemma 3.12. Let k be a positive integer.

(1)
$$
\mathbb{E}(W'-W)^k = |C|^{k/2} \sum_{m=0}^k (-1)^{k-m} {k \choose m} \sum_K \frac{\chi^{\tau}(K)}{\dim(\tau)} \frac{p_m(K)p_{k-m}(K)}{|K|}.
$$

\n(2)
$$
\mathbb{E}(W'-W)^4 \text{ is equal to}
$$

\n
$$
|C|^2 \sum_K \left[8\left(1 - \frac{\chi^{\tau}(C)}{\dim(\tau)}\right) - 6\left(1 - \frac{\chi^{\tau}(K)}{\dim(\tau)}\right)\right] \frac{p_2(K)^2}{|K|}.
$$

Proof. For the first assertion, note that

$$
\mathbb{E}((W'-W)^k|\lambda) = \frac{|C|^{k/2}}{\dim(\lambda)\dim(\tau)} \sum_{\rho} \frac{\dim(\rho)}{|G|} \sum_g \chi^{\lambda}(g) \chi^{\tau}(g) \overline{\chi^{\rho}(g)}
$$

$$
\cdot \sum_{m=0}^k (-1)^{k-m} {k \choose m} \left(\frac{\chi^{\rho}(C)}{\dim(\rho)}\right)^m \left(\frac{\chi^{\lambda}(C)}{\dim(\lambda)}\right)^{k-m}
$$

$$
= \frac{|C|^{k/2}}{\dim(\lambda)\dim(\tau)} \sum_{m=0}^k (-1)^{k-m} {k \choose m} \left(\frac{\chi^{\lambda}(C)}{\dim(\lambda)}\right)^{k-m}
$$

$$
\cdot \sum_g \chi^{\tau}(g) \chi^{\lambda}(g) \sum_{\rho} \frac{\dim(\rho)}{|G|} \left(\frac{\chi^{\rho}(C)}{\dim(\rho)}\right)^m \overline{\chi^{\rho}(g)}
$$

$$
= \frac{|C|^{k/2}}{\dim(\lambda)\dim(\tau)} \sum_{m=0}^k (-1)^{k-m} {k \choose m} \left(\frac{\chi^{\lambda}(C)}{\dim(\lambda)}\right)^{k-m}
$$

$$
\cdot \sum_K \chi^{\tau}(K) \chi^{\lambda}(K) p_m(K),
$$

where the final equality is by Lemma [3.4.](#page-6-1) Thus $\mathbb{E}((W'-W)^k)$ is equal to

$$
\mathbb{E}(\mathbb{E}((W'-W)^k|\lambda))
$$
\n
$$
= \frac{|C|^{k/2}}{\dim(\tau)} \sum_{m=0}^k (-1)^{k-m} {k \choose m}
$$
\n
$$
\cdot \sum_{K} p_m(K) \chi^{\tau}(K) \sum_{\lambda} \frac{\dim(\lambda)^2}{|G|} \frac{\chi^{\lambda}(K)}{\dim(\lambda)} \left(\frac{\chi^{\lambda}(C)}{\dim(\lambda)}\right)^{k-m}.
$$

The first assertion now follows from Lemma [3.4](#page-6-1) and the fact that $\chi^{\lambda}(C)$ is real for all λ .

For the second assertion, note by the first assertion that

$$
\mathbb{E}(W'-W)^4 = |C|^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_K \frac{\chi^{\tau}(K)}{\dim(\tau)} \frac{p_m(K)p_{4-m}(K)}{|K|}.
$$

If τ is the trivial representation, then $W' = W$, which implies that

$$
0 = |C|^2 \sum_{m=0}^{4} (-1)^m {4 \choose m} \sum_{K} \frac{p_m(K)p_{4-m}(K)}{|K|}.
$$

Thus for general τ ,

$$
\mathbb{E}(W'-W)^4 = -|C|^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_K \left(1 - \frac{\chi^{\tau}(K)}{\dim(\tau)}\right) \frac{p_m(K)p_{4-m}(K)}{|K|}.
$$

Observe that the $m = 0.4$ terms in this sum vanish, since the only contribution could come from the identity, which contributes 0. The $m = 2$ term is

$$
-6|C|^2\sum_K\left(1-\frac{\chi^\tau(K)}{dim(\tau)}\right)\frac{p_2(K)^2}{|K|}.
$$

The $m = 1, 3$ terms are equal and together contribute

$$
8|C|^2 \sum_K \left(1 - \frac{\chi^{\tau}(K)}{\dim(\tau)}\right) \frac{p_1(K)p_3(K)}{|K|}
$$

= 8|C| $\left(1 - \frac{\chi^{\tau}(C)}{\dim(\tau)}\right) p_3(C)$
= 8|C|² $\left(1 - \frac{\chi^{\tau}(C)}{\dim(\tau)}\right) p_4(id)$
= 8|C|² $\left(1 - \frac{\chi^{\tau}(C)}{\dim(\tau)}\right) \sum_K \frac{p_2(K)^2}{|K|},$

where id is the identity and the last two equalities can be seen either directly or from Lemma [3.4.](#page-6-1) This completes the proof of the second assertion. \Box

Putting the pieces together, one obtains the following theorem.

Theorem 3.13. Let C be a conjugacy class of a finite group G such that $C =$ C^{-1} and fix a nontrivial irreducible representation τ of G whose character is real valued. Let λ be a random irreducible representation, chosen from the Plancherel measure of G. Let $W = \frac{|C|^{1/2} \chi^{\lambda}(C)}{dim(\lambda)}$ $\frac{d}{dim(\lambda)}$. Then for all real x_0 ,

$$
\begin{split} \left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \\ &\le \frac{|C|}{a} \sqrt{\sum_{K \ne id} \frac{p_2(K)^2}{|K|} \left(\frac{\chi^{\tau}(K)}{dim(\tau)} + 2a - 1\right)^2} \\ &+ \left[\frac{|C|^2}{\pi} \sum_{K} \left(8 - \frac{6}{a} \left(1 - \frac{\chi^{\tau}(K)}{dim(\tau)}\right)\right) \frac{p_2(K)^2}{|K|} \right]^{1/4}, \\ &= 1 - \frac{\chi^{\tau}(C)}{dim(\tau)}. \end{split}
$$

 $where a$ $dim(\tau)$

Proof. One applies Theorem [2.1](#page-4-1) to the exchangeable pair (W, W') of this subsection. By Lemmas [2.4](#page-5-1) and [3.11,](#page-10-0) the first term in Theorem [2.1](#page-4-1) gives the first term in the theorem. To upper bound the second term in Theorem [2.1,](#page-4-1) note by the Cauchy-Schwarz inequality that

$$
\mathbb{E}|W'-W|^3 \leq \sqrt{\mathbb{E}(W'-W)^2\mathbb{E}(W'-W)^4}.
$$

Now use Lemma [3.9](#page-9-2) and part 2 of Lemma [3.12.](#page-11-0)

3.3. Example: Cayley Graphs of the Symmetric Group. This subsection applies the theory of Subsection [3.2](#page-8-0) to the case where the group is S_n and C is the conjugacy class of *i*-cycles.

It is useful to recall some facts about the symmetric group. Since $K =$ K^{-1} for all conjugacy classes K of S_n , Lemma [3.1](#page-6-3) implies that all irreducible characters of S_n are real valued. Also it is elementary that $|K| = \frac{n!}{\prod_j j^{m_j} m_j!}$, where m_i is the number of cycles of length j of an element of K.

In order to upper bound the error terms in Theorem [3.13,](#page-12-0) the following estimate on the two step transition probabilities of the random walk generated by C will be useful.

Lemma 3.14. Let C be the conjugacy class of cycles of length i of the symmetric group S_n . Then for i fixed and $n \geq 2i$, $\frac{p_2(K)^2}{|K|}$ $\frac{d(K)}{|K|}$ is equal to

- (1) $i^2n^{-2i} + O(n^{-2i-1})$ if K is the identity conjugacy class.
- (2) $2i^2n^{-2i} + O(n^{-2i-1})$ if K is the conjugacy class consisting of two cycles of length i.
- (3) $O(n^{-2i-1})$ otherwise.

Proof. The first assertion is clear since if K is the identity class, $p_2(K) = \frac{1}{|C|}$. For the second assertion, note that $p_2(K) = 1 + O(n^{-1})$ since the only way that a product of two i-cycles is not in K is if the i-cycles have a symbol in common.

For the third assertion, there are two cases. The first case is that K has exactly $n-2i$ fixed points. Then $|K|$ is at least $c_i n^{2i}$ where c_i is a constant depending on i. Since K is not the class consisting of exactly two i-cycles,

 $p_2(K) = O(n^{-1})$ by the second assertion, proving the third assertion in this case. The second case is that K has $n - 2i + r$ fixed points, where $1 \leq r < 2i$. Then K is at least $c_i n^{2i-r}$ where c_i is a constant depending on *i*. However $p_2(K) = O(n^{-r})$, since in the product of the two i-cycles, there are r symbols moved by the first i-cycle each of which is mapped back to itself by the second i-cycle. Thus in this case also, the third assertion is \Box

Theorem 3.15. Let C be the conjugacy class of *i*-cycles in S_n , with $i \geq 2$. Choosing λ from the Plancherel measure of the symmetric group, define a $random\ variable\ W =$ $\sqrt{\binom{n}{i}(i-1)!}\chi^{\lambda}(C)$ $\frac{\lambda^{(1)} - \lambda^{(2)} \lambda^{(3)}}{\lambda^{(4)} + \lambda^{(5)}}$. Then there is a constant A_i such that for all real x_0 ,

$$
\left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \le A_i n^{-1/4}.
$$

Proof. One applies Theorem [3.13,](#page-12-0) choosing τ to be the irreducible representation corresponding to $\tau = (n-1,1)$, whose character value is the number of fixed points -1 . Then for any non-identity conjugacy class K,

$$
\frac{p_2(K)^2}{|K|} \left(\frac{\chi^{\tau}(K)}{\dim(\tau)} + 2a - 1 \right)^2 = O(n^{-2i-3}).
$$

This follows from Lemma [3.14](#page-13-1) and the fact that $\frac{\chi^{\tau}(K)}{dim(\tau)} + 2a - 1$ vanishes if K has $n-2i$ fixed points and is $O(n^{-1})$ for any other class K with $p_2(K) \neq 0$, since such a class has at least $n-2i$ fixed points. Since $|C| = \frac{n!}{(n-i)!}$ $\frac{n!}{(n-i)!i}$ and $a = \frac{i}{n}$ $\frac{i}{n-1}$, it follows that the first error term in Theorem [3.13](#page-12-0) is at most $A'_i n^{-1/2}$, where A'_i is a constant depending on *i*.

To bound the second error term in Theorem [3.13,](#page-12-0) observe that Lemma [3.14](#page-13-1) implies that

$$
\sum_{K} \left(8 - \frac{6}{a} \left(1 - \frac{\chi^{\tau}(K)}{\dim(\tau)} \right) \right) \frac{p_2(K)^2}{|K|}
$$

is $O(n^{-2i-1})$, since the n^{-2i} contributions from the identity class and the class of two i-cycles cancel. Hence the second error term is at most $A''_i n^{-1/4}$ where A''_i is a constant depending only on *i*.

4. Gelfand pairs

If G is a finite group and K is a subgroup of G such that the induced representation 1_K^G is multiplicity free, then the pair (G, K) is called a Gelfand pair. This section shows that the results of Section [3](#page-5-0) have an analog for Gelfand pairs.

Subsection [4.1](#page-15-0) discusses the representation theory of Gelfand pairs. Subsection [4.2](#page-17-0) derives a general central limit theorem for random spherical functions of a Gelfand pair. Subsection [4.3](#page-22-0) illustrates the theory on a toy example, proving a central limit theorem for the spectrum of the hypercube. A more serious example is considered in Subsection [4.4,](#page-24-0) which obtains a new central limit theorem for the spectrum of certain graphs whose vertices are the perfect matchings of 2n symbols.

4.1. Background from representation theory. We discuss some facts about the representation theory of Gelfand pairs. A useful reference is Chapter 7 of [\[Mac\]](#page-51-16).

The induced representation 1_K^G decomposes in a multiplicity free way as $\bigoplus_{r=0}^s V_r$, where V_0 denotes the trivial representation of G. Let d_r denote the dimension of V_r . For $0 \leq r \leq s$, let ω_r denote the corresponding spherical function on G , defined by

$$
\omega_r(x) = \frac{1}{|K|} \sum_{k \in K} \chi^{(r)}(x^{-1}k)
$$

where $\chi^{(r)}$ is the character of V_r . The functions ω_r are a basis of the space of functions on G which are constant on the double cosets of K in G . Let K_0, \dots, K_s denote the double cosets of K in G and let g_0, \dots, g_s be corresponding double coset representatives, so that $K_i = Kg_iK$. It is convenient to take g_0 to be the identity element of G. From page 389 of [\[Mac\]](#page-51-16), one has that $\omega_i(g_0) = 1$ for all i.

It is useful to recall facts which are analogs of those in Subsection [3.1.](#page-6-0)

Lemma 4.1. ([\[Mac\]](#page-51-16), page 389) Let ω be a spherical function of the Gelfand pair (G, K) . Then $\omega(x^{-1}) = \overline{\omega(x)}$.

The following two orthogonality relations are also useful.

Lemma 4.2. ([\[Mac\]](#page-51-16), page 389) For $0 \le i, j \le s$,

$$
\frac{d_i}{|G|} \sum_{r=0}^s |K_r| \omega_i(g_r) \overline{\omega_j(g_r)} = \delta_{i,j}.
$$

Lemma 4.3. For $0 \leq r, t \leq s$,

$$
\sum_{i=0}^{s} d_i \omega_i(g_r) \overline{\omega_i(g_t)} = \delta_{r,t} \frac{|G|}{|K_r|}.
$$

Proof. Consider the $s \times s$ matrix whose entry in the ith column and rth row is $\sqrt{\frac{d_i|K_r|}{n}}$ $\frac{|\mathbf{A}_r|}{|G|}\omega_i(g_r)$. By Lemma [4.2,](#page-15-1) the columns of this matrix are orthonormal. Hence so are its rows, proving the lemma. \square

If P is a K biinvariant probability on G (i.e. constant on double cosets of K in G), let $p_m(K_r)$ denote the probability that the m-fold convolution of P assigns to the double coset Kg_rK . Lemma [4.4](#page-15-2) is an analog of Lemma [3.4](#page-6-1) and could be proved along similar lines, as in [\[HSS\]](#page-50-19). Instead, we use the language of Fourier analysis, as developed on page 395 of [\[Mac\]](#page-51-16).

Lemma 4.4. Let P be the K biinvariant probability on G which associates mass 1 to the double coset K_u and mass 0 to all other K double cosets of G. Then for $0 \leq r \leq s$,

$$
p_m(K_r) = |K_r| \sum_{i=0}^s \frac{d_i}{|G|} \omega_i(g_u)^m \overline{\omega_i(g_r)}.
$$

Proof. If f is a complex valued K biinvariant function on G, define $\hat{f}(\omega_i)$ (the Fourier transform of f at the spherical function ω_i) by

$$
\hat{f}(\omega_i) = \sum_{g \in G} f(g) \overline{\omega_i(g)}.
$$

Then the Fourier inversion theorem gives that

$$
f = \frac{1}{|G|} \sum_{i=0}^{s} \hat{f}(\omega_i) d_i \omega_i.
$$

It is also true that the Fourier transform of the convolution of two K biinvariant functions is the product of their Fourier transforms. Thus the m-fold convolution of f is equal to

$$
\frac{1}{|G|} \sum_{i=0}^{s} \left(\hat{f}(\omega_i) \right)^m d_i \omega_i.
$$

The lemma now follows by taking $f = P$.

To connect the study of random spherical functions with spectral graph theory, suppose for convenience that (G, K) is a symmetric Gelfand pair, which means that $KgK = Kg^{-1}K$ for all g. This condition holds for the examples in Subsections [4.3](#page-22-0) and [4.4.](#page-24-0) Then fixing a double coset K_u , one can define a graph H_u whose vertices are the right cosets of K by connecting Kh_1 to Kh_2 if and only if $Kh_1h_2^{-1}K = K_u$. A more general construction appears in [\[Le\]](#page-51-17). The graph H_u is vertex transitive, since G acts transitively on the right cosets of K by sending Kh to Khg^{-1} , and Kh₁ is connected to Kh_2 if and only if Kh_1g^{-1} is connected to Kh_2g^{-1} . Lemma [4.5](#page-16-0) determines the spectrum of random walk on H_u .

Lemma 4.5. Let (G, K) be a symmetric Gelfand pair. Then for any double coset K_u , random walk on the graph H_u has eigenvalues $\omega_i(g_u)$ occurring with multiplicity d_i for $0 \leq i \leq s$.

Proof. Let M be the transition matrix for random walk on H_u . Since H_u is vertex transitive, the trace of M^k is $|G|/|K|$ multiplied by the probability that the random walk on H_u started at the right coset K is at K after k steps. By Theorem 7.5 of [\[Le\]](#page-51-17), this return probability is

$$
\frac{|K|}{|G|} \sum_{i=0}^{s} d_i \omega_i (g_u)^k,
$$

$$
\Box
$$

so that the trace of M^k is $\sum_{i=0}^s d_i \omega_i (g_u)^k$. Since this is true for all $k \geq 0$, the result follows.

Note that since $d_0 + \cdots + d_s = |G/K|$, one can define a probability measure on $\{0, \dots, s\}$ (or equivalently on the set $\{\omega_0, \dots, \omega_s\}$) by choosing i with probability $\frac{d_i|K|}{|G|}$. This is the Gelfand pair analog of the Plancherel measure of a finite group, and in this paper it will be referred to as Plancherel mea-sure. Lemma [4.5](#page-16-0) showed that if (G, K) is a symmetric Gelfand pair, then the eigenvalues of certain graphs on G/K occur with multiplicity proportional to Plancherel measure.

4.2. Central limit theorem for spherical functions. The aim of this subsection is to prove a central limit theorem for the random variable W defined by $W(i) = \frac{|K_u|^{1/2}}{|K|^{1/2}}$ $\frac{K_u|^{1/2}}{|K|^{1/2}}\omega_i(g_u)$. Here g_u is fixed satisfying $Kg_uK = Kg_u^{-1}K$ and i is random from the Plancherel measure of the Gelfand pair (G, K) . It is not assumed that the Gelfand pair (G, K) is symmetric.

To construct an exchangeable pair to be used for Stein's method, it is helpful to define a Markov chain on the set of spherical functions of (G, K) . To do this fix t with $1 \le t \le s$ such that the spherical function ω_t is realvalued. Let L_t be the Markov chain on the set $\{\omega_0, \dots, \omega_s\}$ which transitions from ω_i to ω_j with probability

$$
L_t(\omega_i, \omega_j) := \frac{d_j}{|G|} \sum_{r=0}^s |K_r| \omega_i(g_r) \omega_t(g_r) \overline{\omega_j(g_r)}.
$$

As in the group case, one typically wants to choose ω_t as close as possible to the trivial spherical function.

Lemma [4.6](#page-17-1) verifies that L_t is a Markov chain which is reversible with respect to Plancherel measure.

Lemma 4.6. Let t with $0 \leq t \leq s$ be such that the spherical function ω_t is real-valued. Then the transition probabilities of L_t are real and nonnegative and sum to 1. Moreover the chain L_t is reversible with respect to the Plancherel measure of the pair (G, K) .

Proof. From page 396 of [\[Mac\]](#page-51-16), if a_{it}^k are defined by

$$
\omega_i\omega_t=\sum_{k=0}^s a_{it}^k\omega_k
$$

(where the notation $\omega_i \omega_j$ denotes the pointwise product), then a_{it}^k are real and non-negative. Hence Lemma [4.2](#page-15-1) implies that $L_t(\omega_i, \omega_j)$ is real and nonnegative. Since $\omega_j(g_0) = 1$ for all j, Lemma [4.3](#page-15-3) gives that

$$
\sum_{j=0}^s L_t(\omega_i, \omega_j) = \sum_{r=0}^s |K_r| \omega_i(g_r) \omega_t(g_r) \frac{1}{|G|} \sum_{j=0}^s d_j \omega_j(g_0) \overline{\omega_j(g_r)} = 1.
$$

Reversibility of L_t with respect to Plancherel measure is equivalent to showing that

$$
\frac{d_i d_j |K|}{|G|} \sum_{r=0}^s |K_r| \omega_i(g_r) \omega_t(g_r) \overline{\omega_j(r)} = \frac{d_i d_j |K|}{|G|} \sum_{r=0}^s |K_r| \omega_j(g_r) \omega_t(g_r) \overline{\omega_i(r)}.
$$

Both sides are real by the previous paragraph, so the result follows by the assumption that ω_t is real valued.

An exchangeable pair (W, W') to be used in a Stein's method approach to studying W can be constructed from the chain L_t in the usual way. First choose *i* from Plancherel measure, then choose *j* with probability $L_t(\omega_i, \omega_j)$, and finally let $(W, W') = (W(\omega_i), W(\omega_j)).$

Lemma 4.7. $\mathbb{E}(W'|W) = \omega_t(g_u)W$.

Proof. From the definitions and Lemma [4.3,](#page-15-3)

$$
\mathbb{E}(W'|\omega_i) = \frac{|K_u|^{1/2}}{|K|^{1/2}} \sum_{j=0}^s \frac{d_j}{|G|} \sum_{r=0}^s |K_r| \omega_i(g_r) \omega_i(g_r) \overline{\omega_j(g_r)} \omega_j(g_u)
$$

\n
$$
= \frac{|K_u|^{1/2}}{|K|^{1/2}} \sum_{r=0}^s |K_r| \omega_i(g_r) \omega_t(g_r) \sum_{j=0}^s \frac{d_j}{|G|} \omega_j(g_u) \overline{\omega_j(g_r)}
$$

\n
$$
= \omega_t(g_u) W(\omega_i).
$$

The result follows since this depends on ω_i only through W. \Box

Corollary [4.8](#page-18-0) is not needed in the sequel but is interesting.

Corollary 4.8. The eigenvalues of L_t are $\omega_t(g_r)$ for $0 \le r \le s$. The functions $\psi_r(\omega_i) = \frac{|K_{gr}|^{1/2}}{|K|^{1/2}}$ $\frac{K_{gr}||\gamma}{|K|^{1/2}}\omega_i(g_r)$ are a basis of eigenvectors of L_{τ} , orthonormal with respect to the inner product

$$
\langle f_1, f_2 \rangle = \sum_{i=0}^s f_1(\omega_i) \overline{f_2(\omega_i)} \frac{d_i|K|}{|G|}.
$$

Proof. By the proof of Lemma [4.7,](#page-18-1) ψ_r is an eigenvector of L_t with eigenvalue $\omega_t(q_r)$. The orthonormality assertion follows from Lemma [4.3,](#page-15-3) and the basis assertion follows since the number of spherical functions is equal to the number of K double cosets of G .

Lemma 4.9. $\mathbb{E}(W'-W)^2 = 2(1-\omega_t(g_u)).$

Proof. This is immediate from Lemmas [2.3](#page-5-2) and [4.7.](#page-18-1) \Box

Lemma 4.10. $\mathbb{E}((W')^{2}|\omega_{i}) = \frac{|K_{u}|}{|K|} \sum_{r=0}^{s} \omega_{i}(g_{r}) \omega_{t}(g_{r}) p_{2}(K_{r}).$

Proof. From the definitions,

$$
\mathbb{E}((W')^2|\omega_i) = \frac{|K_u|}{|K|} \sum_{j=0}^s \frac{d_j}{|G|} \sum_{r=0}^s |K_r|\omega_i(g_r)\omega_t(g_r)\overline{\omega_j(g_r)}\omega_j(g_u)^2
$$

$$
= \frac{|K_u|}{|K|} \sum_{r=0}^s |K_r|\omega_i(g_r)\omega_t(g_r)\frac{1}{|G|} \sum_{j=0}^s d_j\omega_j(g_u)^2\overline{\omega_j(g_r)}.
$$

The result follows from Lemma [4.4.](#page-15-2) \Box

Note that the sum in Lemma [4.11](#page-19-0) begins at $r = 1$ and so excludes the $r = 0$ term which corresponds to the trivial spherical function.

Lemma 4.11.

$$
Var(\mathbb{E}[(W'-W)^2|\omega_i]) = \frac{|K_u|^2}{|K|} \sum_{r=1}^s \frac{p_2(K_r)^2}{|K_r|} (\omega_t(g_r) + 1 - 2\omega_t(g_u))^2.
$$

Proof. It follows from Lemmas [4.10,](#page-18-2) [4.9,](#page-18-3) and [4.7](#page-18-1) that

$$
Var(\mathbb{E}[(W'-W)^2|\omega_i])
$$

= $\mathbb{E}(\mathbb{E}((W'-W)^2|\omega_i)^2) - 4(1 - \omega_t(g_u))^2$
= $\frac{|K_u|^2}{|K|^2} \mathbb{E} \left[\sum_{r=0}^s p_2(K_r)\omega_i(g_r)\omega_t(g_r) + (1 - 2\omega_t(g_u))\omega_i(g_u)^2 \right]^2$
-4(1 - \omega_t(g_u))^2
= $T_1 + T_2 + T_3 - 4(1 - \omega_t(g_u))^2$

where

$$
T_1 = \frac{|K_u|^2}{|K|^2} \mathbb{E} \left(\sum_{r=0}^s p_2(K_r) \omega_i(g_r) \omega_t(g_r) \right)^2
$$

$$
T_2 = \frac{|K_u|^2}{|K|^2} 2(1 - 2\omega_t(g_u)) \sum_{r=0}^s p_2(K_r) \omega_t(g_r) \mathbb{E} \left(\omega_i(g_r) \omega_i(g_u)^2 \right)
$$

$$
T_3 = \frac{|K_u|^2}{|K|^2} (1 - 2\omega_t(g_u))^2 \mathbb{E}(\omega_i(g_u)^4).
$$

For reasons similar to the finite group case, there are useful simplifications. Lemma [4.3](#page-15-3) implies that

$$
T_1 = \frac{|K_u|^2}{|K|} \sum_{r=0}^{s} \frac{p_2(K_r)^2 \omega_t(g_r)^2}{|K_r|}.
$$

Lemma [4.4](#page-15-2) gives that

$$
T_2 = \frac{|K_u|^2}{|K|} 2(1 - 2\omega_t(g_u)) \sum_{r=0}^s \frac{p_2(K_r)^2 \omega_t(g_r)}{|K_r|}.
$$

Also, one can write

$$
T_3 = \frac{|K_u|^2}{|K|^2} (1 - 2\omega_t(g_u))^2 p_4(K_0) = \frac{|K_u|^2}{|K|} (1 - 2\omega_t(g_u))^2 \sum_{r=0}^s \frac{p_2(K_r)^2}{|K_r|}.
$$

The first equality used Lemma [4.4](#page-15-2) and the second equality used the assumption that $Kg_uK = Kg_u^{-1}K$. Hence

$$
T_1 + T_2 + T_3 = \frac{|K_u|^2}{|K|} \sum_{r=0}^s \frac{p_2(K_r)^2}{|K_r|} \left(\omega_t(g_r) + 1 - 2\omega_t(g_u)\right)^2,
$$

which implies the result since the $r = 0$ term is $4(1 - \omega_t(g_u))^2$.

Lemma 4.12. Let k be a positive integer.

(1) $\mathbb{E}(W'-W)^k$ is equal to $\sqrt{|K_u|}$ $|K|$ $\bigwedge^{k/2} \bigotimes^k$ $m=0$ $(-1)^{k-m}$ $\binom{k}{m}$ m $\Big\backslash \frac{s}{\sqrt{2}}$ $r=0$ $|K|$ $\frac{|X_r|}{|K_r|}\omega_t(g_r)p_m(K_r)p_{k-m}(K_r).$ (2) $\mathbb{E}(W'-W)^4 = \frac{|K_u|^2}{|K|}$ $\frac{|K_u|^2}{|K|} \sum_{r=0}^s \left[8\left(1-\omega_t(g_u)\right) - 6\left(1-\omega_t(g_r)\right) \right] \frac{p_2(K_r)^2}{|K_r|}$ $\frac{(K_r)^{-}}{|K_r|}$.

Proof. For the first assertion, note that $\mathbb{E}((W'-W)^k|\omega_i)$ is equal to

$$
\left(\frac{|K_u|}{|K|}\right)^{k/2} \sum_{j=0}^{s} \frac{d_j}{|G|} \sum_{r=0}^{s} |K_r| \omega_i(g_r) \omega_t(g_r) \overline{\omega_j(g_r)} (\omega_j(g_u) - \omega_i(g_u))^{k}
$$
\n
$$
= \left(\frac{|K_u|}{|K|}\right)^{k/2} \sum_{j=0}^{s} \frac{d_j}{|G|} \sum_{r=0}^{s} |K_r| \omega_i(g_r) \omega_t(g_r) \overline{\omega_j(g_r)}
$$
\n
$$
\cdot \sum_{m=0}^{k} (-1)^{k-m} {k \choose m} \omega_j(g_u)^m \omega_i(g_u)^{k-m}
$$
\n
$$
= \left(\frac{|K_u|}{|K|}\right)^{k/2} \sum_{m=0}^{k} (-1)^{k-m} {k \choose m} \omega_i(g_u)^{k-m}
$$
\n
$$
\cdot \sum_{r=0}^{s} |K_r| \omega_i(g_r) \omega_t(g_r) \sum_{j=0}^{s} \frac{d_j}{|G|} \omega_j(g_u)^m \overline{\omega_j(g_r)}
$$
\n
$$
= \left(\frac{|K_u|}{|K|}\right)^{k/2} \sum_{m=0}^{k} (-1)^{k-m} {k \choose m} \omega_i(g_u)^{k-m} \sum_{r=0}^{s} \omega_i(g_r) \omega_t(g_r) p_m(K_r),
$$

 \Box

where the last equality is Lemma [4.4.](#page-15-2) Hence

$$
\mathbb{E}(W'-W)^k = \mathbb{E}(\mathbb{E}((W'-W)^k|\omega_i))
$$

\n
$$
= \left(\frac{|K_u|}{|K|}\right)^{k/2} \sum_{m=0}^k (-1)^{k-m} {k \choose m}
$$

\n
$$
\cdot \sum_{r=0}^s \omega_t(g_r) p_m(K_r) \sum_{i=0}^s \frac{d_i|K|}{|G|} \omega_i(g_r) \omega_i(g_u)^{k-m}.
$$

The first assertion follows from Lemma [4.4](#page-15-2) and the fact that $\omega_i(g_u)$ is real.

For the second assertion, one knows from the first assertion that

$$
\mathbb{E}(W'-W)^4 = \left(\frac{|K_u|}{|K|}\right)^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_{r=0}^s \frac{|K|}{|K_r|} \omega_t(g_r) p_m(K_r) p_{4-m}(K_r).
$$

The special case $t = 0$ gives the equation

$$
0 = \left(\frac{|K_u|}{|K|}\right)^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_{r=0}^s \frac{|K|}{|K_r|} p_m(K_r) p_{4-m}(K_r).
$$

Hence in general, $\mathbb{E}(W'-W)^4$ is equal to

$$
-\frac{|K_u|^2}{|K|^2}\sum_{m=0}^4(-1)^m\binom{4}{m}\sum_{r=0}^s\frac{|K|}{|K_r|}(1-\omega_t(g_r))p_m(K_r)p_{4-m}(K_r).
$$

The $m = 0, 4$ terms both vanish since $p_0(K_r) = 0$ if $r \neq 0$. The $m = 2$ term is equal to

$$
-\frac{6|K_u|^2}{|K|}\sum_{r=0}^s\frac{(1-\omega_t(g_r))p_2(K_r)^2}{|K_r|}.
$$

The $m = 1, 3$ terms are equal and together contribute

$$
\frac{8|K_u|^2}{|K|} \sum_{r=0}^s \frac{(1-\omega_t(g_r))p_1(K_r)p_3(K_r)}{|K_r|} = \frac{8|K_u|}{|K|}(1-\omega_t(g_u))p_3(K_u)
$$

$$
= \frac{8|K_u|^2}{|K|^2}(1-\omega_t(g_u))p_4(K_0)
$$

$$
= \frac{8|K_u|^2}{|K|}(1-\omega_t(g_u))\sum_{r=0}^s \frac{p_2(K_r)^2}{|K_r|}.
$$

This completes the proof of the second assertion.

Arguing as in the proof of Theorem [3.13,](#page-12-0) and using the above lemmas, one obtains the following result.

Theorem 4.13. Let (G, K) be a Gelfand pair, and fix a double coset $K_u =$ Kg_uK of G satisfying $Kg_uK = Kg_u^{-1}K$. Let ω_t be a nontrivial spherical function which is real valued. Choosing ω_i from the Plancherel measure of the pair (G, K) , let $W = \frac{|K_u|^{1/2}}{|K|^{1/2}}$ $\frac{K_{u}+\kappa}{|K|^{1/2}}\omega_i(g_u)$. Then for all real x_0 ,

$$
\begin{split}\n&\quad \left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \\
&\le \frac{|K_u|}{a|K|} \sqrt{\sum_{r=1}^s \frac{|K|p_2(K_r)^2}{|K_r|} (\omega_t(g_r) + 2a - 1)^2} \\
&\quad + \frac{1}{(\pi)^{1/4}} \frac{|K_u|^{1/2}}{|K|^{1/2}} \left[\sum_{r=0}^s \left(8 - \frac{6(1 - \omega_t(g_r))}{a} \right) \frac{|K|p_2(K_r)^2}{|K_r|} \right]^{1/4},\n\end{split}
$$

where $a = 1 - \omega_t(g_u)$.

4.3. Example: The Hypercube. The *n* dimensional hypercube \mathbb{Z}_2^n consists of n-tuples of 0's and 1's. Random walk on it proceeds by picking a random coordinate and changing it. The spectrum of this random walk is well known; for each $0 \leq i \leq n$ there is an eigenvalue $1 - \frac{2i}{n}$ $\frac{2i}{n}$ occurring with multiplicity $\binom{n}{i}$ $\binom{n}{i}$ (see for instance page 28 of [\[Di\]](#page-50-16)). Thus by the usual central limit theorem for the binomial distribution, the spectrum of the hypercube is asymptotically normal with an error term $O(n^{-1/2})$. The purpose of this subsection is to revisit this classical result from the viewpoint of Gelfand pairs, illustrating the construction of Subsection [4.2.](#page-17-0)

To begin, note from the remarks on page 58 of [\[Di\]](#page-50-16), that the hypercube can be viewed as G/K for a certain Gelfand pair (G, K) . Namely G is the semidirect product of \mathbb{Z}_2^n with S_n , where the group multiplication is $(x, \pi)(y, \tau) = (x + \pi(y), \pi\tau)$, where $\tau(x)$ permutes the coordinates of x. K is the subgroup $\{(0, \pi) : \pi \in S_n\}$. The induced module 1_K^G decomposes as $\bigoplus_{r=0}^{n} V_r$ where $d_r = \binom{n}{r}$ ⁿ, Thus the Plancherel measure chooses $i \in \{0, \dots, n\}$ with probability $\frac{\binom{n}{i}}{2^n}$. For $0 \leq r \leq n$, the double coset K_r in G consists of elements (x, π) where x has r coordinates equal to 1. Thus $|K_r| = \binom{n}{r}$ $\binom{n}{r}n!$ and as usual let g_r denote some element of K_r . The spherical function ω_i $(0 \leq i \leq n)$ is given by

$$
\omega_i(g_r) = \frac{1}{\binom{n}{i}} \sum_{m=0}^i (-1)^m \binom{r}{m} \binom{n-r}{i-m},
$$

which is a Krawtchouk polynomial if one overlooks the $\binom{n}{i}$ $\binom{n}{i}$ in the denominator.

The following result emerges from Theorem [4.13.](#page-21-0) Since $\omega_i(g_1) = 1 - \frac{2i}{n}$ $\frac{2i}{n}$ (or by Lemma [4.5\)](#page-16-0), it can be seen as a central limit theorem for the spectrum of random walk on the hypercube.

Theorem 4.14. Let $W = \sqrt{n}\omega_i(g_1)$ where i is chosen from Plancherel measure. Then for all real x_0 ,

$$
\left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \le \left(\frac{8}{\pi n}\right)^{1/4}.
$$

Proof. Apply Theorem [4.13](#page-21-0) with $u = 1$ and $t = 1$. Then $a = \frac{2}{n}$ $\frac{2}{n}$. One computes that $p_2(K_0) = \frac{1}{n}$, $p_2(K_2) = 1 - \frac{1}{n}$ $\frac{1}{n}$, and that $p_2(K_r) = 0$ for all $r \neq 0, 2$. Since $\omega_1(g_2) = 1 - \frac{4}{n}$ $\frac{4}{n}$, one has that $\omega_1(g_2) + 2a - 1 = 0$. Thus the first error term in Theorem [4.13](#page-21-0) is 0. The second error term in Theorem [4.13](#page-21-0) is computed to be $\left(\frac{8}{\pi n}\right)^{1/4}$, implying the result.

To get $O(n^{-1/2})$ bounds by Stein's method, it will be shown that $|W'-W|$ is bounded, so that one can use the version of Stein's method in Theorem [2.2](#page-4-2) instead of that in Theorem [2.1.](#page-4-1) The boundedness of $|W'-W|$ will follow from Lemma [4.15,](#page-23-0) which proves that L_1 is in fact a birth-death chain.

Lemma 4.15. The chain L_1 on the set $\{0, \dots, n\}$ is a birth-death chain with transition probabilities

$$
L_1(i,j) = \begin{cases} \frac{i}{n} & \text{if } j = i - 1 \\ 1 - \frac{i}{n} & \text{if } j = i + 1 \end{cases}
$$

Proof. From the three term recurrence for Krawtchouk polynomials on page 152 of [\[MaSl\]](#page-51-18), it follows that

$$
(i+1)\binom{n}{i+1}\omega_{i+1}(g_r) = (n-2r)\binom{n}{i}\omega_i(g_r) - (n-i+1)\binom{n}{i-1}\omega_{i-1}(g_r).
$$

Since $n - 2r = n\omega_1(g_r)$, one obtains that

$$
n\binom{n}{i}\omega_1(g_r)\omega_i(g_r) = (i+1)\binom{n}{i+1}\omega_{i+1}(g_r) + (n-i+1)\binom{n}{i-1}\omega_{i-1}(g_r).
$$

Simplifying one obtains the relation

$$
\omega_1(g_r)\omega_i(g_r) = \left(1 - \frac{i}{n}\right)\omega_{i+1}(g_r) + \left(\frac{i}{n}\right)\omega_{i-1}(g_r).
$$

The result now follows from Lemma [4.2](#page-15-1) and the definition of L_1 .

Now an $O(n^{-1/2})$ error term is established.

Theorem 4.16. Let $W = \sqrt{n}\omega_i(g_1)$ where i is chosen from Plancherel measure. Then for all real x_0 ,

$$
\left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \le 5n^{-1/2}.
$$

Proof. Apply the variation of Theorem [4.13](#page-21-0) which would arise from using Theorem [2.2](#page-4-2) instead of Theorem [2.1.](#page-4-1) Recall that $a = \frac{2}{n}$ $\frac{2}{n}$. Also $|W' - W|$ ≤ A with $A = \frac{2}{\sqrt{2}}$ $\frac{1}{n}$ since L_1 is a birth-death chain. As explained in the proof of Theorem [4.14,](#page-22-1) $Var(\mathbb{E}[(W'-W)^2|W]) = 0$. The result follows.

A a final remark, Lemma [4.15](#page-23-0) shows that L_1 is closely related to the birth-death chain used by Stein [\[Stn1\]](#page-51-10) in proving a central limit theorem for $X_1 + \dots + X_n$, where the $X's$ are independent and each equal to 0 or 1 with probability 1/2. To form an exchangeable pair Stein chose a random $l \in \{1, \dots, n\}$ and then replaced X_l by a new random variable which is also 0 or 1 with probability $1/2$ and independent of all of the $X's$. The chain L_1 would arise by picking a random index l and switching the value of X_l .

4.4. Example: Graphs on Perfect Matchings. In this example $G = S_{2n}$ and K is the hyperoctahedral group of signed permutations on n symbols, of size $2ⁿn!$. This Gelfand pair is discussed at length in Chapter 7 of [\[Mac\]](#page-51-16) and in Section 3 of [\[HSS\]](#page-50-19), to which we refer the reader for proofs of facts in the next two paragraphs. Another useful reference is [\[DHol\]](#page-50-20).

The induced representation 1_K^G decomposes as $\bigoplus_{|\lambda|=n} V_{\lambda}$, where λ ranges over all partitions of size *n*. An explicit formula for the numbers d_{λ} appears in [\[Mac\]](#page-51-16) and is not needed in what follows. However it is worth remarking that the Plancherel measure of (G, K) , which chooses λ with probability $\frac{2^{n} n! d_{\lambda}}{(2n)!}$, is the $\alpha = 2$ case of the so called Jack_a measure on partitions, which chooses λ with probability

$$
\frac{\alpha^n n!}{\prod_{s\in\lambda}(\alpha a(s) + l(s) + 1)(\alpha a(s) + l(s) + \alpha)}
$$

where the product is over all boxes of the partition. Here $a(s)$ is the number of boxes in the same row of s and to the right of s (the "arm" of s), and $l(s)$ is the number of boxes in the same column of s and below s (the "leg" of s). For example the partition of 5 below

$$
\begin{array}{c}\square\ \square\ \square\\ \square\ \square\end{array}
$$

would have Jack_{α} measure

$$
\frac{60\alpha^2}{(2\alpha+2)(3\alpha+1)(\alpha+2)(2\alpha+1)(\alpha+1)}.
$$

The Jack measure on partitions is of interest to researchers in random matrix theory $[BO1], [O2], [K2].$ $[BO1], [O2], [K2].$ $[BO1], [O2], [K2].$ $[BO1], [O2], [K2].$ $[BO1], [O2], [K2].$

The double cosets K_{μ} of K in G are parameterized by partitions μ of size n and have the following concrete description. A perfect matching of $\{1, \dots, 2n\}$ can be regarded as a 1-regular graph with vertex set $\{1, \dots, 2n\}$. Let ϵ be the "identity matching" in which i is adjacent to $n + i$ for $1 \leq i \leq$ n. Given a permutation w in S_{2n} , let $\delta(w)$ be the perfect matching of $\{1, \dots, 2n\}$ in which i is adjacent to j if and only $|w(i) - w(j)| = n$. For example, $\delta(id) = \epsilon$. Note that the union $\delta_1 \cup \delta_2$ of two 1-regular graphs is a 2regular graph, and thus a disjoint union of even length cycles. Let $\Lambda(\delta_1, \delta_2)$ be the partition whose parts are half the cycle lengths of $\delta_1 \cup \delta_2$. Then $Kw_1K = Kw_2K$ if and only if $\Lambda(\epsilon, \delta(w_1)) = \Lambda(\epsilon, \delta(w_2))$. Thus $Kw^{-1}K =$ KwK for all w, so that the machinery of Subsection [4.2](#page-17-0) is applicable. One also has that $Kw_1 = Kw_2$ if and only if $\delta(w_1) = \delta(w_2)$. It follows that G/K is in bijection with the perfect matchings of $\{1, \dots, 2n\}$ and that $\frac{|K_\mu|}{|K|}$ is equal to the number of perfect matchings δ with $\Lambda(\epsilon, \delta) = \mu$, which by elementary counting is $\frac{2^n n!}{2^n n!}$ $\frac{2^{n}n!}{2^{l(\mu)}\prod_j m_j(\mu)!j^{m_j(\mu)}}$ where $l(\mu)$ is the number of parts of μ and $m_i(\mu)$ is the number of parts of μ of size j.

The main purpose of this subsection is to prove a central limit theorem for the random variable $W = \sqrt{2^{i-1} \binom{n}{i}}$ $\sum_{i=1}^{n} (i-1)! \omega_{\lambda}(g_{(i,1^{n-i})}),$ where i is fixed and λ is chosen from the Plancherel measure of (G, K) . Then Lemma [4.5](#page-16-0) gives a central limit theorem for the spectrum of random walk on the graph $H_{(i,1^{n-i})}$, whose vertices are the perfect matchings of $\{1,\dots,2n\}$, with an edge between matchings δ_1 and δ_2 if and only if $\Lambda(\delta_1, \delta_2) = (i, 1^{n-i})$. In the case $i = 2$, the spectrum of this graph was determined in [\[DHol\]](#page-50-20), and was shown to be asymptotically normal in [\[F2\]](#page-50-8) (with error term $O(n^{-1/4})$) and then in [\[CF\]](#page-50-5), [\[F3\]](#page-50-6) (with error term $O(n^{-1/2})$). But the arguments in those papers used different information, and it was not clear that they could be pushed through to larger i. Theorem [4.13](#page-21-0) will be used to deduce a central limit theorem for any fixed i with error term $O(n^{-1/4})$.

To apply Theorem [4.13,](#page-21-0) ω_t will be taken to be the spherical function $\omega_{(n-1,1)}$. An explicit formula for $\omega_{(n-1,1)}$ is available.

Lemma 4.17. (Mac), p. 411) Let $m_1(\mu)$ denote the number of parts of size 1 of μ . Then

$$
\omega_{(n-1,1)}(g_{\mu}) = \frac{(2n-1)m_1(\mu) - n}{2n(n-1)}
$$

for all μ of size n.

Lemma [4.18](#page-25-0) is helpful.

Lemma 4.18. ([\[HSS\]](#page-50-19), Lemma 3.2) The coefficient of K_{μ} in $K_{\tau}K_{(i,1^{n-i})}$ is equal to $\frac{|K|^2}{|K_\mu|}$ multiplied by the number of pairs of perfect matchings (δ, γ) such that $\Lambda(\epsilon,\delta) = (i,1^{n-i}), \Lambda(\delta,\gamma) = \tau, \text{ and } \Lambda(\epsilon,\gamma) = \mu.$

The final combinatorial ingredient is an analog of Lemma [3.14.](#page-13-1)

Lemma 4.19. Consider the random walk on G generated by $K_{(i,1^{n-i})}$. Then for i fixed and $n \geq 2i$, $\frac{p_2(K_\mu)^2|K|}{|K_\mu|}$ $\frac{K_{\mu} \prod |K|}{|K_{\mu}|}$ is equal to

- $(1) \frac{i^2}{4i}$ $\frac{i^2}{4^i-1}n^{-2i}+O(n^{-2i-1})$ if $K_\mu=K_{(1^n)}=K$.
- $(2) \frac{2i^2}{4i}$ $\frac{2i^2}{4^{i-1}}n^{-2i} + O(n^{-2i-1})$ if $K_\mu = K_{(i,i,1^{n-2i})}$.
- (3) $O(n^{-2i-1})$ otherwise.

Proof. Lemma [4.18](#page-25-0) implies that $\frac{p_2(K_\mu)|K_{(i,1^{n-i})}|^2}{|K|^2}$ $\frac{|K|^2 (k+1)^{1/2}}{|K|^2}$ is equal to the number of pairs of matchings (δ, γ) such that $\Lambda(\epsilon, \delta) = (i, 1^{n-i}), \Lambda(\delta, \gamma) = (i, 1^{n-i}),$ and $\Lambda(\epsilon, \gamma) = \mu$.

For the first assertion, it follows either from the previous paragraph or from Lemmas [4.4](#page-15-2) and [4.3](#page-15-3) that $p_2(K) = \frac{1}{2^{i-1} {n \choose i} (i-1)!}$. For the second assertion, it is straightforward from the previous paragraph that $p_2(K_{(i,i,1^{n-2i})}) =$ $1 + O(n^{-1})$. Then use the formula for $\frac{|K_\mu|}{|K|}$ given earlier in this subsection.

For the third assertion, there are two cases. The first case is that μ has $n-2i$ parts of size 1, but is not equal to $(i, i, 1^{n-2i})$. Then $p_2(K_\mu) = O(n^{-1})$ and the formula for $\frac{|K|}{|K_{\mu}|}$ shows it to be at most $c_i n^{-2i}$ where c_i is a constant depending on i . So in the first case, the result is proved. The second case is that μ has $n - 2i + r$ parts of size 1, where $1 \leq r < 2i$. From the first paragraph of the proof, it is straightforward to see that $p_2(K_\mu) = O(n^{-r}).$ Also the formula for $\frac{|K|}{|K_{\mu}|}$ shows it to be $O(n^{r-2i})$, proving the result. \Box

Combining the ingredients, one deduces the following result.

Theorem 4.20. Choose λ from the Plancherel measure of the Gelfand pair (G, K) , and define a random variable $W = \sqrt{2^{i-1} {n \choose i}}$ $\sum_{i}^{n} (i-1)! \omega_{\lambda}(g_{(i,1^{n-i})}).$ Then there is a constant A_i such that for all real x_0 ,

$$
\left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \le A_i n^{-1/4}.
$$

Proof. One applies Theorem [4.13,](#page-21-0) choosing $u = (i, 1^{n-i})$ and ω_t to be the spherical function $\omega_{(n-1,1)}$. By Lemma [4.17,](#page-25-1) ω_t is real valued and $\alpha =$ $\frac{i(2n-1)}{2n(n-1)}$. Observe that $\omega_t(g_\mu) + 2a - 1$ vanishes if $\mu = (i, i, 1^{n-2i})$ and by Lemma [4.17](#page-25-1) is $O(n^{-1})$ for any other μ such that $p_2(K_{\mu}) \neq 0$, since any such μ has at least $n-2i$ parts of size 1. Thus by Lemma [4.19,](#page-25-2)

$$
\sum_{\mu \neq (1^n)} \frac{|K| p_2(K_\mu)^2}{|K_\mu|} (\omega_t(g_\mu) + 2a - 1)^2 = O(n^{-2i-3}).
$$

Since $\frac{|K_u|}{|K|} = 2^{i-1} {n \choose i}$ $\binom{n}{i}(i-1)!$, the first error term in Theorem [4.13](#page-21-0) is at most $A'_i n^{-1/2}$, where A'_i is a constant depending on *i*.

To bound the second error term in Theorem [4.13,](#page-21-0) one applies Lemma [4.19](#page-25-2) to see that

$$
\sum_{\mu} \left(8 - \frac{6(1 - \omega_t(g_{\mu}))}{a} \right) \frac{|K| p_2(K_{\mu})^2}{|K_{\mu}|} = O(n^{-2i-1}).
$$

Note that cancellation occurred for the coefficient of n^{-2i} coming from the terms $\mu = (1^n)$ and $\mu = (i, i, 1^{n-2i})$. It follows that the second error term is at most $A''_i n^{-1/4}$, where A''_i is another constant depending only on i. \Box

5. Twisted Gelfand Pairs

Let G be a finite group and H a subgroup of G. If ϕ is a linear character of H such that the induced representation $Ind_H^G(\phi)$ is multiplicity free, then (G, H, ϕ) is referred to as a twisted Gelfand pair (this terminology was introduced in [\[Stm\]](#page-51-5)).

The approach taken in this section is a bit different than that in the sections on finite groups and Gelfand pairs. Due both to a lack of interesting examples which are not Gelfand pairs and to technical complications which do not arise for finite groups or Gelfand pairs, a completely general theory is not developed. Instead, we focus on one very interesting example: character values of random projective representations of the symmetric group. It should however be noted that many of the lemmas resemble those of earlier sections, and the calculations are organized in a way which should generalize to other examples.

It is known (see [\[HH\]](#page-50-21) for a friendly exposition) that the character values in question are expressed in terms of the coefficients X_{ρ}^{λ} , where λ is a strict (i.e. all parts distinct) partition of n that parameterizes an irreducible character, and μ is an odd (i.e. all parts odd) partition of n that parameterizes a conjugacy class. Throughout this section $DP(n)$ will denote the set of strict partitions of size n, and $OP(n)$ will denote the set of odd partitions of size $\mathfrak n$.

Central limit theorems for random projective representations of the symmetric group have been studied by Ivanov [\[I\]](#page-50-9). The underlying probability measure on strict partitions chooses λ with probability

$$
\frac{2^{n-l(\lambda)}g_{\lambda}^2}{n!},
$$

where $l(\lambda)$ is the number of parts of a partition λ and $g_{\lambda} = X^{\lambda}_{(1^n)}$ is the number of standard shifted tableaux of shape λ ([\[HH\]](#page-50-21),[\[Mac\]](#page-51-16)). This measure on strict partitions is known as shifted Plancherel measure, and Ivanov proves the following result.

Theorem 5.1. ([\[I\]](#page-50-9)) Fix $i \geq 1$. Let λ be chosen from the shifted Plancherel measure on strict partitions of size n. Then as $n \to \infty$, the random variable $n^{\frac{2i+1}{2}} X^{\lambda}_{(2i+1,1^{n-2i-1})}$

 $\frac{(2i+1,1^{n-2i-1})}{2i\sqrt{2i+1}g_{\lambda}}$ converges in distribution to a normal random variable with mean θ and variance 1.

This section refines the result of Ivanov, which was proved by the method of moments, so as to obtain an error term. In the statement of the result, recall that $z_{\mu} = \prod_{j\geq 1} j^{m_j(\mu)} m_j(\mu)!,$ where $m_j(\mu)$ is the number of parts of μ of size *j*.

Theorem 5.2. Fix $i \geq 1$ and let $\mu = (2i + 1, 1^{n-2i-1})$. Choosing λ from the shifted Plancherel measure on partitions of size n, define a random variable

$$
W = \sqrt{\frac{n!}{z_{\mu}2^{n-l(\mu)}}} \frac{X_{\mu}^{\lambda}}{g_{\lambda}}.
$$

Then there is a constant A_i such that for all real x_0 ,

$$
\left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \le A_i n^{-1/4}.
$$

The organization of this section is as follows. Subsection [5.1](#page-28-0) collects and develops background from representation theory. Subsection [5.2](#page-31-0) defines and studies a Markov chain to be used in the construction of an exchangeable pair for a Stein's method proof of Theorem [5.2.](#page-27-0) This is more subtle than the corresponding treatment in earlier sections and involves interesting combinatorics, since the "obvious" adaptation of the construction for Gelfand pairs does not work. Subsection [5.3](#page-33-0) studies the exchangeable pair arising from the Markov chain in Section [5.2,](#page-31-0) and uses it to prove Theorem [5.2.](#page-27-0)

5.1. Background from representation theory. If G is a finite group, K a subgroup of G, and ϕ a linear character of K such that $Ind_K^G(\phi)$ is multiplicity free, the triple (G, K, ϕ) is called a twisted Gelfand pair. The Hecke algebra of the triple (G, K, ϕ) is the CG subalgebra $e \mathbb{C} G e$, where e is the primitive idempotent of $\mathbb{C}K$ defined by

$$
e = \frac{1}{|K|} \sum_{k \in K} \phi(k^{-1}) k.
$$

One reason that twisted Gelfand pairs are interesting is that the Hecke algebra of the triple (G, K, ϕ) is commutative. The paper [\[Stm\]](#page-51-5) is a good reference for the theory of twisted Gelfand pairs.

The twisted Gelfand pair of interest to us is the one studied in [\[Stm\]](#page-51-5). Thus $G = S_{2n}$ and K is the hyperoctahedral group B_n of signed permutations, imbedded in G as the centralizer of the involution $(1, 2)(3, 4) \cdots (2n-1, 2n)$. To define ϕ , note that B_n is the semidirect product of the groups T_n and Σ_n , where T_n is the subgroup (isomorphic to \mathbb{Z}_2^n) generated by $(1, 2), \cdots, (2n -$ 1, 2n) and Σ_n is the subgroup (isomorphic to S_n) generated by the "double" transpositions" $(2i-1, 2j-1)(2i, 2j)$ for $1 \leq i < j \leq n$. Then ϕ is the linear character of B_n whose restriction to Σ_n is the sign character, and whose restriction to T_n is trivial.

Stembridge [\[Stm\]](#page-51-5) defines twisted spherical functions for twisted Gelfand pairs (the analog of spherical functions for Gelfand pairs). One of his main results is that for (S_{2n}, B_n, ϕ) , their values are (aside from scalar multiples) equal to the X^{λ}_{μ} , where λ is a strict partition and μ is an odd partition.

Next we record orthogonality relations for the quantities X_{μ}^{λ} . These orthogonality relations are a special case of more general orthogonality relations for coefficients of power sum symmetric functions in Hall-Littlewood polynomials, where the parameter in the Hall Littlewood polynomial is −1. What this paper calls X^{λ}_{μ} is written as $X^{\lambda}_{\mu}(-1)$ in Chapter 3 of [\[Mac\]](#page-51-16).

Lemma 5.3. ([\[Mac\]](#page-51-16), page 247) For $\lambda, \rho \in DP(n)$,

$$
\sum_{\in OP(n)} \frac{2^{l(\mu)}}{z_{\mu}} X_{\mu}^{\lambda} X_{\mu}^{\rho} = \delta_{\lambda, \rho} 2^{l(\lambda)}.
$$

Lemma 5.4. ([\[Mac\]](#page-51-16), page 247) For $\mu, \sigma \in OP(n)$,

 μ

$$
\sum_{\lambda \in DP(n)} \frac{1}{2^{l(\lambda)}} X_{\mu}^{\lambda} X_{\sigma}^{\lambda} = \delta_{\mu,\sigma} \frac{z_{\mu}}{2^{l(\mu)}}.
$$

As in Subsection [4.4,](#page-24-0) the double cosets of B_n in S_{2n} are indexed by partitions ν of n. It is useful to specify representatives w_{ν} . For the case $\nu = (n)$, one defines

$$
w_{(n)}=(1,2,\cdots,2n),
$$

and for the general case $\nu = (\nu_1, \dots, \nu_l)$, one defines

$$
\omega_{\nu} = w_{(\nu_1)} \circ \cdots \circ w_{(\nu_l)}
$$

where the operation $x \circ y$ (for $x \in S_{2i}, y \in S_{2j}$) denotes the embedding of $S_{2i} \times S_{2j}$ in S_{2i+2j} with S_{2i} acting on $\{1, \dots, 2i\}$ and S_{2j} acting on ${2i+1, \cdots, 2i+2j}.$

For $\nu \in OP(n)$, define

$$
\tilde{K}_{\nu} = \frac{1}{|B_n|^2} \sum_{x_1, x_2 \in B_n} \phi(x_1 x_2) x_1 w_{\nu} x_2.
$$

Corollary 3.2 of [\[Stm\]](#page-51-5) shows that if $\nu \notin OP(n)$, then $\tilde{K}_{\nu} = 0$, and that $\{\tilde{K}_{\nu} : \nu \in OP(n)\}\$ is a basis for the Hecke algebra of (S_{2n}, B_n, ϕ) . Thus for $\mu, \nu \in OP(n)$, it is natural to study the coefficient of \tilde{K}_{ν} in $(\tilde{K}_{\mu})^m$. Lemma [5.5](#page-29-0) gives a character theoretic expression for this coefficient and is analogous to Lemmas [3.4](#page-6-1) and [4.4.](#page-15-2)

Lemma 5.5. Suppose that $\mu_1, \dots, \mu_m, \nu \in OP(n)$. Then the coefficient of \tilde{K}_{ν} in $\tilde{K}_{\mu_1} \cdots \tilde{K}_{\mu_m}$ is equal to

$$
\frac{2^{l(\mu_1)-n} \cdots 2^{l(\mu_m)-n}}{z_{\nu}} \sum_{\rho \in DP(n)} 2^{n-l(\rho)} g_{\rho}^2 \frac{X_{\nu}^{\rho}}{g_{\rho}} \frac{X_{\mu_1}^{\rho}}{g_{\rho}} \cdots \frac{X_{\mu_m}^{\rho}}{g_{\rho}}
$$

.

In particular, the coefficient of \tilde{K}_{ν} in $(\tilde{K}_{\mu})^m$ is

$$
\frac{n! (2^{l(\mu)-n})^m}{z_{\nu}} \mathbb{E}\left[\frac{X^{\rho}_{\nu}}{g_{\rho}}\left(\frac{X^{\rho}_{\mu}}{g_{\rho}}\right)^m\right],
$$

where ρ is random from shifted Plancherel measure on partitions of size n.

Proof. The equality is immediate from the definition of shifted Plancherel measure, so it is enough to establish the first expression. Stembridge [\[Stm\]](#page-51-5) provides a basis of orthogonal idempotents $\{E_{\rho} : \rho \in DP(n)\}\$ of the Hecke algebra of (G, H, ϕ) in terms of the projective characters of the symmetric

group. More precisely, it follows from Proposition 4.1 and Corollary 6.2 of [\[Stm\]](#page-51-5) that one obtains such a basis of orthogonal idempotents by defining

$$
E_{\rho}=2^{n-l(\rho)}g_{\rho}\sum_{\mu\in OP(n)}\frac{X^{\rho}_{\mu}}{z_{\mu}}\tilde{K}_{\mu}.
$$

Multiplying both sides of this equation by $2^{l(\mu)-n}\frac{X_{\mu}^{\rho}}{g_{\rho}}$ and summing over all $\rho \in DP(n)$, it follows from Lemma [5.4](#page-29-1) that

$$
\tilde{K}_{\mu} = 2^{l(\mu)-n} \sum_{\rho \in DP(n)} \frac{X_{\mu}^{\rho}}{g_{\rho}} E_{\rho}.
$$

Thus

$$
\tilde{K}_{\mu_1} \cdots \tilde{K}_{\mu_m} = 2^{l(\mu_1)-n} \cdots 2^{l(\mu_m)-n} \sum_{\rho \in DP(n)} \frac{X_{\mu_1}^{\rho}}{g_{\rho}} \cdots \frac{X_{\mu_m}^{\rho}}{g_{\rho}} E_{\rho}.
$$

The result now follows by the formula for E_{ρ} in the previous paragraph. \Box

Lemma [5.6](#page-30-0) derives an explicit formula for $X^{(n-1,1)}_{\mu}$. This is crucial to our approach.

Lemma 5.6. Let $m_1(\mu)$ denote the number of parts of size 1 of μ . Suppose that $n \geq 3$, so that $(n-1,1)$ is a strict partition. Then

$$
X_{\mu}^{(n-1,1)} = m_1(\mu) - 2
$$

for all odd partitions μ of size n.

Proof. Recall the definition of power sum symmetric functions: for $i \geq 1$, one sets $p_i = \sum_j x_j^i$, and for μ a partition of n, one sets $p_{\mu} = \prod_i p_i^{m_i(\mu)}$ $i^{m_i(\mu)}$. The argument also uses Schur's Q-functions([\[Mac\]](#page-51-16), Sec. 3.8). More precisely, equation 7.5 on page 247 of [\[Mac\]](#page-51-16) shows that for $\mu \in OP(n)$ and $\lambda \in DP(n)$, the coefficient of the power sum symmetric function p_{μ} in Q_{λ} is equal to $2^{l(\mu)}X^\lambda_\mu$ $\frac{z_1}{z_\mu}$.

To proceed one needs an expression for $Q_{(n-1,1)}$. Using the notation that $[xⁿ]f(x)$ is the coefficient of $xⁿ$ in $f(x)$, one has by page 253 of [\[Mac\]](#page-51-16) that

$$
Q_{(n-1,1)} = [t_1^{n-1}t_2] \left(\frac{1 - t_2/t_1}{1 + t_2/t_1} \right) \prod_{i=1}^2 \prod_{j=1}^\infty \frac{1 + t_i x_j}{1 - t_i x_j}
$$

\n
$$
= [t_1^{n-1}] \prod_{j=1}^\infty \frac{1 + t_1 x_j}{1 - t_1 x_j} [t_2] \prod_{j=1}^\infty \frac{1 + t_2 x_j}{1 - t_2 x_j} - 2[t_1^n] \prod_{j=1}^\infty \frac{1 + t_1 x_j}{1 - t_1 x_j}
$$

\n
$$
= 2p_1 Q_{n-1} - 2Q_n.
$$

By page 248 of [\[Mac\]](#page-51-16), $X_{\mu}^{n} = 1$ for all $\mu \in OP(n)$. It follows from the first paragraph that for $\mu \in OP(n)$, the coefficient of p_{μ} in Q_n is $\frac{2^{l(\mu)}}{z_{\mu}}$ $rac{\mu(\mu)}{z_{\mu}}$ and (by considering separately the cases that $m_1(\mu) = 0$ and $m_1(\mu) \neq 0$) that the coefficient of p_μ in $p_1 Q_{n-1}$ is $\frac{m_1(\mu)2^{l(\mu)-1}}{z_\mu}$. Thus the coefficient of p_μ in $Q_{(n-1,1)}$ is $\frac{(m_1(\mu)-2)2^{l(\mu)}}{z_{\mu}}$, which by the first paragraph proves the result. □

5.2. Markov chains on strict partitions. This subsection discusses a Markov chain on $DP(n)$ to be used in defining an exchangeable pair for the proof of Theorem [5.2.](#page-27-0)

Motivated by constructions in the group and Gelfand pair cases, it would be natural, for τ a fixed element of $DP(n)$, to define a "Markov chain" J_{τ} on $DP(n)$ with transition "probabilities"

$$
J_{\tau}(\lambda,\rho):=\frac{g_{\rho}}{2^{l(\rho)}g_{\lambda}g_{\tau}}\sum_{\nu\in OP(n)}\frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\rho}X_{\nu}^{\tau}}{z_{\nu}}.
$$

Using Lemma [5.4,](#page-29-1) one can see that $\sum_{\rho} J_{\tau}(\lambda, \rho) = 1$ for all λ , and that J_{τ} satisfies the reversibility condition

$$
\frac{2^{n-l(\lambda)}g_{\lambda}^2}{n!}J_{\tau}(\lambda,\rho)=\frac{2^{n-l(\rho)}g_{\rho}^2}{n!}J_{\tau}(\rho,\lambda)
$$

for all $\lambda, \rho \in DP(n)$. Unfortunately, the quantity $J_{\tau}(\lambda, \rho)$ can be negative. For instance one can check from Lemma [5.6](#page-30-0) that $J_{(2,1)}((2,1),(2,1)) = -1$.

To deal with the complication raised in the previous paragraph, it is helpful to introduce a genuine Markov chain L on $DP(n)$, with transition probabilities

$$
L(\lambda, \rho) = \frac{2g_{\rho}}{ng_{\lambda}} \sum_{\substack{\eta \in DP(n-1) \\ \eta \nearrow \lambda, \eta \nearrow \rho}} 2^{l(\eta) - l(\rho)}.
$$

Here $\eta \nearrow \lambda$ means that η is obtained from λ by decreasing the size of some part by exactly one.

Lemma [5.7](#page-31-1) proves that L is a Markov chain which is reversible with respect to shifted Plancherel measure. As is mentioned in the proof, the definition of L was motivated by the theory of harmonic functions on Bratteli diagrams.

Lemma 5.7. The transition probabilities of L are real and non-negative and sum to 1. Moreover the chain L is reversible with respect to shifted Plancherel measure.

Proof. This is a special case of a construction in Section 2 of [\[F1\]](#page-50-3). To see this, one takes the underlying Bratteli diagram to be the Schur graph, the properties of which are discussed in Section 5 of [\[BO2\]](#page-50-22). The vertices of the Schur graph are all partitions of all non-negative integers with distinct parts, and the edge multiplicity between $\eta \in DP(n-1)$ and $\lambda \in DP(n)$ is 1 if $\eta \nearrow \lambda$, and is 0 otherwise. The combinatorial dimension of a shape λ is g_{λ} , and if π denotes shifted Plancherel measure, the function $\frac{\pi(\lambda)}{g_{\lambda}}$ is harmonic on the Schur graph.

Proposition [5.9](#page-32-0) will establish a fundamental relation between the Markov chain L and the "Markov chain" $J_{(n-1,1)}$. The argument involves symmetric function theory, and first a lemma is needed about properties of a certain subring Γ of the ring of symmetric functions, defined on page 252 of [\[Mac\]](#page-51-16).

Recall from page 255 of $[Mac]$ that there is an inner product on Γ which satisfies the properties

(1)
$$
\langle p_{\lambda}, p_{\mu} \rangle = 2^{-l(\lambda)} z_{\lambda} \delta_{\lambda, \mu}
$$
 if $\lambda, \mu \in OP(n)$.

(2)
$$
\langle P_{\lambda}, P_{\mu} \rangle = 2^{-l(\lambda)} \delta_{\lambda, \mu}
$$
 if $\lambda, \mu \in DP(n)$.

Here p_{λ} is a power sum symmetric function and P_{λ} is a Hall-Littlewood polynomial with the parameter $t = -1$. It is also helpful to recall, from page 247 of [\[Mac\]](#page-51-16), that for $\lambda \in DP(n)$,

$$
P_{\lambda} = 2^{-l(\lambda)} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}}{z_{\nu}} X_{\nu}^{\lambda} p_{\nu}.
$$

Lemma 5.8. Let p_1^{\perp} be the adjoint in Γ of multiplication by p_1 .

(1) ([\[Mac\]](#page-51-16), page 265) $p_1^{\perp} = \frac{1}{2}$ 2 ∂ $\frac{\partial}{\partial p_1}$. (2) $p_1^{\perp} P_{\lambda} = \sum_{\substack{\eta \in DP(n-1) \\ \eta \nearrow \lambda}} 2^{l(\eta)-l(\lambda)} P_{\eta}.$

Proof. Only the second assertion needs to be proved. Consider the coefficient of P_{η} in $p_1^{\perp}P_{\lambda}$. It is

$$
\frac{\langle p_1^{\perp} P_{\lambda}, P_{\eta} \rangle}{\langle P_{\eta}, P_{\eta} \rangle} = \frac{\langle P_{\lambda}, p_1 P_{\eta} \rangle}{\langle P_{\eta}, P_{\eta} \rangle} = 2^{l(\eta)} \langle P_{\lambda}, p_1 P_{\eta} \rangle.
$$

From Section 3.8 of [\[Mac\]](#page-51-16),

$$
p_1 P_\eta = \sum_{\substack{\lambda \in DP(n) \\ \eta \nearrow \lambda}} P_\lambda,
$$

which implies the result. \Box

Proposition [5.9](#page-32-0) can now be proved.

Proposition 5.9. Suppose that $n \geq 3$, so that $(n-1,1)$ is a strict partition. Then

$$
L(\lambda, \rho) = \frac{n-2}{n} J_{(n-1,1)}(\lambda, \rho)
$$

for $\lambda, \rho \in DP(n)$ such that $\lambda \neq \rho$.

Proof. By Lemma [5.6,](#page-30-0)

$$
J_{(n-1,1)}(\lambda,\rho) = \frac{g_{\rho}}{2^{l(\rho)}g_{\lambda}(n-2)} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\rho}(m_1(\nu) - 2)}{z_{\nu}}.
$$

Since $\lambda \neq \rho$, Lemma [5.3](#page-28-1) and part 1 of Lemma [5.8](#page-32-1) imply that this is equal to

$$
\frac{g_{\rho}}{2^{l(\rho)}g_{\lambda}(n-2)} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\rho}m_{1}(\nu)}{z_{\nu}}
$$
\n
$$
= \frac{g_{\rho}}{2^{l(\rho)}g_{\lambda}(n-2)} \left\langle \sum_{\nu \in OP(n)} m_{1}(\nu) \frac{2^{l(\nu)}}{z_{\nu}} X_{\nu}^{\lambda} p_{\nu}, \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}}{z_{\nu}} X_{\nu}^{\rho} p_{\nu} \right\rangle
$$
\n
$$
= \frac{2^{l(\lambda)}g_{\rho}}{g_{\lambda}(n-2)} \langle 2p_{1}p_{1}^{\perp}P_{\lambda}, P_{\rho} \rangle
$$
\n
$$
= \frac{2^{l(\lambda)+1}g_{\rho}}{g_{\lambda}(n-2)} \langle p_{1}^{\perp}P_{\lambda}, p_{1}^{\perp}P_{\rho} \rangle.
$$

By part 2 of Lemma [5.8,](#page-32-1) this is

$$
\frac{2^{l(\lambda)+1}g_{\rho}}{g_{\lambda}(n-2)} \left\langle \sum_{\substack{\eta \in DP(n-1) \\ \eta \nearrow \lambda}} 2^{l(\eta)-l(\lambda)} P_{\eta}, \sum_{\substack{\eta \in DP(n-1) \\ \eta \nearrow \rho}} 2^{l(\eta)-l(\rho)} P_{\eta} \right\rangle
$$

=
$$
\frac{2g_{\rho}}{g_{\lambda}(n-2)} \sum_{\substack{\eta \in DP(n-1) \\ \eta \nearrow \lambda, \eta \nearrow \rho}} 2^{l(\eta)-l(\rho)}
$$

=
$$
\frac{n}{n-2}L(\lambda, \rho).
$$

 \Box

5.3. Central limit theorem for shifted Plancherel measure. This subsection studies the statistic $W = \sqrt{\frac{n!}{n \cdot n!}}$ $z_\mu 2^{n-l(\mu)}$ $\frac{X_{\mu}^{\lambda}}{g_{\lambda}},$ where $\mu \in OP(n)$ is fixed and $\lambda \in DP(n)$ is chosen from shifted Plancherel measure. The main goal is a proof of Theorem [5.2.](#page-27-0)

Using L , one constructs an exchangeable pair (W, W') as follows. Choose λ from the shifted Plancherel measure on partitions of size *n*. Then choose ρ with probability $L(\lambda, \rho)$ and let $(W, W') = (W(\lambda), W(\rho))$. Using J_{τ} , one constructs an "exchangeable pair" (W, W^*) as follows. Choose λ from the shifted Plancherel measure on partitions of size n. Then choose ρ with "probability" $J_{\tau}(\lambda, \rho)$ and let $(W, W^*) = (W(\lambda), W(\rho))$. The pair (W, W') is a valid candidate for Stein's method, but the pair (W, W^*) is much easier to work with, and by Proposition [5.9,](#page-32-0) when $\tau = (n-1,1)$, this gives insight into the genuine exchangeable pair (W, W') . Even though the transition probabilities of J_{τ} can be negative, for convenience the usual language of probability theory (expected value, variance, etc.) will be used when working with them.

Lemma 5.10. (1)
$$
\mathbb{E}(W^*|W) = \left(\frac{X_\mu^\tau}{g_\tau}\right)W.
$$

(2)
$$
\mathbb{E}(W'|W) = \frac{m_1(\mu)}{n}W.
$$

Proof. For the first assertion, the definition of $J_{(n-1,1)}$ and Lemma [5.4](#page-29-1) imply that

$$
\mathbb{E}(W^*|\lambda) = \sqrt{\frac{n!}{z_{\mu}2^{n-l(\mu)}}} \sum_{\rho \in DP(n)} \frac{g_{\rho}}{2^{l(\rho)}g_{\lambda}g_{\tau}} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\rho}X_{\nu}^{\tau}}{z_{\nu}} \frac{X_{\mu}^{\rho}}{g_{\rho}}
$$

\n
$$
= \sqrt{\frac{n!}{z_{\mu}2^{n-l(\mu)}}} \frac{1}{g_{\lambda}g_{\tau}} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\tau}}{z_{\nu}} \sum_{\rho \in DP(n)} \frac{X_{\nu}^{\rho}X_{\mu}^{\rho}}{2^{l(\rho)}}
$$

\n
$$
= \sqrt{\frac{n!}{z_{\mu}2^{n-l(\mu)}}} \frac{X_{\mu}^{\lambda}}{g_{\lambda}} \frac{X_{\mu}^{\tau}}{g_{\tau}}
$$

\n
$$
= \left(\frac{X_{\mu}^{\tau}}{g_{\tau}}\right)W(\lambda).
$$

The first assertion follows since this depends on λ only through W. For the second assertion, observe that by Proposition [5.9,](#page-32-0)

$$
\mathbb{E}(W'-W|\lambda) = \sum_{\rho} L(\lambda, \rho) (W(\rho) - W(\lambda))
$$

=
$$
\frac{n-2}{n} \sum_{\rho} J_{(n-1,1)}(\lambda, \rho) (W(\rho) - W(\lambda))
$$

=
$$
- \left(\frac{n-2}{n}\right) W(\lambda) + \frac{n-2}{n} \sum_{\rho} J_{(n-1,1)}(\lambda, \rho) W(\rho).
$$

By the first assertion and Lemma [5.6,](#page-30-0) this is

$$
-\left(\frac{n-2}{n}\right)W(\lambda) + \frac{n-2}{n}\frac{m_1(\mu) - 2}{n-2}W(\lambda) = \left(\frac{m_1(\mu)}{n} - 1\right)W(\lambda),
$$

which implies the result.

Corollary [5.11](#page-34-0) will not be needed but is worth recording.

Corollary 5.11. The eigenvalues of J_{τ} are $\frac{X_{\mu}^{\tau}}{g_{\tau}}$ as μ ranges over $OP(n)$. The functions $\psi_{\mu}(\lambda) = \sqrt{\frac{n!}{z_{\mu}2^{n-l(\mu)}}}$ $\frac{X^{\lambda}_{\mu}}{g_{\lambda}}$ are a basis of eigenvectors of J_{τ} , orthonormal with respect to the inner product

$$
\langle f_1, f_2 \rangle = \sum_{\lambda \in DP(n)} f_1(\lambda) \overline{f_2(\lambda)} \frac{2^{n-l(\lambda)} g_\lambda^2}{n!}.
$$

Proof. The proof of part 1 of Lemma [5.10](#page-33-1) shows that ψ_{μ} is an eigenvector of J_{τ} with eigenvalue $\frac{X_{\mu}^{\tau}}{g_{\tau}}$. The orthonormality assertion follows from Lemma [5.4](#page-29-1) and the fact from [\[Mac\]](#page-51-16) that all X^{λ}_{μ} are real valued. The basis assertion follows since $|DP(n)| = |OP(n)|$.

Lemma 5.12. $\mathbb{E}(W^{*}-W)^{2}=2\left(1-\frac{X_{\mu}^{\tau}}{g_{\tau}}\right)$. Proof. This follows from the proof of Lemma [2.3](#page-5-2) (which does not require non-negative transition probabilities) and part 1 of Lemma [5.10.](#page-33-1) \Box

For the remainder of this subsection, $p_m(\tilde{K}_{\nu})$ will denote the coefficient of \tilde{K}_{ν} in $(\tilde{K}_{\mu})^m$. When $m=0$ this is to be interpreted through Lemma [5.5,](#page-29-0) so that $p_0(\tilde{K}_{(1^n)}) = 1$ and $p_0(\tilde{K}_{\nu}) = 0$ for $\nu \neq (1^n)$. Due to the signs in the definition of \tilde{K}_{ν} , these numbers are not probabilities so care must be taken in working with them. For instance it is not true that $\sum_{\nu \in OP(n)} p_m(\tilde{K}_{\nu}) = 1$. However the following three relations will be useful.

Lemma 5.13. $\sum_{\nu \in OP(n)} p_2(\tilde{K}_{\nu}) 2^{l(\nu)-n} = 2^{2[l(\mu)-n]}.$

Proof. Using Lemma [5.5,](#page-29-0) one has that,

$$
\sum_{\nu \in OP(n)} p_2(\tilde{K}_{\nu}) 2^{l(\nu)} = 2^{2l(\mu)-n} \sum_{\rho \in DP(n)} 2^{-l(\rho)} \frac{(X_{\mu}^{\rho})^2}{g_{\rho}} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}}{z_{\nu}} X_{\nu}^{\rho}.
$$

By page 248 of [\[Mac\]](#page-51-16), $X_{\nu}^{n} = 1$, so that Lemma [5.3](#page-28-1) implies that

$$
\sum_{\nu \in OP(n)} \frac{2^{l(\nu)}}{z_{\nu}} X_{\nu}^{\rho} X_{\nu}^n = 2\delta_{\rho,n}.
$$

The result follows.

Lemma 5.14. $\mathbb{E}\left(\frac{X^{\lambda}_{\mu}}{g_{\lambda}}\right)$ $\bigg)^4 = (2^{n-l(\mu)})^4 \sum_{\nu \in OP(n)} p_2(\tilde{K}_{\nu})^2 \frac{2^{l(\nu)} z_{\nu}}{2^n n!}.$

Proof. Consider the coefficient of $\tilde{K}_{(1^n)}$ in $(\tilde{K}_{\mu})^4$. On one hand, by Lemma [5.5](#page-29-0) it is equal to $(2^{l(\mu)-n})^4 \mathbb{E} \left(\frac{X_{\mu}^{\lambda}}{g_{\lambda}} \right)$ \int_{1}^{4} . On the other hand,

$$
(\tilde{K}_{\mu})^4 = (\tilde{K}_{\mu})^2 (\tilde{K}_{\mu})^2 = \left[\sum_{\nu \in OP(n)} p_2(\tilde{K}_{\nu}) \tilde{K}_{\nu} \right]^2.
$$

From Lemmas [5.5](#page-29-0) and Lemma [5.4,](#page-29-1) it follows that the coefficient of $\tilde{K}_{(1^n)}$ in $\tilde{K}_{\nu_1}\tilde{K}_{\nu_2}$ is 0 if $\nu_1 \neq \nu_2$, and that the coefficient of $\tilde{K}_{(1^n)}$ in $(\tilde{K}_{\nu})^2$ is $\frac{2^{l(\nu)-n}z_{\nu}}{n!}$. Thus the coefficient of $\tilde{K}_{(1^n)}$ in $(\tilde{K}_{\mu})^4$ is equal to

$$
\sum_{\nu \in OP(n)} p_2(\tilde{K}_{\nu})^2 \frac{2^{l(\nu)-n}z_{\nu}}{n!}.
$$

Comparing the two expressions for the coefficient of $\tilde{K}_{(1^n)}$ in $(\tilde{K}_{\mu})^4$ proves the result. \Box

Lemma 5.15. $p_3(\tilde{K}_\mu)=\frac{2^{n-l(\mu)}}{z_\mu}$ $\frac{-l(\mu)}{z_{\mu}}\sum_{\nu\in OP(n)} p_2(\tilde{K}_{\nu})^2 2^{l(\nu)-n}z_{\nu}.$

Proof. By Lemma [5.5,](#page-29-0)

$$
p_3(\tilde{K}_{\mu}) = \frac{n!(2^{l(\mu)-n})^3}{z_{\mu}} \mathbb{E}\left(\frac{X_{\mu}^{\lambda}}{g_{\lambda}}\right)^4.
$$

Now use Lemma [5.14.](#page-35-0) \Box

The next lemmas are crucial.

Lemma 5.16. $\mathbb{E}((W^*)^2|\lambda) = \frac{n!}{z_\mu 2^{l(\mu)}} \sum_{\nu \in OP(n)}$ $\frac{X^{\lambda}_{\nu}}{g_{\lambda}}$ $\frac{X_{\nu}^{\tau}}{g_{\tau}} 2^{l(\nu)} p_2(\tilde{K}_{\nu}).$

Proof. It follows from the definitions that $\mathbb{E}((W^*)^2|\lambda)$ is equal to

$$
\frac{n!}{z_{\mu}2^{n-l(\mu)}} \sum_{\rho \in DP(n)} \frac{g_{\rho}}{2^{l(\rho)}g_{\lambda}g_{\tau}} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\rho}X_{\nu}^{\tau}}{z_{\nu}} \left(\frac{X_{\mu}^{\rho}}{g_{\rho}}\right)^2
$$

$$
= \frac{n!}{z_{\mu}2^{n-l(\mu)}} \sum_{\nu \in OP(n)} \frac{X_{\nu}^{\lambda}}{g_{\lambda}} \frac{X_{\tau}^{\lambda}}{g_{\tau}} 2^{l(\nu)} \left(\frac{1}{z_{\nu}} \sum_{\rho \in DP(n)} \frac{g_{\rho}^2}{2^{l(\rho)}} \frac{X_{\nu}^{\rho}}{g_{\rho}} \left(\frac{X_{\mu}^{\rho}}{g_{\rho}}\right)^2\right).
$$

The result now follows from Lemma [5.5.](#page-29-0) $\hfill \square$

Lemma 5.17.

$$
Var(\mathbb{E}[(W^* - W)^2 | \lambda]) = \frac{n!2^n}{z_{\mu}^2 2^{2l(\mu)}} \sum_{\substack{\nu \in OP(n) \\ \nu \neq (1^n)}} 2^{l(\nu)} p_2(\tilde{K}_{\nu})^2 z_{\nu} \left(\frac{X_{\nu}^{\tau}}{g_{\tau}} + 1 - \frac{2X_{\mu}^{\tau}}{g_{\tau}}\right)^2
$$

Proof. By Lemmas [5.16,](#page-36-0) [5.12,](#page-34-1) and [5.10,](#page-33-1) $Var(\mathbb{E}[(W^* - W)^2 | \lambda])$ is equal to

$$
\mathbb{E}(\mathbb{E}((W^* - W)^2 | \lambda)^2) - 4\left(1 - \frac{X_{\mu}^{\tau}}{g_{\tau}}\right)^2
$$
\n
$$
= \mathbb{E}\left[\frac{n!}{z_{\mu}2^{l(\mu)}} \sum_{\nu \in OP(n)} \frac{X_{\nu}^{\lambda}}{g_{\lambda}} \frac{X_{\nu}^{\tau}}{g_{\tau}} 2^{l(\nu)} p_2(\tilde{K}_{\nu}) + \left(1 - \frac{2X_{\mu}^{\tau}}{g_{\tau}}\right) W^2\right]^2
$$
\n
$$
-4\left(1 - \frac{X_{\mu}^{\tau}}{g_{\tau}}\right)^2
$$
\n
$$
= T_1 + T_2 + T_3 - 4\left(1 - \frac{X_{\mu}^{\tau}}{g_{\tau}}\right)^2
$$

where

$$
T_1 = \left(\frac{n!}{z_\mu 2^{l(\mu)}}\right)^2 \mathbb{E}\left(\sum_{\nu \in OP(n)} p_2(\tilde{K}_\nu) \frac{X_\nu^{\lambda}}{g_{\lambda}} \frac{X_\nu^{\tau}}{g_{\tau}} 2^{l(\nu)}\right)^2
$$

$$
T_2 = 2\left(1 - \frac{2X_\mu^{\tau}}{g_{\tau}}\right) \frac{(n!)^2}{z_\mu^2} \sum_{\nu \in OP(n)} \frac{X_\nu^{\tau} p_2(\tilde{K}_\nu)}{g_{\tau} 2^{n-l(\nu)}} \mathbb{E}\left(\frac{X_\nu^{\lambda}(X_\mu^{\lambda})^2}{(g_{\lambda})^3}\right)
$$

$$
T_3 = \left(1 - \frac{2X_\mu^{\tau}}{g_{\tau}}\right)^2 \left(\frac{n!}{2^{n-l(\mu)}z_\mu}\right)^2 \mathbb{E}\left(\frac{X_\mu^{\lambda}}{g_{\lambda}}\right)^4.
$$

These expressions for T_1, T_2, T_3 can be simplified. Lemma [5.4](#page-29-1) implies that

$$
T_1 = \frac{n!2^n}{(z_\mu)^2 2^{2l(\mu)}} \sum_{\nu \in OP(n)} p_2(\tilde{K}_\nu)^2 2^{l(\nu)} z_\nu \left(\frac{X_\nu^{\tau}}{g_{\tau}}\right)^2.
$$

From Lemma [5.5](#page-29-0) it follows that

$$
T_2 = 2\left(1 - \frac{2X_\mu^{\tau}}{g_\tau}\right) \frac{n!2^n}{(z_\mu)^2 2^{2l(\mu)}} \sum_{\nu \in OP(n)} p_2(\tilde{K}_\nu)^2 2^{l(\nu)} z_\nu \frac{X_\nu^{\tau}}{g_\tau}.
$$

By Corollary [5.14,](#page-35-0)

$$
T_3 = \frac{n! 2^n}{2^{2l(\mu)}(z_{\mu})^2} \left(1 - \frac{2X_{\mu}^{\tau}}{g_{\tau}}\right)^2 \sum_{\nu \in OP(n)} p_2(\tilde{K}_{\nu})^2 2^{l(\nu)} z_{\nu}.
$$

Hence, one can write

$$
T_1 + T_2 + T_3 = \frac{n!2^n}{(z_\mu)^2 2^{2l(\mu)}} \sum_{\nu \in OP(n)} 2^{l(\nu)} z_\nu p_2(\tilde{K}_\nu)^2 \left(\frac{X_\nu^\tau}{g_\tau} + 1 - \frac{2X_\mu^\tau}{g_\tau}\right)^2
$$

and the result follows since by Lemmas [5.5](#page-29-0) and [5.4,](#page-29-1) the $\nu = (1^n)$ term in the above sum is $4\left(1-\frac{X_\mu^{\tau}}{g_{\tau}}\right)$ $\big)^2$.

Lemma [5.18](#page-37-0) will also be helpful.

Lemma 5.18. Let k be a positive integer.

(1) $\mathbb{E}(W^* - W)^k$ is equal to

$$
\left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)^{k/2} \sum_{m=0}^{k} (-1)^{k-m} {k \choose m} \sum_{\nu \in OP(n)} \frac{X_{\nu}^{\tau} z_{\nu}}{g_{\tau} n!2^{n-l(\nu)}} p_m(\tilde{K}_{\nu}) p_{k-m}(\tilde{K}_{\nu}).
$$

(2)
$$
\mathbb{E}(W^* - W)^4 \text{ is equal to}
$$

$$
\left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)^2 \left[\sum_{\nu \in OP(n)} \left(8\left(1 - \frac{X_{\mu}^{\tau}}{g_{\tau}}\right) - 6\left(1 - \frac{X_{\nu}^{\tau}}{g_{\tau}}\right)\right) \frac{z_{\nu}p_2(\tilde{K}_{\nu})^2}{n!2^{n-l(\nu)}}\right].
$$

Proof. To prove the first assertion, observe that

$$
\mathbb{E}((W^* - W)^k | \lambda)
$$
\n
$$
= \left(\frac{n!}{z_{\mu}2^{n-l(\mu)}}\right)^{k/2} \sum_{\rho \in DP(n)} \frac{g_{\rho}}{2^{l(\rho)}g_{\lambda}g_{\tau}} \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\rho}X_{\nu}^{\tau}}{z_{\nu}}
$$
\n
$$
\cdot \sum_{m=0}^{k} (-1)^{k-m} {k \choose m} \left(\frac{X_{\mu}^{\rho}}{g_{\rho}}\right)^{m} \left(\frac{X_{\mu}^{\lambda}}{g_{\lambda}}\right)^{k-m}
$$
\n
$$
= \left(\frac{n!}{z_{\mu}2^{n-l(\mu)}}\right)^{k/2} \frac{1}{g_{\lambda}g_{\tau}} \sum_{m=0}^{k} (-1)^{k-m} {k \choose m} \left(\frac{X_{\mu}^{\lambda}}{g_{\lambda}}\right)^{k-m}
$$
\n
$$
\cdot \sum_{\nu \in OP(n)} \frac{2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\tau}}{z_{\nu}} \sum_{\rho \in DP(n)} \frac{g_{\rho}^2}{2^{l(\rho)}} \frac{X_{\nu}^{\rho}}{g_{\rho}} \left(\frac{X_{\mu}^{\rho}}{g_{\rho}}\right)^{m}
$$
\n
$$
= \left(\frac{n!}{z_{\mu}2^{n-l(\mu)}}\right)^{k/2} \frac{1}{2^{n}g_{\lambda}g_{\tau}} \sum_{m=0}^{k} (-1)^{k-m} {k \choose m} (2^{n-l(\mu)})^{m} \left(\frac{X_{\mu}^{\lambda}}{g_{\lambda}}\right)^{k-m}
$$
\n
$$
\cdot \sum_{\nu \in OP(n)} 2^{l(\nu)}X_{\nu}^{\lambda}X_{\nu}^{\tau}p_{m}(\tilde{K}_{\nu}),
$$

where the last equation is Lemma [5.5.](#page-29-0) Hence $\mathbb{E}(W^* - W)^k$ is equal to

$$
\mathbb{E}(\mathbb{E}((W^* - W)^k|\lambda))
$$
\n
$$
= \left(\frac{n!}{z_{\mu}2^{n-l(\mu)}}\right)^{k/2} \frac{(2^{n-l(\mu)})^k}{2^n} \sum_{m=0}^k (-1)^{k-m} {k \choose m}
$$
\n
$$
\sum_{\nu \in OP(n)} 2^{l(\nu)} \frac{X_{\nu}^{\tau}}{g_{\tau}} \frac{p_m(\tilde{K}_{\nu})}{(2^{n-l(\mu)})^{k-m}} \mathbb{E}\left[\frac{X_{\nu}^{\lambda}}{g_{\lambda}} \left(\frac{X_{\mu}^{\lambda}}{g_{\lambda}}\right)^{k-m}\right].
$$

The first assertion now follows from Lemma [5.5.](#page-29-0)

For the second assertion, the first assertion gives that $\mathbb{E}(W^* - W)^4$ is equal to

$$
\left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_{\nu \in OP(n)} \frac{z_{\nu}}{2^{n-l(\nu)} n!} \frac{X_{\nu}^{\tau}}{g_{\tau}} p_m(\tilde{K}_{\nu}) p_{4-m}(\tilde{K}_{\nu}).
$$

The special case $\tau = (n)$ gives that

$$
0 = \left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_{\nu \in OP(n)} \frac{z_{\nu}}{2^{n-l(\nu)} n!} p_m(\tilde{K}_{\nu}) p_{4-m}(\tilde{K}_{\nu}).
$$

Thus in general, $\mathbb{E}(W^* - W)^4$ is equal to

$$
-\left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)^2\sum_{m=0}^4(-1)^m\binom{4}{m}\sum_{\nu\in OP(n)}\left(1-\frac{X^{\tau}_{\nu}}{g_{\tau}}\right)\frac{z_{\nu}p_m(\tilde{K}_{\nu})p_{4-m}(\tilde{K}_{\nu})}{2^{n-l(\nu)}n!}.
$$

The $m = 0, 4$ terms vanish since only $\nu = (1^n)$ could contribute, but it contributes 0. The $m = 2$ term contributes

$$
-6\left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)^2 \sum_{\nu \in OP(n)} \left(1 - \frac{X_{\nu}^{\tau}}{g_{\tau}}\right) \frac{z_{\nu}p_2(\tilde{K}_{\nu})^2}{n!2^{n-l(\nu)}}.
$$

The $m = 1, 3$ terms are equal and together contribute

$$
8\left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)\left(1-\frac{X_{\mu}^{\tau}}{g_{\tau}}\right)p_3(\tilde{K}_{\mu}).
$$

By Lemma [5.15,](#page-35-1) this is

$$
8\left(\frac{n!2^{n-l(\mu)}}{z_{\mu}}\right)^2 \left(1-\frac{X_{\mu}^{\tau}}{g_{\tau}}\right) \sum_{\nu \in OP(n)} \frac{z_{\nu}p_2(\tilde{K}_{\nu})^2}{n!2^{n-l(\nu)}}.
$$

Adding the terms together proves the second assertion.

The final combinatorial ingredient for the proof of Theorem [5.2](#page-27-0) is an upper bound for $p_2(\tilde{K}_{\nu})$. Lemma [5.19](#page-39-0) reduces this to a result obtained earlier in the section on Gelfand pairs.

Lemma 5.19. Let $p_2(\tilde{K}_{\nu})$ be as in this subsection and let $p_2(K_{\nu})$ be as in Subsection [4.4.](#page-24-0) Then $p_2(\tilde{K}_\nu) \leq p_2(K_\nu)$ for all $\nu \in OP(n)$.

Proof. Clearly $p_2(K_\nu)$ is equal to $|B_n\omega_\nu B_n|$ multiplied by the coefficient of ω_{ν} in

$$
\left(\frac{1}{|B_n|^2}\sum_{x_1,x_2\in B_n}x_1\omega_\mu x_2\right)^2.
$$

Next consider $p_2(\tilde{K}_{\nu})$. By definition it is the coefficient of \tilde{K}_{ν} in $(\tilde{K}_{\mu})^2$. Corollary 3.2 of [\[Stm\]](#page-51-5) gives that the coefficient of ω_{ν} in \tilde{K}_{ν} is $\frac{1}{|B_{n}\omega_{\nu}B_{n}|}$. Hence $p_2(\tilde{K}_{\nu})$ is $|B_n\omega_{\nu}B_n|$ multiplied by the coefficient of ω_{ν} in

$$
\left(\frac{1}{|B_n|^2}\sum_{x_1,x_2\in B_n}\phi(x_1x_2)x_1\omega_\mu x_2\right)^2,
$$

where ϕ is the linear character of B_n described in Subsection [4.1.](#page-15-0) Since $\phi(x_1x_2)$ is always ± 1 , the result follows.

Finally, we prove Theorem [5.2.](#page-27-0)

Proof. (Of Theorem [5.2\)](#page-27-0) One can assume that $n \geq 2(2i+1)$. In the O notation throughout the proof, i is fixed and n is growing.

One applies Theorem [2.1](#page-4-1) to the pair (W, W') , where $\mu = (2i+1, 1^{n-2i-1})$. Then $a = 1 - \frac{m_1(\mu)}{n}$ $\frac{h(\mu)}{n}$ by part 2 of Lemma [5.10.](#page-33-1) Proposition [5.9](#page-32-0) implies that

$$
\mathbb{E}((W'-W)^2|\lambda) = \frac{n-2}{n}\mathbb{E}((W^*-W)^2|\lambda)
$$

for all λ . Thus

$$
Var(\mathbb{E}[(W'-W)^2|\lambda]) = \left(\frac{n-2}{n}\right)^2 Var(\mathbb{E}[(W^*-W)^2|\lambda]).
$$

Lemmas [5.17](#page-36-1) and [5.6](#page-30-0) give that the first error term in applying Theorem [2.1](#page-4-1) to the pair (W, W') is at most

$$
\frac{n-2}{n} \frac{n! 2^{n-l(\mu)}}{\left(1 - \frac{m_1(\mu)}{n}\right) z_{\mu}} \sqrt{\sum_{\substack{\nu \in OP(n) \\ \nu \neq (1^n)}} \frac{z_{\nu} p_2(\tilde{K}_{\nu})^2}{n! 2^{n-l(\nu)}} \left(\frac{n+m_1(\nu)-2m_1(\mu)}{n-2}\right)^2}.
$$

Observe that $\frac{n-2}{n}$ $\frac{n!2^{n-l(\mu)}}{\left(1-\frac{m_1(\mu)}{n}\right)z_\mu}$ is $O(n^{2i+2})$. In the sum over $\nu \neq (1^n)$, the term coming from $\nu = (2i+1, 2i+1, 1^{n-4i-2})$ contributes 0. Since $n \geq 2(2i+1)$ 1), Lemmas [5.19](#page-39-0) and [4.19](#page-25-2) imply that $\frac{z_{\nu}p_2(\tilde{K}_{\nu})^2}{n!2^{n-l(\nu)}} = O(n^{-4i-3})$ for all other ν in the sum, and the number of such ν is bounded by a constant depending on i but not n. Moreover if $p_2(\widetilde{K}_{\nu}) \neq 0$, then $p_2(K_{\nu}) \neq 0$, implying by Lemma [4.18](#page-25-0) that $m_1(\nu) \ge n - 4i - 2$ and thus that $\frac{n + m_1(\nu) - 2m_1(\mu)}{n-2} = O(n^{-1}).$ Combining these observations gives that the first term in the upper bound of Theorem [2.1](#page-4-1) is $O(n^{-1/2})$.

By the Cauchy-Schwarz inequality,

$$
\mathbb{E}|W'-W|^3 \leq \sqrt{\mathbb{E}(W'-W)^2\mathbb{E}(W'-W)^4}.
$$

Proposition [5.9](#page-32-0) implies that $\mathbb{E}(W'-W)^k = \frac{n-2}{n} \mathbb{E}(W^* - W)^k$ for all k. Hence Lemmas [5.12](#page-34-1) and [5.6](#page-30-0) imply that the second error term in Theorem [2.1](#page-4-1) is at most

$$
\left[\frac{(n-2)\mathbb{E}(W^* - W)^4}{\pi n \left(1 - \frac{m_1(\mu)}{n}\right)}\right]^{1/4}.
$$

By part 2 of Lemma [5.18](#page-37-0) and Lemma [5.6,](#page-30-0) this can be written as

$$
\left[\frac{1}{\pi}\frac{(n!2^{n-l(\mu)})^2}{(1-\frac{m_1(\mu)}{n})(z_{\mu})^2}\sum_{\nu\in OP(n)}\left(2-\frac{8m_1(\mu)}{n}+\frac{6m_1(\nu)}{n}\right)\frac{z_{\nu}p_2(\tilde{K}_{\nu})^2}{n!2^{n-l(\nu)}}\right]^{1/4}.
$$

The quantity $\frac{(n!2^{n-l(\mu)})^2}{(1-m_1(\mu))^{2}}$ $\frac{(n!2^{n-\iota(\mu)})^2}{(1-\frac{m_1(\mu)}{n})(z_{\mu})^2}$ is $O(n^{4i+3})$. The contribution from $\nu=(1^n)$ to the sum is $\frac{8(2i+1)}{n}p_2(\tilde{K}_{(1^n)})^2$, which by Lemmas [5.5](#page-29-0) and [5.4](#page-29-1) is $\frac{8(2i+1)^3}{2^{4i}}n^{-4i-3}$ + $O(n^{-4i-4})$. By Lemmas [5.19](#page-39-0) and [4.19,](#page-25-2) the contribution to the sum from ν not equal to either (1^n) or $(2i+1, 2i+1, 1^{n-4i-2})$ is $O(n^{-4i-4})$. The contribution from $\nu = (2i + 1, 2i + 1, 1^{n-4i-2})$ is $-\frac{4(2i+1)}{n}$ n $z_{\nu}p_2(\tilde{K}_{\nu})^2$ $rac{\nu_{\nu}p_2(\mathbf{\Lambda}_{\nu})}{n!2^{n-l(\nu)}}$. Lemma [5.13](#page-35-2) implies that $p_2(\tilde{K}_{(2i+1,2i+1,1^{n-4i-2})}) = 1 + O(n^{-1}),$ since $p_2(\tilde{K}_{\nu}) = O(n^{-1})$ for $\nu \neq (2i + 1, 2i + 1, 1^{n-4i-2})$ by Lemma [5.19.](#page-39-0) Thus the contribution from $\nu = (2i + 1, 2i + 1, 1^{n-4i-2})$ is $-\frac{8(2i+1)^3}{2^{4i}}$ $\frac{(2i+1)^3}{2^{4i}}n^{-4i-3} + O(n^{-4i-4})$. Hence there is useful cancellation and

$$
\sum_{\nu \in OP(n)} \left(2 - \frac{8m_1(\mu)}{n} + \frac{6m_1(\nu)}{n} \right) \frac{z_{\nu}p_2(\tilde{K}_{\nu})^2}{n!2^{n-l(\nu)}} = O(n^{-4i-4}).
$$

It follows that the second term in the upper bound of Theorem [2.1](#page-4-1) is $O(n^{-1/4})$, completing the proof.

6. Association Schemes

This final section adapts techniques from earlier sections to study the spectrum of an adjacency matrix of an association scheme. As is well known (see for instance Chapter 3 of [\[BaI\]](#page-49-2)) this includes the spectrum a distance regular graph as a special case.

Subsection [6.1](#page-41-1) discusses needed facts about association schemes. Subsection [6.2](#page-43-0) derives a general central limit theorem for the spectrum of an association scheme. Subsection [6.3](#page-48-0) illustrates the theory of Subsection [6.2](#page-43-0) for the special case of the Hamming scheme, where the result amounts to a central limit theorem for the spectrum of the Hamming graph, or equivalently for values of q-Krawtchouk polynomials. For an application of tools in this section to a problem with a non-normal limit, see [\[CF\]](#page-50-5).

6.1. Background on association schemes. This subsection gives preliminaries about association schemes, using notation from [\[MaSl\]](#page-51-18). Another useful reference is [\[BaI\]](#page-49-2), and what we call association schemes some authors call symmetric association schemes.

Definition. An association scheme with n classes consists of a finite set X with $n+1$ relations R_0, \dots, R_n defined on X which satisfy:

- (1) Each R_i is symmetric: $(x, y) \in R_i \Rightarrow (y, x) \in R_i$.
- (2) For every $x, y \in X$, one has that $(x, y) \in R_i$ for exactly one *i*.
- (3) $R_0 = \{(x, x) : x \in X\}$ is the identity relation.
- (4) If $(x, y) \in R_k$, the number of $z \in X$ such that $(x, z) \in R_i$ and $(y, z) \in R_j$ is a constant c_{ijk} depending on i, j, k but not on the particular choice of x and y .

The adjacency matrix D_i corresponding to the relation R_i is the $|X| \times |X|$ matrix with rows and columns labeled by the points of X defined by

$$
(D_i)_{x,y} = \begin{cases} 1 & \text{if } (x,y) \in R_i \\ 0 & \text{otherwise.} \end{cases}
$$

The Bose-Melner algebra is defined to be the vector space consisting of all matrices $\sum_{i=0}^{n} a_i D_i$ with a_i real. Since the matrices in the Bose-Melner algebra are symmetric and commute with each other (by parts 1 and 4 of the definition of an association scheme), the Bose-Melner algebra is semisimple and has a distinguished basis of primitive idempotents J_0, \dots, J_n satisfying

- (1) $J_i^2 = J_i$ $i = 0, \cdots, n$. (2) $J_iJ_k = 0$ $i \neq k$.
- (3) $\sum_{i=0}^{n} J_i = I$.

Here I is the identity matrix.

The D's are also a basis of the Bose-Melner algebra, so one can write

$$
D_s = \sum_{i=0}^n \phi_s(i) J_i, \quad s = 0, \cdots, n
$$

where the $\phi_s(i)$ are real numbers. Since $D_s J_i = \phi_s(i) J_i$, the $\phi_s(i)$ are the eigenvalues of D_s . For later use note that since D_0 and $\sum_{i=0}^{n} J_i$ are both the identity matrix, $\phi_0(j) = 1$ for all j.

Let μ_i be the rank of J_i . For all s, this is the multiplicity of the eigenvalue $\phi_s(i)$ of D_s . We define the Plancherel measure of the association scheme to be the probability measure on $\{0, \dots, n\}$ which chooses i with probability μ_i $\frac{\mu_i}{|X|}$.

Lemmas [6.1](#page-42-0) and [6.2](#page-42-1) are the orthogonality relations for eigenvalues of association schemes. To state them it is helpful to define $v_i = c_{ii0}$, so that for any x, one has that v_i is the number of pairs $(x, y) \in R_i$. Clearly $\sum_{i=0}^{n} v_i = |X|.$

Lemma 6.1. ([\[MaSl\]](#page-51-18), page 655) For $0 \leq k, l \leq n$,

$$
\sum_{r=0}^{n} \frac{\phi_r(k)\phi_r(l)}{v_r} = \frac{|X|}{\mu_k} \delta_{k,l}.
$$

Lemma 6.2. ([\[MaSl\]](#page-51-18), page 655) For $0 \leq k, l \leq n$,

$$
\sum_{i=0}^{n} \mu_i \phi_k(i) \phi_l(i) = |X| v_k \delta_{k,l}.
$$

Lemma [6.3](#page-42-2) will be crucial.

Lemma 6.3. The coefficient of D_l in $D_{s_1} \cdots D_{s_m}$ is equal to

$$
\frac{1}{v_l}\sum_{i=0}^n\frac{\mu_i}{|X|}\phi_{s_1}(i)\cdots\phi_{s_m}(i)\phi_l(i).
$$

In particular, the coefficient of D_l in $(D_s)^m$ is

$$
\frac{1}{v_l} \mathbb{E}[\phi_s(i)^m \phi_l(i)],
$$

where s is random from Plancherel measure of the association scheme. *Proof.* Since $D_s = \sum_{i=0}^n \phi_s(i) J_i$, one knows that

$$
D_{s_1}\cdots D_{s_m}=\sum_{i=0}^n\phi_{s_1}(i)\cdots\phi_{s_m}(i)J_i.
$$

By pages 654 and 655 of [\[MaSl\]](#page-51-18),

$$
J_i = \frac{1}{|X|} \sum_{l=0}^n \frac{\mu_i \phi_l(i)}{v_l} D_l.
$$

The result follows.

Let $p_m(r)$ denote $\frac{v_r}{(v_s)^m}$ multiplied by the coefficient of D_r in $(D_s)^m$. It follows from the definitions that $p_m(r)$ admits the following probabilistic interpretation. Start from some point $x_0 \in X$, move to a random $x_1 \in X$ such that $(x_0, x_1) \in R_s$, then to a random $x_2 \in X$ such that $(x_1, x_2) \in$ R_s , and so on until one obtains x_m . Then $p_m(r)$ is the probability that $(x_0, x_m) \in R_r$.

The following fact will be useful.

Lemma 6.4. (1)
$$
p_4(0) = \sum_{r=0}^n \frac{p_2(r)^2}{v_r}
$$
.
(2) $p_3(s) = v_s \sum_{r=0}^n \frac{p_2(r)^2}{v_r}$.

Proof. The first assertion is clear from the probabilistic interpretation of $p_m(r)$. For an analytic proof, note that by Lemma [6.3](#page-42-2) (in the first and fourth equalities) and Lemma [6.1](#page-42-0) (in the second and third equalities)

$$
\sum_{r=0}^{n} \frac{p_2(r)^2}{v_r} = \sum_{r=0}^{n} \frac{1}{v_r} \left(\frac{1}{(v_s)^2} \sum_{i=0}^{n} \frac{\mu_i}{|X|} \phi_s(i)^2 \phi_r(i) \right)^2
$$

$$
= \frac{1}{(v_s)^4} \sum_{r=0}^{n} \frac{1}{v_r} \sum_{i=0}^{n} \frac{(\mu_i)^2}{|X|^2} \phi_s(i)^4 \phi_r(i)^2
$$

$$
= \frac{1}{(v_s)^4} \sum_{i=0}^{n} \frac{\mu_i}{|X|} \phi_s(i)^4
$$

$$
= p_4(0).
$$

The second assertion follows from the first assertion since $p_3(s) = v_s p_4(0)$, as can be seen either by the probabilistic interpretation of $p_m(r)$ or by computing both sides of the equation using Lemma [6.3.](#page-42-2) \Box

6.2. Central limit theorem for the spectrum. Recall that the goal is to study Plancherel measure of association schemes, which chooses $i \in$ $\{0, \dots, n\}$ with probability $\frac{\mu_i}{|X|}$. More precisely, for s fixed, a central limit theorem is proved for the random variable W whose value at $i \in \{0, \dots, n\}$ is $\frac{\phi_s(i)}{\sqrt{v_s}}$.

Given $t \in \{0, \dots, n\}$, we define a Markov chain L_t on the set $\{0, \dots, n\}$ which moves from i to j with probability

$$
L_t(i,j) := \frac{\mu_j}{|X|} \sum_{r=0}^n \frac{\phi_r(i)\phi_r(t)\phi_r(j)}{(v_r)^2}.
$$

Lemma 6.5. The transition probabilities of L_t are real and non-negative and sum to 1. Moreover the chain L_t is reversible with respect to Plancherel measure.

Proof. By Theorems 3.6 and 3.8 of [\[BaI\]](#page-49-2), the $L_t(i, j)$ are non-negative real numbers. Next, observe that

$$
\sum_{j=0}^{n} L_t(i,j) = \frac{1}{|X|} \sum_{r=0}^{n} \frac{\phi_r(i)\phi_r(t)}{(v_r)^2} \sum_{j=0}^{n} \mu_j \phi_r(j).
$$

Since $\phi_0(j) = 1$ for all j, Lemma [6.2](#page-42-1) implies that

$$
\sum_{j=0}^{n} L_t(i,j) = v_0 \sum_{r=0}^{n} \frac{\phi_r(i)\phi_r(t)}{(v_r)^2} \delta_{r,0} = 1.
$$

For the reversibility assertion, it is clear from the definition of L_t that

$$
\frac{\mu_i}{|X|}L_t(i,j) = \frac{\mu_j}{|X|}L_t(j,i)
$$

for all i, j .

One uses the chain L_t to construct an exchangeable pair (W, W') in the usual way. First choose i from Plancherel measure, then choose j with probability $L_t(i, j)$, and finally let $(W, W') = (W(i), W(j))$.

Lemma 6.6. $\mathbb{E}(W'|W) = \left(\frac{\phi_s(t)}{v_s}\right)$ v_s $\big)W.$

Proof. By the definitions and Lemma [6.2,](#page-42-1) one has that

$$
\mathbb{E}(W'|i) = \frac{1}{\sqrt{v_s}} \sum_{j=0}^n L_t(i,j) \phi_s(j)
$$

=
$$
\frac{1}{\sqrt{v_s}} \frac{1}{|X|} \sum_{r=0}^n \frac{\phi_r(i) \phi_r(t)}{(v_r)^2} \sum_{j=0}^n \mu_j \phi_s(j) \phi_r(j)
$$

=
$$
\left(\frac{\phi_s(t)}{v_s}\right) W(i).
$$

The result follows since this depends on i only through W .

Corollary [6.7](#page-44-0) will not be used but is worth recording.

Corollary 6.7. The eigenvalues of L_t are $\frac{\phi_s(t)}{v_s}$ $\frac{s(b)}{v_s}$ for $0 \leq s \leq n$. The functions $\psi_s(i) = \frac{\phi_s(i)}{\sqrt{v_s}}$ are a basis of eigenvectors of L_t , orthonormal with respect to the inner product

$$
\langle f_1, f_2 \rangle = \sum_{i=0}^n f_1(i) \overline{f_2(i)} \frac{\mu_i}{|X|}.
$$

Proof. The proof of Lemma [6.6](#page-44-1) shows that ψ_s is an eigenvector of L_t with eigenvalue $\frac{\phi_s(t)}{v_s}$. The orthonormality assertion follows from Lemma [6.2,](#page-42-1) and the basis assertion follows since there are $n + 1$ eigenvectors.

Lemma 6.8. $\mathbb{E}(W'-W)^2=2\left(1-\frac{\phi_s(t)}{v_s}\right)$ $\overline{v_s}$.

Proof. This is immediate from Lemmas [2.3](#page-5-2) and [6.6.](#page-44-1)

Lemma 6.9. $\mathbb{E}((W')^{2}|i) = v_{s} \sum_{r=0}^{n}$ $\phi_r(i) \phi_r(t)$ $\frac{(v)\varphi_r(v)}{(v_r)^2}p_2(r)$.

Proof. It follows from the definitions that

$$
\mathbb{E}((W')^{2}|i) = \frac{1}{v_{s}} \sum_{j=0}^{n} \frac{\mu_{j}}{|X|} \sum_{r=0}^{n} \frac{\phi_{r}(i)\phi_{r}(t)\phi_{r}(j)}{(v_{r})^{2}} \phi_{s}(j)^{2}
$$

$$
= \frac{1}{v_{s}} \sum_{r=0}^{n} \frac{\phi_{r}(i)\phi_{r}(t)}{(v_{r})^{2}} \sum_{j=0}^{n} \frac{\mu_{j}}{|X|} \phi_{s}(j)^{2} \phi_{r}(j).
$$

The result now follows from Lemma [6.3.](#page-42-2)

Note that in the next lemma, the sum is over non-zero r .

Lemma 6.10.

$$
Var(\mathbb{E}[(W'-W)^2|i]) = (v_s)^2 \sum_{r=1}^n \frac{p_2(r)^2}{v_r} \left(\frac{\phi_r(t)}{v_r} + 1 - \frac{2\phi_s(t)}{v_s}\right)^2.
$$

Proof. By Lemmas [6.6,](#page-44-1) [6.8,](#page-44-2) and [6.9,](#page-45-0) $Var(\mathbb{E}[(W'-W)^2|i])$ is equal to

$$
\mathbb{E}(\mathbb{E}((W'-W)^2|i)^2) - 4\left(1 - \frac{\phi_s(t)}{v_s}\right)^2
$$
\n
$$
= \mathbb{E}\left[v_s \sum_{r=0}^n \frac{\phi_r(i)\phi_r(t)}{(v_r)^2} p_2(r) + \left(1 - \frac{2\phi_s(t)}{v_s}\right)W^2\right]^2
$$
\n
$$
-4\left(1 - \frac{\phi_s(t)}{v_s}\right)^2
$$
\n
$$
= T_1 + T_2 + T_3 - 4\left(1 - \frac{\phi_s(t)}{v_s}\right)^2,
$$

where

$$
T_1 = (v_s)^2 \mathbb{E} \left(\sum_{r=0}^n \frac{\phi_r(i)\phi_r(t)}{(v_r)^2} p_2(r) \right)^2
$$

$$
T_2 = 2 \left(1 - \frac{2\phi_s(t)}{v_s} \right) \sum_{r=0}^n \frac{\phi_r(t)}{(v_r)^2} p_2(r) \mathbb{E} \left(\phi_r(i)\phi_s(i)^2 \right)
$$

$$
T_3 = \left(1 - \frac{2\phi_s(t)}{v_s} \right)^2 \mathbb{E} \left(\frac{\phi_s(i)^4}{(v_s)^2} \right).
$$

Next we simplify these terms. By Lemma [6.2,](#page-42-1)

$$
T_1 = (v_s)^2 \sum_{r=0}^n \frac{\phi_r(t)^2}{(v_r)^3} p_2(r)^2.
$$

Lemma [6.3](#page-42-2) implies that

$$
T_2 = 2(v_s)^2 \left(1 - \frac{2\phi_s(t)}{v_s}\right) \sum_{r=0}^n \frac{\phi_r(t)}{(v_r)^2} p_2(r)^2.
$$

$$
\Box
$$

Lemma [6.3](#page-42-2) and part 1 of Lemma [6.4](#page-43-1) imply that

$$
T_3 = (v_s)^2 \left(1 - \frac{2\phi_s(t)}{v_s}\right)^2 p_4(0) = (v_s)^2 \left(1 - \frac{2\phi_s(t)}{v_s}\right)^2 \sum_{r=0}^n \frac{p_2(r)^2}{v_r}.
$$

Thus

$$
T_1 + T_2 + T_3 = (v_s)^2 \sum_{r=0}^{n} \frac{p_2(r)^2}{v_r} \left(\frac{\phi_r(t)}{v_r} + 1 - \frac{2\phi_s(t)}{v_s}\right)^2
$$

Since $p_2(0) = \frac{1}{v_s}$, the $r = 0$ term of $T_1 + T_2 + T_3$ is equal to $4\left(1 - \frac{\phi_s(t)}{v_s}\right)$ v_s $\Big)^2$, proving the result.

Lemma 6.11. Let k be a positive integer.

$$
(1) \mathbb{E}(W'-W)^k = (v_s)^{k/2} \sum_{m=0}^k (-1)^{k-m} {k \choose m} \sum_{r=0}^n \frac{\phi_r(t)}{v_r} \frac{p_m(r)p_{k-m}(r)}{v_r}.
$$

$$
(2) \mathbb{E}(W'-W)^4 = v_s^2 \left[\sum_{r=0}^n \left(8 \left(1 - \frac{\phi_s(t)}{v_s} \right) - 6 \left(1 - \frac{\phi_r(t)}{v_r} \right) \right) \frac{p_2(r)^2}{v_r} \right].
$$

Proof. For the first assertion, note that

$$
\mathbb{E}((W'-W)^k|i)
$$
\n
$$
= \frac{1}{(v_s)^{k/2}} \sum_{j=0}^n \frac{\mu_j}{|X|} \sum_{r=0}^n \frac{\phi_r(i)\phi_r(t)\phi_r(j)}{(v_r)^2} (\phi_s(j) - \phi_s(i))^k
$$
\n
$$
= \frac{1}{(v_s)^{k/2}} \sum_{m=0}^k (-1)^{k-m} {k \choose m} \phi_s(i)^{k-m} \sum_{r=0}^n \frac{\phi_r(i)\phi_r(t)}{(v_r)^2}
$$
\n
$$
\cdot \sum_{j=0}^n \frac{\mu_j}{|X|} \phi_s(j)^m \phi_r(j)
$$
\n
$$
= \frac{1}{(v_s)^{k/2}} \sum_{m=0}^k (-1)^{k-m} {k \choose m} \phi_s(i)^{k-m} (v_s)^m \sum_{r=0}^n \frac{\phi_r(i)\phi_r(t)p_m(r)}{(v_r)^2},
$$

where the last equality is Lemma [6.3.](#page-42-2) Consequently,

$$
\mathbb{E}((W'-W)^k) = \mathbb{E}(\mathbb{E}((W'-W)^k|i))
$$

= $(v_s)^{k/2} \sum_{m=0}^k (-1)^{k-m} {k \choose m}$

$$
\cdot \sum_{r=0}^n \frac{\phi_r(t)}{(v_r)^2} p_m(r) \sum_{i=0}^n \frac{\mu_i}{|X|} \frac{\phi_s(i)^{k-m} \phi_r(i)}{(v_s)^{k-m}}
$$

The first assertion now follows from Lemma [6.3.](#page-42-2)

For the second part, the first assertion gives that

$$
\mathbb{E}(W'-W)^4 = (v_s)^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_{r=0}^n \frac{\phi_r(t)}{v_r} \frac{p_m(r)p_{4-m}(r)}{v_r}.
$$

.

.

By page 654 of [\[MaSl\]](#page-51-18), $\phi_r(0) = v_r$ for all r. Thus specializing to $t = 0$ shows that

$$
0 = (v_s)^2 \sum_{m=0}^{4} (-1)^m {4 \choose m} \sum_{r=0}^{n} \frac{p_m(r) p_{4-m}(r)}{v_r}.
$$

So for general t , one has that

$$
\mathbb{E}(W'-W)^4 = -(v_s)^2 \sum_{m=0}^4 (-1)^m {4 \choose m} \sum_{r=0}^n \left(1 - \frac{\phi_r(t)}{v_r}\right) \frac{p_m(r)p_{4-m}(r)}{v_r}.
$$

The contribution from the $m = 0, 4$ terms is 0, since $p_0(r) = 0$ if $r \neq 0$. The contribution from the $m = 2$ term is

$$
-6(v_s)^2 \sum_{r=0}^n \left(1 - \frac{\phi_r(t)}{v_r}\right) \frac{p_2(r)^2}{v_r}.
$$

The $m = 1, 3$ terms are equal and together contribute

$$
8(v_s)^2 \sum_{r=0}^n \left(1 - \frac{\phi_r(t)}{v_r}\right) \frac{p_1(r)p_3(r)}{v_r} = 8(v_s)^2 \left(1 - \frac{\phi_s(t)}{v_s}\right) \frac{p_3(s)}{v_s}
$$

=
$$
8(v_s)^2 \left(1 - \frac{\phi_s(t)}{v_s}\right) \sum_{r=0}^n \frac{p_2(r)^2}{v_r},
$$

where the final equality is part 2 of Lemma [6.4.](#page-43-1) Adding the terms completes the proof of the second assertion. \Box

Arguing as in the proof of Theorem [3.13,](#page-12-0) and using the above lemmas, one obtains the following result.

Theorem 6.12. Fix $s \in \{0, \dots, n\}$, and let $W = \frac{\phi_s(i)}{\sqrt{v_s}}$ where i is chosen from the Plancherel measure of the association scheme. Fix $t \in \{1, \dots, n\}$. Then for all real x_0 ,

$$
\begin{split} & \left| \mathbb{P}(W \le x_0) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x_0} e^{-\frac{x^2}{2}} dx \right| \\ &\le \frac{v_s}{a} \sqrt{\sum_{r=1}^n \frac{p_2(r)^2}{v_r} \left(\frac{\phi_r(t)}{v_r} + 1 - \frac{2\phi_s(t)}{v_s}\right)^2} \\ &+ \frac{\sqrt{v_s}}{(\pi)^{1/4}} \left[\sum_{r=0}^n \left(8 - \frac{6}{a} \left(1 - \frac{\phi_r(t)}{v_r}\right)\right) \frac{p_2(r)^2}{v_r} \right]^{1/4}, \end{split}
$$

where $a = 1 - \frac{\phi_s(t)}{v_s}$ $\frac{s(t)}{v_s}$.

6.3. Example: Hamming Scheme. This subsection illustrates the theory of Subsection [6.2](#page-43-0) for the Hamming scheme $H(d, q)$, where d, q are positive integers and $q \geq 2$.

To begin we recall the definition of $H(d, q)$ and its basic properties, refer-ring the reader to Chapter 3 of [\[BaI\]](#page-49-2) for more details. The elements X of $H(d, q)$ are d-tuples of numbers chosen from $\{1, \dots, q\}$; clearly $|X| = q^d$. pair (x, y) is in R_i if x and y differ in exactly i coordinates. For $0 \leq i \leq d$, one has that $v_i = (q-1)^i {d \choose i}$ $\binom{d}{i}$ and $\mu_i = (q-1)^i \binom{d}{i}$ $\binom{d}{i}$. Thus the Plancherel measure on $\{0, \dots, d\}$ chooses i with probability $\frac{(q-1)^i {d \choose i}}{q^d}$ $\frac{1}{q^d}$. The numbers $\phi_s(i)$ are equal to

$$
\sum_{j=0}^s(-1)^j(q-1)^{s-j}\binom{i}{j}\binom{d-i}{s-j}.
$$

The polynomial ϕ_s in the variable i is known as a q-Krawtchouk polynomial.

In what follows $W = \frac{\phi_1(i)}{\sqrt{v_1}}$, where *i* is chosen from Plancherel measure of the Hamming scheme $H(d, q)$. The Hamming graph has vertex set X and an edge between two vertices if they differ in one coordinate. Thus the eigenvalues of the adjacency matrix of the Hamming graph are $\phi_1(i)$ with multiplicity μ_i , which motivates the study of W. Hora [\[Ho2\]](#page-50-14) shows that if $d \to \infty$ and $q/d \to 0$, then W converges in distribution to a normal random variable with mean 0 and variance 1.

In fact since $W(i) = \frac{(q-1)d-qi}{\sqrt{(q-1)d}}$ $\frac{(-1)d - qi}{(q-1)d}$ and $\frac{\mu_i}{|X|} = \frac{(q-1)^i {d \choose i}}{q^d}$ $\frac{q^{d}}{q^{d}}$, it is straightforward that W has the same distribution as

$$
\frac{Y_1 + \dots + Y_d}{\sqrt{(q-1)d}},
$$

where the Y's are independent random variables, each equal to $q - 1$ with probability $\frac{1}{q}$ and to -1 with probability $1-\frac{1}{q}$ $\frac{1}{q}$. Hence the Berry-Esseen theorem $[Du]$ shows that W satisfies a central limit theorem with the error term $C\sqrt{\frac{q}{d}}$ where C is a small explicit constant.

For comparison, let us study W using Theorem [6.12](#page-47-0) with $s = 1$ and $t = 1$. Then $a = \frac{q}{a}$ $\frac{q}{(q-1)d}$ and one computes that $p_2(0) = \frac{1}{(q-1)d}$, $p_2(1) = \frac{(q-2)}{(q-1)d}$, and $p_2(2) = 1 - \frac{1}{d}$ $\frac{1}{d}$. The first term in the error term of Theorem [6.12](#page-47-0) is then computed to equal $\sqrt{\frac{(q-2)^2}{(q-1)d}}$, which is less than $\sqrt{\frac{q}{d}}$. The second term in the error term of Theorem [6.12](#page-47-0) is computed to equal $\left(\frac{2q^2}{\pi(a-1)}\right)$ $\pi(q-1)d$ $\int^{1/4}$, which is at most $\left(\frac{2q}{d}\right)$ d $\int^{1/4}$. Thus one obtains a central limit theorem for W with error term $\sqrt{\frac{q}{d}} + \left(\frac{2q}{d}\right)$ d $\big)^{1/4}.$

To close this section, it is shown how our exchangeable pair can be used to give $O(d^{-1/2})$ bounds for q fixed. The key is Lemma [6.13,](#page-49-3) which shows that

the Markov chain L_1 is actually a birth-chain and computes its transition probabilities.

Lemma 6.13. The Markov chain L_1 on the set $\{0, \dots, d\}$ is a birth-death chain with transition probabilities

$$
L_1(i,j) = \begin{cases} \frac{i}{d(q-1)} & \text{if } j = i - 1\\ \frac{i}{d} \left(1 - \frac{1}{q-1} \right) & \text{if } j = i\\ 1 - \frac{i}{d} & \text{if } j = i + 1 \end{cases}
$$

Proof. Recall that $L_1(i, j) = \frac{\mu_j}{|X|} \sum_{r=0}^n$ $\phi_r(i) \phi_r(1) \phi_r(j)$ $\frac{\varphi_r(1)\varphi_r(j)}{(v_r)^2}$. From the formula for $\phi_r(l)$ one sees that

$$
\phi_r(l) = (q-1)^{r-l} \frac{\binom{d}{r}}{\binom{d}{l}} \phi_l(r)
$$

for any $0 \leq r, l \leq d$. Thus

$$
L_1(i,j) = \frac{\mu_j}{|X|d(\substack{d\\i)(\substack{d\\j)(q-1)^{i+j+1}}} \sum_{r=0}^n \binom{d}{r} (q-1)^r \phi_1(r) \phi_i(r) \phi_j(r).
$$

From page 152 of [\[MaSl\]](#page-51-18), there is a recurrence relation

$$
(i+1)\phi_{i+1}(r) = [(d-i)(q-1) + i - qr]\phi_i(r) - (q-1)(d-i+1)\phi_{i-1}(r).
$$

Since $\phi_1(r) = d(q-1) - qr$, the recurrence is equivalent to

$$
\phi_1(r)\phi_i(r) = i(q-2)\phi_i(r) + (i+1)\phi_{i+1}(r) + (q-1)(d-i+1)\phi_{i-1}(r).
$$

Applying this to expression for $L_1(i, j)$ at the end of the previous paragraph, the result follows from Lemma [6.2.](#page-42-1) \Box

Lemma [6.13](#page-49-3) implies that $|W' - W| \leq A$ with $A = \frac{q}{\sqrt{q}}$ $\frac{q}{(q-1)d}$. Thus one can apply the version of Theorem [6.12](#page-47-0) which would arise by using Theorem [2.2](#page-4-2) instead of Theorem [2.1.](#page-4-1) Recalling that $a = \frac{q}{a}$ $\frac{q}{(q-1)d}$, one obtains an error term of $\sqrt{\frac{q}{d}} + \frac{.41q^2 + 1.5q}{\sqrt{(q-1)d}}$ $\frac{q^2+1.5q}{(q-1)d}$. For q fixed this goes as $d^{-1/2}$.

7. Acknowledgements

The author was supported by NSA grant H98230-05-1-0031 and NSF grant DMS-0503901.

REFERENCES

- [AlD] Aldous, D. and Diaconis, P., Longest increasing subsequences: from patience sorting to the Baik-Deift-Johansson theorem, *Bull. Amer. Math. Soc.* 36 (1999), 413- 432.
- [BaI] Bannai, E. and Ito, T., *Algebraic combinatorics, I. Association schemes*, Benjamin/Cummings Publishing Co., Inc., Menlo Park, CA, 1984.
- [Bi] Biane, P., Permutation model for semi-circular systems and quantum random walks, *Pacific J. Math.* 171 (1995), 373-387.

- [BOO] Borodin, A., Okounkov, A., and Olshanski, G., Asymptotics of Plancherel measure for symmetric groups, *J. Amer. Math. Soc.* 13 (2000), 481-515.
- [BO1] Borodin, A. and Olshanski, G., Z-measures on partitions and their scaling limits, *European J. Combin.* 26 (2005), 795-834.
- [BO2] Borodin, A. and Olshanski, G., Harmonic functions on multiplicative graphs and interpolation polynomials, *Elecron. J. Combin.* 7 (2000), Research paper 28, 39 pages.
- [CF] Chatterjee, S. and Fulman, J., Stein's method for χ^2 approximation and spectral measure of Gelfand pairs, in preparation.
- [De] Deift, P., Integrable systems and combinatorial theory, *Notices Amer. Math. Soc.* 47 (2000), 631-640.
- [Di] Diaconis, P., *Group representations in probability and statistics*, Institute of Mathematical Statistics Lecture Notes, Volume 11, 1988.
- [DHol] Diaconis, P. and Holmes, S., Random walks on trees and matchings, *Elec. J. Probab.* 7 (2002), 17 pages (electronic).
- [DSh] Diaconis, P. and Shahshahani, M., Generating a random permutation with random transpositions, *Z. Wahr. Verw. Gebiete* 57 (1981), 159-179.
- [Du] Durrett, R., *Probability: Theory and examples, Second edition*, Duxbury Press, Belmont, CA, 1996.
- [EO] Eskin, A. and Okounkov, A., Asymptotics of branched coverings of a torus and volumes of moduli spaces of holomorphic differentials, *Invent. Math.* 145 (2001), 59-103.
- [F1] Fulman, J., Stein's method and Plancherel measure of the symmetric group, *Transac. Amer. Math. Soc.* 357 (2005), 555-570.
- [F2] Fulman, J., Stein's method, Jack measure, and the Metropolis algorithm, *J. Combin. Theory Ser. A* 108 (2004), 275-296.
- [F3] Fulman, J., An inductive proof of the Berry-Esseen theorem for character ratios, preprint [math.CO/0503227](http://arxiv.org/abs/math/0503227) at [http://xxx.lanl.gov.](http://xxx.lanl.gov)
- [F4] Fulman, J., Martingales and character ratios, to appear in *Transac. Amer. Math. Soc.*
- [HSS] Hanlon, P., Stanley, R., and Stembridge, J., Some combinatorial aspects of the spectra of normally distributed random matrices, in *Hypergeometric functions on domains of positivity, Jack polynomials, and applications*, Contemporary Mathematics, Volume 138 (1992), 151-174.
- [HHO] Hashimoto, Y., Hora, A., and Obata, N., Central limit theorems for large graphs: method of quantum decomposition, *J. Math. Phys.* 44 (2003), 71-88.
- [HH] Hoffman, P. and Humphreys, J., *Projective representations of the symmetric group*, Oxford University Press, New York, 1992.
- [Ho1] Hora, A., Central limit theorem for the adjacency operators on the infinite symmetric group, *Comm. Math. Phys.* 195 (1998), 405-416.
- [Ho2] Hora, A., Central limit theorems and asymptotic spectral analysis on large graphs, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* 1 (1998), 221-246.
- [I] Ivanov, V., The Gaussian limit for projective characters of large symmetric groups, *J. Math. Sci. (N.Y.)*, 121 (2004), 2330-2344. Translated from *Zap. Nauchn. Sem. POMI* 283 (2001), 73-97.
- [IK] Ivanov, V. and Kerov, S., The algebra of conjugacy classes in symmetric groups, and partial permutations, *J. Math. Sci. (New York)* 107 (2001), 4212-4230. Translated from *Zap. Nauchn. Sem. POMI* 256 (1999), 95-120.
- [IO] Ivanov, V. and Olshanski, G., Kerov's central limit theorem for the Plancherel measure on Young diagrams, in *Symmetric Functions 2001: Surveys of developments and perspectives*, Kluwer Academic Publishers, Dodrecht, 2002.
- [J] Johansson, K., Discrete orthogonal polynomial ensembles and the Plancherel measure, *Ann. of Math.* 153 (2001), 259-296.
- [K1] Kerov. S.V., Gaussian limit for the Plancherel measure of the symmetric group, *Compt. Rend. Acad. Sci. Paris, Serie I*, 316 (1993), 303-308.
- [K2] Kerov, S.V., Anisotropic Young diagrams and Jack symmetric functions, *Funct. Anal. Appl.* 34 (2000), 41-51.
- [Le] Letac, G., Les fonctions sphériques d'un couple de Gelfand symétrique et les chaˆines de Markov, *Adv. Appl. Probab.* 14 (1982), 272-294.
- [Mac] Macdonald, I., *Symmetric functions and Hall polynomials*, Second edition, Oxford University Press, New York, 1995.
- [MaSl] Macwilliams, F. and Sloane, N., *The theory of error-correcting codes*, Elsevier Science B.V., Amsterdam, 1977.
- [Mat] Matsumoto, S., Correlation functions of the shifted Schur measure, preprint [math.CO/0312373](http://arxiv.org/abs/math/0312373) at [http://xxx.lanl.gov.](http://xxx.lanl.gov)
- [O1] Okounkov, A., Random matrices and random permutations, *Internat. Math. Res. Notices* 20 (2000), 1043-1095.
- [O2] Okounkov, A., The uses of random partitions, preprint [math-ph/0309015](http://arxiv.org/abs/math-ph/0309015) at [http://xxx.lanl.gov.](http://xxx.lanl.gov)
- [Re] Reinert, G., Couplings for normal approximations with Stein's method, in *Microsurveys in discrete probability*, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., Volume 41 (1998), 193-207.
- [RiRo1] Rinott, Y. and Rotar, V., On coupling constructions and rates in the CLT for dependent summands with applications to the antivoter model and weighted Ustatistics, *Ann. of Appl. Probab.* 7 (1997), 1080-1105.
- [RiRo2] Rinott, Y. and Rotar, V., Normal approximations by Stein's method, *Decis. Econ. Finance* 23 (2000), 15-29.
- [Sa] Sagan, B., *The symmetric group. Representations, combinatorial algorithms, and symmetric functions*, Second edition, Springer-Verlag, New York, 2001.
- [Se] Serre, J-P., *Linear representations of finite groups*, Springer-Verlag, New York, 1977.
- [ShSu] Shao, Q., and Su, Z., The Berry-Esseen bound for character ratios, preprint (2004).
- [Sn1] Sniady, P., Asymptotics of characters of symmetric groups, Gaussian fluctuations of Young diagrams and genus expansion, preprint [math.CO/0411647](http://arxiv.org/abs/math/0411647) at [http://xxx.lanl.gov.](http://xxx.lanl.gov)
- [Sn2] Sniady, P., Gaussian fluctuations of characters of symmetric groups and of Young diagrams, preprint [math.CO/0501112](http://arxiv.org/abs/math/0501112) at [http://xxx.lanl.gov.](http://xxx.lanl.gov)
- [Stn1] Stein, C., *Approximate computation of expectations*, Institute of Mathematical Statistics Lecture Notes, Volume 7, 1986.
- [Stm] Stembridge, J., On Schur's Q-functions and the primitive idempotents of a commutative Hecke algebra, *J. Algebraic Combin.* 1 (1992), 71-95.
- [Te] Terras, A., *Fourier analysis on finite groups and applications*, London Math. Society Student Texts 43, Cambridge University Press, Cambridge, 1999.
- [TW] Tracy, C. and Widom, H., A limit theorem for shifted Schur measures, *Duke Math. Journal* 123 (2004), 171-208.