EXTRA-LARGE METRICS

IGOR RIVIN

A Brack We show that every two dimensional spherical cone metric with all cone angles greater than 2π and the lengths of all closed geodesics greater than 2π admits a triangulation whose 0skeleton is precisely the set of cone points – this is, in fact, the Delaunay triangulation of the set of cone points.

$1. I$

In this note we study *extra large* spherical cone manifolds in dimension 2 (though many of our resultsz and techniques extend to higher dimensions.

A 2-dimensional *spherical cone manifold* is a metric space where all but finitely many points has a neighborhood isometric to a neighborhood of a point on the round sphere S 2 . The exceptional points (*cone points*) have neighboroods isometric to a spherical cone, the angle of which is the *cone angle* at that point.

If *M* is a cone manifold, we define a *geodesic* to be a locally length minimizing curve on *M*. It is easy to see that such a curve is locally a great circle, except at the cone points. There, the geodesic must have the property that it subtends an angle no smaller than π on either side. Consequently, no geodesics can *pass through* cone points, where the cone angles are smaller than 2π (such cone points are known as *positively curved* cone points, since the curvature of a cone point is defined as 2π less the cone angle at the point).

We say that a spherical cone manifold is *extra large* if

(1) All the cone points are negatively curved.

(2) All closed geodesics are longer than 2π .

Such spaces are of considerable importance in geometry in general (due to work of A.D.Aleksandrov, M. Gromov, and then R. Charney and M. Davis [\[1\]](#page-4-0)), and in three-dimensional hyperbolic geometry in particular, due in large part to the results of the author ([\[2,](#page-4-1) [6,](#page-4-2) [3,](#page-4-3) [5,](#page-4-4) [4\]](#page-4-5)),

Date: July 5, 2018.

¹⁹⁹¹ *Mathematics Subject Classification.* 52B11, 52B10, 57M50.

Key words and phrases. triangulation, extra-large, hyperbolic polyhedra.

2 IGOR RIVIN

who showed that the polar duals of convex compact polyhedra in \mathbb{H}^3 are precisely the extra large spherical cone manifolds homeomorphic to S 2 . In that work, the term *extra large* was not used – it was invented by Gromov, to describe the vertex links in negatively curved spaces.

The main objective of this paper is to show:

Theorem 1. *An extra large spherical cone surface admits a cell decomposition whose* 0*-skeleton is precisely the set of cone points.*

For the impatient reader, we first give the recipe for constructing the cell decomposition whose existence is postulated in Theorem [1:](#page-1-0)

First, recall that the *Voronoi diagram* V*^P* of a metric space *M* with respect to a point set $P = \{p_1, \ldots, p_n, \ldots\}$ is the decomposition of M into *Voronoi cells*

$$
V_i = \{x \in M \big| d(x, p_i) \leq d(x, p_j), \quad \forall j\}.
$$

Clearly,

$$
\bigcup_i V_i = M,
$$

and

$$
\overset{\circ}{V}_i \cap \overset{\circ}{V}_j = \emptyset, \quad i \neq j.
$$

As will be shown below, each *Vⁱ* is a geodesic polygon, and the Voronoi diagram V*^P* is a cell decomposition of *M*.. The *Delaunay tesselation* \mathcal{D}_P of *M* with respect to *P* is the Poincaré dual of \mathcal{V}_P : its edges corrspond to pairs p_i, p_j of sites whose Voronoi cells share an edge, while its faces correspond to points of *M* equidistant from three or more elements of *P*. The cells of \mathcal{D}_P are convex, and so the tesselation D*^P* can be completed to a triangulation of *M*.

To push the program above through, we will need a number of steps.

$2. T$

We define the *injectivity radius* of the space *M at a point p* as the radius of the largest disk in the tangent space of *p* for which the exponential map is an embedding. The injectivity radius of *M* is the infimum over all points *p* of the injectivity radii of *M* at *p*. In simpler terms, the injectivity radius of *M* is the smallest *d* such that there exist at least two distinct curves from p to q realizing $d(p, q) = d$.

Our first result is:

Theorem 2. *A space M is extra large if and only if the cone points of M are negatively curved and the injectivity radius of M is greater than* π.

Proof. Let *p*, *q* be the pair of points realizing the injectivity radius. This means that there are two shortest curves γ_1, γ_2 of length $L =$ $\ell(\gamma_1) = \ell(\gamma_2) \leq \pi$ connecting p to q, and L is the smallest with this property.

Let $\gamma = \gamma_1 \cup \gamma_2$. If γ is geodesic, then $L(\gamma) \leq 2\pi$, so we have a contradiction to extra-largeness. If not, suppose (without loss of generality) that γ has a "corner" at p . That means that on one side, the angle α subtended by γ at p is smaller than π . If γ_1 and γ_2 are both smooth, then take the bisector of α and move p a very small distance ρ along the bisector. By elementary spherical eometry,

$$
\frac{d\ell(\gamma_1)}{d\rho} = \frac{d\ell(\gamma_2)}{d\rho} < 0,
$$

which contradicts the minimality of *L*. If (without loss of generality) γ_1 is *not* smooth, while γ_2 is, let *x* be the cone point of γ_1 closest to *p*. Note that if the angle subtended by γ_1 at x on the side of the corner at p equals π , then x can be treated as a smooth point, so the correct definition of *x* is: the closest point of γ_1 to *p*, where γ_1 is not smooth on the side of the corner (if such a point does not exist between *p* and *q*, then we find ourselves back in the smooth case, which corresponds to $x = q$.). Let $L_x = d(p, x)$.

In any event, now, instead of the bisector of the angle α at *p*, we pick a direction, such that

$$
\frac{d\ell(\gamma_2)}{d\rho} = \frac{dL_x}{d\rho} < 0.
$$

Such a direction exists by the intermediate value theorem. The above argument adapts in the obvious way if both γ_1 and γ_2 are singular. \Box

$$
3. V
$$

Lemma 1. *For all* $x \in V_i$, $d(x, p_i) < \pi/2$..

Proof. Suppose that there exists an *x* contradicting the assertion of the Lemma. Then the distance from *x* to the cone locus of *M* is at least $\pi/2$, and so there is a smooth hemisphere around *x*. The boundary of that hemisphere is a closed geodesic of length 2π .

Corollary 1. *The diameter of the Voronoi cell Vⁱ is less than* π.

We will need the following simple lemma from spherical geometry:

Theorem 3. *The boundary of a Voronoi cell Vⁱ is a convex polygonal curve.*

Proof. Let $x \in V_i$. Let $r = d(x, p_i)$; we know that $r < \pi/2$. Consider the disk *Dx*(*r*) of radius *r* around *x*.

There are the following possibilities:

Firstly, p_i might be the *only* cone point in $D_r(x)$. In that case, a neighborhood of x is in V_i , and so x is in the interior of V_i .

Secondly, there msy be exactly one other point p_j such that $d(p_j)$ = *r*. In that case, a small geodesic segment bisecting the angle *pixp^j* lies in $V_i \cap V_j$.

Thirdly, there can be a number of points pi , $p_{j_1}, p_{j_2}, \ldots, p_{j_k}$ at distance *r* from *x*. In that case a small part of the cone from *x* to the Voronoi region of p_i on the boundary of $D_r(x)$ lies in V_i – note that this argument works in arbitrary dimension.

Theorem 4. A Voronoi cell V_i is star-shaped with respect to p_i .

Proof. Let $x \in V_i$, and let *y* be on the segment $p_i x$. By the triangle inequality, we see that

$$
d(y, \partial D_x(r)) + d(y, x) > r.
$$

Since for any $j \neq i$ we have that

 $d(y, p_i) > d(y, \partial D_x(r))$

the assertion of the Theorem follows. \Box

Theorem 5. *Every Voronoi cell Vⁱ is convex.*

Proof. By the preceding result, every Voronoi cell *Vⁱ* is a stashaped subset of a cone of radius $\pi/2$ centered on p_i , with geodesically convex boundary. Take two points *p* and *q* in *Vⁱ* . If one of the angles *ppiq* does not exceed π , the result follows from elementary spherical geometry. If both the angles pp_iq are at least π , the broken line pp_iq is geodesic.

П

Theorem 6. Let V_i , V_j be two Voronoi cells, then $V_i \cap V_j$ is connected.

Proof. Let $p, q \in V_i \cap V_j$. here is a shortest geodesic γ_i from p to q in *V*_{*i*} and a shortest geodesic γ_j from p to q in V_j . Since the diameters of *V*^{*i*} and *V*^{*j*} are smaller than π, it follows that $γ_i = γ_j$. Thus, $γ = γ_i =$ $\gamma_j \subseteq V_i \cap V_j$, hence $V_i \cap V_j$ is path connected. In fact, the argument (together with the results above) easily shows that $V_i \cap V_j$ is an edge $\overline{\text{of both}}$.

The above results sow that $V_P = \{V_1, \ldots, V_n, \ldots\}$ is a simplicial cell decomposition, and so its dual is a cellulation of *M* with vertices at p_1, \ldots, p_n, \ldots . The cells of cellulation are convex (in fact inscribed in circles; The centers are precisely the corners of the boundaries of the cells *Vⁱ* .), and so the proof of Theorem [1](#page-1-0) is complete.

4. R

An identical argument with the appropriate modification of the extra-largeness hypothesis can be used to show an analogous result for a Riemannian surface with cone singularies. In particular, for Euclidean and Hyperbolic cone surfaces, it is sufficient to require the cone angles to be non-positively curved.

R

- [1] Ruth Charney and Michael Davis. The polar dual of a convex polyhedral set in hyperbolic space. *Michigan Math. J.*, 42(3):479–510, 1995.
- [2] Igor Rivin. *On the Geometry of Convex Polyhedra in Hyperbolic 3-Space*. PhD thesis, Princeton University, July 1986.
- [3] Igor Rivin. On the geometry of ideal polyhedra in hyperbolic 3-space. 32(1):87– 92, January 1993.
- [4] Igor Rivin. Euclidean structures on simplicial surfaces and hyperbolic volume. *Annals of Mathematics (ser. 2)*, 139(3):553–580, May 1994.
- [5] Igor Rivin. Combinatorial optimization in geometry. *Advances in Applied Mathematics*, 31(1):242–271, 2003. arxiv.org preprint math.GT/9907032.
- [6] Igor Rivin and C.D.Hodgson. A characterization of compact convex polyhedra in hyperbolic 3-space. *Inventiones Mathematicae*, pages 77–111, January 1993. Corrigendum, vol 117, page 359.