Rigidity of holomorphic generators and one-parameter semigroups

Mark Elin

Department of Mathematics, ORT Braude College,

P.O. Box 78, Karmiel 21982, Israel

e-mail: mark.elin@gmail.com

Marina Levenshtein

Department of Mathematics,

The Technion — Israel Institute of Technology 32000 Haifa, Israel e-mail: marlev@list.ru

David Shoikhet

Department of Mathematics, ORT Braude College,

P.O. Box 78, Karmiel 21982, Israel Roberto Tauraso

Dipartimento di Matematica, Università di Roma 'Tor Vergata', Via della Ricerca Scientifica, 00133 Roma, Italy e-mail: tauraso@mat.uniroma2.it

Abstract

In this paper we establish a rigidity property of holomorphic generators by using their local behavior at a boundary point τ of the open unit disk Δ . Namely, if $f \in Hol(\Delta, \mathbb{C})$ is the generator of a oneparameter continuous semigroup ${F_t}_{t\geq0}$, we state that the equality $f(z) = o(|z - \tau|^3)$ when $z \to \tau$ in each non-tangential approach region at τ implies that f vanishes identically on Δ . Note, that if F is a self-mapping of Δ then $f = I - F$ is a generator, so our result extends the boundary version of the Schwarz Lemma obtained by D. Burns and

S. Krantz. We also prove that two semigroups ${F_t}_{t\geq0}$ and ${G_t}_{t\geq0}$, with generators f and g respectively, commute if and only if the equality $f = \alpha g$ holds for some complex constant α . This fact gives simple conditions on the generators of two commuting semigroups at their common null point τ under which the semigroups coincide identically on ∆.

1 Introduction.

Let $\Delta = \{z \in \mathbb{C} : |z| < 1\}$ be the open unit disk in the complex plane \mathbb{C} , and let $H = \{z \in \mathbb{C} : \text{Re } z > 0\}$ be the right half-plane. We denote by $\text{Hol}(\Delta, D)$ the set of all holomorphic functions on Δ which map Δ into a set $D \subset \mathbb{C}$, and by Hol(Δ) the set of all holomorphic self-mappings of Δ , i.e., $Hol(\Delta) = Hol(\Delta, \Delta).$

The problem of finding conditions for a holomorphic function F to coincide identically with a given holomorphic function G when they have a similar behavior on some subset of Δ , has been studied by many mathematicians.

The following assertions are classical:

• If F and G are holomorphic in Δ and $F = G$ on a subset of Δ that has a nonisolated point, then $F \equiv G$ on Δ (Vitali's uniqueness principle).

• If F and G are holomorphic in Δ and continuous on $\overline{\Delta}$, and $F = G$ on some arc γ of the boundary $\partial \Delta$, then $F \equiv G$ on Δ .

In the point of view of complex dynamics it is natural to study conditions on derivatives of F and G at specific points to conclude that $F \equiv G$.

If, for example, G is the identity mapping I and $\tau \in \Delta$ is the Denjoy–Wolff point of $F \in Hol(\Delta)$, then the equalities $F(\tau) = G(\tau)$ and $F'(\tau) = G'(\tau)$ provide $F \equiv G$ by the Schwarz Lemma. The same conclusion holds for an arbitrary holomorphic function G on Δ , if F commutes with G and satisfies the conditions $F(\tau) = G(\tau) = \tau$ and $F'(\tau) = G'(\tau) \neq 0$ (see, for instance, $[9], [6]$ $[9], [6]$ $[9], [6]$.

Different "identity principles" have been recently studied by several mathematicians under suitable boundary conditions. In general, the following three cases are considered.

 (A) G is the identity mapping;

(B) G is an arbitrary self-mapping of Δ , and F commutes with G, i.e., $F \circ G = G \circ F$;

 (C) G is a constant mapping.

Regarding Case A the following result is due to D. Burns and S. Krantz.

Theorem A ([\[7\]](#page-18-2)) Let $F \in Hol(\Delta)$ and

$$
F(z) = 1 + (z - 1) + O((z - 1)^4).
$$
 (1)

Then $F \equiv I$.

For Case B a uniqueness theorem was given by R. Tauraso in [\[17\]](#page-19-0) (see also [\[6\]](#page-18-1)). To formulate this result we need the following notation. Let $F \in Hol(\Delta)$ and $\tau \in \partial \Delta$. We say that $F \in C_K^m(\tau)$ if it admits the following representation

$$
F(z) = \tau + F'(\tau)(z - \tau) + \ldots + \frac{F^{(m)}(\tau)}{m!}(z - \tau)^m + o(|z - \tau|^m)
$$

when $z \to \tau$ in each non-tangential approach region at τ . Moreover, we say that $F \in C^m(\tau)$ if the limit is taken in the full disk.

Theorem B ([\[17\]](#page-19-0)) Let $F, G \in Hol(\Delta)$ be commuting functions with a common Denjoy–Wolff point $\tau \in \partial \Delta$. If one of the following conditions holds then $F \equiv G$.

(*i*) $F'(\tau) = G'(\tau) < 1;$

(ii) $F \in C^2(\tau)$, $G \in C^2_K(\tau)$, $F'(\tau) = 1$, $F''(\tau) = G''(\tau) \neq 0$ and $\text{Re } \tau F''(\tau) > 0;$

(iii) F, $G \in C^2(\tau)$, $F'(\tau) = 1$, $F''(\tau) = G''(\tau) \neq 0$ and $\text{Re } \tau F''(\tau) = 0$; (iv) $F \in C^3(\tau)$, $G \in C^3_K(\tau)$, $F'(\tau) = 1$, $F''(\tau) = G''(\tau) = 0$ and $F'''(\tau) = G'''(\tau).$

For Case C, when G is a constant mapping, the following fact is an immediate consequence of the Julia–Wolff–Carathéodory Theorem.

• If $F \in Hol(\Delta, \overline{\Delta})$, then the conditions $\lim_{r \to 1^-} F(r\tau) = \tau$ and $\lim_{r \to 1^-} F'(r\tau) = 0$ for some $\tau \in \partial \Delta$ imply that $F \equiv \tau$.

In fact, the considering of holomorphic functions f which are not necessarily self-mappings is more relevant in this situation. Various results in this direction were established by S. Migliorini and F. Vlacci in [\[13\]](#page-18-3).

In what follows we denote by symbol $\angle \lim_{z \to \tau}$ the angular limit of a function defined in Δ at a boundary point $\tau \in \partial \Delta$.

Theorem C (see [\[13\]](#page-18-3)) Let $\tau \in \partial \Delta$. If $f \in Hol(\Delta, \overline{H})$, then

$$
\angle \lim_{z \to \tau} \frac{f(z)}{z - \tau} = 0 \tag{2}
$$

implies that $f \equiv 0$.

More general, if $f \in Hol(\Delta, \mathbb{C})$, and $f(\Delta)$ is contained in a wedge of angle $\pi\alpha$, $0 < \alpha \leq 2$, with vertex at the origin, then the condition

$$
\angle \lim_{z \to \tau} \frac{f(z)}{(z - \tau)^{\alpha}} = 0
$$
\n(3)

implies that $f \equiv 0$.

Although the classes Hol(Δ) of holomorphic self-mappings of Δ and $Hol(\Delta, H)$ of functions with positive real part are connected by the composition with the Cayley transform, Theorem A is not a direct consequence of Theorem C, and conversely.

In this note we find rigidity principles for some classes of holomorphic functions produced by continuous dynamical systems, which are related to both Hol(Δ) and Hol(Δ , H). In particular, by this way one can establish a bridge between Theorems A and C.

We consider, inter alia, the class of mappings $F \in Hol(\Delta, \mathbb{C})$ which are continuous on $\overline{\Delta}$ and satisfy the boundary flow-invariance condition

$$
\operatorname{Re} F(z)\overline{z} \le 1, \, z \in \partial \Delta. \tag{4}
$$

In particular, each function $F \in Hol(\Delta)$ which is continuous on $\overline{\Delta}$ belongs to this class.

Condition [\(4\)](#page-3-0) can be rewritten in the form

$$
\operatorname{Re} f(z)\overline{z} \ge 0, \, z \in \partial \Delta,\tag{5}
$$

where

$$
f(z) = z - F(z). \tag{6}
$$

Note that each mapping f satisfying [\(5\)](#page-3-1) belongs to the class $\mathcal{G}(\Delta)$ of so-called semigroup generators on Δ .

Our main purpose is to establish boundary conditions for a function $f \in \mathcal{G}(\Delta)$ to vanish on Δ identically.

First, we recall that a family $S = \{F_t\}_{t>0} \subset Hol(\Delta)$ is said to be a one-parameter continuous semigroup on Δ if

- (i) $F_t(F_s(z)) = F_{t+s}(z)$ for all $t, s \ge 0$,
- (ii) $\lim_{t \to 0^+} F_t(z) = z$ for all $z \in \Delta$.

Furthermore, it follows by a result of E. Berkson and H. Porta [\[5\]](#page-18-4) that each continuous semigroup is differentiable in $t \in \mathbb{R}^+ = [0, \infty)$, (see also [\[1\]](#page-17-0) and [\[14\]](#page-18-5)). So, for each continuous semigroup $S = \{F_t\}_{t\geq 0} \subset Hol(\Delta)$, the limit

$$
\lim_{t \to 0^+} \frac{z - F_t(z)}{t} = f(z), \quad z \in \Delta,
$$
\n⁽⁷⁾

exists and defines a holomorphic mapping $f \in Hol(\Delta, \mathbb{C})$. This mapping f is called the (infinitesimal) generator of $S = \{F_t\}_{t \geq 0}$. Moreover, the function $u(=u(t, z))$, $(t, z) \in \mathbb{R}^+ \times \Delta$, defined by $u(t, z) = \widetilde{F}_t(z)$ is the unique solution of the Cauchy problem

$$
\begin{cases}\n\frac{\partial u(t,z)}{\partial t} + f(u(t,z)) = 0, \\
u(0,z) = z, \quad z \in \Delta.\n\end{cases}
$$
\n(8)

The class of all holomorphic generators on Δ is denoted by $\mathcal{G}(\Delta)$.

Note, that if $F \in Hol(\Delta)$, then the function $f = I - F$ belongs to $\mathcal{G}(\Delta)$ (see Corollary 3.3.1 in $|15|$).

The following assertion combines characterizations of the class $\mathcal{G}(\Delta)$ obtained in $[2]$, $[3]$ and $[5]$.

Proposition 1 Let $f \in Hol(\Delta, \mathbb{C})$. The following are equivalent:

(i) f is a semigroup generator on Δ ; (ii) Re $f(z)\overline{z} \geq$ Re $f(0)\overline{z}(1-|z|^2)$ for all $z \in \Delta$;

(iii) there exists a unique point $\tau \in \overline{\Delta}$ such that

$$
f(z) = (z - \tau)(1 - \overline{\tau}z)g(z), \ z \in \Delta,
$$
\n(9)

where $q \in Hol(\Delta, \mathbb{C})$, Re $q(z) \geq 0$.

(iv) f admits the representation

$$
f(z) = a - \overline{a}z^2 + zp(z),
$$

where $a \in \mathbb{C}$ and $p \in Hol(\Delta, \mathbb{C})$ with $\text{Re } p(z) \geq 0$.

Remark 1 The point τ in [\(9\)](#page-4-0) is the Denjoy–Wolff point of the semigroup ${F_t}_{t\geq0}$ generated by f. If $\tau \in \Delta$ then $f(0) = 0$ and $\text{Re } f'(\tau) \geq 0$. If $\tau \in \partial \Delta$ then the angular limit $\angle \lim_{z \to \tau} \frac{f(z)}{z-\tau} =: f'(\tau)$ exists and is a nonnegative real number (see $[10]$).

2 Rigidity of infinitesimal generators.

Theorem 1 Let $f \in \mathcal{G}(\Delta)$. Suppose that for some $\tau \in \partial \Delta$

$$
f(z) = a(z - \tau)^3 + o(|z - \tau|^3)
$$

when $z \to \tau$ in each non-tangential approach region at τ . Then $a\tau^2$ is a nonnegative real number. Moreover, $a = 0$ if and only if $f \equiv 0$.

To prove Theorem 1 we need the following lemma.

Lemma 1 Let $g \in Hol(\Delta, \overline{H})$. Then for each $\tau \in \partial \Delta$ the limit

$$
k = \angle \lim_{z \to \tau} \frac{g(z)}{1 - \overline{\tau}z} \tag{10}
$$

is either a nonnegative real number or infinity. Moreover, $q \equiv 0$ if and only if $k = 0$.

Proof. Denote by $C_{\tau}(z) = \frac{\tau - z}{\tau + z}$ the Cayley transform and set $h = C_{\tau}^{-1} \circ g \in$ $Hol(\Delta, \overline{\Delta})$. By the Julia–Wolff–Carathéodory theorem the limit

$$
\beta_h = \angle \lim_{z \to \tau} \frac{\tau - h(z)}{\tau - z}
$$

exists and is either a nonnegative real number or infinity. Moreover, $\beta_h = 0$ if and only if $h \equiv \tau$.

For any $z \in \Delta$ we have

$$
\frac{g(z)}{1-\overline{\tau}z} = \frac{\tau - h(z)}{\tau - z} \cdot \frac{\tau}{\tau + h(z)}.
$$
\n(11)

Hence, $k = 0$ if and only if $\beta_h = 0$, and therefore $g \equiv 0$.

If β_h is a positive real number, $\beta_h > 0$, then $\angle \lim_{z \to \tau} h(z) = \tau$ and, consequently,

$$
k = \angle \lim_{z \to \tau} \frac{\tau - h(z)}{\tau - z} \cdot \angle \lim_{z \to \tau} \frac{\tau}{\tau + h(z)} = \frac{\beta_h}{2} > 0.
$$

Let $\beta_h = \infty$. Since Re $\frac{\tau}{\tau + h(z)} \geq \frac{1}{2}$ $\frac{1}{2}$, formula [\(11\)](#page-5-0) implies that $k = \infty$.

Alternative proof. If $g \neq 0$, then the function p defined by $p(z) := \frac{1}{g(z)}$ belongs to Hol(Δ , H). It is easy to see that for all $\zeta \in \partial \Delta$ the expression $(1-z\overline{\tau})(1+z\overline{\zeta})$ $\frac{1}{1-z\overline{\zeta}}$ is bounded on each non-tangential approach region at τ . Then it follows by the Riesz–Herglots formula that

$$
\angle \lim_{z \to \tau} (1 - z\overline{\tau}) p(z) = \angle \lim_{z \to \tau} \oint_{\partial \Delta} \frac{(1 - z\overline{\tau})(1 + z\overline{\zeta})}{1 - z\overline{\zeta}} dm_p(\zeta) = 2m_p(\tau) \ge 0,
$$

where dm_p is a probability measure on $\partial\Delta$. Setting $k = \frac{1}{2m_p(\tau)}$ we get our assertion.

Proof of Theorem 1. Since

$$
\angle \lim_{z \to \tau} \frac{f(z)}{z - \tau} = 0,
$$

it follows from [\[10\]](#page-18-8) that $\tau \in \partial \Delta$ is the Denjoy–Wolff point for the semigroup ${F_t}_{t\geq0}$ generated by f. Then by Proposition 1 the function f admits the representation [\(9\)](#page-4-0):

$$
f(z) = (z - \tau)(1 - z\overline{\tau})g(z)
$$

with some $g \in Hol(\Delta, \overline{H})$. Hence, by Lemma 1

$$
a\tau^2 = \tau^2 \angle \lim_{z \to \tau} \frac{f(z)}{(z - \tau)^3} = \angle \lim_{z \to \tau} \frac{g(z)}{1 - \overline{\tau}z} = k \ge 0.
$$

Obviously, $a = 0$ if and only if $k = 0$. In this case $g \equiv 0$, so $f \equiv 0$.

Corollary 1 (cf. Theorem 5 in [\[6\]](#page-18-1).) Let $F \in Hol(\Delta, \mathbb{C})$ be continuous on $\overline{\Delta}$ and satisfy the boundary condition

$$
\operatorname{Re} F(z)\overline{z} \le 1, \, z \in \partial \Delta.
$$

If F admit the representation

$$
F(z) = \tau + (z - \tau) + b(z - \tau)^3 + o(|z - \tau|^3)
$$

when $z \to \tau$ in each non-tangential approach region at some point $\tau \in \partial \Delta$, then $b\tau^2 \leq 0$. Moreover, $b = 0$ if and only if $F \equiv I$.

As a consequence of Lemma 1 we also obtain the following assertion.

Corollary 2 Let $f \in \mathcal{G}(\Delta)$ be such that $f(\tau) = 0$ for some $\tau \in \partial \Delta$ and $f(0) = a \in \mathbb{C}$. Suppose that f has a finite angular derivative at τ . Then $f'(\tau)$ is a real number with $f'(\tau) \leq -2 \operatorname{Re}(\overline{a}\tau)$. Moreover, $f'(\tau) = -2 \operatorname{Re}(\overline{a}\tau)$ if and only if f generates a group of automorphisms.

Proof. By Proposition 1 (iv) f admits the representation

$$
f(z) = a - \overline{a}z^2 + zp(z), \ z \in \Delta,\tag{12}
$$

where $p \in Hol(\Delta, \mathbb{C})$ with $\text{Re } p(z) \geq 0$.

Since $f(\tau) = 0$, we have $p(\tau) = \overline{\alpha}\tau - a\overline{\tau} = 2i \operatorname{Im}(\overline{\alpha}\tau)$ is pure imaginary. Then it follows from [\(12\)](#page-7-0), that

$$
f'(\tau) = \angle \lim_{z \to \tau} \frac{a - \overline{a}z^2 + zp(z)}{z - \tau} = -2 \operatorname{Re}(a\overline{\tau}) + \angle \lim_{z \to \tau} \frac{p(z) - 2i \operatorname{Im}(\overline{a}\tau)}{z\overline{\tau} - 1}.
$$

Applying Lemma 1 to the function $g(z) = p(z) - 2i \text{Im}(\overline{a}\tau)$, we get $f'(\tau) \leq -2 \operatorname{Re}(a\overline{\tau}).$

Moreover, $f'(\tau) = -2 \operatorname{Re}(a\overline{\tau})$ if and only if $p \equiv 2i \operatorname{Im}(\overline{a}\tau)$, i.e., $f(z) =$ $a + 2i \operatorname{Im}(\overline{a}\tau) \cdot z - \overline{a}z^2$.

By Proposition 3.5.1 in [\[15\]](#page-18-6) each function of the form $f(z) = a + ibz - \overline{a}z^2$, with $a \in \mathbb{C}$ and $b \in \mathbb{R}$, generates a group of automorphisms of Δ . The proof is complete.

Corollary 3 Let $F \in Hol(\Delta)$ be such that $F(\tau) = \tau$ and $F(0) = a, a \in \Delta$. Suppose that F has a finite angular derivative at τ . Then $F'(\tau) \geq 1 + \tau$ $2 \text{Re}(\overline{a}\tau)$.

Proof. By a result in [\[15,](#page-18-6) Corollary 3.3.1] the function $f(z) = z - F(z)$, $z \in$ Δ is a generator of a one-parameter semigroup. By our assumptions we have $f(\tau) = 0$ and $f(0) = -a$. Hence, by Corollary 2 $f'(\tau) \leq -2 \operatorname{Re}(\overline{a}\tau)$, and $F'(\tau) \geq 1 + 2 \operatorname{Re}(\overline{a}\tau)$.

Now let us consider a class of functions $f \in Hol(\Delta, \mathbb{C})$ which are continuous on $\overline{\Delta}$ and satisfy the boundary condition

$$
\operatorname{Re} f(z)\overline{z} \ge |f(z)|\cos\frac{\alpha\pi}{2} \quad \text{for all} \quad z \in \partial\Delta,\tag{13}
$$

for some $\alpha \in (0, 2]$. As we already mentioned if $\alpha \leq 1$ then condition [\(13\)](#page-7-1) implies $f \in \mathcal{G}(\Delta)$ (cf. Proposition 1 (ii)). Conversely, if $f \in \mathcal{G}(\Delta)$ is continuous on $\overline{\Delta}$, then [\(13\)](#page-7-1) holds with $\alpha = 1$. So, this class generalize in a sense the class of holomorphic generators which are continuous on Δ .

Theorem 2 Let $f \in Hol(\Delta, \mathbb{C})$ be continuous on $\overline{\Delta}$ and satisfy the condition [\(13\)](#page-7-1). Then the condition

$$
\lim_{\substack{z \to \tau \\ z \in \Delta}} \frac{f(z)}{(z - \tau)^{2+\alpha}} = 0 \quad \text{for some} \quad \tau \in \partial \Delta \tag{14}
$$

implies that $f \equiv 0$.

Proof. Denote

$$
g(z) = \frac{f(z)}{(z-\tau)(1-\overline{\tau}z)}.
$$

The continuity of f and (14) imply that this function is continuous (consequently, bounded) on Δ .

Now we rewrite [\(13\)](#page-7-1) in the form:

$$
-\operatorname{Re}\left[\overline{\tau}(\tau-z)^2 g(z)\overline{z}\right] \ge |\tau-z|^2 \cdot |g(z)| \cdot \cos\frac{\alpha \pi}{2}, \quad z \in \partial \Delta.
$$

Hence,

$$
\operatorname{Re} g(z) \ge |g(z)| \cdot \cos \frac{\alpha \pi}{2}, \quad z \in \partial \Delta \setminus \{\tau\}.
$$

This inequality also holds at the point τ because of the continuity of g.

It follows from the subordination principle for subharmonic functions (see, for example, [\[11,](#page-18-9) p. 396]) that the latter inequality holds for all $z \in \Delta$. Geometrically this fact means that g maps Δ into the sector $\overline{A_{\alpha}}$, where

$$
A_{\alpha} = \left\{ w \in \mathbb{C} : |\arg w| < \frac{\alpha \pi}{2}, \ \alpha \in (0,2] \right\}.
$$

Suppose that there exists $z \in \Delta$ such that $w = g(z) \in \partial A_{\alpha}$. Then by the maximum principle $g \equiv \text{const} = w$ and $f(z) = w\tau(z - \tau)^2$. In this case w must be zero, since otherwise we get contradiction with [\(14\)](#page-8-0). Hence, either $w = 0$ or $g(\Delta) \subset A_{\alpha}$.

If $w = 0$ then $f \equiv 0$ and we are done.

Let now $g(\Delta) \subset A_{\alpha}$. Equality [\(14\)](#page-8-0) implies that

$$
\angle \lim_{z \to \tau} \frac{g(z)}{(z - \tau)^{\alpha}} = -\tau \angle \lim_{z \to \tau} \frac{f(z)}{(z - \tau)^{2 + \alpha}} = 0.
$$

Applying Theorem C we get $q \equiv 0$, hence $f \equiv 0$.

Corollary 4 Let $F \in Hol(\Delta, \mathbb{C})$ be continuous on $\overline{\Delta}$ and satisfy the boundary condition

$$
\operatorname{Re} F(z)\overline{z} \le 1 - |F(z) - z| \cos \frac{\alpha \pi}{2}, \quad z \in \partial \Delta,\tag{15}
$$

for some $\alpha \in (0, 2]$. If there exists $\tau \in \partial \Delta$ such that

$$
F(z) = \tau + (z - \tau) + o(|z - \tau|^{2+\alpha})
$$

when $z \to \tau$, then $F \equiv I$.

3 Commuting semigroups.

Theorem 3 Let f and g be generators of one-parameter commuting semigroups ${F_t}_{t\geq0}$ and ${G_t}_{t\geq0}$, respectively, and $f(\tau)=0$ at some point $\tau\in\Delta$. (i) Let $\tau \in \Delta$. If $f'(\tau) = g'(\tau)$ then $f \equiv g$.

(ii) Let $\tau \in \partial \Delta$. Suppose f and g admit the following representations

$$
f(z) = f'(\tau)(z - \tau) + \ldots + \frac{f^{(m)}(\tau)}{m!}(z - \tau)^m + o(|z - \tau|^m)
$$
 (16)

and

$$
g(z) = g(\tau) + g'(\tau)(z - \tau) + \ldots + \frac{g^{(m)}(\tau)}{m!}(z - \tau)^m + o(|z - \tau|^m) \tag{17}
$$

when $z \to \tau$ along some curve lying in Δ and ending at τ . If $f^{(m)}(\tau) =$ $g^{(m)}(\tau) \neq 0$, then $f \equiv g$.

Remark 2 If $\tau \in \partial \Delta$ is the Denjoy–Wolff point of a semigroup generated by a mapping $h \in \mathcal{G}(\Delta)$, then h admits the expansion

$$
h(z) = h'(\tau)(z - \tau) + o(z - \tau)
$$

when $z \to \tau$ in each non-tangential approach region at τ and $h'(\tau) = \angle \lim_{z \to \tau} h'(z)$. Moreover, in this case $h'(\tau)$ is a non-negative real number which is zero if

Therefore, if f (or g) in Theorem 3 generates a semigroup of hyperbolic type with the Denjoy–Wolff point $\tau \in \partial \Delta$ then the condition $f'(\tau) = g'(\tau)$ is enough to provide that $f \equiv q$.

and only if h generates a semigroup of parabolic type (see $[10]$).

Remark 3 As a matter of fact, if f and g have expansion (16) and (17) when $z \to \tau$ in each non-tangential approach region at $\tau \in \partial \Delta$ up to the third order $m = 3$, such that $f'(\tau) = g'(\tau)$, $f''(\tau) = g''(\tau)$ and $f'''(\tau) = g'''(\tau)$ then $f \equiv q$.

If, in particular, $f^{(i)}(\tau) = g^{(i)}(\tau) = 0$, $i = 1, 2, 3$, then both f and g are equal zero identically by Theorem 1.

Theorem 3 is a consequence of the following more general assertion.

Define two linear semigroups $\{A_t\}_{t\geq 0}$ and $\{B_t\}_{t\geq 0}$ of composition operators on Hol (Δ, \mathbb{C}) by

$$
A_t(h) = h \circ F_t \quad \text{and} \quad B_t(h) = h \circ G_t, \quad t \ge 0. \tag{18}
$$

The operators Γ_f and Γ_g defined by

$$
\Gamma_f(h) = h'f \quad \text{and} \quad \Gamma_g(h) = h'g \tag{19}
$$

are their generators, respectively.

Theorem 4 Let f and $q \in Hol(\Delta, \mathbb{C})$ be generators of one-parameter semigroups ${F_t}_{t\geq0}$ and ${G_t}_{t\geq0}$, respectively. Let A_t and B_t be defined by [\(18\)](#page-10-0). Then the following are equivalent:

(i) $F_t \circ G_s = G_s \circ F_t$, $s, t \geq 0$, i.e., the semigroups $\{F_t\}_{t \geq 0}$ and $\{G_t\}_{t \geq 0}$ are commuting;

(ii) $A_t \circ B_s = B_s \circ A_t$, $s, t \geq 0$, i.e., the linear semigroups $\{A_t\}_{t \geq 0}$ and ${B_t}_{t\geq0}$ are commuting;

(iii) $\Gamma_f \circ \Gamma_g = \Gamma_g \circ \Gamma_f$, i.e., the linear semigroup generators Γ_f and Γ_g are commuting;

(iv) the Lie commutator

$$
[f,g] = f'g - g'f = 0;
$$

(v) $f = \alpha q$ for some $\alpha \in \mathbb{C}$.

Proof. Suppose that $f \neq 0$. First we prove the equivalence of assertions (i) and (v) .

Let (i) holds. If $f(\tau) = 0, \tau \in \Delta$, then τ is a unique common fixed point for the semigroup ${F_t}_{t\geq0}$ generated by f, i.e., $F_t(\tau) = \tau$ for all $t \geq 0$ (see, for example, $[5]$, $[15]$).

If F_t and G_s are commuting for all $s, t \geq 0$, then we have

$$
G_s(\tau) = G_s(F_t(\tau)) = F_t(G_s(\tau)).
$$

Hence, it follows by the uniqueness of the fixed point τ that $G_s(\tau) = \tau$ for all $s \geq 0$, and so $g(\tau) = 0$.

Consider the function $h \in Hol(\Delta, \mathbb{C})$ defined by the differential equation

$$
\mu h(z) = h'(z)f(z). \tag{20}
$$

It is known that if $\mu = f'(\tau)$ then equation [\(20\)](#page-11-0) has a unique solution $h \in$ Hol(Δ , \mathbb{C}) normalized by the condition $h'(\tau) = 1$ (see [\[15\]](#page-18-6)).

In addition, this function h solves Schroeder's functional equation

$$
h(F_t(z)) = e^{-\mu t} h(z).
$$
 (21)

Now, for any $s, t \geq 0$ we get from [\(21\)](#page-11-1)

$$
h(G_s(F_t(z))) = h(F_t(G_s(z))) = e^{-\mu t} h(G_s(z)).
$$

Denote $h_s = h \circ G_s$. Then we have

$$
h_s(F_t(z)) = e^{-\mu t} h_s(z).
$$
 (22)

Differentiating [\(22\)](#page-11-2) at $t = 0^+$ we get

$$
\mu h_s(z) = h'_s(z) f(z). \tag{23}
$$

Comparing [\(20\)](#page-11-0) and [\(23\)](#page-11-3) implies $h_s(z) = \lambda(s)h(z)$ for some $\lambda(s) \in \mathbb{C}$, or

$$
h(G_s(z)) = \lambda(s)h(z). \tag{24}
$$

Since the left-hand side of the latter equality is differentiable in $s \geq 0$, the scalar function $\lambda(s)$ is differentiable too. Differentiating [\(24\)](#page-11-4) at $s = 0^+$ we get

$$
\lambda'(0)h(z) = -h'(z)g(z). \tag{25}
$$

Note that $h(\tau) = 0$ while $h(z) \neq 0$ for all $z \in \Delta$, $z \neq \tau$. In addition, it can be shown (see [\[15\]](#page-18-6)) that h is univalent. Hence, $h'(z) \neq 0$ for all $z \in \Delta$.

Finally, we obtain from [\(20\)](#page-11-0) and [\(25\)](#page-11-5) that

$$
f(z) = \alpha g(z)
$$
, where $\alpha = -\frac{\mu}{\lambda'(0)}$.

Now, let us suppose that f has no null point in Δ . Then the function $p : \Delta \mapsto \mathbb{C}$ given by

$$
p(z) = -\int_{0}^{z} \frac{d\zeta}{f(\zeta)}
$$
\n(26)

is well defined holomorphic function on Δ with $p(0) = 0$.

Recall that the semigroup $\{F_t\}_{t\geq 0}$ generated by f can be defined by the Cauchy problem

$$
\begin{cases}\n\frac{dF_t(z)}{dt} + f(F_t(z)) = 0, \, t \ge 0 \\
F_0(z) = z, \quad z \in \Delta\n\end{cases}
$$
\n(27)

Substituting here $f(z) = -\frac{1}{n^{\prime}i^{\prime}}$ $\frac{1}{p'(z)}$ we obtain

$$
p'(F_t(z)) dF_t(z) = dt.
$$

Integrating the latter equality on the interval $[0, t]$ we get that p is a solution of Abel's functional equation

$$
p(F_t(z)) = p(z) + t.
$$
 (28)

Now, for any fixed $s \geq 0$ we have

$$
p(G_s(F_t(z))) = p(F_t(G_s(z))) = p(G_s(z)) + t.
$$

Once again, setting $p_s = p \circ G_s$, we have

$$
p_s(F_t(z)) = p_s(z) + t.
$$
 (29)

Differentiating [\(29\)](#page-12-0) at $t = 0^+$ we get

$$
p'_s(z) = -\frac{1}{f(z)},
$$
\n(30)

and by [\(26\)](#page-12-1), $p_s(z) = p(z) + \kappa(s)$, $\kappa(s) \in \mathbb{C}$, or

$$
p(G_s(z)) = p(z) + \kappa(s). \tag{31}
$$

Differentiating [\(31\)](#page-13-0) at $s = 0^+$ we obtain the equality

$$
p'(z) = -\frac{\kappa'(0)}{g(z)}.\tag{32}
$$

Comparing [\(30\)](#page-12-2) and [\(32\)](#page-13-1) gives

$$
f = \alpha g \quad \text{with} \quad \alpha = \frac{1}{\kappa'(0)}.
$$
 (33)

Now we prove that $(v) \Rightarrow (i)$. Let $f = \alpha g$ for some $\alpha \in \mathbb{C}$.

First we assume that g has an interior null-point $\tau \in \Delta$. In this case there is a univalent solution of the differential equation

$$
\mu h(z) = h'(z)g(z) \tag{34}
$$

with some $\mu \in \mathbb{C}$, Re $\mu \geq 0$.

Since $f = \alpha g$, we have that h is also a solution of the equation

$$
\nu h(z) = h'(z)f(z), \quad \nu = \alpha \mu. \tag{35}
$$

In turn, equations [\(34\)](#page-13-2) and [\(35\)](#page-13-3) are equivalent to Schroeder's functional equations

$$
h(G_s(z)) = e^{-\mu s} h(z), \quad s \ge 0
$$
\n(36)

and

$$
h(F_t(z)) = e^{-\nu t} h(z), \quad t \ge 0, \quad \nu = \alpha \mu,
$$
\n(37)

respectively, where ${F_t}_{t\geq 0}$ is the semigroup generated by f.

Consequently,

$$
F_t (G_s(z)) = h^{-1} (e^{-\nu t} h (G_s(z))) = h^{-1} (e^{-\nu t} \cdot e^{-\mu s} h(z))
$$

= $h^{-1} (e^{-\mu s} h (F_t(z))) = G_s (F_t(z))$

for all $s, t \geq 0$ and we are done.

Now let us assume that g has a boundary null-point $\tau \in \partial \Delta$ with $g'(\tau) \geq 0$ (see Remark 1 above). In this case for each $c \in \mathbb{C}$, $c \neq 0$, Abel's equations

$$
p\left(G_s(z)\right) = p(z) + cs
$$

and

$$
p(F_t(z)) = p(z) + c\alpha t
$$

have the same solution

$$
p(z) = -c \int_{0}^{z} \frac{d\zeta}{g(\zeta)} = -c\alpha \int_{0}^{z} \frac{d\zeta}{f(\zeta)},
$$

which is univalent on Δ .

Once again we calculate

$$
F_t(G_s(z)) = p^{-1} (p(G_s(z)) + c\alpha t) = p^{-1} (p(z) + c\alpha t + cs)
$$

= $p^{-1} (p(F_t(z)) + cs) = G_s(F_t(z)).$

The implication $(v) \Rightarrow (i)$ is proved.

The equivalence of (i) and (ii) is obvious.

To verify the equivalence of (iii) and (iv) we just calculate:

$$
\Gamma_f(\Gamma_g(h)) = h''gf + h'g'f,
$$

$$
\Gamma_g(\Gamma_f(h)) = h''fg + h'f'g.
$$

Hence, $\Gamma_f \circ \Gamma_g = \Gamma_g \circ \Gamma_f$ if and only if $f'g - g'f = 0$.

Now, it is clear, that (v) implies (iv).

Finally we prove the implication (iv) \Rightarrow (v). Obviously, (iv) implies that if f has no null points in Δ then q also has no null points in Δ and, hence, (v) follows. If $f(\tau) = 0$ for some $\tau \in \Delta$, then also $g(\tau) = 0$, and by [\(9\)](#page-4-0) one can write $f(z) = (z - \tau)p(z)$ and $g(z) = (z - \tau)q(z)$, where p and q do not vanish in Δ . Now it follows that

$$
[f,g] = (z - \tau)[p,q]
$$

Hence, again we have $p = aq$, and hence $f = ag$ for some $a \in \mathbb{C}$, $a \neq 0$.

Proof of Theorem 3. First we note, that by Theorem 4

$$
f = \alpha g, \quad \alpha \in \mathbb{C}.\tag{38}
$$

(i) Let $f'(\tau) = g'(\tau) = 0$. By Proposition 1 f admits representation

$$
f(z) = (z - \tau)(1 - \overline{\tau}z)p(z), \ z \in \Delta,
$$

where $p \in Hol(\Delta, \mathbb{C})$, Re $p(z) \geq 0$.

Since $f'(\tau) = (1 - |\tau|^2)p(\tau) = 0$, we have $p(\tau) = 0$ and it follows from the maximum principle that $p \equiv 0$. Hence, $f \equiv 0$ and by [\(38\)](#page-14-0) also $q \equiv 0$.

Assume now $f'(\tau) = g'(\tau) \neq 0$. Then it follows from [\(38](#page-14-0)) that $\alpha = 1$ and so $f \equiv g$.

(ii) In general, by [\(38\)](#page-14-0) we have $f^{(k)}(\tau) = \alpha g^{(k)}(\tau)$, $0 < k \leq m$. Hence, the condition $f^{(k)}(\tau) = g^{(k)}(\tau) \neq 0$ for some $0 < k \leq m$ implies that $\alpha = 1$ and, consequently, $f \equiv g$. \blacksquare

Let $\mathcal{S}_f = \{F_t\}_{t\geq 0}$ be the semigroup generated by $f \in \mathcal{G}(\Delta)$. The set $\mathcal{Z}(\mathcal{S}_f)$ of all semigroups $\mathcal{S} = \{G_t\}_{t\geq 0}$ such that

$$
F_t \circ G_s = G_s \circ F_t, \quad t, s \ge 0,
$$

is called the **centralizer** of S_f .

It is clear that for each $f \in \mathcal{G}(\Delta)$ the centralizer $\mathcal{Z}(\mathcal{S}_f)$ contains $\mathcal{S}_{\alpha f}$ for all $\alpha \geq 0$.

Therefore we will say that the centralizer of S_f is trivial when the conclusion $S \in \mathcal{Z}(\mathcal{S}_f)$ implies that $S = \mathcal{S}_{\alpha f}$ for some $\alpha \geq 0$.

Proposition 1 Let f be the generator of a semigroup $S_f = \{F_t\}_{t \geq 0}$, and let $\tau \in \partial \Delta$ be the Denjoy–Wolff point of S_f . Then if one of the following conditions holds then the centralizer $\mathcal{Z}(\mathcal{S}_f)$ is trivial:

(i) S_f is a hyperbolic type semigroup $(f'(\tau) > 0)$ which is not a group;

(ii) f admits the expansion

$$
f(z) = a(z - \tau)^3 + o((z - \tau)^3)
$$
 with $a \neq 0$

when $z \to \tau$ in each non-tangential approach region at τ .

The first statement is based on the following simple lemma.

Lemma 2 Let f and g be generators of two nontrivial (neither f nor g are identically zero) commuting semigroups $S_f = \{F_t\}_{t\geq 0}$ and $S_g = \{G_t\}_{t\geq 0}$, respectively. Then S_f is of hyperbolic type if and only if S_g is. In this case $f = \alpha g$ with real α . Moreover, $\alpha < 0$ implies that S_f and S_g are both groups of hyperbolic automorphisms having 'opposite' fixed points, i.e., the attractive point for S_f is the repelling point for S_g and conversely.

Proof. Since S_f and S_g are commuting then by Theorem 4 there exists $\alpha \in \mathbb{C}$ such that $f = \alpha g$. In our settings α is not zero. If τ is the Denjoy-Wolff point of S_f then $f(\tau) = 0$ and therefore also $g(\tau) = 0$. Now since $f'(\tau) > 0$ then $g'(\tau) = \frac{1}{\alpha} f'(\tau)$ exists finitely and it must be a real number by Corollary 2. So must be α .

Now let us assume that α is negative. Then $g'(\tau) = \frac{1}{\alpha} f'(\tau) < 0$. Hence the semigroup S_g generated by g must have the Denjoy–Wolff point $\sigma \in \overline{\Delta}$ different from τ .

It is clear that σ can not be inside Δ since otherwise it must be a common fixed point of both semigroups S_f and S_g because of the commuting property.

So, $\sigma \in \partial \Delta$ and $g'(\sigma) \ge 0$ (see [\[10\]](#page-18-8)), then $f(\sigma) = 0$ and $f'(\sigma) \le 0$. It follows by a result in [\[16\]](#page-19-1) that

$$
0 < f'(\tau) \le -f'(\sigma) \tag{39}
$$

and the equality is possible if and only if f is the generator of a group of hyperbolic automorphisms. By the same theorem we have the reversed inequality for g

$$
0 \le g'(\sigma) \le -g'(\tau)
$$

that means

$$
0 \le \frac{1}{\alpha} f'(\sigma) \le -\frac{1}{\alpha} f'(\tau).
$$

Comparing this inequality with [\(39\)](#page-16-0) gives us that $f'(\tau) = -f'(\sigma) > 0$ and $g'(\tau) = -g'(\sigma) < 0$ which means that both f and g generate groups of hyperbolic automorphisms with opposite fixed points. \blacksquare

Remark. The last assertion of this lemma follows also by a result of Behan (see [\[4\]](#page-18-10)). Indeed, let $\alpha < 0$. Then the equality $f(z) = \alpha g(z)$ implies that $g'(\tau)$ exists and is a real negative number. So, the Denjoy–Wolff point τ of the semigroup S_f can not be the Denjoy–Wollf point of the semigroup S_g . Hence by [\[4\]](#page-18-10) we conclude that S_f and S_g are groups of hyperbolic automorphisms. Proof of Proposition 1. The statement (i) is a direct consequence of the previous lemma. To prove the second statement we note that by Theorem 1 the number $a\tau^2$ is a non-negative real number. On the other hand, since S_f and S_q commute by Theorem 4 there is a number $\alpha \in \mathbb{C}$ such that $f = \alpha g$.

Therefore, since $\alpha \neq 0$ also g admits the expansion

$$
g(z) = \frac{a}{\alpha} (z - \tau)^3 + o((z - \tau)^3)
$$

and again by Theorem 1 we have that also $\frac{a}{\alpha} \tau^2 \geq 0$. This implies that α is a nonnegative real number.

A natural question which arises in the context of the above theorem is: • If two elements F_{t_0} and G_{s_0} of semigroups $S_f = \{F_t\}_{t\geq 0}$ and $S_g = \{G_t\}_{t\geq 0}$ commute for some positive t_0 and s_0 , do these semigroup S_f and S_g commute in the sense:

$$
F_t \circ G_s = G_s \circ F_t
$$

for each pair $t, s \geq 0$.

The answer is immediately affirmative due to a more general result of C.C. Cowen ([\[8\]](#page-18-11)), Corollary) if neither F_{t_0} nor G_{s_0} , respectively, are of parabolic type.

The situation becomes more complicated if F_{t_0} , respectively G_{s_0} , are parabolic.

Example 4.4 in [\[8\]](#page-18-11) shows that there is a triple of such mappings F , G_1 and G_2 in Hol(Δ) for which G_1 and G_2 commute with F, but they do not commute each other.

Nevertheless, under some additional requirements on smoothness at the boundary Denjoy-Wolff point repeating the arguments using in the proof of Theorem 1.2 in [\[17\]](#page-19-0) one can give an affirmative answer the above question. Namely,

• Let F_{t_0} and G_{s_0} be two commuting elements of semigroups S_f and S_g , respectively, $t_0, s_0 > 0$, and let F_{t_0} is of parabolic type with a Denjoy–Wolff point $\tau \in \partial \Delta$. If both F_{t_0} and G_{s_0} belong to the class $C^2(\tau)$ and $F''_{t_0}(\tau)$ as well as $G''_{t_0}(\tau)$ do not vanish, then $\tilde{f} = ag$ for some $a \in \mathbb{C}$, i.e., the semigroups S_f and S_g commute:

$$
F_t \circ G_s = G_s \circ F_t
$$

for all $t, s \geq 0$.

References

- [1] M. Abate, Converging semigroups of holomorphic maps, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) 82 (1988), 223–227.
- [2] D. Aharonov, M. Elin, S. Reich and D. Shoikhet, Parametric representations of semi-complete vector fields on the open unit balls in \mathbb{C}^n and in Hilbert space, Atti Acad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei(9)Mat. Appl. **10** (1999), 229–253.
- [3] D. Aharonov, S. Reich and D. Shoikhet, Flow invariance conditions for holomorphic mappings in Banach spaces, Math. Proc. R. Ir. Acad. 99A (1999), 93–104.
- [4] D. Behan, Commuting analytic functions without fixed points, Proc. Amer. Math. Soc. **37** (1973), 114–120.
- [5] E. Berkson, H. Porta, Semigroups of analytic functions and composition operators., *Michigan Math. J.* **25** (1978), 101–115.
- [6] F. Bracci, R. Tauraso, and F. Vlacci, Identity principles for commuting holomorphic self-maps of the unit disk, J. Math. Anal. Appl. 270 (2002), 451–473.
- [7] M. Burns and S. G. Krantz, Rigidity of Holomorphic Mappings and a New Schwarz Lemma at the Boundary, J. Amer. Math. Soc. 7 (1994), 661–676.
- [8] C. C. Cowen, Commuting analytic functions, Trans. Amer. Math. Soc. 283 (1984), 685–695.
- [9] C. C. Cowen and B. D. MacCluer, Schroeder's equation in several variables, *Taiwanese J.of Math.* $7(1)$ (2003), 129–154.
- [10] M. Elin and D. Shoikhet, Dynamic Extension of the Julia-Wolff-Carathéodory Theorem, Dynam. Systems Appl. 10 (2001), 421–438.
- [11] T. W. Gamelin, Complex Analysis, Springer, 2000.
- [12] L. A. Harris, S. Reich and D. Shoikhet, Dissipative Holomorphic Functions, Bloch Radii, and the Schwarz Lemma, Journal D'Analyse Mathematique 82 (2000), 221–232.
- [13] S. Migliorini and F. Vlacci, A new rigidity result for holomorphic maps, *Indag. Mathem., N.S.* $13(4)$ (2002), 537-549.
- [14] S. Reich and D. Shoikhet, Metric domains, holomorphic mappings and nonlinear semigroups, $Abstr.$ Appl. Anal. 3 (1998), 203–228.
- [15] D. Shoikhet, Semigroups in Geometrical Function Theory, Kluwer Acad. Publ., 2001.
- [16] D. Shoikhet, Representations of holomorphic generators and distortion theorems for spirallike functions with respect to a boundary point, Int. J. Pure Appl. Math. 5 (2003), 335–361.
- [17] R. Tauraso, Commuting holomorphic maps of the unit disk, Ergod. Theory Dynam. Systems 24 (2004), 945–953.
- [18] K. Yosida, Functional Analysis, Springer, 1980.