

MANIN'S CONJECTURE ON RATIONAL POINTS OF BOUNDED HEIGHT AND ADELIC MIXING

ALEX GORODNIK, FRANÇOIS MAUCOURANT, HEE OH

Dedicated to Prof. Gregory Margulis on the occasion of his sixtieth birthday

ABSTRACT. Let K be a number field. We compute the asymptotics of the number of K -rational points of bounded height on a connected adjoint semisimple K -group \mathbf{G} for any given irreducible representation. This proves Manin's conjecture for the wonderful compactification X of \mathbf{G} . We also determine the explicit asymptotic distribution of the rational points $\mathbf{G}(K)$ on $X(\mathbb{A})$, which verifies the prediction made by Peyre and Batyrev-Tschinkel. Our approach is based on the mixing property of $L^2(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$ which we prove with a rate of convergence.

CONTENTS

1. Introduction	1
2. Adelic Mixing	11
2.1. Definition and properties of $\xi_{\mathbf{G}}$	11
2.2. Uniform bound for matrix coefficients of $\mathbf{G}(\mathbb{A})$	14
2.3. Automorphic bound for $\mathbf{G}(\mathbb{A})$	16
3. Volume asymptotics	20
4. Equidistribution for saturated cases	25
5. Arithmetic fibrations	32
References	37

1. INTRODUCTION

Let K be a number field and R the set of all (inequivalent) normalized absolute values $x \mapsto |x|_v$ of K . We denote by K_v the completion of K with respect to v . The height $H(x)$ of a point $x = (x_0 : x_1 : \cdots : x_n) \in \mathbb{P}^n(K)$ is given by

$$H(x) := \prod_{v \in R} H_v(x)$$

The first and the third authors are partially supported by DMS-0400631 and DMS-0333397 respectively. The second author would like to thank Caltech for the hospitality during his visit where the work was first conceived.

where $H_v(x) = \max_i |x_i|_v$ for each $v \in R$. By the product formula, H is a well defined function on $\mathbb{P}^n(K)$. For example, when $K = \mathbb{Q}$, we have

$$H(x) = \max_i |x_i|$$

where (x_0, \dots, x_n) is a primitive integral vector.

More generally one can consider a height function which differs from above by changing the local height H_v by another norm on K_v^{n+1} for finitely many places v .

It is easy to see that for any $T > 0$,

$$N(T) := \#\{x \in \mathbb{P}^n(K) : H(x) < T\} < \infty.$$

Schanuel [Sc] computed in 1964 that

$$N(T) \sim c \cdot T^{n+1} \quad \text{as } T \rightarrow \infty$$

for some explicit constant $c = c(H) > 0$.

A fundamental problem in modern algebro-arithmetic geometry is to describe the set of rational points of projective varieties in terms of their geometric invariants. One of the main conjectures in this area was made by Manin in [BM] in the late eighties. It describes the asymptotics of the numbers of rational points on projective varieties with ample anti-canonical classes (such varieties are called Fano varieties). Manin's conjecture has been proved for flag varieties ([FMT], [Pe1]), toric varieties [BT1-2], horospherical varieties [ST], equivariant compactifications of unipotent groups (see [CT2], [ST1], [ST2]), etc. In this paper, we settle Manin's conjecture for the wonderful compactification of a general connected semisimple adjoint group \mathbf{G} defined over a number field.

Let \mathbf{G} be a connected adjoint semisimple group over K . Let $\iota : \mathbf{G} \rightarrow \mathrm{GL}_N$ be a faithful representation of \mathbf{G} defined over K . This defines a projective embedding over K :

$$\bar{\iota} : \mathbf{G} \rightarrow \mathbb{P}(\mathrm{M}_N) = \mathbb{P}^{N^2-1}$$

where M_N denotes the space of matrices of order N . Fixing a height function $H = \prod_{v \in R} H_v$ on $\mathbb{P}^{N^2-1}(K)$, we obtain a height function H_ι on $\mathbf{G}(K)$ via $\bar{\iota}$:

$$(1.1) \quad H_\iota(g) := H(\bar{\iota}(g)) = \prod_{v \in R} H_v(\iota(g)).$$

Set

$$N(H_\iota, T) := \#\{g \in \mathbf{G}(K) : H_\iota(g) < T\}.$$

Theorem 1.2. *Let \mathbf{G} be a product of connected adjoint absolutely simple groups over K and $\iota : \mathbf{G} \rightarrow \mathrm{GL}_N$ a faithful absolutely irreducible representation defined over K . Then there exist $a_\iota \in \mathbb{Q}^+$, $b_\iota \in \mathbb{N}$ and $c = c(H_\iota) > 0$ such that for some $\delta > 0$,*

$$N(H_\iota, T) = c \cdot T^{a_\iota} (\log T)^{b_\iota-1} \cdot (1 + O((\log T)^{-\delta}))$$

for all sufficiently large T .

Remark

- (1) When \mathbf{G} is absolutely simple or, more generally, when H_ι is the product of height functions of the simple factors of \mathbf{G} , we can improve the rate of convergence in the above: for some $\delta > 0$,

$$N(H_\iota, T) = c \cdot T^{a_\iota} P(\log T) \cdot (1 + O(T^{-\delta}))$$

where $P(x)$ is a monic polynomial of degree $b_\iota - 1$.

- (2) We note that for any connected semisimple adjoint algebraic group \mathbf{G} defined over K , there exists a finite field extension, say, F , of K such that \mathbf{G} is a direct product of connected adjoint absolutely simple groups defined over F . Therefore Theorem 1.2 holds for any connected semisimple adjoint group over K by passing to a finite field extension.

The constants a_ι and b_ι can be defined explicitly by combinatorial data coming from the root system of \mathbf{G} and the highest weight of ι . Choose a set Δ of simple roots in the root system $\Phi(\mathbf{G}, \mathbf{T})$ of \mathbf{G} with respect to a maximal torus \mathbf{T} defined over K containing a maximal K -split torus. Denote by 2ρ the sum of all positive roots in $\Phi(\mathbf{G}, \mathbf{T})$, and by χ the highest weight of ι . Define $u_\alpha, m_\alpha \in \mathbb{N}$, $\alpha \in \Delta$, by

$$2\rho = \sum_{\alpha \in \Delta} u_\alpha \alpha \quad \text{and} \quad \chi = \sum_{\alpha \in \Delta} m_\alpha \alpha.$$

The fact that $m_\alpha \in \mathbb{N}$ follows since \mathbf{G} is of adjoint type. Consider the twisted action of the Galois group $\Gamma_K := \text{Gal}(\bar{K}/K)$ on Δ . For instance, if the K -form of \mathbf{G} is inner, this action is just trivial. Then

$$(1.3) \quad a_\iota = \max_{\alpha \in \Delta} \frac{u_\alpha + 1}{m_\alpha} \quad \text{and} \quad b_\iota = \#\{\Gamma_K \cdot \alpha : \frac{u_\alpha + 1}{m_\alpha} = a_\iota\}.$$

Note that the exponent a_ι is independent of the field K , and b_ι depends only on the quasi-split K -form of \mathbf{G} . Therefore, by passing to a finite field extension containing the splitting field of \mathbf{G} , b_ι also becomes independent of K .

Refining Manin's conjecture mentioned above, Peyre made a conjecture on the asymptotic distribution of rational points of Fano varieties [Pe1]. We verify Peyre's conjecture for "saturated" projective embeddings of semisimple adjoint groups. We call a representation $\iota : \mathbf{G} \rightarrow \text{GL}_N$ *saturated* if the set

$$(1.4) \quad \{\alpha \in \Delta : \frac{u_\alpha + 1}{m_\alpha} = a_\iota\}$$

is not contained in the root system of a proper normal subgroup of \mathbf{G} . In particular, if \mathbf{G} is absolutely simple, any representation of \mathbf{G} is saturated.

For each $v \in R$, let $X_{\iota, v}$ denote the closure of $\bar{\iota}(\mathbf{G}(K_v))$ in $\mathbb{P}^{N^2-1}(K_v)$, that is, the Satake compactification of $\mathbf{G}(K_v)$ relative to ι , and consider the probability measure

$\mu_{\iota,v}$ on $X_{\iota,v}$ defined by

$$\mu_{\iota,v} := m_{\iota,v}^{-1} \cdot H_v(\iota(g_v))^{-a_\iota} dg_v \quad \text{where} \quad m_{\iota,v} := \int_{\mathbf{G}(K_v)} H_v(\iota(g_v))^{-a_\iota} dg_v$$

for a Haar measure dg_v on $\mathbf{G}(K_v)$. This is well defined since $m_{\iota,v} < \infty$ (see Lemma 3.3).

Now set

$$X_\iota := \prod_{v \in R} X_{\iota,v} \quad \text{and} \quad \mu_\iota := \prod_{v \in R} \mu_{\iota,v}.$$

Then X_ι is a compact space in which the product $\prod_v \mathbf{G}(K_v)$ is embedded densely and μ_ι defines a probability measure on X_ι .

For a topological space X , we denote by $C(X)$ the space of continuous functions on X and by $C_c(X)$ the subset of $C(X)$ consisting of functions of compact support. We may consider $\mathbf{G}(K)$ as a subset of X_ι via the diagonal embedding $g \mapsto (\bar{\iota}(g)) \in X_\iota$. Noting that X_ι is a compactification of $\mathbf{G}(\mathbb{A})$, the following theorem shows that the asymptotic distribution of $\mathbf{G}(K)$ in X_ι is precisely given by the measure μ_ι :

Theorem 1.5. *Let \mathbf{G} be a product of connected adjoint absolutely simple groups over K and $\iota : \mathbf{G} \rightarrow \mathrm{GL}_N$ a faithful absolutely irreducible saturated representation defined over K . Then for any $f \in C(X_\iota)$,*

$$\lim_{T \rightarrow \infty} \frac{1}{N(H_\iota, T)} \sum_{g \in \mathbf{G}(K) : H_\iota(g) < T} f(g) = \int_{X_\iota} f d\mu_\iota.$$

Since $\mu_\iota|_{\prod_v \mathbf{G}(K_v)} = \mu_\iota$, the above theorem implies that the rational points $\mathbf{G}(K)$ do not escape to the boundary of the compactification $X(\mathbb{A})$. It is interesting to compare this result with the distribution of the integral points $\mathbf{G}(\mathbb{Z})$ of bounded height in the Satake compactification of $\mathbf{G}(\mathbb{R})$ where the limiting distribution is all supported on the boundary (see [Mau] or [GOS] for more details).

Note that if ι is not saturated, Theorem 1.2 implies that a positive proportion of $\mathbf{G}(K)$ lies on a proper subgroup of \mathbf{G} . Hence one cannot expect Theorem 1.5 to hold for non-saturated cases.

It follows from the weak approximation property that for any finite subset S of R , $\mathbf{G}(K)$ is dense in $\mathbf{G}_S := \prod_{v \in S} \mathbf{G}(K_v)$ under the diagonal embedding. The following is an immediate corollary of Theorem 1.5:

Corollary 1.6. *Let \mathbf{G} and ι be as in Theorem 1.5 and S a finite subset of R . Then for any $f \in C_c(\mathbf{G}_S)$,*

$$\lim_{T \rightarrow \infty} \frac{1}{N(H_\iota, T)} \sum_{g \in \mathbf{G}(K) : H_\iota(g) < T} f(g) = m_{\iota,S}^{-1} \int_{\mathbf{G}_S} f(g) H_{\iota,S}(g)^{-a_\iota} d\tau_S(g)$$

where $H_{\iota,S} := \prod_{v \in S} H_v \circ \iota$, $\tau_S := \prod_{v \in S} dg_v$ and $m_{\iota,S} := \int_{\mathbf{G}_S} H_{\iota,S}^{-a_\iota}(g) d\tau_S(g)$.

We note in particular that, with respect to the height H_ι , $\mathbf{G}(K)$ is equidistributed on \mathbf{G}_S relative to the measure $m_{\iota,S}^{-1} H_{\iota,S}^{-a_\iota} d\tau_S$, which is not an invariant measure, unless \mathbf{G}_S is compact.

We also state versions of Theorems 1.2 and 1.5 in terms of arithmetic geometry. Let X be a smooth projective variety defined over K . For every line bundle on X over K , there exists an associated height function on $X(K)$ via Weil's height machine ([Si, Theorem B. 3.2]). For example, if L is a very ample line bundle of X with a K -embedding $\psi_L : X \rightarrow \mathbb{P}^N$, then a height function $H_{\mathcal{L}}$ on $X(K)$ is defined as

$$H_{\mathcal{L}} := H \circ \psi_L$$

where H is a height function on $\mathbb{P}^N(K)$ defined as before. For any ample line bundle L of X , mL is very ample for some $m \in \mathbb{N}$. Then a height function $H_{\mathcal{L}}$ is of the form $H_{\mathcal{L}'}^{1/m}$ where $H_{\mathcal{L}'}$ is a height function associated to $L' := mL$.

We call a pair $\mathcal{L} = (L, H_{\mathcal{L}})$ a metrized line bundle. Due to the freedom of choosing a height function H on $\mathbb{P}^N(K)$, $H_{\mathcal{L}}$ is not uniquely determined and this is why we use the subscript \mathcal{L} rather than L .

Let X denote the projective K -variety, which is the wonderful compactification of \mathbf{G} constructed by De Concini and Procesi [DP] and by De Concini and Springer [DS]. For instance, X can be taken to be the Zariski closure of the image of \mathbf{G} in $\mathbb{P}(\mathbb{M}_N)$ under an irreducible faithful representation $\mathbf{G} \rightarrow \mathrm{GL}_N$ whose highest weight is regular. A dominant weight χ is called regular if $\chi = \sum_{\alpha \in \Delta} m_\alpha \omega_\alpha$ with all $m_\alpha > 0$ where $\{\omega_\alpha : \alpha \in \Delta\}$ is the set of fundamental weights.

The Picard group $\mathrm{Pic}(X)$ is freely generated by the line bundles corresponding to the orbits of the fundamental weights under the twisted Galois action. The closed cone $\Lambda_{\mathrm{eff}}(X)$ of the effective line bundles is the positive cone spanned by $D_{\Gamma_K \cdot \alpha}$, $\alpha \in \Delta$, i.e.,

$$\Lambda_{\mathrm{eff}}(X) = \bigoplus \mathbb{R}_{\geq 0} [D_{\Gamma_K \cdot \alpha}]$$

where the sum is taken over the Γ_K -orbits $\Gamma_K \cdot \alpha$ in the set $\{\alpha \in \Delta\}$ of simple roots and $D_{\Gamma_K \cdot \alpha} := \sum_{\beta \in \Gamma_K \cdot \alpha} D_\beta$, and the anticanonical class $[-K_X]$ corresponds to $2\rho + \sum_{\alpha \in \Delta} \alpha$. Moreover any ample line bundle class $[L]$ of X over K corresponds to a regular dominant weight in such a way that if $[L'] := m[L]$ corresponds to $\chi \in X^*(\mathbf{T})$ for $m \in \mathbb{N}$, the restriction of $H_{\mathcal{L}'}$ to $\mathbf{G}(K)$ coincides with a height function H_ι with respect to the irreducible representation ι defined over K with the highest weight χ (cf. [STT2, Proposition 6.3]). We call an ample line bundle L *saturated* if the representation defined by the corresponding dominant weight is saturated. We refer to [BK, Ch 6] for a more detailed account on the wonderful compactification.

The notion of a saturated line bundle is related to the notion of a strongly saturated metrized line bundle \mathcal{L} which was introduced by Batyrev and Tschinkel in [BT3] in order to state a refined version of Manin's conjecture. A metrized line bundle \mathcal{L} on

X is called *strongly saturated* if for any Zariski open dense subset U of X , one has

$$\lim_{T \rightarrow \infty} \frac{\#\{x \in U(K) : H_{\mathcal{L}}(x) < T\}}{\#\{x \in X(K) : H_{\mathcal{L}}(x) < T\}} = 1.$$

If \mathcal{L} is strongly saturated, then L is saturated. It is also clear from the definition (1.4) that the anticanonical line bundle $-K_X$ is always saturated.

Consider the compact space $X(\mathbb{A}) = \prod_{v \in R} X(K_v)$. Peyre defined the Tamagawa measure $\tau_{\mathcal{L}}$ on $X(\mathbb{A}) = \prod_{v \in R} X(K_v)$ associated with the anti-canonical metrized line bundle $-\mathcal{K}_X = (-K_X, H_{-\mathcal{K}_X})$:

$$\tau_{-\mathcal{K}_X} = \lim_{s \rightarrow 1} (s-1)^{\text{rank}(\text{Pic}(X))} L_{S_0}(s, \text{Pic}(X)) \cdot \frac{1}{\sqrt{d_K}^{\dim(X)}} \cdot \prod_v \lambda_v^{-1} \cdot H_{-\mathcal{K}_X, v}(g_v)^{-1} dg_v$$

where $S_0 \subset R$ is a finite subset of places of bad reduction, $\lambda_v = L_v(1, \text{Pic}(X))$ for all $v \in R - S_0$ and 1 otherwise, d_K is the discriminant of K and dg_v is a Haar measure on $\mathbf{G}(K_v)$. We refer to [BT3] for the definition of a Tamagawa measure associated to a general ample line bundle. In particular, $\tau_{\mathcal{L}}$ is of the form $d_L \cdot \prod_v \lambda_v^{-1} \cdot H_v(g_v)^{-a_L} dg_v$ for some constant $d_L > 0$.

The following theorem settles Manin's conjecture (and its refinements due to Peyre, Batyrev and Tschinkel) for the wonderful compactifications X of semisimple adjoint groups. For an metrized ample line bundle $\mathcal{L} = (L, H_{\mathcal{L}})$ on X and a subset U of X , set

$$N_U(\mathcal{L}, T) := \#\{g \in X(K) \cap U : H_{\mathcal{L}}(g) < T\}.$$

Theorem 1.7. *Let X be the wonderful compactification of a group \mathbf{G} which is a product of connected adjoint absolutely simple groups defined over K , and $\mathcal{L} = (L, H_{\mathcal{L}})$ an metrized ample line bundle on X . Then for some $\delta > 0$,*

$$N_{\mathbf{G}}(\mathcal{L}, T) = c_{\mathcal{L}} \cdot T^{a_L} (\log T)^{b_L - 1} (1 + O((\log T)^{-\delta}))$$

for all sufficiently large T . Here

- $a_L = \inf\{a \in \mathbb{Q}^+ : a[L] + [K_X] \in \Lambda_{\text{eff}}(X)\}$, i.e., the Nevanlinna invariant of L ;
- b_L is the dimension of the minimal face of $\Lambda_{\text{eff}}(X)$ containing $a_L[L] + [K_X]$;
- $c_{\mathcal{L}} > 0$, and $c_{\mathcal{L}} = c_L \cdot \tau_{\mathcal{L}}(\mathbf{G}(\mathbb{A}))$, if L is saturated.

In particular, for a metrization $-\mathcal{K}_X = (-K_X, H_{-\mathcal{K}_X})$ of the anti-canonical line bundle, we have

$$N_{\mathbf{G}}(-\mathcal{K}_X, T) = c_{-K_X} \cdot \tau_{-\mathcal{K}_X}(\mathbf{G}(\mathbb{A})) \cdot T (\log T)^{\text{rank}(\text{Pic}(X)) - 1} (1 + O((\log T)^{-\delta})).$$

For a non-saturated line bundle L , the equality $c_{\mathcal{L}} = c_L \cdot \tau_{\mathcal{L}}(\mathbf{G}(\mathbb{A}))$ fails in general, but the variety X has an *asymptotic arithmetic \mathcal{L} -fibration* in the sense of [BT3]. More precisely, there exist a connected semisimple K -group \mathbf{H} and a surjective K -homomorphism $\pi : \mathbf{G} \rightarrow \mathbf{H}$ such that for each $x \in \mathbf{H}(K)$,

$$N_{\pi^{-1}(x)}(\mathcal{L}, T) \sim c_L \cdot \tau_{\mathcal{L}}(\pi^{-1}(x)(\mathbb{A})) \cdot T^{a_L} (\log T)^{b_L - 1}; \quad \text{and}$$

$$N_{\mathbf{G}}(\mathcal{L}, T) \sim c_L \cdot \sum_{x \in \mathbf{H}(K)} \tau_{\mathcal{L}}(\pi^{-1}(x)(\mathbb{A})) \cdot T^{a_L} (\log T)^{b_L - 1} \quad \text{as } T \rightarrow \infty$$

with $\sum_{x \in \mathbf{H}(K)} \tau_{\mathcal{L}}(\pi^{-1}(x)(\mathbb{A})) < \infty$.

Theorem 1.7 is independently proved by Shalika, Takloo-Bighash and Tschinkel in their recent preprint [STT2]. The methods in [STT2] are based on the study of analytic properties of the height zeta function $\mathcal{Z}(s) := \sum_{\gamma \in \mathbf{G}(K)} H_{\iota}(\gamma)^{-s}$ using the spectral decomposition of $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. Our approach is different and allows us to avoid technical difficulties, such as, dealing with the continuous spectrum. To extend Theorem 1.7 to an arbitrary bi-equivariant compactification, say X , of \mathbf{G} using our methods, we only need to verify that the height functions coming from ample line bundles of X satisfy the assumptions stated in Proposition 4.14.

We also state the analogue of Theorem 1.5 in this setup.

Theorem 1.8. *With the same notation as Theorem 1.7, suppose also that L is saturated. Then for any $f \in C(X(\mathbb{A}))$,*

$$\lim_{T \rightarrow \infty} \frac{1}{N_{\mathbf{G}}(\mathcal{L}, T)} \sum_{g \in \mathbf{G}(K): H_{\mathcal{L}}(g) < T} f(g) = \frac{1}{\tau_{\mathcal{L}}(X(\mathbb{A}))} \int_{X(\mathbb{A})} f d\tau_{\mathcal{L}}.$$

To explain our strategy in counting K -rational points of \mathbf{G} , we first recall the analogous results in counting lattice points in a simple real Lie group. Let $G \subset \mathrm{GL}_N$ be a connected non-compact simple real Lie group and Γ be a lattice in G , i.e., a discrete subgroup of finite co-volume. Fixing a norm $\|\cdot\|$ on $M_N(\mathbb{R})$, set $B_T := \{g \in G : \|g\| \leq T\}$. By Duke-Rudnick-Sarnak [DRS] and Eskin-McMullen [EM], it is well known that

$$(1.9) \quad \#\Gamma \cap B_T \sim \int_{B_T} dg \quad \text{as } T \rightarrow \infty,$$

where dg is the Haar measure on G such that $\int_{\Gamma \backslash G} dg = 1$. In particular, the key ingredient in the proof in [EM] is the mixing property of $L^2(\Gamma \backslash G)$.

Coming back to the question of counting rational points $\mathbf{G}(K)$, we note that $\mathbf{G}(K)$ can indeed be embedded as a lattice in the adèle group $\mathbf{G}(\mathbb{A})$ under the diagonal map. Recall that the adèle group $\mathbf{G}(\mathbb{A})$ is the restricted topological product of $\mathbf{G}(K_v)$'s with respect to a family of compact open subgroups, say U_v , of $\mathbf{G}(K_v)$, $v \in R_f$, where R_f is the set of all non-archimedean absolute values on K (cf. [PR], [We]). Choosing a smooth model of \mathbf{G} over $\mathcal{O}[k^{-1}]$ for the ring \mathcal{O} of integers of K and for some $k \in \mathbb{Z}$, we have $U_v = \mathbf{G}(\mathcal{O}_v)$ for almost all $v \in R_f$ where \mathcal{O}_v is the valuation ring of K_v . Moreover the height function $H_{\iota} = \prod_{v \in R} H_v \circ \iota$ on $\mathbf{G}(K)$ defined in (1.1) extends to $\mathbf{G}(\mathbb{A})$ by

$$(1.10) \quad H_{\iota}(g) := \prod_{v \in R} H_v(\iota(g_v)) \quad \text{for any } g = (g_v)_v \in \mathbf{G}(\mathbb{A}).$$

Since $U_v = \mathbf{G}(\mathcal{O}_v)$ for almost all $v \in R_f$, $H_v(\iota(g_v)) = 1$ for almost all v , and hence H_ι is well defined.

Setting

$$B_T := \{g \in \mathbf{G}(\mathbb{A}) : H_\iota(g) < T\},$$

note that B_T is a relatively compact subset of $\mathbf{G}(\mathbb{A})$ (Lemma 3.6) and that

$$N(H_\iota, T) = \#\mathbf{G}(K) \cap B_T.$$

In view of (1.9), one naturally asks whether the following holds:

$$(1.11) \quad \#\mathbf{G}(K) \cap B_T \sim \tau(B_T) \quad \text{as } T \rightarrow \infty,$$

where τ is the Haar measure on $\mathbf{G}(\mathbb{A})$ such that $\tau(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A})) = 1$.

It turns out that the group $\mathbf{G}(\mathbb{A})$ is too big for (1.11) to hold in general, due to the presence of non-trivial characters in $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. To define a right group for (1.11), set

$$W_f := \{w \in \mathbf{G}(\mathbb{A}_f) : H_\iota(wg) = H_\iota(gw) = H_\iota(g) \text{ for all } g \in \mathbf{G}(\mathbb{A})\},$$

where $\mathbf{G}(\mathbb{A}_f)$ is the subgroup of finite adeles. It easily follows from the definition of H_ι that W_f is a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$. Denoting by Λ^{W_f} the set of all W_f -invariant automorphic characters appearing in $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$, set

$$\mathbf{G}_{H_\iota} := \ker(\Lambda^{W_f}) = \cap \{\ker \chi \subset \mathbf{G}(\mathbb{A}) : \chi \in \Lambda^{W_f}\}.$$

The subgroup \mathbf{G}_{H_ι} is a finite index normal subgroup of $\mathbf{G}(\mathbb{A})$ which clearly contains $\mathbf{G}(K)$ (see Lemma 3.1).

Theorem 1.12. *Let \mathbf{G} be a product of connected adjoint absolutely simple group defined over K and $\iota : \mathbf{G} \rightarrow \mathrm{GL}_N$ be a faithful absolutely irreducible saturated representation defined over K . Then*

$$\#\mathbf{G}(K) \cap B_T \sim \tau(B_T \cap \mathbf{G}_{H_\iota}) \quad \text{as } T \rightarrow \infty,$$

where τ is the Haar measure on \mathbf{G}_{H_ι} normalized so that $\tau(\mathbf{G}(K) \backslash \mathbf{G}_{H_\iota}) = 1$.

We remark that one cannot in general replace \mathbf{G}_{H_ι} by $\mathbf{G}(\mathbb{A})$ in However in the case when \mathbf{G} is of non-compact type (i.e., the \mathbb{Q} -group $\mathrm{Res}_{K/\mathbb{Q}} \mathbf{G}$ has no \mathbb{Q} -simple factor which is \mathbb{R} -anisotropic), there exists a height function H_ι on $\mathbf{G}(\mathbb{A})$ such that $\mathbf{G}_{H_\iota} = \mathbf{G}(\mathbb{A})$ for any ι .

As in the proof of Eskin-McMullen of (1.9), our key ingredient in proving Theorem 1.12 is the mixing theorem on $L^2(\mathbf{G}(K) \backslash \mathbf{G}_{H_\iota})$. We remark that the idea of using mixing properties of actions in counting problems goes back to Margulis thesis in 1970 [Mar].

Let $L_{00}^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$ denote the orthogonal complement to the direct sum of all one-dimensional representations occurring in $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. In the case when \mathbf{G} is simply connected, $L_{00}^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$ coincides with the orthogonal complement $L_0^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$ to the constant functions. The terminology that a sequence g_i tends

to infinity as $i \rightarrow \infty$ in $\mathbf{G}(\mathbb{A})$ means that for any compact subset Ω in $\mathbf{G}(\mathbb{A})$, $g_i \notin \Omega$ for all sufficiently large i .

Theorem 1.13 (Adelic Mixing). *Let \mathbf{G} be a connected absolutely almost simple K -group. Then for any $f, h \in L^2_{00}(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$,*

$$\langle f, g.h \rangle \rightarrow 0$$

as $g \in \mathbf{G}(\mathbb{A})$ tends to infinity.

In particular, if f and h are W_f -invariant functions of $L^2(\mathbf{G}(K) \backslash \mathbf{G}_{H_L})$, then as $g \rightarrow \infty$,

$$\langle f, g.h \rangle \rightarrow \int f d\tau \cdot \int h d\tau$$

where τ is the normalized Haar measure on \mathbf{G}_{H_L} .

In fact we prove the above theorem 1.13 in a much stronger form by giving a rate of convergence (see Theorem 2.7 and 2.16). In particular we obtain the following result which is of independent interest. Set $\mathbf{G}_\infty := \prod_{v \in R_\infty} \mathbf{G}(K_v)$ where R_∞ is the subset of R of all archimedean valuations.

Theorem 1.14 (Automorphic bound for \mathbf{G}). *Let \mathbf{G} be connected absolutely almost simple K -group. Let U_∞ be a maximal compact subgroup of \mathbf{G}_∞ and W_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$.*

Then for any U_∞ -finite and W_f -invariant functions $f, h \in L^2_{00}(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$,

$$|\langle f, g.h \rangle| \leq c_{W_f} \cdot (\dim \langle U_\infty f \rangle \cdot \dim \langle U_\infty h \rangle)^{r_0} \cdot \tilde{\xi}_{\mathbf{G}}(g) \cdot \|f\|_2 \cdot \|h\|_2 \quad \text{for all } g \in \mathbf{G}(\mathbb{A}),$$

where $c_{W_f} > 0$ and $r_0 = r_0(\mathbf{G}_\infty) > 0$. Moreover if \mathbf{G}_∞ has no normal subgroup locally isomorphic to $\mathrm{Sp}_{2n}(\mathbb{C})$, then $r_0 = 1$.

Here, $\tilde{\xi}_{\mathbf{G}} : \mathbf{G}(\mathbb{A}) \rightarrow (0, 1]$ is an explicitly constructed proper function which is L^p -integrable for some $p = p(\mathbf{G}) < \infty$. (see Def. 2.15).

For each $v \in R$, denote $\hat{\mathbf{G}}_v^{\mathrm{Aut}} \subset \hat{\mathbf{G}}_v$ the automorphic dual of $\mathbf{G}(K_v)$, i.e., the subset of unitary dual of $\mathbf{G}(K_v)$ consisting of representations which are weakly contained in the representations appearing as $\mathbf{G}(K_v)$ components of $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))^{O_f}$ for some compact open subgroup O_f of $\mathbf{G}(\mathbb{A}_f)$.

Recall that for \mathbf{G} simply connected, it is shown by Clozel [Cl1] that the trivial representation of $\mathbf{G}(K_v)$ is isolated in $\hat{\mathbf{G}}_v^{\mathrm{Aut}}$ for each $v \in R$, that is, \mathbf{G} has property (τ) (cf. [Lu]). The following corollary presents a uniform version of property (τ) of \mathbf{G} over all $v \in R$.

Corollary 1.15. *Let \mathbf{G} be a connected simply connected absolutely almost simple K -group. Let π denote the quasi-regular representation of $\mathbf{G}(\mathbb{A})$ on $L^2_0(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. Let W be a maximal compact subgroup of $\mathbf{G}(\mathbb{A})$. Then there exist an explicit $p = p(\mathbf{G}) < \infty$ such that any W -finite matrix coefficient of π is $L^p(\mathbf{G}(\mathbb{A}))$ -integrable. In particular, π is strongly L^p .*

This corollary implies that for some $m = m(\mathbf{G}) < \infty$,

$$\pi^{\otimes m(\mathbf{G})} \subset \infty \cdot L^2(\mathbf{G}(\mathbb{A}))$$

(see [HT]), and that for any non-amenable closed subgroup $H \subset \mathbf{G}(\mathbb{A})$, the restriction of π to H does not have an almost invariant vector.

Let $K = \mathbb{Q}$. Let S be a finite set of primes including the archimedean prime ∞ . If Γ is an S -arithmetic subgroup of $\mathbf{G}_S := \prod_{p \in S} \mathbf{G}(\mathbb{Q}_p)$ (here $\mathbb{Q}_\infty = \mathbb{R}$) and $a \in \mathbf{G}(\mathbb{Q})$, then the Hecke operator T_a on $L^2(\Gamma \backslash \mathbf{G}_S)$ is defined by

$$T_a(f)(g) = \frac{1}{\deg(a)} \sum_{x \in \Gamma \backslash \Gamma a \Gamma} f(xg)$$

where $\deg(a) = \#\Gamma \backslash \Gamma a \Gamma$. Theorem 1.14 extends the main result in [COU] where some cases of \mathbb{Q} -anisotropic groups were excluded (see [EO]). In fact, the following corollary immediately follows from Theorem 1.14 and Proposition 2.6 in [COU]:

Corollary 1.16. *Let \mathbf{G} be a connected simply connected absolutely almost simple \mathbb{Q} -group and S a finite set of primes including ∞ . Suppose that \mathbf{G}_S is non-compact. Let $\Gamma \subset \mathbf{G}(\mathbb{Q})$ be an S -congruence subgroup of \mathbf{G}_S . There exists a constant $c = c(\Gamma) > 0$ such that*

$$\|T_a\| \leq c \cdot \tilde{\xi}_{\mathbf{G}}(a) \quad \text{for any } a \in \mathbf{G}(\mathbb{Q}).$$

This corollary in particular implies that for any sequence $a_i \in \mathbf{G}(\mathbb{Q})$ with $\deg(a_i) \rightarrow \infty$, and for any $f \in C_c(\mathbf{G}_S)$,

$$\lim_{i \rightarrow \infty} \frac{1}{\deg(a_i)} \sum_{x \in \Gamma a_i \Gamma} f(x) = \int_{\mathbf{G}_S} f(g) d\tau_S$$

where τ_S is the normalized Haar measure on \mathbf{G}_S so that $\tau_S(\Gamma \backslash \mathbf{G}_S) = 1$. It is interesting to note that unlike the rational points $\mathbf{G}(\mathbb{Q})$ of bounded height (Corollary 1.6), the Hecke points are equidistributed in \mathbf{G}_S with respect to the invariant measure.

The proof of Theorem 1.13 goes roughly as follows: if $\tilde{\xi}_v$ is a uniform bound for the matrix coefficients of infinite dimensional representations in $\hat{\mathbf{G}}_v^{\text{Aut}}$, $\tilde{\xi}_{\mathbf{G}}$ is defined to be the product $\prod_{v \in R} \tilde{\xi}_v$. This can be made precise using the language of direct integral of a representation (cf. proof of Theorem 2.7). For those $v \in R$ such that the K_v -rank of \mathbf{G} is at least 2, the uniform bounds, say ξ_v , of matrix coefficients of *all* infinite dimensional unitary representations of $\mathbf{G}(K_v)$ were obtained in [Oh1]. For these cases, one can simply take $\tilde{\xi}_v = \xi_v$. In particular, if K -rank of \mathbf{G} is at least 2, we have $\tilde{\xi}_{\mathbf{G}} = \prod_{v \in R} \xi_v$ and $\tilde{\xi}_{\mathbf{G}}$ works as a uniform bound for all unitary representation of $\mathbf{G}(\mathbb{A})$ without $\mathbf{G}(K_v)^+$ -invariant vectors for each $v \in R$ (see Theorem 2.7 for a precise statement). Moreover $\tilde{\xi}_{\mathbf{G}}$ is fairly sharp in these cases. For instance, one can show that $\tilde{\xi}_{\mathbf{G}}$ is optimal for $\mathbf{G} = \text{SL}_n$ ($n \geq 3$), or Sp_{2n} ($n \geq 2$) by [5.4, COU].

When there is $v \in R$ with K_v -rank of \mathbf{G} one, finding an automorphic bound $\tilde{\xi}_v$ is essentially carried out by Clozel [Cl1]. In particular, several deep theorems in automorphic theory were used such as the Gelbart-Jacquet bound [GJ] toward Ramanujan conjecture, the results of Burger-Sarnak [BS] and Clozel-Ullmo [CU] on lifting automorphic bounds, the base changes by Rogawski [Ro] and Clozel [Cl2], and Jacquet-Langlands correspondence [JL].

The paper is organized as follows. We discuss the mixing property of unitary representations of $\mathbf{G}(\mathbb{A})$ and prove Theorems 1.13 and 1.14 in section 2. In section 3, we derive the volume estimates from the work in [STT2] on height zeta functions. In section 4, we prove Theorems 1.2, 1.5, 1.7, 1.8 and 1.12 for the saturated cases. The rate of convergence is obtained in section 5, where we also prove Theorems 1.2 and 1.7 for the unsaturated cases.

Acknowledgement We would like to thank Wee Teck Gan, Dinakar Ramakrishnan and Yehuda Shalom for helpful conversations. We also thank Ramin Takloo-Bighash for the useful comments on our preliminary version of this paper.

2. ADELIC MIXING

Let K be a number field. Let \mathbf{G} be a connected absolutely almost simple group defined over K . We keep the same notation R, R_f, K_v as in the introduction. Let \mathcal{O} denote the ring of integers of K and \mathcal{O}_v the valuation ring of K_v . Set $R_\infty = R - R_f$. For $v \in R_f$, let q_v denote the order of the residue field of \mathcal{O}_v . Denote by \mathbb{A} the adèle ring over K and by $\mathbf{G}(\mathbb{A})$ the adèle group associated to \mathbf{G} .

Denote by $\mathbf{G}(\mathbb{A}_f)$ (resp. \mathbf{G}_∞) the subgroup of finite (resp. infinite) adèles, i.e., $((g_v)_v) \in \mathbf{G}(\mathbb{A})$ with $g_v = e$ for all $v \in R_\infty$ (resp. for all $v \in R_f$). Then

$$\mathbf{G}(\mathbb{A}) = \mathbf{G}_\infty \times \mathbf{G}(\mathbb{A}_f).$$

2.1. Definition and properties of $\xi_{\mathbf{G}}$. We fix a smooth model of \mathbf{G} over $\mathcal{O}[k^{-1}]$ for some $k \in \mathbb{Z}$. There exists a finite subset $S_0 \subset R_f$ such that for any $v \in R_f - S_0$, \mathbf{G} is unramified over K_v and $\mathbf{G}(\mathcal{O}_v)$ is a hyperspecial compact subgroup (cf. [Ti2]). We set $U_v = \mathbf{G}(\mathcal{O}_v)$ for each $v \in R_f - S_0$. Then for each $v \in R_f - S_0$, there exists the group A_v of K_v -rational points of a maximal K_v -split torus of \mathbf{G} so that the following Cartan decomposition holds:

$$\mathbf{G}(K_v) = U_v A_v^+ U_v$$

where A_v^+ is a closed positive Weyl chamber of A_v .

For $v \in S_0 \cup R_\infty$, there exists a good maximal compact subgroup U_v of $\mathbf{G}(K_v)$ such that

$$\mathbf{G}(K_v) = U_v A_v^+ \Omega_v U_v$$

where A_v is the group of K_v -rational points of a maximal K_v -split torus of \mathbf{G} and Ω_v is a finite subset in the centralizer of A_v in $\mathbf{G}(K_v)$.

In particular for any $g \in \mathbf{G}(K_v)$, there exist unique $a_v \in A_v^+$ and $d_v \in \Omega_v$ such that $g \in U_v a_v d_v U_v$. For $v \in R_\infty$, any maximal compact subgroup of $\mathbf{G}(K_v)$ is a good maximal compact subgroup and $\Omega_v = \{e\}$.

Let \mathcal{T} denote the set of $v \in R$ such that $\mathbf{G}(K_v)$ is compact, that is, $U_v = \mathbf{G}(K_v)$. It is well known that \mathcal{T} is a finite set.

Denote by Φ_v^+ the system of positive roots in the set of all non-multipliable roots of $\mathbf{G}(K_v)$ relative to A_v^+ and choose a maximal strongly orthogonal system \mathcal{S}_v in Φ_v^+ in the sense of [Oh2] (where an explicit construction is also given). If $v \in R - \mathcal{T}$ and $K_v \neq \mathbb{C}$, then define the bi- U_v -invariant function ξ_v on $\mathbf{G}(K_v)$ (cf. [Oh1]): for each $g = kadk' \in U_v A_v^+ \Omega_v U_v$,

$$\xi_v(g) = \prod_{\alpha \in \mathcal{S}_v} \Xi_v \begin{pmatrix} \alpha(a) & 0 \\ 0 & 1 \end{pmatrix}$$

where Ξ_v is the Harish-Chandra function of $\mathrm{PGL}_2(K_v)$. If $K_v = \mathbb{C}$, set

$$\xi_v(g) = \prod_{\alpha \in \mathcal{S}_v} \Xi_v \begin{pmatrix} \alpha(a) & 0 \\ 0 & 1 \end{pmatrix}^{n_\alpha}$$

where $n_\alpha = 1/2$ if α is a long root of \mathbf{G} which is locally isomorphic to $\mathrm{Sp}_{2n}(\mathbb{C})$ and $n_\alpha = 1$ for all other cases. We set $\xi_v = 1$ for $v \in \mathcal{T}$.

Definition 2.1. Define the function $\xi_{\mathbf{G}}$ on $\mathbf{G}(\mathbb{A})$ by

$$\xi_{\mathbf{G}}(g) = \prod_{v \in R} \xi_v(g_v) \quad \text{for } g = (g_v)_v \in \mathbf{G}(\mathbb{A}).$$

Since $0 < \xi_v(g_v) \leq 1$ for all $v \in R$ and $\xi_v(g_v) = 1$ for almost all v , $\xi_{\mathbf{G}}$ is a well defined function on $\mathbf{G}(\mathbb{A})$ and $0 < \xi_{\mathbf{G}} \leq 1$. Set

$$U = \prod_{v \in R} U_v.$$

Note also that $\xi_{\mathbf{G}}$ is bi- U -invariant.

For $v \in R - \mathcal{T}$, denoting by η_v the product of all positive roots in \mathcal{S}_v , set

$$\eta_v(kadk') := \eta_v(a)$$

where $kadk' \in U_v A_v^+ \Omega_v U_v$ for all v with $K_v \neq \mathbb{C}$. As in the case of the definition of ξ_v , if $K_v = \mathbb{C}$, we set $\eta_v = \prod_{\alpha \in \mathcal{S}_v} \alpha^{n_\alpha}$ with the same n_α defined as before. If $v \in \mathcal{T}$, we set $\eta_v = 1$.

Lemma 2.2. For any $\epsilon > 0$, there is a constant $C_\epsilon > 0$ such that for any $g = (g_v)_v \in \mathbf{G}(\mathbb{A})$,

$$(2.3) \quad \prod_{v \in R} |\eta_v(g_v)|^{-1/2} \leq \xi_{\mathbf{G}}(g) \leq C_\epsilon \cdot \prod_{v \in R} |\eta_v(g_v)|^{-1/2+\epsilon}.$$

In particular,

$$\xi_{\mathbf{G}}(g) \rightarrow 0 \quad \text{as } g \rightarrow \infty \text{ in } \mathbf{G}(\mathbb{A}).$$

Proof. For $v \in R - \mathcal{T}$, it follows from the explicit formula for Ξ_v (cf. [Oh1, 3.7.1]) that for any $\epsilon > 0$, there is a constant $C_{v,\epsilon} > 0$ such that for any $g_v \in \mathbf{G}(K_v)$,

$$|\eta_v(g_v)|^{-1/2} \leq \xi_v(g_v) \leq C_{v,\epsilon} \cdot |\eta_v(g_v)|^{-1/2+\epsilon}.$$

Moreover $C_{v,\epsilon} = 1$ for almost all v . This implies (2.3).

To see the second claim, first note that for any $g \in \mathbf{G}(\mathbb{A})$,

$$(2.4) \quad \xi_{\mathbf{G}}(g) \leq \xi_v(g_v) \leq C_{\epsilon} \cdot |\eta_v(g_v)|^{-1/2+\epsilon}.$$

Now suppose on the contrary that there exists a sequence $g_i \rightarrow \infty$ such that

$$\xi_{\mathbf{G}}(g_i) \not\rightarrow 0.$$

Then by passing to a subsequence we may assume either that there is a place $v \in R - \mathcal{T}$ such that $g_{i,v} \rightarrow \infty$ in $\mathbf{G}(K_v)$ or that there is v_i with $q_{v_i} \rightarrow \infty$ such that $g_{i,v_i} \notin U_{v_i}$ and $\Omega_{v_i} = \{e\}$ for all i . If $g_{i,v} \rightarrow \infty$ as $i \rightarrow \infty$, then $|\eta_v(g_{i,v})| \rightarrow \infty$ as $i \rightarrow \infty$ and hence $\xi_{\mathbf{G}}(g_i) \rightarrow 0$ by (2.3). Therefore the first case cannot happen.

In the second case, note that since $g_{i,v_i} \notin U_{v_i}$ and $\Omega_{v_i} = \{e\}$, we have $|\eta_v(g_{i,v_i})| \geq q_{v_i}$ for each i . Hence by (2.4),

$$\xi_{\mathbf{G}}(g_i) \leq C_{\epsilon} \cdot q_{v_i}^{-1/2+\epsilon}.$$

This gives a contradiction since $q_{v_i} \rightarrow \infty$. \square

Lemma 2.5. *Let $\iota : \mathbf{G} \rightarrow \mathrm{GL}_N$ be a faithful absolutely irreducible representation defined over K and H_{ι} be a height function on $\mathbf{G}(\mathbb{A})$ associated to ι . Then there exist $m \in \mathbb{N}$ and $C > 0$ such that*

$$\xi_{\mathbf{G}}(g) \leq C \cdot H_{\iota}^{-1/m}(g) \quad \text{for any } g \in \mathbf{G}(\mathbb{A}).$$

Proof. Let χ denote the highest weight of ι . Let $l \in \mathbb{N}$ be such that $\chi|_{A_v^+} \leq l \cdot \log_{q_v} \eta_v$ for each $v \in R$. Here $q_v = e$ if $v \in R_{\infty}$. Without loss of generality, we may assume

$$H_v(\iota(a)) = q_v^{\chi(a)} \quad \text{for each } a \in A_v^+ \text{ and } v \in R_f.$$

Since $|\eta_v(a_v)| = q_v^{\log_{q_v} \eta_v(a_v)}$ for $a_v \in A_v^+$, by Lemma 2.2, there exist some $c_1, c_2 > 0$ such that for any $g = (g_v) \in \mathbf{G}(\mathbb{A})$,

$$\xi_{\mathbf{G}}^{Al}(g) \leq c_1 \cdot \prod_v |\eta_v(g_v)|^{-l} \leq c_2 \cdot \prod_v H_v^{-1}(\iota(g_v)) = c_2 \cdot H_{\iota}^{-1}(g).$$

\square

Even though we do not need the following fact in this paper, it is of independent interest:

Proposition 2.6. *There exists $0 < p = p(\mathbf{G}) < \infty$ such that $\xi_{\mathbf{G}} \in L^p(\mathbf{G}(\mathbb{A}))$.*

Proof. Choose any absolutely irreducible representation of \mathbf{G} defined over K , for instance, the adjoint representation, and let H_ι be a height function on $\mathbf{G}(\mathbb{A})$ associated to ι (see (1.10)). By Theorem 3.4, the height zeta function

$$\mathcal{Z}(s) := \int_{\mathbf{G}(\mathbb{A})} H_\iota(g)^{-s} d\tau(g)$$

converges for $\operatorname{Re}(s) > a_\iota$ where a_ι is defined as in (1.3). Since $\xi_{\mathbf{G}}^m(g) \leq C \cdot H_\iota^{-1}(g)$ for some $m \in \mathbb{N}$ by Lemma 2.5, $\xi_{\mathbf{G}}$ is L^p -integrable for any $p > ma_\iota$. \square

2.2. Uniform bound for matrix coefficients of $\mathbf{G}(\mathbb{A})$. Let $W_f \subset \mathbf{G}(\mathbb{A}_f)$ be a compact open subgroup. Write $W_v = W_f \cap \mathbf{G}(K_v)$ for each $v \in R$. Then $W_v = U_v$ for almost all $v \in R_f$. For each $v \in R_f$, by [Be], there exists $d_{W_v} < \infty$ such that for any irreducible unitary representation ρ of $\mathbf{G}(K_v)$, the dimension of W_v -invariant vectors of ρ is at most d_{W_v} . Moreover d_{W_v} can be taken to 1 whenever W_v is a hyper-special compact subgroup. Hence the following number is well-defined:

$$d_{W_f} := \prod_{v \in R_f} d_{W_v} < \infty.$$

Set $U_f = \prod_{v \in R_f} U_v$, and $U_\infty := \prod_{v \in R_\infty} U_v$.

Theorem 2.7. *Let \mathbf{G} be a connected absolutely almost simple K -group with K -rank at least 2. Let W_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$. Let π be any unitary representation of $\mathbf{G}(\mathbb{A})$ without $\mathbf{G}(K_v)^+$ -invariant vector for every $v \in R$. Then for any U_∞ -finite and W_f -invariant unit vectors x and y ,*

$$(2.8) \quad |\langle \pi(g)x, y \rangle| \leq d_0 \cdot c_{W_f} \cdot (\dim \langle U_\infty x \rangle \cdot \dim \langle U_\infty y \rangle)^{(r+1)/2} \cdot \xi_{\mathbf{G}}(g) \quad \text{for all } g \in \mathbf{G}(\mathbb{A})$$

where $c_{W_f} := d_{W_f} \cdot \prod_v [U_v : U_v \cap W_v] \cdot (\max_{d \in \Omega_v} [U_v : dU_v d^{-1}])$ and $d_0, r \geq 1$ depend only on \mathbf{G} . Moreover if $\mathbf{G}(K_v) \not\cong \operatorname{Sp}_{2n}(\mathbb{C})$ locally for any $v \in R_\infty$, $d_0 = 1$ and $r = 1$.

The proof of above theorem is based on theorems in [Oh1]. More precisely, recall:

Theorem 2.9 (Theorem 1.1-2, [Oh1]). *Suppose that the K_v -rank of \mathbf{G} is at least 2. Let π_v be a unitary representation of $\mathbf{G}(K_v)$ without $\mathbf{G}(K_v)^+$ -invariant vectors. Then for any U_v -finite unit vectors x and y ,*

$$|\langle \pi_v(g)x, y \rangle| \leq d_v \cdot c_v \cdot (\dim \langle U_v x \rangle \cdot \dim \langle U_v y \rangle)^{r_v/2} \cdot \xi_v(g) \quad \text{for any } g \in \mathbf{G}(K_v)$$

where $c_v = \max_{d \in \Omega_v} [U_v : dU_v d^{-1}]$ and $d_v, r_v \geq 1$ depend only on $\mathbf{G}(K_v)$. Moreover whenever $\mathbf{G}(K_v) \not\cong \operatorname{Sp}_{2n}(\mathbb{C})$ locally, $d_v = 1$ and $r_v = 1$.

In the case when $\mathbf{G}(K_v) \cong \operatorname{Sp}_{2n}(\mathbb{C})$ locally, the above theorem was stated only for U_v -invariant vectors in [Oh1]. However using the remark following Prop. 2.7 in [Oh1], the proof can be modified for the above claim.

Proof of Theorem 2.7 For $g = (g_v)_v \in \mathbf{G}(\mathbb{A})$, choose a finite subset S_g of places containing

$$\{v \in R_f : g_v \notin U_v\} \cup R_\infty.$$

Note that for $v \in R - S_g$, we have $g_v \in U_v$ and hence $\xi_v(g_v) = 1$. Therefore for $g = (g_v)_v \in \mathbf{G}(\mathbb{A})$,

$$\xi(g) = \prod_{v \in S_g} \xi_v(g_v).$$

Let $G_g = \prod_{v \in S_g} \mathbf{G}(K_v)$ and $W_g = \prod_{v \in S_g \cap R_f} W_v$ for $W_v = W_f \cap \mathbf{G}(K_v)$. As a G_g representation, π has a Hilbert integral decomposition:

$$\pi = \int_{z \in Z_g} \oplus^{m_z} \rho_z d\nu(z)$$

where Z_g is the unitary dual of G_g and ρ_z is irreducible, m_z is a multiplicity for each $z \in Z_g$ and ν is a measure on Z_g (see [Di] or [Section 2.3, Zi]). We may assume that for all z , ρ_z has no $\mathbf{G}(K_v)^+$ -invariant vector (see [Prop. 2.3.2, Zi]).

If we write $\mathcal{L}_z = \oplus^{m_z} \rho_z$, $x = \int x_z d\nu(z)$ and $y = \int y_z d\nu(z)$ with

$$x_z = \sum_{i=1}^{m_z} x_{zi} \quad \text{and} \quad y_z = \sum_{i=1}^{m_z} y_{zi} \in \mathcal{L}_z,$$

we have

$$\langle x, y \rangle = \int_{Z_g} \sum_i \langle x_{zi}, y_{zi} \rangle d\nu(z).$$

It follows from the definition of a Hilbert direct integral that

$$\dim\langle U_\infty x_{zi} \rangle \leq \dim\langle U_\infty x_z \rangle \leq \dim\langle U_\infty x \rangle,$$

x_{zi} is W_g -invariant for almost all z and all i , and similarly for y . Without loss of generality, we assume the above holds for all z . We claim that

(2.10)

$$|\langle \rho_z(g) x_{zi}, y_{zi} \rangle| \leq c_{W_f} \cdot d_0 \cdot \xi_{\mathbf{G}}(g) \cdot (\dim\langle U_\infty x \rangle \cdot \dim\langle U_\infty y \rangle)^{(r+1)/2} \cdot \|x_{zi}\| \cdot \|y_{zi}\|$$

where $r = \max_v r_v$ and $c_0 = d_{W_f} \prod_v (c_v \cdot [U_v : U_v \cap W_v]) < \infty$, $d_0 = \prod_v d_v < \infty$ with c_v, d_v, r_v as in Theorem 2.9. Since G_g is a type (I) group [Be], we may write $\rho_z = \otimes_{v \in S_g} \rho_{z(v)}$ where $\rho_{z(v)}$ is an irreducible representation of $\mathbf{G}(K_v)$ without $\mathbf{G}(K_v)^+$ -invariant vectors. Since the finite linear combinations of pure tensor vectors is dense, it suffices to prove (2.10) assuming x_{zi} and y_{zi} are finite sum of pure tensors. Hence we can write

$$x_{zi} = \sum_j \otimes_{v \in S_g} x_{zij(v)}; \quad y_{zi} = \sum_k \otimes_{v \in S_g} y_{zik(v)}$$

where for each $v \in S_g$, $x_{zij(v)}$ (resp. $y_{zik(v)}$) are mutually orthogonal and the number of summands for x_{zi} (resp. y_{zi}) is at most $\dim\langle U_\infty x \rangle \cdot d_{W_f}$ (resp. $\dim\langle U_\infty y \rangle \cdot d_{W_f}$). Hence by Cauchy-Schwartz inequality, for $x_{zij} = \prod_{v \in S_g} x_{zij(v)}$ and $y_{zij} = \prod_{v \in S_g} y_{zij(v)}$

$$\sum_j \|x_{zij}\| \leq (\dim\langle U_\infty x \rangle \cdot d_{W_f})^{1/2} \|x_{zi}\|; \quad \text{and} \quad \sum_k \|y_{zik}\| \leq (\dim\langle U_\infty y \rangle \cdot d_{W_f})^{1/2} \|y_{zi}\|.$$

Since for $v \in R_f$

$$\dim\langle U_v x \rangle \leq [U_v : W_v \cap U_v] \quad \text{and} \quad \dim\langle U_v y \rangle \leq [U_v : W_v \cap U_v],$$

by Theorem 2.9, we have for $c_0 = \prod_v c_v$,

(2.11)

$$\begin{aligned} |\langle \rho_z(g)x_{zi}, y_{zi} \rangle| &\leq \sum_{j,k} \prod_{v \in S_g} |\langle \rho_z(v)(g_v)x_{zij(v)}, y_{zik(v)} \rangle| \\ &\leq c_0 \cdot d_0 \cdot \prod_{v \in S_g} \xi_v(g_v) \cdot (\dim\langle U_\infty x \rangle \cdot \dim\langle U_\infty y \rangle)^{r/2} \left(\prod_{v \in R_f} [U_v : W_v \cap U_v] \cdot \left(\sum_{j,k} \|x_{zij}\| \cdot \|y_{zik}\| \right) \right) \\ &\leq c_0 \cdot d_0 \cdot \xi_{\mathbf{G}}(g) \cdot (\dim\langle U_\infty x \rangle \cdot \dim\langle U_\infty y \rangle)^{(r+1)/2} \cdot \prod_{v \in R_f} [U_v : W_v \cap U_v] \cdot d_{W_f}(\|x_{zi}\| \cdot \|y_{zi}\|) \\ &= c_{W_f} \cdot d_0 \cdot \xi_{\mathbf{G}}(g) \cdot (\dim\langle U_\infty x \rangle \cdot \dim\langle U_\infty y \rangle)^{(r+1)/2} \cdot (\|x_{zi}\| \cdot \|y_{zi}\|) \end{aligned}$$

proving (2.10). Therefore again by Cauchy-Schwartz inequality,

(2.12)

$$\begin{aligned} |\langle (\oplus^{m_z} \rho_z)(g)(x_z), y_z \rangle| &\leq \sum_i |\langle \rho_z(g)x_{zi}, y_{zi} \rangle| \\ &\leq c_{W_f} \cdot d_0 \cdot \xi_{\mathbf{G}}(g) \cdot (\dim\langle U_\infty x \rangle \cdot \dim\langle U_\infty y \rangle)^{(r+1)/2} \cdot \|x_z\| \cdot \|y_z\|. \end{aligned}$$

By integrating over Z_g , we obtain (2.8).

Since U -finite vectors form a dense subset by Peter-Weyl theorem, the above implies an adelic version of Howe-Moore theorem [HM] on the vanishing of matrix coefficients:

Corollary 2.13. *Let \mathbf{G} and π be as in Theorem 2.7. Then for any vectors x and y ,*

$$\langle \pi(g)x, y \rangle \rightarrow 0 \quad \text{as } g \rightarrow \infty \text{ in } \mathbf{G}(\mathbb{A}).$$

2.3. Automorphic bound for $\mathbf{G}(\mathbb{A})$. If \mathbf{G} has K -rank at most one, the analogue of Theorem 2.7 does not hold in general. However if we look at those infinite dimensional representations occurring in $L^2(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$, we still obtain a similar upper bound.

We now consider the unitary representation of $\mathbf{G}(\mathbb{A})$ on the space $L^2(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$ by right translations. Let $L_{00}^2(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$ denote the orthogonal complement to the direct sum of all one-dimensional representations occurring in $L^2(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$. If \mathbf{G} is simply connected, it follows from the strong approximation property that $L_{00}^2(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$ is the complement to the space of constant functions (cf. Lemma 2.18).

We first state the following conjecture:

Conjecture 2.14. *Let \mathbf{G} be a connected absolutely almost simple K -group. Let W_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$. Then for any $U_\infty \times W_f$ -invariant unit vectors*

$f, h \in L_{00}^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$,

$$|\langle f, g \cdot h \rangle| \leq c_{W_f} \cdot \xi_{\mathbf{G}}(g) \quad \text{for all } g \in \mathbf{G}(\mathbb{A})$$

where $c_{W_f} > 0$ is a constant depending only on \mathbf{G} and W_f .

The above holds for groups of K -rank at least 2 by Theorem 2.7. For $\mathbf{G} = \mathrm{PGL}_2$, Conjecture 2.14 is essentially equivalent to the Ramanujan conjecture. We will prove a weaker statement of Conjecture 2.14 where the function $\xi_{\mathbf{G}}$ is replaced by a function $\tilde{\xi}_{\mathbf{G}}$ with slower decay such that $\xi_{\mathbf{G}} \leq \tilde{\xi}_{\mathbf{G}} \leq \xi_{\mathbf{G}}^{1/2}$.

Definition 2.15. Set $R_1 := \{v \in R : \mathrm{rank}_{K_v}(\mathbf{G}) = 1\}$. Define

$$\tilde{\xi}_{\mathbf{G}}(g) := \prod_{v \in R_1} \xi_v(g_v)^{1/2} \prod_{v \in R - R_1} \xi_v(g_v)$$

where $g = (g_v)_v \in \mathbf{G}(\mathbb{A})$.

Theorem 2.16 (Automorphic bounds). *Let \mathbf{G} be a connected absolutely almost simple K -group. Let W_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$. Then for any U_{∞} -finite and W_f -invariant unit vectors $x, y \in L_{00}^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$,*

$$|\langle x, g \cdot y \rangle| \leq c_{W_f} \cdot (\dim \langle U_{\infty} x \rangle \cdot \dim \langle U_{\infty} y \rangle)^{(r+1)/2} \cdot \tilde{\xi}_{\mathbf{G}}(g) \quad \text{for all } g \in \mathbf{G}(\mathbb{A})$$

where $c_{W_f} > 0, r = r(\mathbf{G}) \geq 1$. Moreover $r = 1$ provided for any $v \in R$, $\mathbf{G}(K_v) \not\cong \mathrm{Sp}_{2n}(\mathbb{C})$ ($n \geq 2$) locally.

Recall that for unitary representations ρ_1 and ρ_2 of $\mathbf{G}(K_v)$, ρ_1 is said to be weakly contained in ρ_2 if every diagonal matrix coefficients of ρ_1 can be approximated uniformly on compact subsets by convex combinations of diagonal matrix coefficients of ρ_2 . For each $v \in R$, denote by $\hat{\mathbf{G}}_v$ the unitary dual of $\mathbf{G}(K_v)$ and by $\hat{\mathbf{G}}_v^{\mathrm{Aut}} \subset \hat{\mathbf{G}}_v$ the automorphic dual of $\mathbf{G}(K_v)$, i.e., the subset of unitary dual of $\mathbf{G}(K_v)$ consisting of representations which are weakly contained in the representations appearing as $\mathbf{G}(K_v)$ components of $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))^{O_f}$ for some compact open subgroup O_f of $\mathbf{G}(\mathbb{A}_f)$.

Theorem 2.17 (Burger-Sarnak [BS], Clozel-Ullmo [CU]). *Let \mathbf{G} be a connected absolutely almost simple K -group. Let $\mathbf{H} \subset \mathbf{G}$ be a connected semisimple K -subgroup. Then for any $v \in R$ and for any $\rho_v \in \hat{\mathbf{G}}_v^{\mathrm{Aut}}$, any irreducible representation of $\mathbf{H}(K_v)$ weakly contained in $\rho_v|_{\mathbf{H}(K_v)}$ is contained in $\hat{\mathbf{H}}_v^{\mathrm{Aut}}$.*

Lemma 2.18. *For any $v \in R$ such that $\mathbf{G}(K_v)$ is non-compact, $L_{00}^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$ has no non-zero $\mathbf{G}(K_v)^+$ -invariant function.*

Proof. (cf. proof of Lemma 3.8 in [GO].) Let \mathcal{L}_v denote the set of $f \in L_{00}^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$ fixed by $\mathbf{G}(K_v)^+$. We need to show that $\mathcal{L}_v = \{0\}$. Since the continuous functions form a dense subset of \mathcal{L}_v , it suffices to show that any continuous function $f \in \mathcal{L}_v$ is

trivial. Let $f \in \mathcal{L}_v$ be continuous. Let $\tilde{\mathbf{G}}$ be the simply connected cover of \mathbf{G} and denote by $\text{pr} : \tilde{\mathbf{G}} \rightarrow \mathbf{G}$ the covering map. Consider the projection map

$$\tilde{\mathbf{G}}(K) \backslash \tilde{\mathbf{G}}(\mathbb{A}) \rightarrow \mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}).$$

Let \tilde{f} be the pull back of f . Since the image of $\tilde{\mathbf{G}}(K_v)$ is $\mathbf{G}(K_v)^+$ under the map pr , the function \tilde{f} is left $\tilde{\mathbf{G}}(K)$ -invariant and right $\tilde{\mathbf{G}}(K_v)$ -invariant. On the other hand, the strong approximation property implies that $\tilde{\mathbf{G}}(K)\tilde{\mathbf{G}}(K_v)$ is dense in $\tilde{\mathbf{G}}(\mathbb{A})$ (cf. [Theorem 7.12, PR]). Therefore \tilde{f} is constant. It follows that f is a sum of characters of $\mathbf{G}(\mathbb{A})$, and hence 0 since $f \in L_{00}^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. \square

Proof of Theorem 2.16 The case when K -rank is at least 2 follows from Theorem 2.7 and Lemma 2.18. Suppose first that \mathbf{G} has K -rank one. Then there is a K -embedding of a K -group \mathbf{H} where \mathbf{H} is either SL_2 or PGL_2 . Using a Gelbart-Jacquet [GJ] estimate toward the Ramanujan conjecture for SL_2 or PGL_2 and Theorem 2.17, we obtain for $v \in R_1$, any infinite dimensional $\rho_v \in \hat{\mathbf{G}}_v^{\text{Aut}}$, and U_v -finite vectors x_v, y_v , (cf. Theorem 3.4 [COU])

$$(2.19) \quad |\langle \rho_v(g)(x_v), y_v \rangle| \leq c_v \cdot \xi_v(g)^{1/2} \cdot (\dim \langle U_v x_v \rangle \cdot \dim \langle U_v y_v \rangle)^{1/2}$$

for any $g \in \mathbf{G}(K_v)$. Combining this with Theorem 2.9, we can derive the desired bound by the same argument as in the proof of Theorem 2.7.

Now suppose \mathbf{G} is K -anisotropic. If $R_1 \neq \emptyset$, it follows from the classification theorem by Tits [Ti] that \mathbf{G} is of Dynkin type \mathcal{A} . Applying [Theorem 1.1, Cl], we deduce that there exists a K -embedding of K -subgroup \mathbf{H} of type \mathcal{A} such that \mathbf{H} has K_v -rank one whenever $v \in R_1$. Let $v \in R_1$. Then up to isogeny, we have either that $\mathbf{H} = \text{PGL}_1(D)$ for a quaternion algebra D over K and $\mathbf{H} = \text{PGL}_2$ over K_v , or $\mathbf{H} = \text{PGU}(D, *)$ for a division algebra D of prime degree d over a quadratic extension F of K with a second kind involution $*$, and $\mathbf{H} = \text{PGU}(n-1, 1)$ over K_v (with $n \geq 3$). In the former case we use the Jacquet-Langlands correspondence [JL] to transfer the Gelbart-Jacquet automorphic bound of PGL_2 to $\mathbf{H}(K_v)$ via Theorem 2.17. In the second case which is hardest, it is explained in [Cl] that by the base changes obtained by Rogawski [Ro] and Clozel [Cl2], we can use the bound of $\text{PGL}_n(F_w)$ to get a bound for $\mathbf{H}(K_v)$ where w is a place of F lying above v and F_w is a quadratic extension of K_v . This gives us (2.19) for $v \in R_1$ again in view of Theorem 2.17 and Theorem 2.9. Combining with Theorem 2.9 for those places $v \in R - (R_1 \cup \mathcal{T})$ as in the proof of Theorem 2.7, we obtain the desired bound. This finishes the proof.

Let X_1, \dots, X_m be an orthonormal basis of the Lie algebra $\text{Lie}(U_\infty)$ with respect to an Ad-invariant scalar product. Then the elliptic operator

$$(2.20) \quad \mathcal{D} := 1 - \sum_{i=1}^m X_i^2$$

lies in the center of the universal enveloping algebra of $\text{Lie}(U_\infty)$. We say a function f on $\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A})$ is smooth if f is W_f -invariant for some compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$ and smooth for the action of \mathbf{G}_∞ .

In Theorem 2.16, we can relax U_∞ -finite conditions to smooth conditions provided we replace the L^2 -norms by L^2 -Sobolev norms:

Theorem 2.21. *Let \mathbf{G} be a connected absolutely almost simple K -group. Let W_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$. Then for any W_f -invariant smooth functions $f, h \in L^2_{00}(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$ with compact support,*

$$|\langle f, g.h \rangle| \leq c_{W_f} \cdot \tilde{\xi}_{\mathbf{G}}(g) \cdot \|\mathcal{D}^l(f)\| \cdot \|\mathcal{D}^l(h)\| \quad \text{for all } g \in \mathbf{G}(\mathbb{A})$$

where $c_{W_f} > 0$ and l is any sufficiently large integer.

Proof. Deducing this from Theorem 2.16 is quite standard in view of the results of Harish-Chandra explained in [Ch 4, Wa]. We give a sketch of the proof. Denote by π the representation $L^2_{00}(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$. Then $\pi = \bigoplus_{\nu \in \hat{U}_\infty} \pi_\nu$ where π_ν is the ν -isotypic component of π and \mathcal{D} acts as a scalar, say, c_ν on each π_ν . We write $f = \sum_{\nu \in \hat{U}_\infty} f_\nu$ and $h = \sum_{\nu \in \hat{U}_\infty} h_\nu$. One has $\|f_\nu\| = c_\nu^{-l} \|\mathcal{D}^l f_\nu\|$ and similarly for h . Then

$$|\langle f, g.h \rangle| \leq \sum_{(\nu_1, \nu_2) \in \hat{U}_\infty \times \hat{U}_\infty} |\langle f_{\nu_1}, g.h_{\nu_2} \rangle|$$

Using Theorem 2.16, we then obtain

$$\begin{aligned} & |\langle f, g.h \rangle| \\ & \leq c_{W_f} \cdot \tilde{\xi}_{\mathbf{G}}(g) \left(\sum_{\nu \in \hat{U}_\infty} \|f_\nu\| \dim \langle U_\infty f_\nu \rangle^{(r+1)/2} \right) \left(\sum_{\nu \in \hat{U}_\infty} \|h_\nu\| \dim \langle U_\infty h_\nu \rangle^{(r+1)/2} \right) \\ & \leq c_{W_f} \cdot \tilde{\xi}_{\mathbf{G}}(g) \cdot \|\mathcal{D}^l(f)\| \cdot \|\mathcal{D}^l(h)\| \cdot \sum_{\nu \in \hat{U}_\infty} c_\nu^{-2l} \dim(\nu)^{r+1} \end{aligned}$$

Now if $l \in \mathbb{N}$ is sufficiently large, then $\sum_{\nu} c_\nu^{-2l} \dim(\nu)^{r+1} < \infty$ [Wa]. This proves the claim. \square

If \mathbf{G} is a connected semisimple K -group, we say that a sequence $\{g_i \in \mathbf{G}(\mathbb{A})\}$ tends to infinity strongly if for any non-trivial connected simple normal K -subgroup \mathbf{H} of \mathbf{G} , $\pi(g_i)$ tends to ∞ as $i \rightarrow \infty$, where $\pi : \mathbf{G}(\mathbb{A}) \rightarrow \mathbf{G}(\mathbb{A})/\mathbf{H}(\mathbb{A})$ denotes the canonical projection.

Theorem 2.22 (Mixing for $L^2(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$). *Let \mathbf{G} be a product of connected absolutely almost simple K -groups. Then for any $f, h \in L^2_{00}(\mathbf{G}(K)\backslash\mathbf{G}(\mathbb{A}))$,*

$$\langle f, g.h \rangle \rightarrow 0$$

as $g \in \mathbf{G}(\mathbb{A})$ tends to infinity strongly.

Proof. Write $\mathbf{G} = \mathbf{G}_1 \times \cdots \times \mathbf{G}_m$ where each \mathbf{G}_i is a connected absolutely almost simple K -group. By Theorem 2.16 and Peter-Weyl theorem, for each $1 \leq i \leq m$, and for any $f_i, h_i \in L^2_{00}(\mathbf{G}_i(K) \backslash \mathbf{G}_i(\mathbb{A}))$,

$$(2.23) \quad \langle f_i, g_i \cdot h_i \rangle \rightarrow 0$$

as $g_i \rightarrow \infty$ in $\mathbf{G}_i(\mathbb{A})$.

Consider $\otimes_{i=1}^m L^2(\mathbf{G}_i(K) \backslash \mathbf{G}_i(\mathbb{A}))$ as a subset of $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. Then the finite sums of the functions of the form $h = \otimes_{i=1}^m h_i \in L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$, $h_j \in L^2(\mathbf{G}_j(K) \backslash \mathbf{G}_j(\mathbb{A}))$, such that for at least one j , $h_j \in L^2_{00}(\mathbf{G}_j(K) \backslash \mathbf{G}_j(\mathbb{A}))$ form a dense subset of the space $L^2_{00}(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. Hence it suffices to prove the claim for $f = \otimes_{i=1}^m f_i$ and $h = \otimes_{i=1}^m h_i$ of such type. Suppose $h_j \in L^2_{00}(\mathbf{G}_j(K) \backslash \mathbf{G}_j(\mathbb{A}))$ for some $1 \leq j \leq m$. If $g = (g_1, \cdots, g_m)$ with $g_i \in \mathbf{G}_i(\mathbb{A})$, then

$$|\langle f, g \cdot h \rangle| = \prod_{i=1}^m |\langle f_i, g_i \cdot h_i \rangle| \leq |\langle f_j, g_j \cdot h_j \rangle| \cdot \left(\prod_{i \neq j} \|f_i\| \cdot \|h_i\| \right).$$

If f'_j denotes the projection of f_j to $L^2_{00}(\mathbf{G}_j(K) \backslash \mathbf{G}_j(\mathbb{A}))$, then

$$\langle f_j, g_j \cdot h_j \rangle = \langle f'_j, g_j \cdot h_j \rangle.$$

Since $g \rightarrow \infty$ strongly and hence $g_j \rightarrow \infty$, we obtain $\langle f'_j, g_j \cdot h_j \rangle \rightarrow 0$ by (2.23). This proves the claim. \square

3. VOLUME ASYMPTOTICS

Let \mathbf{G} be a connected adjoint semisimple algebraic group over K . As in the previous section, let $L^2_{00}(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$ be the orthogonal complement to the direct sum of all one dimensional representations in $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$. Denote by Λ an orthogonal basis for $L^2_{00}(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))^\perp$ consisting of continuous unitary characters of $\mathbf{G}(\mathbb{A})$. Then

$$L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A})) = L^2_{00}(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A})) \oplus \sum_{\chi \in \Lambda} \mathbb{C}\chi.$$

For a compact open subgroup W_f of $\mathbf{G}(\mathbb{A}_f)$, set

$$\Lambda^{W_f} := \{\chi \in \Lambda : u \cdot \chi = \chi \text{ for any } u \in W_f\};$$

$$\text{Ker}(\Lambda^{W_f}) := \cap \{\ker(\chi) : \chi \in \Lambda^{W_f}\}.$$

Note that $\text{Ker}(\Lambda^{W_f})$ is a normal subgroup of $\mathbf{G}(\mathbb{A})$. Moreover:

Lemma 3.1. *We have*

- (1) $\mathbf{G}(K)\mathbf{G}_\infty^\circ W_f \subset \text{Ker}(\Lambda^{W_f})$.
- (2) $\#\Lambda^{W_f} = [\mathbf{G}(\mathbb{A}) : \text{Ker}(\Lambda^{W_f})] < \infty$.

(3) For any $g \in \mathbf{G}(\mathbb{A})$,

$$\sum_{\chi \in \Lambda^{W_f}} \chi(g) = \begin{cases} \#\Lambda^{W_f} & \text{if } g \in \text{Ker}(\Lambda^{W_f}) \\ 0 & \text{otherwise} \end{cases}.$$

Proof. Note that an element of Λ is precisely a continuous homomorphism $\chi : \mathbf{G}(\mathbb{A}) \rightarrow S^1$ which contains $\mathbf{G}(K)$ in the kernel, where S^1 is the unit circle. Since \mathbf{G}_∞° is a connected group, $\mathbf{G}_\infty^\circ \subset \text{ker}(\chi)$ for any $\chi \in \Lambda$. Hence

$$\mathbf{G}(K)\mathbf{G}_\infty^\circ W_f \subset \text{Ker}(\Lambda^{W_f}).$$

Since \mathbf{G}_∞° has a finite index in \mathbf{G}_∞ , it follows from [Theorem 5.1, PR] that there exist finitely many $u_1, \dots, u_h \in \mathbf{G}(\mathbb{A})$ such that

$$\mathbf{G}(\mathbb{A}) = \cup_{i=1}^h \mathbf{G}(K)u_i\mathbf{G}_\infty^\circ W_f.$$

It follows $[\mathbf{G}(\mathbb{A}) : \text{Ker}(\Lambda^{W_f})] < \infty$. Clearly the quotient $\mathbf{G}(\mathbb{A})/\text{Ker}(\Lambda^{W_f})$ is a finite abelian group whose dual is isomorphic to Λ^{W_f} . Hence (2) and (3) easily follow from the duality of finite abelian groups. \square

Let $\iota : \mathbf{G} \rightarrow \text{GL}_N$ be an absolutely irreducible faithful representation defined over K . We give a definition of a height function on $\mathbf{G}(\mathbb{A})$ associated to ι which is slightly more general than those considered in the introduction (see (1.10)). It is this class of the functions for which we prove our main theorems.

Definition 3.2. A height function H_ι on $\mathbf{G}(\mathbb{A})$ is defined by the product $\prod_{v \in R} H_{\iota,v}$ where $H_{\iota,v}$ is a function on $\mathbf{G}(K_v)$ for $v \in R$ satisfying the following:

(1) there exists a finite subset $S \subset R$ such that

$$H_{\iota,v}(g) = \max_{ij} |\iota(g)_{ij}|_v \quad \text{for all } v \in R - S;$$

(2) for $v \in S$, there exists $C > 0$ such that

$$C^{-1} \cdot \max_{ij} |\iota(g)_{ij}|_v \leq H_{\iota,v}(g) \leq C \cdot \max_{ij} |\iota(g)_{ij}|_v;$$

(3) for any $v \in S \cap R_\infty$, there exists $b > 0$ such that for any small $\epsilon > 0$,

$$(1 - b \cdot \epsilon)H_{\iota,v}(x) \leq H_{\iota,v}(gxh) \leq (1 + b \cdot \epsilon)H_{\iota,v}(x)$$

for any $x \in \mathbf{G}(K_v)$ and any g, h in the ϵ -neighborhood of e in $\mathbf{G}(K_v)$ with respect to a Riemannian metric;

(4) for any $v \in S \cap R_f$, $H_{\iota,v}$ is bi-invariant under a compact open subgroup of $\mathbf{G}(K_v)$.

Fix a height function H_ι on $\mathbf{G}(\mathbb{A})$. Note that the height function H_ι is bi-invariant under a compact open subgroup, say W_f , of $\mathbf{G}(\mathbb{A}_f)$.

For $T > 0$, set

$$B_T := \{g \in \mathbf{G}(\mathbb{A}) : H_\iota(g) < T\}.$$

For a finite subset S of R , let \mathbf{G}^S denote the subgroup of $\mathbf{G}(\mathbb{A})$ consisting of (g_v) , with $g_v = e$ for all $v \in S$, and set $\mathbf{G}_S := \prod_{v \in S} \mathbf{G}(K_v)$. Note that $\mathbf{G}^\emptyset = \mathbf{G}(\mathbb{A})$ and $\mathbf{G}(\mathbb{A}) = \mathbf{G}_S \mathbf{G}^S$. Let τ_S and τ^S denote Haar measures on \mathbf{G}_S and \mathbf{G}^S respectively.

The following lemma easily follows from the decomposition of a Haar measures dg_v in terms of Cartan decompositions.

Lemma 3.3. *There exists $\epsilon > 0$ such that for every finite set $S \subset R$,*

$$\tau_S(B_T \cap \mathbf{G}_S) = O(T^{a_\iota - \epsilon}).$$

where the implied constant depends on S . In particular, for any $v \in R$,

$$\int_{\mathbf{G}(K_v)} H_{\iota, v}(g_v)^{-a_\iota} dg_v < \infty.$$

Consider the following height zeta function

$$\mathcal{Z}_{W_f}^S(s) := \sum_{\chi \in \Lambda^{W_f}} \int_{\mathbf{G}^S} H_\iota(g)^{-s} \chi(g) d\tau^S(g).$$

The following is shown in [STT2]:

Theorem 3.4. *Let a_ι and b_ι be as in (1.3). Then for any finite subset S of R , there exists $\epsilon > 0$ such that $\mathcal{Z}_{W_f}^S(s)$ has meromorphic continuations to $\operatorname{Re}(s) > a_\iota - \epsilon$ with unique poles at $s = a_\iota$ of order b_ι and of positive residues. Moreover For some constants $\kappa \in \mathbb{R}$ and $k > 0$,*

$$\left| \frac{(s - a_\iota)^{b_\iota} \mathcal{Z}_{W_f}^S(s)}{s^{b_\iota}} \right| \leq k \cdot |1 + \operatorname{Im}(s)|^\kappa$$

for $\operatorname{Re}(s) > a_\iota - \epsilon$.

We use the following version of Ikehara Tauberian theorem to deduce the volume asymptotics for $B_T \cap \operatorname{Ker}(\Lambda^{W_f}) \cap \mathbf{G}^S$ from Theorem 3.4.

Theorem 3.5. *Fix $a > 0$ and $\delta > 0$. Let $\alpha(t)$ be a non-negative non-decreasing function on (δ, ∞) such that*

$$f(s) := \int_\delta^\infty t^{-s} d\alpha$$

converges for $\operatorname{Re}(s) > a$. Suppose that

- $f(s)$ has a meromorphic continuation to the half plane $\operatorname{Re}(s) > a - \epsilon > 0$ and has a unique pole at $s = a$ with order b ;
- For some $\kappa \in \mathbb{R}$ and $k > 0$,

$$\left| \frac{f(s)(s - a)^b}{s^b} \right| \leq k \cdot |1 + \operatorname{Im}(s)|^\kappa$$

for $\operatorname{Re}(s) > a - \epsilon$.

Then for some $\delta > 0$,

$$\int_{\delta}^T d\alpha = \alpha(T) - \alpha(\delta) = \frac{c}{a(b-1)!} \cdot T^a P(\log T) + O(T^{a-\delta}) \quad \text{as } T \rightarrow \infty$$

where $c = \lim_{s \rightarrow a} (s-a)^b f(s)$ and $P(x)$ is a monic polynomial of degree $b-1$.

Proof. This can be proven by repeating the same argument as in the appendix of [CT1] simply replacing the sum $\sum_n n^{-s} \alpha_n$ by the integral $\int_{\delta}^{\infty} t^{-s} d\alpha(t)$. \square

Lemma 3.6. • *There exists $\delta_0 > 0$ such that*

$$\inf_{g \in \mathbf{G}(\mathbb{A})} H_{\iota}(g) \geq \delta_0.$$

- *For each $T > 0$, B_T is a relatively compact subset of $\mathbf{G}(\mathbb{A})$. In particular, the height function $H_{\iota} : \mathbf{G}(\mathbb{A}) \rightarrow [\delta_0, \infty)$ is proper.*

Proof. We note that there exists a finite subset, say, R_0 , such that for all $v \in R - R_0$, $H_v(\iota(g)) \geq 1$ for any $g \in \mathbf{G}(K_v)$. Let $0 < \delta \leq 1$ be such that $H_v(\iota(g)) \geq \delta$ for $v \in R_0$ and $\delta_0 = \delta^{\#R_0}$. Then $H_{\iota}(g) \geq \delta_0$ for all $g \in \mathbf{G}(\mathbb{A})$. Note that

$$B_T \subset \mathbf{G}(\mathbb{A}) \cap \prod_v \{g_v \in \mathbf{G}(K_v) : H_v(\iota(g_v)) \leq \delta_0^{-1} T\}.$$

Since for almost all $v \in R_f$, $H_v(\iota(g_v)) \geq q_v$ whenever $g_v \notin \mathbf{G}(\mathcal{O}_v)$, it follows that for some finite subset $R_1 \subset R$, we have

$$B_T \subset \{(g_v)_v \in \mathbf{G}(\mathbb{A}) : H_v(\iota(g_v)) \leq \delta_0^{-1} T \text{ for } v \in R_1, \ g_v \in \mathbf{G}(\mathcal{O}_v) \text{ otherwise}\}$$

and hence B_T is a relatively compact subset of $\mathbf{G}(\mathbb{A})$. \square

Theorem 3.7. *Let $a_{\iota} \in \mathbb{Q}^+$ and $b_{\iota} \in \mathbb{N}$ be as in (1.3). Then for any finite subset $S \subset R$,*

$$\tau^S(B_T \cap \text{Ker}(\Lambda^{W_f}) \cap \mathbf{G}^S) = \frac{c_{S, W_f}}{\#\Lambda^{W_f} \cdot a_{\iota}(b_{\iota} - 1)!} \cdot T^{a_{\iota}} P(\log T) + O(T^{a_{\iota} - \delta}) \quad \text{as } T \rightarrow \infty.$$

where $c_{S, W_f} = \lim_{s \rightarrow a_{\iota}} (s - a_{\iota})^{b_{\iota}} \mathcal{Z}_{W_f}^S(s)$, $P(x)$ is a monic polynomial of degree $b_{\iota} - 1$, and $\delta > 0$.

Proof. By the above lemma, B_T is a relatively compact subset of $\mathbf{G}(\mathbb{A})$ and hence $\tau^S(B_T \cap \mathbf{G}^S) < \infty$ for each $T \geq 1$ and for any finite S . Let δ_0 be as in the above lemma. Let

$$\alpha(t) = \tau^S(B_t \cap \text{Ker}(\Lambda^{W_f}) \cap \mathbf{G}^S) \quad \text{for } t \in [\delta_0, \infty),$$

and

$$f(s) = \int_{\delta_0}^{\infty} t^{-s} d\alpha.$$

By Lemma 3.1(3),

$$\mathcal{Z}_{W_f}^S(s) = \#\Lambda^{W_f} \cdot \int_{\text{Ker}(\Lambda^{W_f}) \cap \mathbf{G}^S} H_\iota(g)^{-s} d\tau^S(g) = \#\Lambda^{W_f} \cdot \int_{\delta_0}^{\infty} t^{-s} d\alpha.$$

Note that

$$\lim_{s \rightarrow a_\iota} (s - a_\iota)^{b_\iota} f(s) = \frac{c_{S, W_f}}{\#\Lambda^{W_f}} > 0.$$

Hence applying Theorems 3.4 and 3.5 to f finishes the proof. \square

Proposition 3.8. *Let $S \subset R$ be a finite subset. For any subgroup $V_f < W_f$ of finite index,*

$$\tau^S(B_T \cap \text{Ker}(\Lambda^{W_f}) \cap \mathbf{G}^S) \sim [\text{Ker}(\Lambda^{W_f}) : \text{Ker}(\Lambda^{V_f})] \cdot \tau^S(B_T \cap \text{Ker}(\Lambda^{V_f}) \cap \mathbf{G}^S).$$

Proof. For $\chi \in \Lambda$, set

$$d(\chi, s) := \int_{\mathbf{G}^S} H_\iota^{-s}(g) \chi(g) d\tau^S(g).$$

Note that if $\chi \in \Lambda^{V_f} - \Lambda^{W_f}$ then $\chi(w) \neq 1$ for some $w \in W_f$. Since H_ι is W_f -invariant,

$$d(\chi, s) = \int_{\mathbf{G}^S} H_\iota^{-s}(wg) \chi(wg) d\tau^S(g) = \chi(w) \cdot d(\chi, s)$$

and hence $d(\chi, s) = 0$. Therefore

$$\mathcal{Z}_{V_f}^S(s) = \sum_{\chi \in \Lambda^{V_f}} d(\chi, s) = \sum_{\psi \in \Lambda^{W_f}} d(\chi, s) = \mathcal{Z}_{W_f}^S(s)$$

This in particular implies that $c_{S, W_f} = c_{S, V_f}$ which were defined in Theorem 3.7. Now the claim follows from Theorem 3.7 and Lemma 3.1(2). \square

Proposition 3.9. *In the notation of Theorem 1.7, let \mathcal{L} and \mathcal{L}' be metrizations of the line bundle L and (B_T, W_f) , and (B'_T, W'_f) be defined as above with respect to \mathcal{L} and \mathcal{L}' respectively. Then*

$$\lim_{T \rightarrow \infty} \frac{\tau(B_T \cap \text{Ker}(\Lambda^{W_f}))}{\tau(B'_T \cap \text{Ker}(\Lambda^{W'_f}))} = \frac{[\mathbf{G}(\mathbb{A}) : \text{Ker}(\Lambda^{W_f})] \cdot \tau_{\mathcal{L}}(\mathbf{G}(\mathbb{A}))}{[\mathbf{G}(\mathbb{A}) : \text{Ker}(\Lambda^{W'_f})] \cdot \tau_{\mathcal{L}'}(\mathbf{G}(\mathbb{A}))}$$

where τ is a Haar measure on $\mathbf{G}(\mathbb{A})$.

Proof. Let $V_f = W_f \cap W'_f$. By Proposition 3.8, it suffices to show that

$$(3.10) \quad \lim_{T \rightarrow \infty} \frac{\tau(B_T \cap \text{Ker}(\Lambda^{V_f}))}{\tau(B'_T \cap \text{Ker}(\Lambda^{V_f}))} = \frac{\tau_{\mathcal{L}}(\mathbf{G}(\mathbb{A}))}{\tau_{\mathcal{L}'}(\mathbf{G}(\mathbb{A}))}.$$

Let S be a finite set such that $H_{\mathcal{L}, v} = H_{\mathcal{L}', v}$ for every $v \in R - S$. If we set $H_{\mathcal{L}, S} = \prod_{v \in S} H_{\mathcal{L}, v}$, then

$$(3.11) \quad \frac{\tau_{\mathcal{L}}(\mathbf{G}(\mathbb{A}))}{\tau_{\mathcal{L}'}(\mathbf{G}(\mathbb{A}))} = \frac{\int_{\mathbf{G}_S} H_{\mathcal{L}, S}(g)^{-\alpha_L} d\tau_S}{\int_{\mathbf{G}_S} H_{\mathcal{L}', S}(g)^{-\alpha_L} d\tau_S}.$$

Theorem 3.7 with $S = \emptyset$ and (3.11) imply that (3.10) stays the same when $H_{\mathcal{L},S}$ and $H_{\mathcal{L}',S}$ are replaced by constant multiples. Hence, we can assume that

$$(3.12) \quad H_{\mathcal{L},S}(e) = H_{\mathcal{L}',S}(e) = 1.$$

By the weak approximation theorem, the canonical projection map $\pi_S : \text{Ker}(\Lambda^{V_f}) \rightarrow \mathbf{G}_S$ is surjective. Denote by τ and τ^S the Haar measures on $\text{Ker}(\Lambda^{V_f})$ and $\text{ker}(\pi_S)$ respectively, so that the triple (τ, τ_S, τ^S) is compatible, i.e., for any $h \in C_c(\text{Ker}(\Lambda^{V_f}))$,

$$\int_{\text{Ker}(\Lambda^{V_f})} h d\tau = \int_{g_1 \in \mathbf{G}_S} \int_{g_2 \in \text{ker}(\pi_S)} h(g_1 g_2) d\tau^S(g_2) d\tau_S(g_1).$$

It now follows from the dominated convergence theorem that

$$\begin{aligned} \tau(B_T \cap \text{Ker}(\Lambda^{V_f})) &= \int_{g \in \mathbf{G}_S} \tau^S(B_{T H_{\mathcal{L},S}(g)^{-1}} \cap \text{Ker}(\Lambda^{V_f}) \cap \mathbf{G}^S) d\tau_S(g) \\ &\sim_T \left(\int_{g \in \mathbf{G}_S} H_{\mathcal{L},S}(g)^{-a_L} d\tau_S(g) \right) \cdot \tau^S(B_T \cap \text{Ker}(\Lambda^{V_f}) \cap \mathbf{G}^S). \end{aligned}$$

Similarly,

$$\tau(B'_T \cap \text{Ker}(\Lambda^{V_f})) \sim_T \left(\int_{g \in \mathbf{G}_S} H_{\mathcal{L}',S}(g)^{-a_L} d\tau_S(g) \right) \cdot \tau^S(B'_T \cap \text{Ker}(\Lambda^{V_f}) \cap \mathbf{G}^S).$$

Since by (3.12),

$$B_T \cap \mathbf{G}^S = B'_T \cap \mathbf{G}^S,$$

this finishes the proof. \square

4. EQUIDISTRIBUTION FOR SATURATED CASES

Let \mathbf{G} be a product of connected adjoint absolutely simple groups defined over K and ι be a faithful absolutely irreducible representation of \mathbf{G} . Recall the compact spaces X_l and $X(\mathbb{A})$ defined in Theorems 1.5 and 1.8 respectively. In the discussions below, we treat these two compact spaces simultaneously. For simplicity, we use the notation \tilde{X}_l to denote either of X_l and $X(\mathbb{A})$. Then \tilde{X}_l is of the form $\prod_v \tilde{X}_{l,v}$ where $\tilde{X}_{l,v}$ is a closed subset of $\mathbb{P}^l(K_v)$ for each $v \in R$ for a fixed l . We put $\tilde{X} = \tilde{X}_l$ for simplicity. Without loss of generality, we may consider $\mathbf{G}(K_v)$ as a subset of $\tilde{X}_{l,v}$ for each v and $\mathbf{G}(\mathbb{A})$ as a subset of \tilde{X} . Fix a height function $H_l = \prod_{v \in R} H_{l,v}$ on the associated adèle group $\mathbf{G}(\mathbb{A})$ relative to ι as in the definition 3.2.

Let $a_l \in \mathbb{Q}^+$ and b_l be as defined in (1.3). Let dg_v denote a Haar measure on $\mathbf{G}(K_v)$ for each $v \in R$. For a finite set $S \subset R$, we define $\mathbf{G}_S := \prod_{v \in S} \mathbf{G}(K_v)$ and $\tau_S := \prod_{v \in S} dg_v$.

By Lemma 3.3,

$$m_{l,v} := \int_{\mathbf{G}(K_v)} H_{l,v}(g_v)^{-a_l} dg_v < \infty.$$

and hence $\mu_{\iota,v} := m_{\iota,v}^{-1} H_{\iota,v}^{-a_\iota}(g_v) dg_v$ is a probability measure on each \tilde{X}_v . We denote by μ_ι the probability measure on \tilde{X} given by

$$\mu_\iota = \prod_{v \in R} \mu_{\iota,v}.$$

As before, we set

$$B_T := \{g \in \mathbf{G}(\mathbb{A}) : H_\iota(g) < T\}.$$

For a compact open subgroup W_f of $\mathbf{G}(\mathbb{A}_f)$ under which H_ι is bi-invariant, consider the subgroup $\text{Ker}(\Lambda^{W_f})$ of $\mathbf{G}(\mathbb{A})$ where Λ^{W_f} is the set of all automorphic characters in $L^2(\mathbf{G}(K) \backslash \mathbf{G}(\mathbb{A}))$ which are W_f -invariant, and let τ_{W_f} denote the normalized Haar measure on $\text{Ker}(\Lambda^{W_f})$ so that

$$\tau_{W_f}(\mathbf{G}(K) \backslash \text{Ker}(\Lambda^{W_f})) = 1.$$

In the following, fixing a compact open subgroup W_f of $\mathbf{G}(\mathbb{A}_f)$ under which H_ι is bi-invariant, we set

$$\mathbf{G}_{H_\iota} = \text{ker}(\Lambda^{W_f}) \quad \text{and} \quad \tau = \tau_{W_f}.$$

This section is devoted to a proof of the following:

Theorem 4.1. *Suppose that ι is saturated. Then for any $f \in C(\tilde{X})$,*

$$\lim_{T \rightarrow \infty} \frac{1}{\tau(B_T \cap \mathbf{G}_{H_\iota})} \sum_{g \in \mathbf{G}(K) : H_\iota(g) < T} f(g) = \int_{\tilde{X}} f d\mu_\iota \quad \text{as } T \rightarrow \infty.$$

Note that Theorem 4.1 implies Theorem 1.12 by taking $f = 1$. Combining Theorem 3.7 and Proposition 3.9, we deduce Theorems 1.2, 1.5, 1.7 and 1.8 for the saturated cases. The rate of convergence is obtained in the next section.

Lemma 4.2. *For any $f \in C(\tilde{X})$,*

$$\lim_{T \rightarrow \infty} \frac{1}{\tau(B_T \cap \mathbf{G}_{H_\iota})} \int_{B_T \cap \mathbf{G}_{H_\iota}} f d\tau = \int_{\tilde{X}} f d\mu_\iota$$

Proof. For a finite subset $S \subset R$, set

$$H_S := \prod_{v \in S} H_{\iota,v}; \quad \tilde{X}_S := \prod_{v \in S} \tilde{X}_{\iota,v}; \quad \mu_S := \prod_{v \in S} \mu_{\iota,v}.$$

Consider the canonical projection $\pi_S : \mathbf{G}_{H_\iota} \rightarrow \mathbf{G}_S$. By the weak approximation property, π_S is surjective and hence $\mathbf{G}_{H_\iota} / \text{ker}(\pi_S) \simeq \mathbf{G}_S$. Set $m_S := \prod_{v \in S} m_{\iota,v}$. Denote by τ^S the Haar measure on $\text{ker}(\pi_S)$ which is compatible with (τ, τ_S) , i.e., for any $h \in C_c(\mathbf{G}_{H_\iota})$,

$$\int_{\mathbf{G}_{H_\iota}} h d\tau = \int_{g_1 \in \mathbf{G}_S} \int_{g_2 \in \text{ker}(\pi_S)} h(g_1 g_2) d\tau^S(g_2) d\tau_S(g_1).$$

Denote by C_S the subset of $C(\tilde{X})$ consisting of f of the form

$$f((x_v)_{v \in R}) := f_S((x_v)_{v \in S}) \quad \text{for some } f_S \in C(\tilde{X}_S).$$

Note that for $f \in C_S$, we have

$$\mu_\iota(f) = \mu_S(f).$$

We claim that for $f \in C_S$, as $T \rightarrow \infty$,

$$(4.3) \quad \int_{B_T \cap \mathbf{G}_{H_\iota}} f d\tau \sim \mu_\iota(f) \cdot m_S \cdot \tau^S(\ker(\pi_S) \cap B_T).$$

By Theorem 3.7, there exists a constant $c > 0$ such that for any $g \in \mathbf{G}_S$ with $H_S(g) \leq T/2$,

$$\tau^S(\ker(\pi_S) \cap B_{T \cdot H_S(g)^{-1}}) \leq c H_S(g)^{-a_\iota} T^{a_\iota} (\log T H_S(g)^{-1})^{b_\iota - 1}.$$

With δ_0 as in Lemma 3.6, there exists $d > 0$ such that for any $g \in \mathbf{G}_S$ such that $T/2 \leq H_S(g) \leq T\delta_0^{-1}$

$$\tau^S(\ker(\pi_S) \cap B_{T \cdot H_S(g)^{-1}}) \leq \tau^S(\ker(\pi_S) \cap B_2) \leq d \cdot H_S(g)^{-a_\iota} T^{a_\iota}.$$

Setting

$$y_T(g) := \frac{\tau^S(\ker(\pi_S) \cap B_{T \cdot H_S(g)^{-1}})}{\tau^S(\ker(\pi_S) \cap B_T)},$$

we deduce that for some constant $C > 0$,

$$y_T(g) \leq C \cdot H_S(g)^{-a_\iota} \quad \text{for any } g \in \mathbf{G}_S \text{ with } H_S(g) \leq T\delta_0^{-1}.$$

Since $y_T \cdot \chi_{B_{T\delta_0^{-1}} \cap \mathbf{G}_S}$ is bounded above by a constant multiple of $H_S^{-a_\iota}$ which is in $L^1(\mathbf{G}_S)$, we may use the dominated convergence theorem to deduce that as $T \rightarrow \infty$,

$$\begin{aligned} \int_{B_T \cap \mathbf{G}_{H_\iota}} f d\tau &= \int_{g \in \mathbf{G}_S} f_S(g) \left(\int_{h \in \ker(\pi_S)} \chi_{B_T}(gh) d\tau^S \right) d\tau_S \\ &= \int_{g \in \mathbf{G}_S, H_S(g) \leq T\delta_0^{-1}} f_S(g) (\tau^S(\ker(\pi_S) \cap B_{T \cdot H_S(g)^{-1}})) d\tau_S \\ &\sim \int_{g \in \mathbf{G}_S} f_S(g) H_S(g)^{-a_\iota} d\tau_S \cdot \tau^S(B_T \cap \ker(\pi_S)) \\ &= \mu_S(f_S) \cdot m_S \cdot \tau^S(B_T \cap \ker(\pi_S)) \end{aligned}$$

proving the claim (4.3).

Since the constant function $1 \in C_S$ and $\mu_\iota(1) = 1$, in particular, we have for any finite $S \subset R$,

$$(4.4) \quad \tau(B_T \cap \mathbf{G}_{H_\iota}) \sim m_S \cdot \tau^S(\ker(\pi_S) \cap B_T) \quad \text{as } T \rightarrow \infty.$$

Therefore for any $f \in C_S$,

$$\int_{B_T \cap \mathbf{G}_{H_\iota}} f d\tau \sim \mu_\iota(f) \cdot \tau(B_T \cap \mathbf{G}_{H_\iota}) \quad \text{as } T \rightarrow \infty.$$

Let $f \in C(\tilde{X})$. Fix any $\epsilon > 0$. We can find a finite subset $S \subset R$ and $f^+, f^- \in C_S$ such that

$$f^- \leq f \leq f^+ \quad \text{and} \quad \|f^+ - f^-\|_\infty < \epsilon.$$

By (4.4)

$$\int_{B_T \cap \mathbf{G}_{H_i}} f^\pm d\tau \sim \mu_\iota(f^\pm) \cdot \tau(B_T \cap \mathbf{G}_{H_i}) \quad \text{as } T \rightarrow \infty.$$

Since $\mu(f^+ - f^-) \leq \epsilon$ and $\epsilon > 0$ is arbitrary, it is easy to deduce that

$$\int_{B_T \cap \mathbf{G}_{H_i}} f d\tau \sim \mu_\iota(f) \cdot \tau(B_T \cap \mathbf{G}_{H_i}) \quad \text{as } T \rightarrow \infty.$$

This finishes the proof. \square

Corollary 4.5. *Let V_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$ under which H_i is bi-invariant. Then*

- (1) $\tau(B_T \cap \ker(\mathbf{G}_{H_i})) \sim_{T \rightarrow \infty} \tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))$;
- (2) for any $f \in C(\tilde{X})$,

$$\int_{B_T \cap \mathbf{G}_{H_i}} f d\tau \sim_{T \rightarrow \infty} \int_{B_T \cap \ker(\Lambda^{V_f})} f d\tau_{V_f}.$$

Proof. It suffices to prove the claim for the case when $V_f \subset W_f$. Since the restriction of τ to $\ker(\Lambda^{V_f})$ is equal to $[\ker(\Lambda^{W_f}) : \ker(\Lambda^{V_f})] \cdot \tau_{V_f}$, the first claim follows from Proposition 3.8. The second claim follows from the claim (1) and Lemma 4.2. \square

For a fixed $f \in C(\tilde{X})$, we define a function F_T on $\mathbf{G}_{H_i} \times \mathbf{G}_{H_i}$ by

$$F_T(g, h) = \sum_{\gamma \in \mathbf{G}(K)} f(g^{-1}\gamma h) \cdot \chi_{B_T}(g^{-1}\gamma h)$$

Clearly F_T is well defined as a function on $Y \times Y$ where $Y = \mathbf{G}(K) \backslash \mathbf{G}_{H_i}$.

Note that

$$F_T(e, e) = \sum_{\gamma \in \mathbf{G}(K): H_i(\gamma) \leq T} f(\gamma).$$

Proposition 4.6 (Weak-convergence). *Suppose that ι is saturated. Let $f \in C(\tilde{X})$. For $i = 1, 2$, let $\alpha_i \in C(Y)$ be a W_f -invariant function and $\int_Y \alpha_i d\tau = 1$. If $\alpha(x, y) := \alpha_1(x)\alpha_2(y)$, then*

$$\lim_{T \rightarrow \infty} \frac{1}{\tau(B_T \cap \mathbf{G}_{H_i})} \int_{Y \times Y} F_T \cdot \alpha d(\tau \times \tau) = \lim_{T \rightarrow \infty} \frac{1}{\tau(B_T \cap \mathbf{G}_{H_i})} \int_{B_T \cap \mathbf{G}_{H_i}} f d\tau.$$

Proof. Observe that

$$\begin{aligned}
(4.7) \quad \langle F_T, \alpha \rangle_{Y \times Y} &= \int_{x \in Y} \int_{y \in Y} \left(\sum_{\gamma \in \mathbf{G}(K)} f(x^{-1}\gamma y) \chi_{B_T}(x^{-1}\gamma y) \right) \alpha_1(x) \alpha_2(y) d\tau(y) d\tau(x) \\
&= \int_{x \in Y} \int_{h \in \mathbf{G}_{H_\iota}} f(x^{-1}h) \chi_{B_T}(x^{-1}h) \alpha_1(x) \alpha_2(h) d\tau(h) d\tau(x) \\
&= \int_{g \in \mathbf{G}_{H_\iota}} f(g) \chi_{B_T}(g) \left(\int_{x \in Y} \alpha_1(x) \alpha_2(xg) d\tau(x) \right) d\tau(g) \\
&= \int_{g \in B_T \cap \mathbf{G}_{H_\iota}} f(g) \langle \alpha_1, g \cdot \alpha_2 \rangle d\tau(g)
\end{aligned}$$

Write $\mathbf{G} = \mathbf{G}_1 \cdots \mathbf{G}_m$ as a product of connected absolutely simple K -groups. Since the height function H_ι is proper, we have that $g \rightarrow \infty$ strongly in $\mathbf{G}(\mathbb{A})$ if and only if $H_\iota(g_i) \rightarrow \infty$ for each $i = 1, \dots, m$ where $g = g_1 \cdots g_m$, $g_i \in \mathbf{G}_i(\mathbb{A})$.

For $C > 0$, define

$$B^C := \{g_1 \cdots g_m \in \mathbf{G}(\mathbb{A}) : H_\iota(g_i) > C \text{ for each } i = 1, \dots, m\}.$$

By Theorem 2.22, for any given $\epsilon > 0$, there exists $C > 0$ such that

$$(4.8) \quad |\langle \alpha_1, g \cdot \alpha_2 \rangle - 1| < \epsilon \quad \text{for all } g \in B^C.$$

Hence

$$\begin{aligned}
&\left| \int_{g \in B_T \cap \mathbf{G}_{H_\iota}} f(g) \langle \alpha_1, g \cdot \alpha_2 \rangle d\tau(g) - \int_{g \in B_T \cap \mathbf{G}_{H_\iota}} f(g) d\tau(g) \right| \\
&< \max f \cdot (\|\alpha_1\| \cdot \|\alpha_2\| + 1) \cdot \tau((B_T - B^C) \cap \mathbf{G}_{H_\iota}) + \max f \cdot \tau(B_T \cap B^C \cap \mathbf{G}_{H_\iota}) \cdot \epsilon.
\end{aligned}$$

By the assumption of ι being saturated together with Theorem 3.7 we deduce that

$$\lim_{T \rightarrow \infty} \frac{\tau((B_T - B^C) \cap \mathbf{G}_{H_\iota})}{\tau(B_T \cap \mathbf{G}_{H_\iota})} = 0.$$

Since $\epsilon > 0$ is arbitrary, by (4.7), this proves the claim. \square

By Lemma 4.2, the following theorem implies Theorem 4.1:

Theorem 4.9. *Suppose that ι is saturated. For any $f \in C(\tilde{X})$, we have*

$$\lim_{T \rightarrow \infty} \frac{1}{\tau(B_T \cap \mathbf{G}_{H_\iota})} \sum_{g \in \mathbf{G}(K): H_\iota(g) < T} f(g) = \lim_{T \rightarrow \infty} \frac{1}{\tau(B_T \cap \mathbf{G}_{H_\iota})} \int_{B_T \cap \mathbf{G}_{H_\iota}} f d\tau.$$

Proof. Recall that $\mathbf{G}_{H_i} = \ker(\Lambda^{W_f})$ and $\tau = \tau_{W_f}$. It suffices to prove our theorem for non-negative functions $f \in C(\tilde{X})$. Fix $\epsilon > 0$. Let W_∞ be a symmetric neighborhood of e in \mathbf{G}_∞° such that

$$W_\infty B_T W_\infty \subset B_{(1+\epsilon)T} \quad \text{and} \quad B_{(1-\epsilon)T} \subset \bigcap_{g,h \in W_\infty} h B_T g \quad \text{for all } T > 1.$$

Let V_f be a co-finite subgroup of W_f such that

$$(4.10) \quad f(g^{-1}xh) - \epsilon \leq f(x) \leq f(g^{-1}xh) + \epsilon \quad \text{for all } x \in X \text{ and } g, h \in V_f.$$

By replacing W_∞ by a smaller one if necessary, we may assume that (4.10) holds for any $g, h \in W$ where $W := W_\infty \times V_f$. It follows that for any $T > \delta_0$ and for any $g, h \in W$,

$$(4.11) \quad F_{(1-\epsilon)T}^-(g, h) \leq F_T(e, e) \leq F_{(1+\epsilon)T}^+(g, h)$$

where

$$F_T^\pm(g, h) = \sum_{\gamma \in \mathbf{G}(K)} (f(g^{-1}\gamma h) \pm \epsilon) \cdot \chi_{B_T}(g^{-1}\gamma h).$$

Set $Y = \mathbf{G}(K) \setminus \ker(\Lambda^{V_f})$. Now let $\psi \in C_c(Y)$ be a non-negative V_f -invariant function such that $\text{supp}(\psi) \subset \mathbf{G}(K) \setminus \mathbf{G}(K)W$ and $\int_Y \psi d\tau_{V_f} = 1$. By integrating (4.11) over $Y \times Y$ against the function $\alpha(x, y) = \psi(x) \cdot \psi(y)$, we obtain

$$\langle F_{(1-\epsilon)T}^-, \alpha \rangle \leq F_T(e, e) \leq \langle F_{(1+\epsilon)T}^+, \alpha \rangle.$$

Note that Theorem 3.7 implies the following: there exist constants $a_\epsilon \geq 1$ and $b_\epsilon \leq 1$ tending to 1 as $\epsilon \rightarrow 0$ such that for all sufficiently small $\epsilon > 0$,

$$(4.12) \quad b_\epsilon \leq \liminf_T \frac{\tau_{V_f}(B_{(1-\epsilon)T} \cap \ker(\Lambda^{V_f}))}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))} \leq \limsup_T \frac{\tau_{V_f}(B_{(1+\epsilon)T} \cap \ker(\Lambda^{V_f}))}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))} \leq a_\epsilon.$$

Hence by applying Proposition 4.6,

$$\begin{aligned} & \limsup_T \frac{F_T(e, e)}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))} \leq \limsup_T \frac{\langle F_{(1+\epsilon)T}^+, \alpha \rangle}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))} \\ & \leq \limsup_T \frac{\langle F_{(1+\epsilon)T}^+, \alpha \rangle}{\tau_{V_f}(B_{(1+\epsilon)T} \cap \ker(\Lambda^{V_f}))} \cdot \limsup_T \frac{\tau_{V_f}(B_{(1+\epsilon)T} \cap \ker(\Lambda^{V_f}))}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))} \\ & \leq a_\epsilon \cdot \limsup_T \frac{\int_{B_T \cap \ker(\Lambda^{V_f})} (f + \epsilon)}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))} \leq a_\epsilon \cdot \left(\limsup_T \frac{\int_{B_T \cap \ker(\Lambda^{V_f})} f}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))} + \epsilon \right) \\ & \leq a_\epsilon \cdot \left(\limsup_T \frac{\int_{B_T \cap \ker(\Lambda^{W_f})} f}{\tau(B_T \cap \ker(\Lambda^{W_f}))} + \epsilon \right) \end{aligned}$$

by Corollary 4.5, and similarly,

$$b_\epsilon \cdot \left(\liminf_T \frac{\int_{B_T \cap \ker(\Lambda^{W_f})} f}{\tau(B_T \cap \ker(\Lambda^{W_f}))} - \epsilon \right) \leq \liminf_T \frac{F_T(e, e)}{\tau_{V_f}(B_T \cap \ker(\Lambda^{V_f}))}.$$

Taking $\epsilon \rightarrow 0$, this implies by Corollary 4.5 that

$$\lim_T \frac{F_T(e, e)}{\tau(B_T \cap \mathbf{G}_{H_\epsilon})} = \lim_T \frac{\int_{B_T \cap \mathbf{G}_{H_\epsilon}} f d\tau}{\tau(B_T \cap \mathbf{G}_{H_\epsilon})}.$$

□

Fix a compact open subgroup $W_f \subset \mathbf{G}(\mathbb{A}_f)$. Define an increasing sequence $\{B_T\}$ of relatively compact subsets of $\mathbf{G}(\mathbb{A})$ to be W_f -well rounded if the following holds:

- (1) $W_f B_T W_f = B_T$ for any T
- (2) for any small $\epsilon > 0$, there exists a neighborhood $W_\epsilon \subset \mathbf{G}_\infty^\circ$ such that

$$W_\epsilon B_T W_\epsilon \subset B_{(1+\epsilon)T} \quad \text{and} \quad B_{(1-\epsilon)T} \subset \cap_{g, h \in W_\epsilon} g B_T h$$

for all T

- (3) $\tau_{W_f}(B_T \cap \ker(\Lambda^{W_f})) \rightarrow \infty$ as $T \rightarrow \infty$ and there exist constants $a_\epsilon \geq 1$ and $b_\epsilon \leq 1$ tending to 1 as $\epsilon \rightarrow 0$ such that for all sufficiently small $\epsilon > 0$,

$$b_\epsilon \leq \liminf_T \frac{\tau_{W_f}(B_{(1-\epsilon)T} \cap \ker(\Lambda^{W_f}))}{\tau_{W_f}(B_T \cap \ker(\Lambda^{W_f}))} \leq \limsup_T \frac{\tau_{W_f}(B_{(1+\epsilon)T} \cap \ker(\Lambda^{W_f}))}{\tau_{W_f}(B_T \cap \ker(\Lambda^{W_f}))} \leq a_\epsilon.$$

With this definition, we can easily deduce the following from the proof of Theorem 4.9:

Proposition 4.13. *Suppose that ι is saturated. Let W_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$. Then for any W_f -well rounded sequence $\{B_T\}$ of relatively compact subsets of $\mathbf{G}(\mathbb{A})$,*

$$\#\mathbf{G}(K) \cap B_T \sim_{T \rightarrow \infty} \tau_{W_f}(B_T \cap \ker(\Lambda^{W_f})).$$

Define a proper function $H : \mathbf{G}(\mathbb{A}) \rightarrow \mathbb{R}^+$ to be W_f -well rounded if the following properties hold:

- (1) H is bi- W_f -invariant;
- (2) there exists $b > 0$ such that for any small $\epsilon > 0$, there exists a symmetric neighborhood $W_\epsilon \subset \mathbf{G}_\infty^\circ$ of e such that

$$(1 - b\epsilon)H(x) \leq H(gxh) \leq (1 + b\epsilon)H(x) \quad \text{for any } g, h \in W_\epsilon \text{ and } x \in \mathbf{G}(\mathbb{A});$$
- (3) there exist $a = a(H) > 0$ and $b = b(H) \geq 1$ such that the associated zeta function

$$\mathcal{Z}_{W_f}^S(s) := \sum_{\chi \in \Lambda^{W_f}} \int_{\mathbf{G}(\mathbb{A})} H(g)^{-s} \chi(g) d\tau(g)$$

has a meromorphic continuation to $\text{Re}(s) > a - \epsilon$ with the unique pole at $s = a$ of order b and of positive residue, and for some constants $\kappa \in \mathbb{R}$ and $k > 0$,

$$\left| \frac{(s - a)^b \mathcal{Z}_{W_f}^S(s)}{s^b} \right| \leq k \cdot |1 + \text{Im}(s)|^\kappa$$

for $\text{Re}(s) > a - \epsilon$.

By applying the above proposition to $B_T = \{g \in \mathbf{G}(\mathbb{A}) : H(g) < T\}$, we deduce:

Proposition 4.14. *Suppose that ι is saturated. Let W_f be a compact open subgroup of $\mathbf{G}(\mathbb{A}_f)$. Then for any W_f -well rounded function $H : \mathbf{G}(\mathbb{A}) \rightarrow \mathbb{R}^+$,*

$$\#\{g \in \mathbf{G}(K) : H(g) < T\} \sim_{T \rightarrow \infty} c \cdot T^{a(H)} (\log T)^{b(H)-1}$$

for some constant $c > 0$.

5. ARITHMETIC FIBRATIONS

In this section we prove Theorems 1.2 and 1.7 for a general case, that is, without the saturation assumption on ι . Let \mathbf{G} , ι and H_ι be as in the beginning of Section 4. Fix a compact open subgroup W_f of $\mathbf{G}(\mathbb{A}_f)$ under which H_ι is bi-invariant.

Let \mathbf{M} be the smallest connected normal K -subgroup of \mathbf{G} whose root system contains the set

$$\{\alpha \in \Delta : \frac{u_\alpha + 1}{m_\alpha} = a_\iota\}.$$

Note that ι is saturated if and only if $\mathbf{M} = \mathbf{G}$. There exists a connected normal K -subgroup \mathbf{N} of \mathbf{G} so that $\mathbf{G} = \mathbf{M}\mathbf{N}$ and $\mathbf{M} \cap \mathbf{N} = \{e\}$. Let $\pi : \mathbf{G} \rightarrow \mathbf{N}$ be the canonical projection. Note that any element of $\mathbf{G}(\mathbb{A})$ can be uniquely written as $g_1 g_2$ with $g_1 \in \mathbf{M}(\mathbb{A})$ and $g_2 \in \mathbf{N}(\mathbb{A})$.

As is well known, the restriction $\iota|_{\mathbf{M}}$ of ι is a direct sum of finitely many copies of an irreducible representation, say, ι' , of \mathbf{M} with the highest weight given by the restriction of that of ι . The definition of \mathbf{M} implies that ι' is saturated, $a_{\iota'} = a_\iota$ and $b_{\iota'} = b_\iota$. For $x \in \mathbf{N}(K)$, the function $g \mapsto H_\iota(gx)$ defines a height function on $\mathbf{M}(\mathbb{A})$ as in the definition 3.2. Hence Theorem 1.2 for the saturated cases implies that for each $x \in \mathbf{N}(K)$,

$$(5.1) \quad N_{\pi^{-1}(x)}(H_\iota, T) = \#\{g \in \mathbf{M}(K) : H_\iota(gx) < T\} \sim c_x \cdot T^{a_\iota} (\log T)^{b_\iota-1}$$

for some $c_x > 0$.

Noting that $N(H_\iota, T) = \sum_{x \in \mathbf{N}(K)} N_{\pi^{-1}(x)}(H_\iota, T)$, we prove the following:

Theorem 5.2. *Under the assumptions of Theorem 1.2, we have for some $\delta > 0$,*

$$(5.3) \quad N(H_\iota, T) = c(H_\iota) \cdot T^{a_\iota} (\log T)^{b_\iota-1} (1 + (\log T)^{-\delta})$$

where $c(H_\iota) := \sum_{x \in \mathbf{N}(K)} c_x < \infty$.

The rest of this section is devoted to the proof of the above theorem.

Lemma 5.4. *Let \mathbf{G}_1 and \mathbf{G}_2 be normal algebraic K -subgroups of \mathbf{G} with $\mathbf{G} = \mathbf{G}_1 \mathbf{G}_2$ and $\mathbf{G}_1 \cap \mathbf{G}_2 = \{e\}$. There exists $\kappa > 1$ such that for any $g_1 \in \mathbf{G}_1(\mathbb{A})$ and $g_2 \in \mathbf{G}_2(\mathbb{A})$,*

$$\kappa^{-1} \cdot H_\iota(g_1) H_\iota(g_2) \leq H_\iota(g_1 g_2) \leq \kappa \cdot H_\iota(g_1) H_\iota(g_2).$$

Proof. Let χ denote the highest weight of ι . Then there exists a finite subset $S \subset R$ such that for any $v \in R - S$,

$$\mathbf{G}(K_v) = U_v A_v^+ U_v \quad \text{and} \quad H_v(\iota(g)) = \chi(a) \quad \text{for } g = u_1 a u_2 \in \mathbf{G}(K_v)$$

where U_v and A_v^+ are defined as in the section 2. In particular, it follows that for each $v \in R - S$, and for any $g_1 \in \mathbf{G}_1(K_v)$ and $g_2 \in \mathbf{G}_2(K_v)$,

$$H_v(\iota(g_1 g_2)) = H_v(\iota(g_1)) H_v(\iota(g_2)).$$

On the other hand, for $v \in S$, $H_{\iota, v}$ is equivalent to χ in the sense that there exists $\kappa_v > 1$ such that

$$\kappa_v^{-1} \cdot \chi(a) \leq H_{\iota, v}(g) \leq \kappa_v \cdot \chi(a) \quad \text{for } g = u_1 a u_2 \in U_v A_v^+ \Omega_v U_v = \mathbf{G}(K_v).$$

This implies the lemma. \square

A key ingredient in deducing Theorem 5.2 is the following stronger version of (5.1):

Proposition 5.5. *There exist $\beta, \delta, d > 0$ such that for each $x \in \mathbf{N}(K)$ and for any $T \geq \beta \cdot H_\iota(x)$,*

$$|N_{\pi^{-1}(x)}(H_\iota, T) - c_x \cdot T^{a_\iota} (\log T)^{b_\iota - 1}| \leq d \cdot d_x \cdot T^{a_\iota} (\log T)^{b_\iota - 1 - \delta}$$

where $d_x = H_\iota(x)^{-a_\iota} (\log H_\iota(x))^{b_\iota - 1}$. Moreover $c_x = O(H_\iota(x)^{-a_\iota})$.

Proof. Let $\delta_0 > 0$ be such that $H_\iota(g) \geq \delta_0$ for $g \in \mathbf{G}(\mathbb{A})$ (Lemma 3.6). For each $x \in \mathbf{N}(\mathbb{A})$, we define a function H_ι^x on $\mathbf{M}(\mathbb{A})$ by

$$H_\iota^x(g) := H_\iota(gx), \quad g \in \mathbf{M}(\mathbb{A}).$$

It is easy to see that H_ι^x is a height function as in the definition 3.2 with respect to the representation ι' . Set

$$B_T^x = \{g \in \mathbf{M}(\mathbb{A}) : H_\iota^x(g) < T\}.$$

Since x commutes with $\mathbf{M}(\mathbb{A})$, the group $\mathbf{M}_{H_\iota^x}$ is independent of x , and we denote it by \mathbf{M}_{H_ι} . Let $Y = \mathbf{M}(K) \backslash \mathbf{M}_{H_\iota}$ and τ be the invariant probability measure on Y . For each $x \in \mathbf{N}(K)$, set

$$F_T^x(g, h) := \sum_{\gamma \in \mathbf{M}(K)} \chi_{B_T^x}(g^{-1} \gamma h), \quad g, h \in \mathbf{M}(\mathbb{A}).$$

We may consider F_T^x as a function on $Y \times Y$. Write $\mathbf{M} = \mathbf{M}_1 \cdots \mathbf{M}_r$ as a product of connected absolutely simple K -groups. For a collection of smooth $(W_f \cap \mathbf{M}_i(\mathbb{A}))$ -invariant functions $\psi_i \in C_c(\mathbf{M}_i(K) \backslash \mathbf{M}_i(\mathbb{A}) \cap \mathbf{M}_{H_\iota})$, $1 \leq i \leq r$, define $\psi \in C_c(Y)$ and $\alpha \in C_c(Y \times Y)$ by

$$\psi(z_1, \dots, z_r) := \prod_{i=1}^r \psi_i(z_i) \quad \text{and} \quad \alpha(y_1, y_2) := \psi(y_1) \psi(y_2).$$

Assume that $\int \psi_i d\tau_i = 1$ for each i where τ_i is the invariant probability measure on $\mathbf{M}_i(K) \setminus \mathbf{M}_i(\mathbb{A}) \cap \mathbf{M}_{H_i}$. We claim that for a sufficiently large $l \in \mathbb{N}$, independent of x , we have for any $x \in \mathbf{N}(K)$,

$$\langle F_T^x, \alpha \rangle_{Y \times Y} = c_x \cdot T^{a_\iota} (\log T)^{b_\iota - 1} + O(d_x \cdot C'_\psi \cdot T^{a_\iota} (\log T)^{b_\iota - 1 - \delta}).$$

where $C'_\psi = \max(1, \max_i \|\mathcal{D}^l \psi_i\|^{2r})$ for some large l and \mathcal{D} is the elliptic operator defined in (2.20).

As in the proof of Proposition 4.6, we derive that

$$\langle F_T^x, \alpha \rangle = \int_{g \in B_T^x \cap \mathbf{M}_{H_\iota}} \langle \psi, g \cdot \psi \rangle d\tau(g)$$

Note that

$$\begin{aligned} |\langle \psi, g \cdot \psi \rangle - 1| &= \left| \prod_{i=1}^r \langle \psi_i, g_i \cdot \psi_i \rangle - 1 \right| \\ &= \left| \sum_{i=1}^r \left(\prod_{j=1}^{i-1} \langle \psi_j, g_j \cdot \psi_j \rangle \right) (\langle \psi_i, g_i \cdot \psi_i \rangle - 1) \right| \\ &\leq r \cdot C_\psi \cdot \max_i |\langle \psi_i, g_i \cdot \psi_i \rangle - 1| \\ &= r \cdot C_\psi \cdot \max_i |\langle \psi_i - 1, g_i \cdot (\psi_i - 1) \rangle| \end{aligned}$$

where $C_\psi = \max(1, \max_i \|\psi_i\|^{2r-2})$. Since $\psi_i - 1 \in L_{00}^2(\mathbf{M}_i(K) \setminus \mathbf{M}_i(\mathbb{A}) \cap \mathbf{M}_{H_i})$ for each i , we deduce from Theorem 2.21 that

(5.6)

$$|\langle F_T^x, \alpha \rangle - \tau(B_T^x \cap \mathbf{M}_{H_\iota})| \leq 2r \cdot \left(\prod_i c_{W_f \cap \mathbf{M}_i(\mathbb{A})} \right) \cdot C'_\psi \cdot \int_{g=g_1 \cdots g_r \in B_T^x \cap \mathbf{M}_{H_\iota}} \max_i \tilde{\xi}_{\mathbf{M}_i}(g_i) d\tau(g)$$

where $C'_\psi = \max(1, \max_i \|\mathcal{D}^l \psi_i\|^{2r})$ for some large l .

Since $\tilde{\xi}_{\mathbf{M}_i} \leq \xi_{\mathbf{M}_i}^{1/2}$, it follows from Lemma 2.5 that there exist $m \in \mathbb{N}$ and $C_1 > 0$ such that for any $1 \leq i \leq r$,

$$\tilde{\xi}_{\mathbf{M}_i}(g_i) < C_1 \cdot H_\iota(g_i)^{-1/m} \quad \text{for any } g_i \in \mathbf{M}_i(\mathbb{A}).$$

Define a function on $\mathbf{M}(\mathbb{A})$ by

$$\tilde{H}(g_1 \cdots g_r) := \min_i H_\iota(g_i), \quad g_i \in \mathbf{M}_i(\mathbb{A}).$$

Let κ be as in Lemma 5.4 for $\mathbf{G}_1 = \mathbf{M}$ and $\mathbf{G}_2 = \mathbf{N}$ so that $B_T^x \subset B_{\kappa T \cdot H_\iota(x)^{-1}}$. It then follows from (5.6) that

$$(5.7) \quad |\langle F_T^x, \alpha \rangle - \tau(B_T^x \cap \mathbf{M}_{H_\iota})| < C_2 \cdot C'_\psi \cdot \int_{B_{\kappa T \cdot H_\iota(x)^{-1}} \cap \mathbf{M}_{H_\iota}} \tilde{H}(g)^{-1/m} d\tau(g)$$

for a constant $C_2 > 0$ independent of x .

Since ι' is saturated, for every proper normal K -subgroup \mathbf{L} of \mathbf{M} ,

$$\tau_{\mathbf{L}}(B_T \cap \mathbf{L}_{H_l}) \ll (\log T)^{-1} \tau(B_T \cap \mathbf{M}_{H_l})$$

where $\tau_{\mathbf{L}}$ is a Haar measure on $\mathbf{L}(\mathbb{A})$.

For each $C > 1$, set

$$B^C = \{g \in \mathbf{M}(\mathbb{A}) : \tilde{H}(g) > C\}.$$

Note that

$$(B_T - B^C) \cap \mathbf{M}_{H_l} \subset \cup_{i=1}^r \Omega_i$$

where $\Omega_i = \{g = g_1 \cdots g_r \in \mathbf{M}_{H_l} : H_l(g_i) \leq C, H_l(g) < T\}$. Now denoting by $\mathbf{L}^{(i)}$ the subgroup of \mathbf{M} generated by $\mathbf{M}_1, \dots, \mathbf{M}_{i-1}, \mathbf{M}_{i+1}, \dots, \mathbf{M}_r$, let $\kappa_i > 1$ be a constant as in Lemma 5.4 for $\mathbf{G}_1 = \mathbf{M}_i$ and $\mathbf{G}_2 = \mathbf{L}^{(i)}$. Then for any $C \gg 1$,

$$\begin{aligned} \tau(\Omega_i) &\leq \int_{H_l(g_i) < C} \tau_{\mathbf{L}^{(i)}(\mathbb{A})}(B_{\kappa_i \delta_0^{-1} T} \cap \mathbf{L}_{H_l}^{(i)}) d\tau_{\mathbf{M}_i}(g_i) \\ &\ll C^{a_i} (\log C)^{b_i-1} (\log T)^{-1} \tau(B_{\kappa_0 T} \cap \mathbf{M}_{H_l}) \end{aligned}$$

where $\kappa_0 = \max_i(\kappa_i \delta_0^{-1})$.

Hence for any $C \gg 1$ and $T \gg C$,

$$\tau((B_T - B^C) \cap \mathbf{M}_{H_l}) \ll C^{a_i} (\log C)^{b_i-1} (\log T)^{-1} \tau(B_{\kappa_0 T} \cap \mathbf{M}_{H_l})$$

Therefore

$$\begin{aligned} (5.8) \quad \int_{B_T \cap \mathbf{M}_{H_l}} \tilde{H}^{-1/m} d\tau &= \int_{B_T \cap B^C \cap \mathbf{M}_{H_l}} \tilde{H}^{-1/m} d\tau + \int_{(B_T - B^C) \cap \mathbf{M}_{H_l}} \tilde{H}^{-1/m} d\tau \\ &\ll (C^{-1/m} + \delta_0^{-1/m} \cdot C^{a_i} (\log C)^{b_i-1} (\log T)^{-1}) \cdot \tau(B_{\kappa_0 T} \cap \mathbf{M}_{H_l}) \\ &\ll (\log T)^{-\delta} \cdot \tau(B_{\kappa_0 T} \cap \mathbf{M}_{H_l}) \quad \text{for } C = (\log T)^{1/(2a_i)} \end{aligned}$$

for some $\delta > 0$. We now deduce from (5.7) and (5.8) that

$$(5.9) \quad \langle F_T^x, \alpha \rangle = \tau(B_T^x \cap \mathbf{M}_{H_l}) + O(C'_\psi \cdot (\log T)^{-\delta} \cdot \tau(B_{\kappa_0 \kappa T \cdot H_l(x)^{-1}} \cap \mathbf{M}_{H_l}))$$

for some $\delta > 0$.

For $S \subset R$ as in the proof of Lemma 5.4,

$$(5.10) \quad \tau(B_T^x \cap \mathbf{M}_{H_l}) = \int_{g \in B_{\delta_0^{-1} \kappa T \cdot H_l^{-1}(x)} \cap \mathbf{M}_S} \tau^S(B_{\kappa T \cdot H_l^{-1}(gx)} \cap \mathbf{M}_{H_l} \cap \mathbf{M}^S) d\tau_S(g)$$

where the triple (τ, τ_S, τ^S) is compatibly taken. Since by Theorem 3.7,

$$\tau^S(B_T \cap \mathbf{M}_{H_l} \cap \mathbf{M}^S) = c_{S, W_f} T^{a_i} (\log T)^{b_i-1} + O(T^{a_i} (\log T)^{b_i-2}) \quad \text{for } T \gg 1,$$

we deduce that

$$\begin{aligned} &\tau^S(B_{\kappa T H_l^{-1}(gx)} \cap \mathbf{M}_{H_l} \cap \mathbf{M}^S) \\ &= c \cdot (T \cdot H_l^{-1}(gx))^{a_i} (\log T)^{b_i-1} + O((T \cdot H_l^{-1}(gx))^{a_i} (\log H_l(gx))^{b_i-1} (\log T)^{b_i-2}). \end{aligned}$$

when $H_\iota(gx) \ll T/\delta_0$ and $c = c(S, W_f, \kappa) > 0$. Note that by Lemma 3.3,

$$\tau_S(B_{T \cdot H_\iota^{-1}(x)} \cap \mathbf{M}_S) \ll (TH_\iota^{-1}(x))^{a_\iota - \epsilon}.$$

Hence using Lemmas 3.3 and 5.4, we have

$$\int_{g \in \mathbf{M}_S} H_\iota(gx)^{-a_\iota} (\log H_\iota(gx))^{b_\iota - 1} d\tau_S(g) \ll (\log H_\iota(x))^{b_\iota - 1} H_\iota(x)^{-a_\iota}.$$

Therefore we deduce from (5.10) that for $T \gg H_\iota(x)$,

$$\tau(B_T^x \cap \mathbf{M}_{H_\iota}) = c_x T^{a_\iota} (\log T)^{b_\iota - 1} + O(d_x T^{a_\iota} (\log T)^{b_\iota - 2}),$$

where

$$(5.11) \quad c_x = c \cdot \int_{g \in \mathbf{M}_S} H_\iota(gx)^{-a_\iota} d\tau_S(g) \ll H_\iota(x)^{-a_\iota}.$$

Hence combining (5.9) and (5.10), we have for $T \gg H_\iota(x)$,

$$\langle F_T^x, \alpha \rangle = c_x \cdot T^{a_\iota} (\log T)^{b_\iota - 1} + O(d_x \cdot C'_\psi \cdot T^{a_\iota} (\log T)^{b_\iota - 1 - \delta}).$$

Denote by τ_∞ and τ_f Haar measures on \mathbf{G}_∞ and $\mathbf{G}(\mathbb{A}_f)$ respectively so that $\tau = \tau_\infty \times \tau_f$. Let ϕ_ϵ be a smooth symmetric nonnegative function on \mathbf{M}_∞ , which is a product $\prod_{i=1}^r \phi_{i,\epsilon}$ of smooth functions on the simple factors of \mathbf{M}_∞ , $\int_{\mathbf{M}_\infty} \phi_\epsilon d\tau_\infty = 1$ and $\text{supp}(\phi_\epsilon)$ is contained in the Riemannian ball at e in \mathbf{M}_∞ of radius ϵ , and for some $\rho > 0$, $\max_i \|\mathcal{D}^l \phi_{i,\epsilon}\|^{2r} \ll \epsilon^{-\rho}$ (see, for example, [GO]). By the definition of H_ι in 3.2, there exists $b > 0$ such that

$$\text{supp}(\phi_\epsilon) \cdot B_T^x \cdot \text{supp}(\phi_\epsilon) \subset B_{(1+b\epsilon)T}^x$$

for every $T > 1$ and $x \in \mathbf{N}(K)$.

Define

$$\psi_\epsilon(g) = \frac{1}{\tau_f(W_f)} \sum_{\gamma \in \mathbf{M}(K)} \phi_\epsilon(\gamma g_\infty) \cdot \chi_{W_f}(\gamma g_f), \quad g = g_\infty g_f \in \mathbf{M}_\infty \mathbf{M}(\mathbb{A}_f).$$

Define $\alpha_\epsilon(y_1, y_2) = \psi_\epsilon(y_1) \psi_\epsilon(y_2)$ for $(y_1, y_2) \in Y \times Y$. Then

$$\begin{aligned} N_{\pi^{-1}(x)}(H_\iota, T) &\leq \langle F_{(1+b\epsilon)T}^x, \alpha_\epsilon \rangle \\ &= c_x T^{a_\iota} (\log T)^{b_\iota - 1} + O(c_x \cdot \epsilon \cdot T^{a_\iota} (\log T)^{b_\iota - 1} + d_x \cdot \epsilon^{-\rho} T^{a_\iota} (\log T)^{b_\iota - 1 - \delta}). \end{aligned}$$

Setting $\epsilon = (\log T)^{-\delta/(\rho+1)}$, we derive the upper estimate for $N_{\pi^{-1}(x)}(H_\iota, T)$. The lower estimate is proved similarly. \square

Proof of Theorem 5.2 According to the choice of \mathbf{N} , for any simple root $\alpha \in \Delta$ whose restriction to \mathbf{N} is a root, we have

$$\frac{u_\alpha + 1}{m_\alpha} < a_\iota.$$

Hence it follows from Theorem 3.7 that $N_{\mathbf{N}}(H_\iota, T) = O(T^{a_\iota - \epsilon})$ for some $\epsilon > 0$. Since $c_x \ll H_\iota(x)^{-a_\iota}$ by (5.11) and $d_x = H_\iota(x)^{-a_\iota} (\log H_\iota(x))^{b_\iota - 1}$, it follows that

$$C(H_\iota) := \sum_{x \in \mathbf{N}(K)} c_x < \infty \quad \text{and} \quad \sum_{x \in \mathbf{N}(K)} d_x < \infty.$$

Let $\delta_0 = \inf\{H_\iota(g) : g \in \mathbf{G}(\mathbb{A})\} > 0$. Applying Lemma 5.4 for \mathbf{M} and \mathbf{N} with κ therein, we have

$$\begin{aligned} & \sum_{x \in \mathbf{N}(K): H_\iota(x) > \beta^{-1}T} N_{\pi^{-1}(x)}(H_\iota, T) \\ &= \#\{xy \in \mathbf{N}(K)\mathbf{M}(K) : H_\iota(x) > \beta^{-1}T, H_\iota(xy) < T\} \\ &\leq N_{\mathbf{M}}(H_\iota, \kappa\beta^{-1}) \cdot N_{\mathbf{N}}(H_\iota, \kappa T \delta_0^{-1}) \\ &= O(T^{a_\iota - \epsilon}). \end{aligned}$$

Now applying Proposition 5.5 with β, δ therein, since $\sum_{x \in \mathbf{N}(K)} d_x < \infty$,

$$\begin{aligned} & \sum_{x \in \mathbf{N}(K): H_\iota(x) \leq \beta^{-1}T} N_{\pi^{-1}(x)}(H_\iota, T) \\ &= \left(\sum_{x \in \mathbf{N}(K): H_\iota(x) \leq \beta^{-1}T} c_x \right) T^{a_\iota} (\log T)^{b_\iota - 1} + O(T^{a_\iota} (\log T)^{b_\iota - 1 - \delta}). \end{aligned}$$

Therefore as $T \rightarrow \infty$,

$$\begin{aligned} N(H_\iota, T) &= \sum_{x \in \mathbf{N}(K): H_\iota(x) \leq \beta^{-1}T} N_{\pi^{-1}(x)}(H_\iota, T) + O(T^{a_\iota - \epsilon}) \\ &= \left(\sum_{x \in \mathbf{N}(K): H_\iota(x) \leq \beta^{-1}T} c_x \right) T^{a_\iota} (\log T)^{b_\iota - 1} (1 + O((\log T)^{-\delta})). \end{aligned}$$

Since $\sum_{x \in \mathbf{N}(K): H_\iota(x) \leq \beta^{-1}T} c_x = C(H_\iota) + O(T^{-\epsilon})$, we have

$$N(H_\iota, T) = C(H_\iota) \cdot T^{a_\iota} (\log T)^{b_\iota - 1} (1 + O((\log T)^{-\delta}))$$

finishing the proof.

REFERENCES

- [BM] V. V. Batyrev and Y. I. Manin *Sur le nombre des points rationnels de hauteur bornée des variétés algébriques*, Math. Ann., 286, (1990), pp. 27–43.
- [BT1] V. V. Batyrev and Yu. Tschinkel *Manin's conjecture for toric varieties*, J. Algebraic Geometry 7, (1988), pp. 15–53.
- [BT2] V. V. Batyrev and Yu. Tschinkel *Height zeta functions of toric varieties*, Algebraic Geometry 5 (Manin's Festschrift) Journ. Math Sci 82, pp. 3220–3239 (1998).
- [BT3] V. V. Batyrev and Yu. Tschinkel *Tamagawa measure of polarized algebraic varieties*, Astérisque 251, 299–340 (1998).

- [Be] I. N. Bernstein *All reductive p -adic groups are of type I*, Funkcional. Anal. i Priložen 8 (1974), pp. 3–6, English translation: *Funct. Anal. Appl.* 8 (1974), pp. 91–93.
- [BK] M. Brion and S. Kumar *Frobenius Splitting Methods in Geometry and Representation theory* Progress in Math., 231, Birkhäuser.
- [BT] A. Borel and J. Tits *Groupes réductifs sur un corps local, I*, Publ. Math. IHES 41, (1972), pp. 5–252.
- [BS] M. Burger and P. Sarnak *Ramanujan duals II*, Invent. Math., 106, (1991), pp. 1–11.
- [CT1] A. Chambert-Loir, Yu. Tschinkel *Fonctions zeta des hauteurs des espaces fibrés*, Rational points on algebraic varieties, pp. 71–115, Progress in Math., 199 (2001) Birkhäuser.
- [CT2] A. Chambert-Loir, Yu. Tschinkel *On the distribution of points of bounded height on equivariant compactification of vector groups*, Inventiones Math 48 (2002) pp. 421–452.
- [Cl1] L. Clozel *Démonstration de la conjecture τ* , Invent. Math. 151 (2003) 297–328.
- [Cl2] L. Clozel *Changement de base pour les représentations tempérées des groupes réductifs réels*, Ann. Sci. Éc. Norm. Supér., IV. Sér. 15 (1982) pp. 45–115
- [COU] L. Clozel, H. Oh and E. Ullmo *Hecke operators and equidistribution of Hecke points*, Invent. Math., 144, (2001), pp. 327–351.
- [CU] L. Clozel and E. Ullmo *Equidistribution des points de Hecke*, Contribution to automorphic forms, geometry and number theory, Johns Hopkins Univ. Press, Baltimore, MD, 2004, pp. 193–254.
- [DP] C. De Concini and C. Procesi *Complete symmetric varieties*, in Invariant theory (Montecatini, 1982) LMN V 996, Springer, Berlin, 1983, pp 1–44.
- [DS] C. De Concini and T. A. Springer *Compactification of symmetric varieties*, Dedicated to the memory of Claude Chevalley Transform. Groups 4 (1999), pp. 273–300.
- [Di] J. Dixmier *Les C^* -algèbres et leurs représentations*, Gauthier-Villars, Paris 1964.
- [EM] A. Eskin and C. McMullen. *Mixing, counting and equidistribution on Lie groups*, Duke Math. J. 71 (1993), no. 1, pp. 181–209.
- [EO] A. Eskin and H. Oh *Ergodic theoretic proof of equidistribution of Hecke points*, Erg. The. Dyn. Sys. Vol 26 (2006), pp. 163–167.
- [FMT] J. Franke, Yu. I. Manin and Yu. Tschinkel *Rational points of bounded height on Fano varieties* Inventiones Math. 95 (1989) pp. 421–435.
- [GO] W. T. Gan and H. Oh *Equidistribution of Integer points on a family of homogeneous varieties: A problem of Linnik*, Compos. Math. 136 (2003) pp. 325–352.
- [GOS] A. Gorodnik, H. Oh and N. Shah *Integral points on symmetric varieties and Satake boundary*, Preprint.
- [GJ] S. Gelbart and H. Jacquet *A relation between automorphic representation of GL_2 and GL_3* , Ann. Sci. École Norm. Sup, 11 (1978), pp. 471–552.
- [HS] M. Hindry and J. Silverman *Diophantine geometry: An introduction*, Springer, GTM 201 (2000).
- [HM] R. Howe and C. C. Moore *Asymptotic properties of unitary representations*, J. Functional Analysis 32 (1979) pp. 72–96.
- [HT] R. Howe and E. C. Tan *Non-Abelian Harmonic Analysis*, Springer-Verlag, 1992
- [JL] H. Jacquet and P. Langlands *Automorphic forms on GL_2* , Lecture Notes in Mathematics, 114, Springer-Verlag, Berlin-New York, 1970.
- [Lu] A. Lubotzky *Discrete groups, expanding graphs and invariant measures*, Progress in Mathematics 125, (1994) Birkhäuser.
- [Mar] G. Margulis *On some aspects of the theory of Anosov Systems*, Springer Monographs in Mathematics (2004).

- [Mau] F. Maucourant *Homogeneous asymptotic limits of Haar measures of semisimple linear groups and their lattices* Preprint (2005).
- [Oh1] H. Oh *Uniform pointwise bounds for matrix coefficients of unitary representations and applications to Kazhdan constants*, Duke Math. J. 113 (2002) 133–192.
- [Oh2] H. Oh *Tempered subgroups and representations with minimal decay of matrix coefficients*, Bull. Soc. Math. France 126 (1998) pp. 355–380.
- [Pe1] E. Peyre *Hauteurs et mesures de Tamagawa sur les variétés de Fano*, Duke Math J. 79 (1995) pp. 101–218.
- [Pe2] E. Peyre *Points de hauteur bornée, topologie adélique et mesures de Tamagawa*, Les XXIIèmes Journées Arithmétiques (Lille, 2001) J. Théor. Nombres Bordeaux 15 (2003) pp. 319–349.
- [PR] V. Platonov and A. Rapinchuk *Algebraic groups and Number theory*, Academic Press, New York, (1994).
- [Ro] Rogawski *Automorphic representations of unitary groups in three variables*, Annals of Mathematics Studies, 123. Princeton University Press, Princeton, NJ (1990).
- [Sc] S. Schanuel *On heights in number fields*, Bull. Amer. Math. Soc 70, (1964) pp. 262–263.
- [ST1] J. Shalika and Y. Tschinkel *Height zeta functions of equivariant compactifications of the Heisenberg group*, in Contributions to automorphic forms, geometry and number theory, Johns Hopkins Univ. Press, Baltimore, MD 2004, p. 743–771.
- [ST2] J. Shalika and Y. Tschinkel *Height zeta functions of equivariant compactification of unipotent groups*, In preparation
- [STT1] J. Shalika, R. Takloo-Bighash and Y. Tschinkel *Rational points and automorphic forms*, Contributions to Automorphic Forms, Geometry, and Number Theory (H. Hida, D. Ramakrishnan, F. Shahidi eds.), (2004), 733–742, Johns Hopkins University Press.
- [STT2] J. Shalika, R. Takloo-Bighash and Y. Tschinkel *Rational points on compactifications of semisimple groups*, Preprint.
- [Si] J. Silverman *The theory of Height functions*, in Arithmetic Geometry ed. G. Cornell and J. Silverman, Springer, 1986.
- [ST] M. Strauch and Y. Tschinkel *Height zeta functions of toric bundles over flag varieties*, Selecta Math 5 (1999) pp. 352–396.
- [Ti1] J. Tits *Classification of algebraic semisimple groups*, Algebraic groups and discontinuous groups. Symposium Colorado, Boulder (1965), AMS Proc. Symp. Pure Math., IX
- [Ti2] J. Tits *Reductive groups over local fields*, Proc. Symp. Pure Math., 33, (1979), pp. 29–70 in Automorphic forms, representations and L-functions (eds. A. Borel and W. Casselman).
- [Wa] G. Warner *Harmonic analysis on semisimple Lie groups I*, Springer-Verlag (1972).
- [We] A. Weil *Adeles and algebraic groups*, Boston, Birkhäuser, 1982.
- [Zi] R. Zimmer *Ergodic theory and semisimple groups*, Boston, Birkhäuser, 1984

MATHEMATICS 253-37, CALTECH, PASADENA, CA 91125, USA // IRMAR-CAMPUS DE BEAULIEU,
 35042 RENNES CEDEX, FRANCE // MATHEMATICS 253-37, CALTECH, PASADENA, CA 91125,
 USA

E-mail address: gorodnik@its.caltech.edu

E-mail address: francois.maucourant@math.univ-rennes1.fr

E-mail address: heeoh@its.caltech.edu