

On the concave and convex solutions of mixed convection boundary layer approximation in a porous medium

BERNARD BRIGHI† and JEAN-DAVID HOERNEL‡

Université de Haute-Alsace, Laboratoire de Mathématiques et Applications
4 rue des frères Lumière, 68093 MULHOUSE (France)

Abstract

In this paper we investigate the similarity solutions of a plane mixed convection boundary layer flow near a semi-vertical plate, with a prescribed power law function of the distance from the leading edge for the temperature, that is embedded in a porous medium. We show the existence and uniqueness of convex and concave solutions for positive values of the power law exponent.

1 Introduction

We look at the nonlinear autonomous differential equation established in [1]

$$f''' + (1 + \lambda)ff'' + 2\lambda(1 - f')f' = 0 \quad (1)$$

on $[0, \infty)$ with the boundary conditions

$$f(0) = \alpha, \quad f'(0) = \beta \quad \text{and} \quad f'(\infty) = 1 \quad (2)$$

where $\beta > 0$ and $f'(\infty) := \lim_{t \rightarrow \infty} f'(t)$.

The problem (1)-(2) with $\alpha = 0$ and $\beta = 1 + \varepsilon$, $-1 < \varepsilon < \frac{1}{2}$ is considered in [1] and comes from the study of the mixed convection boundary-layer flow along a semi-infinite vertical plate embedded in a saturated porous medium. We take a rectangular Cartesian co-ordinate system with the origin fixed at the leading edge of the vertical plate, the x -axis directed upward along the plate and the y -axis normal to it and we write the

AMS 2000 Subject Classification: 34B15, 34C11, 76D10.

Key words and phrases: Boundary layer, similarity solution, third order nonlinear differential equation, boundary value problem.

† bernard.brighi@uha.fr ‡ j-d.hoernel@wanadoo.fr

temperature on the plate as $T(x, 0) = T_\infty + Ax^\lambda$ with $A > 0$, $\lambda \in \mathbb{R}$ and where T_∞ denotes the temperature far from the wall. The fluid velocity at the edge of the boundary layer is written as Bx^λ with $B > 0$ and we suppose that there is no normal velocity on the plate. Then, for large Péclet numbers, one can compute the stream function and the temperature of the fluid in the porous medium using the dimensionless similarity function f given by (1)-(2). For more details on the derivation of the model and numerical results, see [1] and [2].

The solution of (1)-(2) depends on two parameters, the power law exponent λ and the mixed convection parameter $\varepsilon = \frac{Ra}{Pe}$ with Ra the Rayleigh number and Pe the Péclet number. For $\varepsilon = 0$ we have forced convection, $\varepsilon > 0$ corresponds to aiding mixed convection and $\varepsilon < 0$ to opposing mixed convection.

For $\lambda = 0$ (isothermal surface), equation (1) reduces to the Blasius equation (see [3]) and is studied with conditions (2) in [4] for $\alpha \geq 0$ and $\beta > 0$, in [5] for $\alpha \in \mathbb{R}$ and $\beta \in (0, 1)$ and in [6] for $\alpha \in \mathbb{R}$ and $\beta > 1$. Blasius equation is also a particular case of the Falkner-Skan equation $f''' + ff'' + \lambda(1 - f'^2) = 0$ (see [5]).

The equivalent problem of free convection over a vertical flat plate embedded in a fluid saturated porous medium is studied in [7], [8], [9] and [10].

In [11] one can find interesting new results about the problem (1)-(2) for $\lambda < 0$ and $\alpha = 0$, in particular that there are infinitely many solutions for $-1 < \lambda < 0$ and $0 < \beta < \frac{3}{2}$, no solutions for $\lambda = -1$ (except the trivial one $f(t) = t$ for $\beta = 1$) and no nonnegative solutions for $\lambda < -1$ and $\beta > \frac{3}{2}$.

In this note, we are interested in the case $\lambda > 0$, and prove existence and uniqueness of the solution of (1)-(2) whose second derivative does not vanish.

2 Existence and uniqueness results

First, let us suppose that f satisfies equation (1) on some interval I and denote by F any anti-derivative of f on I , then we have

$$(f''e^{(1+\lambda)F})' = -2\lambda(1 - f')f'e^{(1+\lambda)F}. \quad (3)$$

As for the Falkner-Skan equation, and contrary to what occurs in [9], the relation (3) shows that oscillatory solutions may exist (see [12]). Nevertheless, for the problem (1)-(2) we could expect convex solutions (i.e. such that $f'' > 0$) if $0 < \beta < 1$, and concave solutions (i.e. such that $f'' < 0$) if $\beta > 1$. Before proving this, let us look briefly at the case $\lambda > 0$ and $\beta = 1$; the function $g(t) = t + \alpha$ is then a solution of (1)-(2). Let f be another solution with $f''(0) = \gamma$ and assume first that $\gamma > 0$. Then, since $f'(0) = f'(\infty) = 1$, there exists $t_0 > 0$ such that $f'(t_0) > 1$, $f''(t_0) = 0$ and $f'''(t_0) \leq 0$. However, from (1) we obtain $f'''(t_0) = -\lambda f'(t_0)(1 - f'(t_0)) > 0$ and thus a contradiction. If $\gamma < 0$, the same approach does not allow us to produce a contradiction, but leads to $f'(t_0) < 0$, and thus g is the unique increasing solution of (1)-(2) when $\lambda > 0$ and $\beta = 1$. Clearly, at this stage, we cannot exclude that oscillatory solutions exist.

In order to obtain solutions of (1)-(2), we consider the following initial value problem:

$$\begin{cases} f''' + (1 + \lambda)ff'' + 2\lambda(1 - f')f' = 0, \\ f(0) = \alpha, \\ f'(0) = \beta, \\ f''(0) = \gamma \end{cases} \quad (4)$$

with $\lambda > 0$, $\alpha \in \mathbb{R}$, $\beta > 0$ and a suitable γ . We denote the solution by f_γ and by $[0, T_\gamma)$ its right maximal interval of existence. Next, integrating (1) on $[0, t]$ for $0 < t < T_\gamma$, we obtain the useful identity

$$f_\gamma''(t) - \gamma + (1 + \lambda)(f_\gamma(t)f_\gamma'(t) - \alpha\beta) + 2\lambda(f_\gamma(t) - \alpha) = (3\lambda + 1) \int_0^t f_\gamma'(s)^2 ds. \quad (5)$$

We also need the following lemma.

Lemma 1 *If f is a solution of (1) on $[0, T_\gamma)$ such that there exists a point t_0 satisfying $f''(t_0) = 0$ and $f'(t_0) = 1$, then $f''(t) = 0$ for every $t \in [0, T_\gamma)$.*

Proof. Let f be a solution of (1) on $[0, T_\gamma)$ such that $f''(t_0) = 0$ and $f'(t_0) = 1$ for some $t_0 \in [0, T_\gamma)$. Since the function $g(t) = t - t_0 + f(t_0)$ is a solution of (1) such that $g(t_0) = f(t_0)$, $g'(t_0) = f'(t_0)$ and $g''(t_0) = f''(t_0)$, we obtain $g = f$ and $f'' \equiv 0$. ■

Theorem 1 *Let $\lambda > 0$. For $\alpha \in \mathbb{R}$ and $0 < \beta < 1$ problem (1)-(2) admits a unique convex solution.*

Proof of existence. Let f_γ be a solution of the initial value problem (4) with $0 < \beta < 1$ and $\gamma \geq 0$. We notice that $f_\gamma(t)$ exists as long as we have $f_\gamma'' > 0$ and $f_\gamma' < 1$. From Lemma 1, f_γ'' cannot vanish at a point where $f_\gamma' = 1$. Therefore, it follows that there are only three possibilities:

- (a) f_γ'' becomes negative from a point such that $f_\gamma' < 1$,
- (b) f_γ' takes the value 1 at some point for which $f_\gamma'' > 0$,
- (c) we always have $0 < f_\gamma' < 1$ and $f_\gamma'' > 0$.

As $f_0'(0) = \beta < 1$, $f_0''(0) = 0$ and $f_0'''(0) = 2\lambda\beta(\beta - 1) < 0$, we have that f_0 is of type (a), and by continuity it must be so for f_γ with $\gamma > 0$ small enough.

On the other hand, as long as $f_\gamma''(t) > 0$ and $f_\gamma'(t) \leq 1$, we have $f_\gamma(t) \leq t + \alpha$, and (5) leads to

$$\begin{aligned} f_\gamma''(t) &\geq -(1 + \lambda)f_\gamma(t)f_\gamma'(t) - 2\lambda f_\gamma(t) + (1 + \lambda)\alpha\beta + 2\lambda\alpha + \gamma \\ &\geq -(3\lambda + 1)t + (1 + \lambda)(\alpha\beta - |\alpha|) + \gamma \end{aligned}$$

and integrating once again we have

$$f_\gamma'(t) \geq -\frac{3\lambda + 1}{2}t^2 + ((1 + \lambda)(\alpha\beta - |\alpha|) + \gamma)t + \beta. \quad (6)$$

Hence, for γ large enough, the polynomial on the right of (6) takes values greater than 1. Therefore, for such a γ , there exists t_0 such that $f_\gamma'(t_0) = 1$ and $f_\gamma''(t) > 0$ for $t \leq t_0$, and f_γ is of type (b).

Defining $A = \{\gamma > 0; f_\gamma \text{ is of type (a)}\}$ and $B = \{\gamma > 0; f_\gamma \text{ is of type (b)}\}$ we have that $A \neq \emptyset$, $B \neq \emptyset$ and $A \cap B = \emptyset$. Both A and B are open sets, so there exists a $\gamma_* > 0$ such that the solution f_{γ_*} of (4) is of type (c) and is defined on the whole interval $[0, \infty)$. For this solution we have that $0 < f'_{\gamma_*} < 1$ and $f''_{\gamma_*} > 0$ which implies that $f'_{\gamma_*}(t) \rightarrow l \in (0, 1]$ as $t \rightarrow \infty$. Suppose that $l \neq 1$, as f'_{γ_*} is increasing we have $f'_{\gamma_*} \leq l$ and

$$f'''_{\gamma_*}(t) \leq -(1 + \lambda)\alpha f''_{\gamma_*}(t) - 2\lambda(1 - l)f'_{\gamma_*}(t).$$

Integrating this inequality leads to

$$f''_{\gamma_*}(t) \leq \gamma_* - (1 + \lambda)\alpha(f'_{\gamma_*}(t) - \beta) - 2\lambda(1 - l)(f_{\gamma_*}(t) - \alpha)$$

and, as $f'_{\gamma_*}(t) \rightarrow l < 1$ and $f_{\gamma_*}(t) \rightarrow \infty$ as $t \rightarrow \infty$, we obtain a contradiction with the positivity of f''_{γ_*} . ■

Proof of uniqueness. Let f be a convex solution of (1)-(2). As $f(t) > 0$ for t large enough, we have that $f'''(t) < 0$ for t large enough. Then $f''(t) \rightarrow 0$ as $t \rightarrow \infty$.

As f' and f'' are positive, we can define a function $v : [\beta^2, 1) \rightarrow [\alpha, \infty)$ such that

$$\forall t \geq 0, \quad v(f'(t)^2) = f(t).$$

Setting $y = f'(t)^2$ leads to

$$f(t) = v(y), \quad f''(t) = \frac{1}{2v'(y)} \quad \text{and} \quad f'''(t) = -\frac{v''(y)\sqrt{y}}{2v'(y)^3} \quad (7)$$

and using (1) we obtain

$$\forall y \in [\beta^2, 1), \quad v'' = (1 + \lambda)\frac{vv'^2}{\sqrt{y}} + 4\lambda(1 - \sqrt{y})v'^3 \quad (8)$$

with

$$v(\beta^2) = v(f'(0)^2) = \alpha, \quad v(1) = \lim_{t \rightarrow \infty} f(t) = \infty \quad \text{and} \quad v'(\beta^2) = \frac{1}{2\gamma} > 0.$$

Suppose now that there are two convex solutions f_1 and f_2 of (1)-(2) with $f''_i(0) = \gamma_i > 0$, $i \in \{1, 2\}$ and $\gamma_1 > \gamma_2$. They give v_1, v_2 solutions of equation (8) defined on $[\beta^2, 1)$ such that

$$v_1(\beta^2) = v_2(\beta^2) = \alpha, \quad v'_1(\beta^2) = \frac{1}{2\gamma_1} \quad \text{and} \quad v'_2(\beta^2) = \frac{1}{2\gamma_2}.$$

Let $w = v_1 - v_2$, we have $w(\beta^2) = 0$ and $w'(\beta^2) < 0$. If w' vanishes, there exists an x in $[\beta^2, 1)$ such that $w'(x) = 0$, $w''(x) \geq 0$ and $w(x) < 0$. But from (8) we then obtain

$$w''(x) = (1 + \lambda)\frac{v'_1(x)^2}{\sqrt{x}}w(x) < 0$$

and this is a contradiction. Therefore $w' < 0$ and $w < 0$ on $[\beta^2, 1)$. Set now $V_i = \frac{1}{v'_i}$ for $i \in \{1, 2\}$ and $W = V_1 - V_2$. We have $W > 0$ and using (8) we obtain

$$W'(y) = -(1 + \lambda)\frac{w(y)}{\sqrt{y}} - 4\lambda(1 - \sqrt{y})w'(y) > 0.$$

But, using (7) we have $V_i(f'(t)^2) = 2f''(t)$ and thus $V_i(y) \rightarrow 0$ as $y \rightarrow 1$. Hence $W(y) \rightarrow 0$ as $y \rightarrow 1$, a contradiction. ■

Theorem 2 Let $\lambda > 0$. For $\alpha \in \mathbb{R}$ and $\beta > 1$, problem (1)-(2) admits a unique concave solution.

Proof of existence. Let f_γ be a solution of the initial value problem (4) with $\beta > 1$ and $\gamma \leq 0$. As long as we have $f'_\gamma > 1$ and $f''_\gamma < 0$, then f_γ exists. Because of Lemma 1, there are only three possibilities

- (a) f''_γ becomes positive from a point such that $f'_\gamma > 1$,
- (b) f'_γ takes the value 1 at some point for which $f''_\gamma < 0$,
- (c) we always have $1 < f'_\gamma$ and $f''_\gamma < 0$.

As $f'_0(0) = \beta > 1$, $f''_0(0) = 0$ and $f'''_0(0) = 2\lambda\beta(\beta - 1) > 0$ we have that $f'_0(t) > 1$ and $f''_0(t) > 0$ on some interval $[0, t_0)$. Then, by continuity for small values of $-\gamma$ we have that f''_γ becomes positive at some point with $f'_\gamma > 1$ and f_γ is of type (a).

As long as $f''_\gamma(t) < 0$ and $f'_\gamma(t) \geq 1$, we have $f_\gamma(t) \geq \alpha$ and (5) leads to

$$f''_\gamma(t) \leq \gamma + (1 + \lambda)\alpha\beta + (3\lambda + 1)\beta^2 t + |\alpha|(1 + \lambda)\beta.$$

Integrating once again we have

$$f'_\gamma(t) \leq \frac{3\lambda + 1}{2}\beta^2 t^2 + (\gamma + (1 + \lambda)\alpha\beta + |\alpha|(1 + \lambda)\beta)t + \beta.$$

Hence, for $-\gamma$ large enough, there exists t_0 such that $f'_\gamma(t_0) = 1$ and $f''_\gamma(t) < 0$ for $t \leq t_0$, and f_γ is of type (b).

Defining $A = \{\gamma < 0; f_\gamma \text{ is of type (a)}\}$ and $B = \{\gamma < 0; f_\gamma \text{ is of type (b)}\}$ we have that $A \neq \emptyset$, $B \neq \emptyset$ and $A \cap B = \emptyset$. Both A and B are open sets, so there exists a $\gamma_* < 0$ such that the solution f_{γ_*} of (4) is of type (c) and is defined on the whole interval $[0, \infty)$. For this solution we have that $f'_{\gamma_*} > 1$ and $f''_{\gamma_*} < 0$ which implies that $f'_{\gamma_*} \rightarrow l \geq 1$ as $t \rightarrow \infty$. Suppose that $l \neq 1$, as f'_{γ_*} is decreasing we have $f'_{\gamma_*} \geq l$ and

$$f'''_{\gamma_*}(t) \geq -(1 + \lambda)\alpha f''_{\gamma_*}(t) - 2\lambda(1 - f'_{\gamma_*}(t))f'_{\gamma_*}(t).$$

Integrating this inequality leads to

$$f''_{\gamma_*}(t) \geq \gamma_* - (1 + \lambda)\alpha(f'_{\gamma_*}(t) - \beta) - 2\lambda(1 - l)(f_{\gamma_*}(t) - \alpha)$$

and, as $f'_{\gamma_*}(t) \rightarrow l > 1$ and $f_{\gamma_*}(t) \rightarrow \infty$ as $t \rightarrow \infty$, we obtain a contradiction with the negativity of f''_{γ_*} . ■

Proof of uniqueness. Let f_1 and f_2 be two concave solutions of (1)-(2) and let $\gamma_i = f''_i(0) < 0$, $i \in \{1, 2\}$ with $\gamma_1 > \gamma_2$. Writing $g = f_1 - f_2$, we have $g'(0) = 0$, $g'(\infty) = 0$ and $g''(0) > 0$. Hence there exists $t_0 > 0$ such that g' admits a positive maximum at t_0 , and this means

$$g'(t_0) > 0, g''(t_0) = 0 \text{ and } g'''(t_0) \leq 0.$$

From (1), and since $f'_i > 1$ and $f''_i < 0$, we obtain

$$g'''(t_0) = -(1 + \lambda)f''_1(t_0)g(t_0) - 2\lambda g'(t_0)(1 - f'_1(t_0) - f'_2(t_0)) > 0$$

and hence a contradiction. ■

References

- [1] E. H. Aly, L. Elliott & D. B. Ingham, Mixed convection boundary-layer flows over a vertical surface embedded in a porous medium, *Eur. J. Mech. B Fluids* **22**, pp. 529-543 (2003).
- [2] R. Nazar, N. Amin & I. Pop, Unsteady mixed convection boundary-layer flow near the stagnation point on a vertical surface in a porous medium, *Int. J. Heat Mass Transfer* **47**, pp. 2681-2688 (2004).
- [3] H. Blasius, Grenzschichten in Flüssigkeiten mit kleiner Reibung, *Z. Math. Phys.* **56**, pp. 1-37 (1908).
- [4] W. A. Coppel, On a differential equation of boundary layer theory, *Phil. Trans. Roy. Soc. London, Ser A* **253**, pp. 101-136 (1960).
- [5] P. Hartmann, *Ordinary Differential Equations*, Wiley, New-York (1964).
- [6] Z. Belhachmi, B. Brighi & K. Taous, On the concave solutions of the Blasius equation, *Acta Math. Univ. Comenianae* **69** (2), pp. 199-214 (2000).
- [7] D. B. Ingham, S. N. Brown, Flow past a suddenly heated vertical plate in a porous medium, *J. Proc. R. Soc. Lond. A* 403 (1986), pp. 51-80.
- [8] Z. Belhachmi, B. Brighi & K. Taous, On a family of differential equations for boundary layer approximations in porous media, *Euro. Jnl of Applied Mathematics*, Vol. 12, Cambridge University Press (2001), pp. 513-528.
- [9] B. Brighi, On a similarity boundary layer equation, *Zeitschrift für Analysis und ihre Anwendungen* **21** (4), pp. 931-948 (2002).
- [10] B. Brighi, T. Sari, Blowing-up coordinates for a similarity boundary layer equation. To appear in *Discrete and Continuous Dynamical Systems (Serie A)*.
- [11] M. Guedda, Multiple solutions of mixed convection boundary-layer approximations in a porous medium. To appear in *Applied Mathematics Letters*.
- [12] S. P. Hastings, W. C. Troy, Oscillating solutions of the Falkner-Skan equations for positive β , *J. Diff. Equations* **71** (1), pp. 123-144 (1988).