

THE BLOCKS OF THE BRAUER ALGEBRA IN CHARACTERISTIC ZERO

ANTON COX, MAUD DE VISSCHER, AND PAUL MARTIN

ABSTRACT. We determine the blocks of the Brauer algebra in characteristic zero. We also give information on the submodule structure of standard modules for this algebra.

1. INTRODUCTION

The Brauer algebra $B_n(\delta)$ was introduced in [Bra37] in the study of the representation theory of orthogonal and symplectic groups. Over \mathbb{C} , and for integral values of δ , its action on tensor space $T = (\mathbb{C}^{|\delta|})^{\otimes n}$ can be identified with the centraliser algebra for the corresponding group action. This generalises the Schur-Weyl duality between symmetric and general linear groups [Wey46].

If n is fixed, then for all $\delta \geq n$ the centraliser algebra $\text{End}_{O(\delta)}(T)$ has multimatrix structure *independent* of δ , and Brauer's algebra $B_n(\delta)$ unifies these algebras, having a basis independent of δ , and a composition which makes sense over any field k and for any $\delta \in k$. The Brauer algebra is well defined in particular for positive integral $\delta < n$, but the action on T is faithful for positive integral δ if and only if $\delta \geq n$.

In classical invariant theory one is interested in the Brauer algebra *per se* only in so far as it coincides with the centraliser of the classical group action on T ; i.e., in the case of δ integral with $|\delta|$ large compared to n . Here we take another view, and consider the stable properties for fixed δ and arbitrarily large n . In such cases $B_n(\delta)$ is not semisimple for δ integral. However it belongs to a remarkable family of algebras arising both in invariant theory and in statistical mechanics for which this view is very natural. (For example when considered from the point of view of transfer matrix algebras in statistical mechanics [Mar91].) Indeed much of the structure of $B_n(\delta)$ can be recovered from a suitable global limit of n by localisation (and in this sense its structure does not depend on n).

This family of algebras can be introduced as follows. Consider the diagram of commuting actions on T , with $|\delta| = N$:

$$\begin{array}{ccccc}
 GL(N) & & & & \mathbb{C}\Sigma_n \\
 & \searrow & & & \nearrow \\
 \cup & & & & \cap \\
 O(N) & \longrightarrow & T & \longleftarrow & B_n(N) \\
 & \nearrow & & & \searrow \\
 \cup & & & & \cap \\
 \Sigma_N & & & & P_n(N)
 \end{array}$$

where the actions of the algebra on the right centralise the action of the group on the left in the same row, and vice versa. The bottom row consists of the diagonal action of Σ_N permuting the standard ordered basis of \mathbb{C}^N on the left, and the partition algebra $P_n(N)$ on the right. The partition algebra $P_n(\delta)$ (for any δ) has a basis of partitions of two rows of n vertices. The Brauer algebra is the subalgebra with basis the subset of pair partitions, and $\mathbb{C}\Sigma_n$ is the subalgebra with basis the pair partitions such that each pair contains a vertex from each row. The Brauer algebra also has a subalgebra with basis the set of pair partitions which can be represented by noncrossing lines drawn vertex-to-vertex in an interval of the plane with the rows of vertices on its boundary. This is the Temperley-Lieb algebra $T_n(\delta)$.

All of these algebras are rather well understood over \mathbb{C} , with the exception of B_n . All their decomposition matrices are known, and all of their blocks can be described by an appropriate geometric linkage principle. For Σ_n both data are trivial, since it is semisimple. For T_n each standard module has either one or two composition factors and its alcove geometry is affine A_1 (affine reflections on the real line). For P_n each standard module has either one or two composition factors and its alcove geometry is affine A_∞ (although locally the block structure looks like affine A_1).

Over \mathbb{C} , the Brauer algebra is semisimple for δ sufficiently large, and is generically semisimple [Bro55]. Hanlon and Wales studied these algebras in a series of papers [HW89b, HW89a, HW90, HW94] and conjectured that $B_n(\delta)$ is semi-simple for all non-integral choices of δ . This was proved by Wenzl [Wen88].

In this paper we determine the blocks of B_n for δ integral. The simple modules of B_n may be indexed by partitions of those natural numbers congruent to n modulo 2 and not exceeding n , and hence by Young diagrams (if $\delta = 0$ then the empty partition is omitted). We will call these indexing objects *weights*. Given $\delta \in R$ a ring we can associate a *charge* $ch(\epsilon) \in R$ to each box ϵ in a Young diagram, as shown in Figure 1. We will also refer later to the usual *content* of boxes which, for the box ϵ in i -th row and j -th column is $c(\epsilon) = j - i$. It is easy to see that $ch(\epsilon) = \delta - 1 + 2c(\epsilon)$. For each pair of diagrams λ and μ we will also need to consider the skew partitions $\lambda/(\lambda \cap \mu)$ and $\mu/(\lambda \cap \mu)$ consisting of those boxes occurring in λ but not μ and in μ but not λ .

With these notations we can now state the two main results of the paper (which are valid without restriction on δ).

Corollary 6.7. *The simple modules $L(\lambda)$ and $L(\mu)$ are in the same block if and only if*

$\delta - 1$	$\delta + 1$	$\delta + 3$	\dots
$\delta - 3$	$\delta - 1$	$\delta + 1$	\dots
$\delta - 5$	$\delta - 3$	$\delta - 1$	\dots
\vdots	\vdots		

FIGURE 1. The charges associated to boxes in a Young diagram

(i) The boxes in $\lambda/(\lambda \cap \mu)$ (respectively $\mu/(\lambda \cap \mu)$) can be put into pairs whose charges sum to zero;

(ii) if $\lambda/(\lambda \cap \mu)$ (respectively $\mu/(\lambda \cap \mu)$) contains $\begin{array}{|c|} \hline 1 \\ \hline -1 \\ \hline \end{array}$ with no $\begin{array}{|c|} \hline 1 \\ \hline \end{array}$ to the right of these boxes then it contains an even number of 1/-1 pairs.

Examples illustrating this result are given in Example 4.9.

Theorem 7.3 (Summary). *For any integral δ and natural number l a standard module can be constructed (for some $B_n(\delta)$) whose socle series length is greater than l . This module also has a socle layer containing at least l simples.*

The second result shows that the structure of standard modules can become arbitrarily complicated. This is in marked contrast to the partition and Temperley-Lieb algebra, and symmetric group, cases.

To prove these results we use the theory of towers of recollement developed in [CMPX]. This approach is already closely modelled, for B_n , in work of Doran, Wales, and Hanlon [DWH99] (since both papers use the methods developed in [Mar96]). This key paper of Doran, Wales and Hanlon will be the starting point for our work, and we will generalise and refine several of their results.

The ‘diagram’ algebras $P_n \supset B_n \supset T_n$ are amenable to many powerful representation theory techniques, and yet the representation theory of the Brauer algebra is highly non-trivial in comparison to the others. We shall see that, in terms of degree of difficulty, the study of Brauer representation theory in characteristic zero is an intermediate between the study of ‘classical’ objects in characteristic zero and the *grand theme* of the representation theory of finite dimensional algebras, the study of Σ_n in characteristic p .

Another such intermediate class of objects are the Hecke algebras of type A at roots of unity, which are Ringel dual to the generalised Lie objects known as quantum groups. The Brauer algebra B_n in characteristic zero has, through its global limit, more Lie-theory-like structure than Σ_n in characteristic p (for which not even a good organisational scheme within which to address the problem is known, for small primes p). This is reminiscent of the virtual algebraic Lie theory discussed for the (generalised) blob algebras in [MRH04, MW03]. However in the Brauer algebra case, any candidate for an alcove geometry formulation will be considerably more complicated [Naz96, OR01]. For these reasons we consider the further

study of the Brauer algebra in characteristic zero to be an important problem in representation theory.

The paper begins with a section defining the various objects of interest, and a review of their basic properties in the spirit of [CMPX]. This is followed by a brief section describing some basic results about Littlewood-Richardson coefficients which will be needed in what follows. In Section 4 we begin the analysis of blocks by giving a necessary condition for two weights to be in the same block. This is based on an analysis of the action of certain central elements in the algebra on standard modules, and inductive arguments using Frobenius reciprocity. Section 5 constructs homomorphisms between standard modules in certain special cases, generalising a result in [DWH99]. Although not necessary for the main block result, this is of independent interest.

The classification of blocks is completed in Section 6. The main idea is to show that every block contains a unique minimal weight, and that there is a homomorphism from any standard labelled by a non-minimal weight to one labelled by a smaller weight. We also describe precisely which weights are minimal in their blocks.

In Section 7 we consider certain explicit choice of weights, and show inductively, via Frobenius reciprocity arguments, that the corresponding standards can have arbitrarily complicated submodule structures. We conclude by outlining the modifications to our arguments required in the case $\delta = 0$.

The structure of the Brauer algebra becomes much more complicated when considered over an arbitrary field k . For general k and δ integral it is expected that this algebra still acts as a centraliser algebra, and this has been shown in a recent series of papers for the symplectic case [Dot98, Oeh01, DDH]. A necessary and sufficient condition for semisimplicity (which holds over arbitrary fields) was given recently by Rui [Rui05]. The study of Young and permutation modules for these algebras has been started in [HP].

2. PRELIMINARIES

In this section we will consider the Brauer algebra defined over a general field k of characteristic $p \geq 0$, although we will later restrict attention to the case $k = \mathbb{C}$. After reviewing the definition of the Brauer algebra, we will show that families of such algebras form towers of recollement in the sense of [CMPX] (which we will see follows from various results of Doran et. al. [DWH99]). This will be the framework in which we base our analysis of these algebras.

Given $n \in \mathbb{N}$ and $\delta \in k$, the *Brauer algebra* $B_n(\delta)$ is a finite dimensional associative k -algebra generated by certain Brauer diagrams. A general (n, t) -*(Brauer) diagram* consists of a rectangular box (or *frame*) with n distinguished points on the northern boundary and t distinguished points on the southern boundary, which we call *nodes*. Each node is joined to precisely one other by a line, and there may also be one or more closed loops inside the frame. Those diagrams without closed loops are called *reduced*. We will label the northern nodes from left to right by $1, 2, \dots, n$ and the southern nodes from left to right by $\bar{1}, \bar{2}, \dots, \bar{t}$. We identify diagrams if they connect the same pairs of labelled nodes, and have the same number of closed loops. Lines which connect two nodes on the northern (respectively southern) boundary will be called *northern* (respectively *southern*) *arcs*; those connecting a northern node to a southern node will be called *propagating lines*.




FIGURE 2. Multiplication of two diagrams in $B_6(\delta)$

Given an (n, t) -diagram A and a (t, u) -diagram B , we define the product AB to be the (n, u) -diagram obtained by concatenation of A above B (where we identify the southern nodes of A with the northern nodes of B and then ignore the section of the frame common to both diagrams). As a set, the Brauer algebra $B_n(\delta)$ consists of linear combinations of (n, n) -diagrams. This has an obvious additive structure, and multiplication is induced by concatenation. We also impose the relation that any non-reduced diagram containing m closed loops equals δ^m times the same diagram with all closed loops removed. A basis is then given by the set of reduced diagrams. An example of a product of two diagrams is given in Figure 2. For convenience, we set $B_0(\delta) = k$. When no confusion is likely to arise, we denote the algebra $B_n(\delta)$ simply by B_n .

We will now apply as much as possible from the general setup of [CMPX] to the Brauer algebra. The labels (A1), (A2), etc., refer to the axioms in that paper. Henceforth, we assume that $\delta \neq 0$; for the case $\delta = 0$ see Section 8.

For $n \geq 2$ consider the idempotent e_n in B_n defined by $1/\delta$ times the Brauer diagram where i is joined to \bar{i} for $i = 1, \dots, n - 2$, and $n - 1$ is joined to n and $\bar{n} - 1$ is joined to \bar{n} . This is illustrated in Figure 3.




FIGURE 3. The idempotent e_n

Lemma 2.1 (A1). *For each $n \geq 2$, we have an algebra isomorphism*

$$\Phi_n : B_{n-2} \longrightarrow e_n B_n e_n$$

which takes a diagram in B_{n-2} to the diagram in B_n obtained by adding an extra northern and southern arc to the righthand end.

This allows us to define, following Green [Gre80], an exact localization functor

$$\begin{aligned} F_n : B_n\text{-mod} &\longrightarrow B_{n-2}\text{-mod} \\ M &\longmapsto e_n M \end{aligned}$$

and a right exact globalization functor

$$\begin{aligned} G_n : B_n\text{-mod} &\longrightarrow B_{n+2}\text{-mod} \\ M &\longmapsto B_{n+2} e_{n+2} \otimes_{B_n} M. \end{aligned}$$

Note that $F_{n+2} G_n(M) \cong M$ for all $M \in B_n\text{-mod}$, and hence G_n is a full embedding.

From this we can quickly deduce an indexing set for the isomorphism classes of simple B_n -modules. It is easy to see that

$$B_n/B_n e_n B_n \cong k\Sigma_n \quad (1)$$

the group algebra of the symmetric group on n symbols. If the simple $k\Sigma_n$ -modules are indexed by the set Λ^n then by [Gre80] and Lemma 2.1, the simple B_n -modules are indexed by the set

$$\Lambda_n = \Lambda^n \sqcup \Lambda_{n-2} = \Lambda^n \sqcup \Lambda^{n-2} \sqcup \dots \sqcup \Lambda^{\min}$$

where $\min = 0$ or 1 depending on the parity of n . If $p = 0$ or $p > n$ then the set Λ^n corresponds to the set of partitions of n ; we write $\lambda \vdash n$ if λ is such a partition.

For $m - n$ even we write Λ_n^m for Λ^m regarded as a subset of Λ_n . (If $m > n$ then $\Lambda_n^m = \emptyset$.) We also write Λ for the disjoint union of all the Λ^n , and call this the set of *weights* for the Brauer algebra. We will henceforth abuse terminology and refer to weights as being in the same block of B_n if the corresponding simple modules are in the same block.

For $n \geq 2$ and $0 \leq t \leq n/2$, define the idempotent $e_{n,t}$ to be 1 if $k = 0$ or $1/\delta^t$ times the Brauer diagram with edges between i and \bar{i} for all $1 \leq i \leq n - 2t$ and between j and $\bar{j} + 1$, and \bar{j} and $\overline{j+1}$ for $n - 2t + 1 \leq j \leq n - 1$. (This is the image of e_t via the isomorphisms arising in Lemma 2.1.) Set $B_{n,t} = B_n/B_n e_{n,t} B_n$.

Lemma 2.2 (A2). *The natural multiplication map*

$$B_{n,t} e_{n,t} \otimes_{e_{n,t} B_{n,t} e_{n,t}} e_{n,t} B_{n,t} \longrightarrow B_{n,t} e_{n,t} B_{n,t}$$

is bijective. If $\delta \neq 0$ and either $p = 0$ or $p > n$ then $B_n/B_n e_n B_n$ is semisimple.

Proof. The second part follows from (1) and standard symmetric group results. For the first part, the map is clearly surjective so we only need to show that it is also injective. It is easy to verify that:

- (i) $B_{n,t}$ has a basis given by all reduced diagrams having at least $n - 2t$ propagating lines,
- (ii) $B_{n,t} e_{n,t} B_{n,t}$ has a basis given by all reduced diagrams having exactly $n - 2t$ vertical edges, and
- (iii) $e_{n,t} B_{n,t} e_{n,t} \cong k\Sigma_{n-2t}$.

Now suppose that X and X' are diagrams in $B_{n,t} e_{n,t}$. Any such diagram has a southern edge where the leftmost $n - 2t$ nodes lie on propagating lines, with the remaining southern nodes paired consecutively. The northern edge has exactly t northern arcs. We will label such a diagram by $X_{v,1,\sigma}$, where v represents the configuration of northern arcs, 1 represents the fixed southern boundary, and $\sigma \in \Sigma_{n-2t}$ is the permutation obtained by setting $\sigma(i) = j$ if the i th propagating northern node from the left is connected to \bar{j} . (For later use we will denote the set of elements v arising thus by $V_{n,t}$, and call such elements *partial one-row diagrams*.) Similarly a diagram Y in $e_{n,t} B_{n,t}$ will be labelled by $Y_{1,v,\sigma}$.

It will be enough to show that the multiplication map is injective on the set of tensor products of diagram elements. Given $X = X_{v,1,\sigma}$ and $X' = X_{v',1,\sigma'}$ in $B_{n,t} e_{n,t}$ and $Y = Y_{1,w,\tau}$, $Y' = Y_{1,w',\tau'}$ in $e_{n,t} B_{n,t}$, assume that $XY = X'Y'$. Then we must have $v = v'$, $w = w'$ and $\sigma \circ \tau = \sigma' \circ \tau'$. It now follows from the identification in (iii) that $X \otimes Y = X' \otimes Y'$ in $B_{n,t} e_{n,t} \otimes_{e_{n,t} B_{n,t} e_{n,t}} e_{n,t} B_{n,t}$. \square

We immediately obtain

Corollary 2.3 (A2'). *If $\delta \neq 0$ and either $p = 0$ or $p > n$ then B_n is a quasi-hereditary algebra, with heredity chain given by*

$$0 \subset \cdots \subset B_n e_{n,k} B_n \subset \cdots \subset B_n e_{n,0} B_n.$$

The partial ordering is given as follows: for $\lambda, \mu \in \Lambda_n$ we have $\lambda \leq \mu$ if and only if either $\lambda = \mu$ or $\lambda \in \Lambda_n^s$ and $\mu \in \Lambda_n^t$ with $s > t$.

Henceforth we assume that p satisfies the conditions in Corollary 2.3. It follows from the quasi-hereditary structure that for each $\lambda \in \Lambda_n$ we have a standard module $\Delta_n(\lambda)$ having simple head $L_n(\lambda)$ and all other composition factor $L_n(\mu)$ satisfying $\mu < \lambda$. Note that if $\lambda \in \Lambda_n^n$ then

$$\Delta_n(\lambda) = L_n(\lambda) \cong S^\lambda$$

the lift to B_n of the Specht module for $B_n/B_n e_n B_n \cong k\Sigma_n$.

Note also that by [Don98, A1] and arguments as in [MRH04, Proposition 3], the quasi-hereditary structure is compatible with the globalization and localization functors. That is, for all $\lambda \in \Lambda_n$ we have

$$G_n(\Delta_n(\lambda)) = \Delta_{n+2}(\lambda) \tag{2}$$

$$F_n(\Delta_n(\lambda)) = \begin{cases} \Delta_{n-2}(\lambda) & \text{if } \lambda \in \Lambda_{n-2} \\ 0 & \text{otherwise} \end{cases} \tag{3}$$

As F_n is exact we also have that

$$F_n(L_n(\lambda)) = \begin{cases} L_{n-2}(\lambda) & \text{if } \lambda \in \Lambda_{n-2} \\ 0 & \text{otherwise} \end{cases} \tag{4}$$

For every partition μ of some $m = n - 2t$ we can give an explicit construction of the modules $\Delta_n(\mu)$. Let $e = e_{n,t} \in B_n$ be as above, so that $eB_n e \cong B_m$. If we denote by S^μ the lift of the Specht module labelled by μ for $k\Sigma_m$ to B_m , then by (2) we have that

$$\Delta_n(\mu) \cong B_n e \otimes_{eB_n e} S^\mu. \tag{5}$$

Using this fact, it is easy to give a basis for this module in terms of some basis $\mathcal{B}(\mu)$ of S^μ , using the notation introduced during the proof of Lemma 2.2.

Lemma 2.4. *If μ is a partition of $n - 2t$ then the module $\Delta_n(\mu)$ has a basis given by*

$$\{X_{v,1,id} \otimes x \mid v \in V_{n,t}, x \in \mathcal{B}(\mu)\}.$$

Via this Lemma we may identify our standard modules $\Delta_n(\lambda)$ with the modules $\mathcal{S}_\lambda(n)$ in [DWH99] (which in turn come from [Bro55]). Note that if we define $\Delta_n(\mu)$ as the tensor product in (5) then we have a definition that makes sense for all values of p . In the non-quasi-hereditary cases these modules still play an important role, as the algebras are cellular [GL96] with the $\Delta_n(\mu)$ as cell modules.

We will frequently need a second way to relate different Brauer algebras.

Lemma 2.5 (A3). *For each $n \geq 1$, the algebra B_n can be identified as a subalgebra of B_{n+1} via the homomorphism which takes a Brauer diagram X in B_n to the Brauer diagram in B_{n+1} obtained by adding two vertices $n + 1$ and $\overline{n + 1}$ with a line between them.*

Lemma 2.5 implies that we can consider the usual restriction and induction functors

$$\begin{aligned} \text{res}_n : B_n\text{-mod} &\longrightarrow B_{n-1}\text{-mod} \\ M &\longmapsto M|_{B_{n-1}} \end{aligned}$$

and

$$\begin{aligned} \text{ind}_n : B_n\text{-mod} &\longrightarrow B_{n+1}\text{-mod} \\ M &\longmapsto B_{n+1} \otimes_{B_n} M. \end{aligned}$$

We can relate these functors to globalisation and localisation via

Lemma 2.6 (A4). (i) For all $n \geq 2$ we have that

$$B_n e_n \cong B_{n-1}$$

as a left B_{n-1} , right B_{n-2} -bimodule.

(ii) For all B_n -modules M we have

$$\text{res}_{n+2}(G_n(M)) \cong \text{ind}_n(M).$$

Proof. (i) Every Brauer diagram in $B_n e_n$ has an edge between $\overline{n-1}$ and \bar{n} . Define a map from $B_n e_n$ to B_{n-1} by sending a diagram X to the diagram with $2(n-1)$ vertices obtained from X by removing the line connecting $\overline{n-1}$ and \bar{n} and the line from n , and pairing the vertex $\overline{n-1}$ to the vertex originally paired with n in X . It is easy to check that this gives an isomorphism.

(ii) Using (i) we have

$$\begin{aligned} \text{res}_{n+2}(G_n(M)) &= (B_{n+2} e_{n+2} \otimes_{B_n} M)|_{B_{n+1}} \\ &\cong B_{n+1} \otimes_{B_n} M \cong \text{ind } M. \end{aligned}$$

□

Let λ be a partition of n and μ be a partition of $n-1$. We write $\lambda \triangleright \mu$ and $\mu \triangleleft \lambda$ if μ is obtained from λ by removing a box from its Young diagram (equivalently if λ is obtained from μ by adding a box to its Young diagram). The following result does *not* require any restriction on the characteristic of our field.

Proposition 2.7 (A5 and 6). For $\lambda \in \Lambda_n$ we have short exact sequences

$$0 \rightarrow \bigoplus_{\mu \triangleleft \lambda} \Delta_{n+1}(\mu) \rightarrow \text{ind}_n \Delta_n(\lambda) \rightarrow \bigoplus_{\mu \triangleright \lambda} \Delta_{n+1}(\mu) \rightarrow 0$$

and

$$0 \rightarrow \bigoplus_{\mu \triangleleft \lambda} \Delta_{n-1}(\mu) \rightarrow \text{res}_n \Delta_n(\lambda) \rightarrow \bigoplus_{\mu \triangleright \lambda} \Delta_{n-1}(\mu) \rightarrow 0.$$

Proof. This was proved for $k = \mathbb{C}$ in [DWH99, Theorem 4.1 and Corollary 6.4] (as the condition $\lambda \vdash n$ in [DWH99, Corollary 6.4] is not needed). However, their proof is valid over any field. □

Wenzl [Wen88] has shown that B_n is semisimple when $k = \mathbb{C}$ and $\delta \notin \mathbb{Z}$. (Over an arbitrary field, a necessary and sufficient condition for semisimplicity has been given by Rui [Rui05].) For this reason we do not consider the case of non-integral δ . As we will regularly need to appeal to the representation theory of the symmetric group, which is not well understood in positive characteristic, we will also only consider the characteristic zero case. In summary:

Henceforth we will assume that $k = \mathbb{C}$ and $\delta \in \mathbb{Z} \setminus \{0\}$, unless otherwise stated.

3. SOME LITTLEWOOD-RICHARDSON COEFFICIENTS

One of the key results used by [DWH99] in their analysis of the Brauer algebra is [HW90, Theorem 4.1] which decomposes standard modules $\Delta_n(\lambda)$ with $\lambda \vdash n$ as symmetric group modules. Recall that a partition is *even* if every part of the partition is even, and that $c_{\mu\eta}^\lambda$ denotes a Littlewood-Richardson coefficient. If $\lambda \vdash n$ and $\mu \vdash m$ then [HW90, Theorem 4.1] states that either $[\text{res}_{\mathbb{C}\Sigma_n} \Delta_n(\mu) : S^\lambda] = 0$ or $m = n - 2t$ for some $t \geq 0$ and

$$[\text{res}_{\mathbb{C}\Sigma_n} \Delta_n(\mu) : S^\lambda] = \sum_{\substack{\eta \vdash 2t \\ \eta \text{ even}}} c_{\mu\eta}^\lambda \quad (6)$$

As this result is stated in terms of Littlewood-Richardson coefficients, we will find it useful to calculate these in certain special cases.

Lemma 3.1. *If $\mu \subset \lambda$ are partitions such that $\nu = \lambda/\mu$ is also a partition then*

$$c_{\mu\eta}^\lambda = \begin{cases} 1 & \text{if } \eta = \nu \\ 0 & \text{otherwise.} \end{cases}$$

Proof. This follows immediately from the definition of Littlewood-Richardson coefficients in terms of rectification of skew tableaux (see [Ful97, Section 5.1, Corollary 2]) □

For our second calculation we will need an alternative definition of Littlewood-Richardson coefficients (which can be found in [JK81, 2.8.14 Corollary]). When considering a configuration of boxes labeled by elements b_{ij} we say that the configuration is *valid* if:

- (i) For all i , if $y < j$ then b_{iy} is in a later column than b_{ij} .
- (ii) For all j , if $x < i$ then b_{xj} is in an earlier row than b_{ij} .

For each box (i, j) of η consider a symbol b_{ij} . Then the Littlewood-Richardson coefficient $c_{\mu\eta}^\lambda$ is the number of ways one can form λ from μ by adding the boxes of η to μ in the following manner. First add $b_{11}, b_{12}, \dots, b_{1\eta_1}$ to η to form a new partition η^1 . Continue inductively by adding $b_{i1}, b_{i2}, \dots, b_{i\eta_i}$ to η^{i-1} to form a new partition η^i . We require that the final configuration of the elements b_{ij} is valid.

Lemma 3.2. *If $\mu \subset \lambda$ are partitions with $\lambda = (a^b)$ for some a and b then there is a unique partition $\eta = (\eta_1, \dots, \eta_r)$ such that $c_{\mu\eta}^\lambda \neq 0$, and for this partition we have $c_{\mu\eta}^\lambda = 1$. Further, $(\lambda/\mu)_i = \eta_{r-i}$.*

Proof. Consider valid extensions of μ by any η to form λ . As λ is a rectangle, the final row of η can only be placed as illustrated in Figure 4(a). Then the penultimate row of η must be placed as illustrated in Figure 4(b). Continuing in this way we see that the choice of η is

unique, and the number of boxes in the final row of λ/μ must equal η_1 , in the penultimate row must equal η_2 , and so on. \square

			$b_{(r-1)\eta_r}$	\cdots	$b_{(r-1)2}$	$b_{(r-1)1}$
$b_{(r-1)\eta_{r-1}}$	\cdots	$b_{(r-1)(\eta_r+1)}$	$b_{r\eta_r}$	\cdots	b_{r2}	b_{r1}

FIGURE 4. The final two rows of η in λ

4. A PARTIAL BLOCK RESULT

Doran, Wales, and Hanlon [DWH99] have given a necessary condition for the existence of a non-zero homomorphism of B_n -modules from $\Delta_n(\lambda)$ to $\Delta_n(\mu)$. We will first elevate this condition to a partial block result, and then give a stronger necessary condition that must also hold for two weights to be in the same block. In section 6 we will see that this stronger condition is also sufficient for two weights to be in the same block.

Let λ be a partition. For a box d in the corresponding Young diagram $[\lambda]$, we denote by $c(d)$ the *content* of d . Recall that if $d = (x, y)$ is in the x -th row (counting from top to bottom) and in the y -th column (counting from left to right) of $[\lambda]$, then $c(d) = y - x$. We denote by $\mathbf{c}(\lambda)$ the *multiset* $\{c(d) : d \in [\lambda]\}$. If μ is a partition with $[\mu] \subseteq [\lambda]$ we write $\mu \subseteq \lambda$, and denote the skew partition obtained by removing μ from λ by λ/μ . We then denote by $\mathbf{c}(\lambda/\mu)$ the multiset $\mathbf{c}(\lambda) \setminus \mathbf{c}(\mu)$.

Write $X_{i,j}$ for the Brauer diagram in B_n with edges between t and \bar{t} for all $t \neq i, j$ and with edges between i and j and between \bar{i} and \bar{j} . Note that B_n is generated by the elements $X_{i,j}$ together with the symmetric group Σ_n (identified with the set of diagrams with n propagating lines). We denote by T_n the element $\sum_{1 \leq i < j \leq n} X_{i,j}$ in B_n . Recall also the definition of partial one-row diagrams in the proof of Lemma 2.2.

Lemma 4.1. *Let μ be a partition of m with $m = n - 2t$. For all $w \in V_{n,t}$ and $x \in S^\mu$ we have that*

$$T_n(X_{w,1,id} \otimes x) = \left(t(\delta - 1) - \sum_{d \in [\mu]} c(d) + \sum_{1 \leq i < j \leq n} (i, j) \right) (X_{w,1,id} \otimes x)$$

where (i, j) denotes the element of Σ_n which transposes i and j . Hence for all $y \in \Delta_n(\mu)$ we have

$$T_n y = \left(t(\delta - 1) - \sum_{d \in [\mu]} c(d) + \sum_{1 \leq i < j \leq n} (i, j) \right) y.$$

Proof. This is essentially [DWH99, Lemma 3.2], together with observations in the proof of [DWH99, Theorem 3.3]. \square

The next result is a slight strengthening of [DWH99, Theorem 3.3] (which in turn generalises [Naz96, formula before (2.13)], which considers the case $\delta \in \mathbb{N}$). The original results provide a necessary condition for the existence of a homomorphism between two standard modules, but can be refined to prove

Proposition 4.2. *Suppose that $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$. Then either $\lambda = \mu$ or $\lambda \in \Lambda_n^r$ and $\mu \in \Lambda_n^s$ for some $r - s = 2t > 0$. Further, we must have*

$$\mu \subseteq \lambda \quad \text{and} \quad t(\delta - 1) + \sum_{d \in [\lambda/\mu]} c(d) = 0.$$

Proof. The first part of the proposition is clear from the quasi-hereditary structure of B_n . For the second part, note that by using the exactness of the localization functor we have

$$[\Delta_n(\mu) : L_n(\lambda)] = [\Delta_r(\mu) : L_r(\lambda)]$$

and hence we may assume that λ is a partition of n . In this case, $L_n(\lambda) = \Delta_n(\lambda) = S^\lambda$, the lift of the Specht module for $\mathbb{C}\Sigma_n$ to $B_n(\delta)$, and so any Brauer diagram having fewer than n propagating lines must act as zero on $L_n(\lambda)$. In particular, all the $X_{i,j}$'s act as zero and hence so does T_n .

The condition that $\mu \subseteq \lambda$ now follows by regarding $\Delta_n(\mu)$ as a $\mathbb{C}\Sigma_n$ -module by restriction and using (6) which describes the multiplicities of composition factors of such a module.

For the final condition, we know by assumption that there must exist a B_n -submodule M of $\Delta_n(\mu)$ and a B_n -homomorphism

$$\phi : L_n(\lambda) \longrightarrow \Delta_n(\mu)/M.$$

Let N be the B_n -submodule of $\Delta_n(\mu)$ containing M such that

$$\phi(L_n(\lambda)) = N/M.$$

As $N|_{\mathbb{C}\Sigma_n}$ is semisimple, we can find a $\mathbb{C}\Sigma_n$ -submodule W of N such that $N = W \oplus M$ and $W \cong S^\lambda$. By Lemma 4.1 we have for all $y \in W$ that

$$T_n y = \left(t(\delta - 1) - \sum_{d \in [\mu]} c(d) + \sum_{1 \leq i < j \leq n} (i, j) \right) y.$$

But $W \cong S^\lambda$ is a simple $\mathbb{C}\Sigma_n$ -module and $\sum_{1 \leq i < j \leq n} (i, j)$ is in the centre of $\mathbb{C}\Sigma_n$, so it must act as a scalar on W . It is well known [Dia, Chapter 1] that this scalar is given by $\sum_{d \in [\lambda]} c(d)$. Hence we have

$$\begin{aligned} T_n y &= \left(t(\delta - 1) - \sum_{d \in [\mu]} c(d) + \sum_{d \in [\lambda]} c(d) \right) y \\ &= \left(t(\delta - 1) + \sum_{d \in [\lambda/\mu]} c(d) \right) y. \end{aligned}$$

But T_n must act as zero on N and hence $t(\delta - 1) + \sum_{d \in [\lambda/\mu]} c(d) = 0$. □

By standard quasi-heredity arguments [Don98, Appendix] we deduce

Corollary 4.3. *Suppose that $\lambda \in \Lambda_n^r$ and $\mu \in \Lambda_n^s$ with $s < r$. If λ and μ are in the same block then $s = r - 2t$ for some $t \in \mathbb{N}$ and*

$$t(\delta - 1) + \sum_{d \in [\lambda]} c(d) - \sum_{d \in [\mu]} c(d) = 0. \quad (7)$$

When $t = 2$ [DWH99] gave a necessary and sufficient condition for the existence of a standard module homomorphism. From their results we obtain

Theorem 4.4. *Suppose that $\mu \subset \lambda$ with $|\lambda/\mu| = 2$. Then*

$$\dim \text{Hom}(\Delta_n(\lambda), \Delta_n(\mu)) \leq 1$$

and is non-zero if and only if λ and μ satisfy (7) with $\lambda/\mu \neq (1^2)$. Indeed, if $\lambda/\mu = (1^2)$ then

$$[\Delta_n(\mu) : L_n(\lambda)] = 0.$$

Proof. It is enough to consider the case when $\lambda \vdash n$, as the general case follows by globalisation. If λ and μ do not satisfy the required conditions then there is no composition factor $L_n(\lambda)$ in $\Delta_n(\mu)$ (and hence no homomorphism) by Corollary 4.3 and the remarks after [DWH99, Theorem 3.1]. In the remaining cases the existence of such a homomorphism was shown in [DWH99, Theorem 3.4]. By the remarks after [DWH99, Theorem 3.1] the multiplicity of the simple module $\Delta_n(\lambda)$ in $\Delta_n(\mu)$ is 1, and the dimension result is now immediate. \square

The next result is a strengthening of Proposition 4.2.

Proposition 4.5. *Suppose that $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$. Then there is a pairing of the boxes in λ/μ such that the sum of the content of the boxes in each pair is equal to $1 - \delta$.*

Proof. We use induction on n ; the case $n = 2$ is covered by Proposition 4.2. Thus we assume that the result holds for $n - 1$ and will show that it holds for n .

If $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$ then by Proposition 4.2 we know that $\mu \subseteq \lambda$ and

$$t(\delta - 1) + \sum_{d \in [\lambda/\mu]} c(d) = 0 \quad (8)$$

where $2t = |\lambda| - |\mu|$. Now suppose, for a contradiction, that there is no pairing of the boxes of $[\lambda/\mu]$ satisfying the condition of the proposition. By localising we may assume that λ is a partition of n , so that $L_n(\lambda) = \Delta_n(\lambda)$. Thus $\Delta_n(\mu)$ has a submodule M such that $\Delta_n(\lambda) \hookrightarrow \Delta_n(\mu)/M$.

The partition λ has a removable box ϵ_i of content s say and by Proposition 2.7 we have a surjection $\text{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon_i) \rightarrow \Delta_n(\lambda)$. Hence we have

$$\text{Hom}(\text{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon_i), \Delta_n(\mu)/M) \neq 0$$

and so by Frobenius reciprocity we have

$$\text{Hom}(\Delta_{n-1}(\lambda - \epsilon_i), \text{res}_n(\Delta_n(\mu)/M)) \neq 0.$$

This implies that $\Delta_{n-1}(\lambda - \epsilon_i) = L_{n-1}(\lambda - \epsilon_i)$ is a composition factor of $\text{res}_n(\Delta_n(\mu))$. Now using Proposition 2.7 we see that either

- (i) the weight μ has a removable box ϵ_j such that $[\Delta_{n-1}(\mu - \epsilon_j) : L_{n-1}(\lambda - \epsilon_i)] \neq 0$, or
 - (ii) the weight μ has an addable box ϵ_j such that $[\Delta_{n-1}(\mu + \epsilon_j) : L_{n-1}(\lambda - \epsilon_i)] \neq 0$.
- We consider each case in turn.

In case (i), Proposition 4.2 implies that $[\mu - \epsilon_j] \subseteq [\lambda - \epsilon_i]$ and

$$t(\delta - 1) + \sum_{d \in [\lambda/\mu]} c(d) - c(\epsilon_i) + c(\epsilon_j) = 0.$$

Hence from (8) we must have

$$c(\epsilon_j) = c(\epsilon_i) = s$$

and by induction we can find a pairing of the boxes in $(\lambda - \epsilon_i)/(\mu - \epsilon_j)$ such that the sum of the content of the boxes in each pair is equal to $1 - \delta$. But as multisets

$$\mathbf{c}((\lambda - \epsilon_i)/(\mu - \epsilon_j)) = \mathbf{c}(\lambda/\mu) - c(\epsilon_i) + c(\epsilon_j) = \mathbf{c}(\lambda/\mu)$$

and hence there is such a pairing for the boxes of λ/μ . This gives the desired contradiction.

Now consider case (ii). Here μ has an addable box ϵ_j such that $[\mu + \epsilon_j] \subseteq [\lambda - \epsilon_i]$ and

$$(t - 1)(\delta - 1) + \sum_{d \in [\lambda/\mu]} c(d) - c(\epsilon_i) - c(\epsilon_j) = 0.$$

Comparing with (8) we deduce that

$$c(\epsilon_j) + c(\epsilon_i) = 1 - \delta.$$

By induction there is a pairing of the boxes of $(\lambda - \epsilon_i)/(\mu + \epsilon_j)$ satisfying the condition of the Proposition. But as multisets

$$\mathbf{c}((\lambda - \epsilon_i)/(\mu + \epsilon_j)) = \mathbf{c}(\lambda/\mu) - c(\epsilon_i) - c(\epsilon_j)$$

and as observed above the $c(\epsilon_i)$ and $c(\epsilon_j)$ can be paired in the right way. Hence the boxes of λ/μ can be paired appropriately, which again gives the desired contradiction. \square

When δ is even we will need a further refinement of Proposition 4.2. Given $\mu \subset \lambda$, consider the boxes with content $-\frac{\delta}{2}$ and $\frac{2-\delta}{2}$ in λ/μ . If $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$ then these must be paired by Proposition 4.5, and so must be in one of the two chain configurations illustrated in Figure 5 (for some length of chain).




FIGURE 5. The two possible configurations of paired boxes of contents $-\frac{\delta}{2}$ and $\frac{2-\delta}{2}$

Proposition 4.6. *Suppose that $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$ and δ is even. If the boxes of content $-\frac{\delta}{2}$ and $\frac{2-\delta}{2}$ are configured as in Figure 5(b) then the number of columns in this configuration must be even.*

Proof. We will show by induction on n that in case (b) the number of columns must be even. The case $n = 2$ is covered by Theorem 4.4.

By repeated applications of F we may assume that $\lambda \vdash n$. Let ϵ_i be a removable box of λ . As in the proof of Proposition 4.5 we have that if

$$[\Delta_n(\mu) : L_n(\lambda)] \neq 0$$

then either

$$[\Delta_{n-1}(\mu - \epsilon_j) : L_{n-1}(\lambda - \epsilon_i)] \neq 0$$

for some removable box ϵ_j of μ with $c(\epsilon_i) = c(\epsilon_j)$ and $\mu - \epsilon_j \subset \lambda - \epsilon_i$, or

$$[\Delta_{n-1}(\mu + \epsilon_j) : L_{n-1}(\lambda - \epsilon_i)] \neq 0$$

for some addable box ϵ_j of μ with $c(\epsilon_i) + c(\epsilon_j) = 1 - \delta$ and $\mu + \epsilon_j \subseteq \lambda - \epsilon_i$.

If $c(\epsilon_i)$ is not equal to either $-\frac{\delta}{2}$ or $\frac{2-\delta}{2}$ then the boxes of $(\lambda - \epsilon_i)/(\mu - \epsilon_j)$ (respectively of $(\lambda - \epsilon_i)/(\mu + \epsilon_j)$) of content $-\frac{\delta}{2}$ and $\frac{2-\delta}{2}$ are the same as those boxes in λ/μ , and so the result follows by induction. Also, by our assumption on the configuration of such boxes the partition λ does not have a removable box of content $\frac{2-\delta}{2}$. Thus we may assume that λ has only one removable box ϵ_i of content $-\frac{\delta}{2}$ (and hence that λ is a rectangle).




FIGURE 6. The partitions $\mu \subset \lambda$, with the configuration as in Figure 5(b) shaded

We have that λ and μ are of the form shown in Figure 6, with $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$. So in particular

$$[\text{res}_{\mathbb{C}\Sigma_n} \Delta_n(\mu) : S^\lambda] \neq 0.$$

By (6) we have

$$[\text{res}_{\mathbb{C}\Sigma_n} \Delta_n(\mu) : S^\lambda] = \sum_{\eta \text{ even}} c_{\mu\eta}^\lambda$$

and hence we must have $c_{\mu\eta}^\lambda \neq 0$ for some even partition $\eta = (\eta_1, \dots, \eta_r)$. As λ is a rectangle Lemma 3.2 implies there is only one possible η , and that each row of λ/μ has length η_i for some $1 \leq i \leq r$. But η was an even partition and hence these lengths are all even, which implies that the number of columns occupied by shaded boxes in Figure 6 is also even as required. \square

Definition 4.7. We say that λ and μ are δ -balanced (or just *balanced* when the context is clear) if: (i) there exists a pairing of the boxes in $\lambda/(\lambda \cap \mu)$ (respectively in $\mu/(\lambda \cap \mu)$) such that the contents of each pair sum to $1 - \delta$, and (ii) if δ is even and the boxes with content $-\frac{\delta}{2}$ and $\frac{2-\delta}{2}$ in $\lambda/(\lambda \cap \mu)$ (respectively in $\mu/(\lambda \cap \mu)$) are configured as in Figure 5(b), then the number of columns in this configuration is even.

Just as for Corollary 4.3 we can immediately deduce from Propositions 4.5 and 4.6 the following block result.

Corollary 4.8. *If λ and μ are in the same block then they are balanced.*




FIGURE 7. The diagrams $[\lambda]$, $[\mu]$ and $[\tau]$ in Example 4.9(i)

Example 4.9. (i) Let $\lambda = (6, 4^2, 2, 1)$, $\mu = (5, 2^2)$, $\tau = \lambda/(\lambda \cap \mu)$, and $\delta = 1$. The diagrams $[\lambda]$, $[\mu]$, and $[\tau]$ are illustrated (with their contents) in Figure 7. Clearly

$$\sum_{d \in [\lambda]} c(d) - \sum_{d \in [\mu]} c(d) = 0$$

and hence λ and μ satisfy the conditions in Corollary 4.3. However, there is no pairing of the boxes in $[\tau]$ such that the content of each pair sums to zero, and hence λ and μ cannot lie in the same block.

(ii) Let $\alpha = (5, 4^4)$, $\beta = (5, 1^4)$, $\gamma = \alpha/(\alpha \cap \beta)$, and $\delta = 2$. The diagrams $[\alpha]$, $[\beta]$, and $[\gamma]$ are illustrated (with their contents) in Figure 7. In this case the boxes in $[\gamma]$ can be put into pairs such that each pair sums to $1 - \delta = -1$, but the boxes with contents 0 and -1 are in configuration (b) from Figure 5, and occupy an odd number of columns. Hence α and β cannot lie in the same block.




FIGURE 8. The diagrams $[\alpha]$, $[\beta]$ and $[\gamma]$ in Example 4.9(ii)

By Corollary 4.8 weights which are not balanced will lie in different blocks. Hence for a B_n -module X we will denote by $\text{pr}_\lambda X$ the direct summand of X with composition factors $L_n(\mu)$ such that μ and λ are balanced.

Lemma 4.10. *Suppose that $\lambda \vdash n$ and $\epsilon_i \in \text{rem}(\lambda)$.*

(i) *There exists a B_n -module X and a short exact sequence*

$$0 \longrightarrow X \longrightarrow \text{pr}_\lambda \text{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon_i) \longrightarrow \Delta_n(\lambda) \longrightarrow 0.$$

Here $X \cong \Delta_n(\lambda - \epsilon_i - \epsilon_j)$ if $(\lambda - \epsilon_i - \epsilon_j, \lambda)$ is a balanced pair or $X = 0$ if no such ϵ_j exists. In the former case the sequence is non-split.

(ii) If

$$\mathrm{Hom}(\mathrm{pr}_\lambda \mathrm{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon_i), \Delta_n(\mu)) \neq 0$$

then $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$.

Proof. (i) The existence of such a sequence, and the form of X , follows from Proposition 2.7 and Corollary 4.8. To see that the sequence is non split, we proceed by induction on $|\lambda|$, the case where $\lambda = \emptyset$ being clear. By Frobenius reciprocity we have

$$\mathrm{Hom}(\Delta_{n-1}(\lambda - \epsilon_i), \mathrm{res} \Delta_n(\lambda - \epsilon_i - \epsilon_j)) \cong \mathrm{Hom}(\mathrm{ind} \Delta_{n-1}(\lambda - \epsilon_i), \Delta_n(\lambda - \epsilon_i - \epsilon_j)) \quad (9)$$

By (2) and Lemma 2.6(ii) the left-hand side equals

$$\mathrm{Hom}(\Delta_{n-1}(\lambda - \epsilon_i), \mathrm{ind} \Delta_{n-2}(\lambda - \epsilon_i - \epsilon_j)).$$

As $\Delta_{n-1}(\lambda - \epsilon_i)$ is simple, we have by the induction hypothesis and Theorem 4.4 that this Hom-space is one dimensional. Hence the right-hand side of (9) is also one dimensional, which by another application of Theorem 4.4 implies that the desired sequence is non-split as required.

Part (ii) is an immediate consequence of (i). \square

5. COMPUTING SOME COMPOSITION MULTIPLICITIES

So far we have concentrated on conditions which imply that weights lie in different blocks of the algebra. In this section we will find certain pairs of weights which do lie in the same block, which we will demonstrate by determining certain composition factors of standard modules, and homomorphisms between such modules.

We first consider the special case where the skew partition λ/μ is itself a partition. For such pairs we will be able to show precisely when $L_n(\lambda)$ is a composition factor of $\Delta_n(\mu)$. We first give a necessary condition, in Proposition 5.1, which is a generalisation of [DWH99, Corollary 9.1] (the latter only considers the case $\mu = \emptyset$ and homomorphisms rather than composition factors).

Proposition 5.1. *Let $\mu \subset \lambda$ are partitions such that $\nu = \lambda/\mu$ is also a partition. If*

$$[\Delta_n(\mu) : L_n(\lambda)] \neq 0$$

then $\nu = (a^b)$ where a is even and $b = \delta + a - 1 + 2c$, where c is the content of the top lefthand box of ν . Moreover, in this case we have

$$[\Delta_n(\mu) : L_n(\lambda)] = 1.$$

Proof. As usual, by localisation we can assume that λ is a partition of n . First suppose that $[\Delta_n(\mu) : L_n(\lambda)] \neq 0$. As $L_n(\lambda)$ is simply the lift of S^λ for $\mathbb{C}\Sigma_n$, we have that

$$[\mathrm{res}_{\mathbb{C}\Sigma_n} \Delta_n(\mu) : S^\lambda] \neq 0.$$

By (6) we have

$$[\mathrm{res}_{\mathbb{C}\Sigma_n} \Delta_n(\mu) : S^\lambda] = \sum_{\substack{\eta \vdash 2k \\ \eta \text{ even}}} c_{\mu\eta}^\lambda.$$

Hence we see that ν must be an even partition, and by Lemma 3.1 that $[\Delta_n(\mu) : L_n(\lambda)] = 1$.

On the other hand, using Proposition 4.5 we know that there is a pairing of the boxes of ν such that the sum of the content of the boxes in each pair is equal to $1 - \delta$. Clearly we have a submodule M of $\Delta_n(\mu)$ and an embedding

$$\Delta_n(\lambda) \hookrightarrow \Delta_n(\mu)/M.$$

If ϵ_i is any removable box of λ then we have a surjective homomorphism

$$\text{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon_i) \rightarrow \Delta_n(\lambda).$$

Composing these maps we see that

$$\text{Hom}(\text{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon_i), \Delta_n(\mu)/M) \neq 0$$

and so by Frobenius reciprocity we have

$$\text{Hom}(\Delta_{n-1}(\lambda - \epsilon_i), \text{res}_n(\Delta_n(\mu)/M)) \neq 0.$$

Thus

$$[\text{res}_n \Delta_n(\mu) : L_{n-1}(\lambda - \epsilon_i)] \neq 0$$

and hence either μ must have a removable box ϵ_j such that

$$[\Delta_{n-1}(\mu - \epsilon_j) : L_{n-1}(\lambda - \epsilon_i)] \neq 0$$

or μ must have an addable box ϵ_j such that

$$[\Delta_{n-1}(\mu + \epsilon_j) : L_{n-1}(\lambda - \epsilon_i)] \neq 0.$$

In the first case we have $\mu - \epsilon_j \subset \lambda - \epsilon_i$ and hence $c(\epsilon_j) = c(\epsilon_i)$. However, as λ/μ is a partition this is impossible, as no removable box in μ can have the same content as some box in λ/μ . Hence we must be in the second case with $\mu + \epsilon_j \subset \lambda - \epsilon_i$, so in fact ϵ_j must be a box in $\nu = \lambda/\mu$. As ν is a partition, there is only one such addable box and its content is given by c . Thus we must have

$$c(\epsilon_i) = 1 - \delta - c.$$

Now, if $\nu = \lambda/\mu$ had another removable box then it would have to have the same content. But different removable boxes have different contents. Hence ν can only have one removable box, i. e. it is a rectangle $\nu = (a^b)$, where a is even as ν must be an even partition. The content of the only removable box of ν inside of λ is given by $c + a - 1 - (b - 1) = c + a - b$ and this must be equal to $1 - \delta - c$. Hence we get

$$b = \delta - 1 + a + 2c$$

as required. \square

We will show that the condition in Proposition 5.1 is also sufficient. This generalises [DWH99, Theorem 9.2], which again only considers homomorphisms and the case $\mu = \emptyset$. Before doing this we will review some standard symmetric groups results which we will require. Details can be found in [Ful97, Chapter 7]

We will need to consider a set of idempotents $\{e_\lambda : \lambda \vdash n\}$ in $\mathbb{C}\Sigma_n$, such that $\mathbb{C}\Sigma_n e_\lambda \cong S^\lambda$. We will choose

$$e_\lambda = \frac{f^\lambda}{n!} \sum_{\sigma \in C_\lambda} \sum_{\tau \in R_\lambda} \text{sgn}(\sigma)\sigma\tau \quad (10)$$

where $f^\lambda = \dim S^\lambda$, C_λ is the column stabiliser of $[\lambda]$ and R_λ is the row stabiliser of $[\lambda]$. For example $e_{(2)}$ and $e_{(1,1)}$ (regarded as elements of B_2) are illustrated in Figure 9.

FIGURE 9. The elements $e_{(2)}$ and $e_{(1,1)}$

We will also need the fact that

$$\operatorname{ind}_{\mathbb{C}(\Sigma_a \times \Sigma_b)}^{\mathbb{C}\Sigma_{a+b}} (S^\mu \otimes S^\nu) \cong \bigoplus_{\lambda \vdash (n+m)} c_{\mu\nu}^\lambda S^\lambda.$$

As all these group algebras are semisimple, this implies by Frobenius reciprocity that

$$\operatorname{res}_{\mathbb{C}(\Sigma_a \times \Sigma_b)}^{\mathbb{C}\Sigma_n} S^\lambda \cong \bigoplus_{\mu \vdash a, \nu \vdash b} c_{\mu\nu}^\lambda (S^\mu \otimes S^\nu). \quad (11)$$

Particular values of $c_{\mu\nu}^\lambda$ which we will need are those where $\nu = (2)$, respectively $\nu = (1,1)$. In these cases $c_{\mu\nu}^\lambda$ is at most 1, and is non-zero precisely when λ/μ consists of two boxes in different columns, respectively different rows.

Theorem 5.2. *Supppose that $\mu \subset \lambda$ and $\lambda/\mu = \nu = (a^b)$. If a is even and $b = \delta - 1 + a + 2c$ where c is the content of the top left box of ν then*

$$[\Delta_n(\mu) : L_n(\lambda)] = 1.$$

Moreover, if $\lambda \vdash n$ then

$$\operatorname{Hom}_{B_n}(L_n(\lambda), \Delta_n(\mu)) = \mathbb{C}.$$

Proof. We can assume without loss of generality that $\lambda \vdash n$. We have seen in the proof of Proposition 5.1 that $[\operatorname{res}_{\mathbb{C}\Sigma_n} \Delta_n(\mu) : S^\lambda] = 1$. Let $W = e_\lambda \Delta_n(\mu)$, which is isomorphic to S^λ as a Σ_n -module. To show this is in fact a B_n -submodule of $\Delta_n(\mu)$, it will be enough to show that $X_{i,j}W = 0$ for all $1 \leq i < j \leq n$. Indeed, it is enough to show that this holds for a single choice of i and j , as

$$\sigma X_{i,j} \sigma^{-1} = X_{\sigma(i), \sigma(j)}$$

for all $\sigma \in \Sigma_n$.

So let us fix i and j with $1 \leq i < j \leq n$ and use the embedding

$$\Sigma_{n-2} \times \Sigma_2 \subset \Sigma_n$$

where Σ_2 is the symmetric group on $\{i, j\}$ and Σ_{n-2} the symmetric group on $\{1, \dots, n\} \setminus \{i, j\}$. By (11) and the remarks following we have

$$\operatorname{res}_{\mathbb{C}(\Sigma_{n-2} \times \Sigma_2)} W \cong \bigoplus_{\alpha \vdash n-2} (S^\alpha \otimes S^{(1,1)}) \bigoplus_{\beta \vdash n-2} (S^\beta \otimes S^{(2)})$$

where we sum over all α 's obtained from λ by removing 2 boxes in different rows and over all β 's obtained from λ by removing two boxes in different columns.

The map $X_{i,j} : \Delta_n(\mu) \longrightarrow \Delta_n(\mu)$ is a $\mathbb{C}\Sigma_{n-2} \times \mathbb{C}\Sigma_2$ -homomorphism. Note that we have $X_{i,j}(\Delta_n(\mu)) \subset U$ where U is the span of all elements of the form $X_{w,1,id} \otimes x$ where w has an arc between i and j and $x \in S^\mu$. Regarding U as a B_{n-2} -module acting on the strings excluding i and j it is easy to see that U is isomorphic to $\Delta_{n-2}(\mu)$, and the restriction of this action to $\mathbb{C}\Sigma_{n-2}$ is the same as restriction to the action of the first component of $\mathbb{C}\Sigma_{n-2} \times \mathbb{C}\Sigma_2$ regarded as a subalgebra of B_n . Also, it is clear that X_{ij} kills the element $e_{(1,1)}$ in Figure 9, and hence kills the simple module $S^{(1,1)}$. Combining these observations with (6) we deduce that, as a $\mathbb{C}\Sigma_{n-2} \times \mathbb{C}\Sigma_2$ -module, U decomposes as

$$U = \bigoplus_{\tau} c_{\tau}(S^{\tau} \otimes S^{(2)})$$

where

$$c_{\tau} = \sum_{\substack{\tau \vdash n-2 \\ \eta \text{ even}}} c_{\mu\eta}^{\tau}.$$

Consider the restriction $X_{i,j} : W \longrightarrow U$. We want to show that $X_{i,j}W = 0$. Look at the simple summands of W . Every summand of the form $S^{\alpha} \otimes S^{(1,1)}$ is sent to zero as it does not appear in U . Moreover, if μ is not contained in β then $S^{\beta} \otimes S^{(2)}$ is sent to zero as U only contains simple modules $S^{\eta} \otimes S^{(2)}$ with $\mu \subset \eta$. So we only need to show that

$$X_{i,j}(S^{\beta} \otimes S^{(2)}) = 0$$

for any $\beta \vdash n-2$ with $\mu \subset \beta$ and β obtained from λ by removing two boxes in different columns. But there is only one such β , namely the partition obtained from λ by removing two boxes from the last row of ν , i.e. $\beta/\mu = (a^{b-1}, a-2)$, and by Lemma 3.1 the coefficient of $S^{\beta} \otimes S^{(2)}$ in U equals 1.

Write $W = V \oplus Y$ where $V = S^{\beta} \otimes S^{(2)}$. As V is simple, either $X_{i,j}$ embeds V into U or $X_{i,j}V = 0$. Label the boxes of the partition λ with the numbers $1, 2, \dots, n$ starting with the first row from left to right, then the second row from left to right, etc., until the last row. Say that the last box of the partition $\nu = (a^b)$ inside of λ is labelled by l . Up until now $X_{i,j}$ was arbitrary; we now fix $i = l-1$ and $j = l$ and we want to show that $X_{l-1,l}V = 0$.



Fix a partial one-row diagram w_0 with t arcs defined as follows: suppose the u -th row of ν inside of λ is labelled by $x_u, x_u + 1, \dots, x_u + a - 1$ for $1 \leq u \leq b$, as illustrated in Figure 10. Then w_0 is defined to have arcs $\{x_u, x_u + 1\}, \{x_u + 2, x_u + 3\}, \dots, \{x_u + a - 2, x_u + a - 1\}$ for $1 \leq u \leq b$. (Note that $x_b + a - 1 = l$.) We will represent elements of $V_{n,t}$ by adding bars to the Young tableau joining each pair of nodes connected by an arc. Thus the element w_0 will be represented by the diagram in Figure 11. Usually we will only represent the boxes of ν in such a diagram.

Now consider the element of $\Delta_n(\mu)$ given by $X_{w_0,1,id} \otimes x$ for some $x \in S^\mu$. Then $e_{\lambda}(X_{w_0,1,id} \otimes x) \in W$, so it decomposes as

$$e_{\lambda}(X_{w_0,1,id} \otimes x) = v + y$$

where $v \in V$ and $y \in Y$. Note that this decomposition is independent of δ . As observed above, we have $X_{l-1,l}e_{\lambda}(X_{w_0,1,id} \otimes x) = X_{l-1,l}v$. Consider the coefficient of $X_{w_0,1,id} \otimes x$ in $X_{l-1,l}v$. We will show that it is a non-zero multiple of

$$\delta - 1 + a - b + 2c.$$

FIGURE 10. The labelling of λ , with ν shaded and μ unshadedFIGURE 11. A diagrammatic representation of the element w_0

Hence, as v is independent of δ we see that $v \neq 0$, but when $\delta - 1 + a - b + 2c = 0$, we have $X_{l-1,l}v = 0$. Thus $X_{l-1,l}$ cannot embed V into U and so it must map V to zero.

Using the labelling of the boxes of λ defined above, we will identify the row and column stabilisers R_λ and C_λ as subgroups of Σ_n , the symmetric group on $\{1, \dots, n\}$. From (10) we have

$$e_\lambda(X_{w_0,1,id} \otimes x) = \frac{f^\lambda}{n!} \sum_{\sigma \in C_\lambda} \sum_{\tau \in R_\lambda} \text{sgn}(\sigma) \sigma \tau (X_{w_0,1,id} \otimes x),$$

and so

$$X_{l-1,l} e_\lambda(X_{w_0,1,id} \otimes x) = \frac{f^\lambda}{n!} \sum_{\sigma \in C_\lambda} \sum_{\tau \in R_\lambda} \text{sgn}(\sigma) X_{l-1,l} \sigma \tau (X_{w_0,1,id} \otimes x).$$

We want to find the coefficient of $X_{w_0,1,id} \otimes x$ in this sum. We consider several cases.

Case 1: Suppose that $\sigma \tau X_{w_0,v_k,id}$ has an arc $\{l-1, l\}$.

In this case $X_{l-1,l}\sigma\tau(X_{w_0,1,id} \otimes x) = \delta\sigma\tau(X_{w_0,1,id} \otimes x)$. If we want $\sigma\tau(X_{w_0,1,id} \otimes x)$ to be in $\text{span}\{X_{w_0,1,id} \otimes S^\mu\}$ then we must have

$$\begin{aligned}\tau &= \tau_1\tau_2 && \text{with } \tau_1 \in R_{\mu\subset\lambda}, \tau_2 \in R_\lambda^0 \\ \sigma &= \sigma_1\sigma_2 && \text{with } \sigma_1 \in C_{\mu\subset\lambda}, \sigma_2 \in C_\lambda^0\end{aligned}$$

where $R_{\mu\subset\lambda}$ denotes the subgroup of R_λ (isomorphic to R_μ) which preserves the rows of μ and fixes everything in ν and R_λ^0 denotes the subgroup of R_λ which fixes $X_{w_0,1,id}$ as a diagram (i.e. fixes all but the t northern arcs, which may be permuted amongst themselves and be reversed). In a similar way we define $C_{\mu\subset\lambda}$ and C_λ^0 .

Set $r = |R_\lambda^0|$. As the a columns of ν are paired by the bars in w_0 , and each pair of such columns may be permuted freely by C_λ^0 we have $|C_\lambda^0| = (b!)^{a/2}$. Moreover $\text{sgn}(\sigma_2) = 1$ as σ_2 is an even permutation (as it is made up of pairs of identical permutations, corresponding to the paired ends of a bar) and so $\text{sgn}(\sigma) = \text{sgn}(\sigma_1)$. Hence in this case we get the contribution

$$\begin{aligned}& \frac{f^\lambda}{n!} \sum_{\sigma_2 \in C_\lambda^0} \sum_{\sigma_1 \in C_{\mu\subset\lambda}} \sum_{\tau_2 \in R_\lambda^0} \sum_{\tau_1 \in R_{\mu\subset\lambda}} \text{sgn}(\sigma_1\sigma_2)\sigma_1\sigma_2\tau_1\tau_2(X_{w_0,1,id} \otimes x) \\ &= \frac{f^\lambda}{n!} \sum_{\sigma_2 \in C_\lambda^0} \sum_{\tau_2 \in R_\lambda^0} \sigma_2\tau_2(X_{w_0,1,id} \otimes \sum_{\sigma_1 \in C_{\mu\subset\lambda}} \sum_{\tau_1 \in R_{\mu\subset\lambda}} \text{sgn}(\sigma_1)\sigma_1\tau_1(x)) \\ &= \frac{f^\lambda}{n!} \frac{|\mu|}{f^\mu} \sum_{\sigma_2 \in C_\lambda^0} \sum_{\tau_2 \in R_\lambda^0} \sigma_2\tau_2(X_{w_0,1,id} \otimes e_\mu(x)) \\ &= \frac{f^\lambda}{n!} \frac{|\mu|}{f^\mu} r(b!)^{a/2}(X_{w_0,1,id} \otimes x)\end{aligned}$$

using for the second equality the isomorphisms $C_{\mu\subset\lambda} \cong C_\mu$ and $R_{\mu\subset\lambda} \cong R_\mu$, and for the final equality the fact that $e_\mu(x) = x$ for all $x \in S^\mu$.

Case 2: Suppose that neither $l-1$ nor l is part of an arc in $\sigma\tau X_{w_0,1,id}$.

In this case $X_{l-1,l}\sigma\tau X_{w_0,1,id}$ has $t+1$ arcs in the top row and so $X_{l-1,l}(X_{w_0,1,id} \otimes x) = 0$.

Case 3: Suppose that in $\sigma\tau X_{w_0,v_k,id}$ there are arcs $\{l-1, i\}$ and $\{l, j\}$.

In this case, $X_{l-1,l}\sigma\tau X_{w_0,1,id}$ is obtained from $\sigma\tau X_{w_0,1,id}$ by replacing the arcs $\{l-1, i\}$ and $\{l, j\}$ by the arcs $\{i, j\}$ and $\{l-1, l\}$. Hence if we want to have $X_{l-1,l}\sigma\tau(X_{w_0,1,id} \otimes x)$ lying in $\text{span}\{X_{w_0,1,id} \otimes S^\mu\}$ then $\{i, j\}$ must be an arc of w and $i = j \pm 1$. Here we consider two subcases.

Subcase 3(a): First assume that the pair $\{i, j\}$ is not in the last double column. Then $\tau = \tau_2\tau_1$ with $\tau_1 \in R_{\mu\subset\lambda}$ and $\tau_2 \in \tilde{\tau}R_\lambda^0$, where $\tilde{\tau} = (u-1, v)$ or (u, v) such that v is a box of ν in the same column as l (possibly l itself) and u is the box of ν in the same row as v and in the same column as $\max(i, j)$. An example of such a situation is illustrated in Figure 12.

Thus we have b choices for v and $(\frac{a}{2} - 1)$ choices for the position of $\{i, j\}$ (and hence of u), and so there are $2b(\frac{a}{2} - 1)$ choices for $\tilde{\tau}$. Hence there are $2rb(\frac{a}{2} - 1)$ choices for τ_2 . Now $\sigma = \sigma_2\sigma_1$ where $\sigma_1 \in C_{\mu\subset\lambda}$, and σ_2 permutes the pairs in all double columns, except the last and the double column containing $\{j-1, j\}$, arbitrarily. In the last double column it must send $v-1$ to $l-1$ and v to l , and in the double column containing $\{j-1, j\}$, it can




FIGURE 12. An example of subcase 3(a)

permute the pairs in any way (as $\{j-1, j\}$ can be any pair in this double column). So we get $(b!)^{\frac{a}{2}-2}(b-1)!b!$ possibilities for σ_2 . Note also that σ_2 is always an even permutation and so $\text{sgn}(\sigma) = \text{sgn}(\sigma_1)$. Thus in this subcase, we get a contribution of

$$\begin{aligned} & \frac{f^\lambda}{n!} 2r b \left(\frac{a}{2} - 1\right) (b!)^{\left(\frac{a}{2}-2\right)} (b-1)! b! X_{w_0,1,id} \otimes \sum_{\sigma_1 \in C_{\mu \subset \lambda}} \sum_{\tau_1 \in R_{\mu \subset \lambda}} \text{sgn}(\sigma_1) \sigma_1 \tau_1(x) \\ &= \frac{f^\lambda}{n!} \frac{|\mu|!}{f^\mu} r (a-2) (b!)^{\frac{a}{2}} X_{w_0,1,id} \otimes x \end{aligned}$$

where the equality follows as in Subcase 1.

Subcase 3(b): Next assume that the pair $\{i, j\}$ is in the last column. We must have $\tau = \tau_2 \tau_1$ where $\tau_1 \in R_{\mu \subset \lambda}$ and $\tau_2 \in R_\lambda^0$. Also $\sigma = \sigma_2 \sigma_1$ where $\sigma_1 \in C_{\mu \subset \lambda}$ and $\sigma_2 \in (j, l)C_\lambda^0$. We have $b-1$ choices for j being a box of ν in the same column as l . Note that in this case $\text{sgn}(\sigma_2) = -1$ and so $\text{sgn}(\sigma) = -\text{sgn}(\sigma_1)$. Hence arguing as in Subcases 1 and 3(a) we get a contribution of

$$-\frac{f^\lambda}{n!} \frac{|\mu|!}{f^\mu} r (b-1) (b!)^{\frac{a}{2}} X_{w_0,1,id} \otimes x.$$

Case 4: Suppose that in $\sigma\tau X_{w_0,1,id}$ there is a link from $l-1$ to i , say, and l is not part of an arc (or vice versa).

In this case $X_{l-1,l}\sigma\tau X_{w_0,1,id}$ is obtained from $\sigma\tau X_{w_0,1,id}$ by replacing the arc $\{i, l-1\}$ (or $\{i, l\}$) with the arc $\{l-1, l\}$ and i is not part of an arc any more. So, if we want to have $X_{l-1,l}\sigma\tau X_{w_0,1,id}$ in $\text{span}\{X_{w_0,1,id} \otimes S^\mu\}$ then i cannot be one of the boxes of ν . There are various potential subcases that can arise. After action by an element of $R_{\mu \subset \lambda}$ the element i may be in any box in the same row of μ . There are three cases: (a) i is now to the left of the first column of ν ; (b) i is above ν but not above $l-1$ or l ; (c) i is above $l-1$ or l .

Subcase 4(a): First, assume that the box i is in a column to the left of ν in λ . In this case, $\tau = \tau_2 \tau_1$ where $\tau_1 \in R_{\mu \subset \lambda}$ (as we have already acted by such an element to put i in this case above) and $\tau_2 \in (v-1, u)R_\lambda^0$, or $\tau_2 \in (v, u)R_\lambda^0$ where v is any box in ν in the same column as l and u is the box of μ in the same row as v and in the same column as i . An example of such a situation is illustrated in Figure 13.

Let c_1 be the number of columns of λ to the left of ν . Then there are $2rbc_1$ possible choices of τ_2 . Now $\sigma = \sigma_2 \sigma_1$ where $\sigma_1 \in C_{\mu \subset \lambda}$ (as i is an arbitrary element in its column of μ) and σ_2 permutes the pairs in each of the first $(\frac{a}{2}-1)$ double columns of ν arbitrarily, and in the last double column sends $v-1$ to $l-1$ and v to l and then permutes the other




FIGURE 13. An example of subcase 4(a)

pairs arbitrarily. Note that $\text{sgn}(\sigma_1) = \text{sgn}(\sigma)$. Hence (arguing as in earlier cases) we get a contribution of

$$\begin{aligned} & \frac{f^\lambda}{n!} \frac{|\mu|!}{f^\mu} 2 r b c_1 (b!)^{\left(\frac{a}{2}-1\right)} (b-1)! X_{w_0,1,id} \otimes x \\ &= \frac{f^\lambda}{n!} \frac{|\mu|!}{f^\mu} r (b!)^{\frac{a}{2}} 2c_1 X_{w_0,1,id} \otimes x. \end{aligned}$$

Subcase 4(b): Suppose that i is a box of μ which is above some column of ν but to the left of $l-1$. Then the only way to use row and column permutations not involving $R_{\mu \subset \lambda}$ (which we have already used to position i) to connect i and l (or $l-1$) is by some pair τ and σ similar to that shown in Figure 14. But (as illustrated) any such pair does not preserve the remaining edges in ν . Hence this subcase cannot arise.




FIGURE 14. An example of the impossibility of subcase 4(b)

Subcase 4(c): Finally we are left with the subcase where after action by $R_{\mu \subset \lambda}$ the element i is in a box of μ which is either in the same column as $l-1$ or in the same column as l . In this case $\tau = \tau_2 \tau_1$ where $\tau_1 \in R_{\mu \subset \lambda}$ and $\tau_2 \in R_\lambda^0$. Also, $\sigma = \sigma_2 \sigma_1$ where $\sigma_1 \in C_{\mu \subset \lambda}$ (as i is an arbitrary element in its column of μ) and either $\sigma_2 \in (i, l)C_\lambda^0$ or $\sigma_2 \in (i, l-1)C_\lambda^0$. If c_2 is the number of columns above ν in λ then there are $2c_2$ choices for the position of i . Note that here $\text{sgn}(\sigma_2) = -1$ and so $\text{sgn}(\sigma) = -\text{sgn}(\sigma_1)$. Hence, in this case we get a contribution of

$$-\frac{f^\lambda}{n!} \frac{|\mu|!}{f^\mu} r (b!)^{\frac{a}{2}} 2c_2 X_{w_0,1,id} \otimes x.$$

Note that the final sets of permutations obtained in Subcases 4(a) and 4(c) are disjoint, so there is no double counting in these contributions. Now on adding up all contributions

from Cases 1–4 we see that the coefficient of $X_{w_0,1,id} \otimes x$ inside of $X_{l-1,l}e_\lambda(X_{w_0,1,id} \otimes x)$ is given by

$$\frac{f^\lambda}{n!} \frac{|\mu|!}{f^\mu} r(b!)^{\frac{a}{2}} (\delta - 1 + a - b + 2(c_1 - c_2)).$$

The content of the top left box of the partition ν inside the partition λ is given by $c = (c_1 + 1) - (c_2 + 1) = c_1 - c_2$. Thus we have proved that this coefficient is a non-zero multiple of $(\delta - 1) + a - b + 2c$ as required. \square

6. THE BLOCKS OF THE BRAUER ALGEBRA

In section 4 we saw that a necessary condition for two weights λ and μ to be in the same block was that the pair was balanced. We will now show that this condition is also sufficient. The key idea will be to construct from any partition λ in a balanced pair with some $\mu \subset \lambda$ a partition $\nu \subset \lambda$ and a homomorphism connecting $\Delta_n(\lambda)$ and $\Delta_n(\nu)$. This will allow us to proceed by induction.

Given a partition λ we denote by $\text{add}(\lambda)$ the set of *addable* boxes of λ (i.e. the set of boxes which may be added to λ such that the new shape is still a partition). Similarly we denote by $\text{rem}(\lambda)$ the set of *removable* boxes of λ . If $\mu \subset \lambda$ then we denote the set of boxes in $\text{rem}(\lambda)$ which are also boxes of λ/μ by $\text{rem}(\lambda/\mu)$. Distinct boxes in $\text{add}(\lambda)$ (respectively in $\text{rem}(\lambda)$) have distinct contents, and we will identify such boxes by their contents. We will order the boxes in λ with a given content by saying that box ϵ is *smaller* than box ϵ' if ϵ appears on an earlier row than ϵ' .

Definition 6.1. Suppose that $\mu \subset \lambda$ is a balanced pair. For each $\epsilon_i \in \text{rem}(\lambda/\mu)$ we wish to consider μ^i , the *i-maximal balanced subpartition between μ and λ* . This is the maximal partition $\mu^i \subset \lambda$ such that μ^i does not contain ϵ_i and λ and μ^i form a balanced pair. We will construct μ^i by recursively defining a series of skew partitions $(\lambda/\mu^i)_j$ which will eventually equal the skew partition λ/μ^i . There is by the pairing condition a maximal box (i.e. all others smaller) with content $c(\epsilon'_i)$ such that $c(\epsilon_i) + c(\epsilon'_i) = 1 - \delta$. Let $(\lambda/\mu^i)_0 = \{\epsilon_i, \epsilon'_i\}$. Given $(\lambda/\mu^i)_m$, we set

$$(\lambda/\mu^i)_{m+1} = (\lambda/\mu^i)_m \cup A_{m+1} \cup A'_{m+1}$$

where A_{m+1} is the set of boxes ϵ in λ such that ϵ is to the right of or below a box in $(\lambda/\mu^i)_m$, and A'_{m+1} is the set of boxes ϵ' in (λ/μ) such that $c(\epsilon) + c(\epsilon') = 1 - \delta$ for some $\epsilon \in A_{m+1}$ and ϵ' is maximal with such content among the boxes of λ/μ not already in $(\lambda/\mu^i)_m$.

This iterative process eventually stabilises, and we obtain $(\lambda/\mu^i)_t$ which is a (possibly disconnected) subset of the edge of λ/μ , having width one. (In particular it does not contain two boxes with the same content.) If δ is even and $(\lambda/\mu^i)_t$ does not contain a vertical pair of boxes with content $\frac{2-\delta}{2}$ and $-\frac{\delta}{2}$, or δ is odd and $(\lambda/\mu^i)_t$ does not contain a box of content $\frac{1-\delta}{2}$ then we set $\lambda/\mu^i = (\lambda/\mu^i)_t$. Otherwise if δ is even we set

$$(\lambda/\mu^i)_{t+1} = (\lambda/\mu^i)_t \cup \{x, y\} \tag{12}$$

where x, y are the maximal boxes in λ of content $\frac{2-\delta}{2}$ and $-\frac{\delta}{2}$ not in $(\lambda/\mu^i)_t$, and if δ is odd we set

$$(\lambda/\mu^i)_{t+1} = (\lambda/\mu^i)_t \cup \{z\} \tag{13}$$

where z is the maximal box in λ of content $\frac{1-\delta}{2}$ not in $(\lambda/\mu^i)_t$. This new skew partition is not necessarily stable under the addition of boxes A and A' as above, and we repeat that process again until the skew partition eventually stabilises at some step s . We then set $\lambda/\mu^i = (\lambda/\mu^i)_s$. Thus λ/μ^i is a removable subset of λ/μ having width at most two (so at most two boxes with any given content).




FIGURE 15. Two examples of the λ/μ^i construction

Example 6.2. We will now consider several examples of this construction. First let $\lambda = (6, 5, 5, 2, 1)$ and $\mu = (6, 4, 1)$; this is a balanced pair for $\delta = 2$. If ϵ_i is any of the removable boxes in Figure 15(a), then λ/μ^i is the shaded region shown. For an example where the resulting skew partition is connected, consider $\lambda = (7, 6, 5, 5, 2, 2)$ and $\mu = (7, 4, 4, 1, 1)$. This is a balanced pair for $\delta = 2$. If ϵ_i is any of the removable boxes in λ/μ then the skew partition λ/μ^i is the shaded region shown in Figure 15(b). In this case there is a pair of boxes in the skew partition with contents $\frac{2-\delta}{2}$ and $-\frac{\delta}{2}$ (i.e. 0 and -1), but we do not get a strip of width 2 because these boxes are not vertically aligned.

For an example of the full iterative process consider $\lambda = (7, 6, 4^4, 1^2)$ and $\mu = (4, 3^4)$. This is a balanced pair for $\delta = 2$, and after the first part of the iterative process the skew partition stabilises into the lightly shaded region shown in Figure 16(a). However, we now have a vertical pair in the skew partition with contents $\frac{2-\delta}{2}$ and $-\frac{\delta}{2}$ (i.e. 0 and -1). Thus we have to apply (12), and add the darkly shaded boxes with content 0 and -1 to this skew partition. The complement of this is no longer a partition, so we remove the remaining darkly shaded region by one further application of the iterative procedure.




FIGURE 16. More examples of the λ/μ^i construction

Definition 6.3. We now wish to define a *maximal balanced subpartition between μ and λ* , which we will denote by λ/μ' . Having constructed a skew partition λ/μ^i for each removable box ϵ_i of λ , we partially order this collection by inclusion. We then take λ/μ' to be some minimal element of this set.

Example 6.4. To see a non-trivial example of this choice, consider $\lambda = (7, 6^2, 5, 4^2, 2)$ and $\mu = (5, 3, 2^3, 1)$. This is a balanced pair for $\delta = 1$, but has several different associated skew partitions. If we take ϵ_i to be one of the removable boxes labelled by 6 or -5 then λ/μ^i equals the entire shaded region in Figure 16(b). However, if we take ϵ_j to be any of the other removable boxes then λ/μ^j consists of the six darkly shaded boxes. As $\lambda/\mu^j \subset \lambda/\mu^i$, we take λ/μ' to equal λ/μ^j in this case, and hence $\mu' = (7, 6, 4^2, 3, 2^2)$. (Note that if this example had one additional box of content 0 between the two darkly shaded regions, then we would have to apply (13) and this box would have associated skew partition all of the darkly shaded region together with itself and the diagonally adjacent box with content 0.)

The importance of this construction is given by

Theorem 6.5. *If $\mu \subset \lambda$ is a balanced pair, then for any maximal balanced subpartition μ' between μ and λ we have*

$$\text{Hom}(\Delta_n(\lambda), \Delta_n(\mu')) \neq 0.$$

Proof. As usual, we may assume that λ is a partition of n . Pick $\epsilon \in \text{rem}(\lambda/\mu')$ with $|c(\epsilon) - \frac{1-\delta}{2}|$ maximal. (Note that there are at most two such boxes.) If δ is even and $c(\epsilon) = \frac{-\delta}{2}$ or $c(\epsilon) = \frac{2-\delta}{2}$ then λ/μ' is one of the two cases in Figure 17(a) or (b), while if δ is odd and $c(\epsilon) = \frac{1-\delta}{2}$ then λ/μ' is as in Figure 17(c). In each of these cases there is a non-zero homomorphism from $\Delta_n(\lambda)$ to $\Delta_n(\mu')$ by Theorem 5.2 (or more directly by repeated applications of Frobenius reciprocity). Thus we henceforth assume we are not in any of these cases.




FIGURE 17. Some small ϵ cases, with matched box denoted by ϵ'

Suppose that ϵ is paired with a maximal ϵ' of content $1 - \delta - c(\epsilon)$. We will assume that ϵ is above, or to the right of, ϵ' , and leave the (obvious) modifications required for the other case to the reader.

We will be able to proceed by induction using the following claim.

Claim 6.6. (i) *There is no box of content $c(\epsilon)$ in $\text{rem}(\mu')$.*
(ii) *There is a unique box ϵ' of content $1 - \delta - c(\epsilon)$ in $\text{add}(\mu')$.*
(iii) *If $|\lambda/\mu'| > 2$ then the pair $\lambda - \epsilon$ and $\mu' + \epsilon'$ is balanced, and the associated skew partition is minimal in the set of those of the form $(\lambda - \epsilon)/(\mu + \epsilon')^k$, with ϵ_k in $\text{rem}((\lambda - \epsilon)/(\mu' + \epsilon'))$. Equivalently, for every ϵ_k in $\text{rem}((\lambda - \epsilon)/(\mu' + \epsilon'))$ we have*

$$(\lambda - \epsilon)/(\mu + \epsilon')^k = (\lambda - \epsilon)/(\mu + \epsilon').$$

Before proving this claim, we show how it can be used to complete the proof of Theorem 6.5. Note that if $\lambda - \epsilon$ has a removable box τ with content $1 - \delta - c(\epsilon)$ then by minimality $\lambda/\mu' = \{\epsilon, \tau\}$, and we are done by Theorem 4.4 and our assumptions on λ . Thus we assume that there is no such removable box. By Frobenius reciprocity, Corollary 4.8, and Lemma 4.10, we have

$$\begin{aligned} \text{Hom}(\Delta_n(\lambda), \Delta_n(\mu')) &\cong \text{Hom}(\text{pr}_\lambda \text{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon), \Delta_n(\mu')) \\ &\cong \text{Hom}(\Delta_{n-1}(\lambda - \epsilon), \text{pr}_{\lambda - \epsilon} \text{res}_n \Delta_n(\mu')). \end{aligned}$$

By the first two parts of Claim 6.6 this latter Homspace is isomorphic to

$$\text{Hom}(\Delta_{n-1}(\lambda - \epsilon), \Delta_{n-1}(\mu' + \epsilon'))$$

and by the final part of Claim 6.6 (and induction) this is non-zero as required.

Thus it only remains to prove Claim 6.6.

Proof of Claim 6.6: (i) First suppose that there is only one box in λ/μ' with content $c(\epsilon)$. By construction, if there are any boxes above ϵ in λ/μ' then the one with largest content, or its matched pair, is removable. But this contradicts the choice of ϵ . The other possibility is that there is a second box τ in λ/μ' with content $c(\epsilon)$, occupying the opposite corner of a two by two square. Arguing as in the previous case, if there are any boxes in λ/μ' above this square then this contradicts the choice of ϵ . These two cases are illustrated in Figure 18(a) and (b). In both these cases we deduce that μ' cannot have a removable box of content $c(\epsilon)$, as there must be boxes to the right of any such box in μ' .




FIGURE 18. Two corner cases

(ii) Note that if λ/μ' consists of two boxes then the result is obvious, so we assume this is not the case. It is also clear that any addable box of a given content must be unique. Let ϵ' be the maximal box in λ/μ' with content $1 - \delta - c(\epsilon)$.

First suppose that λ/μ' has only one box with content $c(\epsilon)$, so that we are in the case shown in Figure 18(a). The box ϵ' paired with ϵ in Figure 18(a) must be to the right or above ϵ' , and hence we are in one of the two configurations shown in Figure 19.




FIGURE 19. The first corner case

The case in Figure 19(a) is impossible by our assumption on the size of λ/μ' (and minimality), as both ϵ and ϵ' are removable boxes. In the remaining case it is clear that μ' has addable box ϵ' , as required.

Next suppose that λ/μ' has two boxes with content $c(\epsilon)$, so that we are in the case shown in Figure 18(b). As in the previous case, the box α' paired with α must be to the right or above ϵ' . If it is above then we have a configuration similar to that in Figure 19(a), and hence ϵ' is a removable box. But this is impossible exactly as for the case in Figure 19(a). Hence α' must be to the right of ϵ' , and we must have a configuration as in Figure 20. But this configuration clearly has an addable box, τ' , of content $c(\epsilon')$.




FIGURE 20. The second corner case

(iii) The two partitions $\lambda - \epsilon$ and $\mu' + \epsilon'$ are clearly balanced. For minimality we consider the various cases that can arise. If we are in the case shown in Figure 18(a), then paired boxes are as shown in Figure 19(b). Suppose for a contradiction that $(\lambda - \epsilon)/(\mu' + \epsilon')$ is not minimal, and hence contains a smaller skew partition η . If η does not involve $*$ and $*'$ then it is also contained in λ/μ' , which contradicts the minimality of this original pair. If η does involve $*$ and $*'$ then this contradicts λ/μ' being minimal, as λ/μ' contains $\eta \cup \{*, *'\}$, which is a smaller sub-skew partition of λ/μ' .

Now consider the case shown in Figure 18(b), where the paired boxes are as in Figure 20. As before, suppose for a contradiction that $(\lambda - \epsilon)/(\mu' + \tau')$ is not minimal, and hence contains a smaller skew partition η . If η does not involve α and α' then it is also contained in λ/μ' . If η does involve α and α' but not τ and ϵ' , then $\eta \cup \{\epsilon, \epsilon'\}$ is a removable skew inside λ/μ' . Finally, if η involves all of α , α' , τ , and ϵ' , then η must also involve β and β' . Now the skew obtained from η by replacing τ by ϵ can be removed from λ/μ' . In each of these three cases we have found a proper removable skew inside λ/μ' , which contradicts the minimality of λ/μ' . Thus $(\lambda - \epsilon)/(\mu' + \tau')$ must be minimal, which completes the proof of Claim 6.6, and hence also of Theorem 6.5. \square

Corollary 6.7. *Two weights λ and μ are in the same block of B_n if and only if they are balanced. Each block contains a unique minimal weight.*

Proof. In Corollary 4.8 we proved that two weights in the same block must be balanced. For the reverse implication, we will proceed by induction. By Theorem 6.5, if λ contains a smaller partition μ with which it is balanced, then there exists some $\mu' \subset \lambda$ with a non-zero homomorphism from $\Delta_n(\lambda)$ to $\Delta_n(\mu')$. In particular, λ and μ' will lie in the same block of B_n . Thus it is enough to show that there is a unique minimal partition in the set of partitions which are balanced with λ .

But if there are two such minimal partitions μ and ν , then set $\eta = \mu \cap \nu$. Clearly η is a partition, and it forms a balanced pair with both μ and ν (and hence with λ). This contradicts our assumption of minimality \square

We conclude this section with a description of the minimal partitions in each block (and hence give a parametrisation of the blocks). We begin by constructing inductively a skew partition $\hat{\lambda}$ related to λ . Let $\lambda(0) = \lambda$. Given $\lambda(i)$, consider $\epsilon \in \text{rem}(\lambda(i))$ such that $|c(\epsilon) - \frac{1-\delta}{2}|$ is maximal. Suppose that there does not exist $\epsilon' \in [\lambda]$ with $c(\epsilon) + c(\epsilon') = 1 - \delta$ and $\epsilon' \neq \epsilon$. Hence either the set of rows λ^t above and including the row containing ϵ (if $c(\epsilon) - \frac{1-\delta}{2} > 0$) or the set of columns λ^l to the left of and including the column containing ϵ (if $c(\epsilon) - \frac{1-\delta}{2} < 0$) cannot be removed. In this case set $\lambda(i+1) = \lambda(i)/\lambda^t$, respectively $\lambda(i+1) = \lambda(i)/\lambda^l$. If there exists $\epsilon' \in [\lambda]$ with $c(\epsilon) + c(\epsilon') = 1 - \delta$ and $\epsilon' \neq \epsilon$ then $\hat{\lambda} = \lambda(i)$. This procedure will eventually terminate in the construction of $\hat{\lambda}$.




FIGURE 21. An example of the construction of $\hat{\lambda}$

Example 6.8. As an example of this construction, consider $\delta = 1$ and $\lambda = (7^2, 6, 5, 4, 2, 1^2)$, as illustrated in Figure 21. At the first stage, we take ϵ to be the box labelled -7 , and hence remove the first column. Next we take the box labelled 5 , and remove the first two rows. This is followed by the removal of the second column, then the third row, leaving the skew partition illustrated in the figure. As the two remaining removable nodes both have a paired partner (in this case each other) no more rows or columns need be removed, and we have constructed $\hat{\lambda}$.

Proposition 6.9. *The minimal partitions in each block are precisely those for which either $\hat{\lambda} = \emptyset$ or a single row or column, or δ is even and $\hat{\lambda}$ consists of two rows, the second of which has final box of content $-\frac{\delta}{2}$.*

Proof. Clearly if $\hat{\lambda} = \emptyset$ then λ is minimal in its block. In the remaining cases, removal of any part of λ can only involve boxes in $\hat{\lambda}$, and hence to be balanced must involve either a single unpaired box of content $\frac{1-\delta}{2}$ or a single vertical pair in the configuration shown in Figure 5(b). But this is impossible. Hence we assume that $\hat{\lambda}$ is not of the form given in the proposition, and will show that λ is not minimal.

First suppose that δ is odd. If $\hat{\lambda}$ contains two boxes of content $\frac{1-\delta}{2}$ then we can construct a maximal balanced subpartition of λ , mimicking the process in Definitions 6.1 and 6.3 by starting with ϵ . Hence by Theorem 6.5 λ is non minimal. If $\hat{\lambda}$ only contains one box ω with content $\frac{1-\delta}{2}$ then, again by considering Definitions 6.1 and 6.3 and Theorem 6.5, any removable balanced skew-partition must involve ω . The assumption also implies that ϵ is in the first row or column of $\hat{\lambda}$.

Suppose that ϵ is in the first row of $\hat{\lambda}$ and there is more than one row (the case where ϵ is in the first column is similar). If λ is minimal, then no final segment of this row has a removable paired segment in $\hat{\lambda}$; this can only arise if $\hat{\lambda}$ is of the form show in Figure 22 (where shaded areas indicate boxes definitely not in $\hat{\lambda}$), where τ is not paired with any box to the right of ω . But this means that τ has content $\frac{1-\delta}{2}$ which is impossible, and hence λ is not minimal.




FIGURE 22. Possible configuration of $\hat{\lambda}$ when δ is odd

Now suppose that δ is even. If $\hat{\lambda}$ contains either of the configurations shown in Figure 23(a) and (b) then we can again construct a maximal balanced subpartition, and by Theorem 6.5 λ is not minimal.




FIGURE 23. Possible configurations in $\hat{\lambda}$ when δ is even

If $\hat{\lambda}$ contains only one box with content either $-\frac{\delta}{2}$ or $1 - \frac{\delta}{2}$ then this box is either at the end of the first row or bottom of the first column, which contradicts the definition of $\hat{\lambda}$. Thus we must have one of the configurations in Figure 23(c) or (d).

In case (c) ϵ must lie at the end of the first column, and in case (d) at the end of the first row. Arguing as in the δ odd case, we see in case (c) that if λ is minimal then $\hat{\lambda}$ must consist of a single column. However, in case (d), if λ is minimal then we either have a single row or we are in a similar situation to that in Figure 22 and τ must have content $-\frac{\delta}{2}$. But this implies that $\hat{\lambda}$ consists of two rows with the final box having of the second having content $-\frac{\delta}{2}$, which contradicts our assumptions on λ .

Thus the only cases where λ is a minimal partition are those described in the theorem, and so we are done. \square

Example 6.10. To illustrate the last result, consider $\delta = 1$ with $\lambda = (7, 6^2, 5, 2^2)$ as shown in Figure 24. The associated $\hat{\lambda}$ is also shown, and has only one row, and it is easy to see that λ is indeed minimal inside its block.




FIGURE 24. A minimal weight λ and the associated $\hat{\lambda}$

7. ON THE SUBMODULE STRUCTURE OF CERTAIN STANDARD MODULES

In this section we will show that the structure of standard modules can become arbitrarily complicated (as measured by their Loewy length and number of simples in each Loewy layer). For this it will be sufficient to consider certain special partitions which can be more easily analysed.

Lemma 7.1. *If $\epsilon_i \in \text{rem}(\lambda)$ then*

$$[\text{res}_n L_n(\lambda) : L_{n-1}(\lambda - \epsilon_i)] \neq 0.$$

Proof. By (2) and (4) we may assume that $\lambda \vdash n$; the result then follows from Proposition 2.7. □

When considering a multi-skew-partition of differences these skew partitions will be listed in the order from top right to bottom left. We will extend the power notation for partitions to *multipartitions*, so $((2)^2, (21^3))$ will denote the triple of partitions (2) , (2) , and (21^3) .

Example 7.2. To illustrate these definitions we return to the partitions λ and μ considered in Figure 7. In this case we have $\text{add}(\lambda) = \{-5, -3, -1, 3, 6\}$ and $\text{rem}(\lambda) = \{-4, -2, 1, 5\}$. Similarly $\text{add}(\mu) = \{-3, 1, 5\}$ and $\text{rem}(\mu) = \{-1, 4\}$. The pair (λ, μ) is not δ -balanced for any δ , and $\lambda/(\lambda \cap \mu)$ has shape $((1), (2^2), (2, 1))$.

We will be interested in δ -balanced pairs $\mu \subset \lambda$ such that the associated skew partition consists entirely of isolated boxes. If $\mu \subset \lambda$ are balanced with $\lambda/\mu = ((1)^{2m})$, denote the matched pairs of boxes in λ/μ by $\epsilon_1, \epsilon'_1, \dots, \epsilon_m, \epsilon'_m$ with respective contents $a_1, a'_1, \dots, a_m, a'_m$. Let $\mathcal{P}(m)$ denote the power set of $\{1, 2, \dots, m\}$, and for $x \in \mathcal{P}(m)$ set

$$\lambda - x = \lambda - \sum_{i \in x} (\epsilon_i + \epsilon'_i).$$

For example, $\lambda - \{1, \dots, m\} = \mu$.


Theorem 7.3. *Let $\lambda \vdash n$ and $\mu \subset \lambda$ be a balanced pair with $\lambda/\mu = ((1)^{2m})$. Then*

$$\dim \text{Hom}(\Delta_n(\lambda), \Delta_n(\mu)) = 1$$

and

$$[\Delta_n(\mu) : L_n(\lambda - x)] = 1$$

for all $x \in \mathcal{P}(m)$.

FIGURE 25. An example of $\mathcal{L}(\mu, \lambda)$

Further, denote by $\mathcal{L}(\mu, \lambda)$ the induced lattice in the full submodule lattice of $\Delta_n(\mu)$ with vertices those simple modules of the form $L_n(\lambda - x)$ for some $x \in \mathcal{P}(m)$. Then $\mathcal{L}(\mu, \lambda)$ is isomorphic to the superset lattice on $\mathcal{P}(m)$; i.e. every submodule of $\Delta_n(\mu)$ which contains $L_n(\lambda - x)$ contains $L_n(\lambda - y)$ for all $y \subset x$.

In particular the length of the socle series of $\Delta_n(\mu)$ is at least $m + 1$ and there is a socle series layer containing at least m simple modules.

Remark 7.4. (i) Note that for the induced lattice we are only considering factors of the form $L_n(\lambda - x)$. In general the module $\Delta_n(\mu)$ will have many other composition factors. Thus an arrow $A \rightarrow B$ in our induced lattice structure is to be understood as representing some non-trivial extension in $\Delta_n(\mu)$ with A in the head and B in the socle.

(ii) Clearly the final part of the theorem can be strengthened, but is already enough to show that standard modules can have arbitrarily large socle series lengths (and layers of arbitrary width).

Example 7.5. If λ and μ are balanced with $\lambda/\mu = ((1)^6) = \{\epsilon_1, \epsilon'_1, \epsilon_2, \epsilon'_2, \epsilon_3, \epsilon'_3\}$ then the lattice $\mathcal{L}(\mu, \lambda)$ is illustrated in Figure 25.

Proof. We proceed by induction on m , the result being obvious for $m = 0$. By Frobenius reciprocity we have

$$\mathrm{Hom}(\mathrm{ind}_{n-1} \Delta_{n-1}(\lambda - \epsilon_i), \Delta_n(\mu)) \cong \mathrm{Hom}(\Delta_{n-1}(\lambda - \epsilon_i), \mathrm{res}_n \Delta_n(\mu)). \quad (14)$$

By Proposition 2.7 and Corollary 4.8, the only submodule of $\mathrm{res}_n \Delta_n(\mu)$ which can lie in the same block as $\Delta_{n-1}(\lambda - \epsilon_i)$ is isomorphic to $\Delta_{n-1}(\mu + \epsilon'_i)$, and hence by the inductive hypothesis the right-hand side of (14) is one dimensional. Lemma 4.10 now implies that $L_n(\lambda)$ is a composition factor of $\Delta_n(\mu)$. To show that $\dim \mathrm{Hom}(\Delta_n(\lambda), \Delta_n(\mu)) = 1$ it will be enough to show that there is precisely one copy of this composition factor in $\Delta_n(\mu)$ (which will necessarily lie in the socle).

By assumption the pair (λ, μ) is balanced. We will define the *bias* of a pair (λ, τ) with $|\lambda \Delta \tau| = 2t$ to be

$$b(\lambda, \tau) = \left(\sum_{d \in \lambda \Delta \tau} c(d) \right) - t(1 - \delta).$$

Thus a balanced pair has zero bias. Consider the restriction $\text{res}_n \Delta_n(\mu)$. By Proposition 2.7 we have a short exact sequence

$$0 \rightarrow \bigoplus_{\tau \triangleleft \mu} \Delta_{n-1}(\tau) \rightarrow \text{res}_n \Delta_n(\mu) \rightarrow \bigoplus_{\tau \triangleright \mu} \Delta_{n-1}(\tau) \rightarrow 0. \quad (15)$$

Note that μ has no removable boxes with content $\pm a_i$ for $1 \leq i \leq m$, as this would contradict the existence of an addable node with such a content. Thus the only modules $\Delta_{n-1}(\tau)$ in the sequence (15) with bias $\pm a_i$ are $\Delta_{n-1}(\mu + \epsilon_i)$ and $\Delta_{n-1}(\mu + \epsilon'_i)$

By Lemma 7.1 we have that

$$[\text{res}_n L_n(\lambda - x) : L_{n-1}(\lambda - x - \epsilon_i)] = 1$$

provided that $i \notin x$. But (by the observations on bias above) $L_{n-1}(\lambda - x - \epsilon_i)$ can only occur in $\Delta_{n-1}(\mu + \epsilon'_i)$, and by the inductive hypothesis it occurs there precisely once. By varying i we deduce that there is at most one copy of each $L_n(\lambda - x)$ in $\Delta_n(\mu)$. But by induction we know that there is a homomorphism from $\Delta_{n'}(\lambda - x)$ to $\Delta_{n'}(\mu)$ where $n' = |\lambda - x|$, and hence by repeated applications of G that there is a homomorphism from $\Delta_n(\lambda - x)$ to $\Delta_n(\mu)$. Hence we see that $L_n(\lambda - x)$ occurs exactly once in $\Delta_n(\mu)$.

Now consider the summand $\Delta_{n-1}(\mu + \epsilon'_i)$ in $\text{res}_n \Delta_n(\mu)$. This is the only summand of the restriction in which $L_{n-1}(\lambda - x - \epsilon_i)$ (with $i \notin x$) can arise, and this simple appears in an extension below $L_{n-1}(\lambda - y - \epsilon_i)$ for all $y \supset x$ (with $i \notin y$), by the inductive hypothesis. In particular the copy of $L_{n-1}(\lambda - x - \epsilon_i)$ appearing in $\text{res}_n L_n(\lambda - x)$ appears below $L_{n-1}(\lambda - x - \epsilon_i - \epsilon_j - \epsilon'_j)$ in an extension, and this latter simple must come from $\text{res}_n L_n(\lambda - x - \epsilon_j - \epsilon'_j)$. It follows that $L_n(\lambda - x)$ must occur in some extension beneath $L_n(\lambda - x - \epsilon_j - \epsilon'_j)$. This argument works for all j and x , and hence verifies the claimed submodule structure except for the top two layers. However, these are forced by the structure of standard modules. \square

8. THE CASE $\delta = 0$

In this section we will sketch the modifications to the preceding arguments which are required when $\delta = 0$. The most obvious change is that the idempotents e_n considered thus far no longer exist. This is easily remedied — however a more serious complication is the failure of the algebras to be quasihereditary when n is even.




FIGURE 26. The element \bar{e}_n in B_n

For $n \geq 3$ let \bar{e}_n be the element illustrated in Figure 26. This is an idempotent for every value of δ , and satisfies (A1), i.e.

$$\bar{e}_n B_n \bar{e}_n \cong B_{n-2}.$$

Unfortunately we can no longer prove an analogue of (A2) in general, as the algebras are not quasihereditary. If n is odd then there are no problems, and the arguments in the $\delta \neq 0$ case for (A1-6) go through unchanged. The results in Sections 4-7 also generalise, as the various

results needed from [DWH99] include the case $\delta = 0$, and we thus deduce the block result in this case.

For n even, we can no longer appeal directly to the general machinery in [CMPX]. However, the algebras in this case are cellular, and the modules considered by [DWH99] are precisely the cell modules for these algebras. The necessary results coming from the general theory in [CMPX] now have to be verified on an *ad hoc* basis, but this has been carried out in [DWH99]. Thus, again, the results in Sections 4-7 go through unchanged (noting that it is enough to analyse cell modules when determining blocks by [GL96, (3.9.8)] (see [Mat99, 2.22 Corollary])).

REFERENCES

- [Bra37] R. Brauer, *On algebras which are connected with the semisimple continuous groups*, Ann. of Math. **38** (1937), 857–872.
- [Bro55] W. Brown, *An algebra related to the orthogonal group*, Michigan Math. J. **3** (1955), 1–22.
- [CMPX] A. G. Cox, P. P. Martin, A. E. Parker, and C. Xi, *Representation theory of towers of recollement: theory, notes, and examples*, J. Algebra, to appear.
- [DDH] R. Dipper, S. Doty, and J. Hu, *Brauer’s centralizer algebras, symplectic Schur algebras and Schur-Weyl duality*, preprint.
- [Dia] P. Diaconis, *Group representations in probability and statistics*, Institute of Mathematical Statistics Lecture Notes, Monograph Series 11.
- [Don98] S. Donkin, *The q -Schur algebra*, LMS Lecture Notes Series, vol. 253, Cambridge University Press, 1998.
- [Dot98] S. Doty, *Polynomial representations, algebraic monoids, and Schur algebras of classical type*, J. Pure Appl. Algebra **123** (1998), 165–199.
- [DWH99] W. F. Doran, D. B. Wales, and P. J. Hanlon, *On the semisimplicity of the Brauer centralizer algebras*, J. Algebra **211** (1999), 647–685.
- [Ful97] W. Fulton, *Young tableaux*, LMS Student Texts, vol. 35, Cambridge, 1997.
- [GL96] J. J. Graham and G. I. Lehrer, *Cellular algebras*, Invent. Math. **123** (1996), 1–34.
- [Gre80] J. A. Green, *Polynomial representations of GL_n* , Lecture Notes in Mathematics 830, Springer, 1980.
- [HP] R. Hartmann and R. Paget, *Young modules and filtration multiplicities for Brauer algebras*, preprint.
- [HW89a] P. J. Hanlon and D. B. Wales, *Eigenvalues connected with Brauer’s centralizer algebras*, J. Algebra **121** (1989), 446–476.
- [HW89b] ———, *On the decomposition of Brauer’s centralizer algebras*, J. Algebra **121** (1989), 409–445.
- [HW90] ———, *Computing the discriminants of Brauer’s centralizer algebras*, Math. Comp. **54** (1990), 771–796.
- [HW94] ———, *A tower construction for the radical in Brauer’s centralizer algebras*, J. Algebra **164** (1994), 773–830.
- [JK81] G. D. James and A. Kerber, *The representation theory of the Symmetric group*, Encyclopedia of Mathematics and its Applications, vol. 16, Addison-Wesley, 1981.
- [Mar91] P. P. Martin, *Potts models and related problems in statistical mechanics*, World Scientific, Singapore, 1991.
- [Mar96] ———, *The structure of the Partition algebras*, J. Algebra **183** (1996), 319–358.
- [Mat99] A. Mathas, *Iwahori-Hecke algebras and Schur algebras of the symmetric groups*, University lecture series, vol. 15, American Mathematical Society, 1999.
- [MRH04] P. P. Martin and S. Ryom-Hansen, *Virtual algebraic Lie theory: Tilting modules and Ringel duals for blob algebras*, Proc. LMS **89** (2004), 655–675.
- [MW03] P. P. Martin and D. Woodcock, *Generalized blob algebras and alcove geometry*, LMS J. of Comp. and Math. **6** (2003), 249–296.

- [Naz96] M. Nazarov, *Young's orthogonal form for Brauer's centralizer algebra*, J. Algebra **182** (1996), 664–693.
- [Oeh01] S. Oehms, *Centralizer coalgebras, FRT-construction and symplectic monoids*, J. Algebra **244** (2001), 19–44.
- [OR01] R. Orellana and A. Ram, *Affine braids, markov traces and the category \mathcal{O}* , preprint, 2001.
- [Rui05] H. Rui, *A criterion on the semisimple Brauer algebras*, J. Comb. Theory Ser. A **111** (2005), 78–88.
- [Wen88] H. Wenzl, *On the structure of Brauer's centralizer algebra*, Ann. Math. **128** (1988), 173–193.
- [Wey46] H. Weyl, *The classical groups, their invariants and representations*, Princeton University Press, 1946.

E-mail address: A.G.Cox@city.ac.uk, M.Devisscher@city.ac.uk, P.P.Martin@city.ac.uk

CENTRE FOR MATHEMATICAL SCIENCE, CITY UNIVERSITY, NORTHAMPTON SQUARE, LONDON, EC1V 0HB, ENGLAND.