# EQUIVARIANT K-THEORY OF AFFINE FLAG MANIFOLDS AND AFFINE GROTHENDIECK POLYNOMIALS

#### MASAKI KASHIWARA AND MARK SHIMOZONO

ABSTRACT. We study the equivariant K-group of the affine flag manifold with respect to the Borel group action. We prove that the structure sheaf of the (infinite-dimensional) Schubert variety in the K-group is represented by a unique polynomial, which we call the affine Grothendieck polynomial.

#### 1. Introduction

Let G be a simply connected semisimple group, B its Borel subgroup, and X = G/B the flag manifold. Its B-orbits are of the form BwB/B for some w in the Weyl group W. Its closure  $X_w$  is called the Schubert variety. It is well-known that the equivariant K-group  $K_B(X)$ , which is the Grothendieck group of the abelian category of coherent B-equivariant  $\mathscr{O}_X$ -modules, is a free  $K_B(\operatorname{pt})$ -module with  $\{[\mathscr{O}_{X_w}]\}_{w\in W}$  as a basis. Note that  $K_B(\operatorname{pt})$  is isomorphic to the group ring  $\mathbb{Z}[P]$  of the weight lattice P of a maximal torus of B. On the other hand,  $K_B(X) \simeq K_{B\times B}(G)$  gives another structure of  $K_B(\operatorname{pt})$ -module on  $K_B(X)$  and we have a morphism  $\mathbb{Z}[P]\otimes\mathbb{Z}[P]\simeq K_B(\operatorname{pt})\otimes K_B(\operatorname{pt})\to K_B(X)$ , which factors through a homomorphism

(1.1) 
$$\mathbb{Z}[P] \otimes_{\mathbb{Z}[P]^W} \mathbb{Z}[P] \to K_B(X)$$

called the equivariant Borel map. Here  $\mathbb{Z}[P]^W$  is the ring of invariants with respect to the action of the Weyl group W. It is also well-known that  $\mathbb{Z}[P] \otimes_{\mathbb{Z}[P]^W} \mathbb{Z}[P] \to K_B(X)$  is an isomorphism. An element in  $\mathbb{Z}[P] \otimes \mathbb{Z}[P]$  whose image is  $[\mathscr{O}_{X_w}]$ , is known as a double Grothendieck polynomial when G = SL(n) [9].

The purpose of this paper is to generalize these facts to the affine case. Contrary to the finite-dimensional case, there are two kinds of flag manifolds; the inductive limit of Schubert varieties  $\overline{BwB/B}$ , each of which is a finite-dimensional projective variety (see [8] and the references there), and an infinite-dimensional scheme whose Schubert varieties  $\overline{BwB_-/B_-}$  are finite-codimensional subschemes. Here,  $B_-$  is the opposite Borel subgroup. In [7, 8], Kostant-Kumar considered the first flag manifold and studied its equivariant cohomology and K-theory.

In this paper we use the latter flag manifold, which is studied in [3]. We take the affine flag manifold  $X = G/B_{-}$  (see § 2). It is an infinite-dimensional (not quasi-compact) scheme over  $\mathbb{C}$ . Its B-orbits are parameterized by the elements of the Weyl group W. Each B-orbit  $\overset{\circ}{X}_{w}$  is a locally closed subscheme with finite codimension. As a scheme it is isomorphic to  $\mathbb{A}^{\infty} = \operatorname{Spec}(\mathbb{C}[x_{1}, x_{2}, \ldots])$ . The flag

<sup>1991</sup> Mathematics Subject Classification. Primary:19L47; Secondary:14M17, 17B67, 22E65. MS is partially supported by NSF DMS-0401012.

manifold X is a union of B-stable quasi-compact open subsets  $\Omega$ . We define  $K_B(X)$ as the projective limit of  $K_B(\Omega)$ . Then we have  $K_B(X) \cong \prod_{w \in W} K_B(\operatorname{pt})[\mathscr{O}_{X_w}]$ . Similarly to the finite-dimensional case, we have a homomorphism

(1.2) 
$$\mathbb{Z}[P] \otimes_{\mathbb{Z}[P]W} \mathbb{Z}[P] \to K_B(X).$$

In the affine case this morphism is injective but is not surjective; not all  $[\mathscr{O}_{X_w}]$  are in the image of this morphism. However, as we shall see in this paper,  $[\mathscr{O}_{X_w}]$  is in the image after a localization.

Let  $\delta$  be the generator of null roots and let R be the subring of  $\mathbb{Q}(e^{\delta})$  generated by  $e^{\pm \delta}$  and  $(e^{n\delta} - 1)^{-1}$   $(n \neq 0)$ . Tensoring R with (1.2), we have the morphism

$$(1.3) R \underset{\mathbb{Z}[e^{\pm \delta}]}{\otimes} \mathbb{Z}[P] \underset{\mathbb{Z}[P]^W}{\otimes} \mathbb{Z}[P] \to R \underset{\mathbb{Z}[e^{\pm \delta}]}{\otimes} K_B(X).$$

Our main result is the following theorem.

**Theorem 4.4.** For all  $w \in W$ ,  $[\mathscr{O}_{X_w}] \in K_B(X)$ , considered as an element of  $R \otimes_{\mathbb{Z}[e^{\pm\delta}]} K_B(X)$ , is in the image of (1.3).

Note that we have  $\mathbb{Z}[P]^W \simeq \mathbb{Z}[P^W]$  in the affine case.

Roughly speaking, we call the element of  $R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[P] \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P]$  corresponding to  $[\mathscr{O}_{X_w}]$  the affine Grothendieck polynomial (see Proposition 4.5).

In order to prove Theorem 4.4, we use the following vanishing theorem of the first group cohomology.

### Theorem 5.2.

- (i) If |I| > 2 then  $H^1(W, R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[P]) = 0$ . (ii) For any affine Lie algebra  $\mathfrak{g}$ ,  $H^1(W, R \otimes_{\mathbb{Z}[e^{\pm \delta}]} (\bigoplus_{\lambda \in P, |\langle c, \lambda \rangle| < \kappa^*} \mathbb{Z}e^{\lambda})) = 0$ , where  $\kappa^*$  is the dual Coxeter number.

Here I is the index set of simple roots and c is the canonical central element of  $\mathfrak{g}$ .

The plan of this paper is as follows. In § 2 we review the flag manifold of Kac-Moody Lie algebras. In §3 we study the Demazure operators. We also give a simple proof of the fact that the Schubert varieties are normal and Cohen-Macaulay. This proof seems to be new even in the finite-dimensional case. In §4 we study the affine flag manifolds. After the preparation in § 5, we prove Theorem 5.2 in §6. As its application, we give in §7 the proof of Theorem 4.4, the existence of affine Grothendieck polynomials. In § 8 we give the character formula of the global cohomology groups of  $\mathcal{O}_{X_w}(\lambda)$  using the affine Grothendieck polynomials. In § 9 we explain an analogous result for the equivariant cohomology groups of the affine flag manifolds. In § 10 we shall give examples of the affine Grothendieck polynomials.

## 2. Flag manifolds

Let us recall in this section the definition and properties of the flag manifold of a symmetrizable Kac-Moody Lie algebra following [3].

Let  $(a_{ij})_{i,j\in I}$  be a symmetrizable generalized Cartan matrix,  $\mathfrak{g}$  an associated Kac-Moody Lie algebra, and  $\mathfrak{t}$  its Cartan subalgebra. Let  $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{t} \oplus \mathfrak{n}_{-}$  be the triangular decomposition and  $\mathfrak{g} = \bigoplus_{\alpha \in \mathfrak{t}^*} \mathfrak{g}_{\alpha}$  the root decomposition. Let  $\Delta :=$  $\{\alpha \in \mathfrak{t}^* : \mathfrak{g}_{\alpha} \neq 0\} \setminus \{0\}$  be the set of roots and  $\Delta^{\pm}$  the set of positive and negative roots.

Let  $\{\alpha_i\}_{i\in I}$  be the set of simple roots in  $\mathfrak{t}^*$  and  $\{h_i\}_{i\in I}$  the set of simple co-roots in  $\mathfrak{t}$ . Hence we have  $\langle h_i, \alpha_j \rangle = a_{ij}$ . Let us take an integral weight lattice  $P \subset \mathfrak{t}^*$ . We assume the following conditions:

$$(2.1) \begin{cases} \text{(i) } \alpha_i \in P \text{ for all } i \in I, \\ \text{(ii) } \{\alpha_i\}_{i \in I} \text{ is linearly independent,} \\ \text{(iii) } h_i \in P^* \text{ for all } i \in I, \text{ where } P^* \text{ is the dual lattice } \operatorname{Hom}(P, \mathbb{Z}) \subset \mathfrak{t}, \\ \text{(iv) there exists } \Lambda_i \in P \text{ such that } \langle h_j, \Lambda_i \rangle = \delta_{ij}. \end{cases}$$

Let T be the algebraic torus with P as its character lattice. Let W be the Weyl group. It is the subgroup of  $\operatorname{Aut}(\mathfrak{t}^*)$  generated by the simple reflections  $s_i$   $(i \in I)$   $:s_i(\lambda) = \lambda - \langle h_i, \lambda \rangle \alpha_i$ . Let  $U_{\pm}$  be the group scheme with  $\mathfrak{n}_{\pm}$  as its Lie algebra. Let  $B_{\pm} = T \times U_{\pm}$  be the Borel subgroup, whose Lie algebra is  $\mathfrak{b}_{\pm} = \mathfrak{t} \oplus \mathfrak{n}_{\pm}$ . For any  $i \in I$ , let us denote by  $P_i^{\pm}$  the parabolic group whose Lie algebra is  $\mathfrak{b}_{\pm} \oplus \mathfrak{g}_{\mp \alpha_i}$ . Then  $P_i^{\pm}/B_{\pm}$  is isomorphic to the projective line  $\mathbb{P}^1$ .

Let  $P^+ := \{\lambda \in P : \langle h_i, \lambda \rangle \geqslant 0 \ (i \in I) \}$  be the set of dominant integral weights. For  $\lambda \in P^+$ , let  $V(\lambda)$  (resp.  $V(-\lambda)$ ) be the irreducible  $\mathfrak{g}$ -module with highest weight  $\lambda$  (resp. lowest weight  $-\lambda$ ). Then  $A(\mathfrak{g}) := \bigoplus_{\lambda \in P^+} V(\lambda) \otimes V(-\lambda)$  has an algebra

structure and we denote  $\operatorname{Spec}(A(\mathfrak{g}))$  by  $G_{\infty}$ . The scheme  $G_{\infty}$  contains a canonical point e and is endowed with a left action of  $P_i$  and a right action of  $P_i^-$ . The union of  $P_{i_1} \cdots P_{i_m} e P_{j_1}^- \cdots P_{j_m}^- \subset G_{\infty}$  is an open subset of  $G_{\infty}$  and we denote it by G. Then  $P_i$  and  $P_i^-$  act freely on G. The flag manifold X is defined as the quotient  $G/B_-$ . It is a separated (not quasi-compact in general) scheme over  $\mathbb{C}$ . It is covered by affine open subsets isomorphic to  $\mathbb{A}^{\infty} := \operatorname{Spec}(\mathbb{C}[x_1, x_2, \ldots])$  (or  $\mathbb{A}^n$ ), and its structure sheaf  $\mathscr{O}_X$  is coherent. Let  $x_0 \in X$  be the image of  $e \in G$ . Then for  $w \in W$ ,  $wx_0 \in X$  has a sense. The set X has a Bruhat decomposition  $X = \bigsqcup_{w \in W} \overset{\circ}{X}_w$ , where  $\overset{\circ}{X}_w$  is the locally closed subscheme  $Bwx_0$  of X. Let  $X_w$  be the closure of  $\overset{\circ}{X}_w$  endowed with the reduced scheme structure. It is called the  $Schubert \ variety$ . It has codimension  $\ell(w)$ , the length of w, and its structure sheaf  $\mathscr{O}_{X_w}$  is coherent. As a set we have  $X_w = \bigsqcup_{x \geq w} \overset{\circ}{X}_x$ .

- **Remark 2.1.** (i) For any w,  $\overline{B_-wB_-/B_-}$  is a finite-dimensional projective subscheme of X and its union  $\bigcup_{w\in W}\overline{B_-wB_-/B_-}$  is an ind-scheme. This is another flag manifold which we do not use here.
- (ii) For a regular dominant integral weight  $\lambda$ , set  $\widehat{V}(-\lambda) = \prod_{\mu \in P} V(-\lambda)_{\mu}$  where  $V(-\lambda)_{\mu}$  is the weight space of  $V(-\lambda)$  of weight  $\mu$ . Then  $\mathbb{P}(\widehat{V}(-\lambda)) = (\widehat{V}(-\lambda) \setminus \{0\})/\mathbb{C}^{\times}$  has a scheme structure, and  $P_i$  acts on  $\widehat{V}(-\lambda)$  and  $\mathbb{P}(\widehat{V}(-\lambda))$ . Then X is embedded in  $\mathbb{P}(\widehat{V}(-\lambda))$  by  $x_0 \mapsto \overline{u_{-\lambda}}$ , where  $\overline{u_{-\lambda}}$  is the line containing the lowest weight vector  $u_{-\lambda}$ .

Let S be a finite subset of W such that  $x \in S$  as soon as  $x \leqslant y$  for some  $y \in S$ . Then  $\Omega_S := \bigcup_{w \in S} \overset{\circ}{X}_w$  is a B-stable quasi-compact open subset which coincides with  $\bigcup_{w \in S} wBx_0$ . Conversely, any B-stable quasi-compact open subset is of this form. Let  $\operatorname{Coh}_B(\mathscr{O}_{\Omega_S})$  be the abelian category of coherent B-equivariant  $\mathscr{O}_{\Omega_S}$ -modules and let  $K_B(\Omega_S)$  be the Grothendieck group of  $\operatorname{Coh}_B(\mathscr{O}_{\Omega_S})$ . For  $\mathscr{F} \in \operatorname{Coh}_B(\mathscr{O}_{\Omega_S})$ , let us denote by  $[\mathscr{F}]$  the corresponding element of  $K_B(\Omega_S)$ . For  $w \in W$ , the sheaf  $\mathcal{O}_{X_w}$  is a coherent B-equivariant  $\mathcal{O}_X$ -module and gives an element  $[\mathcal{O}_{X_w}]$  of  $K_B(\Omega_S)$ . The equivariant K-group  $K_B(\Omega_S)$  is a module over the ring  $K_B(\operatorname{pt})$ . For  $\lambda \in P$ , we denote by  $e^{\lambda}$  the element of  $K_B(\operatorname{pt})$  represented by the one-dimensional representation of B given by  $B \to T \xrightarrow{e^{\lambda}} \mathbb{C}^{\times}$ . By  $P \ni \lambda \mapsto e^{\lambda}$ ,  $K_B(\operatorname{pt})$  is isomorphic to the group ring  $\mathbb{Z}[P]$ . We also denote by  $e^{\lambda}$  the element of  $\mathbb{Z}[P]$  corresponding to  $\lambda \in P$ .

For  $w \in S$ ,  $wx_0 \in \Omega_S$  is a T-fixed point. It defines a T-equivariant inclusion  $i_w \colon \operatorname{pt} \to \Omega_S$ . Since any coherent  $\mathscr{O}_{\Omega_S}$ -module  $\mathscr{F}$  has locally a finite resolution by locally free modules of finite rank (see Lemma 8.1), the k-th left derived functor  $L_k i_w^* \mathscr{F}$  vanishes for  $k \gg 0$ . Hence we can define the  $K_T(\operatorname{pt})$ -linear homomorphism

$$i_w^* \colon K_B(\Omega_S) \to K_T(\Omega_S) \to K_T(\mathrm{pt}) \simeq K_B(\mathrm{pt})$$

by  $[\mathscr{F}] \mapsto \sum_{k=0}^{\infty} (-1)^k [L_k i_w^* \mathscr{F}]$ . Note that, similarly to the finite-dimensional case, we have

(2.2) 
$$i_x^*([\mathscr{O}_{X_w}]) = \begin{cases} \prod_{\alpha \in \Delta^+ \cap w\Delta^-} (1 - e^{\alpha}) & \text{if } x = w, \\ 0 & \text{unless } x \geqslant w. \end{cases}$$

**Remark 2.2.** The function  $W \to \mathbb{Z}[P]$  given by the equivariant localization  $x \mapsto i_x^*([\mathscr{O}_{X_w}])$ , coincides with the function  $\psi^w$  in [7].

**Lemma 2.3.**  $K_B(\Omega_S)$  is a free  $K_B(\text{pt})$ -module with basis  $\{[\mathscr{O}_{X_w}]\}_{w\in S}$ .

*Proof.* Let us argue by induction on the cardinality of S. Let w be a maximal element of S. Set  $S' = S \setminus \{w\}$ . Then we have  $\Omega_S = \Omega_{S'} \sqcup \overset{\circ}{X}_w$ . Hence we have an exact sequence

$$K_B(\overset{\circ}{X}_w) \to K_B(\Omega_S) \to K_B(\Omega_{S'}) \to 0.$$

By induction  $K_B(\Omega_{S'})$  is a free  $K_B(\operatorname{pt})$ -module with a basis  $\{[\mathscr{O}_{X_x}]\}_{x\in S'}$ . Also  $K_B(\overset{\circ}{X}_w)$  is a free  $K_B(\operatorname{pt})$ -module generated by  $[\mathscr{O}_{X_w}]$ . Hence  $K_B(\Omega_S)$  is generated by  $\{[\mathscr{O}_{X_x}]\}_{x\in S}$ . By (2.2) the image of  $\{[\mathscr{O}_{X_x}]\}_{x\in S}$  under the map

$$K_B(\Omega_S) \xrightarrow{\prod_{x \in S} i_x^*} K_B(\text{pt})^{\prod S}$$

is linearly independent over  $K_B(\operatorname{pt})$ . Here  $K_B(\operatorname{pt})^{\prod S}$  is the product of the copies of  $K_B(\operatorname{pt})$  parameterized by elements of S.

Remark 2.4. For  $\ell \in \mathbb{Z}_{\geqslant 0}$ , let  $\mathfrak{n}_{\ell}$  be the direct sum of  $\mathfrak{g}_{\alpha}$  where  $\alpha = \sum_{i} m_{i} \alpha_{i} \in \Delta^{+}$  ranges over positive roots such that  $\sum_{i} m_{i} > \ell$ . Then  $\mathfrak{n}_{\ell}$  is an ideal of  $\mathfrak{n}$ . Let  $U_{\ell}$  be the normal subgroup of U with  $\mathfrak{n}_{\ell}$  as its Lie algebra. Then for any S as above,  $U_{\ell}$  acts on  $\Omega_{S}$  freely for  $\ell \gg 0$ , and the quotient space  $U_{\ell} \backslash \Omega_{S}$  is a finite-dimensional scheme. Hence  $\Omega_{S}$  is the projective limit of  $\{U_{\ell} \backslash \Omega_{S}\}_{\ell}$  and  $K_{B}(\Omega_{S})$  is the inductive limit of  $\{K_{B/U_{\ell}}(U_{\ell} \backslash \Omega_{S})\}_{\ell}$ .

We set  $K_B(X) = \varprojlim_S K_B(\Omega_S)$ . Hence we have  $K_B(X) = \prod_{w \in W} K_B(\operatorname{pt})[\mathscr{O}_{X_w}]$ .

For  $w \in W$ , the homomorphism  $i_w^* : K_B(\Omega_S) \to K_T(\text{pt})$  induces a homomorphism

$$i_w^* \colon K_B(X) \to K_T(\mathrm{pt}).$$

By (2.2) they induce a monomorphism

(2.3) 
$$K_B(X) \xrightarrow{\prod_{w \in W} i_w^*} K_T(\operatorname{pt})^{\prod W}.$$

For  $i \in I$ , set  $X_i = G/P_i^-$ . Then there is a canonical projection  $p_i \colon X \to X_i$  which is a  $\mathbb{P}^1$ -bundle. We have  $p_i(\overset{\circ}{X}_w) = p_i(\overset{\circ}{X}_{ws_i})$  and  $p_i^{-1}p_i(\overset{\circ}{X}_w) = \overset{\circ}{X}_w \sqcup \overset{\circ}{X}_{ws_i}$  for any  $w \in W$ , and we have the *B*-orbit decomposition  $X_i = \underset{w \in W, ws_i > w}{\sqcup} p_i(\overset{\circ}{X}_w)$ . Similarly to  $K_B(X)$ , we define  $K_B(X_i)$  as  $\varprojlim_S K_B(p_i(\Omega_S))$ . It is isomorphic to

$$\prod_{w \in W, w s_i > w} K_B(\mathrm{pt})[\mathscr{O}_{p_i(X_w)}].$$

There exist homomorphisms  $p_{i*}: K_B(X) \to K_B(X_i)$  and  $p_i^*: K_B(X_i) \to K_B(X)$ , defined by  $[\mathscr{F}] \mapsto \sum_{k=0}^{\infty} (-1)^k [R^k p_{i*} \mathscr{F}] = [p_{i*} \mathscr{F}] - [R^1 p_{i*} \mathscr{F}]$  and  $[\mathscr{E}] \mapsto [p_i^* \mathscr{E}]$ , respectively.

Any element  $\lambda \in P$  induces a group homomorphism  $B_- \to T \xrightarrow{e^{\lambda}} \mathbb{C}^{\times}$ . Let  $\mathscr{O}_X(\lambda)$  be the invertible  $\mathscr{O}_X$ -module on  $X = G/B_-$  induced by this character of  $B_-$ . Then we have a homomorphism of abelian groups  $\mathbb{Z}[P] \to K_B(X)$  given by  $e^{\lambda} \mapsto [\mathscr{O}_X(\lambda)]$ . Note that, for  $\lambda \in P^+$ , we have  $\Gamma(X; \mathscr{O}_X(\lambda)) \simeq V(\lambda)$  and  $H^k(X; \mathscr{O}_X(\lambda)) = 0$  for  $k \neq 0$  (see [4]).

Similarly to the finite-dimensional case ([1, 8]), we have a commutative diagram

$$\mathbb{Z}[P] \longrightarrow K_B(X)$$

$$\downarrow^{p_i^* \circ p_{i_*}}$$

$$\mathbb{Z}[P] \longrightarrow K_B(X).$$

Here  $\mathcal{D}_i$  is given by

(2.4) 
$$\mathscr{D}_i(e^{\lambda}) = \frac{e^{\lambda} - e^{s_i \lambda - \alpha_i}}{1 - e^{-\alpha_i}},$$

and is called the *Demazure operator*.

The  $K_B(\operatorname{pt})$ -module structure on  $K_B(X)$  induces a  $K_B(\operatorname{pt})$ -linear homomorphism  $K_B(\operatorname{pt}) \otimes \mathbb{Z}[P] \to K_B(X)$ . Let  $\mathbb{Z}[P]^W$  be the ring of W-invariants of  $\mathbb{Z}[P]$ . Then  $K_B(\operatorname{pt}) \simeq \mathbb{Z}[P]$  are  $\mathbb{Z}[P]^W$ -algebras. The morphism  $K_B(\operatorname{pt}) \otimes \mathbb{Z}[P] \to K_B(X)$  decomposes as the composition of  $K_B(\operatorname{pt}) \otimes \mathbb{Z}[P] \to K_B(\operatorname{pt}) \otimes_{\mathbb{Z}[P]^W} \mathbb{Z}[P]$  and

$$\beta \colon K_B(\mathrm{pt}) \otimes_{\mathbb{Z}[P]^W} \mathbb{Z}[P] \to K_B(X).$$

It is sometimes called the *equivariant Borel map*.

**Remark 2.5.** By [7], the equivariant Borel map  $K_B(\operatorname{pt}) \otimes_{\mathbb{Z}[P]^W} \mathbb{Z}[P] \to K_B(X)$  is an isomorphism if G is a finite-dimensional simply connected semisimple group.

The composition

(2.5) 
$$K_B(\operatorname{pt}) \otimes_{\mathbb{Z}[P]^W} \mathbb{Z}[P] \xrightarrow{\beta} K_B(X) \xrightarrow{i_w^*} K_B(\operatorname{pt}) \simeq \mathbb{Z}[P]$$

is given by  $a \otimes b \mapsto a \cdot (wb)$ .

#### 3. Demazure operators

Let  $p_i : X \to X_i$  be the  $\mathbb{P}^1$ -bundle as in § 2. In this section we shall show

$$p_i^* p_{i*}([\mathscr{O}_{X_w}]) = \begin{cases} [\mathscr{O}_{X_{ws_i}}] & \text{if } ws_i < w, \\ [\mathscr{O}_{X_w}] & \text{if } ws_i > w. \end{cases}$$

Note that if  $ws_i > w$ , then  $X_w = p_i^{-1}p_i(X_w)$  and  $p_i(X_w) = p_i(X_{ws_i})$ . Moreover,  $\overset{\circ}{X}_{ws_i} \to p_i(\overset{\circ}{X}_{ws_i}) = p_i(\overset{\circ}{X}_w)$  is an isomorphism.

**Lemma 3.1.** We have  $R^1p_{i_*}\mathscr{O}_{X_w}=0$  for all  $w\in W$ .

Proof. Assume that  $ws_i < w$ . Since  $X_{ws_i} = p_i^{-1}p_i(X_w)$ , we have  $\mathscr{O}_{X_{ws_i}} = p_i^*\mathscr{O}_{p_i(X_w)}$ , which implies that  $R^1p_{i_*}\mathscr{O}_{X_{ws_i}} = 0$ . Applying the right exact functor  $R^1p_{i_*}$  to the exact sequence  $\mathscr{O}_{X_{ws_i}} \to \mathscr{O}_{X_w} \to 0$ , we obtain  $R^1p_{i_*}\mathscr{O}_{X_w} = 0$ .

As shown in [3], for any point  $p \in X_w$ , there exist an open neighborhood  $\Omega$  of p and a closed subset S of  $\mathbb{A}^n$  for some n such that we have a commutative diagram

$$X_w \cap \Omega \xrightarrow{\sim} S \times \mathbb{A}^{\infty}$$

$$Q \xrightarrow{\sim} \mathbb{A}^n \times \mathbb{A}^{\infty}.$$

Hence various properties of  $X_w$  (such as normality) make sense.

**Proposition 3.2.** For any  $w \in W$ , we have

- (i)  $X_w$  is normal.
- (ii)  $p_{i*}\mathscr{O}_{X_w} \simeq \mathscr{O}_{p_i(X_w)}$ .

*Proof.* Let us show (i) and (ii) by induction on the length  $\ell(w)$ . Note that when  $ws_i > w$ , (ii) follows from  $X_w = p_i^{-1} p_i(X_w)$ .

When w = e, (i) and (ii) are obvious. Assume that  $\ell(w) > 0$ .

Let us first show (ii). We may assume  $ws_i < w$ . We have a monomorphism  $j \colon \mathscr{O}_{p_i(X_w)} = p_{i*}\mathscr{O}_{X_{ws_i}} \rightarrowtail p_{i*}\mathscr{O}_{X_w}$ , which is an isomorphism on  $p_i(\overset{\circ}{X}_w)$ . By the induction hypothesis,  $X_{ws_i}$  as well as  $p_i(X_w) = p_i(X_{ws_i})$  is normal. Hence j is globally an isomorphism since  $p_{i*}\mathscr{O}_{X_w}$  is a coherent  $\mathscr{O}_{X_i}$ -module.

Next let us show (i). Let  $\widetilde{\mathscr{O}}_{X_w}$  be the normalization of  $\mathscr{O}_{X_w}$ . Then  $\widetilde{\mathscr{O}}_{X_w}$  is a coherent  $\mathscr{O}_X$ -module. Let S be the support of  $\widetilde{\mathscr{O}}_{X_w}/\mathscr{O}_{X_w}$ . Then S is a B-stable closed subset contained in  $X_w \setminus \overset{\circ}{X}_w$ . We shall show that S is an empty set.

Otherwise let  $x \in W$  be a minimal element such that  $X_x \subset S$ . Then x > w. Let us take  $i \in I$  such that  $xs_i < x$ . Assume first  $ws_i > w$ . Then  $\mathscr{O}_{X_w} = p_i^* \mathscr{O}_{p_i(X_w)}$ . Hence  $\widetilde{\mathscr{O}_{X_w}}$  is the inverse image of the normalization of  $\mathscr{O}_{p_i(X_w)}$ . Hence  $S = p_i^{-1} p_i(S)$ . This contradicts  $X_{xs_i} \not\subset S$ . Hence we have  $ws_i < w$ .

We have monomorphisms  $\mathscr{O}_{p_i(X_w)} \rightarrow p_{i_*}\mathscr{O}_{X_w} \rightarrow p_{i_*}\mathscr{O}_{X_w}$ . By the induction hypothesis,  $X_{ws_i}$  as well as  $p_i(X_w) = p_i(X_{ws_i})$  is normal. Since  $p_{i_*}\mathscr{O}_{X_w}$  is a coherent  $\mathscr{O}_{p_i(X_w)}$ -module and  $\mathscr{O}_{p_i(X_w)} \rightarrow p_{i_*}\mathscr{O}_{X_w} \rightarrow p_{i_*}\mathscr{O}_{X_w}$  are isomorphisms on  $p_i(\mathring{X}_w)$ , the

normality of  $p_i(X_w)$  implies that  $\mathscr{O}_{p_i(X_w)} \to p_{i*} \widetilde{\mathscr{O}_{X_w}}$  is an isomorphism on  $X_i$ . Hence we have isomorphisms  $\mathscr{O}_{p_i(X_w)} \xrightarrow{\sim} p_{i*} \mathscr{O}_{X_w} \xrightarrow{\sim} p_{i*} \widetilde{\mathscr{O}_{X_w}}$ .

We have an exact sequence  $p_{i*}\mathscr{O}_{X_w} \to p_{i*}\widetilde{\mathscr{O}_{X_w}} \to p_{i*}(\widetilde{\mathscr{O}_{X_w}}/\mathscr{O}_{X_w}) \to R^1 p_{i*}\mathscr{O}_{X_w}$ . Since  $R^1 p_{i*}\mathscr{O}_{X_w} = 0$  by Lemma 3.1, we obtain  $p_{i*}(\widetilde{\mathscr{O}_{X_w}}/\mathscr{O}_{X_w}) = 0$ . On the other hand, since  $S \to X_i$  is an isomorphism on  $p_i(\mathring{X}_x)$ , the support of  $p_{i*}(\widetilde{\mathscr{O}_{X_w}}/\mathscr{O}_{X_w})$  contains  $p_i(\mathring{X}_x)$ , which is a contradiction.

Corollary 3.3. For any  $w \in W$ , we have

$$p_i^* p_{i*}([\mathscr{O}_{X_w}]) = \begin{cases} [\mathscr{O}_{X_{ws_i}}] & \text{if } ws_i < w, \\ [\mathscr{O}_{X_w}] & \text{if } ws_i > w. \end{cases}$$

**Proposition 3.4.** The module  $\mathcal{O}_{X_w}$  is Cohen-Macaulay for any  $w \in W$ .

*Proof.* Since  $X_w$  has codimension  $\ell(w)$ ,  $\mathscr{E}xt_{\mathscr{O}_X}^k(\mathscr{O}_{X_w},\mathscr{O}_X)=0$  for  $k<\ell(w)$ . Hence it is enough to show that  $\mathscr{E}xt_{\mathscr{O}_X}^k(\mathscr{O}_{X_w},\mathscr{O}_X)=0$  for  $k>\ell(w)$ . We shall prove it by induction on  $\ell(w)$ . When w=e, it is obvious. Assume that  $\ell(w)>0$ . Set

(3.1) 
$$\mathscr{F} = \tau^{>\ell(w)} R \mathscr{H}om_{\mathscr{O}_X}(\mathscr{O}_{X_w}, \mathscr{O}_X),$$

where  $\tau^{>\ell(w)}$  is the truncation functor (see e.g. [6]). Let us set  $S = \text{Supp}(\mathscr{F})$ . Then S is a B-stable closed subset of  $X_w$ . Let  $x \in W$  be a minimal element of  $\{x \in W : X_x \subset S\}$ . Let us take  $i \in I$  such that  $xs_i < x$ . If  $ws_i > w$ , then we have

$$\mathscr{F} \simeq p_i^* \tau^{>\ell(w)} R \mathscr{H} om_{\mathscr{O}_{X_i}}(\mathscr{O}_{p_i(X_w)}, \mathscr{O}_{X_i}),$$

which implies that  $p_i^{-1}p_i(S) = S$ . Hence  $X_{xs_i} \subset S$ , which is a contradiction. Hence we have  $ws_i < w$ .

Let  $\Omega_{X/X_i}$  be the relative canonical sheaf, which is isomorphic to  $\mathcal{O}_X(-\alpha_i)$ . Then the Grothendieck-Serre duality theorem says that

$$(3.2) Rp_{i_*}R\mathcal{H}om_{\mathscr{O}_X}(\mathscr{O}_{X_w}, \Omega_{X/X_i}[1]) \simeq R\mathcal{H}om_{\mathscr{O}_{X_i}}(Rp_{i_*}\mathscr{O}_{X_w}, \mathscr{O}_{X_i}).$$

Since  $Rp_{i_*}\mathscr{O}_{X_w} \simeq \mathscr{O}_{p_i(X_w)}$  by Lemma 3.1 and Proposition 3.2 and since the induction hypothesis implies that  $p_i(X_w) = p_i(X_{ws_i})$  is Cohen-Macaulay, we have

$$R\mathscr{H}om_{\mathscr{O}_{X_i}}(Rp_{i*}\mathscr{O}_{X_w},\mathscr{O}_{X_i}) \simeq \mathscr{E}xt_{\mathscr{O}_{X_i}}^{\ell(w)-1}(\mathscr{O}_{p_i(X_w)},\mathscr{O}_{X_i})[1-\ell(w)].$$

Hence (3.2) implies that

$$\mathrm{R}p_{i_*}\mathrm{R}\mathscr{H}om_{\mathscr{O}_X}(\mathscr{O}_{X_w},\Omega_{X/X_i})\simeq \mathscr{E}xt_{\mathscr{O}_{X_i}}^{\ell(w)-1}(\mathscr{O}_{p_i(X_w)},\mathscr{O}_{X_i})[-\ell(w)].$$

Applying  $Rp_{i_*}$  to the distinguished triangle

$$\mathscr{E}xt^{\ell(w)}_{\mathscr{O}_X}(\mathscr{O}_{X_w},\Omega_{X/X_i})[-\ell(w)] \to R\mathscr{H}om_{\mathscr{O}_X}(\mathscr{O}_{X_w},\Omega_{X/X_i}) \to \mathscr{F} \otimes \Omega_{X/X_i} \stackrel{+1}{\longrightarrow},$$
 we obtain a distinguished triangle

$$Rp_{i*}\mathscr{E}xt_{\mathscr{O}_{X}}^{\ell(w)}(\mathscr{O}_{X_{w}},\Omega_{X/X_{i}})[-\ell(w)] \to \mathscr{E}xt_{\mathscr{O}_{X_{i}}}^{\ell(w)-1}(\mathscr{O}_{p_{i}(X_{w})},\mathscr{O}_{X_{i}})[-\ell(w)] \\ \to Rp_{i*}(\mathscr{F}\otimes\Omega_{X/X_{i}}) \xrightarrow{+1}.$$

Hence, taking cohomology groups, we obtain

$$R^k p_{i*}(\mathscr{F} \otimes \Omega_{X/X_i}) \xrightarrow{\sim} R^{k-\ell(w)+1} p_{i*} \mathscr{E}xt_{\mathscr{O}_{Y}}^{\ell(w)}(\mathscr{O}_{X_w}, \Omega_{X/X_i}) \simeq 0$$

for  $k > \ell(w)$ . On a neighborhood of  $p_i(\overset{\circ}{X}_x)$ ,  $S \to X_i$  is an embedding, and we have  $R^k p_{i*}(\mathscr{F} \otimes \Omega_{X/X_i}) \simeq p_{i*} H^k(\mathscr{F} \otimes \Omega_{X/X_i})$ . Hence  $H^k(\mathscr{F} \otimes \Omega_{X/X_i}) = 0$  for  $k > \ell(w)$  on a neighborhood of  $\overset{\circ}{X}_x$ , which is a contradiction.

### 4. Affine flag manifolds

In this section we shall consider affine flag manifolds. Let  $(a_{ij})_{ij\in I}$  be an affine Cartan matrix. Let  $Q = \bigoplus_{i\in I} \mathbb{Z}\alpha_i$  and  $L = \bigoplus_{i\in I} \mathbb{Z}\Lambda_i$ . We take  $L \oplus Q$  as the integral weight lattice P. In the Cartan subalgebra  $\mathfrak{t} = \operatorname{Hom}(P, \mathbb{C})$ , we give the simple coroots  $h_i \in \mathfrak{t}$  by

$$\langle h_i, \alpha_i \rangle = a_{ii}$$
, and  $\langle h_i, \Lambda_i \rangle = \delta_{ii}$ .

- **Remark 4.1.** (i) We have taken  $L \oplus Q$  as the integral weight lattice P. Let B denote the associated Borel subgroup. Let P' be another integral weight lattice satisfying (2.1) and let B' be its associated Borel subgroup. Then there is a map  $P \to P'$  and therefore a morphism  $B' \to B$ . Hence we have morphisms  $K_B(\operatorname{pt}) \to K_{B'}(\operatorname{pt})$  and  $K_{B'}(\operatorname{pt}) \otimes_{K_B(\operatorname{pt})} K_B(X) \xrightarrow{\sim} K_{B'}(X)$ . In this sense, our choice of P is universal.
  - (ii) However our choice of P can often be realized as a direct sum  $P' \oplus S$ , where  $S \subset P^W$  and  $Q \subset P'$ . Then we have  $K_B(X) \simeq \mathbb{Z}[S] \otimes_{\mathbb{Z}} K_{B'}(X)$ .

The Weyl group W acts on P and Q, and we have an exact sequence of Wmodules

$$0 \to Q \to P \to L \to 0$$
,

where L is endowed with the trivial action of W.

Let us set  $Q_+ = \sum_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i$  and let  $\delta \in Q_+$  be the imaginary root such that

$$\{\alpha \in Q ; \langle h_i, \alpha \rangle = 0 \text{ for all } i \} = \mathbb{Z}\delta.$$

Similarly, let us choose  $c \in \sum_{i \in I} \mathbb{Z}_{\geq 0} h_i$  such that

$$\left\{h \in \sum_{i \in I} \mathbb{Z} h_i; \langle h, \alpha_i \rangle = 0 \text{ for all } i\right\} = \mathbb{Z} c.$$

We write

$$\delta = \sum_{i \in I} a_i \alpha_i \text{ and } c = \sum_{i \in I} a_i^{\vee} h_i.$$

Then there exists a unique symmetric bilinear form ( , ):  $P \times P \to \mathbb{Q}$  such that

$$(\lambda, \lambda') = 0 \text{ for any } \lambda, \lambda' \in L,$$

$$\langle h_i, \lambda \rangle = \frac{2(\alpha_i, \lambda)}{(\alpha_i, \alpha_i)} \text{ for any } \lambda \in P,$$

$$\langle c, \lambda \rangle = (\delta, \lambda) \text{ for any } \lambda \in P.$$

Set  $Q_{\rm cl} = Q/\mathbb{Z}\delta$ . Then W acts on  $Q_{\rm cl}$ . Let us choose  $0 \in I$  such that the image  $W_{\rm cl}$  of W in  ${\rm Aut}(Q_{\rm cl})$  is generated by the image of  $\{s_i\}_{i\in I\setminus\{0\}}$  and  $a_0=1$ . These conditions are equivalent to saying that  $a_0=1$  and  $\delta-\alpha_0$  is a constant multiple of a root. Such a 0 exists uniquely up to a Dynkin diagram automorphism. Note that

$$(\alpha_0, \alpha_0) = \begin{cases} 2 & \text{if } \mathfrak{g} \not\simeq A_{2n}^{(2)}, \\ 4 & \text{if } \mathfrak{g} \simeq A_{2n}^{(2)}, \end{cases} \quad \text{and} \quad \theta := \delta - \alpha_0 \in \frac{(\alpha_0, \alpha_0)}{2} \Delta^+.$$

Let  $P^W$  denote the space of W-invariant integral weights. We have in the affine case

$$\mathbb{Z}[P^W] \xrightarrow{\sim} \mathbb{Z}[P]^W$$
,

since, for any  $\lambda \in P$ , either  $\lambda \in P^W$  or  $W\lambda$  is an infinite set (see Remark 11.2).

We see easily that  $P^W = \{\lambda \in P ; \langle h_i, \lambda \rangle = 0 \text{ for all } i \}$  is isomorphic to Q by the map  $\eta: Q \xrightarrow{\sim} P^W$  given by

(4.2) 
$$\eta(\beta) = \beta - \sum_{i \in I} \langle h_i, \beta \rangle \Lambda_i.$$

Hence we have  $P = L \bigoplus P^W$  and  $\mathbb{Z}[P] \cong \mathbb{Z}[L] \otimes_{\mathbb{Z}} \mathbb{Z}[P^W]$ , which implies

$$K_B(\mathrm{pt}) \otimes_{\mathbb{Z}[P]^W} \mathbb{Z}[P] \cong \mathbb{Z}[P] \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \cong \mathbb{Z}[P] \otimes_{\mathbb{Z}} \mathbb{Z}[L] \cong \mathbb{Z}[L] \otimes_{\mathbb{Z}} \mathbb{Z}[P].$$

For any  $w \in W$ , let  $i_w^* \circ \beta \colon \mathbb{Z}[P] \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \to \mathbb{Z}[P]$  be as in (2.5).

**Lemma 4.2.** The homomorphism induced by the  $i_w^* \circ \beta$ 's

$$\mathbb{Z}[P] \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \to \mathbb{Z}[P]^{\prod W}$$

is injective. Here  $\mathbb{Z}[P]^{\prod W}$  is the product of the copies of  $\mathbb{Z}[P]$  parameterized by the elements of W.

We shall give the proof of this lemma in  $\S 11$ .

As a corollary (together with (2.3)), we have

Corollary 4.3. The homomorphism

$$K_B(\operatorname{pt}) \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \xrightarrow{\beta} K_B(X)$$

is injective.

Let R be the subring of  $\mathbb{Q}(e^{\delta})$  generated by  $e^{\pm \delta}$  and  $(e^{n\delta} - 1)^{-1}$  (n > 0). Then we have an injective homomorphism

$$(4.3) \beta \colon R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_B(\mathrm{pt}) \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \rightarrowtail R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_B(X).$$

Our main result is the following theorem.

**Theorem 4.4.** For all  $w \in W$ ,  $[\mathscr{O}_{X_w}] \in K_B(X)$ , considered as an element of  $R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_B(X)$ , is in the image of the map  $\beta$  in (4.3).

Hence,  $\bigoplus_{w \in W} K_B(\operatorname{pt})[\mathscr{O}_{X_w}]$  may be regarded as a submodule of  $R \underset{\mathbb{Z}[e^{\pm \delta}]}{\otimes} K_B(\operatorname{pt}) \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P]$ .

We shall prove a slightly more precise result.

Let  $\xi \colon \mathbb{Z}[P] \to K_B(\operatorname{pt}) \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \simeq \mathbb{Z}[L] \otimes_{\mathbb{Z}} \mathbb{Z}[P]$  be the homomorphism given by

(4.4) 
$$\xi(e^{\lambda+\beta}) = e^{-\lambda} \otimes e^{\lambda+\beta} \text{ for } \lambda \in L \text{ and } \beta \in Q.$$

We extend this to

$$\xi \colon R \otimes_{\mathbb{Z}[\mathbf{e}^{\pm \delta}]} \mathbb{Z}[P] \rightarrowtail R \otimes_{\mathbb{Z}[\mathbf{e}^{\pm \delta}]} K_B(\mathrm{pt}) \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P].$$

Set  $\rho = \sum_{i \in I} \Lambda_i$ . Then  $\kappa^* := \langle c, \rho \rangle = \sum_{i \in I} a_i^{\vee}$  is called the *dual Coxeter number* of  $\mathfrak{g}$ . Let us introduce the W-submodule of  $\mathbb{Z}[P]$ :

(4.5) 
$$\mathbb{Z}[P]_{[0,-\kappa^*)} := \bigoplus_{\lambda \in P, \, 0 \geqslant \langle c, \lambda \rangle > -\kappa^*} \mathbb{Z}e^{\lambda}.$$

**Proposition 4.5.** For any  $w \in W$ , there exists a unique

$$\mathcal{G}_w \in R \otimes_{\mathbb{Z}[\mathrm{e}^{\pm \delta}]} \mathbb{Z}[P]_{[0,-\kappa^*)}$$

such that  $\beta \circ \xi(\mathcal{G}_w) = [\mathscr{O}_{X_w}].$ 

We call  $\mathcal{G}_w$  the affine Grothendieck polynomial. The proof of this proposition is given in § 7 as an application of Theorem 5.2 below.

**Remark 4.6.** In the finite-dimensional case,  $Z[P]^W$  is much bigger than  $\mathbb{Z}[P^W]$ , and the choice of Grothendieck polynomials is not unique. However, in the affine case  $\mathcal{G}_w$  is uniquely determined.

**Remark 4.7.** Let  $\psi \colon \mathbb{Z}[P] \to \mathbb{Z}[P]$  be the homomorphism given by

$$\psi(e^{\lambda+\alpha}) = e^{\lambda-\eta(\alpha)} \text{ for } \lambda \in L \text{ and } \alpha \in Q.$$

Then we have

(4.6) 
$$\mathcal{G}_{w^{-1}} = \psi(\mathcal{G}_w) \text{ for any } w \in W.$$

Indeed, let  $\varphi \colon \mathfrak{g} \to \mathfrak{g}$  be the Lie algebra homomorphism such that  $\varphi(e_i) = f_i$ ,  $\varphi(f_i) = e_i$  and  $\varphi(h) = -h$  for  $h \in \mathfrak{t}$ . It induces group scheme morphisms  $\varphi \colon B \to B_-$  and  $\varphi \colon B_- \to B$ . Note that  $\varphi$  induces an isomorphism

$$\mathbb{Z}[P] \simeq K_T(\mathrm{pt}) \simeq K_B(\mathrm{pt}) \xrightarrow{\varphi^*} K_{B_-}(\mathrm{pt}) \simeq K_T(\mathrm{pt}) \simeq \mathbb{Z}[P],$$

which is given by  $e^{\lambda} \mapsto e^{-\lambda}$ . There exists a unique scheme isomorphism  $a: G \to G$  such that a(e) = e and  $a(bgb_{-}^{-1}) = \varphi(b_{-})a(g)\varphi(b)^{-1}$  for  $b \in B$ ,  $b_{-} \in B_{-}$  and  $g \in G$ . We have  $K_{B}(X) \simeq K_{B \times B_{-}}(G)$ , and  $a: G \to G$  induces a commutative diagram

$$\mathbb{Z}[P] \xrightarrow{\xi} K_B(\mathrm{pt}) \otimes K_{B_-}(\mathrm{pt}) \xrightarrow{\beta} K_{B \times B_-}(G) \xrightarrow{\sim} K_B(X)$$

$$\downarrow^{\psi} \qquad \qquad \downarrow^{\bar{\psi}} \qquad \qquad \downarrow^{\bar{\psi}}$$

$$\mathbb{Z}[P] \xrightarrow{\xi} K_B(\mathrm{pt}) \otimes K_{B_-}(\mathrm{pt}) \xrightarrow{\beta} K_{B \times B_-}(G) \xrightarrow{\sim} K_B(X).$$

Here  $k(e^{\lambda} \otimes e^{\mu}) = e^{-\mu} \otimes e^{-\lambda}$ . Indeed,  $k\xi(e^{\lambda+\alpha}) = k(e^{-\lambda} \otimes e^{\lambda+\alpha}) = e^{-\lambda-\alpha} \otimes e^{\lambda} = e^{-\lambda-\alpha+\eta(\alpha)} \otimes e^{\lambda-\eta(\alpha)} = \xi(e^{\lambda-\eta(\alpha)})$ .

Since  $a(BwB_{-}) = Bw^{-1}B_{-}$ , we have  $\tilde{\psi}([\mathscr{O}_{X_w}]) = [\mathscr{O}_{X_{w-1}}]$ . which implies (4.6).

## 5. Vanishing of the Weyl group cohomology

Let (W, S) be a Coxeter group, where S is a system of generators. For a subset S' of S, let us denote by  $W_{S'}$  the subgroup of W generated by S'.

**Lemma 5.1.** For any W-module V, we have

$$H^1(W,V) = \frac{\{\{v_s\}_{s \in S} : v_s \in V \text{ satisfy conditions (i) and (ii) below}\}}{\{\{v_s\}_{s \in S} : \text{there exists } v \in V \text{ such that } v_s = (1-s)v\}},$$

- (i)  $(1+s)v_s = 0$ ,
- (ii) for any pair of distinct elements  $s, t \in S$  such that  $W_{\{s,t\}}$  is a finite group,

$$\sum_{w \in W_{\{s,t\}}, ws > w} (-1)^{\ell(w)} w v_s = \sum_{w \in W_{\{s,t\}}, wt > w} (-1)^{\ell(w)} w v_t.$$

Here  $\ell \colon W \to \mathbb{Z}_{\geqslant 0}$  is the length function.

*Proof.* Let  $\mathbb{Z}[W] \to \mathbb{Z}$  be the augmentation homomorphism  $W \ni w \mapsto 1$ . Then its kernel is the image of the homomorphism  $\varphi \colon \bigoplus_{s \in S} \mathbb{Z}[W]e_s \to \mathbb{Z}[W]$  given by  $\varphi(e_s) = 1 - s$ . By definition,  $H^1(W, V) = \operatorname{Ext}^1_{\mathbb{Z}[W]}(\mathbb{Z}, V)$  is the cohomology of

$$\operatorname{Hom}_{\mathbb{Z}[W]}(\mathbb{Z}[W], V) \to \operatorname{Hom}_{\mathbb{Z}[W]}(\bigoplus_{s \in S} \mathbb{Z}[W]e_s, V) \to \operatorname{Hom}_{\mathbb{Z}[W]}(\operatorname{Ker}(\varphi), V).$$

Hence it is enough to show that  $Ker(\varphi)$  coincides with the  $\mathbb{Z}[W]$ -submodule N generated by  $(1+s)e_s$   $(s \in W)$  and  $\sum_{x \in W_{\{s,t\}}, xs > x} (-1)^{\ell(x)} x e_s - \sum_{x \in W_{\{s,t\}}, xt > x} (-1)^{\ell(x)} x e_t$ where  $(s,t) \in S \times S$  ranges over the pairs as in (ii). It is easy to see that those elements are in  $Ker(\varphi)$ . Indeed, the last elements belong to  $Ker(\varphi)$  because  $\sum_{x \in W_{\{s,t\}}, xs > x} (-1)^{\ell(x)} x (1-s) = \sum_{x \in W_{\{s,t\}}} (-1)^{\ell(x)} x.$  Let  $z = \sum_{w \in W, s \in S} a_{w,s} w e_s$  be an element of  $\operatorname{Ker}(\varphi)$  where  $a_{w,s} \in \mathbb{Z}$ .

Since  $\sum_{w \in W, s \in S} a_{w,s} w(1-s) = 0$ , we have

(5.1) 
$$\sum_{s \in S} a_{w,s} = \sum_{s \in S} a_{ws,s} \quad \text{for all } w \in W.$$

We shall show  $z \in N$  by induction on  $\ell$ , the largest number among the  $\ell(w)$ 's such that  $a_{w,s} \neq 0$  for some  $s \in S$ . Then we shall show  $z \in N$  by induction on the cardinality of  $\{(w,s) \in W \times S : a_{w,s} \neq 0 \text{ and } \ell = \ell(w) \}$ . Let us take  $(w_1, s_1)$ such that  $a_{w_1,s_1} \neq 0$  and  $\ell = \ell(w_1)$ . Subtracting  $a_{w_1,s_1}w_1(1+s_1)e_{s_1} \in N$  from z when  $w_1s_1 < w_1$ , we may assume from the beginning that  $w_2 := w_1s_1 > w_1$ . Then  $\ell(w_2) = \ell + 1$ . Applying (5.1) for  $w = w_2$ , we have  $0 = \sum_{s \in S} a_{w_2,s} = \sum_{s \in S} a_{w_2s,s} = a_{w_1,s_1} + \sum_{s \in S, s \neq s_1} a_{w_2s,s}$ . Hence there exists  $s_2 \neq s_1$  such that  $a_{w_2s_2,s_2} \neq 0$ . Hence we have  $\ell(w_2s_2) \leqslant \ell < \ell(w_2) = \ell + 1$ . Hence  $w_2$  is the longest element in  $w_2W_{s_1,s_2}$ . Therefore  $W_{s_1,s_2}$  is a finite group. Let  $w_3$  be the shortest element of  $w_1W_{s_1,s_2}$ . Subtracting  $\pm a_{w_1,s_1} \Big( \sum_{x \in W_{\{s_1,s_2\}}, \, xs_1 > x} (-1)^{\ell(x)} w_3 x e_{s_1} - \sum_{x \in W_{\{s_1,s_2\}}, \, xs_2 > x} (-1)^{\ell(x)} w_3 x e_{s_2} \Big)$  in N from z, we can erase the term  $w_1 e_{s_1}$  in z, and the induction proceeds.

Now let us return to the affine case where W is the Weyl group. Recall that Ris the ring generated by  $e^{\pm \delta}$  and  $(e^{n\delta} - 1)^{-1}$   $(n \neq 0)$ .

Note that  $\mathbb{Z}[P]$  is a direct sum of W-submodules of the form  $\bigoplus_{\lambda \in W\lambda_0} \mathbb{Z}e^{\lambda}$  ( $\lambda_0 \in P$ ).

**Theorem 5.2.** (i) If 
$$|I| > 2$$
, then  $H^1(W, R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[P]) = 0$ .

(ii) For any affine Lie algebra  $\mathfrak{g}$ ,  $H^1(W, R \otimes_{\mathbb{Z}[e^{\pm \delta}]} (\bigoplus_{|\langle c, \lambda \rangle| < \langle c, \rho \rangle} \mathbb{Z}e^{\lambda})) = 0$  where  $\rho =$  $\sum_{i \in I} \Lambda_i$ .

In fact, we shall prove more precise results. For  $J \subset I$ , let  $W_J$  be the subgroup of W generated by  $\{s_i : i \in J\}$ .

**Proposition 5.3.** (i) If  $J \neq I$ , then  $H^1(W_J, \mathbb{Z}[P]) = 0$ ,

(ii) if  $\langle c, \lambda_0 \rangle \neq 0$  and if  $\lambda_0$  satisfies one of the conditions below (a)  $\lambda_0$  is not regular (i.e.,  $(\beta, \lambda_0) = 0$  for some  $\beta \in \Delta$ ), (b) |I| > 2, then  $H^1(W, \bigoplus_{\lambda \in W \lambda_0} \mathbb{Z} e^{\lambda}) = 0$ ,

(iii) if  $\langle c, \lambda_0 \rangle = 0$ , and  $\langle h_i, \lambda_0 \rangle \geqslant 0$  for all  $i \in I \setminus \{0\}$ , then  $1 - e^{\langle h_0, \lambda_0 \rangle \delta} \colon H^1(W, \bigoplus_{\lambda \in W \lambda_0} \mathbb{Z} e^{\lambda}) \to H^1(W, \bigoplus_{\lambda \in W \lambda_0} \mathbb{Z} e^{\lambda})$ 

is the zero map.

Note that in (iii), we have  $W\lambda_0 + \mathbb{Z}\langle h_0, \lambda_0 \rangle \delta = W\lambda_0$ . Also note that, if  $0 < \infty$  $|\langle c, \lambda_0 \rangle| < \langle c, \rho \rangle$ , then  $\lambda_0$  is not regular. Together with  $H^1(W, \mathbb{Z}) = 0$ , it is easy to see that Proposition 5.3 implies Theorem 5.2. The proof of Proposition 5.3 will be given in the next section.

**Remark 5.4.** If |I| = 2 and  $\lambda_0$  is regular dominant, then  $H^1(W, \bigoplus_{\lambda \in W \lambda_0} \mathbb{Z} e^{\lambda}) \simeq \mathbb{Z}$ .

#### 6. Proof of Proposition 5.3

In this section, we shall prove Proposition 5.3.

6.1. Proof of Proposition 5.3 (i), (ii). In the case (ii), we may assume that  $\langle c, \lambda_0 \rangle > 0$ . In such a case,  $W\lambda_0$  contains a dominant weight. Hence, in order to prove (i) and (ii), it is enough to show that

(6.1) 
$$H^{1}(W_{J}, \bigoplus_{\lambda \in W_{J}\lambda_{0}} \mathbb{Z}e^{\lambda}) = 0$$

under the condition

if  $J \subset I$  and  $\langle h_i, \lambda_0 \rangle \geqslant 0$  for any  $i \in J$ . Moreover, when J = I(6.2)and |I|=2, we assume further that  $\lambda_0$  is not regular.

We shall show this by induction on the cardinality of J. If  $|J| \leq 1$ , then it is obvious. Assuming that |J| > 1, let us take  $i_0 \in J$ , and set  $J_0 = J \setminus \{i_0\}$ . Then (6.1) is true for  $J_0$  by the induction hypothesis.

Assuming that  $(v_i)_{i\in J}$  with  $v_i \in \bigoplus_{\lambda \in W_J \lambda_0} \mathbb{Z}e^{\lambda}$  satisfies (i) and (ii) in Lemma 5.1, let us show the existence of  $v \in \bigoplus_{\lambda \in W_J \lambda_0} \mathbb{Z}e^{\lambda}$  such that  $v_i = (1 - s_i)v$ . By the induction hypothesis, there exists  $v' \in \bigoplus_{\lambda \in W_J \lambda_0} \mathbb{Z}e^{\lambda}$  such that  $v_i = (1 - s_i)v'$  for all  $i \in J_0$ .

Hence replacing  $v_i$  with  $v_i - (1 - s_i)v'$ , we may assume from the beginning that  $v_i = 0$  for all  $i \in J_0$ . On the other hand, since  $(1 + s_{i_0})v_{i_0} = 0$ , there exists  $u_0$  such that  $v_{i_0} = (1 - s_{i_0})u_0$ . By (ii) in Lemma 5.1, we have

(6.3) 
$$\sum_{w \in W_{\{i_0,j\}}} (-1)^{\ell(w)} w u_0 = 0 \text{ if } j \in J_0 \text{ and } W_{\{i_0,j\}} \text{ is a finite group.}$$

It is enough to show the following:

(6.4) there exists a decomposition 
$$u_0 = z_0 + z_1$$
 where  $z_0, z_1 \in \bigoplus_{\lambda \in W_J \lambda_0} \mathbb{Z} e^{\lambda}$  and  $s_{i_0} z_0 = z_0$  and  $z_1$  is  $W_{J_0}$ -invariant.

Indeed, we then have  $v_{i_0} = (1 - s_{i_0})u_0 = (1 - s_{i_0})z_1$  and  $v_i = 0 = (1 - s_i)z_1$  for  $i \in J_0$ .

We shall show (6.4) under the conditions (6.2) and (6.3). Let  $d: W_J \lambda_0 \to \mathbb{Z}_{\geq 0}$  be the function given by  $d(\lambda) = \sum_{i \in J} m_i$  writing  $\lambda_0 - \lambda = \sum_{i \in J} m_i \alpha_i$   $(m_i \in \mathbb{Z}_{\geq 0})$ . Let us write  $u_0 = \sum_{\lambda \in W_J \lambda_0} a_{\lambda} e^{\lambda}$ . Set  $\operatorname{supp}(u_0) := \{\lambda \in W_J \lambda_0 ; a_{\lambda} \neq 0\}$ , and we argue by induction on  $d(u_0) := \max \{d(\lambda); \lambda \in \text{supp}(u_0)\}$ . Then we argue by induction on the cardinality of supp<sup>max</sup> $(u_0) := \{\lambda \in \text{supp}(u_0) ; d(\lambda) = d(u_0) \}.$ 

Let us take  $\lambda_1 \in \operatorname{supp}^{\max}(u_0)$ . If  $\langle h_{i_0}, \lambda_1 \rangle < 0$ , then  $d(s_{i_0}\lambda_1) < d(\lambda_1)$  and, subtracting the  $s_{i_0}$ -invariant  $a_{\lambda_1}(e^{\lambda_1}+e^{s_{i_0}\lambda_1})$  from  $u_0$ , we can delete the term  $e^{\lambda_1}$  in  $u_0$ . If  $\langle h_{i_0}, \lambda_1 \rangle = 0$ , then subtracting the  $s_{i_0}$ -invariant  $a_{\lambda_1} e^{\lambda_1}$  from  $u_0$ , we can delete the term  $e^{\lambda_1}$  in  $u_0$ . Hence we can assume that  $\langle h_{i_0}, \lambda_1 \rangle > 0$ .

Now assume that  $\langle h_i, \lambda_1 \rangle > 0$  for some  $i \in J_0$ . Then  $\lambda_1$  is regular dominant with respect to  $\{i_0, i\}$ . If  $I = \{i_0, i\}$ , then  $\lambda_1 = \lambda_0$  and it contradicts the hypothesis that  $\lambda_0$  is not regular. Hence  $I \neq \{i_0, i\}$  and  $W_{\{i_0, i\}}$  is a finite group. Therefore (6.3) implies that  $\sum_{w \in W_{\{i_0, i\}}} (-1)^{\ell(w)} a_{w\lambda_1} = 0$ . For  $w \in W_{\{i_0, i\}} \setminus \{e\}$ , we have  $a_{w\lambda_1} = 0$ , because  $d(w\lambda_1) > d(\lambda_1)$ . Thus we obtain the contradiction  $a_{\lambda_1} = 0$ .

We thus conclude  $\langle h_i, \lambda_1 \rangle \leqslant 0$  for all  $i \in J_0$ . Hence  $d(\lambda) < d(\lambda_1)$  for all  $\lambda \in W_{J_0} \lambda_1 \setminus \{\lambda_1\}$ . Then subtracting the  $W_{J_0}$ -invariant  $a_{\lambda_1} \left(\sum_{\lambda \in W_{J_0} \lambda_1} e^{\lambda}\right)$  from  $u_0$ , we can

erase the term  $e^{\lambda_1}$  in  $u_0$ , and the induction proceeds. Note that  $W_{J_0}$  is a finite group.

Thus we have proved (6.4) under the conditions (6.2) and (6.3).

6.2. **Proof of Proposition 5.3 (iii).** We may assume that  $\langle h_0, \lambda_0 \rangle < 0$ . Set  $I_0 = I \setminus \{0\}$ , and let  $W_0$  be the subgroup of W generated by  $\{s_i\}_{i\neq 0}$ . Then  $W\lambda_0 \subset W_0\lambda_0 + \mathbb{Z}\delta$ . Hence, as in the proof of (ii), it is enough to show that if  $u_0 \in \bigoplus_{\lambda \in W_0\lambda_0} \mathbb{Z}[e^{\pm \delta}]e^{\lambda}$  satisfies the condition

(6.5) 
$$\sum_{w \in W_{\{0,j\}}} (-1)^{\ell(w)} w u_0 = 0 \text{ if } j \in I_0 \text{ and } W_{\{0,j\}} \text{ is a finite group,}$$

then

(6.6) there exists a decomposition 
$$(1 - e^{\langle h_0, \lambda_0 \rangle \delta})u_0 = z_0 + z_1$$
  
where  $z_0, z_1 \in \bigoplus_{\lambda \in W_0 \lambda_0} \mathbb{Z}[e^{\pm \delta}]e^{\lambda}$  and  $s_0 z_0 = z_0$  and  $z_1$  is  $W_0$ -invariant.

Let us write  $u_0 = \sum_{\lambda \in W_0 \lambda_0} a_\lambda e^\lambda$  with  $a_\lambda \in \mathbb{Z}[e^{\pm \delta}]$ . Let us set  $\theta = \delta - \alpha_0 \in \frac{(\alpha_0, \alpha_0)}{2} \Delta^+$ . Let  $s_\theta$  be the reflection with respect to  $\theta$ :  $s_\theta(\lambda) = \lambda - \frac{2(\theta, \lambda)}{(\theta, \theta)} \theta$ . Then  $s_\theta$  belongs to  $W_0$ , and  $s_0 \lambda_0 = s_\theta \lambda_0 - \langle h_0, \lambda_0 \rangle \delta$ . We have an  $s_0$ -invariant  $z_0 := e^{\lambda_0} + e^{s_0 \lambda_0} = e^{\lambda_0} + e^{-\langle h_0, \lambda_0 \rangle \delta} e^{s_\theta \lambda_0}$ . Subtracting a constant multiple of  $z_0$  from  $u_0$ , we may assume that  $a_{s_\theta \lambda_0}$  vanishes. On the other hand,  $z_1 := \sum_{\lambda \in W_0 \lambda_0} e^{\lambda}$  is a  $W_0$ -invariant. Their linear combination  $e^{-\langle h_0, \lambda_0 \rangle \delta} z_1 - z_0$  has no term  $e^{s_\theta \lambda_0}$  and the coefficient of  $e^{\lambda_0}$  is  $e^{-\langle h_0, \lambda_0 \rangle \delta} - 1$ . Hence subtracting a constant multiple of it from  $(1 - e^{\langle h_0, \lambda_0 \rangle \delta})u_0$ , we may assume that  $a_{\lambda_0}$  and  $a_{s_\theta \lambda_0}$  vanish. Let us set  $\sup(u_0) = \{\lambda \in W_0 \lambda_0 ; a_\lambda \neq 0\}$ . By subtracting an  $s_0$ -invariant from  $u_0$ , we may assume further that

(6.7) 
$$\langle \theta, \lambda \rangle = -\langle h_0, \lambda \rangle > 0 \text{ for any } \lambda \in \text{supp}(u_0).$$

Hence we have reduced the problem to proving

(6.8) if 
$$a_{\lambda_0} = a_{s_{\theta}\lambda_0} = 0$$
 and if  $u_0$  satisfies (6.5) and (6.7), then  $u_0 = 0$ .

If |I| is 2, it is obvious, since  $W_0\lambda_0 = \{\lambda_0, s_\theta\lambda_0\}$ . Let us assume |I| > 2. Hence  $W_{\{0,i\}}$  is a finite group for all  $i \in I_0$ .

For  $\lambda \in W_0 \lambda_0$ , we set

(6.9) 
$$a_{\lambda+n\delta} = e^{-n\delta} a_{\lambda},$$

so that we have  $a_{\lambda+n\delta}e^{\lambda+n\delta}=a_{\lambda}e^{\lambda}$ . Then (6.5) reads as

(6.10) 
$$\sum_{w \in W_{\{0,i\}}} (-1)^{\ell(w)} a_{w\lambda} = 0 \text{ for any } \lambda \in W_0 \lambda_0 + \mathbb{Z}\delta \text{ and } i \in I_0.$$

Note that  $s_0\lambda = s_\theta\lambda - \langle h_0, \lambda \rangle \delta$  and

(6.11) 
$$a_{s_0\lambda} = e^{\langle h_0, \lambda \rangle \delta} a_{s_{\theta}\lambda}.$$

**Sublemma 6.1.** Let  $\lambda \in W_0 \lambda_0$ . If  $k \in I_0$  satisfies  $(\alpha_0, \alpha_k) = 0$ , then  $a_{\lambda} = a_{s_k \lambda}$ .

*Proof.* We may assume that  $\langle h_0, \lambda \rangle < 0$ . Then, (6.7) implies that  $W_{\{0,k\}}\lambda \cap (\sup(u_0) + \mathbb{Z}\delta) \subset \{\lambda, s_k\lambda\}$ , and (6.10) implies the desired result.

Set

$$I_1 := \{k \in I ; (\alpha_0, \alpha_k) = 0 \},$$

and let  $W_1$  be the subgroup of W generated by  $\{s_k; k \in I_1\}$ . Then Sublemma 6.1 implies that

(6.12) 
$$a_{\lambda} = a_{w\lambda} \text{ for any } w \in W_1.$$

Now we shall divide the proof into two cases:

- (A) there exists a  $1 \in I_0 \setminus I_1$  such that  $(\alpha_1, \alpha_1) \neq (\alpha_0, \alpha_0)$ ,
- (B) for all  $i \in I_0 \setminus I_1$ , we have  $(\alpha_i, \alpha_i) = (\alpha_0, \alpha_0)$ .

Case(A) In this case,  $I = \{1\} \sqcup I_1$  and  $\langle h_0, \alpha_1 \rangle \langle h_1, \alpha_0 \rangle = 2$  as seen by the classification of affine Dynkin diagrams. Note that  $\{0, 1\}$  is a Dynkin diagram of type  $C_2$ . Then  $\theta := \delta - \alpha_0 = \sum_{i \neq 0} a_i \alpha_i$  satisfies  $\langle h_0, \theta \rangle = -2$ , and hence we have  $\langle h_0, \alpha_1 \rangle a_1 = -2$ , which implies  $a_1 + \langle h_1, \alpha_0 \rangle = 0$ . Since  $\langle h_1, \theta \rangle = -\langle h_1, \alpha_0 \rangle$ , we have  $\beta := s_1 \theta = (a_1 + \langle h_1, \alpha_0 \rangle) \alpha_1 + \sum_{i \in I_1} a_i \alpha_i = \sum_{i \in I_1} a_i \alpha_i$ . Hence  $\beta$  is a constant multiple of a root in  $\Delta \cap (\sum_{i \in I_1} \mathbb{Z} \alpha_i)$ , and  $s_\beta$  belongs to  $W_1$ . Assuming that  $u_0$  does not vanish, let us choose an element  $\mu$  in supp $(u_0)$ , highest with respect to  $I_0$  (i.e., maximal with respect to the ordering  $\geqslant : \mu \geqslant \mu'$  if  $\mu - \mu' \in \sum_{i \in I_0} \mathbb{Z}_{\geqslant 0} \alpha_i$ ). By (6.12), we have

(6.13) 
$$\langle h_k, \mu \rangle \geqslant 0 \text{ for any } k \in I_1.$$

Let us show that  $\langle h_1, \mu \rangle \geqslant 0$ . Otherwise,  $\mu$  is regular and anti-dominant with respect to  $\{0,1\}$ . By (6.7), we have  $W_{\{0,1\}}\mu \cap (\operatorname{supp}(u_0) + \mathbb{Z}\delta) \subset \{\mu, s_1\mu, s_1s_0\mu, s_1s_0s_1\mu\}$ . We have  $s_1\mu > \mu$ , and hence  $a_{s_1\mu} = 0$ . Since  $s_\beta = s_1s_\theta s_1$ , we have  $a_{s_1s_\theta\mu} = a_{s_\beta s_1\mu} = a_{s_1\mu} = 0$  by (6.12). Hence we have  $a_{s_1s_0\mu} = 0$ . Thus we obtain  $W_{\{0,1\}}\mu \cap (\operatorname{supp}(u_0) + \mathbb{Z}\delta) \subset \{\mu, s_1s_0s_1\mu\}$ . Hence (6.10) implies that  $a_\mu - a_{s_1s_0s_1\mu} = 0$ . On the other hand, we have  $a_{s_1s_0s_1\mu} = e^{\langle h_0, s_1\mu \rangle \delta} a_{s_1s_\theta s_1\mu} = e^{\langle h_0, s_1\mu \rangle \delta} a_{s_\beta\mu} = e^{\langle h_0, s_1\mu \rangle \delta} a_\mu$ . Hence  $(1 - e^{\langle h_0, s_1\mu \rangle \delta}) a_\mu = 0$ . Since  $\langle h_0, s_1\mu \rangle = \langle s_1h_0, \mu \rangle < 0$ , we obtain  $a_\mu = 0$ , which is a contradiction.

We thus conclude that  $\langle h_1, \mu \rangle \geqslant 0$ . Along with (6.13),  $\mu \in W_0 \lambda_0$  is dominant with respect to  $I_0$ , and hence we conclude  $\mu = \lambda_0$ , which contradicts  $a_{\lambda_0} = 0$ .

Case(B) The proof in this case is similar to the one in Case(B), but slightly more complicated. In this case,  $|I_0 \setminus I_1|$  is one or two by the classification of affine Dynkin diagrams. The case  $|I_0 \setminus I_1| = 2$  is exactly the case  $A_n^{(1)}$   $(n \ge 2)$ . Set  $I_0 \setminus I_1 = \{i_1, i_2\}$ 

(when  $|I_0 \setminus I_1| = 1$ , by convention  $i_1 = i_2$ ). We have  $(\alpha_0, \alpha_i) = -1$  for  $i \in I_0 \setminus I_1$ ,  $\theta := \delta - \alpha_0$  is a root, and

$$\delta = \alpha_0 + \alpha_{i_1} + \alpha_{i_2} + \sum_{i \in I_1} a_i \alpha_i, \quad \theta = \alpha_{i_1} + \alpha_{i_2} + \sum_{i \in I_1} a_i \alpha_i$$

(when  $|I_0 \setminus I_1| = 1$ ,  $\delta = \alpha_0 + 2\alpha_{i_1} + \sum_{i \in I_1} a_i \alpha_i$ ). Let  $w \in W_1$  be the longest element of  $W_1$ .

Sublemma 6.2. We have  $ws_{i_1}\theta = \alpha_{i_2}$ .

Proof. We have  $s_{i_1}\theta = \theta - \alpha_{i_1} = \alpha_{i_2} + \sum_{i \in I_1} a_i \alpha_i$ . Moreover, we have  $\langle h_k, s_{i_1}\theta \rangle = -\langle h_k - \langle h_k, \alpha_{i_1} \rangle h_{i_1}, \alpha_0 \rangle \geqslant 0$  for  $k \in I_1$ . Hence  $s_{i_1}\theta$  is dominant with respect to  $I_1$ . Hence  $ws_{i_1}\theta$  is anti-dominant with respect to  $I_1$ :  $\langle h_k, ws_{i_1}\theta \rangle \leqslant 0$  for any  $k \in I_1$ . Write

$$ws_{i_1}\theta = \alpha_{i_2} + \beta.$$

Since  $ws_{i_1}\theta$  is a root,  $\beta$  has the form

$$\beta = \sum_{i \in I_1} m_i \alpha_i \text{ with } m_i \in \mathbb{Z}_{\geqslant 0}.$$

Hence we have  $(\beta, ws_{i_1}\theta) \leq 0$ . On the other hand,  $(\alpha_0, \alpha_0) = (\alpha_{i_2} + \beta, \alpha_{i_2} + \beta)$  implies that  $(\beta, \beta) + 2(\beta, \alpha_{i_2}) = 0$ . Hence we obtain  $(\beta, \beta) = 2(\beta, \alpha_{i_2} + \beta) \leq 0$ , which implies  $\beta = 0$ .

As a corollary, we have  $ws_{i_1}s_{\theta}s_{i_1}w = s_{i_2}$  and

$$(6.14) s_{i_2} w s_{i_1} s_{\theta} = w s_{i_1}.$$

$$(6.15) s_{i_2} s_{\theta} w s_{i_1} s_{\theta} = w$$

Indeed, the last equality follows from  $(ws_{i_2}s_{\theta})(ws_{i_1}s_{\theta}) = (s_{i_1}ws_{i_2})(s_{i_2}ws_{i_1}) = e$ .

Assuming that  $u_0$  does not vanish, let us choose  $\mu$  in supp $(u_0)$ , highest with respect to  $I_0$ . By (6.12), we have

(6.16) 
$$\langle h_k, \mu \rangle \geqslant 0 \text{ for any } k \in I_1.$$

Let us show that  $\langle h_{i_1}, \mu \rangle \geq 0$ . Assume the contrary:  $\langle h_{i_1}, \mu \rangle < 0$ . Then  $\mu$  is regular anti-dominant with respect to  $\{0, i_1\}$ . Since  $s_{i_1}\mu > \mu$ , we have

$$(6.17) a_{s_{i_1}\mu} = 0.$$

The property (6.7) implies that  $W_{\{0,i_1\}}\mu \cap (\operatorname{supp}(u_0) + \mathbb{Z}\delta) \subset \{\mu, s_{i_1}\mu, s_{i_1}s_0\mu\}$ , and hence by (6.10), together with (6.17), we have

$$(6.18) a_{\mu} + a_{s_{i_1}s_0\mu} = 0.$$

Set  $\mu_1 := w s_{i_1} s_{\theta} \mu$ . Then we have  $(\alpha_0, \mu_1) = (\alpha_0, s_{i_1} s_{0} \mu) = (s_0 s_{i_1} \alpha_0, \mu) = (\alpha_{i_1}, \mu) < 0$ and  $(\alpha_{i_2}, \mu_1) = (s_{i_1} w \alpha_{i_2}, s_{\theta} \mu) = (\theta, s_{\theta} \mu) = -(\theta, \mu) < 0$ . Hence  $\mu_1$  is also regular anti-dominant with respect to  $\{0, i_2\}$ . By (6.7), we have  $W_{\{0, i_2\}} \mu_1 \cap (\text{supp}(u_0) + \mathbb{Z}\delta) \subset \{\mu_1, s_{i_2} \mu_1, s_{i_2} s_{0} \mu\}$ . Since  $s_{i_2} \mu_1 = s_{i_2} w s_{i_1} s_{\theta} \mu = w s_{i_1} \mu$  by (6.14), we have  $a_{s_{i_2} \mu_1} = a_{w s_{i_1} \mu} = a_{s_{i_1} \mu} = 0$  by (6.17). Here we used (6.12) in the second equality. Hence we have  $W_{\{0, i_2\}} \mu_1 \cap (\text{supp}(u_0) + \mathbb{Z}\delta) \subset \{\mu_1, s_{i_2} s_{0} \mu_1\}$ , and (6.10) implies that

$$a_{\mu_1} + a_{s_{i_2}s_0\mu_1} = 0.$$

By (6.15), we have  $s_{i_2}s_{\theta}\mu_1 = s_{i_2}s_{\theta}ws_{i_1}s_{\theta}\mu = w\mu$ , which implies  $a_{s_{i_2}s_{\theta}\mu_1} = a_{\mu}$  by (6.12). By (6.9), we have  $a_{s_{i_2}s_{\theta}\mu_1} = e^{\langle h_{0,\mu_1} \rangle}a_{s_{i_2}s_{\theta}\mu_1} = e^{\langle h_{i_1}, \mu \rangle}a_{\mu}$ , and  $a_{\mu_1} = a_{s_{i_1}s_{\theta}\mu} = e^{-\langle h_{0,\mu} \rangle}a_{s_{i_1}s_{\theta}\mu}$ . Thus we obtain  $e^{\langle h_{i_1}, \mu \rangle}a_{\mu} + e^{-\langle h_{0,\mu} \rangle}a_{s_{i_1}s_{\theta}\mu} = 0$ . Together with (6.19) and  $\langle h_{0}, \mu \rangle + \langle h_{i_1}, \mu \rangle < 0$ , we conclude that  $a_{\mu} = 0$ . It is a contradiction.

Hence we have obtained  $\langle h_{i_1}, \mu \rangle \geqslant 0$ . Similarly we have  $\langle h_{i_2}, \mu \rangle \geqslant 0$ . Thus  $\mu$  is dominant with respect to  $I_0$ , and hence  $\mu = \lambda_0$ , which is a contradiction.

## 7. Proof of Proposition 4.5

In this section we shall prove Proposition 4.5 as an application of Theorem 5.2. Corollary 3.3 implies that for any  $w \in W$  we have

(7.1) 
$$p_i^* p_{i*}([\mathscr{O}_{X_w}]) = \begin{cases} [\mathscr{O}_{X_{ws_i}}] & \text{if } ws_i < w, \\ [\mathscr{O}_{X_w}] & \text{if } ws_i > w. \end{cases}$$

**Lemma 7.1.** Let  $J \subset I$ . If  $A \in R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_B(X)$  satisfies the conditions:

- (i)  $p_i^* p_{i*} A = 0$  for all  $i \in J$ ,
- (ii)  $p_i^* p_{i*} A = A \text{ for all } i \in I \setminus J$ ,
- (iii)  $i_e^* A = 0$ ,

then A = 0.

*Proof.* Write  $A = \sum_{w \in W} a_w[\mathscr{O}_{X_w}]$  (infinite sum). Then the condition (iii) implies that  $a_e = 0$ . Let us show  $a_w = 0$  by induction on  $\ell(w)$ . By (7.1), we have

$$p_i^* p_{i*} A = \sum_{x \in W, x s_i > x} (a_x + a_{x s_i}) [\mathscr{O}_{X_x}].$$

If  $xs_i < x$  for some  $i \in I \setminus J$ , then (ii) implies  $a_x = 0$ . The condition (i) implies that  $a_x + a_{xs_i} = 0$  for any x and  $i \in J$ .

Let us take i such that  $ws_i < w$ . If  $i \notin J$ , then  $a_w = 0$ . If  $i \in J$ , then the induction hypothesis implies that  $a_w = -a_{ws_i}$  vanishes.

Let us recall that we have a monomorphism (Corollary 4.3):

$$\beta \colon R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_B(\mathrm{pt}) \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \rightarrowtail R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_B(X).$$

Let us also recall  $1 \otimes \mathcal{D}_i \in \operatorname{End}(R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_B(\operatorname{pt}) \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P])$  which acts on the last factor  $\mathbb{Z}[P]$  as in (2.4). Then, we have a commutative diagram (for  $\xi$  see (4.4)):

Let  $j_w \colon R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[P] \to R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[Q]$  be the homomorphism given by

(7.2) 
$$j_w(e^{\lambda+\alpha}) = e^{w(\lambda+\alpha)-\lambda} \text{ for } \lambda \in L \text{ and } \alpha \in Q.$$

Then we have a commutative diagram

(7.3) 
$$R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[P] \xrightarrow{\xi} R \otimes_{\mathbb{Z}[e^{\pm \delta}]} K_{B}(\text{pt}) \otimes_{\mathbb{Z}[P^{W}]} \mathbb{Z}[P]$$

$$\downarrow^{j_{w}} \qquad \qquad i_{w}^{*} \circ \beta \downarrow$$

$$R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[Q] \hookrightarrow R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[P].$$

We have  $i_e^*[\mathscr{O}_{X_w}] = 0$  for  $w \neq e$ .

Hence in order to prove Proposition 4.5, it is enough to construct  $\mathcal{G}_w \in R \otimes_{\mathbb{Z}[e^{\pm \delta}]}$  $\mathbb{Z}[P]_{[0,-\kappa^*)}$  which satisfies

(7.4) 
$$\begin{cases} (i) & \mathcal{G}_e = 1, \\ (ii) & \mathcal{D}_i(\mathcal{G}_w) = \begin{cases} \mathcal{G}_{ws_i} & \text{if } ws_i < w, \\ \mathcal{G}_w & \text{if } ws_i > w, \end{cases} & \text{for } w \neq e, \\ (iii) & j_e(\mathcal{G}_w) = 0 & \text{for } w \neq e. \end{cases}$$

Then Lemma 7.1 guarantees that

$$\beta \circ \xi(\mathcal{G}_w) = [\mathscr{O}_{X_w}].$$

Hence  $\mathcal{G}_w$  is the affine Grothendieck polynomial.

We shall construct such  $\mathcal{G}_w$ 's by induction on  $\ell(w)$ . Assuming that  $\mathcal{G}_x \in R \otimes_{\mathbb{Z}[e^{\pm \delta}]}$  $\mathbb{Z}[P]_{[0,-\kappa^*)}$  has been constructed for x < w satisfying (7.4), let us construct  $\mathcal{G}_w$ . Note that  $\mathcal{G}_x$  is  $s_i$ -invariant if  $xs_i > x$ , x < w, and  $\mathcal{D}_i \mathcal{G}_x = \mathcal{G}_{xs_i}$  if  $xs_i < x < w$ . Let us set  $J = \{i \in I : ws_i < w\}$ , and  $\rho_J := \sum_{i \in J} \Lambda_i$ . Set  $B = e^{\rho_J} \mathcal{G}_w$ . We have

(7.5) 
$$\mathcal{D}_{i} \circ e^{-\rho_{J}} = e^{-\rho_{J}} (1 - e^{-\alpha_{i}})^{-1} \circ (1 - s_{i}) \text{ for } i \in J,$$

$$(\mathcal{D}_{i} - 1) \circ e^{-\rho_{J}} = e^{-\rho_{J} - \alpha_{i}} (1 - e^{-\alpha_{i}})^{-1} \circ (1 - s_{i}) \text{ for } i \in I \setminus J.$$

Hence the condition (7.4) (ii) reads as

(7.6) 
$$(1 - s_i)B = \begin{cases} e^{\rho_J} (1 - e^{-\alpha_i}) \mathcal{G}_{ws_i} & \text{if } i \in J, \\ 0 & \text{otherwise.} \end{cases}$$

Assume that this equation is solved with

(7.7) 
$$B \in R \underset{\mathbb{Z}[e^{\pm \delta}]}{\otimes} \left( \underset{\langle c, \rho_J \rangle \geqslant \langle c, \lambda \rangle > \langle c, \rho_J - \rho \rangle}{\oplus} \mathbb{Z}e^{\lambda} \right).$$

Set  $C = j_e(B) \in R \otimes_{\mathbb{Z}[e^{\pm \delta}]} \mathbb{Z}[Q]$ . Then  $\eta(C)$  (see (4.2)) belongs to  $R \otimes_{\mathbb{Z}[e^{\pm \delta}]}$  $\mathbb{Z}[P^W]$  and satisfies  $j_e(\eta(C)) = C$ . Therefore,  $\mathcal{G}_w := e^{-\rho_J}(B - \eta(C))$  belongs to  $R \underset{\mathbb{Z}[e^{\pm \delta}]}{\otimes} \left( \underset{0 \geqslant \langle c, \lambda \rangle > -\langle c, \rho \rangle}{\oplus} \mathbb{Z}^{e^{\lambda}} \right)$  and satisfies all the conditions in (7.4).

Thus we reduced the problem to solving the equation (7.6) with (7.7).

In order to solve (7.6) with (7.7), let us apply Theorem 5.2 (ii). Note that  $e^{\rho_J}(1-e^{-\alpha_i})\mathcal{G}_{ws_i} \in R \underset{\mathbb{Z}[e^{\pm \delta}]}{\otimes} (\bigoplus_{\langle c,\rho_J\rangle \geqslant \langle c,\lambda\rangle > \langle c,\rho_J-\rho\rangle} \mathbb{Z}^{e^{\lambda}})$ . Since  $J \neq I$ , we have  $\langle c,\rho\rangle > \langle c,\rho_J\rangle$  and  $\langle c,\rho_J-\rho\rangle \geqslant -\langle c,\rho\rangle$ . Hence, by Theorem 5.2 (ii), it is enough to show the compatibility conditions (i) and (ii) in Lemma 5.1: namely

(7.8) 
$$\begin{cases} (i) \ (1+s_{i})e^{\rho_{J}}(1-e^{-\alpha_{i}})\mathcal{G}_{ws_{i}} = 0 \text{ for all } i \in J, \\ (ii) \ \text{if } i, j \in J, \text{ then} \\ \sum_{x \in W_{\{i,j\}}, xs_{i} > x} (-1)^{\ell(x)} x e^{\rho_{J}} (1-e^{-\alpha_{i}}) \mathcal{G}_{ws_{i}} = \sum_{x \in W_{\{i,j\}}, xs_{j} > x} (-1)^{\ell(x)} x e^{\rho_{J}} (1-e^{-\alpha_{j}}) \mathcal{G}_{ws_{j}}, \\ (iii) \ \text{if } i \in J, j \in I \setminus J \text{ and } W_{\{i,j\}} \text{ is a finite group, then} \\ \sum_{x \in W_{\{i,j\}}, xs_{i} > x} (-1)^{\ell(x)} x e^{\rho_{J}} (1-e^{-\alpha_{i}}) \mathcal{G}_{ws_{i}} = 0. \end{cases}$$

Proof of (7.8) (i) This follows from the fact that  $\mathcal{G}_{ws_i}$  is  $s_i$ -invariant, which implies that  $e^{\rho_J}(1 - e^{-\alpha_i})\mathcal{G}_{ws_i} = (1 - s_i)(e^{\rho_J}\mathcal{G}_{ws_i})$ .

In order to prove (7.8) (ii), (iii), let us recall the following well-known results on Demazure operators. For  $x \in W$ ,  $\mathcal{D}_{i_1} \cdots \mathcal{D}_{i_m}$  does not depend on the choice of reduced expressions  $x = s_{i_1} \cdots s_{i_m}$ . We denote it by  $\mathcal{D}_x$ . We have  $\mathcal{D}_x \mathcal{D}_i = \mathcal{D}_x$  if  $xs_i < x$ .

Proof of (7.8) (ii) Since  $ws_i < w$  and  $ws_j < w$ , w is the longest element in  $wW_{\{i,j\}}$ . Let  $w_1$  be the shortest element of  $wW_{\{i,j\}}$ , and  $w_0$  the longest element of  $W_{\{i,j\}}$ . Hence  $w = w_1w_0$ . Let  $\Delta_{\{i,j\}}^+ = \Delta^+ \cap (\mathbb{Z}\alpha_i + \mathbb{Z}\alpha_j)$ , and  $D = \prod_{\alpha \in \Delta_{\{i,j\}}^+} (1 - e^{-\alpha})$ . Then we have (see e.g. [1, 8])

$$\mathcal{D}_{w_0} = e^{-\rho_J} D^{-1} \sum_{x \in W_{\{i,j\}}} (-1)^{\ell(x)} x \circ e^{\rho_J}.$$

Since  $\mathscr{D}_{w_0}\mathcal{G}_{ws_i} = \mathcal{G}_{w_1} = \mathscr{D}_{w_0}\mathcal{G}_{ws_j}$ , we have

$$\sum_{x \in W_{\{i,j\}}} (-1)^{\ell(x)} x e^{\rho_J} \mathcal{G}_{ws_i} = \sum_{x \in W_{\{i,j\}}} (-1)^{\ell(x)} x e^{\rho_J} \mathcal{G}_{ws_j}.$$

It remains to remark that

$$\sum_{x \in W_{\{i,j\}}} (-1)^{\ell(x)} x e^{\rho_J} \mathcal{G}_{ws_i} = \sum_{x \in W_{\{i,j\}}, xs_i > x} (-1)^{\ell(x)} x (1 - s_i) e^{\rho_J} \mathcal{G}_{ws_i}$$

$$= \sum_{x \in W_{\{i,j\}}, xs_i > x} (-1)^{\ell(x)} x e^{\rho_J} (1 - e^{-\alpha_i}) \mathcal{G}_{ws_i}.$$

Proof of (7.8) (iii) Let  $w_0$  be the longest element of  $W_{\{i,j\}}$ , and let  $w_1$  be the shortest element of  $wW_{\{i,j\}}$ . Hence  $w_1w_0$  is the longest element of  $wW_{\{i,j\}}$ .

It is enough to show that

$$\sum_{x \in W_{\{i,j\}}, xs_i > x} (-1)^{\ell(x)} x e^{\Lambda_i} (1 - e^{-\alpha_i}) \mathcal{G}_{ws_i} = 0.$$

Since  $e^{\Lambda_i}(1 - e^{-\alpha_i})\mathcal{G}_{ws_i} = (1 - s_i)e^{\Lambda_i}\mathcal{G}_{ws_i}$ , it is enough to show

(7.9) 
$$\sum_{x \in W_{\{i,j\}}} (-1)^{\ell(x)} x e^{\Lambda_i} \mathcal{G}_{ws_i} = 0.$$

Since  $ws_j > w > ws_i$ , we have  $\mathcal{D}_{w_0s_i}\mathcal{G}_{ws_i} = \mathcal{G}_{w_1} = \mathcal{D}_{w_0}\mathcal{G}_{ws_i}$ . Hence we have

$$(\mathscr{D}_{w_0 s_i} - \mathscr{D}_{w_0}) \mathcal{G}_{w s_i} = 0.$$

Setting  $K = (\mathcal{D}_{w_0 s_i} - \mathcal{D}_{w_0}) \circ e^{-\Lambda_i}$ , we obtain

$$(7.10) Ke^{\Lambda_i} \mathcal{G}_{ws_i} = 0.$$

Since  $K = (\mathscr{D}_{w_0 s_j} - \mathscr{D}_{w_0}) \circ \mathscr{D}_i \circ e^{-\Lambda_i}$  and  $\mathscr{D}_i \circ e^{-\Lambda_i} = e^{-\Lambda_i} (1 - e^{-\alpha_i})^{-1} (1 - s_i)$ , we have

$$(7.11) K \circ (1+s_i) = 0.$$

On the other hand, we have  $K \circ \mathcal{D}_j = (\mathcal{D}_{w_0 s_j} - \mathcal{D}_{w_0}) \circ \mathcal{D}_j \circ e^{-\Lambda_i} = 0$ . Since  $\mathcal{D}_j = (1+s_j) \circ (1-e^{-\alpha_j})^{-1}$ , we have  $0 = K \circ (1+s_j) \circ (1-e^{-\alpha_j})^{-1}$ , which implies  $K \circ (1+s_j) = 0$ . Together with (7.11), K can be written as  $K = \psi \circ E$  for some  $\psi$  in the quotient field of  $\mathbb{Z}[P]$ . Here,  $E = \sum_{x \in W_{\{i,j\}}} (-1)^{\ell(x)} x$ . Since  $K \neq 0$ ,  $\psi$  does not vanish and (7.10) implies the desired result  $E e^{\Lambda_i} \mathcal{G}_{ws_i} = 0$ .

This completes the proof of (7.8).

#### 8. Global Cohomology Character formulas

The affine Grothendieck polynomials give the character formula for the cohomologies of  $\mathscr{O}_{X_w}(\mu) := \mathscr{O}_{X_w} \otimes \mathscr{O}_X(\mu)$  under certain conditions on  $\mu \in P$ .

For  $w \in W$ , the *B*-orbit  $X_w$  is contained in a *T*-stable open affine set  $V_w := wBx_0$  as a closed subset. As a scheme with *T*-action,  $V_w$  is isomorphic to the group scheme  ${}^wU$  whose Lie algebra is  $\bigoplus_{\alpha \in w\Delta^+} \mathfrak{g}_{\alpha}$ . We have a commutative diagram

$$\overset{\circ}{X}_{w} \xrightarrow{\sim} \prod_{\alpha \in \Delta^{+} \cap w \Delta^{+}} \mathfrak{g}_{\alpha}$$

$$\downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$V_{w} \xrightarrow{\sim} \prod_{\alpha \in w \Delta^{+}} \mathfrak{g}_{\alpha}.$$

Let  $\operatorname{Coh}_T(\mathscr{O}_{V_w})$  be the abelian category of coherent T-equivariant  $\mathscr{O}_{V_w}$ -modules.

**Lemma 8.1.** Any  $\mathscr{F} \in \operatorname{Coh}_T(\mathscr{O}_{V_w})$  admits a free resolution in  $\operatorname{Coh}_T(\mathscr{O}_{V_w})$ :

$$(8.1) 0 \to F_n \otimes \mathscr{O}_{V_w} \to \cdots \to F_1 \otimes \mathscr{O}_{V_w} \to F_0 \otimes \mathscr{O}_{V_w} \to \mathscr{F} \to 0,$$

where  $F_k$  are finite-dimensional T-modules.

Proof. Set  $E = \bigoplus_{\alpha \in w\Delta^+} (\mathfrak{g}_{\alpha})^*$ . Then  $V_w$  is isomorphic to  $\operatorname{Spec}(S(E))$ . Hence, there exists a finite-dimensional T-stable subspace  $E' \subset E$  such that  $\mathscr{F}$  is the pull back of a coherent T-equivariant sheaf on  $\operatorname{Spec}(S(E'))$  by the faithfully flat projection  $\operatorname{Spec}(S(E)) \to \operatorname{Spec}(S(E'))$ . Hence the assertion is a consequence of the following well-known lemma.

**Lemma 8.2.** Let E be a finite-dimensional T-module whose weights are contained in  $\{\lambda \in P : \langle h, \lambda \rangle > 0\}$  for some  $h \in P^*$ . Then for any T-equivariant  $\mathscr{O}_E$ -module  $\mathscr{F}$ , there exists a free resolution of  $\mathscr{F}$  in  $Coh_T(\mathscr{O}_E)$ :

$$0 \to F_n \otimes \mathscr{O}_E \to \cdots \to F_1 \otimes \mathscr{O}_E \to F_0 \otimes \mathscr{O}_E \to \mathscr{F} \to 0,$$

where  $F_k$  are finite-dimensional T-modules.

For a locally closed subset S of a topological space Z, we denote by  $H_S^k(Z; \bullet)$  the k-th relative cohomology, and by  $\mathscr{H}_{S}^{k}(\bullet)$  the k-th local cohomology (see e.g. [2, 5]). The following results are proved in [4]

**Lemma 8.3.** For  $w \in W$  and  $\mu \in P$ , we have

- (i)  $H^k_{\overset{\circ}{X}^w}(X;\mathscr{O}_X(\mu)) = 0$  for  $k \neq \ell(w)$ , (ii)  $H^{\ell(w)}_{\overset{\circ}{Y}}(X;\mathscr{O}_X(\mu))$  is isomorphic to the dual Verma module with highest weight

$$\operatorname{ch}\left(H_{\mathring{X}_w}^{\ell(w)}(X;\mathscr{O}_X(\mu))\right) = \frac{\mathrm{e}^{w(\mu+\rho)-\rho}}{\prod_{\alpha\in\Delta^+}(1-\mathrm{e}^{-\alpha})^{\dim\mathfrak{g}_\alpha}}.$$

Here, for a T-module M such that its weight space  $M_{\lambda}$  of weight  $\lambda$  is finitedimensional for any  $\lambda \in P$ , we set

(8.2) 
$$\operatorname{ch}(M) = \sum_{\lambda \in P} (\dim M_{\lambda}) e^{\lambda}.$$

**Lemma 8.4.** For any  $w \in W$  and any coherent T-equivariant  $\mathcal{O}_{V_w}$ -module  $\mathscr{F}$ , we

- (i)  $H_{\overset{\circ}{X}_w}^k(V_w; \mathscr{F}) = 0 \text{ for } k > \ell(w),$
- (ii) For any  $\xi \in P$ , dim  $H_{\overset{\circ}{X}_w}^k(V_w; \mathscr{F})_{\xi} < \infty$ ,
- (iii) there exists a finite subset S of P such that the set of weights of  $H_{\stackrel{\circ}{\mathcal{V}}}^k$   $(V_w;\mathscr{F})$

is contained in 
$$S + Q_-$$
, where  $Q_- := \sum_i \mathbb{Z}_{\leqslant 0} \alpha_i$ ,  
(iv)  $\sum_k (-1)^k \operatorname{ch}(H^k_{\overset{\circ}{X}_w}(V_w; \mathscr{F})) = (-1)^{\ell(w)} \operatorname{ch}(H^{\ell(w)}_{\overset{\circ}{X}_w}(X; \mathscr{O}_X)) \cdot \left(\sum_k (-1)^k \operatorname{ch}(L_k i_w^* \mathscr{F})\right)$ .

*Proof.* Since  $X_w$  is a closed subscheme of the affine scheme  $V_w$  defined as the intersection of the zero loci of  $f_i$   $(1 \leq i \leq \ell(w))$  for some  $f_i \in \mathcal{O}_{V_w}(V_w)$ , we obtain (i).

Let us prove the other statements. Let us take a free resolution as in (8.1). Then  $H^k_{\overset{\circ}{X}_w}(V_w;\mathscr{F})$  is the cohomology group of  $F_{\bullet}\otimes H^{\ell(w)}_{\overset{\circ}{V}}(V_w;\mathscr{O}_X)$ . Hence the results follow from the corresponding fact for  $H_{X_w}^k(V_w; \mathscr{O}_X)$  in Lemma 8.3 and  $\sum_{k} (-1)^k \operatorname{ch}(L_k i_w^* \mathscr{F}) = \sum_{k} (-1)^k \operatorname{ch}(F_k).$ 

By this lemma, we obtain the following result.

**Proposition 8.5.** (i) For  $w \in W$  and a B-stable quasi-compact open subset  $\Omega$  of X such that  $X_w \subset \Omega$ , we have homomorphisms

(8.3) 
$$K_B(X) \longrightarrow K_B(\Omega) \longrightarrow \prod_{\lambda \in P} \mathbb{Z} e^{\lambda}$$

given by 
$$[\mathscr{F}] \mapsto \sum_{k=0}^{\infty} (-1)^k \operatorname{ch} \left( H_{\mathring{X}_w}^k(\Omega; \mathscr{F}) \right).$$

(ii)  $\sum_{k} (-1)^k \operatorname{ch}(H^k_{\overset{\circ}{X}_w}(X;\mathscr{F})) = (-1)^{\ell(w)} \operatorname{ch}(H^{\ell(w)}_{\overset{\circ}{X}_w}(X;\mathscr{O}_X)) \cdot \operatorname{ch}(i_w^*([\mathscr{F}])) \text{ for any } \mathscr{F} \in \operatorname{Coh}_B(\mathscr{O}_X).$ 

**Lemma 8.6.** Let  $w, x \in W$  and  $\mu \in P$ . Then, we have

$$\mathscr{H}^k_{\overset{\circ}{X}_x}(\mathscr{O}_{X_w}) = 0 \ and \ H^k_{\overset{\circ}{X}_x}(X;\mathscr{O}_{X_w}(\mu)) = 0 \ unless \ x \geqslant w \ and \ k = \ell(x) - \ell(w).$$

*Proof.* We may assume that  $X_x \subset X_w$ . Set  $\omega_{X_w} = \mathscr{E}xt^{\ell(w)}_{\mathscr{O}_X}(\mathscr{O}_{X_w}, \mathscr{O}_X)$ . Since  $\mathscr{O}_{X_w}$  is Cohen-Macaulay by Proposition 3.4, we have  $\mathscr{O}_{X_w} = R\mathscr{H}om_{\mathscr{O}_X}(\omega_{X_w}, \mathscr{O}_X)[\ell(w)]$ , and

$$\mathscr{H}^{k}_{\overset{\circ}{X}_{x}}(\mathscr{O}_{X_{w}}) = \mathscr{E}xt^{k+\ell(w)-\ell(x)}_{\mathscr{O}_{X}}(\omega_{X_{w}}, \mathscr{H}^{\ell(x)}_{\overset{\circ}{X}_{x}}(\mathscr{O}_{X})).$$

Let  $\xi$  be a generic point of  $\overset{\circ}{X}_x$ . Since  $\mathscr{H}^{\ell(x)}_{\overset{\circ}{X}_x}(\mathscr{O}_X)_{\xi}$  is an injective  $(\mathscr{O}_X)_{\xi}$ -module (see [2]),  $\mathscr{H}^k_{\overset{\circ}{X}_x}(\mathscr{O}_{X_w})_{\xi} = 0$  for  $k \neq \ell(x) - \ell(w)$ . Since  $\mathscr{H}^k_{\overset{\circ}{X}_x}(\mathscr{O}_{X_w})$  is a quasi-coherent B-equivariant  $\mathscr{O}_X$ -module and  $\overset{\circ}{X}_x$  is a B-orbit,  $\mathscr{H}^k_{\overset{\circ}{X}_x}(\mathscr{O}_{X_w})|_{V_w} = 0$ . Let  $j: V_w \hookrightarrow X$  be the inclusion. Since j is affine,  $\mathscr{H}^k_{\overset{\circ}{X}_x}(\mathscr{O}_{X_w}) = j_*j^{-1}\mathscr{H}^k_{\overset{\circ}{X}_x}(\mathscr{O}_{X_w}) = 0$  for  $k \neq \ell(x) - \ell(w)$ .

Since 
$$\overset{\circ}{X}_x$$
 is affine,  $H^k_{\overset{\circ}{X}_x}(X;\mathscr{O}_{X_w}(\mu)) = \Gamma(\overset{\circ}{X}_x;\mathscr{H}^k_{\overset{\circ}{X}_x}(\mathscr{O}_{X_w}(\mu))) = 0$  for  $k \neq \ell(x) - \ell(w)$ .

Note that Lemmas 8.1–8.6 still hold for any symmetrizable Kac-Moody Lie algebra  $\mathfrak{g}$ . Now we shall use the fact that  $\mathfrak{g}$  is affine.

**Lemma 8.7.** For  $w \in W$ , let us write

(8.4) 
$$\mathcal{G}_w = \sum_{(\lambda,\alpha)\in L\times Q} a_{\lambda,\alpha} e^{\lambda+\alpha} \quad \text{with } a_{\lambda,\alpha} \in R.$$

Then

$$\operatorname{ch}(H_{X_x}^{\ell(x)-\ell(w)}(X; \mathscr{O}_{X_w}(\mu))) = (-1)^{\ell(w)} \frac{e^{x(\mu+\rho)-\rho} j_x(\mathcal{G}_w)}{\prod_{\alpha \in \Delta^+} (1 - e^{-\alpha})^{\dim \mathfrak{g}_\alpha}}$$

$$= (-1)^{\ell(w)} \frac{\sum_{\alpha \in \Delta^+} a_{\lambda,\alpha} e^{x(\mu+\lambda+\alpha+\rho)-\lambda-\rho}}{\prod_{\alpha \in \Delta^+} (1 - e^{-\alpha})^{\dim \mathfrak{g}_\alpha}}.$$

*Proof.* By Lemma 8.3 and Proposition 8.5, we have

$$\begin{split} (-1)^{\ell(x)-\ell(w)} & \operatorname{ch}(H_{\mathring{X}_x}^{\ell(x)-\ell(w)}(X;\mathscr{O}_{X_w}(\mu))) \\ &= \sum_k (-1)^k \operatorname{ch}(H_{\mathring{X}_x}^k(X;\mathscr{O}_{X_w}(\mu))) \\ &= \sum_k (-1)^k \sum_{(\lambda,\alpha) \in L \times Q} a_{\lambda,\alpha} \operatorname{e}^{-\lambda} \operatorname{ch}\big(H_{\mathring{X}_x}^k(X;\mathscr{O}_X(\mu+\lambda+\alpha))\big) \\ &= (-1)^{\ell(x)} \sum_{(\lambda,\alpha) \in L \times Q} a_{\lambda,\alpha} \operatorname{e}^{-\lambda} \operatorname{ch}\big(H_{\mathring{X}_x}^{\ell(x)}(X;\mathscr{O}_X(\mu+\lambda+\alpha))\big), \end{split}$$

which implies the desired result.

For  $\ell \in \mathbb{Z}_{\geq 0}$ , let  $(X_w)_{\ell} = \cup_x X_x$  where x ranges over the elements of W such that  $x \geq w$  and  $\ell(x) \geq \ell(w) + \ell$ . Then  $\{(X_w)_{\ell}\}_{\ell \in \mathbb{Z}_{\geq 0}}$  is a decreasing sequence of B-stable

closed subsets of  $X_w$ . Moreover,  $(X_w)_{\ell} \setminus (X_w)_{\ell+1}$  is a disjoint union of  $X_x$  where  $X_x$  ranges over the elements of  $X_x$  where  $X_x$ 

(8.5) 
$$x \geqslant w \text{ and } \ell(x) = \ell(w) + \ell.$$

By Lemma 8.6, we have

$$H_{(X_w)_{\ell}\setminus (X_w)_{\ell+1}}^k(X; \mathscr{O}_{X_w}(\mu)) = 0 \text{ for } k \neq \ell,$$

because it is the direct sum of  $H_{\overset{\circ}{X}_x}^k(X; \mathscr{O}_{X_w}(\mu))$  where  $x \in W$  ranges over the elements satisfying (8.5). Hence, by a general argument (see [2, 8]),  $H^k(X; \mathscr{O}_{X_w}(\mu))$  is the k-th cohomology group of

(8.6) 
$$H^{0}_{X_{w}\setminus(X_{w})_{1}}(X;\mathscr{O}_{X_{w}}(\mu)) \rightarrow H^{1}_{(X_{w})_{1}\setminus(X_{w})_{2}}(X;\mathscr{O}_{X_{w}}(\mu)) \rightarrow H^{2}_{(X_{w})_{2}\setminus(X_{w})_{3}}(X;\mathscr{O}_{X_{w}}(\mu)) \rightarrow \cdots$$

Corollary 8.8. Let  $\mu \in P$  and  $w \in W$ . Assume that  $\langle c, \mu \rangle \geqslant 0$ . Then, for any  $\xi \in P$ ,  $\sum_{k} \dim H^{k}(X; \mathscr{O}_{X_{w}}(\mu))_{\xi}$  is finite.

Proof. Let us set  $d(\alpha) = \sum_i m_i$  for  $\alpha = \sum_i m_i \alpha_i \in Q$ . With the notation as in (8.4), if  $a_{\lambda,\alpha} \neq 0$ , then  $\langle c, \lambda + \alpha + \rho \rangle > 0$  by Proposition 4.5. Hence, we have  $\langle c, \mu + \lambda + \alpha + \rho \rangle > 0$ . Therefore, for any integer n, there are only finitely many  $x \in W$  such that  $d(\mu + \lambda + \alpha + \rho - x(\mu + \lambda + \alpha + \rho)) < n$ . Hence,  $\sum_{x \in W} \dim H_{\hat{X}_x}^{\ell(x) - \ell(w)}(X; \mathscr{O}_{X_w}(\mu))_{\xi}$  is finite by Lemma 8.7. Since  $H^k(X; \mathscr{O}_{X_w}(\mu))$  is a subquotient of  $\bigoplus_{x \in W, \ell(x) = k + \ell(w)} H_{\hat{X}_x}^k(X; \mathscr{O}_{X_w}(\mu))$ , we obtain the desired result.  $\square$ 

Thus we obtain the following proposition.

**Proposition 8.9.** Let  $w \in W$  and  $\mu \in P$ . Assume that  $\langle c, \mu \rangle \geqslant 0$ . Then we have, with the notation (8.4),

(8.7) 
$$\sum_{k} (-1)^{k} \operatorname{ch} \left( H^{k}(X; \mathscr{O}_{X_{w}}(\mu)) \right) = \sum_{(\lambda, \alpha) \in L \times Q} a_{\lambda, \alpha} e^{-\lambda} \chi_{\mu + \lambda + \alpha},$$

where 
$$\chi_{\mu} = \frac{\sum_{x \in W} (-1)^{\ell(x)} e^{x(\mu+\rho)-\rho}}{\prod_{\alpha \in \Delta^{+}} (1 - e^{-\alpha})^{\dim \mathfrak{g}_{\alpha}}}.$$

Note that  $\sum_{k} (-1)^k \operatorname{ch}(H^k(X; \mathcal{O}_{X_w}(\mu)))$  has a sense by Corollary 8.8.

*Proof.* We have

$$\sum_{k} (-1)^{k} \operatorname{ch} \left( H^{k}(X; \mathscr{O}_{X_{w}}(\mu)) \right) = \sum_{k} (-1)^{k} \operatorname{ch} \left( H^{k}_{(X_{w})_{k} \setminus (X_{w})_{k+1}}(X; \mathscr{O}_{X_{w}}(\mu)) \right) \\
= \sum_{x \in W} (-1)^{\ell(x) - \ell(w)} \operatorname{ch} \left( H^{\ell(x) - \ell(w)}_{X_{x}}(X; \mathscr{O}_{X_{w}}(\mu)) \right),$$

and Lemma 8.7 implies the desired result.

Conjecture 8.10. We conjecture that, if  $\mu$  is dominant, then  $H^k(X; \mathcal{O}_{X_w}(\mu)) = 0$  for  $k \neq 0$  and

(8.8) 
$$\Gamma(X; \mathscr{O}_X(\mu)) \to \Gamma(X; \mathscr{O}_{X_w}(\mu))$$

is surjective.

Note that  $\Gamma(X; \mathscr{O}_X(\mu))$  is isomorphic to the irreducible  $\mathfrak{g}$ -module  $V(\mu)$  with highest weight  $\mu$ , and the kernel N of (8.8) is equal to  $\{v \in V(\mu); U(\mathfrak{b})v \cap V(\mu)_{w\mu} = 0\}$ . The module  $V(\mu)$  has a non-degenerate symmetric bilinear form with respect to which the  $e_i$ 's and the  $f_i$ 's are adjoint to each other, and N is orthogonal to  $U(\mathfrak{b}_-)u_{w\mu}$ . Here  $u_{w\mu}$  is a non-zero vector in the one-dimensional weight space  $V(\mu)_{w\mu}$ . Hence if the conjecture is true,  $\bigoplus_{\xi \in P} (\Gamma(X; \mathscr{O}_{X_w}(\mu))_{\xi})^*$  is isomorphic to  $U(\mathfrak{b}_-)u_{w\mu} \subset V(\mu)$ . By Proposition 8.9, Conjecture 8.10 implies

(8.9)  $\operatorname{ch}(U(\mathfrak{b}_{-})u_{w\mu}) = \sum_{(\lambda,\alpha)\in L\times Q} a_{\lambda,\alpha} e^{-\lambda} \chi_{\mu+\lambda+\alpha}$  for any dominant integral weight  $\mu$ .

### 9. Equivariant cohomology

Theorem 4.4 implies a similar result on the equivariant cohomology of affine flag manifolds. For a B-stable quasi-compact open subset  $\Omega$  of X, the equivariant cohomology  $H_B^*(\Omega, \mathbb{C})$  is a free module over the ring  $H_B^*(\operatorname{pt}, \mathbb{C}) \simeq \mathbb{C}[\mathfrak{t}] = S(\mathfrak{t}^*)$  generated by the B-equivariant cohomology classes  $[X_w]$  ( $X_w \subset \Omega$ ). For any k,  $H_B^k(X,\mathbb{C}) \simeq H_B^k(\Omega,\mathbb{C})$  if  $\Omega$  is large enough (if  $2\operatorname{codim}(X \setminus \Omega) > k+1$ ). Hence we have

$$H_B^*(X,\mathbb{C}) = \bigoplus_{w \in W} H_B^*(\mathrm{pt},\mathbb{C})[X_w].$$

We have a homomorphism  $S(\mathfrak{t}^*) \to H_B^*(X,\mathbb{C})$  by  $P \ni \lambda \mapsto c_2(\mathscr{O}_X(\lambda)) \in H_B^2(X,\mathbb{C})$ , where  $c_2$  is the second Chern class. It induces an  $H_B^*(\mathrm{pt})$ -linear homomorphism

$$H_B^*(\mathrm{pt}) \otimes_{S(\mathfrak{t}^*)^W} S(\mathfrak{t}^*) \to H_B^*(X,\mathbb{C}).$$

Let us write  $\widehat{H}_B^*(X,\mathbb{C}) = \prod_k H_B^k(X,\mathbb{C})$ . The equivariant Chern character defines a homomorphism

$$\operatorname{ch}_B \colon K_B(X) \to \widehat{H}_B^*(X, \mathbb{C}).$$

Hence we have a commutative diagram

$$K_B(\operatorname{pt}) \otimes \mathbb{Z}[P] \longrightarrow K_B(X)$$
 $\exp \otimes \exp \bigvee \qquad \qquad \qquad \downarrow \operatorname{ch}_B$ 
 $\widehat{H}_B^*(\operatorname{pt}, \mathbb{C}) \otimes \widehat{S}(\mathfrak{t}^*) \longrightarrow \widehat{H}_B^*(X, \mathbb{C})$ 

where  $\widehat{S}(\mathfrak{t}^*) = \prod_n S^n(\mathfrak{t}^*)$  and  $\exp \colon \mathbb{Z}[P] \to \widehat{S}(\mathfrak{t}^*)$  is given by  $P \ni \lambda \mapsto \sum_n \lambda^n/n! \in \widehat{S}(\mathfrak{t}^*)$ .

Since the component of  $\operatorname{ch}_B(\mathscr{O}_{X_w})$  of degree  $2\ell(w)$  coincides with  $[X_w]$ , we can translate Theorem 4.4 as follows.

**Theorem 9.1.**  $\mathbb{C}[\delta, \delta^{-1}] \otimes_{\mathbb{C}[\delta]} H_B^*(\mathrm{pt}, \mathbb{C}) \otimes_{S(\mathfrak{t}^*)^W} S(\mathfrak{t}^*) \xrightarrow{\sim} \mathbb{C}[\delta, \delta^{-1}] \otimes_{\mathbb{C}[\delta]} H_B^*(X, \mathbb{C}).$ 

Note that  $S(\mathfrak{t}^*)^W \simeq S(\mathfrak{t}^{*W})[\Delta]$ , where  $\Delta$  is the Casimir operator

$$\Delta = \sum_{i \in I} \frac{1}{(\alpha_i, \alpha_i)} \Lambda_i \cdot \alpha_i - \sum_{i,j \in I} \frac{(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)(\alpha_j, \alpha_j)} \Lambda_i \cdot \Lambda_j$$
$$= \frac{1}{2} \sum_{i \in I} \frac{1}{(\alpha_i, \alpha_i)} \Lambda_i \cdot (\alpha_i + \eta(\alpha_i)).$$

**Remark 9.2.** The absolute K-group K(X) is similarly defined. It is isomorphic to  $\prod_{w \in W} \mathbb{Z}[\mathscr{O}_{X_w}]$  and there is a Borel map  $\mathbb{Z}[P] \to K(X)$ . However, Theorem 4.4 gives no information on this map because  $1 - e^{n\delta}$  vanishes in K(X).

### 10. Examples of Affine Grothendieck Polynomials

It is easy to verify directly that

$$\mathcal{G}_{s_i} = 1 - e^{-\Lambda_i}$$
 for all  $i \in I$ .

Indeed, we have an exact sequence in  $\operatorname{Coh}_B(\mathscr{O}_X)$ :

$$0 \to \mathbb{C}_{\Lambda_i} \otimes \mathscr{O}_X(-\Lambda_i) \to \mathscr{O}_X \to \mathscr{O}_{X_{s_i}} \to 0,$$

where  $\mathbb{C}_{\Lambda_i}$  is the one-dimensional *B*-module with weight  $\Lambda_i$ .

Slightly more generally, if  $\{s_j; j \in J\}$  is a collection of mutually commuting simple reflections for some  $J \subset I$ , then

$$\mathcal{G}_{\prod_{j\in J} s_j} = \prod_{j\in J} (1 - e^{-\Lambda_j}).$$

The proof of the existence of the elements  $\mathcal{G}_w$  given in this paper yields an algorithm to compute them. We have implemented this algorithm and used it to provide the examples below, using the notation  $q = e^{\delta}$  and  $E^{\lambda} = \sum_{\mu \in W_{I \setminus \{0\}} \cdot \lambda} e^{\mu}$  for the sum of the exponentials of the weights in the orbit of the element  $\lambda \in P$  under the Weyl group  $W_{I \setminus \{0\}}$  of the classical subalgebra. Usually  $\lambda$  is taken to be anti-dominant with respect to  $W_{I \setminus \{0\}}$ . We write only the subscripts of the simple reflections to indicate Weyl group elements, so that  $\mathcal{G}_{10}$  means  $\mathcal{G}_{s_1s_0}$ , for example.

For 
$$A_{n-1}^{(1)}$$
  $(n \ge 2)$   $(I = \mathbb{Z}/n\mathbb{Z} = \{0, 1, \dots, n-1\})$ :

$$\mathcal{G}_{10} = 1 + (1 - q)^{-1} \{ q E^{-\Lambda_0} - E^{-\Lambda_1} + \sum_{k=1}^{n-1} E^{-\Lambda_1 - \Lambda_k + \Lambda_{k+1} + \alpha_1 + \dots + \alpha_k} \}.$$

For  $A_1^{(1)}$ :

$$\mathcal{G}_{010} = 1 + ((1-q)(1-q^2))^{-1} \{ -(1+q)(1+q^2) E^{-\Lambda_0} - q E^{-\Lambda_0 - \alpha_1} - q^2 E^{\Lambda_0 - 2\Lambda_1 + \alpha_1} + (q+q^2) E^{-\Lambda_1} + (q+q^2) E^{-2\Lambda_0 + \Lambda_1} - q^2 E^{-3\Lambda_0 + 2\Lambda_1 - \alpha_1} \},$$

$$\mathcal{G}_{1010} = 1 + ((1-q)(1-q^2)(1-q^3))^{-1} \{ q(1+q+q^2)(1+q+q^3) E^{-\Lambda_0} + (q^2+q^3+q^4) E^{-\Lambda_0 - \alpha_1} + q^2 E^{3\Lambda_0 - 4\Lambda_1 + 2\alpha_1} - (q+q^2+q^3) E^{2\Lambda_0 - 3\Lambda_1 + \alpha_1} + (q+q^2+q^3) E^{\Lambda_0 - 2\Lambda_1} + (1+q+q^2)(1+q+q^3) E^{\Lambda_0 - 2\Lambda_1 + \alpha_1} - (1+q+q^2)(1+q^2+q^3) E^{-\Lambda_1} - q^2 E^{-\Lambda_1 - \alpha_1} - (q^3+q^4+q^5) E^{-2\Lambda_0 + \Lambda_1} + q^5 E^{-3\Lambda_0 + 2\Lambda_1 - \alpha_1} \},$$

$$\begin{split} \mathcal{G}_{01010} &= 1 + ((1-q)(1-q^2)(1-q^3)(1-q^4))^{-1} \big\{ \\ &- (1+q^2)(1+q+q^2)(1+q^2+2q^3+q^4+q^6) \mathcal{E}^{-\Lambda_0} - q^4 \mathcal{E}^{-\Lambda_0-2\alpha_1} \\ &- q(1+q)(1+q^2)(1+q^2+q^3+q^4) \mathcal{E}^{-\Lambda_0-\alpha_1} - q^6 \mathcal{E}^{3\Lambda_0-4\Lambda_1+2\alpha_1} \\ &+ q^4(1+q)(1+q^2) \mathcal{E}^{2\Lambda_0-3\Lambda_1+\alpha_1} - q^3(1+q^2)(1+q+q^2) \mathcal{E}^{\Lambda_0-2\Lambda_1} \\ &- q^2(1+q)(1+q^2)(1+q+q^2+q^4) \mathcal{E}^{-\Lambda_0-2\Lambda_1+\alpha_1} \\ &+ q(1+q)^2(1+q^2)(1+q^2+q^4) \mathcal{E}^{-\Lambda_1} + q^3(1+q)(1+q^2) \mathcal{E}^{-\Lambda_1-\alpha_1} \\ &+ q(1+q)^2(1+q^2)(1+q^2+q^4) \mathcal{E}^{-2\Lambda_0+\Lambda_1} + q^3(1+q)(1+q^2) \mathcal{E}^{-2\Lambda_0+\Lambda_1-2\alpha_1} \\ &- q^3(1+q^2)(1+q+q^2) \mathcal{E}^{-3\Lambda_0+2\Lambda_1} \\ &- q^2(1+q)(1+q^2)(1+q+q^2+q^4) \mathcal{E}^{-3\Lambda_0+2\Lambda_1-\alpha_1} \\ &+ q^4(1+q)(1+q^2) \mathcal{E}^{-4\Lambda_0+3\Lambda_1-2\alpha_1} - q^6 \mathcal{E}^{-5\Lambda_0+4\Lambda_1-2\alpha_1} \big\}. \end{split}$$

For  $A_2^{(1)}$ :

$$\begin{split} \mathcal{G}_{010} &= 1 - \mathrm{e}^{-\Lambda_0 - \Lambda_1} + \left(1 - q\right)^{-1} \left\{ -q \mathrm{e}^{-\Lambda_0} (\mathrm{e}^{-\alpha_1} + \mathrm{e}^{-\alpha_1 - \alpha_2}) \right. \\ &- \mathrm{e}^{-\Lambda_1} (\mathrm{e}^{\alpha_1} + \mathrm{e}^{\alpha_1 + \alpha_2}) + \mathrm{e}^{\Lambda_0 - \Lambda_1 - \Lambda_2 + \alpha_1 + \alpha_2} + q \mathrm{e}^{-\Lambda_0 + \Lambda_1 - \Lambda_2 - \alpha_1} \\ &+ q \mathrm{e}^{-2\Lambda_0 + \Lambda_2 - \alpha_1 - \alpha_2} + \mathrm{e}^{-2\Lambda_1 + \Lambda_2 + \alpha_1} \right\}, \\ \mathcal{G}_{210} &= 1 + ((1 - q)(1 - q^2))^{-1} \left\{ -(q^2 + q^3) \mathrm{E}^{-\Lambda_0} + (q + q^2) \mathrm{E}^{-\Lambda_1} \right. \\ &- \mathrm{E}^{2\Lambda_0 - 3\Lambda_2 + \alpha_1 + 2\alpha_2} + (1 + q) \mathrm{E}^{\Lambda_0 - 2\Lambda_2 + \alpha_2} - (q + q^2) \mathrm{E}^{\Lambda_1 - 2\Lambda_2 + \alpha_2} \\ &- (1 + q) \mathrm{E}^{-\Lambda_2} - q \mathrm{E}^{-\Lambda_2 - \alpha_1} - (q + q^2) \mathrm{E}^{\Lambda_0 - \Lambda_1 - \Lambda_2 + \alpha_1 + \alpha_2} \\ &+ (q + q^2) \mathrm{E}^{-\Lambda_0 + \Lambda_1 - \Lambda_2 - \alpha_1} - q^2 \mathrm{E}^{-2\Lambda_0 + 2\Lambda_1 - \Lambda_2 - \alpha_1} - q^2 \mathrm{E}^{-2\Lambda_1 + \Lambda_2 + \alpha_1} \right\}, \\ \mathcal{G}_{1210} &= 1 + q^{-1} \mathrm{E}^{-\Lambda_1 - \Lambda_2 + \alpha_1 + \alpha_2} + ((1 - q)(1 - q^2))^{-1} \left\{ -(1 + q + q^2 + q^3) \mathrm{E}^{-\Lambda_0} \right. \\ &- \mathrm{E}^{2\Lambda_0 - 3\Lambda_1 + 2\alpha_1 + \alpha_2} + (1 + q) \mathrm{E}^{\Lambda_0 - 2\Lambda_1 + \alpha_1} - q \mathrm{E}^{-\Lambda_1 - \alpha_2 - \alpha_3} - \mathrm{E}^{2\Lambda_0 - 3\Lambda_2 + \alpha_1 + 2\alpha_2} \\ &+ (1 + q) \mathrm{E}^{\Lambda_0 - 2\Lambda_2 + \alpha_2} - (1 + q + q^2) \mathrm{E}^{\Lambda_1 - 2\Lambda_2 + \alpha_2} - q \mathrm{E}^{-\Lambda_2 - \alpha_1} \\ &- q^{-1} (1 + q + q^2 + q^3) \mathrm{E}^{\Lambda_0 - \Lambda_1 - \Lambda_2 + \alpha_1 + \alpha_2} + (q + q^2) \mathrm{E}^{-\Lambda_0 + \Lambda_1 - \Lambda_2 - \alpha_1} \\ &- q^2 \mathrm{E}^{-2\Lambda_0 + 2\Lambda_1 - \Lambda_2 - \alpha_1} - (1 + q + q^2) \mathrm{E}^{-2\Lambda_1 + \Lambda_2 + \alpha_1} \\ &+ (q + q^2) \mathrm{E}^{-\Lambda_0 - \Lambda_1 + \Lambda_2 - \alpha_2 - \alpha_3} - q^2 \mathrm{E}^{-2\Lambda_0 - \Lambda_1 + 2\Lambda_2 - \alpha_1 - 2\alpha_2 - \alpha_3} \right\}. \end{split}$$

For  $A_3^{(1)}$ :

$$\begin{split} \mathcal{G}_{210} &= 1 + ((1-q)(1-q^2))^{-1} \Big\{ - (q^2 + q^3) \mathbf{E}^{-\Lambda_0} + (q+q^2) \mathbf{E}^{-\Lambda_1} \\ &- (1+q) \mathbf{E}^{\Lambda_0 - 2\Lambda_2 + \alpha_1 + 2\alpha_2 + \alpha_3} - (1+q) \mathbf{E}^{-\Lambda_2} - q \mathbf{E}^{-\Lambda_2 - \alpha_1} \\ &+ (q+q^2) \mathbf{E}^{-\Lambda_0 + \Lambda_1 - \Lambda_2 - \alpha_1} - q^2 \mathbf{E}^{-2\Lambda_0 + 2\Lambda_1 - \Lambda_2 - \alpha_1} - q^2 \mathbf{E}^{-2\Lambda_1 - \Lambda_2 + \alpha_1} \\ &- \mathbf{E}^{2\Lambda_0 - \Lambda_2 - 2\Lambda_3 + \alpha_1 + 2\alpha_2 + 2\alpha_3} - (q+q^2) \mathbf{E}^{\Lambda_0 - \Lambda_1 - \Lambda_3 + \alpha_1 + \alpha_2 + \alpha_3} \\ &+ (1+q) \mathbf{E}^{\Lambda_0 - \Lambda_2 - \Lambda_3 + \alpha_2 + \alpha_3} - (q+q^2) \mathbf{E}^{\Lambda_1 - \Lambda_2 - \Lambda_3 + \alpha_2 + \alpha_3} \\ &+ (1+q) \mathbf{E}^{-2\Lambda_2 + \Lambda_3 + \alpha_2} - (q+q^2) \mathbf{E}^{-\Lambda_0 + \Lambda_1 - 2\Lambda_2 + \Lambda_3 + \alpha_2} \\ &- (q+q^2) \mathbf{E}^{-\Lambda_1 - \Lambda_2 + \Lambda_3 + \alpha_1 + \alpha_2} - \mathbf{E}^{-3\Lambda_2 + 2\Lambda_3 + \alpha_1 + 2\alpha_2} \Big\}. \end{split}$$

For  $D_4^{(1)}$   $(I = \{0, 1, 2, 3, 4\})$  where 2 is the node connected to all others):

$$\mathcal{G}_{12} = (1 - e^{-\Lambda_1})(1 - e^{-\Lambda_1 + \alpha_1}),$$

$$\mathcal{G}_{121} = (1 - e^{-\Lambda_1})(1 - e^{-\Lambda_1 + \alpha_1})(1 - e^{\Lambda_1 - \Lambda_2 - \alpha_1}),$$

$$\mathcal{G}_{321} = 1 - e^{-2\Lambda_3 - \alpha_1 + \alpha_3} + (1 - q)^{-1} \{qe^{-\Lambda_1} + qe^{\Lambda_1 - \Lambda_2 - \alpha_1} + e^{\Lambda_1 - 2\Lambda_3 - \alpha_1 + \alpha_3} + e^{-\Lambda_1 + \Lambda_2 - 2\Lambda_3 + \alpha_3} - e^{-\Lambda_3} - qe^{-\Lambda_3 - \alpha_1} - qe^{-\Lambda_3 - \alpha_1 - \alpha_2} - e^{-\Lambda_3 + \alpha_3} - qe^{-\Lambda_3 - \alpha_1 + \alpha_3} - e^{-\Lambda_3 + \alpha_2 + \alpha_3} - qe^{-\Lambda_3 - \alpha_1 - \alpha_2 - \alpha_4} - e^{-\Lambda_3 + \alpha_2 + \alpha_3 + \alpha_4} + e^{\Lambda_0 - \Lambda_3 - \Lambda_4 + \alpha_2 + \alpha_3 + \alpha_4} + qe^{-\Lambda_0 - \Lambda_3 - \Lambda_4 + \alpha_2 + \alpha_3 + \alpha_4} + qe^{-\Lambda_0 - \Lambda_3 + \Lambda_4 - \alpha_1 - \alpha_2 - \alpha_4} + e^{\Lambda_0 - \Lambda_2 - \Lambda_3 + \Lambda_4 + \alpha_2 + \alpha_3} \}.$$

For  $C_n^{(1)}$  let  $I = \{0, 1, ..., n\}$  and let  $\epsilon_i$  be the *i*-th standard basis vector in  $\mathbb{Z}^n$  for  $1 \le i \le n$  and let  $\alpha_i = \epsilon_i - \epsilon_{i+1}$  for  $1 \le i \le n-1$  and  $\alpha_n = 2\epsilon_n$ . Then we have

$$\mathcal{G}_{10} = 1 + (1 - q)^{-1} \left\{ q E^{-\Lambda_0} - E^{-\Lambda_1} + \sum_{k=1}^{n} E^{-\Lambda_1 + \Lambda_{k-1} - \Lambda_k + \epsilon_1 + \epsilon_k} + \sum_{k=1}^{n-1} E^{-\Lambda_1 - \Lambda_k + \Lambda_{k+1} + \epsilon_1 - \epsilon_{k+1}} \right\}.$$

For  $C_2^{(1)}$ :

$$\mathcal{G}_{010} = 1 + ((1-q)(1-q^2))^{-1} \left\{ -(1+q)^3 E^{-\Lambda_0} - (q+q^2) E^{-\Lambda_0-\alpha_1} - q E^{-\Lambda_0-\alpha_2} - q^2 E^{\Lambda_0-2\Lambda_1+2\alpha_1+\alpha_2} + (q+q^2) E^{-\Lambda_1} + (q+q^2) E^{-2\Lambda_0+\Lambda_1-2\alpha_1-\alpha_2} - q^2 E^{-3\Lambda_0+2\Lambda_1-2\alpha_1-\alpha_2} - q^2 E^{-\Lambda_0+2\Lambda_1-2\Lambda_2+\alpha_2} - (q+q^2) E^{-\Lambda_2+\alpha_1+\alpha_2} + (q+q^2) E^{-\Lambda_0+\Lambda_1-\Lambda_2-\alpha_1} - (q+q^2) E^{-2\Lambda_0+2\Lambda_1-\Lambda_2-\alpha_1} - (q+q^2) E^{-2\Lambda_0+\Lambda_2-\alpha_1-\alpha_2} - (q+q^2) E^{-2\Lambda_1+\Lambda_2+\alpha_1} + (q+q^2) E^{-\Lambda_0-\Lambda_1+\Lambda_2-\alpha_1-\alpha_2} - q^2 E^{-\Lambda_0-2\Lambda_1+2\Lambda_2-\alpha_2} \right\}.$$

## 11. Proof of Lemma 4.2

In this section, we shall give a proof of Lemma 4.2. Let us set  $\mathfrak{t}_{\mathrm{cl}}^* = \mathfrak{t}^*/(\mathfrak{t}^*)^W$  and let cl:  $\mathfrak{t}^* \to \mathfrak{t}_{\mathrm{cl}}^*$  be the canonical projection. Let us recall that  $Q_{\mathrm{cl}} = Q/\mathbb{Z}\delta = \mathrm{cl}(Q)$  and  $W_{\mathrm{cl}}$  is the image of W in  $\mathrm{Aut}(Q_{\mathrm{cl}})$ . Note that  $\mathbb{C} \otimes Q_{\mathrm{cl}}$  is a hyperplane of  $\mathfrak{t}_{\mathrm{cl}}^*$ . For  $\xi_0 \in \mathbb{C} \otimes Q_{\mathrm{cl}}$ , we define  $t(\xi_0) \in \mathrm{Aut}(\mathfrak{t}^*)$  by

(11.1) 
$$t(\xi_0)(\lambda) = \lambda + (\lambda, \delta)\xi - (\lambda, \xi)\delta - \frac{(\lambda, \delta)(\xi, \xi)}{2}\delta$$

for  $\xi \in \mathbb{C} \otimes Q$  such that  $\operatorname{cl}(\xi) = \xi_0$ . It does not depend on the choice of  $\xi$ . Let  $\widetilde{Q}$  be the sublattice  $Q_{\operatorname{cl}} \cap \operatorname{cl}\left(\bigoplus_{i} \frac{2}{(\alpha_i, \alpha_i)} \alpha_i\right)$ . Then we have an exact sequence

$$1 \longrightarrow \widetilde{Q} \stackrel{t}{\longrightarrow} W \longrightarrow W_{cl} \longrightarrow 1.$$

Now we shall prove Lemma 4.2. Assume that  $u \in \mathbb{Z}[P] \otimes_{\mathbb{Z}[P^W]} \mathbb{Z}[P] \simeq \mathbb{Z}[L] \otimes_{\mathbb{Z}} \mathbb{Z}[P]$  satisfies  $i_w^*\beta(u) = 0$  for all  $w \in W$ . We can write uniquely

$$u = \sum_{\lambda \in L} (e^{\lambda} \otimes 1) \cdot \xi(u_{\lambda})$$
 with  $u_{\lambda} \in \mathbb{Z}[P]$ .

Then we have  $\sum_{\lambda \in L} e^{\lambda} j_w(u_{\lambda}) = 0$  for any  $w \in W$  (see (7.2) and the commutative diagram (7.3)). Since  $j_w(u_{\lambda}) \in \mathbb{Z}[Q]$ , we have  $j_w(u_{\lambda}) = 0$  for any  $\lambda \in L$  and  $w \in W$ . Hence we have reduced Lemma 4.2 to the following lemma.

**Lemma 11.1.** Let  $u \in \mathbb{Z}[P]$ . If  $j_{t(\xi_0)}(u) = 0$  for any  $\xi_0 \in \widetilde{Q}$ , then u = 0.

*Proof.* Write  $u = \sum_{(\lambda,\alpha)\in L\times Q} a_{\lambda,\alpha} e^{\lambda+\alpha}$ . For any  $\xi \in Q \cap \operatorname{cl}^{-1}(\widetilde{Q}\setminus\{0\})$  and any integer n, we have  $(\xi,\xi)>0$ , and  $j_{t(\operatorname{cl}(n\xi))}(u)=0$  reads as

$$\sum_{(\lambda,\alpha)\in L\times Q} a_{\lambda,\alpha} e^{\alpha+n\{(\lambda,\delta)\xi-(\lambda+\alpha,\xi)\delta\}-n^2\frac{(\lambda,\delta)(\xi,\xi)}{2}\delta} = 0.$$

Hence we have for all n and  $\ell$ 

$$\sum_{(\lambda,\alpha)\in L\times Q,\,(\lambda,\delta)=\ell} a_{\lambda,\alpha} \mathrm{e}^{\alpha+n\{(\lambda,\delta)\xi-(\lambda+\alpha,\xi)\delta\}} = 0,$$

which implies that

$$\sum_{(\lambda,\alpha)\in L\times Q,\, (\lambda,\delta)=\ell} a_{\lambda,\alpha} \mathrm{e}^{\alpha-n(\lambda+\alpha,\xi)\delta} = 0.$$

Hence for all  $\ell$ , m and  $\xi$ , we have

$$\sum_{(\lambda,\alpha)\in L\times Q,\,(\lambda,\delta)=\ell,\,(\lambda+\alpha,\xi)=m}a_{\lambda,\alpha}e^{\alpha}=0,$$

which implies that

$$\sum_{\lambda \in L, (\lambda, \xi) = m} a_{\lambda, \alpha} = 0$$

for any  $\alpha$ .

Set supp $(u) = \{\lambda \in L : a_{\lambda,\alpha} \neq 0 \text{ for some } \alpha \in Q \}$ . Since there exists  $\xi \in Q$  such that  $(\lambda, \xi) \neq (\lambda', \xi)$  for any pair of distinct elements  $\lambda$ ,  $\lambda'$  in supp(u), we have  $a_{\lambda,\alpha} = 0$  for all  $\lambda$ ,  $\alpha$ .

**Remark 11.2.** If  $\lambda \in P \setminus P^W$ , then  $W\lambda$  is an infinite set. Indeed, there exists  $\xi \in Q \cap \operatorname{cl}^{-1}(\widetilde{Q} \setminus \{0\})$  such that  $(\lambda, \xi) \neq 0$ , and we can easily see that  $\{t(\operatorname{cl}(n\xi))\lambda : n \in \mathbb{Z} \}$  is an infinite set.

#### References

- [1] Demazure, Michel, Désingularisation des variétés de Schubert généralisées, Collection of articles dedicated to Henri Cartan on the occasion of his 70th birthday, I. Ann. Sci. École Norm. Sup. (4) 7 (1974), 53–88.
- [2] Hartshorne, Robin, *Residues and Duality*, Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics **20**, Springer-Verlag, Berlin-New York (1966).
- [3] Kashiwara, Masaki, *The flag manifold of Kac-Moody Lie algebra*, Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), Johns Hopkins Univ. Press, Baltimore, MD, (1989) 161–190.
- [4] \_\_\_\_\_, Kazhdan-Lusztig conjecture for a symmetrizable Kac-Moody Lie algebra, The Grothendieck Festschrift, Vol. II, Progr. Math. 87, Birkhäuser Boston, Boston, MA, (1990) 407–433.
- [5] Kashiwara, Masaki and Schapira, Pierre, *Sheaves on manifolds*, Grundlehren der Mathematischen Wissenschaften **292**, Springer-Verlag, Berlin, (1990).

- [6] \_\_\_\_\_\_, Categories and Sheaves, Grundlehren der Mathematischen Wissenschaften **332**, Springer-Verlag Berlin Heiderberg (2006).
- [7] Kostant, Bertram and Kumar, Shrawan, *T-equivariant K-theory of generalized flag varieties*, J. Differential Geom. **32** (1990), no. 2, 549–603.
- [8] Kumar, Shrawan, Kac-Moody Groups, their Flag Varieties and Representation Theory, Progr. Math., 204, Birkhäuser Boston, Inc., Boston, MA, (2002).
- [9] Lascoux, Alain, and Schützenberger, Marcel-Paul, Structure de Hopf de l'anneau de cohomologie et l'anneau de Grothendieck d'une variété de drapeaux, C. R. Acad. Sci. Paris Sér. I Math. 295 (1982), no. 11, 629–633.

(Masaki KASHIWARA) RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606–8502, JAPAN

(Mark SHIMOZONO) 460 McBryde Hall, Department of Mathematics, Virginia Tech, Blacksburg, VA 24061-0123, USA