# Complete submanifolds of $\mathbb{R}^n$ with finite topology

G. Pacelli Bessa

Luquésio Jorge

J. Fábio Montenegro

#### Abstract

We show that a complete *m*-dimensional immersed submanifold Mof  $\mathbb{R}^n$  with a(M) < 1 is properly immersed and have finite topology, where  $a(M) \in [0, \infty]$  is an scaling invariant number that gives the rate that the norm of the second fundamental form decays to zero at infinity. The class of submanifolds M with a(M) < 1 contains all complete minimal surfaces in  $\mathbb{R}^n$  with finite total curvature, all *m*dimensional minimal submanifolds M of  $\mathbb{R}^n$  with finite total scalar curvature  $\int_M |\alpha|^m dV < \infty$  and all complete 2-dimensional complete surfaces with  $\int_M |\alpha|^2 dV < \infty$  and nonpositive curvature with respect to every normal direction, since a(M) = 0 for them.

## 1 Introduction

Let M be a complete surface minimally immersed in  $\mathbb{R}^n$  and let K be its Gaussian curvature. Osserman [7] for n = 3 and Chern-Osserman [2] for  $n \geq 3$  proved that  $\int_M |K| dV < \infty$  if and only if M is conformally equivalent to a compact Riemann surface  $\overline{M}$  punctured at a finite number of points  $\{p_1, \ldots, p_r\}$  and the Gauss map  $\Phi : M \to \mathbb{G}_{2,n}$  extends to a holomorphic map  $\overline{\Phi} : \overline{M} \to \mathbb{G}_{2,n}$ , see [5] for a beautiful exposition. M. Anderson [1] proved a higher dimension version of Chern-Osserman finite total curvature theorem, i.e. a complete m-dimensional minimally immersed submanifold M of  $\mathbb{R}^n$  has finite total scalar curvature  $\int_M |\alpha|^m dV < \infty$  if and only if M is  $C^{\infty}$ - diffeomorphic to a compact smooth Riemannian manifold  $\overline{M}$  punctured at a finite number of points  $\{p_1, \ldots, p_r\}$  and the Gauss map  $\Phi$  on M extends to a  $C^{\infty}$ - map  $\overline{\Phi}$  on  $\overline{M}$ , where  $|\alpha|$  is the norm of the second fundamental form of M.

It is interesting that these results have appropriate versions in the nonminimal setting. B. White [8], proved that a complete 2-dimensional surface M immersed in  $\mathbb{R}^n$  with  $\int_M |\alpha|^2 dV < \infty$  and nonpositive curvature with respect to every normal direction<sup>1</sup> is homeomorphic to a compact Riemann surface  $\overline{M}$  punctured at finite number of points  $\{p_1, \ldots, p_r\}$ , its Gauss map  $\Phi$  extends continuously to all of  $\overline{M}$  and M is properly immersed. It should be observed that the properness of M in White's Theorem is consequence of from the first two statements about the immersion, i.e. Jorge and Meeks [4], proved that a complete *m*-dimensional immersed submanifold M of  $\mathbb{R}^n$ , homeomorphic to a compact Riemann manifold  $\overline{M}$  punctured at finite number of points  $\{\underline{p}_1, \ldots, p_r\}$  and such that the Gauss map  $\Phi$  extends continuously to all of  $\overline{M}$  is properly immersed.

S. Muller and V. Sverak [6], answering a question of White, proved that a complete 2-dimensional surface M immersed in  $\mathbb{R}^n$  with  $\int_M |\alpha|^2 dV < \infty$ is properly immersed. For higher dimension submanifolds, Do Carmo and Elbert [3] proved that a complete hypersurface M of  $\mathbb{R}^{n+1}$  with strong finite total scalar curvature,  $(\int_M (|\alpha|^{n-1} + |\nabla \alpha|^{n-2}) dV < \infty)$  is proper and it is  $C^{\infty}$ - diffeomorphic to a compact smooth manifold  $\overline{M}$  punctured at a finite number of points  $\{p_1, \ldots, p_r\}$ . Moreover, they showed that if the Gauss-Kronecker curvature does not change sign in a punctured neighborhood of each  $p_i$ ,  $i = 1, \ldots, r$  then the Gauss map  $\Phi$  on M extends continuously to all of  $\overline{M}$ .

The purpose of this paper is to put another piece on this puzzle showing that a complete *m*-dimensional submanifold of  $\mathbb{R}^n$  with the norm of the second fundamental form uniformly decaying to zero  $|\alpha(x)| \to 0$  as  $x \to \infty$ in a certain rate is proper and has finite topological type. The decaying rate of  $|\alpha(x)| \to 0$  considered is not fast enough to make  $\int_M |\alpha(x)|^m dV < \infty$ .

To be more precise, let M be a complete m-dimensional submanifold of  $\mathbb{R}^n$  and let  $K_1 \subset K_2 \subset \ldots$  be an exhaustion sequence of M by compact sets. Fix a point  $p \in K_1$  and set  $a_i = \sup\{\rho(x) \cdot |\alpha(x)|, x \in M \setminus K_i\}$ , where  $\rho(x) = \operatorname{dist}_M(p, x)$  and  $|\alpha(x)|$  is the norm of the second fundamental form of M at x. The  $a_i$ 's form a non-increasing sequence  $\infty \geq a_1 \geq a_2 \geq \cdots \geq 0$  with  $a_1 = \infty$  iff  $a_l = \infty$  for all  $l \geq 1$ . Define the (possibly extended) scaling invariant number  $a(M) = \lim_{i \to \infty} a_i \in [0, \infty]$ . It can be shown that a(M) does not depend on the exhaustion sequence nor on the point p. It follows from the work of Jorge-Meeks [4] that complete m-dimensional Riemannian submanifolds M of  $\mathbb{R}^n$ homeomorphic to a compact Riemann manifold  $\overline{M}$  punctured at finite number of points  $\{p_1, \ldots, p_r\}$  and well defined normal vector at infinity have a(M) = 0. In particular, complete minimal surfaces in  $\mathbb{R}^n$  with finite total curvature, complete 2-dimensional complete surfaces

 $<sup>{}^{1}</sup>M \subset \mathbb{R}^{n}$  is nonpositively curved with respect to each normal direction at x if det $(\eta \cdot \alpha(, )) \leq 0$  for all normals  $\eta$  to M at x, see [8].

with  $\int_M |\alpha|^2 dV < \infty$  and nonpositive curvature with respect to every normal direction considered by White or the *m*-dimensional minimal submanifolds M of  $\mathbb{R}^n$  with finite total scalar curvature considered by Anderson have a(M) = 0. In our main result, we prove that the larger class of complete *m*-dimensional immersed submanifolds of  $\mathbb{R}^n$  with a(M) < 1 share some properties with these submanifolds with a(M) = 0. We prove the following theorem.

#### theorem1

**Theorem 1.1** Let M be a complete m-dimensional submanifold of  $\mathbb{R}^n$  with a(M) < 1. Then M is properly immersed and it is  $C^{\infty}$ - diffeomorphic to a compact smooth manifold  $\overline{M}$  with boundary.

Observe that  $\int_M |\alpha|^m dV < \infty$  is not equivalent to a(M) < 1. However, one might ask if Theorem (1.1) holds under finite total scalar curvature  $\int_M |\alpha|^m dV < \infty$ .

For complete *m*-dimensional minimal submanifolds M of  $\mathbb{R}^n$  we define the increasing sequence  $b_i = \inf\{\rho^2(x) \cdot Ric(x)(\nu,\nu), |\nu| = 1, x \in M \setminus K_i\}$ with  $b_1 = -\infty$  iff  $b_l = -\infty$  for all  $l \ge 1$ . Define the scaling invariant number  $b(M) = \lim_{i\to\infty} b_i \in [-\infty, 0]$ . Again, it can be shown that b(M)does not depend on the exhaustion sequence nor on the point p. The proof of Theorem (1.1) can be slightly modified to prove the following version for minimal submanifolds.

theorem2 Theorem 1.2 Let M be a complete m-dimensional minimal submanifold of  $\mathbb{R}^n$  with b(M) > -1. Then M is properly immersed and it is  $C^{\infty}$ - diffeomorphic to a compact smooth manifold  $\overline{M}$  with boundary.

### 2 Proof of Theorem 1.1

#### 2.1 *M* is properly immersed

Let  $\varphi : M^m \hookrightarrow \mathbb{R}^n$  be a complete submanifold with a(M) < 1 and let  $p \in M$ be a fixed point such that  $\varphi(p) = 0 \in \mathbb{R}^n$ . There exist a geodesic ball  $B_M(p, R_0)$  centered at p with radius  $R_0$  such that for all  $x \in M \setminus B_M(p, R_0)$ we have that  $\rho(x)|\alpha(x)| \leq c < 1$ . Let  $f: M^m \to \mathbb{R}$  given by  $f(x) = |\varphi(x)|^2$ . Fix a point  $x \in M \setminus B_M(p, R_0)$  then for  $\nu \in T_x M$ ,  $|\nu| = 1$  we have that

$$\frac{1}{2} \operatorname{Hess} f(x)(\nu, \nu) = 1 + \langle \varphi(x), \, \alpha(x)(\nu, \nu) \rangle$$

$$\geq 1 - |\varphi(x)| \cdot |\alpha(x)| \qquad (1) \quad \text{eqHess}$$

$$\geq 1 - \rho(x)|\alpha(x)|$$

$$\geq 1 - c$$

Let  $\sigma : [0, \rho(x)] \to M^m$  be a minimal geodesic from p to x. We have from (1) that for all  $t \ge R_0$  that  $(f \circ \sigma)''(t)$  = Hess  $f(\sigma(t))(\sigma', \sigma') \ge 2(1-c)$  and for  $t < R_0$  that  $(f \circ \sigma)''(t) \ge b$ ,  $b = \inf_{x \in B_M(p,R_0)} \{ \text{Hess } f(x)(\nu,\nu), |\nu| = 1 \}$ . Thus

$$(f \circ \sigma)'(s) = \int_0^s (f \circ \sigma)''(\tau) d\tau$$
  

$$\geq \int_0^{R_0} b \, d\tau + \int_{R_0}^s (1-c) d\tau \qquad (2) \quad \text{eq2}$$
  

$$\geq b R_0 + (1-c)(s-R_0)$$

$$f(x) = \int_{0}^{\rho(x)} (f \circ \sigma)'(s) \, ds$$
  

$$\geq \int_{0}^{\rho(x)} b \, R_0 + (1 - c)(s - R_0) \, ds \qquad (3) \quad \text{eq3}$$
  

$$= b \, R_0 \, \rho(x) + (1 - c) \left(\frac{\rho(x)^2}{2} - R_0 \rho(x)\right)$$

Thus  $\varphi(x)|^2 \ge (b-1+c)R_0\rho(x) + (1-c)\rho(x)^2/2$  for all  $x \in M \setminus B_M(p, R)$ . In fact, this proof that f is proper proves that following proposition.

**PropositionProper Proposition 2.1** Let  $f: M \to \mathbb{R}$  be a  $C^2$ -function defined on a complete Riemannian manifold such that Hess  $f(x) \ge g(\rho(x))$ , where  $\rho$  is the distance function to  $x_o$  and  $g: [0, \infty) \to \mathbb{R}$  is a piecewise continuous function. Setting  $G(t) = f(x_o) - |\text{grad } f(x_o)| t + \int_0^t \int_0^s g(u) duds, t \in [0, \infty)$  we have that if G is proper and bounded from below then f is proper.

#### 2.2 *M* has finite topology

Let  $\varphi : M^m \hookrightarrow \mathbb{R}^n$  be a complete properly immersed submanifold with a(M) < 1. Let  $p \in M$  be such that  $\varphi(p) = 0 \in \mathbb{R}^n$  and suppose without

loss of generality that  $R(x) = |\varphi(x)|, x \in M$  is a Morse function. Let r > 0 be such that  $\Gamma_r = \varphi(M) \cap \mathbb{S}^{n-1}(r)$  is a compact submanifold of  $\mathbb{S}^{n-1}(r)$  and  $\rho(x) \cdot |\alpha(x)| \leq c < 1$  for all  $x \in M \setminus \varphi^{-1}(B_{\mathbb{R}^n}(r))$ . For  $y \in \Gamma_r$ , let  $\eta(y) = y/r$ , the unit vector perpendicular to  $T_y \mathbb{S}^{n-1}(r)$ . There is only one unit vector  $\nu(y) \in T_y \varphi(M)$  such that  $T_y \varphi(M) = T_y \Gamma_r \oplus [[\nu(y)]]$  and  $\langle \nu(y), \eta(y) \rangle > 0$ . Here  $T_y \varphi(M), T_y \mathbb{S}^{n-1}(r)$  are the tangent spaces of  $\varphi(M)$  and  $\mathbb{S}^{n-1}(r)$  at y respectively and  $[[\nu(y)]]$  is the vector space generated by  $\nu(y)$ . This procedure defines a smooth vector field  $\nu(y)$  on the neighborhood  $\varphi^{-1}(B_{\mathbb{R}^n}(r+\epsilon) \setminus B_{\mathbb{R}^n}(r-\delta))$  of  $\varphi^{-1}(\Gamma_r)$  for some  $\epsilon, \delta > 0$ . Consider the function  $\psi$  defined on that neighborhood given by

$$\psi(y) = \langle \nu(y), \eta(y) \rangle = \cos \theta(y)$$

Identify  $X \in TM$  with  $d\varphi X$ . We have that  $X(R) = \langle X, \eta \rangle$  and writing  $\eta(y) = \sin \theta(y) \nu^*(y) + \cos \theta(y) \nu(y), \nu^*(y) \perp \nu(y)$ , we have that grad  $R = \psi \nu$ . Now for each  $y \in \Gamma_r$  consider  $\xi(t, y)$  the solution of the following problem on  $\varphi(M)$ 

$$\xi_t = \frac{1}{\psi} \nu(\xi)$$

$$(4) \quad eqData$$

$$\xi(0, y) = y$$

We have for  $R(t, y) = |\xi(t, y)|$  that

$$R_t = \langle \operatorname{grad} R, \frac{1}{\psi} \nu \rangle = \langle \psi \nu, \frac{1}{\psi} \nu \rangle = 1 \Leftrightarrow R = R(t, y) = t + r$$

We will derive a differential equation that the function  $\psi \circ \xi(t, y)$  satisfies.

$$\psi_t = \xi_t \langle \nu, \eta \rangle = \langle D_{\frac{1}{\psi}\nu}\nu, \eta \rangle + \langle \nu, D_{\xi_t}\eta \rangle = \langle \nabla_\nu \nu + \alpha(\nu, \nu), \eta \rangle + \langle \nu, D_{\xi_t}(\frac{\xi}{R}) \rangle$$

But  $\langle \nu, \nu \rangle = 1 \Rightarrow \langle \nu, \nabla_{\nu} \nu \rangle = 0$  and  $\nabla_{\nu} \nu \in T_x M \Rightarrow \nabla_{\nu} \nu \in (T_x M \cap T_x S_R^n)$  $\Rightarrow \langle \nabla_{\nu} \nu, \eta \rangle = 0$ . On other hand

$$D_{\xi_t}(\frac{\xi}{R}) = \frac{\frac{1}{\psi}\nu}{R} - \frac{R_t}{R^2}\varphi = \frac{1}{R\psi}\nu - \frac{1}{R}\eta$$

then

$$\psi_t = \frac{1}{\psi} \langle \alpha(\nu, \nu), \eta \rangle + \frac{1}{\psi R} - \frac{\psi}{R} = \frac{\sqrt{1 - \psi^2}}{\psi} \langle \alpha(\nu, \nu), \nu^* \rangle + \frac{1 - \psi^2}{\psi R} \quad (5) \quad \text{eqdf}$$

To determine a differential equation satisfied by  $\sin \theta(t, y) = \sqrt{1 - \psi^2}$  we proceed as follows. By (5) we have

$$\frac{\psi\psi_t}{\sqrt{1-\psi^2}} = \langle \alpha(\nu,\nu),\nu^* \rangle + \frac{\sqrt{1-\psi^2}}{R}$$
(6) eqdf2

Observing that R(t, y) = t + r, equation (6) can be written as

$$-(t+r)(\sqrt{1-\psi^2})_t = (t+r)\langle \alpha(\nu,\nu),\nu^* \rangle + \sqrt{1-\psi^2}$$
(7) eqdf3

and rewritten as

$$\left[(t+r)\sqrt{1-\psi^2}\right]_t + (t+r)\langle\alpha(\nu,\nu),\nu^*\rangle = 0 \tag{8} \quad \text{eqdf4}$$

Integrating the equation (8) we have the following equation

$$\sqrt{1-\psi^2} = \frac{r}{t+r}\sqrt{1-\psi_o^2} - \frac{1}{t+r}\int_0^t (s+r)\langle \alpha(\nu,\nu),\nu^* \rangle \, ds \qquad (9) \quad \text{eqdf5}$$

where  $\psi_o = \psi(\xi(0, y))$ . Since  $\sin \theta(\xi(t, y)) = \sqrt{1 - \psi^2}$  we rewrite (9) in the following form

$$\sin\theta(\xi(t,y)) = \frac{r}{t+r}\sin\theta(\xi(0,y)) - \frac{1}{t+r}\int_0^t (s+r)\langle\alpha(\nu,\nu),\nu^*\rangle\,ds \quad (10) \quad \text{eqdf6}$$

Now,

$$-\langle \alpha(\nu,\nu),\nu^*\rangle(\xi(s,y)) \le |\alpha|(\xi(s,y)) \le c/\rho(\xi(s,y)) \le c/R(s,y).$$

Substituting in (10) and noticing that R(s, y) = s + r we have that

$$\sin \theta(\xi(t,y)) \leq \frac{r}{t+r} \sin \theta(\xi(0,y)) + \frac{1}{t+r} \int_0^t (s+r) \frac{c}{s+r} ds$$
$$= \frac{ct+r \sin \theta(\xi(0,y))}{t+r} < 1, \ \forall t \ge 0.$$
(11) eqdf7

Recall that the critical points of R are those x such that  $\psi(x) = 0$ , or those points where  $\sin \theta(x) = 1$ . Thus, along the integral curves  $\xi(t, y), y \in \Gamma_r$ there is no critical point for the function  $R(x) = |\varphi(x)|$ . This shows that outside the compact set  $M \setminus B_M(p, r)$  there are no critical points for R. Since R is a Morse function, its critical points are isolated thus there are finitely many of them. This shows that M has finite topology.

## 3 Sketch of proof for Theorem 1.2

Let  $\varphi : M^m \hookrightarrow \mathbb{R}^n$  be a complete minimal submanifold and let  $x \in M$ ,  $\nu \in T_x M$  and  $\{e_1, \ldots e_m = \nu\}$  an orthonormal basis for  $T_x M$ . Using the Gauss equation we can compute the Ricci curvature in the direction  $\nu$  by

$$\begin{aligned} Ric(x)(\nu) &= \langle \sum_{i=1}^{m} \alpha_{ii}, \alpha_{mm} \rangle - \sum_{i=1}^{m-1} |\alpha_{im}|^2 \\ &= \langle mH - \alpha_{mm}, \alpha_{mm} \rangle - \sum_{i=1}^{m-1} |\alpha_{im}|^2 \end{aligned} \tag{12} \quad eqRic \\ &= -\sum_{i=1}^{m} |\alpha_{im}|^2, \end{aligned}$$

where  $\alpha_{ij} = \alpha(e_i, e_j)$ . Let  $f : M^m \to \mathbb{R}$  given by  $f(x) = |\varphi(x)|^2$ . The Hessian of f at  $x \in M$  and  $\nu \in T_x M$ ,  $|\nu| = 1$  satisfies

$$\frac{1}{2} \operatorname{Hess} f(x)(\nu, \nu) = + \langle \varphi(x), \alpha(\nu, \nu) \rangle \\
\geq [1 - |\varphi(x)| \cdot |\alpha_{mm}|] \\
\geq [1 - |\varphi(x)| \sqrt{-Ric(x)(\nu)}] \\
\geq [1 - \rho(x) \sqrt{-Ric(x)(\nu)}] \\
\geq 1 - c$$
(13)

The proof of Theorem (1.1) from equation (1) on shows that M is is properly immersed in Theorem (1.2). To show that M has finite topological type observe that

$$|\alpha|(\xi(s,y)) \le \sqrt{-Ric(\xi(s,y))(\nu,\nu)} \le c/\rho(\xi(s,y)) \le c/R(s,y)$$

after (10) and we still have (11).

## References

- M. Anderson: The compactification of a minimal subamnifold by the Gauss Map. Preprint IEHS (1985).
- [2] S. S. Chern & R. Osserman: Complete minimal surfaces in euclidean n-space. J. d'Analyse Math. vol. 19, 15-34, (1967).

- [3] M. P. do Carmo & M. F. Elbert: Complete hypersurfaces in Euclidean spaces with strong finite total curvature. Preprint.
- [4] Luquésio P. Jorge & W. Meeks III: The topology of complete minimal surfaces of finite total Gaussian curvature. Topology, vol. 22 (2), 203-221, (1983).
- [5] H. B. Lawson: Lectures on minimal submanifolds. Monografias de Matemática, IMPA, Rio de Janeiro, Brazil.
- S. Muller & V. Sverak: On surfaces of finite total curvature.J. Diff. Geometry, 42, 229-258 (1995).
- [7] R. Osserman: Global properties of minimal surfaces in E<sup>3</sup> and E<sup>n</sup>. Ann. of Math. vol. 80, 340-364, (1964).
- [8] B. White: Complete surfaces of finite total curvature J. Diff. Geometry, 26, 315-326 (1987)