

# Schur-Weyl reciprocity for the $q$ -analogue of the alternating group

Hideo Mitsuhashi

Department of Information Technology

Kanagawa Prefectural Junior College for Industrial Technology

2-4-1 Nakao, Asahi-ku, Yokohama-shi, Kanagawa-ken 241-0815, Japan

## Abstract

In this paper, we establish Schur-Weyl reciprocity for the  $q$ -analogue of the alternating group. We analyze the sign  $q$ -permutation representation of the Hecke algebra  $\mathcal{H}_{\mathbb{Q}(q),r}(q)$  on the  $r$ th tensor product of  $\mathbb{Z}_2$ -graded  $\mathbb{Q}$ -vector space  $V = V_{\overline{0}} \otimes V_{\overline{1}}$  in detail, and examine its restriction to the  $q$ -analogue of the alternating group  $\mathcal{H}_{\mathbb{Q}(q),r}^1(q)$ . In consequence, we find out that if  $\dim V_{\overline{0}} = \dim V_{\overline{1}}$ , then the centralizer of  $\mathcal{H}_{\mathbb{Q}(q),r}^1(q)$  is a  $\mathbb{Z}_2$ -crossed product of the centralizer of  $\mathcal{H}_{\mathbb{Q}(q),r}(q)$  and obtain Schur-Weyl reciprocity between  $\mathcal{H}_{\mathbb{Q}(q),r}^1(q)$  and its centralizer. Though the structure of the centralizer is more complicated for the case  $\dim V_{\overline{0}} \neq \dim V_{\overline{1}}$ , we obtain some results about the case. When  $q = 1$ , Regev has proved Schur-Weyl reciprocity for alternating groups in [12]. Therefore, our result can be regarded as an extension of Regev's work.

## 1 Introduction

The purpose of this study is to research Schur-Weyl reciprocity for the  $q$ -analogue of the alternating group. In our previous paper [10], we established Schur-Weyl reciprocity between the Hecke algebra  $\mathcal{H}_{\mathbb{Q}(q),r}(q)$  and the quantum super Lie algebra  $U_q^\sigma(\mathfrak{gl}(m, n))$ . In that paper, we defined the  $q$ -permutation representation of  $\mathcal{H}_{\mathbb{Q}(q),r}(q)$ , and showed that the image of the  $q$ -permutation representation is the centralizer of the image of the vector representation of the quantum super Lie algebra  $U_q^\sigma(\mathfrak{gl}(m, n))$  on the  $r$ th tensor product of a  $\mathbb{Z}_2$ -graded  $(m+n)$ -dimensional  $\mathbb{Q}(q)$ -vector space  $V = V_{\overline{0}} \oplus V_{\overline{1}}$ . In this paper, we find out the centralizer of the  $q$ -analogue of the alternating group as the restriction of the  $q$ -permutation representation. When  $q = 1$ , Regev has already shown Schur-Weyl reciprocity for the alternating group in [12]. Hence our result is regarded as an extension of Regev's work.

Let  $q$  be an indeterminate and  $K = \mathbb{Q}(q)$ . Let  $(\pi_r, V^{\otimes r})$  be the  $q$ -permutation representation of  $\mathcal{H}_{K,r}(q)$  (definition of the  $q$ -permutation representation is at (4.1)) and  $(\rho_r, V^{\otimes r})$  the vector representation of  $U_q^\sigma(\mathfrak{gl}(m, n))$  (definition of the vector representation is at (4.4)). We have proved in [10] that  $\mathcal{A}_q = \pi_r(\mathcal{H}_{K,r}(q))$  and  $\mathcal{B}_q = \rho_r(U_q^\sigma(\mathfrak{gl}(m, n)))$  are full centralizers of each other, namely:

$$\mathcal{B}_q = \text{End}_{\mathcal{A}_q} V^{\otimes r} \quad \text{and} \quad \mathcal{A}_q = \text{End}_{\mathcal{B}_q} V^{\otimes r}. \quad (1.1)$$

Let  $R_0$  be a commutative domain which includes an invertible element  $q$ . We further assume that 2 and  $q + q^{-1}$  are invertible elements of  $R_0$ . Then we can define the  $q$ -analogue of the alternating group  $\mathcal{H}_{R_0,r}^1(q)$  (see Definition 3.1 and Proposition 3.6) in  $\mathcal{H}_{R_0,r}(q)$ . In [9], we defined the  $q$ -analogue of the alternating group as a subalgebra of Iwahori-Hecke algebra of type  $A$  and obtained defining relations (see Proposition 3.7) for the first time. In this paper, we show that  $\mathcal{H}_{R_0,r}(q)$  is isomorphic to the  $\mathbb{Z}_2$ -crossed product which is obtained from the crossed system  $(\mathcal{H}_{R_0,r}^1(q), \mathbb{Z}_2, \psi_0, \alpha_0)$  (definition of  $\psi_0$  and  $\alpha_0$  are at (3.2) and (3.3) respectively).

**Theorem 3.9.**  *$\mathcal{H}_{R_0,r}(q)$  is isomorphic to the  $\mathbb{Z}_2$ -crossed product  $\mathcal{H}_{R_0,r}^1(q)_{\alpha_0}^{\psi_0}[\mathbb{Z}_2]$  as  $R_0$ -algebras.*

Let  $\mathcal{C}_q = \pi_r(\mathcal{H}_{K,r}^1(q))$  and  $\mathcal{D}_q = \text{End}_{\mathcal{C}_q} V^{\otimes r}$ . Our main subject is to solve the relation between  $\mathcal{D}_q$  and  $\mathcal{B}_q$ . From (1.1), one can immediately see that  $\mathcal{B}_q \subseteq \mathcal{D}_q$ . But the structure of  $\mathcal{D}_q$  is not trivial. Indeed, the structure of  $\mathcal{D}_q$  depends on the dimensions  $m = \dim_K V_{\bar{0}}$  and  $n = \dim_K V_{\bar{1}}$ . In this paper, we show that if  $m = n$ , then  $\mathcal{D}_q$  is isomorphic to  $\mathbb{Z}_2$ -crossed product which is obtained from the crossed system  $(\mathcal{B}_q, \mathbb{Z}_2, \psi_1, \alpha_1)$  (definition of  $\psi_0$  and  $\alpha_0$  are at (5.4) and (5.5) respectively).

**Theorem 5.6.** *If  $m = n$ , then  $\mathcal{D}_q$  is isomorphic to the  $\mathbb{Z}_2$ -crossed product  $\mathcal{B}_{q\alpha_1}^{\psi_1}[\mathbb{Z}_2]$  as  $K$ -algebras.*

From this theorem, we immediately obtain that  $\dim_K \mathcal{D}_q = 2 \dim_K \mathcal{B}_q$ . Moreover, we show Schur-Weyl reciprocity for  $\mathcal{H}_{K,r}^1(q)$ .

**Theorem 5.8.**  *$\text{End}_{\mathcal{C}_q} V^{\otimes r} = \mathcal{D}_q$  and  $\text{End}_{\mathcal{D}_q} V^{\otimes r} = \mathcal{C}_q$  hold.*

In the general case, the matter is more complicated, but we can find out to some extent if we exchange the base field from  $K$  to its algebraic closure  $\bar{K}$ . Let  $\bar{U}_q^\sigma(\mathfrak{gl}(m, n)) = U_q^\sigma(\mathfrak{gl}(m, n)) \otimes_K \bar{K}$ ,  $\bar{\mathcal{A}}_q = \mathcal{A}_q \otimes_K \bar{K}$ ,  $\bar{\mathcal{B}}_q = \mathcal{B}_q \otimes_K \bar{K}$  and  $\bar{\mathcal{C}}_q = \mathcal{C}_q \otimes_K \bar{K}$ . Then we have the following theorem by the theory of semisimple algebras.

**Theorem 6.1.**  *$\bar{\mathcal{A}}_q$  and  $\bar{\mathcal{C}}_q$  have direct sum decompositions  $\bar{\mathcal{A}}_q = \bar{\mathcal{A}}_q^0 \oplus \bar{\mathcal{A}}_q^1$  and  $\bar{\mathcal{C}}_q = \bar{\mathcal{C}}_q^0 \oplus \bar{\mathcal{C}}_q^1$  respectively, which are satisfy the following relations.*

$$(1) \quad \bar{\mathcal{C}}_q^0 \subseteq \bar{\mathcal{A}}_q^0 \quad \text{and} \quad \dim_{\bar{K}} \bar{\mathcal{A}}_q^0 = 2 \dim_{\bar{K}} \bar{\mathcal{C}}_q^0$$

(2)  $\bar{\mathcal{C}}_q^1 = \bar{\mathcal{A}}_q^1$ . Especially  $\dim_{\bar{K}} \bar{\mathcal{A}}_q^1 = \dim_{\bar{K}} \bar{\mathcal{C}}_q^1$ .

As a corollary, we obtain an anomalous phenomenon for non-super case as follows.

**Corollary 6.2.** *Let  $n = 0$ . If  $m^2 < r$ , then  $\bar{\mathcal{A}}_q = \bar{\mathcal{C}}_q$  and  $\text{End}_{\bar{\mathcal{C}}_q} \bar{V}^{\otimes r} = \text{End}_{\bar{\mathcal{A}}_q} \bar{V}^{\otimes r}$ .*

The details of the decompositions of  $\bar{\mathcal{A}}_q$  and  $\bar{\mathcal{C}}_q$  are described in section 6. We also obtain the similar result to Theorem 6.1 about the endomorphism algebras  $\text{End}_{\bar{\mathcal{A}}_q} \bar{V}^{\otimes r}$  and  $\text{End}_{\bar{\mathcal{C}}_q} \bar{V}^{\otimes r}$ ; there exist two  $\mathcal{H}_{\bar{K},r}(q)$ -submodules  $W_0$  and  $W_1$  which satisfy the following properties.

**Corollary 6.3.**  *$\text{End}_{\bar{\mathcal{C}}_q} W_0 \supseteq \text{End}_{\bar{\mathcal{A}}_q} W_0$  and  $\text{End}_{\bar{\mathcal{C}}_q} W_1 = \text{End}_{\bar{\mathcal{A}}_q} W_1$ .*

Although the relation between  $\bar{\mathcal{A}}_q$  and  $\bar{\mathcal{C}}_q$  is made clear by (6.4)-(6.7), that between  $\text{End}_{\bar{\mathcal{A}}_q} W$  and  $\text{End}_{\bar{\mathcal{C}}_q} W$  is not clear except for Corollary 6.2 at this point.

## 2 Preliminaries

Let  $R$  be a commutative ring with 1 and  $G$  a group. In this section, we shall review the definition and some properties about  $G$ -crossed products. A full account about  $G$ -graded algebras and  $G$ -crossed products is given in [11].

**Definition 2.1 ( $G$ -graded algebra).** An  $R$ -algebra  $A$  is said to be  $G$ -graded if there exist a family of  $R$ -submodules  $\{A_\sigma | \sigma \in G\}$  of  $A$  indexed by elements of  $G$  which satisfies the following two conditions:

$$(G1) \quad A = \bigoplus_{\sigma \in G} A_\sigma,$$

$$(G2) \quad A_\sigma A_\tau \subseteq A_{\sigma\tau} \text{ for } \sigma, \tau \in G.$$

Moreover,  $A$  is said to be *strongly  $G$ -graded* when (G2) is replaced by the following condition:

$$(G'2) \quad A_\sigma A_\tau = A_{\sigma\tau} \text{ for } \sigma, \tau \in G.$$

We notice that if  $A = \bigoplus_{\sigma \in G} A_\sigma$  is a  $G$ -graded algebra, then  $A_{1_G}$  ( $1_G$  means the identity element of  $G$ ) is a subalgebra of  $A$  and  $1_A \in A_{1_G}$ .

**Definition 2.2 ( $G$ -crossed product).** A  $G$ -graded  $R$ -algebra  $A = \bigoplus_{\sigma \in G} A_\sigma$  is said to be a  $G$ -crossed product if each  $A_\sigma$  has an invertible element.

We notice that a  $G$ -crossed product is a strongly  $G$ -graded algebra. Indeed, if  $A$  is a  $G$ -crossed product, then for an invertible element  $u_\sigma \in A_\sigma$ ,  $u_\sigma^{-1} \in A_{\sigma^{-1}}$  and  $1_A = u_\sigma u_\sigma^{-1} \in A_\sigma A_{\sigma^{-1}}$ . So, we have  $A_{\sigma\tau} = 1_A A_{\sigma\tau} \subseteq (A_\sigma A_{\sigma^{-1}}) A_{\sigma\tau} = A_\sigma (A_{\sigma^{-1}} A_{\sigma\tau}) \subseteq A_\sigma A_\tau$ .

**Definition 2.3 (crossed system).** Let  $A$  be an algebra and  $G$  a group. Suppose that there exist two maps

$$\psi : G \longrightarrow \text{Aut}(A),$$

(we denote  $\psi(\sigma)(a)$  by  ${}^\sigma a$  for brevity), and

$$\alpha : G \times G \longrightarrow A^\times,$$

where  $A^\times$  is the multiplicative group of units of  $A$ , which satisfy the relations:

$${}^\sigma({}^\tau a) = \alpha(\sigma, \tau)^{(\sigma\tau)} a \alpha(\sigma, \tau)^{-1} \quad (2.1)$$

$${}^{\sigma_1} \alpha(\sigma_2, \sigma_3) \alpha(\sigma_1, \sigma_2 \sigma_3) = \alpha(\sigma_1, \sigma_2) \alpha(\sigma_1 \sigma_2, \sigma_3) \quad (2.2)$$

$$\alpha(\sigma, 1) = \alpha(1, \sigma) = 1, \quad (2.3)$$

for  $\sigma, \tau, \sigma_1, \sigma_2, \sigma_3 \in G, a \in A$ .  $(A, G, \psi, \alpha)$  is said to be a *crossed system*.  $\psi$  is called a *weak action* of  $G$  on  $A$ , and  $\alpha$  is called a  *$\psi$ -cocycle*.

We denote by  $A_\alpha^\psi[G]$  the free left  $A$ -module with the basis  $\{u_\sigma | \sigma \in G\}$  and the following multiplication:

$$(a_1 u_\sigma)(a_2 u_\tau) = a_1 {}^\sigma a_2 \alpha(\sigma, \tau) u_{\sigma\tau}, \quad (2.4)$$

for  $a_1, a_2 \in A, \sigma, \tau \in G$ .

**Proposition 2.4 ([11]Proposition 1.4.1).**  $A_\alpha^\psi[G]$  is a  $G$ -crossed product.

**Proposition 2.5 ([11]Proposition 1.4.2).** Every  $G$ -crossed product is of the form  $A_\alpha^\psi[G]$  for some algebra  $A$ , some weak action  $\psi$  and some  $\psi$ -cocycle  $\alpha$ .

When  $G$  is finite and a strongly  $G$ -graded algebra  $A = \bigoplus_{\sigma \in G} A_\sigma$  is finitely generated over  $R$  as modules,  $A$  is said to have a  *$G$ -graded Clifford system*  $\{A_\sigma | \sigma \in G\}$  if  $A$  satisfies (C1).

(C1) For each  $\sigma \in G$ , there exists an invertible element  $a_\sigma \in A$  such that  $A_\sigma = a_\sigma A_{1_G} = A_{1_G} a_\sigma$ .

It is clear that such  $a_\sigma$  is in  $A_\sigma$ . An exposition about group graded Clifford systems can be found in [1], section 11C.

### 3 The $q$ -analogue of the alternating group and its representation

Let  $(W, S = \{s_1, \dots, s_r\})$  be a Coxeter system of rank  $r$ . Let  $R_0$  be a commutative domain with 1, and let  $q_i (i = 1, \dots, r)$  be any invertible elements of  $R_0$  such that  $q_i = q_j$  if  $s_i$  is conjugate to  $s_j$  in  $W$ . Further

we assume that 2 and  $q_i + q_i^{-1}$  ( $i = 1, 2, \dots, r$ ) are invertible elements of  $R_0$ . The Iwahori-Hecke algebra  $\mathcal{H}_{R_0}(W, S)$  is an  $R_0$ -algebra generated by  $\{T_{s_i} | s_i \in S\}$  with the defining relations:

$$(H1) \quad T_{s_i}^2 = (q_i - q_i^{-1})T_{s_i} + 1 \quad \text{if } i = 1, 2, \dots, r,$$

$$(H2) \quad (T_{s_i}T_{s_j})^{k_{ij}} = (T_{s_j}T_{s_i})^{k_{ij}} \quad \text{if } m_{ij} = 2k_{ij},$$

$$(H3) \quad (T_{s_i}T_{s_j})^{k_{ij}}T_{s_i} = (T_{s_j}T_{s_i})^{k_{ij}}T_{s_j} \quad \text{if } m_{ij} = 2k_{ij} + 1,$$

where  $m_{ij}$  is the order of  $s_i s_j$  in  $W$ . We write  $T_i = T_{s_i}$  for brevity.

If  $(W, S)$  is of type  $A$  and of rank  $r-1$ , then  $W$  is isomorphic to the symmetric group  $\mathfrak{S}_r$ . Furthermore, all the elements of  $S$  are conjugate to each other, hence we may assume  $q_1 = \dots = q_{r-1} = q$ . The Iwahori-Hecke algebra  $\mathcal{H}_{R_0, r}(q) = \mathcal{H}_{R_0}(W, S)$  of type  $A$  has the defining relations:

$$(A1) \quad T_i^2 = (q - q^{-1})T_i + 1 \quad \text{if } i = 1, 2, \dots, r-1,$$

$$(A2) \quad T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1} \quad \text{if } i = 1, 2, \dots, r-2,$$

$$(A3) \quad T_i T_j = T_j T_i \quad \text{if } |i - j| > 1.$$

Let  $\hat{\cdot}$  be the Goldman involution. This is an involution on  $\mathcal{H}_{R_0, r}(q)$  defined by

$$\hat{T}_i = (q - q^{-1}) - T_i.$$

**Definition 3.1.** We define  $\mathcal{H}_{R_0, r}^{\pm 1}(q)$  to be the eigenspaces of  $\mathcal{H}_{R_0, r}(q)$  corresponding to the eigenvalues  $\pm 1$  of  $\hat{\cdot}$  respectively.

We notice that  $\mathcal{H}_{R_0, r}^1(q)$  is a subalgebra of  $\mathcal{H}_{R_0, r}(q)$ . Let  $T'_i$  ( $i = 1, 2, \dots, r-1$ ) be the elements of  $\mathcal{H}_{R_0, r}(q)$  defined by

$$T'_i = \frac{T_i - \hat{T}_i}{q + q^{-1}} = \frac{2T_i - (q - q^{-1})}{q + q^{-1}} \quad \text{for } i = 1, 2, \dots, r-1.$$

Then one can immediately check  $\hat{T}'_i = -T'_i$ .

**Proposition 3.2.**  $T'_i$  ( $i = 1, 2, \dots, r$ ) generate  $\mathcal{H}_{R_0, r}(q)$  and satisfy the following defining relations:

$$(A'1) \quad T_i'^2 = 1 \quad \text{if } i = 1, 2, \dots, r-1,$$

$$(A'2) \quad T'_i T'_{i+1} T'_i = T'_{i+1} T'_i T'_{i+1} - \left( \frac{q - q^{-1}}{q + q^{-1}} \right)^2 (T'_i - T'_{i+1}) \quad \text{if } i = 1, 2, \dots, r-2,$$

$$(A'3) \quad T'_i T'_j = T'_j T'_i \quad \text{if } |i - j| > 1.$$

*Proof.* From the equations

$$T_i = \frac{1}{2}\{(q + q^{-1})T'_i + (q - q^{-1})\} \quad \text{for } i = 1, 2, \dots, r-1,$$

which are obtained from the definition of  $T'_i$ , we see that  $T'_i (i = 1, 2, \dots, r)$  generate  $\mathcal{H}_{R_0, r}(q)$ . The defining relations are obtained from a direct computation.  $\square$

Consider the following sets of monomials:

$$\begin{aligned} \mathcal{C}_1 &= \{1, T_1\} \\ \mathcal{C}_2 &= \{1, T_2, T_2T_1\} \\ \mathcal{C}_3 &= \{1, T_3, T_3T_2, T_3T_2T_1\} \\ &\vdots \\ \mathcal{C}_{r-1} &= \{1, T_{r-1}, T_{r-1}T_{r-2}, \dots, T_{r-1}T_{r-2}\cdots T_1\} \end{aligned}$$

We shall say that  $M_1M_2\cdots M_{r-1}$  is a *monomial in  $T_i$ -normal form* in  $\mathcal{H}_{R_0, r}(q)$  if  $M_i \in \mathcal{C}_i$  for  $i = 1, 2, \dots, r-1$ . The following fact is well-known in the theory of the Iwahori-Hecke algebra.

**Proposition 3.3.**  $\text{rank}_{R_0} \mathcal{H}_{R_0, r}(q) = r!$  and

$$\mathcal{H}_{R_0, r}(q) = \bigoplus_{M_i \in \mathcal{C}_i} R_0 M_1 M_2 \cdots M_{r-1}$$

We derive from this fact that all monomials in  $T'_i$ -normal form also constitute a basis of  $\mathcal{H}_{R_0, r}(q)$ .

**Proposition 3.4.** Let  $\mathcal{C}'_i = \{1, T'_i, T'_i T'_{i-1}, \dots, T'_i T'_{i-1} \cdots T'_1\}$  for  $i = 1, 2, \dots, r-1$ . Then we have

$$\mathcal{H}_{R_0, r}(q) = \bigoplus_{M'_i \in \mathcal{C}'_i} R_0 M'_1 M'_2 \cdots M'_{r-1}.$$

*Proof.* Consider the map  $f$  from  $\{T_i\}_{i=1,2,\dots,r-1}$  to  $\{T'_i\}_{i=1,2,\dots,r-1}$  which is defined by  $f(T_i) = T'_i$ . This map induces the  $R_0$ -endomorphism  $\bar{f}$  of  $\mathcal{H}_{R_0, r}(q)$ .  $\bar{f}$  is an  $R_0$ -isomorphism because the inverse  $\bar{g}$  which is induced from the map  $g$  from  $\{T'_i\}_{i=1,2,\dots,r-1}$  to  $\{T_i\}_{i=1,2,\dots,r-1}$  defined by

$$g(T'_i) = \frac{1}{2}\{(q + q^{-1})T_i + (q - q^{-1})\}$$

exists. Hence we conclude that  $M'_1 M'_2 \cdots M'_{r-1} (M'_i \in \mathcal{C}'_i)$ , which are images of  $M_1 M_2 \cdots M_{r-1} (M_i \in \mathcal{C}_i)$ , are linearly independent and constitute a basis of  $\mathcal{H}_{R_0, r}(q)$ .  $\square$

Let  $\mathcal{E}_r$  (respectively  $\mathcal{O}_r$ ) be the set of all monomials in  $T'_i$ -normal form in  $\mathcal{H}_{R_0, r}(q)$  which are products of even (respectively odd) numbers of  $T'_i$ 's. Then the following holds.

**Lemma 3.5.**  $|\mathcal{E}_r| = |\mathcal{O}_r| = 2^{-1}r!$  for  $r > 1$ .

*Proof.* The proof is done by induction on  $r$ . It is trivial for  $r = 2$ . Let  $M'_1 M'_2 \cdots M'_{r-1} \in \mathcal{E}_r$  with  $r > 2$  and  $M'_i \in \mathcal{C}'_i$ . Then  $M'_1 M'_2 \cdots M'_{r-2}$  is considered as a monomial in  $T'_i$ -normal form in  $\mathcal{H}_{R_0, r-1}(q)$ . Let  $(\mathcal{C}'_{r-1})^e$  (respectively  $(\mathcal{C}'_{r-1})^o$ ) be the subset of  $\mathcal{C}'_{r-1}$  which consists of products of even (respectively odd) numbers of  $T'_i$ 's. We can readily see that  $M'_1 M'_2 \cdots M'_{r-2} \in \mathcal{E}_{r-1}$  if and only if  $M'_{r-1} \in (\mathcal{C}'_{r-1})^e$ . By induction,  $|\mathcal{E}_{r-1}| = |\mathcal{O}_{r-1}| = 2^{-1}(r-1)!$  and hence we obtain the following

$$|\mathcal{E}_r| = |(\mathcal{C}'_{r-1})^e| |\mathcal{E}_{r-1}| + |(\mathcal{C}'_{r-1})^o| |\mathcal{O}_{r-1}| = r \times 2^{-1}(r-1)! = 2^{-1}r!$$

as desired.  $\square$

Now we characterize  $\mathcal{H}_{R_0, r}^1(q)$  as a  $q$ -analogue of the alternating group.

**Proposition 3.6.**  $\text{rank}_{R_0} \mathcal{H}_{R_0, r}(q) = 2 \text{rank}_{R_0} \mathcal{H}_{R_0, r}^1(q)$ . Moreover  $\mathcal{H}_{R_0, r}^1(q)$  is the subalgebra which consists of all the products of even numbers of  $T'_i$ 's.

*Proof.* Let  $\mathcal{H}_{R_0, r}^e(q) = \oplus_{M \in \mathcal{E}_r} R_0 M$  and  $\mathcal{H}_{R_0, r}^o(q) = \oplus_{M \in \mathcal{O}_r} R_0 M$ . Then we can see immediately that  $\mathcal{H}_{R_0, r}(q) = \mathcal{H}_{R_0, r}^e(q) \oplus \mathcal{H}_{R_0, r}^o(q)$ . Furthermore we obtain  $\mathcal{H}_{R_0, r}^e(q) = \mathcal{H}_{R_0, r}^1(q)$  and  $\mathcal{H}_{R_0, r}^o(q) = \mathcal{H}_{R_0, r}^{-1}(q)$  from the property  $\hat{T}'_i = -T'_i$ . Combined with Lemma 3.5 we have  $\text{rank}_{R_0} \mathcal{H}_{R_0, r}(q) = 2 \text{rank}_{R_0} \mathcal{H}_{R_0, r}^1(q)$ . Let  $\bar{\mathcal{H}}_{R_0, r}^1(q)$  be the set of all the linear combinations of products of even numbers of  $T'_i$ 's. Obviously  $\mathcal{H}_{R_0, r}^1(q) \subseteq \bar{\mathcal{H}}_{R_0, r}^1(q)$ . From the defining relations (A'1)-(A'3), one can see that if a monomial in  $T'_i$ -normal form which consists of even (respectively odd) number of  $T'_i$ 's is expressed in a linear combination of other expressions, then each term consists of even (respectively odd) number of  $T'_i$ 's. Hence if we express an element of  $\bar{\mathcal{H}}_{R_0, r}^1(q)$  by a linear combination of monomials in  $T'_i$ -normal form, each term is in  $\mathcal{H}_{R_0, r}^1(q)$ . Consequently  $\bar{\mathcal{H}}_{R_0, r}^1(q) = \mathcal{H}_{R_0, r}^1(q)$ .  $\square$

When we suppose that  $R_0 = \mathbb{C}$  and take a limit  $q \rightarrow 1$ ,  $\mathcal{H}_{\mathbb{C}, r}^1(1)$  is isomorphic to the group algebra  $\mathbb{C}[\mathfrak{A}_n]$  of the alternating group  $\mathfrak{A}_n$ .

**Theorem 3.7 ([9]).**  $\mathcal{H}_{R_0, r}^1(q)$  is isomorphic to the  $R_0$ -algebra which is generated by  $r-2$  elements  $X_1, X_2, \dots, X_{r-2}$  with the defining relations:

$$(B1) \quad X_1^3 = -\left(\frac{q-q^{-1}}{q+q^{-1}}\right)^2 (X_1^2 - X_1) + 1,$$

$$(B2) \quad X_i^2 = 1 \quad \text{for } i > 1,$$

$$(B3) \quad (X_{i-1} X_i)^3 = -\left(\frac{q-q^{-1}}{q+q^{-1}}\right)^2 \left\{ (X_{i-1} X_i)^2 - X_{i-1} X_i \right\} + 1 \quad \text{for } i = 2, 3, \dots, r-2,$$

(B4)  $(X_i X_j)^2 = 1$  whenever  $|i - j| > 1$ .

An isomorphism is given by  $X_i \longrightarrow T'_1 T'_{i+1}$ .

Next, we shall show that  $\mathcal{H}_{R_0, r}(q)$  is a  $\mathbb{Z}_2$ -crossed product. Since  $T'_i$  has an inverse as itself, we can readily see from Proposition 3.6 that  $\mathcal{H}_{R_0, r}^1(q) T'_1$  is an  $R_0$ -submodule of  $\mathcal{H}_{R_0, r}(q)$  which consists of all linear combinations of products of odd numbers of  $T'_i$ 's. Therefore, we obtain a direct sum decomposition of left  $\mathcal{H}_{R_0, r}^1(q)$ -modules:

$$\mathcal{H}_{R_0, r}(q) = \mathcal{H}_{R_0, r}^1(q) \oplus \mathcal{H}_{R_0, r}^1(q) T'_1. \quad (3.1)$$

Let  $\mathbb{Z}_2 = \langle 1, -1 \rangle$  be a multiplicative group. We define two maps  $\psi_0$  and  $\alpha_0$  to be

$$\psi_0 : \mathbb{Z}_2 \longrightarrow \text{Aut}(\mathcal{H}_{R_0, r}^1(q)), \quad \psi_0(1)(T) = T, \quad \psi_0(-1)(T) = T'_1 T T'_1, \quad T \in \mathcal{H}_{R_0, r}^1(q). \quad (3.2)$$

and

$$\alpha_0 : \mathbb{Z}_2 \times \mathbb{Z}_2 \longrightarrow (\mathcal{H}_{R_0, r}^1(q))^\times, \quad \alpha_0(\sigma, \tau) = 1 \quad \text{for all } \sigma, \tau \in \mathbb{Z}_2. \quad (3.3)$$

Then we have the following immediately.

**Lemma 3.8.**  $\psi_0$  and  $\alpha_0$  satisfy (2.1)–(2.3).

Thus, we obtain a  $\mathbb{Z}_2$ -crossed product  $\mathcal{H}_{R_0, r}^1(q)_{\alpha_0}^{\psi_0}[\mathbb{Z}_2]$  from the crossed system  $(\mathcal{H}_{R_0, r}^1(q), \mathbb{Z}_2, \psi_0, \alpha_0)$ , where  $\psi_0$  and  $\alpha_0$  are given by (3.2) and (3.3) respectively.

**Theorem 3.9.**  $\mathcal{H}_{R_0, r}(q)$  is isomorphic to  $\mathcal{H}_{R_0, r}^1(q)_{\alpha_0}^{\psi_0}[\mathbb{Z}_2]$  as  $R_0$ -algebras.

*Proof.* Since both  $\mathcal{H}_{R_0, r}(q)$  and  $\mathcal{H}_{R_0, r}^1(q)_{\alpha_0}^{\psi_0}[\mathbb{Z}_2]$  are free left  $\mathcal{H}_{R_0, r}^1(q)$ -modules, we may define an isomorphism of  $\mathcal{H}_{R_0, r}^1(q)$ -modules

$$\iota_0 : \mathcal{H}_{R_0, r}^1(q)_{\alpha_0}^{\psi_0}[\mathbb{Z}_2] \longrightarrow \mathcal{H}_{R_0, r}(q), \quad \iota_0(u_1) = 1, \quad \iota_0(u_{-1}) = T'_1$$

From (2.4) and (3.2) and (3.3), we can determine the multiplication law as follows.

$$(a_1 u_\sigma)(a_2 u_\tau) = \begin{cases} a_1 a_2 u_{\sigma\tau} & \text{if } \sigma = 1, \\ a_1 T'_1 a_2 T'_1 u_{\sigma\tau} & \text{if } \sigma = -1, \end{cases}$$

where  $a_1, a_2 \in \mathcal{H}_{R_0, r}^1(q)$  and  $\sigma, \tau \in \mathbb{Z}_2$ . Therefore, we get four formulas in  $\mathcal{H}_{R_0, r}(q)$

$$(a_1 1)(a_2 1) = a_1 a_2 1, \quad (a_1 1)(a_2 T'_1) = a_1 a_2 T'_1, \quad (a_1 T'_1)(a_2 1) = a_1 T'_1 a_2 T'_1 T'_1, \quad (a_1 T'_1)(a_2 T'_1) = a_1 T'_1 a_2 T'_1 1,$$

which derive the conclusion that  $\iota_0$  is an isomorphism of  $R_0$ -algebras.  $\square$



We denote by  $\bar{K}$  an algebraic closure of a field  $K$ . Let  $q$  be an indeterminate and  $K = \mathbb{Q}(q)$ . We shall show (split) semisimplicity of  $\mathcal{H}_{\bar{K},r}^1(q)$  and the branching rule from  $\mathcal{H}_{\bar{K},r}(q)$  to  $\mathcal{H}_{\bar{K},r}^1(q)$ . The manner of proof given here is credited to K.Uno, who sent me a letter enclosing the outline of this proof. It is well-known that  $\mathcal{H}_{\bar{K},r}(q)$  is split semisimple and that isomorphism classes of simple left  $\mathcal{H}_{\bar{K},r}(q)$ -modules are parametrized by Young diagrams of total size  $n$ . Let  $\Lambda_r$  be the set of all Young diagrams of total size  $r$ . Then, provided that  $\{M_{q,\lambda} | \lambda \in \Lambda_r\}$  is a set of all isomorphism classes of simple left  $\mathcal{H}_{\bar{K},r}(q)$ -modules and that  $d_\lambda = \deg M_{q,\lambda}$ , we may write

$$\mathcal{H}_{\bar{K},r}(q) = \bigoplus_{\lambda \in \Lambda_r} I_{q,\lambda} \quad (I_{q,\lambda} \cong \text{Mat}_{d_\lambda}(\bar{K})), \quad (3.4)$$

where each  $I_{q,\lambda}$  is the homogeneous component corresponding to  $\lambda$ .  $I_{q,\lambda}$  is isomorphic to a  $d_\lambda \times d_\lambda$  matrix algebra  $\text{Mat}_{d_\lambda}(\bar{K})$  whose entries lie in  $\bar{K}$ . Since  $\hat{\phantom{x}}$  is an involution of  $\mathcal{H}_{\bar{K},r}(q)$ , for each  $\lambda \in \Lambda_r$  there exists  $\mu \in \Lambda_r$  such that  $\hat{I}_{q,\lambda} = I_{q,\mu}$ . Especially,  $d_\lambda = d_\mu$  follows. Dipper and James defined Specht modules  $S_K^\lambda$  for Hecke algebras as irreducible submodules of regular modules in [4], and improved the theory of representations of Hecke algebras in the series of articles such as [4, 5, 6]. In particular, they showed in [6] that if  $K$  is a field and  $\mathcal{H}_{K,r}(q)$  is semisimple, then  $\hat{S}_K^\lambda \cong_{\mathcal{H}_{K,r}(q)} S_K^{\lambda'}$  where  $\lambda'$  denotes the transpose of  $\lambda$ . In this case,  $M_{q,\lambda}$  is equivalent to  $S_K^\lambda$ , thus  $\mu = \lambda'$  follows. We divide into two cases depending on whether  $\lambda$  is self-conjugate or not.

(Case1)  $\lambda \neq \lambda'$ :

In this case,  $\hat{\phantom{x}}$  induces an involution on  $I_{q,\lambda} \oplus I_{q,\lambda'}$ . Let  $\tilde{I}_{q,\lambda} = \{X + Y \in I_{q,\lambda} \oplus I_{q,\lambda'} | (X + Y)^\hat{=} X + Y\}$  for  $\lambda \neq \lambda'$ . Then we have

$$\tilde{I}_{q,\lambda} = \{X + \hat{X} \in I_{q,\lambda} \oplus I_{q,\lambda'} | X \in I_{q,\lambda}\} \cong I_{q,\lambda} (\cong \text{Mat}_{d_\lambda}(\bar{K})), \quad (3.5)$$

Thereby, the image of the regular representation of  $\mathcal{H}_{\bar{K},r}^1(q)$  on  $S_{\bar{K}}^\lambda$  is isomorphic to  $I_{q,\lambda}$ , so  $\text{res}_{\mathcal{H}_{\bar{K},r}^1(q)}^{\mathcal{H}_{\bar{K},r}(q)} S_{\bar{K}}^\lambda$  is a simple left  $\mathcal{H}_{\bar{K},r}^1(q)$ -module. If  $g \in \mathcal{H}_{\bar{K},r}(q)$  satisfies  $\hat{g} = g$ , then the matrix coefficients with respect to the basis  $x_1, x_2, \dots, x_{d_\lambda}$  of  $S_{\bar{K}}^\lambda$  is the same as those with respect to the basis  $\hat{x}_1, \hat{x}_2, \dots, \hat{x}_{d_\lambda}$  of  $\hat{S}_{\bar{K}}^\lambda \cong S_{\bar{K}}^{\lambda'}$ . Hence we have the isomorphism of simple left  $\mathcal{H}_{\bar{K},r}^1(q)$ -modules as follows.

$$\text{res}_{\mathcal{H}_{\bar{K},r}^1(q)}^{\mathcal{H}_{\bar{K},r}(q)} S_{\bar{K}}^\lambda \cong \text{res}_{\mathcal{H}_{\bar{K},r}^1(q)}^{\mathcal{H}_{\bar{K},r}(q)} S_{\bar{K}}^{\lambda'}.$$

(Case2)  $\lambda = \lambda'$ :

In this case,  $\hat{\phantom{x}}$  induces an involution on  $I_{q,\lambda} \cong \text{Mat}_{d_\lambda}(\bar{K})$ . By Skolem-Noether Theorem, there exists an invertible element  $P$  of  $\text{Mat}_{d_\lambda}(\bar{K})$  such that  $\hat{X} = PXP^{-1}$  for all  $X \in \text{Mat}_{d_\lambda}(\bar{K})$ . Since eigenvalues of  $P$

are  $\pm 1$ , We may also assume that

$$P = \begin{bmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & -1 & & \\ & & & & \ddots & \\ & & & & & -1 \end{bmatrix}.$$

Then we may write

$$P \begin{bmatrix} X_1 & X_2 \\ X_3 & X_4 \end{bmatrix} P^{-1} = \begin{bmatrix} X_1 & -X_2 \\ -X_3 & X_4 \end{bmatrix},$$

for some submatrices  $X_1, X_2, X_3, X_4$ . Therefore, we obtain

$$\{X \in \text{Mat}_{d_\lambda}(\bar{K}) \mid \hat{X} = X\} = \left\{ X \in \text{Mat}_{d_\lambda}(\bar{K}) \mid X = \begin{bmatrix} X_1 & 0 \\ 0 & X_4 \end{bmatrix} \right\}.$$

Assuming that 1 appears  $m$  times in  $P$ . Then  $\dim_{\bar{K}} \{X \in \text{Mat}_{d_\lambda}(\bar{K}) \mid \hat{X} = X\} = m^2 + (d_\lambda - m)^2$ . We easily see that

$$m^2 + (d_\lambda - m)^2 \geq \frac{d_\lambda^2}{2} \quad \text{equality holds iff } m = \frac{d_\lambda}{2}.$$

Combining this with (3.5) and the fact that  $\dim_{\bar{K}} \mathcal{H}_{\bar{K},r}^1(q) = \frac{1}{2} \dim_{\bar{K}} \mathcal{H}_{\bar{K},r}(q)$  (Proposition 3.6), we deduce that  $m = d_\lambda/2$ . Thereby,

$$\{X \in \text{Mat}_{d_\lambda}(\bar{K}) \mid \hat{X} = X\} \cong \text{Mat}_{d_\lambda/2}(\bar{K}) \oplus \text{Mat}_{d_\lambda/2}(\bar{K}) \quad (3.6)$$

holds. This means that  $\text{res}_{\mathcal{H}_{\bar{K},r}^1(q)}^{\mathcal{H}_{\bar{K},r}(q)} S_{\bar{K}}^\lambda$  decomposes into two simple left  $\mathcal{H}_{\bar{K},r}^1(q)$ -modules  $\bar{S}_{\bar{K}}^{\lambda+}$  and  $\bar{S}_{\bar{K}}^{\lambda-}$  which are mutually non-isomorphic. Let  $\tilde{I}_{q,\lambda} = \{X \in I_{q,\lambda} \mid \hat{X} = X\}$  and  $\tilde{I}_{q,\lambda}^+$  (resp.  $\tilde{I}_{q,\lambda}^-$ ) be the homogeneous component corresponding to  $\bar{S}_{\bar{K}}^{\lambda+}$  (resp.  $\bar{S}_{\bar{K}}^{\lambda-}$ ) for  $\lambda \neq \lambda'$ . Then (3.6) implies that

$$\tilde{I}_{q,\lambda} = \tilde{I}_{q,\lambda}^+ \oplus \tilde{I}_{q,\lambda}^-, \quad (3.7)$$

Summarizing our argument, we conclude that  $\mathcal{H}_{\bar{K},r}^1(q)$  is isomorphic to the direct sum of minimal two-sided ideals as follows,

$$\mathcal{H}_{\bar{K},r}^1(q) = \left\{ \bigoplus_{\lambda \in \Lambda_r, \lambda > \lambda'} \tilde{I}_{q,\lambda} \right\} \oplus \left[ \bigoplus_{\lambda \in \Lambda_r, \lambda = \lambda'} \{ \tilde{I}_{q,\lambda}^+ \oplus \tilde{I}_{q,\lambda}^- \} \right],$$

where  $<$  denotes the lexicographic order on  $\Lambda_r$ ;  $\lambda = (\lambda_1, \lambda_2, \dots) < \mu = (\mu_1, \mu_2, \dots)$  iff  $\lambda_k < \mu_k$  for the smallest  $k$  such that  $\lambda_k \neq \mu_k$ . Consequently we have proved the following result.

**Theorem 3.10.** *Let  $q$  be an indeterminate and  $K = \mathbb{Q}(q)$ .*

*If  $\lambda' \neq \lambda$ , then  $\bar{S}_{\bar{K}}^\lambda$  is simple and  $\bar{S}_{\bar{K}}^\lambda \cong \bar{S}_{\bar{K}}^{\lambda'}$  where  $\bar{S}_{\bar{K}}^\lambda$  denotes  $\text{res}_{\mathcal{H}_{\bar{K},r}^1(q)}^{\mathcal{H}_{\bar{K},r}(q)} S_{\bar{K}}^\lambda$ .*

*If  $\lambda' = \lambda$ , then  $\bar{S}_{\bar{K}}^\lambda$  decomposes into mutually non-isomorphic two simple left  $\mathcal{H}_{\bar{K},r}^1(q)$ -modules  $\bar{S}_{\bar{K}}^{\lambda+}$  and  $\bar{S}_{\bar{K}}^{\lambda-}$ .*

*The simple left  $\mathcal{H}_{\bar{K},r}^1(q)$ -modules  $\bar{S}_{\bar{K}}^\lambda, \bar{S}_{\bar{K}}^{\mu+}, \bar{S}_{\bar{K}}^{\mu-}$  ( $\lambda, \mu \in \Lambda_r, \lambda > \lambda', \mu = \mu'$ ) constitute a basic set of simple left  $\mathcal{H}_{\bar{K},r}^1(q)$ -modules. Moreover,  $\mathcal{H}_{\bar{K},r}^1(q)$  is a split semisimple  $\bar{K}$ -algebra and its homogeneous decomposition is as follows:*

$$\mathcal{H}_{\bar{K},r}^1(q) = \left\{ \bigoplus_{\lambda \in \Lambda_r, \lambda > \lambda'} \tilde{I}_{q,\lambda} \right\} \bigoplus \left[ \bigoplus_{\lambda \in \Lambda_r, \lambda = \lambda'} \{ \tilde{I}_{q,\lambda}^+ \oplus \tilde{I}_{q,\lambda}^- \} \right],$$

*where each  $\tilde{I}_{q,\lambda}$  is the homogeneous component corresponding to  $\lambda$  which is isomorphic to  $\text{Mat}_{d_\lambda}(\bar{K})$  and each direct sum  $\tilde{I}_{q,\lambda}^+ \oplus \tilde{I}_{q,\lambda}^-$  consists of two homogeneous components  $\tilde{I}_{q,\lambda}^+$  and  $\tilde{I}_{q,\lambda}^-$  both corresponding to  $\lambda$  which are isomorphic to  $\text{Mat}_{d_\lambda/2}(\bar{K})$ .*

## 4 Schur-Weyl reciprocity between $U_q^\sigma(\mathfrak{gl}(m, n))$ and $\mathcal{H}_{K,r}(q)$

In our previous paper [10], we defined the  $q$ -permutation representation and established Schur-Weyl reciprocity between the quantum superalgebra  $U_q^\sigma(\mathfrak{gl}(m, n))$  and  $\mathcal{H}_{K,r}(q)$ . Schur-Weyl reciprocity between the general Lie superalgebra  $\mathfrak{gl}(m, n)$  and the symmetric group  $\mathfrak{S}_r$  was established in, for example, [3, 13]. In this section, we shall review the  $q$ -permutation representation and the vector representation of  $U_q^\sigma(\mathfrak{gl}(m, n))$ .

Let  $V = \bigoplus_{k=1}^{m+n} K v_k$  be a  $\mathbb{Z}_2$ -graded  $K$ -module of rank  $m+n$ . By  $\mathbb{Z}_2$ -graded, we mean that  $V$  is a direct sum of two submodule  $V_{\bar{0}} = \bigoplus_{k=1}^m K v_k$  and  $V_{\bar{1}} = \bigoplus_{k=m+1}^{m+n} K v_k$ , and that for each homogeneous element the degree map  $|\cdot|$

$$|v| = \begin{cases} 0 & \text{if } v \in V_{\bar{0}}, \\ 1 & \text{if } v \in V_{\bar{1}}, \end{cases}$$

is given.

Let  $\pi_r$  be the  $q$ -permutation representation of  $\mathcal{H}_{K,r}(q)$  on the tensor space  $V^{\otimes r}$ .  $\pi_r$  is given by  $\pi_r(T_i) = \text{Id}^{\otimes i-1} \otimes T \otimes \text{Id}^{\otimes r-i-1}$  ( $i = 1, 2, \dots, r-1$ ) where  $T$  is the operator on  $V \otimes V$  defined by

$$T v_k \otimes v_l = \begin{cases} \frac{(-1)^{|v_k|} (q + q^{-1}) + q - q^{-1}}{2} v_k \otimes v_l & \text{if } k = l, \\ (-1)^{|v_k||v_l|} v_l \otimes v_k + (q - q^{-1}) v_k \otimes v_l & \text{if } k < l, \\ (-1)^{|v_k||v_l|} v_l \otimes v_k & \text{if } k > l. \end{cases} \quad (4.1)$$

and  $\text{Id}$  is the identity operator on  $V$ . This representation  $\pi_r$  is reduced to the (normal)  $q$ -permutation representation of  $\mathcal{H}_{K,r}(q)$  with  $n = 0$  and to the sign permutation representation of  $\mathfrak{S}_r$  with  $q \rightarrow 1$ . Let  $T'$  be the operator defined by

$$T' = \frac{2T - (q - q^{-1})}{q + q^{-1}}$$

Then  $T'$  is determined by

$$T'v_k \otimes v_l = \begin{cases} (-1)^{|v_k|} v_k \otimes v_l & \text{if } k = l, \\ \frac{2(-1)^{|v_k||v_l|}}{q + q^{-1}} v_l \otimes v_k + \frac{q - q^{-1}}{q + q^{-1}} v_k \otimes v_l & \text{if } k < l, \\ \frac{2(-1)^{|v_k||v_l|}}{q + q^{-1}} v_l \otimes v_k - \frac{q - q^{-1}}{q + q^{-1}} v_k \otimes v_l & \text{if } k > l. \end{cases} \quad (4.2)$$

$\pi_r$  is also given by  $\pi_r(T'_i) = \text{Id}^{\otimes i-1} \otimes T' \otimes \text{Id}^{\otimes r-i-1}$  ( $i = 1, 2, \dots, r-1$ )

Next we shall review quantum superalgebras and their vector representations. Several definitions of quantum superalgebras appear in, for example, [2, 8, 14]. The much complete definition and detailed observations of quantum superalgebras can be found in [14]. In this paper, we obey the manner of definition of  $U_q^\sigma(\mathfrak{gl}(m, n))$  in [2]. The method of construction of superalgebra depends on [7] basically. Let  $\Pi = \{\alpha_i\}_{i \in I}$  be a set of simple roots with the index set  $I = \{1, \dots, r\}$ . We assume that  $I$  is a disjoint union of two subsets  $I_{\text{even}}$  and  $I_{\text{odd}}$ . We define a map  $p : I \rightarrow \{0, 1\}$  to be such that

$$p(i) = \begin{cases} 0 & \text{if } i \in I_{\text{even}}, \\ 1 & \text{if } i \in I_{\text{odd}}. \end{cases}$$

Let  $P$  be a free  $\mathbb{Z}$ -module which includes all  $\alpha_i \in P$  ( $i \in I$ ). We assume that a  $\mathbb{Q}$ -valued symmetric bilinear form on  $P$   $(\cdot, \cdot) : P \times P \rightarrow \mathbb{Q}$  is defined and that the simple coroots  $h_i \in P^*$  ( $i \in I$ ) are given as data. The natural pairing  $\langle \cdot, \cdot \rangle : P^* \times P \rightarrow \mathbb{Z}$  between  $P$  and  $P^*$  is assumed to satisfy

$$\langle h_i, \alpha_j \rangle = \begin{cases} 2 & \text{if } i = j \text{ and } i \in I_{\text{even}}, \\ 0 \text{ or } 2 & \text{if } i = j \text{ and } i \in I_{\text{odd}}, \\ \leq 0 & \text{if } i \neq j. \end{cases}$$

We denote by  $\Pi^\vee = \{h_i | i \in I\}$  the set of all coroots. Furthermore, for each  $i \in I$  we assume that there exists a nonzero integer  $\ell_i$  such that  $\ell_i \langle h_i, \lambda \rangle = (\alpha_i, \lambda)$  for every  $\lambda \in P$ . Then we immediately have the Cartan matrix  $A = [\langle h_i, \alpha_j \rangle]_{ij}$  is symmetrizable because  $\ell_i \langle h_i, \alpha_j \rangle = (\alpha_i, \alpha_j) = (\alpha_j, \alpha_i) = \ell_j \langle h_j, \alpha_i \rangle$ . We mention that the symmetrized matrix is  $A^{\text{sym}} = \text{diag}(\ell_1, \dots, \ell_r)A = [(\alpha_i, \alpha_j)]_{ij}$ . Let  $\mathfrak{h} = P^* \otimes_{\mathbb{Z}} \mathbb{Q}$ . Then  $\Phi = (\mathfrak{h}, \Pi^\vee, \Pi)$  is said to be a fundamental root data associated to  $A$ . Let  $\mathfrak{g} = \mathfrak{g}(\Phi)$  be the contragredient

Lie superalgebra obtained from  $\Phi$  and  $p$ . The quantized enveloping algebra  $U_q^\sigma(\mathfrak{g})$  is the unital associative algebra over  $K = \mathbb{Q}(q)$  with generators  $q^h (h \in P^*), e_i, f_i (i \in I)$  and an additional element  $\sigma$  which satisfy the following defining relations:

$$(Q1) \quad q^h = 1 \quad \text{for } h = 0,$$

$$(Q2) \quad q^{h_1} q^{h_2} = q^{h_1+h_2} \quad \text{for } h_1, h_2 \in P^*,$$

$$(Q3) \quad q^h e_i = q^{\langle h, \alpha_j \rangle} e_i q^h \quad \text{for } h \in P^* \text{ and } i \in I,$$

$$(Q4) \quad q^h f_i = q^{-\langle h, \alpha_j \rangle} f_i q^h \quad \text{for } h \in P^* \text{ and } i \in I,$$

$$(Q5) \quad [e_i, f_j] = \delta_{ij} \frac{q^{\ell_i h_i} - q^{-\ell_i h_i}}{q^{\ell_i} - q^{-\ell_i}} \quad \text{for } i, j \in I,$$

$$(Q6) \quad \sigma^2 = 1,$$

$$(Q7) \quad q^h \sigma = \sigma q^h \quad \text{for } h \in P^*,$$

$$(Q8) \quad e_i \sigma = (-1)^{p(i)} \sigma e_i \quad \text{for } i \in I,$$

$$(Q9) \quad f_i \sigma = (-1)^{p(i)} \sigma f_i \quad \text{for } i \in I,$$

where  $[e_i, f_j]$  means the supercommutator

$$[e_i, f_j] = e_i f_j - (-1)^{p(i)p(j)} f_j e_i.$$

We assume further conditions:

$$(Q10) \quad \text{If } a \in \sum_{i \in I} U_q(\mathfrak{n}_+) e_i U_q(\mathfrak{n}_+) \text{ satisfies } f_i a \in U_q(\mathfrak{n}_+) f_i \text{ for all } i \in I, \text{ then } a = 0,$$

$$(Q11) \quad \text{If } a \in \sum_{i \in I} U_q(\mathfrak{n}_-) f_i U_q(\mathfrak{n}_-) \text{ satisfies } e_i a \in U_q(\mathfrak{n}_-) e_i \text{ for all } i \in I, \text{ then } a = 0,$$

where  $U_q(\mathfrak{n}_+)$  (respectively  $U_q(\mathfrak{n}_-)$ ) is the subalgebra of  $U_q^\sigma(\mathfrak{g})$  generated by  $\{e_i | i \in I\}$  (respectively  $\{f_i | i \in I\}$ ).  $U_q^\sigma(\mathfrak{g})$  is a Hopf algebra whose comultiplication  $\Delta_\sigma$ , counit  $\varepsilon_\sigma$ , antipode  $S_\sigma$  are as follows.

$$\Delta_\sigma(\sigma) = \sigma \otimes \sigma,$$

$$\Delta_\sigma(q^h) = q^h \otimes q^h \quad \text{for } h \in P^*,$$

$$\Delta_\sigma(e_i) = e_i \otimes q^{-\ell_i h_i} + \sigma^{p(i)} \otimes e_i \quad \text{for } i \in I,$$

$$\Delta_\sigma(f_i) = f_i \otimes 1 + \sigma^{p(i)} q^{\ell_i h_i} \otimes f_i \quad \text{for } i \in I,$$

$$\varepsilon_\sigma(\sigma) = \varepsilon_\sigma(q^h) = 1 \quad \text{for } h \in P^*, \quad \varepsilon_\sigma(e_i) = \varepsilon_\sigma(f_i) = 0 \quad \text{for } i \in I,$$

$$S_\sigma(\sigma) = \sigma, \quad S_\sigma(q^{\pm h}) = q^{\mp h} \quad \text{for } h \in P^*,$$

$$S_\sigma(e_i) = -\sigma^{p(i)} e_i q^{\ell_i h_i}, \quad S_\sigma(f_i) = -\sigma^{p(i)} q^{-\ell_i h_i} f_i \quad \text{for } i \in I.$$

The quantized enveloping algebra  $U_q^\sigma(\mathfrak{gl}(m, n))$  is obtained from the fundamental root data as follows.

- $I = I_{\text{even}} \cup I_{\text{odd}}$  is defined by  $I_{\text{even}} = \{1, 2, \dots, m-1, m+1, \dots, m+n-1\}$  and  $I_{\text{odd}} = \{m\}$ ,
- $P = \bigoplus_{b \in B} \mathbb{Z}\epsilon_b$ , where  $B = B_+ \cup B_-$  with  $B_+ = \{1, \dots, m\}$  and  $B_- = \{m+1, \dots, m+n\}$ ,
- $(\cdot, \cdot) : P \times P \rightarrow \mathbb{Q}$  is the symmetric bilinear form on  $P$  defined by

$$(\epsilon_a, \epsilon_{a'}) = \begin{cases} 1 & \text{if } a = a' \in B_+, \\ -1 & \text{if } a = a' \in B_-, \\ 0 & \text{otherwise,} \end{cases}$$

- $\Pi = \{\alpha_i | i \in I\}$  is defined by  $\alpha_i = \epsilon_i - \epsilon_{i+1}$ ,
- $\Pi^\vee = \{h_i | i \in I\}$  is uniquely determined by the formula  $\ell_i \langle h_i, \lambda \rangle = (\alpha_i, \lambda)$  for any  $\lambda \in P$ ,

where

$$\ell_i = \begin{cases} 1 & \text{if } i = 1, \dots, m, \\ -1 & \text{if } i = m+1, \dots, m+n-1. \end{cases}$$

The vector representation  $(\rho, V)$  of  $U_q^\sigma(\mathfrak{gl}(m, n))$  on the  $\mathbb{Z}_2$ -graded vector space  $V = V_0 \oplus V_1$  (recall that  $V_0 = \bigoplus_{i=1}^m K v_i$ ,  $V_1 = \bigoplus_{i=m+1}^{m+n} K v_i$ ) is defined by (see [2])

$$\begin{aligned} \rho(\sigma)v_j &= (-1)^{|v_j|} v_j \quad \text{for } j = 1, \dots, m+n, \\ \rho(q^h)v_j &= q^{\epsilon_j(h)} v_j \quad \text{for } h \in P^*, j = 1, \dots, m+n, \\ \rho(e_j)v_{j+1} &= v_j \quad \text{for } j = 1, \dots, m+n-1, \\ \rho(f_j)v_j &= v_{j+1} \quad \text{for } j = 1, \dots, m+n-1, \\ &\text{otherwise } 0. \end{aligned} \tag{4.3}$$

The vector representation  $\rho_r$  on  $V^{\otimes r}$  is given by  $\rho_r(x) = \rho^{\otimes r} \circ \Delta^{(r-1)}(x)$  where  $\Delta^{(1)} = \Delta_\sigma$  and  $\Delta^{(k)} = (\Delta_\sigma \otimes \text{Id}^{\otimes k-1}) \Delta^{(k-1)}$  inductively. One can readily see that  $\rho_r$  is of the following form:

$$\begin{aligned} \rho_r(\sigma) &= \rho(\sigma)^{\otimes r}, \\ \rho_r(q^h) &= \rho(q^h)^{\otimes r} \quad \text{for } h \in P^*, \\ \rho_r(e_i) &= \sum_{k=1}^N \rho(\sigma^{p(i)})^{\otimes k-1} \otimes \rho(e_i) \otimes \rho(q^{-\ell_i h_i})^{\otimes r-k} \quad \text{for } i \in I, \\ \rho_r(f_i) &= \sum_{k=1}^r \rho(\sigma^{p(i)} q^{\ell_i h_i})^{\otimes k-1} \otimes \rho(f_i) \otimes \text{Id}^{\otimes r-k} \quad \text{for } i \in I. \end{aligned} \tag{4.4}$$

Our precedent works in [10] are Theorem 4.1 and Theorem 4.2.

**Theorem 4.1 (Schur-Weyl reciprocity [10]).** *Let  $\mathcal{A}_q = \pi_r(\mathcal{H}_{K,r}(q))$  and  $\mathcal{B}_q = \rho_r(U_q^\sigma(\mathfrak{gl}(m, n)))$ . Then we have  $\text{End}_{\mathcal{B}_q} V^{\otimes r} = \mathcal{A}_q$  and  $\text{End}_{\mathcal{A}_q} V^{\otimes r} = \mathcal{B}_q$ .*

Let  $\bar{U}_q^\sigma(\mathfrak{gl}(m, n)) = U_q^\sigma(\mathfrak{gl}(m, n)) \otimes_K \bar{K}$ ,  $\bar{\mathcal{A}}_q = \mathcal{A}_q \otimes_K \bar{K}$  and  $\bar{\mathcal{B}}_q = \mathcal{B}_q \otimes_K \bar{K}$ . Then,  $\pi_r(\mathcal{H}_{\bar{K},r}(q)) = \bar{\mathcal{A}}_q$  and  $\rho_r(\bar{U}_q^\sigma(\mathfrak{gl}(m, n))) = \bar{\mathcal{B}}_q$  as  $\bar{K}$ -algebras of operators on  $\bar{V}^{\otimes r} = (V \otimes_K \bar{K})^{\otimes r}$ . We notice that  $\text{End}_{\bar{\mathcal{B}}_q} \bar{V}^{\otimes r} = \bar{\mathcal{A}}_q$  and  $\text{End}_{\bar{\mathcal{A}}_q} \bar{V}^{\otimes r} = \bar{\mathcal{B}}_q$  hold.

Let  $H(m, n; r) = \{\lambda = (\lambda_1, \lambda_2, \dots) \in \Lambda_r \mid \lambda_j \leq n \text{ if } j > m\}$ . Diagrams of elements of  $H(m, n; r)$  are exactly those contained in the  $(m, n)$ -hooks. Then the following holds.

**Theorem 4.2 (Decomposition of the tensor space [10]).**  *$\bar{\mathcal{A}}_q = \bigoplus_{\lambda \in H(m, n; r)} \bar{\mathcal{A}}_{q, \lambda}$  where each  $\bar{\mathcal{A}}_{q, \lambda} = \pi_r(I_{q, \lambda})(\cong I_{q, \lambda})$  is the image of the homogeneous component  $I_{q, \lambda}$  corresponding to  $\lambda \in H(m, n; r)$  as in (3.4). Moreover, we have  $\bar{V}^{\otimes r} = \bigoplus_{\lambda \in H(m, n; r)} H_\lambda \otimes V_\lambda$  where  $H_\lambda$ 's are mutually non-isomorphic simple left  $\mathcal{H}_{\bar{K},r}(q)$ -modules indexed by the elements of  $H(m, n; r)$ , and  $V_\lambda$ 's are mutually non-isomorphic simple left  $\bar{U}_q^\sigma(\mathfrak{gl}(m, n))$ -modules indexed by the elements of  $H(m, n; r)$ .*

## 5 Schur-Weyl reciprocity for $\mathcal{H}_{K,r}^1(q)$ in case of $m = n$

Schur-Weyl reciprocity for the alternating group  $\mathfrak{A}_r$  has been researched by Regev in [12]. The paper [12] showed that if  $\dim V_{\bar{0}} = \dim V_{\bar{1}}$  under the base field  $\mathbb{C}$ , then the centralizer algebra  $\text{End}_{\mathfrak{A}_r} V^{\otimes r}$  has remarkable property;  $\mathbb{Z}_2$ -crossed product for  $\text{End}_{\mathfrak{S}_r} V^{\otimes r}$ . In this paper, we establish a  $q$ -analogue extension of Regev's result. We also show that Schur-Weyl reciprocity is valid even if the base field is  $\mathbb{Q}(q)$ .

We set  $K = \mathbb{Q}(q)$  in succession, and denote  $\mathcal{C}_q = \pi_r(\mathcal{H}_{K,r}^1(q))$ . Let us consider the relation between  $\text{End}_{\mathcal{A}_q} V^{\otimes r}$  and  $\text{End}_{\mathcal{C}_q} V^{\otimes r}$ . Since  $\mathcal{H}_{K,r}^1(q) \subsetneq \mathcal{H}_{K,r}(q)$ , we immediately have

$$\mathcal{C}_q \subseteq \mathcal{A}_q \quad \text{and} \quad \text{End}_{\mathcal{A}_q} V^{\otimes r} \subseteq \text{End}_{\mathcal{C}_q} V^{\otimes r}.$$

Recall that  $V$  is an  $m + n$ -dimensional  $\mathbb{Z}_2$ -graded vector space over  $K$ . In this section, we analyze the structure of  $\text{End}_{\mathcal{C}_q} V^{\otimes r}$  in case of  $m = n$ . We will consider the general case in the next section.

Assume that  $m = n$ . Recall that

$$\mathcal{B}_q = \{f \in \text{End}_K V^{\otimes r} \mid \pi_r(T'_i) f = f \pi_r(T'_i) \text{ for } i = 1, 2, \dots, r-1\} = \text{End}_{\mathcal{A}_q} V^{\otimes r}.$$

We define  $\mathcal{B}_q^\dagger, \mathcal{C}_q$  to be the subspaces of  $\text{End}_K V^{\otimes r}$  as follows.

$$\mathcal{B}_q^\dagger = \{f \in \text{End}_K V^{\otimes r} \mid \pi_r(T'_i) f = -f \pi_r(T'_i) \text{ for } i = 1, 2, \dots, r-1\},$$

$$\mathcal{D}_q = \{f \in \text{End}_K V^{\otimes r} \mid \pi_r(T'_1 T'_{i+1}) f = f \pi_r(T'_1 T'_{i+1}) \text{ for } i = 1, 2, \dots, r-2\} = \text{End}_{\mathcal{C}_q} V^{\otimes r}.$$

**Lemma 5.1.**  $\mathcal{D}_q = \mathcal{B}_q \oplus \mathcal{B}_q^\dagger$ .

*Proof.* It is clear that the sum is direct.  $\supseteq$  is also obvious. We notice that if  $f \in \mathcal{D}_q$ , then  $\pi_r(T'_j)f\pi_r(T'_j) = \pi_r(T'_1)f\pi_r(T'_1)$  for  $j = 2, 3, \dots, r-1$  from the definition of  $\mathcal{D}_q$ .

In general, we may write

$$f = \frac{1}{2}(f + \pi_r(T'_1)f\pi_r(T'_1)) + \frac{1}{2}(f - \pi_r(T'_1)f\pi_r(T'_1)).$$

If  $f \in \mathcal{D}_q$ , then one can readily see

$$\begin{aligned} \pi_r(T'_i)\frac{1}{2}(f + \pi_r(T'_1)f\pi_r(T'_1)) &= \frac{1}{2}(\pi_r(T'_i)f\pi_r(T'_i)^2 + \pi_r(T'_i)\pi_r(T'_1)f\pi_r(T'_1)) \\ &= \frac{1}{2}(\pi_r(T'_1)f\pi_r(T'_1)\pi_r(T'_i) + f\pi_r(T'_i)\pi_r(T'_1)^2) \\ &= \frac{1}{2}(\pi_r(T'_1)f\pi_r(T'_1) + f)\pi_r(T'_i). \end{aligned}$$

Hence we have  $2^{-1}(f + \pi_r(T'_1)f\pi_r(T'_1)) \in \mathcal{B}_q$ . In the same fashion, we also get  $2^{-1}(f - \pi_r(T'_1)f\pi_r(T'_1)) \in \mathcal{B}_q^\dagger$ . Thus we have proved the reverse inclusion  $\subseteq$ .  $\square$

We notice that if  $V$  is  $\mathbb{Z}_2$ -graded, then  $\text{End}_K V$  is also  $\mathbb{Z}_2$ -graded. Namely,

$$\text{End}_K V = (\text{End}_K V)_{\bar{0}} \oplus (\text{End}_K V)_{\bar{1}},$$

where

$$\begin{aligned} (\text{End}_K V)_{\bar{0}} &= \{f \in \text{End}_K V \mid f(V_i) \subseteq V_i, i \in \{0, 1\}\}, \\ (\text{End}_K V)_{\bar{1}} &= \{f \in \text{End}_K V \mid f(V_i) \subseteq V_{i+1}, i \in \{0, 1\}\}. \end{aligned}$$

Let  $\varphi \in \text{End}_K V$  be given by  $\varphi(v_i) = v_{2m-i+1}$ . Obviously,  $\varphi \in (\text{End}_K V)_{\bar{1}}$ . Let  $\varphi^{\otimes r}$  be the tensor product of  $\varphi$ . In general, for homogeneous elements  $f_1, f_2, \dots, f_r$  of  $\text{End}_K V$ ,  $f_1 \otimes f_2 \otimes \dots \otimes f_r \in \text{End}_K V^{\otimes r}$  is given by

$$f_1 \otimes \dots \otimes f_r(u_1 \otimes \dots \otimes u_r) = (-1)^{\sum_{i=2}^r (|f_i| \sum_{j=1}^{i-1} |u_j|)} f_1(u_1) \otimes \dots \otimes f_r(u_r),$$

where  $u_1, \dots, u_r$  are homogeneous elements of  $V$ . Hence we have

$$\varphi^{\otimes r}(u_1 \otimes \dots \otimes u_r) = (-1)^{\sum_{i=2}^r (\sum_{j=1}^{i-1} |u_j|)} \varphi(u_1) \otimes \dots \otimes \varphi(u_r). \quad (5.1)$$

**Lemma 5.2.**  $\varphi^{\otimes r}$  is an isomorphism of  $K$ -vector space  $V^{\otimes r}$  which satisfies following properties:

$$(\varphi^{\otimes r})^2 = (-1)^{r(r-1)/2} \mathbf{I} \quad (5.2)$$

and

$$\pi_r(T'_i)\varphi^{\otimes r} = -\varphi^{\otimes r}\pi_r(T'_i) \quad \text{for } i = 1, 2, \dots, r-1, \quad (5.3)$$

where  $\mathbf{I}$  is the identity operator on  $V^{\otimes r}$ .



*Proof.* For homogeneous elements  $u_1, \dots, u_r$  of  $V$ , one can readily see that

$$\begin{aligned}
(\varphi^{\otimes r})^2(u_1 \otimes \dots \otimes u_r) &= (-1)^{\sum_{i=2}^r (\sum_{j=1}^{i-1} |u_j|)} \varphi(u_1) \otimes \dots \otimes \varphi(u_r) \\
&= (-1)^{\sum_{i=2}^r (\sum_{j=1}^{i-1} (|u_j| + |\varphi(u_j)|))} u_1 \otimes \dots \otimes u_r \\
&= (-1)^{\sum_{i=2}^r \sum_{j=1}^{i-1} 1} u_1 \otimes \dots \otimes u_r \\
&= (-1)^{(r-1)r/2} u_1 \otimes \dots \otimes u_r.
\end{aligned}$$

Hence  $(\varphi^{\otimes r})^2 = (-1)^{r(r-1)/2} \mathbf{I}$ , and  $\varphi^{\otimes r}$  has the inverse  $(\varphi^{\otimes r})^{-1} = (-1)^{(r-1)r/2} \varphi^{\otimes r}$ .

To prove the last statement, we check three cases of the definition of  $\pi_r$  which has appeared in (4.2). We notice that it suffice to prove only the case  $r = 2$ .

Case1 :  $k = l$

$$\begin{aligned}
\varphi^{\otimes 2} T'(v_k \otimes v_l) &= (-1)^{|v_k|} \varphi^{\otimes 2}(v_k \otimes v_k) \\
&= v_{2m-k+1} \otimes v_{2m-k+1} \\
T' \varphi^{\otimes 2}(v_k \otimes v_l) &= (-1)^{|v_k|} T'(v_{2m-k+1} \otimes v_{2m-k+1}) \\
&= (-1)^{|v_k|} (-1)^{|v_{2m-k+1}|} v_{2m-k+1} \otimes v_{2m-k+1} \\
&= -v_{2m-k+1} \otimes v_{2m-k+1}
\end{aligned}$$

Case2 :  $k < l$

$$\begin{aligned}
\varphi^{\otimes 2} T'(v_k \otimes v_l) &= \frac{2(-1)^{|v_k||v_l|}}{q+q^{-1}} \varphi^{\otimes 2}(v_l \otimes v_k) + \frac{q-q^{-1}}{q+q^{-1}} \varphi^{\otimes 2}(v_k \otimes v_l) \\
&= \frac{2(-1)^{|v_k||v_l|+|v_l|}}{q+q^{-1}} (v_{2m-l+1} \otimes v_{2m-k+1}) + \frac{q-q^{-1}}{q+q^{-1}} (-1)^{|v_k|} (v_{2m-k+1} \otimes v_{2m-l+1})
\end{aligned}$$

$$\begin{aligned}
T' \varphi^{\otimes 2}(v_k \otimes v_l) &= (-1)^{|v_k|} T'(v_{2m-k+1} \otimes v_{2m-l+1}) \\
&= \frac{2(-1)^{|v_k|+|v_{2m-k+1}|+|v_{2m-l+1}|}}{q+q^{-1}} (v_{2m-l+1} \otimes v_{2m-k+1}) - \frac{q-q^{-1}}{q+q^{-1}} (-1)^{|v_k|} (v_{2m-k+1} \otimes v_{2m-l+1})
\end{aligned}$$

If  $|v_k| = |v_l| = 0$ , then  $(-1)^{|v_k||v_l|+|v_l|} = 1$  and  $(-1)^{|v_k|+|v_{2m-k+1}|+|v_{2m-l+1}|} = -1$ .

If  $|v_k| = 0, |v_l| = 1$ , then  $(-1)^{|v_k||v_l|+|v_l|} = -1$  and  $(-1)^{|v_k|+|v_{2m-k+1}|+|v_{2m-l+1}|} = 1$ .

If  $|v_k| = 1, |v_l| = 0$ , then  $(-1)^{|v_k||v_l|+|v_l|} = 1$  and  $(-1)^{|v_k|+|v_{2m-k+1}|+|v_{2m-l+1}|} = -1$ .

If  $|v_k| = |v_l| = 1$ , then  $(-1)^{|v_k||v_l|+|v_l|} = 1$  and  $(-1)^{|v_k|+|v_{2m-k+1}|+|v_{2m-l+1}|} = -1$ .

After all,  $\varphi^{\otimes 2} T'(v_k \otimes v_l) = -T' \varphi^{\otimes 2}(v_k \otimes v_l)$  holds.

Case3 :  $k > l$

$$\begin{aligned}
\varphi^{\otimes 2} T'(v_k \otimes v_l) &= \frac{2(-1)^{|v_k||v_l|}}{q+q^{-1}} \varphi^{\otimes 2}(v_l \otimes v_k) - \frac{q-q^{-1}}{q+q^{-1}} \varphi^{\otimes 2}(v_k \otimes v_l) \\
&= \frac{2(-1)^{|v_k||v_l|+|v_l|}}{q+q^{-1}} (v_{2m-l+1} \otimes v_{2m-k+1}) - \frac{q-q^{-1}}{q+q^{-1}} (-1)^{|v_k|} (v_{2m-k+1} \otimes v_{2m-l+1})
\end{aligned}$$

$$\begin{aligned}
T' \varphi^{\otimes 2}(v_k \otimes v_l) &= (-1)^{|v_k|} T'(v_{2m-k+1} \otimes v_{2m-l+1}) \\
&= \frac{2(-1)^{|v_k|+|v_{2m-k+1}|+|v_{2m-l+1}|}}{q+q^{-1}} (v_{2m-l+1} \otimes v_{2m-k+1}) + \frac{q-q^{-1}}{q+q^{-1}} (-1)^{|v_k|} (v_{2m-k+1} \otimes v_{2m-l+1})
\end{aligned}$$

In the same manner as case 2, we have  $\varphi^{\otimes 2} T'(v_k \otimes v_l) = -T' \varphi^{\otimes 2}(v_k \otimes v_l)$ .  $\square$

Let  $\Phi$  be the endomorphism of the vector space  $\text{End}_K V^{\otimes r}$  which is given by:

$$\Phi(f) = \varphi^{\otimes r} f \quad \text{for } f \in \text{End}_K V^{\otimes r}.$$

Then we have the following.

**Proposition 5.3.**  $\Phi$  is an automorphism of the vector space  $\text{End}_K V^{\otimes r}$ . The restriction of  $\Phi$  to  $\mathcal{D}_q$  gives an automorphism of  $\mathcal{D}_q$  which satisfies the following properties:

$$\Phi(\mathcal{B}_q) = \mathcal{B}_q^\dagger, \quad \Phi(\mathcal{B}_q^\dagger) = \mathcal{B}_q.$$

Especially, we have  $\dim_K \mathcal{B}_q = \dim_K \mathcal{B}_q^\dagger$  and  $\mathcal{D}_q = \mathcal{B}_q \oplus \Phi(\mathcal{B}_q) = \mathcal{B}_q \oplus \varphi^{\otimes r} \mathcal{B}_q$ .

*Proof.* From (5.2), it immediately follows that  $\Phi^2 f = (-1)^{r(r-1)/2} f$  for  $f \in \text{End}_K V^{\otimes r}$ . Hence  $\Phi$  is an isomorphism of  $\text{End}_K V^{\otimes r}$ . Let  $f \in \mathcal{B}_q$ . From (5.3), it follows that

$$\begin{aligned}
\pi_r(T'_i) \Phi(f) &= \pi_r(T'_i) \varphi^{\otimes r} f \\
&= -\varphi^{\otimes r} \pi_r(T'_i) f \\
&= -\varphi^{\otimes r} f \pi_r(T'_i) \\
&= -\Phi(f) \pi_r(T'_i).
\end{aligned}$$

Hence  $\Phi(f) \in \mathcal{B}_q^\dagger$ . In the same way, we also obtain  $\Phi(f) \in \mathcal{B}_q$  for  $f \in \mathcal{B}_q^\dagger$ . Combining with Lemma 5.1, we can conclude that  $\Phi$  defines an automorphism of  $\mathcal{D}_q = \mathcal{B}_q \oplus \mathcal{B}_q^\dagger$  satisfying  $\Phi(\mathcal{B}_q) = \mathcal{B}_q^\dagger$  and  $\Phi(\mathcal{B}_q^\dagger) = \mathcal{B}_q$ .  $\square$

Let  $\omega$  be the endomorphism of the algebra  $\text{End}_K V^{\otimes r}$  which is given by:

$$\omega(f) = (-1)^{r(r-1)/2} \varphi^{\otimes r} f \varphi^{\otimes r} \quad \text{for } f \in \text{End}_K V^{\otimes r}.$$

Indeed, one can easily see that the following.

$$\begin{aligned}
\omega(fg) &= (-1)^{r(r-1)/2} \varphi^{\otimes r} fg \varphi^{\otimes r} \\
&= \varphi^{\otimes r} f \varphi^{\otimes r} \varphi^{\otimes r} g \varphi^{\otimes r} \\
&= ((-1)^{r(r-1)/2} \varphi^{\otimes r} f \varphi^{\otimes r}) ((-1)^{r(r-1)/2} \varphi^{\otimes r} g \varphi^{\otimes r}) \\
&= \omega(f) \omega(g).
\end{aligned}$$

So,  $\omega$  is algebraic.

**Proposition 5.4.**  $\omega$  is an automorphism of order 2 of the algebra  $\mathcal{B}_q$ .

*Proof.* If  $f \in \mathcal{B}_q$ , then it follows from (5.3) that

$$\begin{aligned}\pi_r(T'_i)\omega(f) &= (-1)^{r(r-1)/2}\pi_r(T'_i)\varphi^{\otimes r}f\varphi^{\otimes r} \\ &= (-1)^{r(r-1)/2}\pi_r(T'_i)\varphi^{\otimes r}f\varphi^{\otimes r} \\ &= (-1)^{r(r-1)/2}\varphi^{\otimes r}f\varphi^{\otimes r}\pi_r(T'_i) \\ &= \omega(f)\pi_r(T'_i)\end{aligned}$$

Therefore  $\omega(f) \in \mathcal{B}_q$ . From (5.2), it follows that

$$\begin{aligned}\omega^2(f) &= (-1)^{r(r-1)/2}\omega(\varphi^{\otimes r}f\varphi^{\otimes r}) \\ &= (-1)^{r(r-1)}(\varphi^{\otimes r})^2f(\varphi^{\otimes r})^2 \\ &= f\end{aligned}$$

Therefore  $\omega$  is an automorphism of  $\mathcal{B}_q$  of order 2.  $\square$

Let  $H = \langle 1, \omega \rangle$ . Then  $H$  is a subgroup of  $\text{Aut}(\mathcal{B}_q)$ .  $H$  is naturally isomorphic to  $\mathbb{Z}_2 = \langle 1, -1 \rangle$  as (multiplicative) groups. We define two maps

$$\psi_1 : \mathbb{Z}_2 \longrightarrow \text{Aut}(\mathcal{B}_q), \quad \psi_1(1) = 1, \psi_1(-1) = \omega, \quad (5.4)$$

and

$$\alpha_1 : \mathbb{Z}_2 \times \mathbb{Z}_2 \longrightarrow (\mathcal{B}_q)^\times, \quad \alpha_1(\sigma, \tau) = \begin{cases} 1 & \text{if } \sigma = 1 \text{ or } \tau = 1, \\ (-1)^{r(r-1)/2} & \text{otherwise.} \end{cases} \quad (5.5)$$

Then we have the following lemma.

**Lemma 5.5.**  $\psi_1$  and  $\alpha_1$  satisfy (2.1)–(2.3).

*Proof.* (2.3) is trivial. Since  $\alpha(\sigma, \tau) = \pm 1$ , (2.1) may be reduced to  $\sigma(\tau a) = (\sigma\tau)a$ , so holds obviously. If  $\sigma_1 = 1$ , then both sides of (2.2) equal  $\alpha(\sigma_2, \sigma_3)$ . In the same manner, if  $\sigma_2 = 1$  (respectively  $\sigma_3 = 1$ ), then those equal  $\alpha(\sigma_1, \sigma_3)$  (respectively  $\alpha(\sigma_1, \sigma_2)$ ). If  $\sigma_1 = \sigma_2 = \sigma_3 = -1$ , then both sides of (2.2) become  $-1$ . Thus (2.2) holds.  $\square$

Hence, we obtain a crossed product  $\mathcal{B}_{q, \alpha_1}^{\psi_1}[\mathbb{Z}_2]$  from the crossed system  $(\mathcal{B}_q, \mathbb{Z}_2, \psi_1, \alpha_1)$ , where  $\psi_1$  and  $\alpha_1$  are given in (5.4) and (5.5) respectively.

**Theorem 5.6.** If  $m = n$ , then  $\mathcal{D}_q$  is isomorphic to the  $\mathbb{Z}_2$ -crossed product  $\mathcal{B}_{q, \alpha_1}^{\psi_1}[\mathbb{Z}_2]$  as  $K$ -algebras.

*Proof.* Because  $\mathcal{B}_q^\dagger = \varphi^{\otimes r} \mathcal{B}_q = \mathcal{B}_q \varphi^{\otimes r}$  holds from (5.3) and Proposition 5.3,

$$\mathcal{D}_q = \mathcal{B}_q 1 \oplus \mathcal{B}_q \varphi^{\otimes r}$$

follows. Since both  $\mathcal{B}_{q_{\alpha_1}^{\psi_1}}[\mathbb{Z}_2]$  and  $\mathcal{D}_q$  are free left  $\mathcal{B}_q$ -modules, we may define an isomorphism of  $\mathcal{B}_q$ -modules:

$$\iota_1 : \mathcal{B}_{q_{\alpha_1}^{\psi_1}}[\mathbb{Z}_2] \longrightarrow \mathcal{D}_q, \quad \iota_1(u_1) = 1, \iota_1(u_{-1}) = \varphi^{\otimes r}.$$

From (2.4) and (5.4) and (5.5), one can deduce the multiplication law for  $\mathcal{B}_{q_{\alpha_1}^{\psi_1}}[\mathbb{Z}_2]$  as follows.

$$(a_1 u_\sigma)(a_2 u_\tau) = \begin{cases} a_1 a_2 u_1 & \text{if } \sigma = 1 \text{ and } \tau = 1, \\ a_1 a_2 u_{-1} & \text{if } \sigma = 1 \text{ and } \tau = -1, \\ (-1)^{r(r-1)/2} a_1 \varphi^{\otimes r} a_2 \varphi^{\otimes r} u_{-1} & \text{if } \sigma = -1 \text{ and } \tau = 1, \\ a_1 \varphi^{\otimes r} a_2 \varphi^{\otimes r} u_1 & \text{if } \sigma = -1 \text{ and } \tau = -1, \end{cases}$$

where  $a_1, a_2 \in \mathcal{B}_q$ . Therefore, we get four formulas in  $\mathcal{D}_q$ ,

$$\begin{aligned} (a_1 1)(a_2 1) &= a_1 a_2 1, \\ (a_1 1)(a_2 \varphi^{\otimes r}) &= a_1 a_2 \varphi^{\otimes r}, \\ (a_1 \varphi^{\otimes r})(a_2 1) &= (-1)^{r(r-1)/2} a_1 \varphi^{\otimes r} a_2 \varphi^{\otimes r} \varphi^{\otimes r}, \\ (a_1 \varphi^{\otimes r})(a_2 \varphi^{\otimes r}) &= a_1 \varphi^{\otimes r} a_2 \varphi^{\otimes r} 1, \end{aligned}$$

which derive the conclusion that  $f$  is an isomorphism of  $K$ -algebras.  $\square$

For a commutative domain  $R$  and a subalgebra  $A$  of  $\text{Mat}(m, R)$ , we set  $\tilde{A} = \{X \in \text{Mat}(m, R) | XY = YX \text{ for all } Y \in A\}$ . Let  $R_1 = \mathbb{Q}[q, q^{-1}]$  be the algebra of Laurent polynomials. Since matrix elements of  $\rho_r$  are in  $R_1$  by (4.3) and (4.4), we may define  $\mathcal{D}'_q$  to be the subalgebra of  $\text{Mat}((2m)^r, R_1)$  generated by the set  $\{\rho_r(\sigma), \rho_r(q^h), \rho_r(e_i), \rho_r(f_i), \varphi^{\otimes r} | h \in P^*, i \in I\}$ . Similarly by (4.2), we may also define  $\mathcal{C}'_q$  to be the one generated by  $\{I_{(2m)^r}, (q+q^{-1})^2 \pi_r(T'_i T'_i) | i = 2, \dots, r-1\}$  where  $I_{(2m)^r}$  is the identity matrix. Now we shall complete Schur-Weyl reciprocity for the  $q$ -analogue of the alternating group, namely,  $\mathcal{C}_q = \text{End}_{\mathcal{D}_q} V^{\otimes r}$ .

$\mathcal{D}_q = \tilde{\mathcal{C}}_q$  is by definition. We shall show  $\mathcal{C}_q = \tilde{\mathcal{D}}_q$ . The specialization to a nonzero complex number  $t$  is a ring homomorphism  $\varphi_t : R_1 \longrightarrow \mathbb{C}$  with the condition  $\varphi_t(q) = t$ .  $\mathbb{C}$  becomes  $(\mathbb{C}, R_1)$ -bimodule, with  $R_1$  acting from the right via  $\varphi_t$ . If  $t$  is a transcendental number, we can extend the specialization from  $R_1$  to its quotient field  $K$ , namely  $\varphi_t : K \longrightarrow \mathbb{C}$ . Applying the specialization  $\varphi_t$ , we obtain the specialized algebras  $\mathcal{C}_t = \mathbb{C} \otimes_{R_1} \mathcal{C}'_q$  and  $\mathcal{D}_t = \mathbb{C} \otimes_{R_1} \mathcal{D}_q$  which are subalgebras of  $\text{Mat}((2m)^r, \mathbb{C})$ .  $\mathcal{C}_t$  and  $\mathcal{D}_t$  act on the specialized vector space  $\mathbb{C} \otimes_K V^{\otimes r}$  in obvious ways.

**Proposition 5.7.**  $\mathcal{C}_q = \tilde{\mathcal{D}}_q$ .

*Proof.* Obviously  $\mathcal{C}'_q \subseteq \tilde{\mathcal{D}}'_q$ , so  $\text{rank}_{R_1} \mathcal{C}'_q \leq \text{rank}_{R_1} \tilde{\mathcal{D}}'_q$  holds.

Since  $R_1$  is a principal ideal domain, the submodules  $\mathcal{C}'_q$  and  $\mathcal{D}'_q$  of the free  $R_1$ -module  $\text{Mat}((2m)^r, R_1)$  are also free. Assume  $N = \text{rank}_{R_1} \mathcal{C}'_q$ . Let  $X_i(q) (i = 1, \dots, N)$  be a basis of  $\mathcal{C}'_q$  and  $x_i^{k,l}(q) \in R_1$  the  $(k, l)$ -entry of  $X_i(q)$ . Then we immediately have that the specialized elements  $X_i(t) = (x_i^{k,l}(t)) (i = 1, \dots, N)$  generate  $\mathcal{C}_t$  and  $\dim_{\mathbb{C}} \mathcal{C}_t \leq \text{rank}_{R_1} \mathcal{C}'_q$ . Because  $X_i(q)$  are linearly independent,  $\sum_{i=1}^N \alpha_i(q) X_i(q) = 0$  for  $\alpha_1(q), \dots, \alpha_N(q) \in R_1$  implies  $\alpha_1(q) = \dots = \alpha_N(q) = 0$ . But the equation  $\sum_{i=1}^N \alpha_i X_i(t) = 0$  for  $\alpha_1, \dots, \alpha_N \in \mathbb{C}$  may admit a nonzero  $\alpha_i$ . Consider the system of linear equations:

$$\begin{bmatrix} x_1^{1,1}(t) & x_2^{1,1}(t) & \cdots & x_N^{1,1}(t) \\ x_1^{1,2}(t) & x_2^{1,2}(t) & \cdots & x_N^{1,2}(t) \\ \dots & \dots & \dots & \dots \\ x_1^{(2m)^r, (2m)^r-1}(t) & x_2^{(2m)^r, (2m)^r-1}(t) & \cdots & x_N^{(2m)^r, (2m)^r-1}(t) \\ x_1^{(2m)^r, (2m)^r}(t) & x_2^{(2m)^r, (2m)^r}(t) & \cdots & x_N^{(2m)^r, (2m)^r}(t) \end{bmatrix} \times \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \dots \\ \alpha_N \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \dots \\ 0 \\ 0 \end{bmatrix}.$$

Reducing to a common denominator for each row, we may assume that each  $x_i^{k,l}(t) \in \mathbb{Q}[t]$ . Suppose there exists a non-trivial solution. Then the rank of above matrix is less than  $N$ , which implies that every  $N$ th minor determinant equals 0. This is impossible if  $t$  is transcendental. Hence  $\dim_{\mathbb{C}} \mathcal{C}_t = \text{rank}_{R_1} \mathcal{C}'_q$  for a transcendental  $t$ . In the same manner,  $\dim_{\mathbb{C}} \mathcal{D}_t = \text{rank}_{R_1} \mathcal{D}'_q$  if  $t$  is a transcendental number. This argument is valid even if  $R_1$  is replaced by  $K$ , thus we also obtain  $\dim_{\mathbb{C}} \mathcal{C}_t = \dim_K \mathcal{C}_q$ ,  $\dim_{\mathbb{C}} \mathcal{D}_t = \dim_K \mathcal{D}_q$  for a transcendental number  $t$ .

Next we shall show that  $\text{rank}_{R_1} \tilde{\mathcal{D}}'_q = \dim_{\mathbb{C}} \tilde{\mathcal{D}}_t$ . Assume that  $Y(q) = (y^{k,l}(q)) \in \tilde{\mathcal{D}}'_q$ . Then  $Y(q)$  commutes with all  $\rho_r(\sigma), \rho_r(q^h), \rho_r(e_i), \rho_r(f_i), \varphi^{\otimes r}(h \in P^*, i \in I)$ . Hence the matrix elements  $y^{k,l}(q) (k, l = 1, \dots, (2m)^r)$  determine polynomial equations with coefficients in  $R_1$ . We notice that  $\text{rank}_{R_1} \tilde{\mathcal{D}}'_q \leq \dim_{\mathbb{C}} \tilde{\mathcal{D}}_t$  in general; besides the elements of specialized algebra  $\mathbb{C} \otimes_{R_1} \tilde{\mathcal{D}}'_q$ , possibly  $\tilde{\mathcal{D}}_t$  contains another element which commutes with specializations of  $\rho_r(\sigma), \rho_r(q^h), \rho_r(e_i), \rho_r(f_i), \varphi^{\otimes r}(h \in P^*, i \in I)$ . Assume the case. Reducing to a common denominator, we may deduce that  $t$  is a common solution of certain polynomial equations with coefficients in  $\mathbb{Q}$ . This is impossible if  $t$  is transcendental. Thus  $\text{rank}_{R_1} \tilde{\mathcal{C}}'_q = \dim_{\mathbb{C}} \tilde{\mathcal{C}}_t$  for a transcendental  $t$ . In the same manner, we also have  $\text{rank}_{R_1} \tilde{\mathcal{D}}'_q = \dim_{\mathbb{C}} \tilde{\mathcal{D}}_t$  for a transcendental  $t$ . Replacing  $R_1$  by  $K$ , we obtain  $\dim_K \tilde{\mathcal{C}}_q = \dim_{\mathbb{C}} \tilde{\mathcal{C}}_t$  and  $\dim_K \tilde{\mathcal{D}}_q = \dim_{\mathbb{C}} \tilde{\mathcal{D}}_t$ . Considering the specialization  $\varphi_1$ , we obtain the following:

$$\dim_{\mathbb{C}} \mathcal{C}_1 \leq \dim_{\mathbb{C}} \mathcal{C}_t = \text{rank}_{R_1} \mathcal{C}'_q \leq \text{rank}_{R_1} \tilde{\mathcal{D}}'_q = \dim_{\mathbb{C}} \tilde{\mathcal{D}}_t \leq \dim_{\mathbb{C}} \tilde{\mathcal{D}}_1.$$

In [12], Regev has shown  $\mathcal{D}_1 = \tilde{\mathcal{C}}_1$ . Applying double centralizer theorem, we have  $\mathcal{C}_1 = \tilde{\mathcal{D}}_1$ . Therefore

$\text{rank}_{R_1} C'_q = \text{rank}_{R_1} \tilde{D}'_q$  and  $\dim_K C_q = \dim_K \tilde{D}_q$ , which imply  $C_q = \tilde{D}_q$ .  $\square$

From Proposition 5.7 we immediately obtain Schur-Weyl reciprocity for  $\mathcal{H}_{K,r}^1(q)$  as follows.

**Theorem 5.8.**  $\text{End}_{C_q} V^{\otimes r} = \mathcal{D}_q$  and  $\text{End}_{\mathcal{D}_q} V^{\otimes r} = C_q$  hold.

## 6 Schur-Weyl reciprocity for $\mathcal{H}_{\bar{K},r}^1(q)$ in the general case

Henceforth, we do not assume  $m \neq n$  and consider the general case. In this case, results can be obtained in the same way as  $q = 1$  case which is used in [12]. Recall Theorem 4.2:

$$\bar{A}_q = \bigoplus_{\lambda \in H(m,n;r)} \bar{A}_{q,\lambda}, \quad (6.1)$$

where each  $\bar{A}_{q,\lambda} = \pi_r(I_{q,\lambda})(\cong I_{q,\lambda})$  is the image of the homogeneous component  $I_{q,\lambda}$  of  $\mathcal{H}_{\bar{K},r}(q)$  corresponding to  $\lambda \in H(m,n;r)$ . In the same manner as [12], we define  $H_0(m,n;r) = \{\lambda \in \Lambda_r \mid \lambda, \lambda' \in H(m,n;r)\}$  and  $H_1(m,n;r) = H(m,n;r) \setminus H_0(m,n;r)$ . Then we obtain the following from (6.1).

$$\bar{A}_q = \left[ \bigoplus_{\lambda \in H_0(m,n;r), \lambda > \lambda'} \{\bar{A}_{q,\lambda} \oplus \bar{A}_{q,\lambda'}\} \right] \bigoplus \left\{ \bigoplus_{\lambda \in H_0(m,n;r), \lambda = \lambda'} \bar{A}_{q,\lambda} \right\} \bigoplus \left\{ \bigoplus_{\lambda \in H_1(m,n;r)} \bar{A}_{q,\lambda} \right\}. \quad (6.2)$$

Let  $\bar{C}_q = C_q \otimes_{\mathbb{Q}(q)} \bar{K} = \pi_r(\mathcal{H}_{\bar{K},r}^1(q))$ . From Theorem 3.10 we have:

$$\mathcal{H}_{\bar{K},r}^1(q) = \left\{ \bigoplus_{\lambda \in \Lambda_r, \lambda > \lambda'} \tilde{I}_{q,\lambda} \right\} \bigoplus \left[ \bigoplus_{\lambda \in \Lambda_r, \lambda = \lambda'} \{\tilde{I}_{q,\lambda}^+ \oplus \tilde{I}_{q,\lambda}^-\} \right].$$

So we immediately obtain:

$$\bar{C}_q = \left\{ \bigoplus_{\lambda \in H_0(m,n;r), \lambda > \lambda'} \pi_r(\tilde{I}_{q,\lambda}) \right\} \bigoplus \left[ \bigoplus_{\lambda \in H_0(m,n;r), \lambda = \lambda'} \{\pi_r(\tilde{I}_{q,\lambda}^+) \oplus \pi_r(\tilde{I}_{q,\lambda}^-)\} \right] \bigoplus \left\{ \bigoplus_{\lambda \in H_1(m,n;r)} \pi_r(\tilde{I}_{q,\lambda}) \right\}. \quad (6.3)$$

If  $\lambda \in H_0(m,n;r)$  and  $\lambda > \lambda'$ , then  $\pi_r(I_{q,\lambda}) \cong I_{q,\lambda}$  and  $\pi_r(I_{q,\lambda'}) \cong I_{q,\lambda'}$ . Therefore  $\pi_r(I_{q,\lambda} \oplus I_{q,\lambda'}) \cong I_{q,\lambda} \oplus I_{q,\lambda'}$ , which implies  $\pi_r(\tilde{I}_{q,\lambda}) \cong \pi_r(I_{q,\lambda}) \cong I_{q,\lambda}$  because  $I_{q,\lambda} \cong \tilde{I}_{q,\lambda} \subseteq I_{q,\lambda} \oplus I_{q,\lambda'}$ .

If  $\lambda \in H_0(m,n;r)$  and  $\lambda = \lambda'$ , then  $\pi_r(\tilde{I}_{q,\lambda}^+) \cong \tilde{I}_{q,\lambda}^+$  and  $\pi_r(\tilde{I}_{q,\lambda}^-) \cong \tilde{I}_{q,\lambda}^-$  hold because  $\pi_r(\tilde{I}_{q,\lambda}) \cong \tilde{I}_{q,\lambda}$  and  $\tilde{I}_{q,\lambda}^\pm \subseteq \tilde{I}_{q,\lambda}$ .

If  $\lambda \in H_1(m,n;r)$  then  $\pi_r(I_{q,\lambda}) \cong I_{q,\lambda}$  and  $\pi_r(I_{q,\lambda'}) = \{0\}$ . Since  $\tilde{I}_{q,\lambda} = \{X + \hat{X} \in I_{q,\lambda} \oplus I_{q,\lambda'} \mid X \in I_{q,\lambda}\}$ , we have  $\pi_r(X + \hat{X}) = \pi_r(X)$  for  $X + \hat{X} \in \tilde{I}_{q,\lambda}$ . Thereby  $\pi_r(\tilde{I}_{q,\lambda}) = \pi_r(I_{q,\lambda}) \cong I_{q,\lambda}$  follows.

Let us define two-sided ideals of  $\bar{\mathcal{A}}_q$  and those of  $\bar{\mathcal{C}}_q$  as follows.

$$\begin{aligned}\bar{\mathcal{A}}_q \supseteq \bar{\mathcal{A}}_q^0 &= \left[ \bigoplus_{\lambda \in H_0(m,n;r), \lambda > \lambda'} \{ \bar{\mathcal{A}}_{q,\lambda} \oplus \bar{\mathcal{A}}_{q,\lambda'} \} \right] \oplus \left\{ \bigoplus_{\lambda \in H_0(m,n;r), \lambda = \lambda'} \bar{\mathcal{A}}_{q,\lambda} \right\}, \\ \bar{\mathcal{A}}_q \supseteq \bar{\mathcal{A}}_q^1 &= \bigoplus_{\lambda \in H_1(m,n;r)} \bar{\mathcal{A}}_{q,\lambda}, \\ \bar{\mathcal{C}}_q \supseteq \bar{\mathcal{C}}_q^0 &= \left\{ \bigoplus_{\lambda \in H_0(m,n;r), \lambda > \lambda'} \pi_r(\tilde{I}_{q,\lambda}) \right\} \oplus \left[ \bigoplus_{\lambda \in H_0(m,n;r), \lambda = \lambda'} \{ \pi_r(\tilde{I}_{q,\lambda}^+) \oplus \pi_r(\tilde{I}_{q,\lambda}^-) \} \right] \\ \bar{\mathcal{C}}_q \supseteq \bar{\mathcal{C}}_q^1 &= \bigoplus_{\lambda \in H_1(m,n;r)} \pi_r(\tilde{I}_{q,\lambda})\end{aligned}$$

Comparing dimensions of components between  $\bar{\mathcal{A}}_q$  and  $\bar{\mathcal{C}}_q$ , we obtain the following theorem.

**Theorem 6.1.**  $\bar{\mathcal{A}}_q$  and  $\bar{\mathcal{C}}_q$  have direct sum decompositions  $\bar{\mathcal{A}}_q = \bar{\mathcal{A}}_q^0 \oplus \bar{\mathcal{A}}_q^1$  and  $\bar{\mathcal{C}}_q = \bar{\mathcal{C}}_q^0 \oplus \bar{\mathcal{C}}_q^1$  respectively, which satisfy the following relations.

- (1)  $\bar{\mathcal{C}}_q^0 \subseteq \bar{\mathcal{A}}_q^0$  and  $\dim_{\bar{K}} \bar{\mathcal{A}}_q^0 = 2 \dim_{\bar{K}} \bar{\mathcal{C}}_q^0$
- (2)  $\bar{\mathcal{C}}_q^1 = \bar{\mathcal{A}}_q^1$ . Especially  $\dim_{\bar{K}} \bar{\mathcal{A}}_q^1 = \dim_{\bar{K}} \bar{\mathcal{C}}_q^1$ .

As a special case, when  $m = n$ , we readily see that  $H_1(m, n; r) = \emptyset$ . Hence both  $\bar{\mathcal{A}}_q^1$  and  $\bar{\mathcal{C}}_q^1$  are zero and  $\dim_{\bar{K}} \bar{\mathcal{A}}_q = 2 \dim_{\bar{K}} \bar{\mathcal{C}}_q$ . Moreover, if  $n = 0$ , namely non-super case, then we readily see that if  $m^2 < r$ , then all  $\lambda \in \Lambda_r$  are contained in  $H_1(m, 0; r)$  and  $\bar{\mathcal{A}}_q^0 = \bar{\mathcal{C}}_q^0 = 0$ . The Corollary below is immediately obtained from this argument and (6.2), (6.3).

**Corollary 6.2.** Let  $n = 0$ . If  $m^2 < r$ , then  $\bar{\mathcal{A}}_q = \bar{\mathcal{C}}_q$  and  $\text{End}_{\bar{\mathcal{C}}_q} \bar{V}^{\otimes r} = \text{End}_{\bar{\mathcal{A}}_q} \bar{V}^{\otimes r}$ .

We denote by  $W_0$  and  $W_1$  direct summands of  $\bar{V}^{\otimes r}$  which are defined as follows.

$$\begin{aligned}W_0 &= \left[ \bigoplus_{\lambda \in H_0(m,n;r), \lambda > \lambda'} \{ (H_\lambda \otimes V_\lambda) \oplus (H_{\lambda'} \otimes V_{\lambda'}) \} \right] \oplus \left\{ \bigoplus_{\lambda \in H_0(m,n;r), \lambda = \lambda'} (H_\lambda \otimes V_\lambda) \right\} \\ W_1 &= \bigoplus_{\lambda \in H_1(m,n;r)} (H_\lambda \otimes V_\lambda)\end{aligned}$$

We notice that  $\text{End}_{\bar{\mathcal{A}}_q} \bar{V}^{\otimes r} = \text{End}_{\bar{\mathcal{A}}_q} W_0 \oplus \text{End}_{\bar{\mathcal{A}}_q} W_1$  and that  $\text{End}_{\bar{\mathcal{C}}_q} \bar{V}^{\otimes r} = \text{End}_{\bar{\mathcal{C}}_q} W_0 \oplus \text{End}_{\bar{\mathcal{C}}_q} W_1$ . From Theorem 6.1, we immediately have the following.

**Corollary 6.3.**  $\text{End}_{\bar{\mathcal{C}}_q} W_0 \supseteq \text{End}_{\bar{\mathcal{A}}_q} W_0$  and  $\text{End}_{\bar{\mathcal{C}}_q} W_1 = \text{End}_{\bar{\mathcal{A}}_q} W_1$ .

*Proof.* Since  $\bar{\mathcal{A}}_q^1 W_0 = 0$  and  $\bar{\mathcal{A}}_q^0 W_1 = 0$ ,  $\text{End}_{\bar{\mathcal{A}}_q} W_0 = \text{End}_{\bar{\mathcal{A}}_q^0} W_0$  and  $\text{End}_{\bar{\mathcal{A}}_q} W_1 = \text{End}_{\bar{\mathcal{A}}_q^1} W_1$  hold. Similarly, we have  $\text{End}_{\bar{\mathcal{C}}_q} W_0 = \text{End}_{\bar{\mathcal{C}}_q^1} W_0$  and  $\text{End}_{\bar{\mathcal{C}}_q} W_1 = \text{End}_{\bar{\mathcal{C}}_q^1} W_1$ . Hence we immediately obtain from Theorem 6.1(1) that  $\text{End}_{\bar{\mathcal{C}}_q} W_0 = \text{End}_{\bar{\mathcal{C}}_q^0} W_0 \supseteq \text{End}_{\bar{\mathcal{A}}_q^0} W_0 = \text{End}_{\bar{\mathcal{A}}_q} W_0$  and from Theorem 6.1(2) that  $\text{End}_{\bar{\mathcal{C}}_q} W_1 = \text{End}_{\bar{\mathcal{C}}_q^1} W_1 = \text{End}_{\bar{\mathcal{A}}_q^1} W_1 = \text{End}_{\bar{\mathcal{A}}_q} W_1$  also.  $\square$

## References

- [1] C. W. Curtis and I. Reiner, “Methods of Representation Theory”, Vol.1, John Wiley & Sons, 1981.
- [2] G. Benkart, S. Kang and M. Kashiwara, Crystal bases for the quantum superalgebra  $U_q(\mathfrak{gl}(m, n))$ , *J. Amer. Math. Soc.* **13(2)** (2000), 295–331.
- [3] A. Berele and A. Regev, Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras, *Adv. Math.* **64** (1987), 118–175.
- [4] R. Dipper and G.D. James, Representations of Hecke algebras of general linear groups, *Proc. London Math. Soc.* **52** (1986), 20–52.
- [5] R. Dipper and G.D. James, Blocks and idempotents of Hecke algebras of general linear groups, *Proc. London Math. Soc.* **54** (1987), 57–82.
- [6] R. Dipper and G.D. James, The  $q$ -Schur algebra, *Proc. London Math. Soc.* **59** (1989), 23–50.
- [7] V. G. Kac, Lie superalgebras, *Adv. in Math.* **26** (1977), 8–96.
- [8] S. M. Khoroshkin and V. N. Tolstoy, Universal  $R$ -matrix for quantized (super)algebras, *Commun. Math. Phys.* **141** (1991), 599–617.
- [9] H. Mitsuhashi, The  $q$ -analogue of the alternating group and its representations, *J. Algebra* **240** (2001), 535–558.
- [10] H. Mitsuhashi, Schur-Weyl reciprocity between the quantum superalgebra and the Iwahori-Hecke algebra, *preprint* .
- [11] C. Nastasescu and F. V. Oystaeyen, Methods of Graded Rings, Lecture Notes in Math., No. 1836, Springer-Verlag, Berlin, 2004.
- [12] A. Regev, Double centralizing theorems for the alternating groups, *J. Algebra* **250** (2002), 335–352.
- [13] A. N. Sergeev, The tensor algebra of the identity representation as a module over the Lie superalgebras  $\mathfrak{gl}(n, m)$  and  $Q(n)$ , *Math. USSR Sbornik.* **51** (1985), No.2, 419–427.
- [14] H. Yamane, Quantized enveloping algebras associated with simple Lie superalgebras and their universal  $R$ -matrices, *Publ. RIMS.* **30** (1994), 15–87.